Person with disabilities experiencing problems accessing NCHS birth and death tables should contact **nchsed@cdc.gov**, or call 301-458-4688

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
United States												
All races, both sexes	45	-	-	-	1	-	1	2	-	-	-	1
Male	19	-	-	-	1	-	-	1	-	-	-	1
Female	26	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	40	-	-	-	1	-	1	2	-	-	-	1
Male	17	-	-	-	1	-	-	1	-	-	-	1
Female	23	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Arkansas												
All races, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	5	-	-	-	-	-	-	1	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	1	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Florida												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Hawaii												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Illinois												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Iowa												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Maine												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	4	-	-	-	1	-	-	-	-	-	-	-
Male	2	-	-	-	1	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Nebraska												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
New Hampshire												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	3	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
North Carolina												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Ohio												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Texas												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Washington												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Wisconsin												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
United States												
All races, both sexes	-	-	2	3	3	8	7	7	3	6	1	-
Male	-	-	2	1	2	4	3	2	-	2	-	-
Female	-	-	-	2	1	4	4	5	3	4	1	-
White, both sexes	-	-	2	2	3	6	5	7	3	6	1	-
Male	-	-	2	1	2	3	2	2	-	2	-	-
Female	-	-	-	1	1	3	3	5	3	4	1	-
Black, both sexes	-	-	-	1	-	2	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	1	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	1	-	-	-
White, both sexes	-	-	-	-	1	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Florida												
All races, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	1	-	1	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	1	-
White, both sexes	-	-	-	-	-	-	-	1	-	1	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Iowa												
All races, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Maine												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	1	1	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Nebraska												
All races, both sexes	-	-	-	-	1	-	-	-	-	1	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	1	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
North Carolina												
All races, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Ohio												
All races, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Texas												
All races, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Washington												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
United States												
All races, both sexes	65	3	2	-	1	2	1	1	1	3	5	4
Male	22	-	2	-	-	-	-	1	-	2	1	2
Female	43	3	-	-	1	2	1	-	1	1	4	2
White, both sexes	54	3	-	-	1	2	1	1	-	1	4	3
Male	17	-	-	-	-	-	-	1	-	1	1	1
Female	37	3	-	-	1	2	1	-	-	-	3	2
Black, both sexes	9	-	2	-	-	-	-	-	1	1	1	1
Male	4	-	2	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	1	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Arkansas												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Colorado												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Florida												
All races, both sexes	3	-	-	-	-	-	-	-	1	1	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	1	1	1	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	2	-	-	-	-	-	-	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Illinois												
All races, both sexes	4	-	1	-	-	-	-	-	-	-	-	1
Male	2	-	1	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	1	-	-	-	-	-	-	-	-	1
Male	2	-	1	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Iowa												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	2	-	-	-	-	-	1	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	1	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	2	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Massachusetts												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Minnesota												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	2	-	1	-	1	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Missouri												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Nebraska												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
North Carolina												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Ohio												
All races, both sexes	6	1	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	6	1	-	-	-	-	-	-	-	-	1	-
White, both sexes	4	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Oregon												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Rhode Island												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Texas												
All races, both sexes	3	-	-	-	-	1	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	1	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Washington												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Wisconsin												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
United States												
All races, both sexes	2	3	3	4	6	7	6	5	3	1	2	-
Male	1	1	1	2	2	2	3	2	-	-	-	-
Female	1	2	2	2	4	5	3	3	3	1	2	-
White, both sexes	2	3	3	4	5	7	5	4	3	1	1	-
Male	1	1	1	2	2	2	3	1	-	-	-	-
Female	1	2	2	2	3	5	2	3	3	1	1	-
Black, both sexes	-	-	-	-	1	-	-	1	-	-	1	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	2	-	-	1	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	2	-	-	1	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Colorado												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Florida												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	1	-	1	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	1	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Illinois												
All races, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Iowa												
All races, both sexes	1	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	2	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	2	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Massachusetts												
All races, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Minnesota												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Missouri												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Nebraska												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	1	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
North Carolina												
All races, both sexes	-	-	-	-	2	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	2	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Ohio												
All races, both sexes	-	-	-	-	-	1	-	-	2	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	2	-	1	-
White, both sexes	-	-	-	-	-	1	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	1	-	1	1	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	1	-	1	1	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Rhode Island												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Texas												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Washington												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Wisconsin												
All races, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
United States												
All races, both sexes	810,257	624	233	155	189	465	944	1,572	2,498	5,088	10,101	18,669
Male	389,951	354	105	86	99	298	598	1,002	1,632	3,402	6,893	12,802
Female	420,306	270	128	69	90	167	346	570	866	1,686	3,208	5,867
White, both sexes	694,619	399	154	112	133	309	623	1,011	1,609	3,422	7,106	13,580
Male	333,083	223	66	61	74	189	382	658	1,080	2,343	4,991	9,676
Female	361,536	176	88	51	59	120	241	353	529	1,079	2,115	3,904
Black, both sexes	96,752	187	59	30	48	144	285	483	777	1,464	2,607	4,536
Male	47,046	115	32	19	21	101	191	291	472	919	1,624	2,727
Female	49,706	72	27	11	27	43	94	192	305	545	983	1,809
American Indian 1/, both sexes	3,499	13	8	1	-	6	11	24	41	51	122	169
Male	1,919	5	5	1	-	4	5	17	30	32	79	121
Female	1,580	8	3	-	-	2	6	7	11	19	43	48
Asian or Pacific Islander 2/, both sexes	15,387	25	12	12	8	6	25	54	71	151	266	384
Male	7,903	11	2	5	4	4	20	36	50	108	199	278
Female	7,484	14	10	7	4	2	5	18	21	43	67	106
Alabama												
All races, both sexes	15,786	15	4	5	2	8	35	45	66	149	275	492
Male	7,583	8	4	3	-	5	24	26	36	78	174	325
Female	8,203	7	-	2	2	3	11	19	30	71	101	167
White, both sexes	11,920	9	-	3	1	4	22	23	36	82	166	314
Male	5,729	6	-	2	-	3	17	13	22	49	106	207
Female	6,191	3	-	1	1	1	5	10	14	33	60	107
Black, both sexes	3,815	6	4	2	1	4	13	20	28	67	106	177
Male	1,828	2	4	1	-	2	7	12	14	29	67	117
Female	1,987	4	-	1	1	2	6	8	14	38	39	60
American Indian 1/, both sexes	19	-	-	-	-	-	-	1	-	-	3	1
Male	10	-	-	-	-	-	-	1	-	-	1	1
Female	9	-	-	-	-	-	-	-	-	-	2	-
Asian or Pacific Islander 2/, both sexes	32	-	-	-	-	-	-	1	2	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	1	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Alaska												
All races, both sexes	818	-	1	2	-	2	-	1	6	3	25	31
Male	468	-	1	1	-	1	-	-	4	1	18	21
Female	350	-	-	1	-	1	-	1	2	2	7	10
White, both sexes	630	-	-	2	-	2	-	-	1	2	19	23
Male	359	-	-	1	-	1	-	-	1	1	15	16
Female	271	-	-	1	-	1	-	-	-	1	4	7
Black, both sexes	24	-	-	-	-	-	-	-	-	-	2	1
Male	14	-	-	-	-	-	-	-	-	-	1	1
Female	10	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	136	-	1	-	-	-	-	1	5	-	3	5
Male	81	-	1	-	-	-	-	-	3	-	1	3
Female	55	-	-	-	-	-	-	1	2	-	2	2
Asian or Pacific Islander 2/, both sexes	28	-	-	-	-	-	-	-	-	1	1	2
Male	14	-	-	-	-	-	-	-	-	-	1	1
Female	14	-	-	-	-	-	-	-	-	1	-	1
Arizona												
All races, both sexes	13,586	14	3	3	4	10	22	28	45	82	166	298
Male	6,952	5	1	2	2	7	15	20	32	57	116	215
Female	6,634	9	2	1	2	3	7	8	13	25	50	83
White, both sexes	12,805	11	3	2	3	7	15	26	36	70	134	265
Male	6,566	4	1	2	1	5	12	19	25	48	98	193
Female	6,239	7	2	-	2	2	3	7	11	22	36	72
Black, both sexes	372	1	-	1	-	1	4	-	3	6	10	20
Male	180	-	-	-	-	-	2	-	3	5	6	11
Female	192	1	-	1	-	1	2	-	-	1	4	9
American Indian 1/, both sexes	303	2	-	-	-	2	3	2	6	4	20	10
Male	157	1	-	-	-	2	1	1	4	2	11	8
Female	146	1	-	-	-	-	2	1	2	2	9	2
Asian or Pacific Islander 2/, both sexes	106	-	-	-	1	-	-	-	-	2	2	3
Male	49	-	-	-	1	-	-	-	-	2	1	3
Female	57	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Arkansas												
All races, both sexes	9,687	9	6	-	3	7	15	23	37	74	167	283
Male	4,623	7	3	-	3	4	8	13	26	54	121	194
Female	5,064	2	3	-	-	3	7	10	11	20	46	89
White, both sexes	8,288	5	3	-	2	4	10	13	23	61	115	219
Male	3,974	3	1	-	2	2	4	8	20	46	83	155
Female	4,314	2	2	-	-	2	6	5	3	15	32	64
Black, both sexes	1,343	4	3	-	1	3	4	10	13	13	47	64
Male	625	4	2	-	1	2	3	5	6	8	36	39
Female	718	-	1	-	-	1	1	5	7	5	11	25
American Indian 1/, both sexes	26	-	-	-	-	-	-	-	1	-	2	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	1	-	2	-
Asian or Pacific Islander 2/, both sexes	30	-	-	-	-	-	1	-	-	-	3	-
Male	16	-	-	-	-	-	1	-	-	-	2	-
Female	14	-	-	-	-	-	-	-	-	-	1	-
California												
All races, both sexes	82,836	61	24	14	17	48	95	168	246	493	920	1,752
Male	40,323	35	9	8	10	34	68	111	177	337	664	1,246
Female	42,513	26	15	6	7	14	27	57	69	156	256	506
White, both sexes	68,781	46	18	11	15	33	70	108	173	345	673	1,276
Male	33,261	22	8	6	9	23	47	77	126	234	488	913
Female	35,520	24	10	5	6	10	23	31	47	111	185	363
Black, both sexes	6,872	9	5	-	1	10	15	35	48	86	147	297
Male	3,385	8	1	-	-	8	12	17	30	62	98	194
Female	3,487	1	4	-	1	2	3	18	18	24	49	103
American Indian 1/, both sexes	400	-	-	-	-	1	1	3	2	3	11	23
Male	223	-	-	-	-	-	1	3	2	3	10	17
Female	177	-	-	-	-	1	-	-	-	-	1	6
Asian or Pacific Islander 2/, both sexes	6,783	6	1	3	1	4	9	22	23	59	89	156
Male	3,454	5	-	2	1	3	8	14	19	38	68	122
Female	3,329	1	1	1	-	1	1	8	4	21	21	34

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Colorado												
All races, both sexes	8,512	13	5	5	-	4	11	17	22	56	120	187
Male	4,165	8	3	4	-	2	6	10	14	41	77	141
Female	4,347	5	2	1	-	2	5	7	8	15	43	46
White, both sexes	8,107	11	5	5	-	3	8	13	20	53	105	170
Male	3,962	7	3	4	-	1	3	8	12	40	72	131
Female	4,145	4	2	1	-	2	5	5	8	13	33	39
Black, both sexes	294	2	-	-	-	-	3	3	1	3	12	14
Male	148	1	-	-	-	-	3	1	1	1	4	8
Female	146	1	-	-	-	-	-	2	-	2	8	6
American Indian 1/, both sexes	37	-	-	-	-	1	-	-	1	-	1	1
Male	25	-	-	-	-	1	-	-	1	-	1	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	74	-	-	-	-	-	-	1	-	-	2	2
Male	30	-	-	-	-	-	-	1	-	-	-	1
Female	44	-	-	-	-	-	-	-	-	-	2	1
Connecticut												
All races, both sexes	9,454	6	4	3	-	6	11	10	28	50	92	172
Male	4,325	4	1	2	-	5	6	4	15	39	66	121
Female	5,129	2	3	1	-	1	5	6	13	11	26	51
White, both sexes	8,755	6	2	2	-	2	8	10	16	37	67	143
Male	3,970	4	-	1	-	2	5	4	9	29	47	108
Female	4,785	2	2	1	-	-	3	6	7	8	20	35
Black, both sexes	628	-	1	1	-	3	3	-	12	12	23	27
Male	314	-	1	1	-	3	1	-	6	9	17	12
Female	314	-	-	-	-	-	2	-	6	3	6	15
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	63	-	1	-	-	1	-	-	-	1	2	2
Male	35	-	-	-	-	-	-	-	-	1	2	1
Female	28	-	1	-	-	1	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Delaware												
All races, both sexes	2,413	3	2	-	1	1	5	3	6	28	39	75
Male	1,204	1	-	-	1	1	3	1	3	18	25	56
Female	1,209	2	2	-	-	-	2	2	3	10	14	19
White, both sexes	2,007	3	1	-	1	1	3	2	3	22	28	48
Male	986	1	-	-	1	1	1	1	2	14	17	37
Female	1,021	2	1	-	-	-	2	1	1	8	11	11
Black, both sexes	381	-	1	-	-	-	2	1	2	5	10	27
Male	202	-	-	-	-	-	2	-	1	3	7	19
Female	179	-	1	-	-	-	-	1	1	2	3	8
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	1	1	-
Male	9	-	-	-	-	-	-	-	-	1	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	1	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	1	-	-	-
Dist. of Columbia												
All races, both sexes	1,783	1	-	1	-	-	3	3	8	16	27	59
Male	850	1	-	1	-	-	1	1	6	11	18	42
Female	933	-	-	-	-	-	2	2	2	5	9	17
White, both sexes	391	1	-	-	-	-	1	2	-	1	2	5
Male	203	1	-	-	-	-	-	1	-	1	1	3
Female	188	-	-	-	-	-	1	1	-	-	1	2
Black, both sexes	1,374	-	-	1	-	-	2	1	8	14	25	54
Male	640	-	-	1	-	-	1	-	6	9	17	39
Female	734	-	-	-	-	-	1	1	2	5	8	15
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	-	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	1	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Florida												
All races, both sexes	55,198	42	6	11	11	28	58	77	142	304	601	1,106
Male	27,983	21	4	6	6	21	41	47	101	221	412	751
Female	27,215	21	2	5	5	7	17	30	41	83	189	355
White, both sexes	48,910	16	5	10	4	18	37	49	76	193	427	820
Male	24,908	9	3	6	1	14	25	31	56	145	301	586
Female	24,002	7	2	4	3	4	12	18	20	48	126	234
Black, both sexes	5,921	24	1	1	6	10	21	26	63	108	160	276
Male	2,872	11	1	-	4	7	16	14	43	74	100	159
Female	3,049	13	-	1	2	3	5	12	20	34	60	117
American Indian 1/, both sexes	46	-	-	-	-	-	-	-	1	1	2	3
Male	31	-	-	-	-	-	-	-	1	1	1	2
Female	15	-	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	321	2	-	-	1	-	-	2	2	2	12	7
Male	172	1	-	-	1	-	-	2	1	1	10	4
Female	149	1	-	-	-	-	-	-	1	1	2	3
Georgia												
All races, both sexes	21,779	35	12	7	13	25	46	54	81	241	422	720
Male	10,703	22	2	6	7	15	33	34	47	157	267	454
Female	11,076	13	10	1	6	10	13	20	34	84	155	266
White, both sexes	15,513	10	5	4	6	16	20	27	41	127	217	416
Male	7,634	8	-	3	3	10	11	19	25	79	139	282
Female	7,879	2	5	1	3	6	9	8	16	48	78	134
Black, both sexes	6,102	25	4	3	6	9	25	27	40	109	203	298
Male	2,996	14	2	3	4	5	21	15	22	74	127	168
Female	3,106	11	2	-	2	4	4	12	18	35	76	130
American Indian 1/, both sexes	17	-	1	-	-	-	-	-	-	-	2	2
Male	4	-	-	-	-	-	-	-	-	-	1	1
Female	13	-	1	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	147	-	2	-	1	-	1	-	-	5	-	4
Male	69	-	-	-	-	-	1	-	-	4	-	3
Female	78	-	2	-	1	-	-	-	-	1	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Hawaii												
All races, both sexes	3,093	3	2	1	-	1	2	8	12	28	47	71
Male	1,687	1	1	1	-	1	1	8	4	16	31	44
Female	1,406	2	1	-	-	-	1	-	8	12	16	27
White, both sexes	816	1	1	-	-	-	1	2	1	6	17	19
Male	467	-	1	-	-	-	-	2	-	5	11	14
Female	349	1	-	-	-	-	1	-	1	1	6	5
Black, both sexes	26	-	1	-	-	-	-	1	-	-	-	1
Male	18	-	-	-	-	-	-	1	-	-	-	1
Female	8	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2,245	2	-	1	-	1	1	5	11	22	30	51
Male	1,197	1	-	1	-	1	1	5	4	11	20	29
Female	1,048	1	-	-	-	-	-	-	7	11	10	22
Idaho												
All races, both sexes	3,318	4	2	-	-	2	3	5	9	19	30	78
Male	1,665	3	-	-	-	1	1	4	5	8	16	50
Female	1,653	1	2	-	-	1	2	1	4	11	14	28
White, both sexes	3,259	4	2	-	-	2	3	5	8	16	27	77
Male	1,627	3	-	-	-	1	1	4	4	5	14	49
Female	1,632	1	2	-	-	1	2	1	4	11	13	28
Black, both sexes	9	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	27	-	-	-	-	-	-	-	-	3	2	-
Male	16	-	-	-	-	-	-	-	-	3	1	-
Female	11	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	23	-	-	-	-	-	-	-	1	-	-	1
Male	16	-	-	-	-	-	-	-	1	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Illinois												
All races, both sexes	33,974	27	10	7	7	16	36	84	113	218	436	856
Male	15,991	15	5	3	4	13	26	57	69	148	291	577
Female	17,983	12	5	4	3	3	10	27	44	70	145	279
White, both sexes	28,428	17	5	5	7	12	30	57	66	120	266	543
Male	13,199	10	3	2	4	10	22	37	43	88	189	390
Female	15,229	7	2	3	3	2	8	20	23	32	77	153
Black, both sexes	5,135	9	4	1	-	4	4	25	45	94	162	306
Male	2,591	5	2	1	-	3	2	20	25	57	97	182
Female	2,544	4	2	-	-	1	2	5	20	37	65	124
American Indian 1/, both sexes	29	-	-	-	-	-	-	-	-	1	1	1
Male	13	-	-	-	-	-	-	-	-	-	1	1
Female	16	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	382	1	1	1	-	-	2	2	2	3	7	6
Male	188	-	-	-	-	-	2	-	1	3	4	4
Female	194	1	1	1	-	-	-	2	1	-	3	2
Indiana												
All races, both sexes	18,141	16	6	1	7	9	23	40	67	120	264	462
Male	8,589	12	1	1	3	6	16	19	42	78	179	329
Female	9,552	4	5	-	4	3	7	21	25	42	85	133
White, both sexes	16,700	12	6	1	6	8	16	30	51	98	207	391
Male	7,845	9	1	1	3	6	13	14	32	67	143	280
Female	8,855	3	5	-	3	2	3	16	19	31	64	111
Black, both sexes	1,409	4	-	-	-	1	7	10	16	21	56	70
Male	727	3	-	-	-	-	3	5	10	10	35	48
Female	682	1	-	-	-	1	4	5	6	11	21	22
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	27	-	-	-	1	-	-	-	-	1	1	-
Male	14	-	-	-	-	-	-	-	-	1	1	-
Female	13	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Iowa												
All races, both sexes	9,288	5	1	1	-	2	5	10	18	39	86	153
Male	4,337	1	1	-	-	2	3	8	10	26	61	107
Female	4,951	4	-	1	-	-	2	2	8	13	25	46
White, both sexes	9,121	4	1	1	-	2	5	7	17	36	83	139
Male	4,247	1	1	-	-	2	3	6	9	23	60	97
Female	4,874	3	-	1	-	-	2	1	8	13	23	42
Black, both sexes	133	1	-	-	-	-	-	3	1	2	2	12
Male	73	-	-	-	-	-	-	2	1	2	-	9
Female	60	1	-	-	-	-	-	1	-	-	2	3
American Indian 1/, both sexes	14	-	-	-	-	-	-	-	-	1	-	1
Male	8	-	-	-	-	-	-	-	-	1	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	20	-	-	-	-	-	-	-	-	-	1	1
Male	9	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	-	-	-	-	1
Kansas												
All races, both sexes	8,020	5	2	-	2	5	13	9	24	48	87	161
Male	3,670	3	1	-	1	3	8	6	15	37	50	115
Female	4,350	2	1	-	1	2	5	3	9	11	37	46
White, both sexes	7,530	3	2	-	2	4	12	6	21	34	74	144
Male	3,440	1	1	-	1	3	8	5	13	27	45	105
Female	4,090	2	1	-	1	1	4	1	8	7	29	39
Black, both sexes	387	1	-	-	-	-	1	2	3	11	9	11
Male	185	1	-	-	-	-	-	-	2	7	3	5
Female	202	-	-	-	-	-	1	2	1	4	6	6
American Indian 1/, both sexes	67	1	-	-	-	1	-	1	-	2	1	6
Male	33	1	-	-	-	-	-	1	-	2	-	5
Female	34	-	-	-	-	1	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	36	-	-	-	-	-	-	-	-	1	3	-
Male	12	-	-	-	-	-	-	-	-	1	2	-
Female	24	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Kentucky												
All races, both sexes	12,900	6	3	2	4	4	20	35	53	105	238	344
Male	6,195	3	2	-	1	-	14	22	31	60	155	233
Female	6,705	3	1	2	3	4	6	13	22	45	83	111
White, both sexes	11,986	5	2	2	4	4	11	30	43	93	208	300
Male	5,752	2	2	-	1	-	7	19	26	54	142	208
Female	6,234	3	-	2	3	4	4	11	17	39	66	92
Black, both sexes	886	1	1	-	-	-	9	5	10	12	30	43
Male	426	1	-	-	-	-	7	3	5	6	13	24
Female	460	-	1	-	-	-	2	2	5	6	17	19
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	24	-	-	-	-	-	-	-	-	-	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	13,002	15	7	2	9	13	25	34	59	128	235	434
Male	6,360	9	5	2	4	11	17	25	33	81	155	271
Female	6,642	6	2	-	5	2	8	9	26	47	80	163
White, both sexes	9,105	5	3	-	6	7	10	14	28	67	115	232
Male	4,409	4	1	-	3	6	7	10	16	46	75	152
Female	4,696	1	2	-	3	1	3	4	12	21	40	80
Black, both sexes	3,830	10	3	1	3	6	15	17	30	60	119	199
Male	1,917	5	3	1	1	5	10	13	16	35	79	117
Female	1,913	5	-	-	2	1	5	4	14	25	40	82
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	54	-	1	1	-	-	-	3	1	1	1	2
Male	29	-	1	1	-	-	-	2	1	-	1	2
Female	25	-	-	-	-	-	-	1	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Maine												
All races, both sexes	3,763	2	1	1	2	2	5	4	5	16	41	69
Male	1,855	1	1	-	1	-	2	3	3	12	34	48
Female	1,908	1	-	1	1	2	3	1	2	4	7	21
White, both sexes	3,733	2	-	1	2	2	5	3	5	16	38	63
Male	1,834	1	-	-	1	-	2	3	3	12	32	44
Female	1,899	1	-	1	1	2	3	-	2	4	6	19
Black, both sexes	12	-	1	-	-	-	-	1	-	-	2	3
Male	9	-	1	-	-	-	-	-	-	-	1	3
Female	3	-	-	-	-	-	-	1	-	-	1	-
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	-	1	3
Male	8	-	-	-	-	-	-	-	-	-	1	1
Female	3	-	-	-	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	14,655	6	3	4	6	12	19	33	54	126	221	389
Male	6,961	3	2	3	-	8	14	20	35	83	150	246
Female	7,694	3	1	1	6	4	5	13	19	43	71	143
White, both sexes	10,621	3	2	2	4	4	8	16	21	59	107	203
Male	4,991	2	1	1	-	2	6	9	15	40	77	140
Female	5,630	1	1	1	4	2	2	7	6	19	30	63
Black, both sexes	3,793	2	1	2	2	8	11	16	32	66	110	178
Male	1,838	1	1	2	-	6	8	10	19	42	69	101
Female	1,955	1	-	-	2	2	3	6	13	24	41	77
American Indian 1/, both sexes	15	-	-	-	-	-	-	-	-	-	-	2
Male	10	-	-	-	-	-	-	-	-	-	-	2
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	226	1	-	-	-	-	-	1	1	1	4	6
Male	122	-	-	-	-	-	-	1	1	1	4	3
Female	104	1	-	-	-	-	-	-	-	-	-	3

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Massachusetts												
All races, both sexes	16,822	7	4	1	4	15	13	29	38	72	153	306
Male	7,714	3	2	-	2	8	7	17	26	52	115	219
Female	9,108	4	2	1	2	7	6	12	12	20	38	87
White, both sexes	16,004	6	3	1	3	11	11	22	32	61	125	268
Male	7,295	3	1	-	1	6	6	12	21	43	95	194
Female	8,709	3	2	1	2	5	5	10	11	18	30	74
Black, both sexes	614	-	1	-	1	4	2	4	6	7	23	29
Male	307	-	1	-	1	2	1	3	5	5	16	18
Female	307	-	-	-	-	2	1	1	1	2	7	11
American Indian 1/, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	189	1	-	-	-	-	-	3	-	4	5	9
Male	103	-	-	-	-	-	-	2	-	4	4	7
Female	86	1	-	-	-	-	-	1	-	-	1	2
Michigan												
All races, both sexes	31,138	21	3	1	3	21	26	57	105	212	348	718
Male	15,025	11	2	-	2	13	15	37	72	143	250	503
Female	16,113	10	1	1	1	8	11	20	33	69	98	215
White, both sexes	26,332	17	2	1	2	13	15	32	57	114	227	517
Male	12,595	9	1	-	2	6	9	21	39	82	170	373
Female	13,737	8	1	1	-	7	6	11	18	32	57	144
Black, both sexes	4,517	3	-	-	1	8	10	25	46	91	108	195
Male	2,285	2	-	-	-	7	6	16	31	58	71	124
Female	2,232	1	-	-	1	1	4	9	15	33	37	71
American Indian 1/, both sexes	140	1	-	-	-	-	-	-	1	4	7	2
Male	75	-	-	-	-	-	-	-	1	2	5	2
Female	65	1	-	-	-	-	-	-	-	2	2	-
Asian or Pacific Islander 2/, both sexes	149	-	1	-	-	-	1	-	1	3	6	4
Male	70	-	1	-	-	-	-	-	1	1	4	4
Female	79	-	-	-	-	-	1	-	-	2	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Minnesota												
All races, both sexes	10,611	14	8	3	2	7	12	20	24	43	97	203
Male	5,117	9	4	3	1	2	5	15	17	28	66	140
Female	5,494	5	4	-	1	5	7	5	7	15	31	63
White, both sexes	10,218	11	7	2	1	7	9	10	17	33	77	171
Male	4,897	6	4	2	1	2	3	8	13	22	52	120
Female	5,321	5	3	-	-	5	6	2	4	11	25	51
Black, both sexes	218	1	1	-	1	-	2	7	3	6	12	19
Male	126	1	-	-	-	-	2	4	2	5	10	12
Female	92	-	1	-	1	-	-	3	1	1	2	7
American Indian 1/, both sexes	95	1	-	-	-	-	-	2	3	3	5	8
Male	52	1	-	-	-	-	-	2	2	-	2	4
Female	43	-	-	-	-	-	-	-	1	3	3	4
Asian or Pacific Islander 2/, both sexes	80	1	-	1	-	-	1	1	1	1	3	5
Male	42	1	-	1	-	-	-	1	-	1	2	4
Female	38	-	-	-	-	-	1	-	1	-	1	1
Mississippi												
All races, both sexes	10,316	11	2	3	5	12	19	27	40	86	208	328
Male	4,873	6	1	2	2	5	12	18	24	54	133	208
Female	5,443	5	1	1	3	7	7	9	16	32	75	120
White, both sexes	6,791	5	-	3	2	4	6	5	18	37	93	149
Male	3,164	3	-	2	1	1	4	4	13	22	58	98
Female	3,627	2	-	1	1	3	2	1	5	15	35	51
Black, both sexes	3,486	6	2	-	3	8	13	22	22	48	112	175
Male	1,687	3	1	-	1	4	8	14	11	31	74	107
Female	1,799	3	1	-	2	4	5	8	11	17	38	68
American Indian 1/, both sexes	23	-	-	-	-	-	-	-	-	1	2	2
Male	14	-	-	-	-	-	-	-	-	1	1	2
Female	9	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	-	-	-	1	2
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Missouri												
All races, both sexes	18,785	8	6	9	2	9	16	32	70	102	235	459
Male	8,805	4	3	4	1	8	10	21	47	72	147	315
Female	9,980	4	3	5	1	1	6	11	23	30	88	144
White, both sexes	16,753	4	6	7	1	7	14	23	46	75	186	346
Male	7,857	2	3	3	1	6	9	16	34	49	121	251
Female	8,896	2	3	4	-	1	5	7	12	26	65	95
Black, both sexes	1,930	3	-	1	1	2	2	9	22	25	44	110
Male	902	1	-	1	-	2	1	5	12	21	22	62
Female	1,028	2	-	-	1	-	1	4	10	4	22	48
American Indian 1/, both sexes	31	-	-	-	-	-	-	-	1	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	71	1	-	1	-	-	-	-	1	2	5	3
Male	34	1	-	-	-	-	-	-	1	2	4	2
Female	37	-	-	1	-	-	-	-	-	-	1	1
Montana												
All races, both sexes	2,537	1	1	-	1	2	2	5	2	8	27	56
Male	1,319	-	1	-	1	2	-	2	-	6	18	40
Female	1,218	1	-	-	-	-	2	3	2	2	9	16
White, both sexes	2,442	1	-	-	1	2	1	5	1	8	24	52
Male	1,267	-	-	-	1	2	-	2	-	6	17	37
Female	1,175	1	-	-	-	-	1	3	1	2	7	15
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	89	-	1	-	-	-	1	-	1	-	3	4
Male	48	-	1	-	-	-	-	-	-	-	1	3
Female	41	-	-	-	-	-	1	-	1	-	2	1
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Nebraska												
All races, both sexes	4,882	3	3	-	1	1	3	5	7	25	60	81
Male	2,234	-	-	-	-	-	2	3	2	18	44	54
Female	2,648	3	3	-	1	1	1	2	5	7	16	27
White, both sexes	4,678	3	2	-	1	1	3	3	7	21	53	69
Male	2,133	-	-	-	-	-	2	1	2	16	40	46
Female	2,545	3	2	-	1	1	1	2	5	5	13	23
Black, both sexes	168	-	-	-	-	-	-	2	-	3	7	10
Male	83	-	-	-	-	-	-	2	-	2	4	7
Female	85	-	-	-	-	-	-	-	-	1	3	3
American Indian 1/, both sexes	21	-	1	-	-	-	-	-	-	1	-	2
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	1	-	-	-	-	-	-	1	-	1
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	5,869	8	2	2	4	3	16	23	22	55	110	168
Male	3,355	5	-	2	2	1	9	15	16	39	76	106
Female	2,514	3	2	-	2	2	7	8	6	16	34	62
White, both sexes	5,116	6	-	2	3	3	10	14	17	40	79	129
Male	2,957	3	-	2	2	1	6	10	13	30	56	90
Female	2,159	3	-	-	1	2	4	4	4	10	23	39
Black, both sexes	481	2	1	-	-	-	6	7	3	13	20	26
Male	264	2	-	-	-	-	3	4	1	7	13	12
Female	217	-	1	-	-	-	3	3	2	6	7	14
American Indian 1/, both sexes	52	-	-	-	-	-	-	1	1	1	3	2
Male	32	-	-	-	-	-	-	1	1	1	2	1
Female	20	-	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	220	-	1	-	1	-	-	1	1	1	8	11
Male	102	-	-	-	-	-	-	-	1	1	5	3
Female	118	-	1	-	1	-	-	1	-	-	3	8

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
New Hampshire												
All races, both sexes	3,221	3	1	-	1	1	1	5	4	12	30	58
Male	1,539	1	1	-	-	1	-	3	1	8	23	41
Female	1,682	2	-	-	1	-	1	2	3	4	7	17
White, both sexes	3,197	3	1	-	1	1	1	5	4	12	29	55
Male	1,532	1	1	-	-	1	-	3	1	8	23	39
Female	1,665	2	-	-	1	-	1	2	3	4	6	16
Black, both sexes	10	-	-	-	-	-	-	-	-	-	1	2
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	23,866	15	2	4	6	9	18	25	47	100	228	459
Male	11,111	11	-	2	3	6	8	15	40	66	146	325
Female	12,755	4	2	2	3	3	10	10	7	34	82	134
White, both sexes	20,667	7	1	2	4	4	13	13	29	59	147	319
Male	9,558	6	-	1	3	3	6	8	25	39	106	241
Female	11,109	1	1	1	1	1	7	5	4	20	41	78
Black, both sexes	2,737	7	1	1	2	5	5	9	15	36	76	124
Male	1,307	5	-	1	-	3	2	5	12	22	37	72
Female	1,430	2	1	-	2	2	3	4	3	14	39	52
American Indian 1/, both sexes	21	-	-	-	-	-	-	-	-	-	-	2
Male	7	-	-	-	-	-	-	-	-	-	-	2
Female	14	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	441	1	-	1	-	-	-	3	3	5	5	14
Male	239	-	-	-	-	-	-	2	3	5	3	10
Female	202	1	-	1	-	-	-	1	-	-	2	4

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
New Mexico												
All races, both sexes	4,411	6	1	-	-	4	4	9	12	29	41	88
Male	2,239	2	1	-	-	4	1	8	8	18	35	68
Female	2,172	4	-	-	-	-	3	1	4	11	6	20
White, both sexes	4,112	4	-	-	-	3	3	6	8	28	36	80
Male	2,060	2	-	-	-	3	-	6	4	18	30	62
Female	2,052	2	-	-	-	-	3	-	4	10	6	18
Black, both sexes	85	-	-	-	-	1	-	-	1	-	2	1
Male	48	-	-	-	-	1	-	-	1	-	2	-
Female	37	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	195	2	1	-	-	-	-	3	3	1	3	7
Male	121	-	1	-	-	-	-	2	3	-	3	6
Female	74	2	-	-	-	-	-	1	-	1	-	1
Asian or Pacific Islander 2/, both sexes	19	-	-	-	-	-	1	-	-	-	-	-
Male	10	-	-	-	-	-	1	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	59,268	37	12	9	14	32	52	69	123	286	594	998
Male	27,104	24	6	3	5	17	36	47	92	197	416	701
Female	32,164	13	6	6	9	15	16	22	31	89	178	297
White, both sexes	49,588	23	6	6	7	15	31	37	80	184	384	661
Male	22,660	15	2	2	3	8	21	27	61	137	284	493
Female	26,928	8	4	4	4	7	10	10	19	47	100	168
Black, both sexes	8,268	12	5	2	6	17	18	32	40	92	185	304
Male	3,689	8	4	1	2	9	12	20	28	52	109	178
Female	4,579	4	1	1	4	8	6	12	12	40	76	126
American Indian 1/, both sexes	71	-	-	-	-	-	-	-	-	1	2	5
Male	40	-	-	-	-	-	-	-	-	1	2	5
Female	31	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1,341	2	1	1	1	-	3	-	3	9	23	28
Male	715	1	-	-	-	-	3	-	3	7	21	25
Female	626	1	1	1	1	-	-	-	-	2	2	3

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
North Carolina												
All races, both sexes	23,637	25	12	8	5	9	27	53	95	195	338	630
Male	11,450	14	6	5	3	7	12	33	71	125	216	423
Female	12,187	11	6	3	2	2	15	20	24	70	122	207
White, both sexes	18,232	13	7	7	2	3	16	23	50	109	210	377
Male	8,881	7	2	4	1	2	8	16	39	76	145	272
Female	9,351	6	5	3	1	1	8	7	11	33	65	105
Black, both sexes	5,097	10	4	1	3	6	8	28	41	81	120	243
Male	2,401	7	3	1	2	5	2	15	28	46	65	145
Female	2,696	3	1	-	1	1	6	13	13	35	55	98
American Indian 1/, both sexes	211	1	1	-	-	-	2	1	2	3	6	8
Male	116	-	1	-	-	-	1	1	2	1	5	5
Female	95	1	-	-	-	-	1	-	-	2	1	3
Asian or Pacific Islander 2/, both sexes	97	1	-	-	-	-	1	1	2	2	2	2
Male	52	-	-	-	-	-	1	1	2	2	1	1
Female	45	1	-	-	-	-	-	-	-	-	1	1
North Dakota												
All races, both sexes	1,898	3	2	-	-	-	2	6	4	8	16	36
Male	920	2	-	-	-	-	1	3	2	5	10	28
Female	978	1	2	-	-	-	1	3	2	3	6	8
White, both sexes	1,838	3	2	-	-	-	-	5	4	3	13	33
Male	885	2	-	-	-	-	-	2	2	2	7	26
Female	953	1	2	-	-	-	-	3	2	1	6	7
Black, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	56	-	-	-	-	-	2	-	-	4	3	3
Male	31	-	-	-	-	-	1	-	-	2	3	2
Female	25	-	-	-	-	-	1	-	-	2	-	1
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Ohio												
All races, both sexes	35,497	13	14	5	7	15	40	58	115	218	399	838
Male	16,754	8	8	3	4	9	28	36	73	155	263	588
Female	18,743	5	6	2	3	6	12	22	42	63	136	250
White, both sexes	31,608	6	10	4	5	12	27	40	88	170	312	677
Male	14,820	4	6	2	2	6	17	24	57	121	212	492
Female	16,788	2	4	2	3	6	10	16	31	49	100	185
Black, both sexes	3,758	7	4	1	1	3	13	17	27	44	81	159
Male	1,859	4	2	1	1	3	11	11	16	31	48	94
Female	1,899	3	2	-	-	-	2	6	11	13	33	65
American Indian 1/, both sexes	19	-	-	-	-	-	-	-	-	-	1	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	112	-	-	-	1	-	-	1	-	4	5	2
Male	64	-	-	-	1	-	-	1	-	3	3	2
Female	48	-	-	-	-	-	-	-	-	1	2	-
Oklahoma												
All races, both sexes	12,423	8	3	3	3	4	7	30	44	71	161	337
Male	5,918	5	1	2	3	2	3	20	34	48	94	236
Female	6,505	3	2	1	-	2	4	10	10	23	67	101
White, both sexes	10,941	5	1	2	3	3	5	22	30	59	130	246
Male	5,172	2	-	1	3	1	1	14	22	42	73	174
Female	5,769	3	1	1	-	2	4	8	8	17	57	72
Black, both sexes	798	1	1	-	-	1	2	3	8	5	17	63
Male	390	1	-	-	-	1	2	2	7	2	12	44
Female	408	-	1	-	-	-	-	1	1	3	5	19
American Indian 1/, both sexes	633	2	1	1	-	-	-	5	6	7	13	28
Male	341	2	1	1	-	-	-	4	5	4	8	18
Female	292	-	-	-	-	-	-	1	1	3	5	10
Asian or Pacific Islander 2/, both sexes	51	-	-	-	-	-	-	-	-	-	1	-
Male	15	-	-	-	-	-	-	-	-	-	1	-
Female	36	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Oregon												
All races, both sexes	9,287	7	1	1	1	6	8	10	27	42	86	181
Male	4,609	3	-	1	1	3	4	9	22	26	67	133
Female	4,678	4	1	-	-	3	4	1	5	16	19	48
White, both sexes	8,978	6	1	1	1	5	7	7	24	35	81	166
Male	4,430	3	-	1	1	2	3	6	19	20	62	122
Female	4,548	3	1	-	-	3	4	1	5	15	19	44
Black, both sexes	108	-	-	-	-	1	-	2	1	5	3	3
Male	69	-	-	-	-	1	-	2	1	4	3	2
Female	39	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	62	-	-	-	-	-	1	-	-	-	1	4
Male	32	-	-	-	-	-	1	-	-	-	1	2
Female	30	-	-	-	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	139	1	-	-	-	-	-	1	2	2	1	8
Male	78	-	-	-	-	-	-	1	2	2	1	7
Female	61	1	-	-	-	-	-	-	-	-	-	1
Pennsylvania												
All races, both sexes	42,735	13	7	8	3	17	44	60	97	166	359	783
Male	19,913	5	1	1	3	12	26	36	62	102	260	523
Female	22,822	8	6	7	-	5	18	24	35	64	99	260
White, both sexes	38,913	11	6	3	3	11	32	42	65	125	282	621
Male	18,080	4	1	-	3	8	20	25	40	77	207	430
Female	20,833	7	5	3	-	3	12	17	25	48	75	191
Black, both sexes	3,584	1	-	5	-	6	11	16	30	40	70	157
Male	1,708	1	-	1	-	4	5	10	20	24	47	89
Female	1,876	-	-	4	-	2	6	6	10	16	23	68
American Indian 1/, both sexes	16	-	-	-	-	-	-	-	-	-	1	1
Male	10	-	-	-	-	-	-	-	-	-	1	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	222	1	1	-	-	-	1	2	2	1	6	4
Male	115	-	-	-	-	-	1	1	2	1	5	3
Female	107	1	1	-	-	-	-	1	-	-	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Rhode Island												
All races, both sexes	3,418	3	1	-	-	1	-	1	10	14	41	48
Male	1,487	2	1	-	-	-	-	1	8	9	29	40
Female	1,931	1	-	-	-	1	-	-	2	5	12	8
White, both sexes	3,291	2	1	-	-	-	-	1	9	12	36	43
Male	1,425	1	1	-	-	-	-	1	7	8	26	36
Female	1,866	1	-	-	-	-	-	-	2	4	10	7
Black, both sexes	98	-	-	-	-	1	-	-	-	2	5	3
Male	48	-	-	-	-	-	-	-	-	1	3	2
Female	50	-	-	-	-	1	-	-	-	1	2	1
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	1	-	-	1
Male	4	-	-	-	-	-	-	-	1	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	19	1	-	-	-	-	-	-	-	-	-	1
Male	10	1	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	12,336	10	3	2	7	8	14	31	61	116	223	339
Male	6,177	7	1	1	5	5	5	18	28	78	149	213
Female	6,159	3	2	1	2	3	9	13	33	38	74	126
White, both sexes	8,698	5	2	1	5	4	7	15	29	49	132	173
Male	4,362	4	1	1	3	1	2	8	15	33	93	116
Female	4,336	1	1	-	2	3	5	7	14	16	39	57
Black, both sexes	3,585	5	1	1	2	4	7	16	32	66	90	162
Male	1,781	3	-	-	2	4	3	10	13	44	55	93
Female	1,804	2	1	1	-	-	4	6	19	22	35	69
American Indian 1/, both sexes	18	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	35	-	-	-	-	-	-	-	-	1	1	3
Male	23	-	-	-	-	-	-	-	-	1	1	3
Female	12	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
South Dakota												
All races, both sexes	2,220	1	2	-	-	4	1	4	7	6	20	34
Male	1,097	-	2	-	-	1	1	3	3	5	16	23
Female	1,123	1	-	-	-	3	-	1	4	1	4	11
White, both sexes	2,096	-	2	-	-	3	1	3	6	3	11	25
Male	1,032	-	2	-	-	-	1	2	2	2	10	16
Female	1,064	-	-	-	-	3	-	1	4	1	1	9
Black, both sexes	8	-	-	-	-	-	-	1	-	-	1	1
Male	6	-	-	-	-	-	-	1	-	-	1	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	108	1	-	-	-	1	-	-	1	3	8	8
Male	57	-	-	-	-	1	-	-	1	3	5	6
Female	51	1	-	-	-	-	-	-	-	-	3	2
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	19,006	11	4	2	1	7	29	46	77	143	326	567
Male	9,066	8	1	-	-	5	17	29	48	95	220	394
Female	9,940	3	3	2	1	2	12	17	29	48	106	173
White, both sexes	16,087	5	3	1	1	6	20	32	47	98	238	431
Male	7,655	3	1	-	-	5	12	21	29	66	163	302
Female	8,432	2	2	1	1	1	8	11	18	32	75	129
Black, both sexes	2,844	6	1	1	-	1	9	14	29	44	86	135
Male	1,373	5	-	-	-	-	5	8	19	28	55	91
Female	1,471	1	1	1	-	1	4	6	10	16	31	44
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	66	-	-	-	-	-	-	-	1	1	2	1
Male	33	-	-	-	-	-	-	-	-	1	2	1
Female	33	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Texas												
All races, both sexes	52,022	58	22	12	12	36	78	145	180	407	755	1,399
Male	25,552	28	10	5	6	19	52	86	106	270	526	950
Female	26,470	30	12	7	6	17	26	59	74	137	229	449
White, both sexes	44,932	46	18	7	9	27	55	106	124	301	562	1,089
Male	22,092	23	9	2	5	13	33	64	76	205	396	763
Female	22,840	23	9	5	4	14	22	42	48	96	166	326
Black, both sexes	6,487	11	4	4	3	9	22	37	53	100	175	288
Male	3,157	5	1	3	1	6	19	22	28	61	115	171
Female	3,330	6	3	1	2	3	3	15	25	39	60	117
American Indian 1/, both sexes	80	-	-	-	-	-	1	1	-	1	-	7
Male	48	-	-	-	-	-	-	-	-	-	-	6
Female	32	-	-	-	-	-	1	1	-	1	-	1
Asian or Pacific Islander 2/, both sexes	523	1	-	1	-	-	-	1	3	5	18	15
Male	255	-	-	-	-	-	-	-	2	4	15	10
Female	268	1	-	1	-	-	-	1	1	1	3	5
Utah												
All races, both sexes	4,061	7	2	-	2	1	3	13	15	31	48	83
Male	1,992	3	-	-	2	1	2	7	10	22	33	55
Female	2,069	4	2	-	-	-	1	6	5	9	15	28
White, both sexes	3,937	7	2	-	2	1	3	12	14	29	40	74
Male	1,925	3	-	-	2	1	2	7	9	20	28	49
Female	2,012	4	2	-	-	-	1	5	5	9	12	25
Black, both sexes	27	-	-	-	-	-	-	-	-	-	4	2
Male	14	-	-	-	-	-	-	-	-	-	1	2
Female	13	-	-	-	-	-	-	-	-	-	3	-
American Indian 1/, both sexes	35	-	-	-	-	-	-	1	1	1	4	3
Male	23	-	-	-	-	-	-	-	1	1	4	2
Female	12	-	-	-	-	-	-	1	-	-	-	1
Asian or Pacific Islander 2/, both sexes	62	-	-	-	-	-	-	-	-	1	-	4
Male	30	-	-	-	-	-	-	-	-	1	-	2
Female	32	-	-	-	-	-	-	-	-	-	-	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Vermont												
All races, both sexes	1,566	1	-	-	-	-	1	3	4	1	11	18
Male	739	-	-	-	-	-	-	2	3	-	8	16
Female	827	1	-	-	-	-	1	1	1	1	3	2
White, both sexes	1,559	1	-	-	-	-	1	3	4	1	11	18
Male	737	-	-	-	-	-	-	2	3	-	8	16
Female	822	1	-	-	-	-	1	1	1	1	3	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	18,319	19	3	5	6	15	29	54	73	131	255	468
Male	8,919	16	2	3	2	12	17	37	50	88	171	314
Female	9,400	3	1	2	4	3	12	17	23	43	84	154
White, both sexes	14,158	9	2	4	2	9	17	31	38	78	156	300
Male	6,858	8	1	3	1	6	9	23	27	56	108	218
Female	7,300	1	1	1	1	3	8	8	11	22	48	82
Black, both sexes	3,875	9	1	-	4	6	10	22	33	51	90	158
Male	1,914	8	1	-	1	6	7	14	22	30	57	88
Female	1,961	1	-	-	3	-	3	8	11	21	33	70
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	273	1	-	1	-	-	2	1	2	2	9	10
Male	141	-	-	-	-	-	1	-	1	2	6	8
Female	132	1	-	1	-	-	1	1	1	-	3	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Washington												
All races, both sexes	15,015	9	4	4	5	8	7	30	29	61	161	281
Male	7,326	4	1	4	3	5	5	21	18	39	111	184
Female	7,689	5	3	-	2	3	2	9	11	22	50	97
White, both sexes	13,980	6	1	4	5	7	7	24	20	52	135	236
Male	6,796	4	1	4	3	4	5	17	12	32	96	157
Female	7,184	2	-	-	2	3	2	7	8	20	39	79
Black, both sexes	383	-	-	-	-	1	-	3	4	2	15	30
Male	217	-	-	-	-	1	-	2	3	2	10	18
Female	166	-	-	-	-	-	-	1	1	-	5	12
American Indian 1/, both sexes	172	2	1	-	-	-	-	1	2	2	5	7
Male	84	-	-	-	-	-	-	-	1	1	2	4
Female	88	2	1	-	-	-	-	1	1	1	3	3
Asian or Pacific Islander 2/, both sexes	480	1	2	-	-	-	-	2	3	5	6	8
Male	229	-	-	-	-	-	-	2	2	4	3	5
Female	251	1	2	-	-	-	-	-	1	1	3	3
West Virginia												
All races, both sexes	6,849	4	2	2	1	4	7	12	22	38	79	164
Male	3,259	2	2	2	1	3	3	8	12	25	59	111
Female	3,590	2	-	-	-	1	4	4	10	13	20	53
White, both sexes	6,617	3	1	2	1	4	6	12	21	32	75	156
Male	3,150	2	1	2	1	3	3	8	12	21	56	106
Female	3,467	1	-	-	-	1	3	4	9	11	19	50
Black, both sexes	219	1	1	-	-	-	1	-	1	5	4	7
Male	101	-	1	-	-	-	-	-	-	3	3	5
Female	118	1	-	-	-	-	1	-	1	2	1	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	1	-	1
Male	7	-	-	-	-	-	-	-	-	1	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Wisconsin												
All races, both sexes	14,960	9	1	1	5	8	12	17	37	60	133	315
Male	7,192	7	-	-	4	7	9	11	25	42	101	235
Female	7,768	2	1	1	1	1	3	6	12	18	32	80
White, both sexes	14,201	6	-	1	5	7	6	14	29	49	109	256
Male	6,805	4	-	-	4	6	5	9	21	33	84	198
Female	7,396	2	-	1	1	1	1	5	8	16	25	58
Black, both sexes	609	3	1	-	-	1	5	3	5	9	20	52
Male	305	3	-	-	-	1	4	2	2	7	13	32
Female	304	-	1	-	-	-	1	1	3	2	7	20
American Indian 1/, both sexes	83	-	-	-	-	-	-	-	1	2	3	3
Male	46	-	-	-	-	-	-	-	-	2	3	2
Female	37	-	-	-	-	-	-	-	1	-	-	1
Asian or Pacific Islander 2/, both sexes	67	-	-	-	-	-	1	-	2	-	1	4
Male	36	-	-	-	-	-	-	-	2	-	1	3
Female	31	-	-	-	-	-	1	-	-	-	-	1
Wyoming												
All races, both sexes	1,276	1	1	-	-	1	1	2	6	7	20	34
Male	657	1	1	-	-	-	1	2	4	4	11	30
Female	619	-	-	-	-	1	-	-	2	3	9	4
White, both sexes	1,249	1	1	-	-	1	1	1	5	7	18	33
Male	640	1	1	-	-	-	1	1	3	4	10	29
Female	609	-	-	-	-	1	-	-	2	3	8	4
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	17	-	-	-	-	-	-	1	1	-	2	-
Male	10	-	-	-	-	-	-	1	1	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
United States												
All races, both sexes	28,181	36,874	44,509	50,797	65,378	95,622	134,316	148,961	109,479	45,761	9,803	38
Male	19,571	25,488	29,488	31,722	37,989	50,717	62,659	58,855	34,269	10,372	1,522	28
Female	8,610	11,386	15,021	19,075	27,389	44,905	71,657	90,106	75,210	35,389	8,281	10
White, both sexes	20,633	27,727	35,161	40,633	53,822	82,338	119,310	135,445	100,666	41,769	8,631	26
Male	14,819	19,679	23,763	25,890	31,808	44,299	56,362	54,101	31,801	9,493	1,305	20
Female	5,814	8,048	11,398	14,743	22,014	38,039	62,948	81,344	68,865	32,276	7,326	6
Black, both sexes	6,772	8,062	8,149	8,649	9,737	10,936	11,982	10,658	6,919	3,253	1,004	11
Male	4,193	5,076	4,956	4,915	5,149	5,195	4,923	3,545	1,788	623	164	7
Female	2,579	2,986	3,193	3,734	4,588	5,741	7,059	7,113	5,131	2,630	840	4
American Indian 1/, both sexes	221	300	292	361	357	428	426	360	198	88	22	-
Male	157	200	185	217	203	236	196	147	53	18	8	-
Female	64	100	107	144	154	192	230	213	145	70	14	-
Asian or Pacific Islander 2/, both sexes	555	785	907	1,154	1,462	1,920	2,598	2,498	1,696	651	146	1
Male	402	533	584	700	829	987	1,178	1,062	627	238	45	1
Female	153	252	323	454	633	933	1,420	1,436	1,069	413	101	-
Alabama												
All races, both sexes	666	932	988	1,184	1,482	1,986	2,497	2,533	1,665	606	151	-
Male	447	630	641	704	880	994	1,080	916	468	117	23	-
Female	219	302	347	480	602	992	1,417	1,617	1,197	489	128	-
White, both sexes	405	626	681	870	1,098	1,535	2,004	2,111	1,356	474	100	-
Male	272	426	453	522	678	789	890	788	377	95	14	-
Female	133	200	228	348	420	746	1,114	1,323	979	379	86	-
Black, both sexes	258	301	302	309	380	446	485	419	306	131	50	-
Male	173	200	184	181	199	201	188	127	89	22	9	-
Female	85	101	118	128	181	245	297	292	217	109	41	-
American Indian 1/, both sexes	-	2	2	1	1	2	2	1	1	1	1	-
Male	-	1	2	-	1	2	-	-	1	-	-	-
Female	-	1	-	1	-	-	2	1	-	1	1	-
Asian or Pacific Islander 2/, both sexes	3	3	3	4	3	3	6	2	2	-	-	-
Male	2	3	2	1	2	2	2	1	1	-	-	-
Female	1	-	1	3	1	1	4	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Alaska												
All races, both sexes	47	73	53	73	86	106	98	112	72	23	4	-
Male	35	50	43	43	63	67	51	43	21	4	1	-
Female	12	23	10	30	23	39	47	69	51	19	3	-
White, both sexes	33	61	48	54	59	74	81	87	62	19	3	-
Male	25	41	40	35	42	46	40	34	18	3	-	-
Female	8	20	8	19	17	28	41	53	44	16	3	-
Black, both sexes	4	1	1	1	3	5	-	3	3	-	-	-
Male	4	1	-	1	3	2	-	-	1	-	-	-
Female	-	-	1	-	-	3	-	3	2	-	-	-
American Indian 1/, both sexes	9	7	2	16	19	25	16	18	5	3	1	-
Male	6	6	1	6	14	17	11	9	2	-	1	-
Female	3	1	1	10	5	8	5	9	3	3	-	-
Asian or Pacific Islander 2/, both sexes	1	4	2	2	5	2	1	4	2	1	-	-
Male	-	2	2	1	4	2	-	-	-	1	-	-
Female	1	2	-	1	1	-	1	4	2	-	-	-
Arizona												
All races, both sexes	460	638	839	922	1,174	1,688	2,273	2,498	1,655	657	105	2
Male	326	449	557	584	721	950	1,093	1,044	571	165	18	2
Female	134	189	282	338	453	738	1,180	1,454	1,084	492	87	-
White, both sexes	407	559	784	854	1,101	1,606	2,181	2,413	1,598	628	100	2
Male	292	399	525	548	685	907	1,056	1,013	557	156	18	2
Female	115	160	259	306	416	699	1,125	1,400	1,041	472	82	-
Black, both sexes	27	41	27	36	37	40	35	38	29	12	4	-
Male	17	24	18	17	18	17	16	14	7	5	-	-
Female	10	17	9	19	19	23	19	24	22	7	4	-
American Indian 1/, both sexes	21	27	19	22	26	31	43	32	18	15	-	-
Male	14	18	9	14	14	20	18	13	5	2	-	-
Female	7	9	10	8	12	11	25	19	13	13	-	-
Asian or Pacific Islander 2/, both sexes	5	11	9	10	10	11	14	15	10	2	1	-
Male	3	8	5	5	4	6	3	4	2	2	-	-
Female	2	3	4	5	6	5	11	11	8	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Arkansas												
All races, both sexes	384	513	585	692	837	1,162	1,536	1,606	1,167	469	112	-
Male	254	345	359	428	497	621	674	566	337	92	17	-
Female	130	168	226	264	340	541	862	1,040	830	377	95	-
White, both sexes	273	389	493	584	727	1,012	1,373	1,439	1,043	414	86	-
Male	190	269	314	359	442	553	610	516	299	85	13	-
Female	83	120	179	225	285	459	763	923	744	329	73	-
Black, both sexes	110	120	86	107	104	140	157	161	119	52	25	-
Male	63	74	44	68	53	62	61	49	35	6	4	-
Female	47	46	42	39	51	78	96	112	84	46	21	-
American Indian 1/, both sexes	1	3	3	1	2	4	4	4	1	-	-	-
Male	1	1	-	1	1	2	2	-	-	-	-	-
Female	-	2	3	-	1	2	2	4	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	3	-	4	6	2	2	4	3	1	-
Male	-	1	1	-	1	4	1	1	3	1	-	-
Female	-	-	2	-	3	2	1	1	1	2	1	-
California												
All races, both sexes	2,828	3,584	4,376	4,826	6,324	9,312	13,867	15,848	11,757	5,137	1,135	4
Male	2,027	2,507	2,919	3,023	3,673	4,921	6,388	6,645	3,978	1,348	192	3
Female	801	1,077	1,457	1,803	2,651	4,391	7,479	9,203	7,779	3,789	943	1
White, both sexes	2,081	2,683	3,413	3,748	4,950	7,690	11,798	13,732	10,405	4,532	978	3
Male	1,523	1,945	2,294	2,391	2,903	4,107	5,483	5,796	3,531	1,168	165	2
Female	558	738	1,119	1,357	2,047	3,583	6,315	7,936	6,874	3,364	813	1
Black, both sexes	493	590	588	591	708	729	821	852	504	265	78	-
Male	308	350	369	340	401	348	335	308	131	57	8	-
Female	185	240	219	251	307	381	486	544	373	208	70	-
American Indian 1/, both sexes	22	26	37	39	45	49	47	51	26	12	2	-
Male	20	21	21	27	28	20	19	20	9	2	-	-
Female	2	5	16	12	17	29	28	31	17	10	2	-
Asian or Pacific Islander 2/, both sexes	232	285	338	448	621	844	1,201	1,213	822	328	77	1
Male	176	191	235	265	341	446	551	521	307	121	19	1
Female	56	94	103	183	280	398	650	692	515	207	58	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Colorado												
All races, both sexes	319	402	452	513	677	998	1,368	1,633	1,137	469	104	-
Male	240	276	302	319	408	529	609	658	378	117	23	-
Female	79	126	150	194	269	469	759	975	759	352	81	-
White, both sexes	295	365	425	485	636	949	1,315	1,578	1,109	456	101	-
Male	223	252	285	305	384	507	590	634	363	116	22	-
Female	72	113	140	180	252	442	725	944	746	340	79	-
Black, both sexes	20	32	18	21	33	37	29	35	18	10	3	-
Male	14	21	13	11	20	14	10	14	10	1	1	-
Female	6	11	5	10	13	23	19	21	8	9	2	-
American Indian 1/, both sexes	2	2	8	4	3	5	3	2	3	1	-	-
Male	1	2	4	3	2	5	1	2	1	-	-	-
Female	1	-	4	1	1	-	2	-	2	1	-	-
Asian or Pacific Islander 2/, both sexes	2	3	1	3	5	7	21	18	7	2	-	-
Male	2	1	-	-	2	3	8	8	4	-	-	-
Female	-	2	1	3	3	4	13	10	3	2	-	-
Connecticut												
All races, both sexes	234	314	406	440	598	1,097	1,516	2,079	1,511	712	164	1
Male	173	227	259	264	344	590	733	817	468	161	25	1
Female	61	87	147	176	254	507	783	1,262	1,043	551	139	-
White, both sexes	188	265	339	365	534	1,009	1,446	1,997	1,469	691	158	1
Male	143	195	219	222	306	540	702	794	458	158	23	1
Female	45	70	120	143	228	469	744	1,203	1,011	533	135	-
Black, both sexes	39	46	57	69	57	79	63	74	37	19	6	-
Male	26	29	33	39	35	45	28	18	6	3	2	-
Female	13	17	24	30	22	34	35	56	31	16	4	-
American Indian 1/, both sexes	1	2	1	-	1	1	1	-	-	1	-	-
Male	1	2	1	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	6	1	9	6	6	8	6	8	5	1	-	-
Male	3	1	6	3	2	4	3	5	4	-	-	-
Female	3	-	3	3	4	4	3	3	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Delaware												
All races, both sexes	93	112	124	154	218	293	404	392	303	126	29	2
Male	67	80	94	102	144	143	187	152	101	20	4	1
Female	26	32	30	52	74	150	217	240	202	106	25	1
White, both sexes	66	80	95	112	173	246	355	358	277	106	25	2
Male	48	59	71	73	116	122	169	139	93	18	2	1
Female	18	21	24	39	57	124	186	219	184	88	23	1
Black, both sexes	26	31	29	41	41	43	45	31	23	19	4	-
Male	18	20	23	28	25	20	18	10	5	1	2	-
Female	8	11	6	13	16	23	27	21	18	18	2	-
American Indian 1/, both sexes	1	-	-	1	2	2	2	1	1	1	-	-
Male	1	-	-	1	1	1	-	1	1	1	-	-
Female	-	-	-	-	1	1	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	2	2	2	2	2	-	-	-
Male	-	1	-	-	2	-	-	2	2	-	-	-
Female	-	-	-	-	-	2	2	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	83	106	134	166	157	193	249	251	191	102	33	-
Male	57	60	87	96	85	100	96	96	57	27	8	-
Female	26	46	47	70	72	93	153	155	134	75	25	-
White, both sexes	7	14	14	22	27	41	64	70	68	37	15	-
Male	6	9	11	15	16	29	36	36	24	10	4	-
Female	1	5	3	7	11	12	28	34	44	27	11	-
Black, both sexes	76	91	120	143	127	148	182	180	121	63	18	-
Male	51	51	76	80	66	71	59	60	33	16	4	-
Female	25	40	44	63	61	77	123	120	88	47	14	-
American Indian 1/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	3	4	2	1	2	2	-	-
Male	-	-	-	-	3	-	1	-	-	1	-	-
Female	-	1	-	-	-	4	1	1	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Florida												
All races, both sexes	1,750	2,181	2,753	3,310	4,340	6,825	9,549	10,650	7,606	3,197	648	3
Male	1,210	1,494	1,828	2,135	2,576	3,844	4,820	4,629	2,818	870	126	2
Female	540	687	925	1,175	1,764	2,981	4,729	6,021	4,788	2,327	522	1
White, both sexes	1,264	1,718	2,256	2,719	3,651	6,079	8,763	10,003	7,201	3,009	590	2
Male	932	1,218	1,540	1,789	2,201	3,499	4,488	4,418	2,704	828	113	1
Female	332	500	716	930	1,450	2,580	4,275	5,585	4,497	2,181	477	1
Black, both sexes	467	444	473	551	642	689	737	606	382	176	57	1
Male	264	261	273	319	347	317	310	197	107	35	12	1
Female	203	183	200	232	295	372	427	409	275	141	45	-
American Indian 1/, both sexes	1	6	3	4	4	7	4	5	3	1	1	-
Male	1	4	2	4	2	4	2	4	1	1	1	-
Female	-	2	1	-	2	3	2	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes	18	13	21	36	43	50	45	36	20	11	-	-
Male	13	11	13	23	26	24	20	10	6	6	-	-
Female	5	2	8	13	17	26	25	26	14	5	-	-
Georgia												
All races, both sexes	1,076	1,401	1,575	1,738	1,964	2,439	3,168	3,291	2,325	953	191	2
Male	732	956	1,031	1,082	1,131	1,266	1,420	1,168	667	181	24	1
Female	344	445	544	656	833	1,173	1,748	2,123	1,658	772	167	1
White, both sexes	606	829	978	1,157	1,360	1,810	2,501	2,654	1,890	709	129	1
Male	425	585	667	753	830	955	1,166	969	553	133	18	1
Female	181	244	311	404	530	855	1,335	1,685	1,337	576	111	-
Black, both sexes	461	563	581	569	588	603	635	626	423	242	61	1
Male	301	365	357	322	293	301	241	196	111	48	6	-
Female	160	198	224	247	295	302	394	430	312	194	55	1
American Indian 1/, both sexes	1	-	2	-	2	3	3	1	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	1	-	1	-	1	3	3	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	9	14	12	14	23	29	10	12	2	1	-
Male	6	6	6	7	7	10	13	3	3	-	-	-
Female	2	3	8	5	7	13	16	7	9	2	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Hawaii												
All races, both sexes	98	156	170	222	214	349	501	608	408	156	36	-
Male	73	113	116	141	137	200	277	278	169	59	16	-
Female	25	43	54	81	77	149	224	330	239	97	20	-
White, both sexes	29	44	59	65	55	86	114	160	103	41	12	-
Male	23	31	48	51	37	51	60	70	48	10	5	-
Female	6	13	11	14	18	35	54	90	55	31	7	-
Black, both sexes	2	6	2	-	3	3	1	5	-	-	1	-
Male	2	4	1	-	1	2	1	4	-	-	1	-
Female	-	2	1	-	2	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	1	1	1	-	-	1	-	1	1	-	-
Male	-	1	1	1	-	-	1	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	67	105	108	156	156	260	385	443	304	114	23	-
Male	48	77	66	89	99	147	215	204	121	48	10	-
Female	19	28	42	67	57	113	170	239	183	66	13	-
Idaho												
All races, both sexes	99	142	169	209	272	381	493	684	497	191	29	-
Male	71	103	126	134	168	197	261	287	172	56	2	-
Female	28	39	43	75	104	184	232	397	325	135	27	-
White, both sexes	95	138	165	204	266	374	483	676	495	191	28	-
Male	69	102	124	130	165	194	253	281	171	56	1	-
Female	26	36	41	74	101	180	230	395	324	135	27	-
Black, both sexes	1	1	-	1	-	3	-	2	-	-	-	-
Male	-	1	-	1	-	2	-	1	-	-	-	-
Female	1	-	-	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	2	2	2	2	4	2	3	2	2	-	1	-
Male	1	-	1	1	3	1	2	1	1	-	1	-
Female	1	2	1	1	1	1	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	2	2	2	2	7	4	-	-	-	-
Male	1	-	1	2	-	-	6	4	-	-	-	-
Female	-	1	1	-	2	2	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Illinois												
All races, both sexes	1,202	1,632	1,744	2,060	2,692	3,822	5,615	6,116	4,767	2,066	445	3
Male	818	1,104	1,129	1,362	1,556	1,936	2,596	2,380	1,424	415	62	1
Female	384	528	615	698	1,136	1,886	3,019	3,736	3,343	1,651	383	2
White, both sexes	830	1,136	1,294	1,545	2,084	3,189	4,919	5,559	4,437	1,903	403	1
Male	593	787	857	1,040	1,223	1,635	2,303	2,183	1,336	390	54	-
Female	237	349	437	505	861	1,554	2,616	3,376	3,101	1,513	349	1
Black, both sexes	360	471	423	488	566	581	620	490	289	153	38	2
Male	215	299	257	303	309	276	259	172	75	24	7	1
Female	145	172	166	185	257	305	361	318	214	129	31	1
American Indian 1/, both sexes	3	4	1	1	5	4	3	4	-	-	1	-
Male	2	1	1	-	3	1	1	1	-	-	1	-
Female	1	3	-	1	2	3	2	3	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	21	26	26	37	48	73	63	41	10	3	-
Male	8	17	14	19	21	24	33	24	13	1	-	-
Female	1	4	12	7	16	24	40	39	28	9	3	-
Indiana												
All races, both sexes	637	883	990	1,159	1,502	2,132	3,020	3,194	2,394	996	219	-
Male	449	587	643	691	903	1,104	1,373	1,242	694	192	25	-
Female	188	296	347	468	599	1,028	1,647	1,952	1,700	804	194	-
White, both sexes	522	763	865	1,043	1,368	1,971	2,810	3,057	2,304	956	215	-
Male	372	509	568	625	824	1,021	1,285	1,186	676	185	25	-
Female	150	254	297	418	544	950	1,525	1,871	1,628	771	190	-
Black, both sexes	110	119	123	113	129	157	206	136	88	39	4	-
Male	74	77	74	65	75	80	87	56	18	7	-	-
Female	36	42	49	48	54	77	119	80	70	32	4	-
American Indian 1/, both sexes	1	-	-	1	-	-	1	1	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	1	2	2	5	4	3	-	2	1	-	-
Male	2	1	1	1	4	3	-	-	-	-	-	-
Female	2	-	1	1	1	1	3	-	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Iowa												
All races, both sexes	251	316	385	453	689	1,048	1,536	1,781	1,614	735	160	-
Male	186	231	260	288	398	580	779	717	509	153	17	-
Female	65	85	125	165	291	468	757	1,064	1,105	582	143	-
White, both sexes	236	300	369	439	669	1,035	1,519	1,766	1,601	732	160	-
Male	178	225	254	279	386	570	769	712	502	153	17	-
Female	58	75	115	160	283	465	750	1,054	1,099	579	143	-
Black, both sexes	13	14	11	10	17	10	12	12	11	2	-	-
Male	7	5	4	6	11	8	8	4	6	-	-	-
Female	6	9	7	4	6	2	4	8	5	2	-	-
American Indian 1/, both sexes	1	1	1	2	1	1	2	1	1	1	-	-
Male	1	-	1	2	-	1	1	-	-	-	-	-
Female	-	1	-	-	1	-	1	1	1	1	-	-
Asian or Pacific Islander 2/, both sexes	1	1	4	2	2	2	3	2	1	-	-	-
Male	-	1	1	1	1	1	1	1	1	-	-	-
Female	1	-	3	1	1	1	2	1	-	-	-	-
Kansas												
All races, both sexes	239	255	346	427	576	893	1,312	1,560	1,305	594	157	-
Male	165	168	230	257	340	461	622	619	420	130	19	-
Female	74	87	116	170	236	432	690	941	885	464	138	-
White, both sexes	208	220	321	379	533	826	1,257	1,505	1,248	579	152	-
Male	146	145	216	229	318	429	601	599	405	125	18	-
Female	62	75	105	150	215	397	656	906	843	454	134	-
Black, both sexes	28	31	20	32	35	53	45	43	46	11	5	-
Male	18	20	12	20	18	25	18	17	13	5	1	-
Female	10	11	8	12	17	28	27	26	33	6	4	-
American Indian 1/, both sexes	1	3	5	10	4	9	6	8	5	4	-	-
Male	1	2	2	6	3	5	2	2	1	-	-	-
Female	-	1	3	4	1	4	4	6	4	4	-	-
Asian or Pacific Islander 2/, both sexes	2	1	-	6	4	5	4	4	6	-	-	-
Male	-	1	-	2	1	2	1	1	1	-	-	-
Female	2	-	-	4	3	3	3	3	5	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Kentucky												
All races, both sexes	522	707	809	989	1,163	1,605	2,074	2,077	1,447	582	111	-
Male	372	479	513	608	660	829	940	734	394	129	16	-
Female	150	228	296	381	503	776	1,134	1,343	1,053	453	95	-
White, both sexes	452	617	752	904	1,073	1,510	1,955	1,990	1,366	564	101	-
Male	337	416	478	561	614	786	888	699	369	127	16	-
Female	115	201	274	343	459	724	1,067	1,291	997	437	85	-
Black, both sexes	68	87	54	81	87	93	114	85	78	18	10	-
Male	35	60	33	44	44	41	50	34	24	2	-	-
Female	33	27	21	37	43	52	64	51	54	16	10	-
American Indian 1/, both sexes	-	2	-	1	-	-	-	-	1	-	-	-
Male	-	2	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	3	3	3	2	5	2	2	-	-	-
Male	-	1	2	3	2	2	2	1	-	-	-	-
Female	2	-	1	-	1	-	3	1	2	-	-	-
Louisiana												
All races, both sexes	630	829	900	1,000	1,178	1,504	2,098	1,972	1,329	486	114	1
Male	425	571	555	607	657	765	940	721	375	106	24	1
Female	205	258	345	393	521	739	1,158	1,251	954	380	90	-
White, both sexes	338	477	530	625	803	1,103	1,629	1,585	1,063	388	77	-
Male	247	335	345	375	463	593	734	583	314	86	14	-
Female	91	142	185	250	340	510	895	1,002	749	302	63	-
Black, both sexes	287	346	365	370	365	394	463	382	261	97	36	1
Male	174	232	208	227	191	170	204	135	60	20	10	1
Female	113	114	157	143	174	224	259	247	201	77	26	-
American Indian 1/, both sexes	-	3	2	1	1	2	1	1	1	-	-	-
Male	-	1	1	1	-	1	-	1	-	-	-	-
Female	-	2	1	-	1	1	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	5	3	3	4	9	5	5	4	4	1	1	-
Male	4	3	1	4	3	1	2	2	1	-	-	-
Female	1	-	2	-	6	4	3	2	3	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Maine												
All races, both sexes	109	136	209	217	305	490	594	726	543	233	53	-
Male	85	93	152	145	191	273	273	281	185	61	11	-
Female	24	43	57	72	114	217	321	445	358	172	42	-
White, both sexes	108	135	207	216	303	486	591	725	541	231	53	-
Male	84	92	151	145	189	269	272	280	184	59	11	-
Female	24	43	56	71	114	217	319	445	357	172	42	-
Black, both sexes	1	1	1	-	-	1	-	-	-	1	-	-
Male	1	1	-	-	-	1	-	-	-	1	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	2	1	2	1	-	-	-	-
Male	-	-	1	-	2	1	1	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	2	1	-	2	1	-	-
Male	-	-	-	-	-	2	-	-	1	1	-	-
Female	-	-	-	1	-	-	1	-	1	-	-	-
Maryland												
All races, both sexes	586	700	867	927	1,247	1,680	2,320	2,629	1,845	816	164	1
Male	394	469	581	553	700	861	1,048	1,023	555	186	26	1
Female	192	231	286	374	547	819	1,272	1,606	1,290	630	138	-
White, both sexes	320	379	500	579	799	1,244	1,813	2,168	1,577	680	132	1
Male	229	258	353	361	460	661	846	852	496	160	21	1
Female	91	121	147	218	339	583	967	1,316	1,081	520	111	-
Black, both sexes	263	307	344	326	420	398	477	418	254	128	30	-
Male	162	201	212	176	223	183	192	150	51	25	4	-
Female	101	106	132	150	197	215	285	268	203	103	26	-
American Indian 1/, both sexes	-	-	1	-	2	5	3	1	1	-	-	-
Male	-	-	-	-	2	3	2	1	-	-	-	-
Female	-	-	1	-	-	2	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	3	14	22	22	26	33	27	42	13	8	2	-
Male	3	10	16	16	15	14	8	20	8	1	1	-
Female	-	4	6	6	11	19	19	22	5	7	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Massachusetts												
All races, both sexes	438	567	699	835	1,190	1,890	2,912	3,544	2,681	1,154	269	1
Male	315	410	464	532	739	1,017	1,311	1,384	805	249	36	1
Female	123	157	235	303	451	873	1,601	2,160	1,876	905	233	-
White, both sexes	392	520	631	754	1,095	1,795	2,813	3,452	2,622	1,123	263	1
Male	286	383	424	483	683	972	1,264	1,355	787	242	33	1
Female	106	137	207	271	412	823	1,549	2,097	1,835	881	230	-
Black, both sexes	32	35	53	58	71	71	71	73	46	22	5	-
Male	20	20	30	36	38	36	37	20	14	2	2	-
Female	12	15	23	22	33	35	34	53	32	20	3	-
American Indian 1/, both sexes	4	1	2	2	2	2	1	-	-	1	-	-
Male	2	1	2	1	2	1	-	-	-	-	-	-
Female	2	-	-	1	-	1	1	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	10	11	13	21	22	22	27	19	13	8	1	-
Male	7	6	8	12	16	8	10	9	4	5	1	-
Female	3	5	5	9	6	14	17	10	9	3	-	-
Michigan												
All races, both sexes	1,208	1,429	1,755	1,913	2,386	3,699	5,170	5,778	4,247	1,675	363	-
Male	844	991	1,180	1,197	1,326	1,950	2,423	2,300	1,329	382	55	-
Female	364	438	575	716	1,060	1,749	2,747	3,478	2,918	1,293	308	-
White, both sexes	826	984	1,341	1,484	1,939	3,165	4,529	5,271	3,920	1,557	319	-
Male	612	705	922	958	1,106	1,683	2,173	2,100	1,224	352	48	-
Female	214	279	419	526	833	1,482	2,356	3,171	2,696	1,205	271	-
Black, both sexes	365	425	396	404	419	500	592	474	305	110	40	-
Male	220	272	246	224	201	249	238	187	98	30	5	-
Female	145	153	150	180	218	251	354	287	207	80	35	-
American Indian 1/, both sexes	12	10	11	9	14	20	22	16	8	1	2	-
Male	9	7	7	5	9	11	6	8	2	-	1	-
Female	3	3	4	4	5	9	16	8	6	1	1	-
Asian or Pacific Islander 2/, both sexes	5	10	7	16	14	14	27	17	14	7	2	-
Male	3	7	5	10	10	7	6	5	5	-	1	-
Female	2	3	2	6	4	7	21	12	9	7	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Minnesota												
All races, both sexes	295	375	457	518	691	1,132	1,662	2,234	1,870	793	151	-
Male	217	289	308	336	419	659	850	946	607	176	20	-
Female	78	86	149	182	272	473	812	1,288	1,263	617	131	-
White, both sexes	262	326	427	485	658	1,092	1,622	2,212	1,852	788	149	-
Male	196	253	293	313	402	640	832	938	602	175	20	-
Female	66	73	134	172	256	452	790	1,274	1,250	613	129	-
Black, both sexes	24	29	17	14	13	20	24	9	12	2	2	-
Male	15	21	9	11	6	8	12	4	4	-	-	-
Female	9	8	8	3	7	12	12	5	8	2	2	-
American Indian 1/, both sexes	5	12	10	13	10	11	6	2	2	2	-	-
Male	3	8	5	8	7	7	2	-	-	1	-	-
Female	2	4	5	5	3	4	4	2	2	1	-	-
Asian or Pacific Islander 2/, both sexes	4	8	3	6	10	9	10	11	4	1	-	-
Male	3	7	1	4	4	4	4	4	1	-	-	-
Female	1	1	2	2	6	5	6	7	3	1	-	-
Mississippi												
All races, both sexes	511	604	740	780	1,020	1,203	1,555	1,485	1,087	462	124	4
Male	343	430	479	463	559	606	651	500	261	90	22	4
Female	168	174	261	317	461	597	904	985	826	372	102	-
White, both sexes	254	317	422	474	669	840	1,147	1,122	826	333	63	2
Male	173	238	283	294	391	432	496	388	193	59	9	2
Female	81	79	139	180	278	408	651	734	633	274	54	-
Black, both sexes	254	287	315	301	347	358	404	358	261	128	60	2
Male	167	192	194	166	166	171	154	110	68	31	12	2
Female	87	95	121	135	181	187	250	248	193	97	48	-
American Indian 1/, both sexes	3	-	2	5	2	1	2	2	-	1	-	-
Male	3	-	1	3	1	1	1	-	-	-	-	-
Female	-	-	1	2	1	-	1	2	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	2	4	2	3	-	-	1	-
Male	-	-	1	-	1	2	-	2	-	-	1	-
Female	-	-	-	-	1	2	2	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Missouri												
All races, both sexes	644	772	1,002	1,192	1,511	2,249	3,095	3,454	2,550	1,127	240	1
Male	444	518	659	743	875	1,180	1,438	1,304	750	228	33	1
Female	200	254	343	449	636	1,069	1,657	2,150	1,800	899	207	-
White, both sexes	527	606	857	1,039	1,311	1,985	2,835	3,224	2,386	1,058	209	1
Male	366	424	563	657	759	1,061	1,340	1,243	709	211	28	1
Female	161	182	294	382	552	924	1,495	1,981	1,677	847	181	-
Black, both sexes	110	160	141	147	186	247	247	219	154	69	31	-
Male	74	93	93	84	108	109	94	58	40	17	5	-
Female	36	67	48	63	78	138	153	161	114	52	26	-
American Indian 1/, both sexes	6	1	1	4	5	6	2	3	2	-	-	-
Male	3	-	1	1	4	3	-	-	-	-	-	-
Female	3	1	-	3	1	3	2	3	2	-	-	-
Asian or Pacific Islander 2/, both sexes	1	5	3	2	9	11	11	8	8	-	-	-
Male	1	1	2	1	4	7	4	3	1	-	-	-
Female	-	4	1	1	5	4	7	5	7	-	-	-
Montana												
All races, both sexes	85	107	161	144	200	278	404	483	392	159	19	-
Male	67	81	108	99	119	167	198	214	146	46	4	-
Female	18	26	53	45	81	111	206	269	246	113	15	-
White, both sexes	80	97	150	132	192	265	392	479	384	158	18	-
Male	62	75	102	90	113	160	194	214	143	45	4	-
Female	18	22	48	42	79	105	198	265	241	113	14	-
Black, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	10	11	11	7	13	10	3	8	1	1	-
Male	4	6	6	8	5	7	3	-	3	1	-	-
Female	-	4	5	3	2	6	7	3	5	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	2	1	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Nebraska												
All races, both sexes	129	158	188	255	332	502	783	959	879	412	96	-
Male	97	113	133	171	191	289	390	392	246	77	12	-
Female	32	45	55	84	141	213	393	567	633	335	84	-
White, both sexes	115	137	169	236	307	482	765	937	867	407	93	-
Male	84	101	122	164	177	279	385	381	245	77	11	-
Female	31	36	47	72	130	203	380	556	622	330	82	-
Black, both sexes	11	19	14	15	18	18	14	18	11	5	3	-
Male	10	10	10	5	10	9	3	9	1	-	1	-
Female	1	9	4	10	8	9	11	9	10	5	2	-
American Indian 1/, both sexes	3	1	2	3	3	1	2	1	1	-	-	-
Male	3	1	-	2	2	1	1	-	-	-	-	-
Female	-	-	2	1	1	-	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	3	1	4	1	2	3	-	-	-	-
Male	-	1	1	-	2	-	1	2	-	-	-	-
Female	-	-	2	1	2	1	1	1	-	-	-	-
Nevada												
All races, both sexes	280	387	489	569	590	765	946	796	471	136	26	1
Male	208	269	348	375	356	454	523	356	150	38	6	1
Female	72	118	141	194	234	311	423	440	321	98	20	-
White, both sexes	223	311	418	476	504	680	869	740	443	124	24	1
Male	171	219	300	329	306	412	489	338	140	33	6	1
Female	52	92	118	147	198	268	380	402	303	91	18	-
Black, both sexes	34	51	48	59	57	53	48	27	15	10	1	-
Male	21	34	32	32	33	29	23	8	6	4	-	-
Female	13	17	16	27	24	24	25	19	9	6	1	-
American Indian 1/, both sexes	5	4	3	5	5	5	9	5	3	-	-	-
Male	4	3	3	2	4	3	5	2	-	-	-	-
Female	1	1	-	3	1	2	4	3	3	-	-	-
Asian or Pacific Islander 2/, both sexes	18	21	20	29	24	27	20	24	10	2	1	-
Male	12	13	13	12	13	10	6	8	4	1	-	-
Female	6	8	7	17	11	17	14	16	6	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
New Hampshire												
All races, both sexes	78	118	182	155	256	358	540	648	509	209	52	-
Male	65	86	131	106	160	193	245	267	149	48	10	-
Female	13	32	51	49	96	165	295	381	360	161	42	-
White, both sexes	77	117	178	152	255	352	540	648	505	209	52	-
Male	65	85	129	104	160	193	245	267	149	48	10	-
Female	12	32	49	48	95	159	295	381	356	161	42	-
Black, both sexes	1	-	1	3	-	1	-	-	1	-	-	-
Male	-	-	1	2	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	1	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	2	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	1	5	-	-	3	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	5	-	-	3	-	-	-
New Jersey												
All races, both sexes	609	867	1,016	1,280	1,784	2,797	4,260	4,874	3,580	1,546	338	2
Male	425	601	685	796	1,041	1,474	2,055	1,893	1,107	348	64	-
Female	184	266	331	484	743	1,323	2,205	2,981	2,473	1,198	274	2
White, both sexes	446	643	788	962	1,429	2,408	3,796	4,504	3,347	1,438	307	1
Male	321	472	544	613	840	1,285	1,836	1,764	1,055	330	60	-
Female	125	171	244	349	589	1,123	1,960	2,740	2,292	1,108	247	1
Black, both sexes	146	200	191	274	298	336	390	311	192	89	28	1
Male	89	112	117	150	163	167	185	106	43	13	3	-
Female	57	88	74	124	135	169	205	205	149	76	25	1
American Indian 1/, both sexes	-	1	-	5	2	1	3	2	4	1	-	-
Male	-	-	-	2	2	-	1	-	-	-	-	-
Female	-	1	-	3	-	1	2	2	4	1	-	-
Asian or Pacific Islander 2/, both sexes	17	23	37	39	55	52	71	57	37	18	3	-
Male	15	17	24	31	36	22	33	23	9	5	1	-
Female	2	6	13	8	19	30	38	34	28	13	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
New Mexico												
All races, both sexes	142	223	273	279	381	538	741	799	559	230	52	-
Male	101	157	175	175	235	298	346	327	209	57	14	-
Female	41	66	98	104	146	240	395	472	350	173	38	-
White, both sexes	132	198	253	258	341	506	691	763	535	219	48	-
Male	92	142	161	160	213	276	317	310	198	53	13	-
Female	40	56	92	98	128	230	374	453	337	166	35	-
Black, both sexes	4	7	8	4	12	9	19	7	7	2	1	-
Male	4	4	6	3	7	8	9	1	2	-	-	-
Female	-	3	2	1	5	1	10	6	5	2	1	-
American Indian 1/, both sexes	5	16	12	17	28	21	26	24	15	8	3	-
Male	4	10	8	12	15	14	18	12	8	4	1	-
Female	1	6	4	5	13	7	8	12	7	4	2	-
Asian or Pacific Islander 2/, both sexes	1	2	-	-	-	2	5	5	2	1	-	-
Male	1	1	-	-	-	-	2	4	1	-	-	-
Female	-	1	-	-	-	2	3	1	1	1	-	-
New York												
All races, both sexes	1,639	2,184	2,751	3,339	4,447	6,951	10,066	11,406	9,077	4,212	970	-
Male	1,106	1,495	1,801	2,068	2,486	3,698	4,647	4,373	2,854	889	143	-
Female	533	689	950	1,271	1,961	3,253	5,419	7,033	6,223	3,323	827	-
White, both sexes	1,131	1,554	2,014	2,471	3,485	5,714	8,694	10,122	8,275	3,847	847	-
Male	804	1,120	1,359	1,577	1,986	3,107	4,100	3,954	2,648	827	125	-
Female	327	434	655	894	1,499	2,607	4,594	6,168	5,627	3,020	722	-
Black, both sexes	463	549	655	750	824	1,036	1,141	1,082	652	302	101	-
Male	271	317	382	412	412	487	445	331	155	43	11	-
Female	192	232	273	338	412	549	696	751	497	259	90	-
American Indian 1/, both sexes	5	7	7	11	7	7	7	7	3	2	-	-
Male	3	3	6	5	5	4	2	4	-	-	-	-
Female	2	4	1	6	2	3	5	3	3	2	-	-
Asian or Pacific Islander 2/, both sexes	40	74	75	107	131	194	224	195	147	61	22	-
Male	28	55	54	74	83	100	100	84	51	19	7	-
Female	12	19	21	33	48	94	124	111	96	42	15	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
North Carolina												
All races, both sexes	943	1,224	1,521	1,687	2,178	2,891	3,851	3,950	2,717	1,032	244	2
Male	633	854	1,030	1,053	1,260	1,526	1,710	1,472	764	207	25	1
Female	310	370	491	634	918	1,365	2,141	2,478	1,953	825	219	1
White, both sexes	585	784	1,069	1,192	1,646	2,298	3,176	3,295	2,341	835	194	-
Male	412	561	753	784	967	1,244	1,452	1,273	674	171	18	-
Female	173	223	316	408	679	1,054	1,724	2,022	1,667	664	176	-
Black, both sexes	340	410	423	457	505	555	640	619	362	191	48	2
Male	208	272	257	248	280	258	241	188	89	34	6	1
Female	132	138	166	209	225	297	399	431	273	157	42	1
American Indian 1/, both sexes	13	17	18	28	21	23	27	27	9	3	1	-
Male	9	12	14	17	10	14	13	8	1	1	1	-
Female	4	5	4	11	11	9	14	19	8	2	-	-
Asian or Pacific Islander 2/, both sexes	5	13	11	10	6	15	8	9	5	3	1	-
Male	4	9	6	4	3	10	4	3	-	1	-	-
Female	1	4	5	6	3	5	4	6	5	2	1	-
North Dakota												
All races, both sexes	53	49	76	106	120	193	282	412	335	166	29	-
Male	43	38	47	60	79	119	153	173	117	38	2	-
Female	10	11	29	46	41	74	129	239	218	128	27	-
White, both sexes	48	43	65	101	117	187	279	408	333	165	29	-
Male	38	34	41	60	78	116	151	171	116	37	2	-
Female	10	9	24	41	39	71	128	237	217	128	27	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	6	11	5	3	6	3	4	2	-	-	-
Male	4	4	6	-	1	3	2	2	1	-	-	-
Female	-	2	5	5	2	3	1	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Ohio												
All races, both sexes	1,196	1,596	1,951	2,125	2,887	4,213	6,095	6,699	4,851	1,784	378	-
Male	800	1,124	1,314	1,306	1,603	2,199	2,786	2,588	1,431	377	51	-
Female	396	472	637	819	1,284	2,014	3,309	4,111	3,420	1,407	327	-
White, both sexes	930	1,271	1,623	1,792	2,460	3,757	5,602	6,259	4,562	1,655	346	-
Male	650	926	1,112	1,121	1,388	1,973	2,546	2,418	1,347	350	46	-
Female	280	345	511	671	1,072	1,784	3,056	3,841	3,215	1,305	300	-
Black, both sexes	261	314	314	323	413	440	475	426	279	126	30	-
Male	145	193	193	179	210	214	230	164	80	24	5	-
Female	116	121	121	144	203	226	245	262	199	102	25	-
American Indian 1/, both sexes	2	1	4	2	-	2	2	1	2	1	1	-
Male	2	-	3	1	-	2	2	-	-	1	-	-
Female	-	1	1	1	-	-	-	1	2	-	1	-
Asian or Pacific Islander 2/, both sexes	3	10	10	8	14	14	16	13	8	2	1	-
Male	3	5	6	5	5	10	8	6	4	2	-	-
Female	-	5	4	3	9	4	8	7	4	-	1	-
Oklahoma												
All races, both sexes	471	597	774	906	1,114	1,538	2,018	2,107	1,479	611	137	-
Male	322	391	493	538	619	774	923	800	467	115	28	-
Female	149	206	281	368	495	764	1,095	1,307	1,012	496	109	-
White, both sexes	374	473	648	767	973	1,337	1,832	1,962	1,384	565	120	-
Male	263	309	412	456	556	683	841	753	439	104	23	-
Female	111	164	236	311	417	654	991	1,209	945	461	97	-
Black, both sexes	55	64	63	68	71	109	99	79	50	27	12	-
Male	32	46	41	35	33	44	40	18	18	8	4	-
Female	23	18	22	33	38	65	59	61	32	19	8	-
American Indian 1/, both sexes	40	53	55	66	62	86	80	63	44	16	5	-
Male	26	34	37	46	29	46	40	28	9	2	1	-
Female	14	19	18	20	33	40	40	35	35	14	4	-
Asian or Pacific Islander 2/, both sexes	2	7	8	5	8	6	7	3	1	3	-	-
Male	1	2	3	1	1	1	2	1	1	1	-	-
Female	1	5	5	4	7	5	5	2	-	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Oregon												
All races, both sexes	267	393	447	544	713	1,023	1,529	1,840	1,478	570	112	1
Male	197	253	301	361	425	575	784	745	525	152	21	1
Female	70	140	146	183	288	448	745	1,095	953	418	91	-
White, both sexes	250	369	425	514	679	994	1,479	1,805	1,461	555	112	1
Male	185	237	283	343	406	556	760	736	517	146	21	1
Female	65	132	142	171	273	438	719	1,069	944	409	91	-
Black, both sexes	7	15	6	12	12	10	12	8	7	4	-	-
Male	5	11	6	10	9	7	5	-	3	-	-	-
Female	2	4	-	2	3	3	7	8	4	4	-	-
American Indian 1/, both sexes	7	3	5	9	9	4	9	9	-	1	-	-
Male	4	2	5	5	3	2	5	2	-	-	-	-
Female	3	1	-	4	6	2	4	7	-	1	-	-
Asian or Pacific Islander 2/, both sexes	3	6	11	9	13	15	29	18	10	10	-	-
Male	3	3	7	3	7	10	14	7	5	6	-	-
Female	-	3	4	6	6	5	15	11	5	4	-	-
Pennsylvania												
All races, both sexes	1,137	1,620	2,088	2,283	3,095	5,200	7,646	8,834	6,357	2,430	485	3
Male	803	1,138	1,406	1,407	1,822	2,718	3,543	3,495	1,909	569	70	2
Female	334	482	682	876	1,273	2,482	4,103	5,339	4,448	1,861	415	1
White, both sexes	877	1,305	1,775	1,983	2,741	4,723	7,138	8,365	6,050	2,302	451	2
Male	633	944	1,217	1,231	1,619	2,506	3,332	3,337	1,839	541	65	1
Female	244	361	558	752	1,122	2,217	3,806	5,028	4,211	1,761	386	1
Black, both sexes	250	297	301	279	326	450	472	428	288	123	33	1
Male	162	183	183	159	188	199	199	140	62	26	5	1
Female	88	114	118	120	138	251	273	288	226	97	28	-
American Indian 1/, both sexes	-	2	1	1	1	1	2	4	1	-	1	-
Male	-	2	-	1	1	1	2	1	-	-	-	-
Female	-	-	1	-	-	-	-	3	1	-	1	-
Asian or Pacific Islander 2/, both sexes	10	16	11	20	27	26	34	37	18	5	-	-
Male	8	9	6	16	14	12	10	17	8	2	-	-
Female	2	7	5	4	13	14	24	20	10	3	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Rhode Island												
All races, both sexes	82	127	132	158	208	352	590	740	588	276	46	-
Male	61	89	92	103	124	171	253	268	169	66	1	-
Female	21	38	40	55	84	181	337	472	419	210	45	-
White, both sexes	73	116	117	149	192	337	576	725	584	272	46	-
Male	57	82	80	100	113	166	249	265	168	64	1	-
Female	16	34	37	49	79	171	327	460	416	208	45	-
Black, both sexes	5	8	13	7	13	12	12	12	3	2	-	-
Male	2	6	11	2	9	4	3	3	1	1	-	-
Female	3	2	2	5	4	8	9	9	2	1	-	-
American Indian 1/, both sexes	2	1	2	-	1	1	-	-	1	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	1	1	1	-	1	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	-	2	2	2	2	3	-	2	-	-
Male	1	1	-	1	2	1	1	-	-	1	-	-
Female	1	1	-	1	-	1	1	3	-	1	-	-
South Carolina												
All races, both sexes	578	739	952	1,053	1,180	1,527	1,896	1,845	1,212	453	84	3
Male	374	522	640	684	676	794	879	675	324	86	10	3
Female	204	217	312	369	504	733	1,017	1,170	888	367	74	-
White, both sexes	318	429	600	680	804	1,114	1,464	1,474	985	348	57	3
Male	212	301	420	454	473	607	693	572	271	72	7	3
Female	106	128	180	226	331	507	771	902	714	276	50	-
Black, both sexes	255	305	346	364	371	409	426	367	224	105	27	-
Male	157	218	217	224	198	185	183	103	52	14	3	-
Female	98	87	129	140	173	224	243	264	172	91	24	-
American Indian 1/, both sexes	2	4	3	2	1	-	3	1	1	-	-	-
Male	2	3	1	1	1	-	2	-	-	-	-	-
Female	-	1	2	1	-	-	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	3	1	3	7	4	4	3	3	2	-	-	-
Male	3	-	2	5	4	2	1	-	1	-	-	-
Female	-	1	1	2	-	2	2	3	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
South Dakota												
All races, both sexes	58	75	83	106	158	236	361	444	389	186	45	-
Male	43	55	56	72	97	141	191	194	137	52	5	-
Female	15	20	27	34	61	95	170	250	252	134	40	-
White, both sexes	50	61	73	98	144	226	347	433	382	184	44	-
Male	39	45	52	66	92	134	185	191	136	52	5	-
Female	11	16	21	32	52	92	162	242	246	132	39	-
Black, both sexes	1	1	2	-	-	-	-	-	-	-	1	-
Male	1	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	7	10	6	7	14	10	12	11	7	2	-	-
Male	3	8	3	5	5	7	6	3	1	-	-	-
Female	4	2	3	2	9	3	6	8	6	2	-	-
Asian or Pacific Islander 2/, both sexes	-	3	2	1	-	-	2	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	2	2	-	-	-	2	-	-	-	-	-
Tennessee												
All races, both sexes	833	1,060	1,203	1,441	1,723	2,363	2,986	3,139	2,062	824	159	-
Male	556	738	780	871	955	1,255	1,302	1,080	548	144	20	-
Female	277	322	423	570	768	1,108	1,684	2,059	1,514	680	139	-
White, both sexes	618	799	973	1,193	1,423	2,065	2,617	2,803	1,851	727	136	-
Male	430	570	643	731	824	1,094	1,152	968	494	131	16	-
Female	188	229	330	462	599	971	1,465	1,835	1,357	596	120	-
Black, both sexes	210	255	225	238	293	287	356	330	207	94	23	-
Male	124	166	135	133	128	152	145	109	54	12	4	-
Female	86	89	90	105	165	135	211	221	153	82	19	-
American Indian 1/, both sexes	-	-	1	1	1	2	3	-	-	1	-	-
Male	-	-	1	1	-	2	1	-	-	-	-	-
Female	-	-	-	-	1	-	2	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	5	6	4	9	6	9	10	6	4	2	-	-
Male	2	2	1	6	3	7	4	3	-	1	-	-
Female	3	4	3	3	3	2	6	3	4	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Texas												
All races, both sexes	2,177	2,807	3,414	3,634	4,563	6,446	8,327	8,694	5,888	2,467	501	-
Male	1,480	1,893	2,204	2,210	2,623	3,355	3,857	3,467	1,784	535	86	-
Female	697	914	1,210	1,424	1,940	3,091	4,470	5,227	4,104	1,932	415	-
White, both sexes	1,654	2,209	2,789	3,004	3,848	5,652	7,460	7,975	5,368	2,204	425	-
Male	1,151	1,528	1,829	1,875	2,239	2,985	3,493	3,217	1,644	477	65	-
Female	503	681	960	1,129	1,609	2,667	3,967	4,758	3,724	1,727	360	-
Black, both sexes	493	558	572	573	656	717	778	647	468	246	73	-
Male	312	340	345	301	351	335	328	223	120	52	18	-
Female	181	218	227	272	305	382	450	424	348	194	55	-
American Indian 1/, both sexes	5	6	4	15	9	9	7	11	4	-	-	-
Male	3	5	3	9	7	6	2	6	1	-	-	-
Female	2	1	1	6	2	3	5	5	3	-	-	-
Asian or Pacific Islander 2/, both sexes	25	34	49	42	50	68	82	61	48	17	3	-
Male	14	20	27	25	26	29	34	21	19	6	3	-
Female	11	14	22	17	24	39	48	40	29	11	-	-
Utah												
All races, both sexes	116	186	185	244	311	458	767	786	550	228	25	-
Male	78	129	130	156	173	232	353	338	192	71	5	-
Female	38	57	55	88	138	226	414	448	358	157	20	-
White, both sexes	107	171	172	235	301	449	754	774	540	226	24	-
Male	72	118	121	152	169	226	348	333	189	71	5	-
Female	35	53	51	83	132	223	406	441	351	155	19	-
Black, both sexes	2	2	3	3	-	2	3	3	2	1	-	-
Male	2	1	2	2	-	2	-	1	1	-	-	-
Female	-	1	1	1	-	-	3	2	1	1	-	-
American Indian 1/, both sexes	3	6	2	3	2	2	4	1	2	-	-	-
Male	3	5	2	1	-	1	2	1	-	-	-	-
Female	-	1	-	2	2	1	2	-	2	-	-	-
Asian or Pacific Islander 2/, both sexes	4	7	8	3	8	5	6	8	6	1	1	-
Male	1	5	5	1	4	3	3	3	2	-	-	-
Female	3	2	3	2	4	2	3	5	4	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Vermont												
All races, both sexes	47	55	77	99	101	183	289	303	231	123	19	-
Male	35	37	55	60	63	105	125	124	69	33	4	-
Female	12	18	22	39	38	78	164	179	162	90	15	-
White, both sexes	47	54	77	97	101	183	285	303	231	123	19	-
Male	35	37	55	59	63	105	124	124	69	33	4	-
Female	12	17	22	38	38	78	161	179	162	90	15	-
Black, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Virginia												
All races, both sexes	695	966	1,053	1,247	1,572	2,242	3,014	3,190	2,219	870	193	-
Male	474	667	705	775	927	1,184	1,372	1,216	663	197	27	-
Female	221	299	348	472	645	1,058	1,642	1,974	1,556	673	166	-
White, both sexes	452	636	712	888	1,171	1,711	2,475	2,702	1,886	732	147	-
Male	313	456	495	573	711	909	1,142	1,041	575	165	18	-
Female	139	180	217	315	460	802	1,333	1,661	1,311	567	129	-
Black, both sexes	236	314	322	335	365	496	495	452	302	129	45	-
Male	157	202	194	187	196	258	214	160	74	30	8	-
Female	79	112	128	148	169	238	281	292	228	99	37	-
American Indian 1/, both sexes	1	1	-	-	1	4	2	-	2	2	-	-
Male	-	1	-	-	1	1	1	-	1	1	-	-
Female	1	-	-	-	-	3	1	-	1	1	-	-
Asian or Pacific Islander 2/, both sexes	6	15	19	24	35	31	42	36	29	7	1	-
Male	4	8	16	15	19	16	15	15	13	1	1	-
Female	2	7	3	9	16	15	27	21	16	6	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Washington												
All races, both sexes	462	602	792	824	1,185	1,716	2,418	2,951	2,334	946	185	1
Male	332	424	537	537	713	914	1,151	1,255	789	242	36	1
Female	130	178	255	287	472	802	1,267	1,696	1,545	704	149	-
White, both sexes	399	531	715	735	1,080	1,581	2,251	2,830	2,251	927	182	1
Male	289	374	487	480	654	851	1,084	1,205	762	238	36	1
Female	110	157	228	255	426	730	1,167	1,625	1,489	689	146	-
Black, both sexes	30	28	30	37	42	56	47	29	21	5	3	-
Male	20	21	19	25	27	27	19	13	9	1	-	-
Female	10	7	11	12	15	29	28	16	12	4	3	-
American Indian 1/, both sexes	13	18	18	16	16	23	20	21	4	3	-	-
Male	8	10	14	7	6	10	9	10	2	-	-	-
Female	5	8	4	9	10	13	11	11	2	3	-	-
Asian or Pacific Islander 2/, both sexes	20	25	29	36	47	56	100	71	58	11	-	-
Male	15	19	17	25	26	26	39	27	16	3	-	-
Female	5	6	12	11	21	30	61	44	42	8	-	-
West Virginia												
All races, both sexes	240	310	395	510	595	866	1,213	1,162	833	326	64	-
Male	174	214	264	308	347	433	581	423	223	55	9	-
Female	66	96	131	202	248	433	632	739	610	271	55	-
White, both sexes	230	292	378	493	580	836	1,187	1,121	810	317	60	-
Male	165	202	250	296	339	418	571	414	218	54	8	-
Female	65	90	128	197	241	418	616	707	592	263	52	-
Black, both sexes	10	17	15	13	15	30	25	39	22	9	4	-
Male	9	11	12	9	8	15	10	9	4	1	1	-
Female	1	6	3	4	7	15	15	30	18	8	3	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	4	-	-	1	2	1	-	-	-
Male	-	-	2	3	-	-	-	-	1	-	-	-
Female	-	-	-	1	-	-	1	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Wisconsin												
All races, both sexes	404	586	705	806	1,080	1,658	2,590	2,956	2,366	987	224	-
Male	294	438	486	536	674	919	1,252	1,190	740	200	22	-
Female	110	148	219	270	406	739	1,338	1,766	1,626	787	202	-
White, both sexes	342	501	644	737	998	1,581	2,505	2,896	2,324	971	220	-
Male	252	391	450	502	629	879	1,221	1,169	728	198	22	-
Female	90	110	194	235	369	702	1,284	1,727	1,596	773	198	-
Black, both sexes	57	68	50	51	67	59	64	43	35	13	3	-
Male	37	34	29	27	36	33	21	14	8	2	-	-
Female	20	34	21	24	31	26	43	29	27	11	3	-
American Indian 1/, both sexes	3	13	5	12	7	13	10	7	2	1	1	-
Male	3	9	3	5	5	4	5	4	1	-	-	-
Female	-	4	2	7	2	9	5	3	1	1	1	-
Asian or Pacific Islander 2/, both sexes	2	4	6	6	8	5	11	10	5	2	-	-
Male	2	4	4	2	4	3	5	3	3	-	-	-
Female	-	-	2	4	4	2	6	7	2	2	-	-
Wyoming												
All races, both sexes	51	79	74	84	112	152	222	209	150	61	9	-
Male	34	52	52	58	71	87	107	80	44	16	2	-
Female	17	27	22	26	41	65	115	129	106	45	7	-
White, both sexes	51	75	70	83	110	148	219	208	148	60	9	-
Male	34	49	48	57	70	84	106	80	44	16	2	-
Female	17	26	22	26	40	64	113	128	104	44	7	-
Black, both sexes	-	-	-	-	1	3	-	-	1	1	-	-
Male	-	-	-	-	-	3	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	1	-	-
American Indian 1/, both sexes	-	4	3	1	-	1	2	1	1	-	-	-
Male	-	3	3	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	2	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	-	1	-	-	-	-	-
Male	-	-	1	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
United States												
All races, both sexes	3,201	2	1	-	1	4	8	15	13	24	46	81
Male	1,026	-	1	-	-	2	4	7	7	13	14	33
Female	2,175	2	-	-	1	2	4	8	6	11	32	48
White, both sexes	2,853	1	1	-	-	3	6	8	10	19	32	58
Male	907	-	1	-	-	2	3	3	6	9	11	22
Female	1,946	1	-	-	-	1	3	5	4	10	21	36
Black, both sexes	245	-	-	-	1	1	2	3	1	3	11	19
Male	84	-	-	-	-	-	1	2	-	2	2	9
Female	161	-	-	-	1	1	1	1	1	1	9	10
American Indian 1/, both sexes	17	1	-	-	-	-	-	1	-	1	1	1
Male	7	-	-	-	-	-	-	-	-	1	1	1
Female	10	1	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	86	-	-	-	-	-	-	3	2	1	2	3
Male	28	-	-	-	-	-	-	2	1	1	-	1
Female	58	-	-	-	-	-	-	1	1	-	2	2
Alabama												
All races, both sexes	36	-	-	-	-	1	-	-	1	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	26	-	-	-	-	1	-	-	1	-	-	1
White, both sexes	29	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	6	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Alaska												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	40	1	-	-	-	-	-	1	1	1	-	2
Male	17	-	-	-	-	-	-	1	1	1	-	-
Female	23	1	-	-	-	-	-	-	-	-	-	2
White, both sexes	40	1	-	-	-	-	-	1	1	1	-	2
Male	17	-	-	-	-	-	-	1	1	1	-	-
Female	23	1	-	-	-	-	-	-	-	-	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Arkansas												
All races, both sexes	29	-	-	-	-	-	1	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	1	-	-	-	-	1
White, both sexes	26	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	382	-	-	-	-	1	1	2	3	2	6	10
Male	151	-	-	-	-	1	-	-	2	-	2	6
Female	231	-	-	-	-	-	1	2	1	2	4	4
White, both sexes	321	-	-	-	-	1	1	1	2	2	5	8
Male	127	-	-	-	-	1	-	-	1	-	2	5
Female	194	-	-	-	-	-	1	1	1	2	3	3
Black, both sexes	25	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	36	-	-	-	-	-	-	1	1	-	1	1
Male	13	-	-	-	-	-	-	-	1	-	-	-
Female	23	-	-	-	-	-	-	1	-	-	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Colorado												
All races, both sexes	64	-	-	-	-	-	-	-	1	-	-	2
Male	24	-	-	-	-	-	-	-	1	-	-	1
Female	40	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	62	-	-	-	-	-	-	-	1	-	-	2
Male	24	-	-	-	-	-	-	-	1	-	-	1
Female	38	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	49	-	-	-	-	-	-	-	-	-	-	3
Male	17	-	-	-	-	-	-	-	-	-	-	1
Female	32	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	44	-	-	-	-	-	-	-	-	-	-	2
Male	16	-	-	-	-	-	-	-	-	-	-	1
Female	28	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Delaware												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Florida												
All races, both sexes	208	-	-	-	-	-	-	-	-	6	4	6
Male	84	-	-	-	-	-	-	-	-	4	1	2
Female	124	-	-	-	-	-	-	-	-	2	3	4
White, both sexes	190	-	-	-	-	-	-	-	-	5	2	5
Male	77	-	-	-	-	-	-	-	-	3	1	2
Female	113	-	-	-	-	-	-	-	-	2	1	3
Black, both sexes	17	-	-	-	-	-	-	-	-	1	2	1
Male	7	-	-	-	-	-	-	-	-	1	-	-
Female	10	-	-	-	-	-	-	-	-	-	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	55	-	-	-	-	1	-	-	2	1	-	2
Male	12	-	-	-	-	-	-	-	1	-	-	-
Female	43	-	-	-	-	1	-	-	1	1	-	2
White, both sexes	45	-	-	-	-	1	-	-	2	-	-	2
Male	9	-	-	-	-	-	-	-	1	-	-	-
Female	36	-	-	-	-	1	-	-	1	-	-	2
Black, both sexes	9	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Hawaii												
All races, both sexes	17	-	-	-	-	-	-	2	-	1	1	1
Male	7	-	-	-	-	-	-	2	-	1	-	1
Female	10	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	2	-	1	-	1
Male	6	-	-	-	-	-	-	2	-	1	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Illinois												
All races, both sexes	117	-	-	-	-	-	-	1	-	2	3	3
Male	36	-	-	-	-	-	-	-	-	1	1	2
Female	81	-	-	-	-	-	-	1	-	1	2	1
White, both sexes	103	-	-	-	-	-	-	1	-	1	2	2
Male	32	-	-	-	-	-	-	-	-	-	1	2
Female	71	-	-	-	-	-	-	1	-	1	1	-
Black, both sexes	12	-	-	-	-	-	-	-	-	1	-	1
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Indiana												
All races, both sexes	59	-	-	-	-	-	-	-	-	-	1	1
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	42	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	58	-	-	-	-	-	-	-	-	-	1	1
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	41	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Iowa												
All races, both sexes	60	-	-	-	-	-	-	-	-	-	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	41	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	58	-	-	-	-	-	-	-	-	-	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	39	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	36	-	-	-	-	1	-	-	-	-	-	1
Male	12	-	-	-	-	1	-	-	-	-	-	1
Female	24	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	33	-	-	-	-	1	-	-	-	-	-	1
Male	10	-	-	-	-	1	-	-	-	-	-	1
Female	23	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Kentucky												
All races, both sexes	42	-	-	-	-	-	-	-	-	2	-	-
Male	12	-	-	-	-	-	-	-	-	1	-	-
Female	30	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	42	-	-	-	-	-	-	-	-	2	-	-
Male	12	-	-	-	-	-	-	-	-	1	-	-
Female	30	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	31	-	-	-	-	-	-	-	1	1	-	1
Male	12	-	-	-	-	-	-	-	-	1	-	1
Female	19	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	24	-	-	-	-	-	-	-	1	1	-	-
Male	7	-	-	-	-	-	-	-	-	1	-	-
Female	17	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Maine												
All races, both sexes	21	-	-	-	-	-	-	-	-	-	-	2
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	15	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	20	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	53	-	-	-	1	-	-	-	1	-	-	1
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	37	-	-	-	1	-	-	-	1	-	-	1
White, both sexes	40	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	11	-	-	-	1	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	1	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Massachusetts												
All races, both sexes	73	-	-	-	-	-	-	-	-	-	-	2
Male	27	-	-	-	-	-	-	-	-	-	-	1
Female	46	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	69	-	-	-	-	-	-	-	-	-	-	1
Male	25	-	-	-	-	-	-	-	-	-	-	1
Female	44	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	109	-	-	-	-	-	-	-	-	1	1	1
Male	32	-	-	-	-	-	-	-	-	-	-	-
Female	77	-	-	-	-	-	-	-	-	1	1	1
White, both sexes	97	-	-	-	-	-	-	-	-	1	1	1
Male	28	-	-	-	-	-	-	-	-	-	-	-
Female	69	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Minnesota												
All races, both sexes	72	-	-	-	-	-	-	-	-	-	2	-
Male	21	-	-	-	-	-	-	-	-	-	2	-
Female	51	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	69	-	-	-	-	-	-	-	-	-	2	-
Male	21	-	-	-	-	-	-	-	-	-	2	-
Female	48	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	25	-	-	-	-	-	-	-	-	-	-	2
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Missouri												
All races, both sexes	78	-	-	-	-	-	-	1	-	-	1	3
Male	17	-	-	-	-	-	-	-	-	-	-	1
Female	61	-	-	-	-	-	-	1	-	-	1	2
White, both sexes	68	-	-	-	-	-	-	1	-	-	1	1
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	55	-	-	-	-	-	-	1	-	-	1	1
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Nebraska												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	-	3
Male	5	-	-	-	-	-	-	-	-	-	-	2
Female	12	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
New Hampshire												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	88	-	-	-	-	-	-	-	1	-	1	1
Male	30	-	-	-	-	-	-	-	1	-	1	1
Female	58	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	80	-	-	-	-	-	-	-	1	-	1	-
Male	26	-	-	-	-	-	-	-	1	-	1	-
Female	54	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
New Mexico												
All races, both sexes	17	1	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	14	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	165	-	-	-	-	-	1	-	-	-	6	5
Male	51	-	-	-	-	-	1	-	-	-	1	1
Female	114	-	-	-	-	-	-	-	-	-	5	4
White, both sexes	144	-	-	-	-	-	1	-	-	-	2	3
Male	42	-	-	-	-	-	1	-	-	-	-	1
Female	102	-	-	-	-	-	-	-	-	-	2	2
Black, both sexes	19	-	-	-	-	-	-	-	-	-	4	2
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	-	-	-	3	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
North Carolina												
All races, both sexes	83	-	-	-	-	-	1	1	-	1	3	1
Male	23	-	-	-	-	-	1	-	-	1	-	1
Female	60	-	-	-	-	-	-	1	-	-	3	-
White, both sexes	64	-	-	-	-	-	1	1	-	1	2	-
Male	19	-	-	-	-	-	1	-	-	1	-	-
Female	45	-	-	-	-	-	-	1	-	-	2	-
Black, both sexes	17	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	1	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	13	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Ohio												
All races, both sexes	156	-	-	-	-	-	1	1	-	1	3	1
Male	40	-	-	-	-	-	1	-	-	-	2	-
Female	116	-	-	-	-	-	-	1	-	1	1	1
White, both sexes	143	-	-	-	-	-	-	1	-	1	3	1
Male	38	-	-	-	-	-	-	-	-	-	2	-
Female	105	-	-	-	-	-	-	1	-	1	1	1
Black, both sexes	12	-	-	-	-	-	1	-	-	-	-	-
Male	2	-	-	-	-	-	1	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	33	-	-	-	-	-	-	-	-	-	1	2
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	31	-	-	-	-	-	-	-	-	-	1	2
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Oregon												
All races, both sexes	64	-	-	-	-	-	-	-	-	-	-	1
Male	22	-	-	-	-	-	-	-	-	-	-	1
Female	42	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	63	-	-	-	-	-	-	-	-	-	-	1
Male	22	-	-	-	-	-	-	-	-	-	-	1
Female	41	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	168	-	-	-	-	-	-	-	1	-	1	2
Male	42	-	-	-	-	-	-	-	-	-	-	-
Female	126	-	-	-	-	-	-	-	1	-	1	2
White, both sexes	163	-	-	-	-	-	-	-	1	-	1	2
Male	41	-	-	-	-	-	-	-	-	-	-	-
Female	122	-	-	-	-	-	-	-	1	-	1	2
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Rhode Island												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	52	-	-	-	-	-	-	-	-	-	2	2
Male	21	-	-	-	-	-	-	-	-	-	1	1
Female	31	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	42	-	-	-	-	-	-	-	-	-	2	1
Male	17	-	-	-	-	-	-	-	-	-	1	-
Female	25	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
South Dakota												
All races, both sexes	16	-	-	-	-	-	-	-	-	1	1	-
Male	7	-	-	-	-	-	-	-	-	1	1	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	1	1	-
Male	2	-	-	-	-	-	-	-	-	1	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	80	-	-	-	-	-	3	2	-	-	2	4
Male	20	-	-	-	-	-	1	1	-	-	-	1
Female	60	-	-	-	-	-	2	1	-	-	2	3
White, both sexes	65	-	-	-	-	-	3	-	-	-	1	3
Male	15	-	-	-	-	-	1	-	-	-	-	1
Female	50	-	-	-	-	-	2	-	-	-	1	2
Black, both sexes	15	-	-	-	-	-	-	2	-	-	1	1
Male	5	-	-	-	-	-	-	1	-	-	-	-
Female	10	-	-	-	-	-	-	1	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Texas												
All races, both sexes	157	-	1	-	-	-	-	2	-	2	3	8
Male	42	-	1	-	-	-	-	2	-	1	1	5
Female	115	-	-	-	-	-	-	-	-	1	2	3
White, both sexes	135	-	1	-	-	-	-	2	-	2	1	7
Male	36	-	1	-	-	-	-	2	-	1	-	4
Female	99	-	-	-	-	-	-	-	-	1	1	3
Black, both sexes	17	-	-	-	-	-	-	-	-	-	2	1
Male	4	-	-	-	-	-	-	-	-	-	1	1
Female	13	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	25	-	-	-	-	-	-	1	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Vermont												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	65	-	-	-	-	-	-	1	1	-	1	1
Male	17	-	-	-	-	-	-	1	1	-	-	-
Female	48	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	51	-	-	-	-	-	-	-	1	-	-	-
Male	14	-	-	-	-	-	-	-	1	-	-	-
Female	37	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	13	-	-	-	-	-	-	1	-	-	1	1
Male	3	-	-	-	-	-	-	1	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Washington												
All races, both sexes	97	-	-	-	-	-	-	-	-	-	-	1
Male	32	-	-	-	-	-	-	-	-	-	-	1
Female	65	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	94	-	-	-	-	-	-	-	-	-	-	1
Male	32	-	-	-	-	-	-	-	-	-	-	1
Female	62	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	21	-	-	-	-	-	-	-	-	-	1	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	21	-	-	-	-	-	-	-	-	-	1	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Wisconsin												
All races, both sexes	78	-	-	-	-	-	-	-	-	2	-	-
Male	28	-	-	-	-	-	-	-	-	1	-	-
Female	50	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	76	-	-	-	-	-	-	-	-	2	-	-
Male	28	-	-	-	-	-	-	-	-	1	-	-
Female	48	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
United States												
All races, both sexes	96	144	193	218	311	442	613	572	308	93	16	-
Male	45	56	77	76	102	153	181	159	76	20	-	-
Female	51	88	116	142	209	289	432	413	232	73	16	-
White, both sexes	67	98	160	180	278	411	579	539	299	89	15	-
Male	36	39	64	67	91	140	167	153	74	19	-	-
Female	31	59	96	113	187	271	412	386	225	70	15	-
Black, both sexes	24	31	26	25	18	23	23	22	7	4	1	-
Male	7	13	11	6	5	10	9	4	2	1	-	-
Female	17	18	15	19	13	13	14	18	5	3	1	-
American Indian 1/, both sexes	1	1	2	4	3	1	-	-	-	-	-	-
Male	1	-	1	-	2	-	-	-	-	-	-	-
Female	-	1	1	4	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	14	5	9	12	7	11	11	2	-	-	-
Male	1	4	1	3	4	3	5	2	-	-	-	-
Female	3	10	4	6	8	4	6	9	2	-	-	-
Alabama												
All races, both sexes	2	1	1	4	8	3	9	4	1	-	-	-
Male	1	-	-	2	4	-	2	-	1	-	-	-
Female	1	1	1	2	4	3	7	4	-	-	-	-
White, both sexes	-	1	1	4	8	2	8	3	1	-	-	-
Male	-	-	-	2	4	-	2	-	1	-	-	-
Female	-	1	1	2	4	2	6	3	-	-	-	-
Black, both sexes	2	-	-	-	-	1	1	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Alaska												
All races, both sexes	-	2	-	1	1	-	-	-	1	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	2	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	1	-	1	1	-	-	-	1	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1	2	2	5	1	8	7	4	3	1	-	-
Male	1	-	1	2	-	6	1	2	1	-	-	-
Female	-	2	1	3	1	2	6	2	2	1	-	-
White, both sexes	1	2	2	5	1	8	7	4	3	1	-	-
Male	1	-	1	2	-	6	1	2	1	-	-	-
Female	-	2	1	3	1	2	6	2	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Arkansas												
All races, both sexes	1	1	2	4	2	2	6	6	3	-	-	-
Male	1	-	-	1	2	1	1	1	-	-	-	-
Female	-	1	2	3	-	1	5	5	3	-	-	-
White, both sexes	1	1	2	4	2	2	6	4	3	-	-	-
Male	1	-	-	1	2	1	1	1	-	-	-	-
Female	-	1	2	3	-	1	5	3	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	13	23	27	24	43	47	65	59	44	8	4	-
Male	7	13	15	11	13	16	32	20	10	3	-	-
Female	6	10	12	13	30	31	33	39	34	5	4	-
White, both sexes	10	16	23	19	29	45	52	52	44	7	4	-
Male	6	9	14	8	10	15	26	17	10	3	-	-
Female	4	7	9	11	19	30	26	35	34	4	4	-
Black, both sexes	1	2	3	2	7	-	6	2	-	1	-	-
Male	-	2	1	1	1	-	3	2	-	-	-	-
Female	1	-	2	1	6	-	3	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	5	1	3	7	2	7	5	-	-	-	-
Male	1	2	-	2	2	1	3	1	-	-	-	-
Female	1	3	1	1	5	1	4	4	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Colorado												
All races, both sexes	3	2	2	4	8	12	10	15	5	-	-	-
Male	1	1	1	1	3	5	2	6	2	-	-	-
Female	2	1	1	3	5	7	8	9	3	-	-	-
White, both sexes	3	2	2	3	8	11	10	15	5	-	-	-
Male	1	1	1	1	3	5	2	6	2	-	-	-
Female	2	1	1	2	5	6	8	9	3	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	1	2	2	5	14	6	10	4	1	1	-
Male	-	1	-	1	1	5	2	5	1	-	-	-
Female	-	-	2	1	4	9	4	5	3	1	1	-
White, both sexes	-	1	1	1	4	13	6	10	4	1	1	-
Male	-	1	-	1	1	4	2	5	1	-	-	-
Female	-	-	1	-	3	9	4	5	3	1	1	-
Black, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Delaware												
All races, both sexes	-	2	-	1	2	1	1	4	1	1	-	-
Male	-	-	-	-	1	1	-	1	-	-	-	-
Female	-	2	-	1	1	-	1	3	1	1	-	-
White, both sexes	-	2	-	1	2	1	1	4	1	1	-	-
Male	-	-	-	-	1	1	-	1	-	-	-	-
Female	-	2	-	1	1	-	1	3	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	-	2	-	-	1	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	2	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Black, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Florida												
All races, both sexes	5	12	12	8	20	28	38	36	26	7	-	-
Male	4	6	4	4	9	12	9	14	11	4	-	-
Female	1	6	8	4	11	16	29	22	15	3	-	-
White, both sexes	4	6	12	7	18	28	36	34	26	7	-	-
Male	4	3	4	3	8	12	9	13	11	4	-	-
Female	-	3	8	4	10	16	27	21	15	3	-	-
Black, both sexes	1	6	-	1	2	-	1	2	-	-	-	-
Male	-	3	-	1	1	-	-	1	-	-	-	-
Female	1	3	-	-	1	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Georgia												
All races, both sexes	2	5	2	6	6	7	7	9	2	2	1	-
Male	1	1	1	1	1	1	3	1	-	1	-	-
Female	1	4	1	5	5	6	4	8	2	1	1	-
White, both sexes	1	1	1	6	5	6	7	8	2	2	1	-
Male	1	-	-	1	-	1	3	1	-	1	-	-
Female	-	1	1	5	5	5	4	7	2	1	1	-
Black, both sexes	1	3	1	-	1	1	-	1	-	-	-	-
Male	-	1	1	-	1	-	-	-	-	-	-	-
Female	1	2	-	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Hawaii												
All races, both sexes	1	-	-	2	1	1	4	2	1	-	-	-
Male	-	-	-	1	-	1	1	-	-	-	-	-
Female	1	-	-	1	1	-	3	2	1	-	-	-
White, both sexes	-	-	-	1	1	-	3	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	3	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	1	1	2	1	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	1	-	-	1	-	-	-	2	1	-	-	-
Idaho												
All races, both sexes	1	1	2	1	3	2	4	6	4	1	-	-
Male	-	-	-	-	-	-	2	3	-	-	-	-
Female	1	1	2	1	3	2	2	3	4	1	-	-
White, both sexes	1	-	2	1	3	2	4	6	4	1	-	-
Male	-	-	-	-	-	-	2	3	-	-	-	-
Female	1	-	2	1	3	2	2	3	4	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Illinois												
All races, both sexes	2	4	6	10	5	26	21	25	7	1	1	-
Male	2	1	1	7	2	8	6	4	1	-	-	-
Female	-	3	5	3	3	18	15	21	6	1	1	-
White, both sexes	1	3	5	9	5	23	18	24	7	1	1	-
Male	1	1	1	7	2	6	6	4	1	-	-	-
Female	-	2	4	2	3	17	12	20	6	1	1	-
Black, both sexes	1	1	1	1	-	3	3	-	-	-	-	-
Male	1	-	-	-	-	2	-	-	-	-	-	-
Female	-	1	1	1	-	1	3	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Indiana												
All races, both sexes	1	1	7	4	5	7	9	14	9	-	-	-
Male	-	-	2	2	1	2	3	5	2	-	-	-
Female	1	1	5	2	4	5	6	9	7	-	-	-
White, both sexes	1	-	7	4	5	7	9	14	9	-	-	-
Male	-	-	2	2	1	2	3	5	2	-	-	-
Female	1	-	5	2	4	5	6	9	7	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Iowa												
All races, both sexes	1	2	-	3	7	8	11	17	9	2	-	-
Male	1	-	-	1	2	4	3	4	3	1	-	-
Female	-	2	-	2	5	4	8	13	6	1	-	-
White, both sexes	1	1	-	3	6	8	11	17	9	2	-	-
Male	1	-	-	1	2	4	3	4	3	1	-	-
Female	-	1	-	2	4	4	8	13	6	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Kansas												
All races, both sexes	3	3	3	2	5	1	7	5	3	2	-	-
Male	1	-	2	-	1	1	1	3	1	-	-	-
Female	2	3	1	2	4	-	6	2	2	2	-	-
White, both sexes	3	2	2	2	5	1	7	4	3	2	-	-
Male	1	-	1	-	1	1	1	2	1	-	-	-
Female	2	2	1	2	4	-	6	2	2	2	-	-
Black, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Kentucky												
All races, both sexes	1	2	5	6	5	5	10	5	1	-	-	-
Male	-	-	3	1	1	1	3	2	-	-	-	-
Female	1	2	2	5	4	4	7	3	1	-	-	-
White, both sexes	1	2	5	6	5	5	10	5	1	-	-	-
Male	-	-	3	1	1	1	3	2	-	-	-	-
Female	1	2	2	5	4	4	7	3	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	5	1	4	2	3	6	3	3	-	-	-
Male	1	2	1	1	1	1	3	-	-	-	-	-
Female	-	3	-	3	1	2	3	3	3	-	-	-
White, both sexes	-	3	-	2	2	3	6	3	3	-	-	-
Male	-	-	-	1	1	1	3	-	-	-	-	-
Female	-	3	-	1	1	2	3	3	3	-	-	-
Black, both sexes	1	2	1	2	-	-	-	-	-	-	-	-
Male	1	2	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	2	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Maine												
All races, both sexes	2	1	1	1	1	3	1	6	2	1	-	-
Male	-	-	1	-	-	1	-	1	1	1	-	-
Female	2	1	-	1	1	2	1	5	1	-	-	-
White, both sexes	2	1	1	1	1	3	1	6	2	1	-	-
Male	-	-	1	-	-	1	-	1	1	1	-	-
Female	2	1	-	1	1	2	1	5	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	6	4	3	1	9	10	12	4	-	-	-
Male	-	3	2	2	-	3	2	3	1	-	-	-
Female	1	3	2	1	1	6	8	9	3	-	-	-
White, both sexes	-	3	2	2	1	6	10	11	4	-	-	-
Male	-	1	1	1	-	2	2	3	1	-	-	-
Female	-	2	1	1	1	4	8	8	3	-	-	-
Black, both sexes	1	1	2	1	-	3	-	1	-	-	-	-
Male	-	1	1	1	-	1	-	-	-	-	-	-
Female	1	-	1	-	-	2	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Massachusetts												
All races, both sexes	-	2	1	7	2	7	24	12	14	2	-	-
Male	-	1	-	3	-	4	9	4	5	-	-	-
Female	-	1	1	4	2	3	15	8	9	2	-	-
White, both sexes	-	2	1	4	2	7	24	12	14	2	-	-
Male	-	1	-	1	-	4	9	4	5	-	-	-
Female	-	1	1	3	2	3	15	8	9	2	-	-
Black, both sexes	-	-	-	2	-	-	-	-	-	-	-	-
Male	-	-	-	2	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	4	5	7	7	19	9	22	18	13	1	1	-
Male	2	2	-	3	9	3	7	3	3	-	-	-
Female	2	3	7	4	10	6	15	15	10	1	1	-
White, both sexes	3	4	6	6	16	8	21	16	12	1	1	-
Male	2	2	-	3	7	3	6	3	2	-	-	-
Female	1	2	6	3	9	5	15	13	10	1	1	-
Black, both sexes	1	-	1	-	2	-	1	2	1	-	-	-
Male	-	-	-	-	1	-	1	-	1	-	-	-
Female	1	-	1	-	1	-	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Minnesota												
All races, both sexes	3	1	5	5	5	8	12	14	14	3	-	-
Male	2	-	2	2	1	3	3	5	1	-	-	-
Female	1	1	3	3	4	5	9	9	13	3	-	-
White, both sexes	3	-	3	5	5	8	12	14	14	3	-	-
Male	2	-	2	2	1	3	3	5	1	-	-	-
Female	1	-	1	3	4	5	9	9	13	3	-	-
Black, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	2	2	3	2	2	4	4	2	2	-	-	-
Male	1	1	-	1	-	2	2	-	1	-	-	-
Female	1	1	3	1	2	2	2	2	1	-	-	-
White, both sexes	1	-	2	2	2	3	4	2	2	-	-	-
Male	-	-	-	1	-	1	2	-	1	-	-	-
Female	1	-	2	1	2	2	2	2	1	-	-	-
Black, both sexes	1	2	1	-	-	1	-	-	-	-	-	-
Male	1	1	-	-	-	1	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Missouri												
All races, both sexes	1	2	2	6	5	19	13	12	10	2	1	-
Male	-	-	1	2	2	5	3	1	2	-	-	-
Female	1	2	1	4	3	14	10	11	8	2	1	-
White, both sexes	-	2	2	6	4	15	13	11	9	2	1	-
Male	-	-	1	2	1	4	3	1	1	-	-	-
Female	-	2	1	4	3	11	10	10	8	2	1	-
Black, both sexes	1	-	-	-	-	4	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	1	-	-	-	-	3	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Montana												
All races, both sexes	1	1	-	1	1	1	4	5	-	-	-	-
Male	1	-	-	-	-	-	1	1	-	-	-	-
Female	-	1	-	1	1	1	3	4	-	-	-	-
White, both sexes	1	1	-	1	-	1	4	5	-	-	-	-
Male	1	-	-	-	-	-	1	1	-	-	-	-
Female	-	1	-	1	-	1	3	4	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Nebraska												
All races, both sexes	1	1	3	1	2	3	7	1	3	1	-	-
Male	1	-	2	1	2	-	2	1	1	-	-	-
Female	-	1	1	-	-	3	5	-	2	1	-	-
White, both sexes	1	-	3	1	2	3	6	1	3	1	-	-
Male	1	-	2	1	2	-	2	1	1	-	-	-
Female	-	-	1	-	-	3	4	-	2	1	-	-
Black, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	2	-	1	1	5	2	1	-	2	-	-	-
Male	1	-	1	-	1	-	-	-	-	-	-	-
Female	1	-	-	1	4	2	1	-	2	-	-	-
White, both sexes	2	-	1	-	5	2	1	-	2	-	-	-
Male	1	-	1	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	4	2	1	-	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
New Hampshire												
All races, both sexes	-	-	1	1	2	3	1	5	3	-	-	-
Male	-	-	-	-	1	1	-	-	1	-	-	-
Female	-	-	1	1	1	2	1	5	2	-	-	-
White, both sexes	-	-	1	1	2	3	1	5	3	-	-	-
Male	-	-	-	-	1	1	-	-	1	-	-	-
Female	-	-	1	1	1	2	1	5	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	3	9	3	7	18	22	9	11	3	-	-
Male	-	2	2	1	5	6	6	2	2	1	-	-
Female	-	1	7	2	2	12	16	7	9	2	-	-
White, both sexes	-	3	6	3	7	15	22	8	11	3	-	-
Male	-	2	1	1	5	4	6	2	2	1	-	-
Female	-	1	5	2	2	11	16	6	9	2	-	-
Black, both sexes	-	-	2	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	2	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	2	-	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
New Mexico												
All races, both sexes	1	-	-	2	-	3	6	2	1	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	2	-	3	5	2	1	-	-	-
White, both sexes	-	-	-	1	-	3	6	2	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	3	5	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	7	3	8	7	10	27	36	35	12	6	2	-
Male	1	1	5	2	3	10	12	8	4	2	-	-
Female	6	2	3	5	7	17	24	27	8	4	2	-
White, both sexes	4	2	5	5	9	26	33	35	12	5	2	-
Male	1	1	3	1	3	9	9	8	4	1	-	-
Female	3	1	2	4	6	17	24	27	8	4	2	-
Black, both sexes	3	1	2	1	1	1	3	-	-	1	-	-
Male	-	-	2	-	-	1	3	-	-	1	-	-
Female	3	1	-	1	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
North Carolina												
All races, both sexes	2	1	8	10	14	8	15	8	5	5	-	-
Male	-	-	2	2	5	3	5	2	1	-	-	-
Female	2	1	6	8	9	5	10	6	4	5	-	-
White, both sexes	-	-	5	7	11	8	14	6	5	3	-	-
Male	-	-	1	2	3	3	5	2	1	-	-	-
Female	-	-	4	5	8	5	9	4	4	3	-	-
Black, both sexes	2	1	3	2	2	-	1	2	-	2	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	2	1	2	2	1	-	1	2	-	2	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	1	4	1	2	4	-	1	-	-	-
Male	-	-	1	1	1	1	2	-	-	-	-	-
Female	-	-	-	3	-	1	2	-	1	-	-	-
White, both sexes	-	-	-	4	-	2	4	-	1	-	-	-
Male	-	-	-	1	-	1	2	-	-	-	-	-
Female	-	-	-	3	-	1	2	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Ohio												
All races, both sexes	3	8	8	8	11	18	38	33	18	4	-	-
Male	-	4	3	2	2	6	8	8	3	1	-	-
Female	3	4	5	6	9	12	30	25	15	3	-	-
White, both sexes	1	6	7	7	9	17	36	32	18	4	-	-
Male	-	4	3	2	2	5	8	8	3	1	-	-
Female	1	2	4	5	7	12	28	24	15	3	-	-
Black, both sexes	2	2	1	1	2	1	2	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	2	2	1	1	2	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Oklahoma												
All races, both sexes	1	2	-	5	9	3	2	5	3	-	-	-
Male	1	1	-	2	3	2	-	3	-	-	-	-
Female	-	1	-	3	6	1	2	2	3	-	-	-
White, both sexes	1	2	-	3	9	3	2	5	3	-	-	-
Male	1	1	-	1	3	2	-	3	-	-	-	-
Female	-	1	-	2	6	1	2	2	3	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Oregon												
All races, both sexes	2	1	3	3	5	5	15	18	5	6	-	-
Male	2	1	1	1	2	1	5	3	4	1	-	-
Female	-	-	2	2	3	4	10	15	1	5	-	-
White, both sexes	2	1	3	2	5	5	15	18	5	6	-	-
Male	2	1	1	1	2	1	5	3	4	1	-	-
Female	-	-	2	1	3	4	10	15	1	5	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	5	7	12	7	12	20	35	43	14	8	1	-
Male	3	1	6	-	3	4	11	11	3	-	-	-
Female	2	6	6	7	9	16	24	32	11	8	1	-
White, both sexes	5	6	12	7	12	20	33	42	13	8	1	-
Male	3	1	6	-	3	4	10	11	3	-	-	-
Female	2	5	6	7	9	16	23	31	10	8	1	-
Black, both sexes	-	1	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Rhode Island												
All races, both sexes	2	1	-	1	-	3	7	4	-	-	-	-
Male	1	-	-	-	-	-	3	2	-	-	-	-
Female	1	1	-	1	-	3	4	2	-	-	-	-
White, both sexes	1	1	-	1	-	3	7	4	-	-	-	-
Male	1	-	-	-	-	-	3	2	-	-	-	-
Female	-	1	-	1	-	3	4	2	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	2	1	7	3	8	9	3	9	4	1	1	-
Male	-	1	5	2	2	4	2	2	1	-	-	-
Female	2	-	2	1	6	5	1	7	3	1	1	-
White, both sexes	1	1	6	2	8	7	2	6	4	1	1	-
Male	-	1	4	2	2	3	1	2	1	-	-	-
Female	1	-	2	-	6	4	1	4	3	1	1	-
Black, both sexes	1	-	1	1	-	2	1	3	-	-	-	-
Male	-	-	1	-	-	1	1	-	-	-	-	-
Female	1	-	-	1	-	1	-	3	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
South Dakota												
All races, both sexes	-	-	2	1	1	-	5	3	1	1	-	-
Male	-	-	1	1	-	-	1	2	-	-	-	-
Female	-	-	1	-	1	-	4	1	1	1	-	-
White, both sexes	-	-	2	1	1	-	5	3	1	1	-	-
Male	-	-	1	1	-	-	1	2	-	-	-	-
Female	-	-	1	-	1	-	4	1	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	3	3	2	5	9	16	9	14	5	3	-	-
Male	1	2	1	1	2	6	-	3	-	1	-	-
Female	2	1	1	4	7	10	9	11	5	2	-	-
White, both sexes	2	2	1	5	9	15	8	11	2	3	-	-
Male	-	1	-	1	2	6	-	2	-	1	-	-
Female	2	1	1	4	7	9	8	9	2	2	-	-
Black, both sexes	1	1	1	-	-	1	1	3	3	-	-	-
Male	1	1	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	1	2	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Texas												
All races, both sexes	5	11	11	11	18	17	25	29	8	4	2	-
Male	1	3	2	4	4	3	6	8	1	-	-	-
Female	4	8	9	7	14	14	19	21	7	4	2	-
White, both sexes	5	6	9	7	16	16	23	28	7	4	1	-
Male	1	1	1	4	4	3	5	8	1	-	-	-
Female	4	5	8	3	12	13	18	20	6	4	1	-
Black, both sexes	-	3	1	4	1	1	1	1	1	-	1	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	2	-	4	1	1	1	1	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	1	-	1	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	1	1	-	1	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	1	2	4	5	3	3	5	1	-	-	-
Male	-	1	1	1	1	2	-	4	-	-	-	-
Female	-	-	1	3	4	1	3	1	1	-	-	-
White, both sexes	-	1	2	4	5	3	3	5	1	-	-	-
Male	-	1	1	1	1	2	-	4	-	-	-	-
Female	-	-	1	3	4	1	3	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Vermont												
All races, both sexes	-	-	1	-	-	5	5	3	1	-	-	-
Male	-	-	1	-	-	2	1	-	-	-	-	-
Female	-	-	-	-	-	3	4	3	1	-	-	-
White, both sexes	-	-	1	-	-	5	5	3	1	-	-	-
Male	-	-	1	-	-	2	1	-	-	-	-	-
Female	-	-	-	-	-	3	4	3	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	3	3	6	3	7	6	18	9	1	4	1	-
Male	2	2	2	-	1	1	3	3	-	1	-	-
Female	1	1	4	3	6	5	15	6	1	3	1	-
White, both sexes	1	1	3	2	7	4	18	9	-	4	1	-
Male	1	1	2	-	1	1	3	3	-	1	-	-
Female	-	-	1	2	6	3	15	6	-	3	1	-
Black, both sexes	2	1	3	1	-	2	-	-	1	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	3	1	-	2	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Washington												
All races, both sexes	1	3	5	7	12	13	21	16	10	8	-	-
Male	-	1	2	3	3	4	6	6	4	2	-	-
Female	1	2	3	4	9	9	15	10	6	6	-	-
White, both sexes	-	3	5	5	12	13	21	16	10	8	-	-
Male	-	1	2	3	3	4	6	6	4	2	-	-
Female	-	2	3	2	9	9	15	10	6	6	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	4	1	1	3	7	1	1	2	-	-	-
Male	-	3	1	-	2	5	-	-	-	-	-	-
Female	-	1	-	1	1	2	1	1	2	-	-	-
White, both sexes	-	4	1	1	3	7	1	1	2	-	-	-
Male	-	3	1	-	2	5	-	-	-	-	-	-
Female	-	1	-	1	1	2	1	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Wisconsin												
All races, both sexes	2	2	3	5	5	15	18	13	9	4	-	-
Male	2	1	1	2	4	6	6	1	3	1	-	-
Female	-	1	2	3	1	9	12	12	6	3	-	-
White, both sexes	2	2	3	4	5	14	18	13	9	4	-	-
Male	2	1	1	2	4	6	6	1	3	1	-	-
Female	-	1	2	2	1	8	12	12	6	3	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	-	2	-	-	1	4	1	2	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	2	-	-	1	4	1	1	-	-	-
White, both sexes	1	-	2	-	-	1	4	1	2	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	2	-	-	1	4	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
United States												
All races, both sexes	32	-	-	-	-	1	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	26	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	30	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Florida												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Kentucky												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Massachusetts												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
New Hampshire												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
New Mexico												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
North Carolina												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Ohio												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Oregon												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Texas												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
United States												
All races, both sexes	1	2	1	4	-	6	9	4	3	1	-	-
Male	1	-	-	2	-	2	1	-	-	-	-	-
Female	-	2	1	2	-	4	8	4	3	1	-	-
White, both sexes	-	2	1	4	-	6	9	4	3	1	-	-
Male	-	-	-	2	-	2	1	-	-	-	-	-
Female	-	2	1	2	-	4	8	4	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Florida												
All races, both sexes	-	1	-	-	-	-	2	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	2	1	-	-	-	-
White, both sexes	-	1	-	-	-	-	2	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	2	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Kentucky												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	1	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	2	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
New Mexico												
All races, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
North Carolina												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Ohio												
All races, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Oregon												
All races, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Texas												
All races, both sexes	-	-	-	1	-	2	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	2	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
United States												
All races, both sexes	3,169	2	1	-	1	3	8	15	13	24	46	81
Male	1,020	-	1	-	-	2	4	7	7	13	14	33
Female	2,149	2	-	-	1	1	4	8	6	11	32	48
White, both sexes	2,823	1	1	-	-	3	6	8	10	19	32	58
Male	902	-	1	-	-	2	3	3	6	9	11	22
Female	1,921	1	-	-	-	1	3	5	4	10	21	36
Black, both sexes	244	-	-	-	1	-	2	3	1	3	11	19
Male	84	-	-	-	-	-	1	2	-	2	2	9
Female	160	-	-	-	1	-	1	1	1	1	9	10
American Indian 1/, both sexes	16	1	-	-	-	-	-	1	-	1	1	1
Male	6	-	-	-	-	-	-	-	-	1	1	1
Female	10	1	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	86	-	-	-	-	-	-	3	2	1	2	3
Male	28	-	-	-	-	-	-	2	1	1	-	1
Female	58	-	-	-	-	-	-	1	1	-	2	2
Alabama												
All races, both sexes	35	-	-	-	-	-	-	-	1	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	29	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Alaska												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	40	1	-	-	-	-	-	1	1	1	-	2
Male	17	-	-	-	-	-	-	1	1	1	-	-
Female	23	1	-	-	-	-	-	-	-	-	-	2
White, both sexes	40	1	-	-	-	-	-	1	1	1	-	2
Male	17	-	-	-	-	-	-	1	1	1	-	-
Female	23	1	-	-	-	-	-	-	-	-	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Arkansas												
All races, both sexes	29	-	-	-	-	-	1	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	1	-	-	-	-	1
White, both sexes	26	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	381	-	-	-	-	1	1	2	3	2	6	10
Male	151	-	-	-	-	1	-	-	2	-	2	6
Female	230	-	-	-	-	-	1	2	1	2	4	4
White, both sexes	320	-	-	-	-	1	1	1	2	2	5	8
Male	127	-	-	-	-	1	-	-	1	-	2	5
Female	193	-	-	-	-	-	1	1	1	2	3	3
Black, both sexes	25	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	36	-	-	-	-	-	-	1	1	-	1	1
Male	13	-	-	-	-	-	-	-	1	-	-	-
Female	23	-	-	-	-	-	-	1	-	-	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Colorado												
All races, both sexes	64	-	-	-	-	-	-	-	1	-	-	2
Male	24	-	-	-	-	-	-	-	1	-	-	1
Female	40	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	62	-	-	-	-	-	-	-	1	-	-	2
Male	24	-	-	-	-	-	-	-	1	-	-	1
Female	38	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	49	-	-	-	-	-	-	-	-	-	-	3
Male	17	-	-	-	-	-	-	-	-	-	-	1
Female	32	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	44	-	-	-	-	-	-	-	-	-	-	2
Male	16	-	-	-	-	-	-	-	-	-	-	1
Female	28	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Delaware												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Florida												
All races, both sexes	204	-	-	-	-	-	-	-	-	6	4	6
Male	84	-	-	-	-	-	-	-	-	4	1	2
Female	120	-	-	-	-	-	-	-	-	2	3	4
White, both sexes	186	-	-	-	-	-	-	-	-	5	2	5
Male	77	-	-	-	-	-	-	-	-	3	1	2
Female	109	-	-	-	-	-	-	-	-	2	1	3
Black, both sexes	17	-	-	-	-	-	-	-	-	1	2	1
Male	7	-	-	-	-	-	-	-	-	1	-	-
Female	10	-	-	-	-	-	-	-	-	-	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	54	-	-	-	-	1	-	-	2	1	-	2
Male	12	-	-	-	-	-	-	-	1	-	-	-
Female	42	-	-	-	-	1	-	-	1	1	-	2
White, both sexes	44	-	-	-	-	1	-	-	2	-	-	2
Male	9	-	-	-	-	-	-	-	1	-	-	-
Female	35	-	-	-	-	1	-	-	1	-	-	2
Black, both sexes	9	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Hawaii												
All races, both sexes	17	-	-	-	-	-	-	2	-	1	1	1
Male	7	-	-	-	-	-	-	2	-	1	-	1
Female	10	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	2	-	1	-	1
Male	6	-	-	-	-	-	-	2	-	1	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Illinois												
All races, both sexes	117	-	-	-	-	-	-	1	-	2	3	3
Male	36	-	-	-	-	-	-	-	-	1	1	2
Female	81	-	-	-	-	-	-	1	-	1	2	1
White, both sexes	103	-	-	-	-	-	-	1	-	1	2	2
Male	32	-	-	-	-	-	-	-	-	-	1	2
Female	71	-	-	-	-	-	-	1	-	1	1	-
Black, both sexes	12	-	-	-	-	-	-	-	-	1	-	1
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Indiana												
All races, both sexes	57	-	-	-	-	-	-	-	-	-	1	1
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	41	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	56	-	-	-	-	-	-	-	-	-	1	1
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	40	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Iowa												
All races, both sexes	60	-	-	-	-	-	-	-	-	-	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	41	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	58	-	-	-	-	-	-	-	-	-	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	39	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	36	-	-	-	-	1	-	-	-	-	-	1
Male	12	-	-	-	-	1	-	-	-	-	-	1
Female	24	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	33	-	-	-	-	1	-	-	-	-	-	1
Male	10	-	-	-	-	1	-	-	-	-	-	1
Female	23	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Kentucky												
All races, both sexes	41	-	-	-	-	-	-	-	-	2	-	-
Male	12	-	-	-	-	-	-	-	-	1	-	-
Female	29	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	41	-	-	-	-	-	-	-	-	2	-	-
Male	12	-	-	-	-	-	-	-	-	1	-	-
Female	29	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	31	-	-	-	-	-	-	-	1	1	-	1
Male	12	-	-	-	-	-	-	-	-	1	-	1
Female	19	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	24	-	-	-	-	-	-	-	1	1	-	-
Male	7	-	-	-	-	-	-	-	-	1	-	-
Female	17	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Maine												
All races, both sexes	21	-	-	-	-	-	-	-	-	-	-	2
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	15	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	20	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	53	-	-	-	1	-	-	-	1	-	-	1
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	37	-	-	-	1	-	-	-	1	-	-	1
White, both sexes	40	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	11	-	-	-	1	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	1	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Massachusetts												
All races, both sexes	70	-	-	-	-	-	-	-	-	-	-	2
Male	27	-	-	-	-	-	-	-	-	-	-	1
Female	43	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	66	-	-	-	-	-	-	-	-	-	-	1
Male	25	-	-	-	-	-	-	-	-	-	-	1
Female	41	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	108	-	-	-	-	-	-	-	-	1	1	1
Male	31	-	-	-	-	-	-	-	-	-	-	-
Female	77	-	-	-	-	-	-	-	-	1	1	1
White, both sexes	96	-	-	-	-	-	-	-	-	1	1	1
Male	27	-	-	-	-	-	-	-	-	-	-	-
Female	69	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Minnesota												
All races, both sexes	72	-	-	-	-	-	-	-	-	-	2	-
Male	21	-	-	-	-	-	-	-	-	-	2	-
Female	51	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	69	-	-	-	-	-	-	-	-	-	2	-
Male	21	-	-	-	-	-	-	-	-	-	2	-
Female	48	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	2
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Missouri												
All races, both sexes	78	-	-	-	-	-	-	1	-	-	1	3
Male	17	-	-	-	-	-	-	-	-	-	-	1
Female	61	-	-	-	-	-	-	1	-	-	1	2
White, both sexes	68	-	-	-	-	-	-	1	-	-	1	1
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	55	-	-	-	-	-	-	1	-	-	1	1
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Nebraska												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	3
Male	5	-	-	-	-	-	-	-	-	-	-	2
Female	11	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
New Hampshire												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	88	-	-	-	-	-	-	-	1	-	1	1
Male	30	-	-	-	-	-	-	-	1	-	1	1
Female	58	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	80	-	-	-	-	-	-	-	1	-	1	-
Male	26	-	-	-	-	-	-	-	1	-	1	-
Female	54	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
New Mexico												
All races, both sexes	15	1	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	13	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	164	-	-	-	-	-	1	-	-	-	6	5
Male	51	-	-	-	-	-	1	-	-	-	1	1
Female	113	-	-	-	-	-	-	-	-	-	5	4
White, both sexes	143	-	-	-	-	-	1	-	-	-	2	3
Male	42	-	-	-	-	-	1	-	-	-	-	1
Female	101	-	-	-	-	-	-	-	-	-	2	2
Black, both sexes	19	-	-	-	-	-	-	-	-	-	4	2
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	-	-	-	3	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
North Carolina												
All races, both sexes	82	-	-	-	-	-	1	1	-	1	3	1
Male	23	-	-	-	-	-	1	-	-	1	-	1
Female	59	-	-	-	-	-	-	1	-	-	3	-
White, both sexes	63	-	-	-	-	-	1	1	-	1	2	-
Male	19	-	-	-	-	-	1	-	-	1	-	-
Female	44	-	-	-	-	-	-	1	-	-	2	-
Black, both sexes	17	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	1	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	13	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Ohio												
All races, both sexes	154	-	-	-	-	-	1	1	-	1	3	1
Male	40	-	-	-	-	-	1	-	-	-	2	-
Female	114	-	-	-	-	-	-	1	-	1	1	1
White, both sexes	141	-	-	-	-	-	-	1	-	1	3	1
Male	38	-	-	-	-	-	-	-	-	-	2	-
Female	103	-	-	-	-	-	-	1	-	1	1	1
Black, both sexes	12	-	-	-	-	-	1	-	-	-	-	-
Male	2	-	-	-	-	-	1	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	33	-	-	-	-	-	-	-	-	-	1	2
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	31	-	-	-	-	-	-	-	-	-	1	2
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Oregon												
All races, both sexes	62	-	-	-	-	-	-	-	-	-	-	1
Male	22	-	-	-	-	-	-	-	-	-	-	1
Female	40	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	61	-	-	-	-	-	-	-	-	-	-	1
Male	22	-	-	-	-	-	-	-	-	-	-	1
Female	39	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	167	-	-	-	-	-	-	-	1	-	1	2
Male	42	-	-	-	-	-	-	-	-	-	-	-
Female	125	-	-	-	-	-	-	-	1	-	1	2
White, both sexes	162	-	-	-	-	-	-	-	1	-	1	2
Male	41	-	-	-	-	-	-	-	-	-	-	-
Female	121	-	-	-	-	-	-	-	1	-	1	2
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Rhode Island												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	51	-	-	-	-	-	-	-	-	-	2	2
Male	20	-	-	-	-	-	-	-	-	-	1	1
Female	31	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	41	-	-	-	-	-	-	-	-	-	2	1
Male	16	-	-	-	-	-	-	-	-	-	1	-
Female	25	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
South Dakota												
All races, both sexes	16	-	-	-	-	-	-	-	-	1	1	-
Male	7	-	-	-	-	-	-	-	-	1	1	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	1	1	-
Male	2	-	-	-	-	-	-	-	-	1	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	80	-	-	-	-	-	3	2	-	-	2	4
Male	20	-	-	-	-	-	1	1	-	-	-	1
Female	60	-	-	-	-	-	2	1	-	-	2	3
White, both sexes	65	-	-	-	-	-	3	-	-	-	1	3
Male	15	-	-	-	-	-	1	-	-	-	-	1
Female	50	-	-	-	-	-	2	-	-	-	1	2
Black, both sexes	15	-	-	-	-	-	-	2	-	-	1	1
Male	5	-	-	-	-	-	-	1	-	-	-	-
Female	10	-	-	-	-	-	-	1	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Texas												
All races, both sexes	154	-	1	-	-	-	-	2	-	2	3	8
Male	40	-	1	-	-	-	-	2	-	1	1	5
Female	114	-	-	-	-	-	-	-	-	1	2	3
White, both sexes	132	-	1	-	-	-	-	2	-	2	1	7
Male	34	-	1	-	-	-	-	2	-	1	-	4
Female	98	-	-	-	-	-	-	-	-	1	1	3
Black, both sexes	17	-	-	-	-	-	-	-	-	-	2	1
Male	4	-	-	-	-	-	-	-	-	-	1	1
Female	13	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	25	-	-	-	-	-	-	1	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Vermont												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	64	-	-	-	-	-	-	1	1	-	1	1
Male	17	-	-	-	-	-	-	1	1	-	-	-
Female	47	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	50	-	-	-	-	-	-	-	1	-	-	-
Male	14	-	-	-	-	-	-	-	1	-	-	-
Female	36	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	13	-	-	-	-	-	-	1	-	-	1	1
Male	3	-	-	-	-	-	-	1	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Washington												
All races, both sexes	97	-	-	-	-	-	-	-	-	-	-	1
Male	32	-	-	-	-	-	-	-	-	-	-	1
Female	65	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	94	-	-	-	-	-	-	-	-	-	-	1
Male	32	-	-	-	-	-	-	-	-	-	-	1
Female	62	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	21	-	-	-	-	-	-	-	-	-	1	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	21	-	-	-	-	-	-	-	-	-	1	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Wisconsin												
All races, both sexes	78	-	-	-	-	-	-	-	-	2	-	-
Male	28	-	-	-	-	-	-	-	-	1	-	-
Female	50	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	76	-	-	-	-	-	-	-	-	2	-	-
Male	28	-	-	-	-	-	-	-	-	1	-	-
Female	48	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
United States												
All races, both sexes	95	142	192	214	311	436	604	568	305	92	16	-
Male	44	56	77	74	102	151	180	159	76	20	-	-
Female	51	86	115	140	209	285	424	409	229	72	16	-
White, both sexes	67	96	159	176	278	405	570	535	296	88	15	-
Male	36	39	64	65	91	138	166	153	74	19	-	-
Female	31	57	95	111	187	267	404	382	222	69	15	-
Black, both sexes	24	31	26	25	18	23	23	22	7	4	1	-
Male	7	13	11	6	5	10	9	4	2	1	-	-
Female	17	18	15	19	13	13	14	18	5	3	1	-
American Indian 1/, both sexes	-	1	2	4	3	1	-	-	-	-	-	-
Male	-	-	1	-	2	-	-	-	-	-	-	-
Female	-	1	1	4	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	14	5	9	12	7	11	11	2	-	-	-
Male	1	4	1	3	4	3	5	2	-	-	-	-
Female	3	10	4	6	8	4	6	9	2	-	-	-
Alabama												
All races, both sexes	2	1	1	4	8	3	9	4	1	-	-	-
Male	1	-	-	2	4	-	2	-	1	-	-	-
Female	1	1	1	2	4	3	7	4	-	-	-	-
White, both sexes	-	1	1	4	8	2	8	3	1	-	-	-
Male	-	-	-	2	4	-	2	-	1	-	-	-
Female	-	1	1	2	4	2	6	3	-	-	-	-
Black, both sexes	2	-	-	-	-	1	1	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Alaska												
All races, both sexes	-	2	-	1	1	-	-	-	1	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	2	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	1	-	1	1	-	-	-	1	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1	2	2	5	1	8	7	4	3	1	-	-
Male	1	-	1	2	-	6	1	2	1	-	-	-
Female	-	2	1	3	1	2	6	2	2	1	-	-
White, both sexes	1	2	2	5	1	8	7	4	3	1	-	-
Male	1	-	1	2	-	6	1	2	1	-	-	-
Female	-	2	1	3	1	2	6	2	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Arkansas												
All races, both sexes	1	1	2	4	2	2	6	6	3	-	-	-
Male	1	-	-	1	2	1	1	1	-	-	-	-
Female	-	1	2	3	-	1	5	5	3	-	-	-
White, both sexes	1	1	2	4	2	2	6	4	3	-	-	-
Male	1	-	-	1	2	1	1	1	-	-	-	-
Female	-	1	2	3	-	1	5	3	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	13	23	27	24	43	47	64	59	44	8	4	-
Male	7	13	15	11	13	16	32	20	10	3	-	-
Female	6	10	12	13	30	31	32	39	34	5	4	-
White, both sexes	10	16	23	19	29	45	51	52	44	7	4	-
Male	6	9	14	8	10	15	26	17	10	3	-	-
Female	4	7	9	11	19	30	25	35	34	4	4	-
Black, both sexes	1	2	3	2	7	-	6	2	-	1	-	-
Male	-	2	1	1	1	-	3	2	-	-	-	-
Female	1	-	2	1	6	-	3	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	5	1	3	7	2	7	5	-	-	-	-
Male	1	2	-	2	2	1	3	1	-	-	-	-
Female	1	3	1	1	5	1	4	4	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Colorado												
All races, both sexes	3	2	2	4	8	12	10	15	5	-	-	-
Male	1	1	1	1	3	5	2	6	2	-	-	-
Female	2	1	1	3	5	7	8	9	3	-	-	-
White, both sexes	3	2	2	3	8	11	10	15	5	-	-	-
Male	1	1	1	1	3	5	2	6	2	-	-	-
Female	2	1	1	2	5	6	8	9	3	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	1	2	2	5	14	6	10	4	1	1	-
Male	-	1	-	1	1	5	2	5	1	-	-	-
Female	-	-	2	1	4	9	4	5	3	1	1	-
White, both sexes	-	1	1	1	4	13	6	10	4	1	1	-
Male	-	1	-	1	1	4	2	5	1	-	-	-
Female	-	-	1	-	3	9	4	5	3	1	1	-
Black, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Delaware												
All races, both sexes	-	2	-	1	2	1	1	4	1	1	-	-
Male	-	-	-	-	1	1	-	1	-	-	-	-
Female	-	2	-	1	1	-	1	3	1	1	-	-
White, both sexes	-	2	-	1	2	1	1	4	1	1	-	-
Male	-	-	-	-	1	1	-	1	-	-	-	-
Female	-	2	-	1	1	-	1	3	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	-	2	-	-	1	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	2	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Black, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Florida												
All races, both sexes	5	11	12	8	20	28	36	35	26	7	-	-
Male	4	6	4	4	9	12	9	14	11	4	-	-
Female	1	5	8	4	11	16	27	21	15	3	-	-
White, both sexes	4	5	12	7	18	28	34	33	26	7	-	-
Male	4	3	4	3	8	12	9	13	11	4	-	-
Female	-	2	8	4	10	16	25	20	15	3	-	-
Black, both sexes	1	6	-	1	2	-	1	2	-	-	-	-
Male	-	3	-	1	1	-	-	1	-	-	-	-
Female	1	3	-	-	1	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Georgia												
All races, both sexes	2	5	2	6	6	7	6	9	2	2	1	-
Male	1	1	1	1	1	1	3	1	-	1	-	-
Female	1	4	1	5	5	6	3	8	2	1	1	-
White, both sexes	1	1	1	6	5	6	6	8	2	2	1	-
Male	1	-	-	1	-	1	3	1	-	1	-	-
Female	-	1	1	5	5	5	3	7	2	1	1	-
Black, both sexes	1	3	1	-	1	1	-	1	-	-	-	-
Male	-	1	1	-	1	-	-	-	-	-	-	-
Female	1	2	-	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Hawaii												
All races, both sexes	1	-	-	2	1	1	4	2	1	-	-	-
Male	-	-	-	1	-	1	1	-	-	-	-	-
Female	1	-	-	1	1	-	3	2	1	-	-	-
White, both sexes	-	-	-	1	1	-	3	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	3	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	1	1	2	1	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	1	-	-	1	-	-	-	2	1	-	-	-
Idaho												
All races, both sexes	1	1	2	1	3	2	4	6	4	-	-	-
Male	-	-	-	-	-	-	2	3	-	-	-	-
Female	1	1	2	1	3	2	2	3	4	-	-	-
White, both sexes	1	-	2	1	3	2	4	6	4	-	-	-
Male	-	-	-	-	-	-	2	3	-	-	-	-
Female	1	-	2	1	3	2	2	3	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Illinois												
All races, both sexes	2	4	6	10	5	26	21	25	7	1	1	-
Male	2	1	1	7	2	8	6	4	1	-	-	-
Female	-	3	5	3	3	18	15	21	6	1	1	-
White, both sexes	1	3	5	9	5	23	18	24	7	1	1	-
Male	1	1	1	7	2	6	6	4	1	-	-	-
Female	-	2	4	2	3	17	12	20	6	1	1	-
Black, both sexes	1	1	1	1	-	3	3	-	-	-	-	-
Male	1	-	-	-	-	2	-	-	-	-	-	-
Female	-	1	1	1	-	1	3	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Indiana												
All races, both sexes	1	1	7	4	5	6	9	13	9	-	-	-
Male	-	-	2	2	1	1	3	5	2	-	-	-
Female	1	1	5	2	4	5	6	8	7	-	-	-
White, both sexes	1	-	7	4	5	6	9	13	9	-	-	-
Male	-	-	2	2	1	1	3	5	2	-	-	-
Female	1	-	5	2	4	5	6	8	7	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Iowa												
All races, both sexes	1	2	-	3	7	8	11	17	9	2	-	-
Male	1	-	-	1	2	4	3	4	3	1	-	-
Female	-	2	-	2	5	4	8	13	6	1	-	-
White, both sexes	1	1	-	3	6	8	11	17	9	2	-	-
Male	1	-	-	1	2	4	3	4	3	1	-	-
Female	-	1	-	2	4	4	8	13	6	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Kansas												
All races, both sexes	3	3	3	2	5	1	7	5	3	2	-	-
Male	1	-	2	-	1	1	1	3	1	-	-	-
Female	2	3	1	2	4	-	6	2	2	2	-	-
White, both sexes	3	2	2	2	5	1	7	4	3	2	-	-
Male	1	-	1	-	1	1	1	2	1	-	-	-
Female	2	2	1	2	4	-	6	2	2	2	-	-
Black, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Kentucky												
All races, both sexes	1	2	5	5	5	5	10	5	1	-	-	-
Male	-	-	3	1	1	1	3	2	-	-	-	-
Female	1	2	2	4	4	4	7	3	1	-	-	-
White, both sexes	1	2	5	5	5	5	10	5	1	-	-	-
Male	-	-	3	1	1	1	3	2	-	-	-	-
Female	1	2	2	4	4	4	7	3	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	5	1	4	2	3	6	3	3	-	-	-
Male	1	2	1	1	1	1	3	-	-	-	-	-
Female	-	3	-	3	1	2	3	3	3	-	-	-
White, both sexes	-	3	-	2	2	3	6	3	3	-	-	-
Male	-	-	-	1	1	1	3	-	-	-	-	-
Female	-	3	-	1	1	2	3	3	3	-	-	-
Black, both sexes	1	2	1	2	-	-	-	-	-	-	-	-
Male	1	2	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	2	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Maine												
All races, both sexes	2	1	1	1	1	3	1	6	2	1	-	-
Male	-	-	1	-	-	1	-	1	1	1	-	-
Female	2	1	-	1	1	2	1	5	1	-	-	-
White, both sexes	2	1	1	1	1	3	1	6	2	1	-	-
Male	-	-	1	-	-	1	-	1	1	1	-	-
Female	2	1	-	1	1	2	1	5	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	6	4	3	1	9	10	12	4	-	-	-
Male	-	3	2	2	-	3	2	3	1	-	-	-
Female	1	3	2	1	1	6	8	9	3	-	-	-
White, both sexes	-	3	2	2	1	6	10	11	4	-	-	-
Male	-	1	1	1	-	2	2	3	1	-	-	-
Female	-	2	1	1	1	4	8	8	3	-	-	-
Black, both sexes	1	1	2	1	-	3	-	1	-	-	-	-
Male	-	1	1	1	-	1	-	-	-	-	-	-
Female	1	-	1	-	-	2	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Massachusetts												
All races, both sexes	-	2	1	7	2	7	24	11	12	2	-	-
Male	-	1	-	3	-	4	9	4	5	-	-	-
Female	-	1	1	4	2	3	15	7	7	2	-	-
White, both sexes	-	2	1	4	2	7	24	11	12	2	-	-
Male	-	1	-	1	-	4	9	4	5	-	-	-
Female	-	1	1	3	2	3	15	7	7	2	-	-
Black, both sexes	-	-	-	2	-	-	-	-	-	-	-	-
Male	-	-	-	2	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	4	5	7	6	19	9	22	18	13	1	1	-
Male	2	2	-	2	9	3	7	3	3	-	-	-
Female	2	3	7	4	10	6	15	15	10	1	1	-
White, both sexes	3	4	6	5	16	8	21	16	12	1	1	-
Male	2	2	-	2	7	3	6	3	2	-	-	-
Female	1	2	6	3	9	5	15	13	10	1	1	-
Black, both sexes	1	-	1	-	2	-	1	2	1	-	-	-
Male	-	-	-	-	1	-	1	-	1	-	-	-
Female	1	-	1	-	1	-	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Minnesota												
All races, both sexes	3	1	5	5	5	8	12	14	14	3	-	-
Male	2	-	2	2	1	3	3	5	1	-	-	-
Female	1	1	3	3	4	5	9	9	13	3	-	-
White, both sexes	3	-	3	5	5	8	12	14	14	3	-	-
Male	2	-	2	2	1	3	3	5	1	-	-	-
Female	1	-	1	3	4	5	9	9	13	3	-	-
Black, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	2	2	3	2	2	3	4	2	2	-	-	-
Male	1	1	-	1	-	2	2	-	1	-	-	-
Female	1	1	3	1	2	1	2	2	1	-	-	-
White, both sexes	1	-	2	2	2	2	4	2	2	-	-	-
Male	-	-	-	1	-	1	2	-	1	-	-	-
Female	1	-	2	1	2	1	2	2	1	-	-	-
Black, both sexes	1	2	1	-	-	1	-	-	-	-	-	-
Male	1	1	-	-	-	1	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Missouri												
All races, both sexes	1	2	2	6	5	19	13	12	10	2	1	-
Male	-	-	1	2	2	5	3	1	2	-	-	-
Female	1	2	1	4	3	14	10	11	8	2	1	-
White, both sexes	-	2	2	6	4	15	13	11	9	2	1	-
Male	-	-	1	2	1	4	3	1	1	-	-	-
Female	-	2	1	4	3	11	10	10	8	2	1	-
Black, both sexes	1	-	-	-	-	4	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	1	-	-	-	-	3	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Montana												
All races, both sexes	1	1	-	1	1	1	4	5	-	-	-	-
Male	1	-	-	-	-	-	1	1	-	-	-	-
Female	-	1	-	1	1	1	3	4	-	-	-	-
White, both sexes	1	1	-	1	-	1	4	5	-	-	-	-
Male	1	-	-	-	-	-	1	1	-	-	-	-
Female	-	1	-	1	-	1	3	4	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Nebraska												
All races, both sexes	1	1	3	1	2	3	7	1	3	1	-	-
Male	1	-	2	1	2	-	2	1	1	-	-	-
Female	-	1	1	-	-	3	5	-	2	1	-	-
White, both sexes	1	-	3	1	2	3	6	1	3	1	-	-
Male	1	-	2	1	2	-	2	1	1	-	-	-
Female	-	-	1	-	-	3	4	-	2	1	-	-
Black, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	2	-	1	1	5	1	1	-	2	-	-	-
Male	1	-	1	-	1	-	-	-	-	-	-	-
Female	1	-	-	1	4	1	1	-	2	-	-	-
White, both sexes	2	-	1	-	5	1	1	-	2	-	-	-
Male	1	-	1	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	4	1	1	-	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
New Hampshire												
All races, both sexes	-	-	1	1	2	3	1	5	2	-	-	-
Male	-	-	-	-	1	1	-	-	1	-	-	-
Female	-	-	1	1	1	2	1	5	1	-	-	-
White, both sexes	-	-	1	1	2	3	1	5	2	-	-	-
Male	-	-	-	-	1	1	-	-	1	-	-	-
Female	-	-	1	1	1	2	1	5	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	3	9	3	7	18	22	9	11	3	-	-
Male	-	2	2	1	5	6	6	2	2	1	-	-
Female	-	1	7	2	2	12	16	7	9	2	-	-
White, both sexes	-	3	6	3	7	15	22	8	11	3	-	-
Male	-	2	1	1	5	4	6	2	2	1	-	-
Female	-	1	5	2	2	11	16	6	9	2	-	-
Black, both sexes	-	-	2	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	2	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	2	-	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
New Mexico												
All races, both sexes	-	-	-	2	-	3	5	2	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	2	-	3	4	2	1	-	-	-
White, both sexes	-	-	-	1	-	3	5	2	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	3	4	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	7	2	8	7	10	27	36	35	12	6	2	-
Male	1	1	5	2	3	10	12	8	4	2	-	-
Female	6	1	3	5	7	17	24	27	8	4	2	-
White, both sexes	4	1	5	5	9	26	33	35	12	5	2	-
Male	1	1	3	1	3	9	9	8	4	1	-	-
Female	3	-	2	4	6	17	24	27	8	4	2	-
Black, both sexes	3	1	2	1	1	1	3	-	-	1	-	-
Male	-	-	2	-	-	1	3	-	-	1	-	-
Female	3	1	-	1	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
North Carolina												
All races, both sexes	2	1	8	9	14	8	15	8	5	5	-	-
Male	-	-	2	2	5	3	5	2	1	-	-	-
Female	2	1	6	7	9	5	10	6	4	5	-	-
White, both sexes	-	-	5	6	11	8	14	6	5	3	-	-
Male	-	-	1	2	3	3	5	2	1	-	-	-
Female	-	-	4	4	8	5	9	4	4	3	-	-
Black, both sexes	2	1	3	2	2	-	1	2	-	2	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	2	1	2	2	1	-	1	2	-	2	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	1	4	1	2	4	-	1	-	-	-
Male	-	-	1	1	1	1	2	-	-	-	-	-
Female	-	-	-	3	-	1	2	-	1	-	-	-
White, both sexes	-	-	-	4	-	2	4	-	1	-	-	-
Male	-	-	-	1	-	1	2	-	-	-	-	-
Female	-	-	-	3	-	1	2	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Ohio												
All races, both sexes	3	8	8	8	11	18	37	32	18	4	-	-
Male	-	4	3	2	2	6	8	8	3	1	-	-
Female	3	4	5	6	9	12	29	24	15	3	-	-
White, both sexes	1	6	7	7	9	17	35	31	18	4	-	-
Male	-	4	3	2	2	5	8	8	3	1	-	-
Female	1	2	4	5	7	12	27	23	15	3	-	-
Black, both sexes	2	2	1	1	2	1	2	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	2	2	1	1	2	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Oklahoma												
All races, both sexes	1	2	-	5	9	3	2	5	3	-	-	-
Male	1	1	-	2	3	2	-	3	-	-	-	-
Female	-	1	-	3	6	1	2	2	3	-	-	-
White, both sexes	1	2	-	3	9	3	2	5	3	-	-	-
Male	1	1	-	1	3	2	-	3	-	-	-	-
Female	-	1	-	2	6	1	2	2	3	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Oregon												
All races, both sexes	2	1	2	3	5	5	14	18	5	6	-	-
Male	2	1	1	1	2	1	5	3	4	1	-	-
Female	-	-	1	2	3	4	9	15	1	5	-	-
White, both sexes	2	1	2	2	5	5	14	18	5	6	-	-
Male	2	1	1	1	2	1	5	3	4	1	-	-
Female	-	-	1	1	3	4	9	15	1	5	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	5	7	12	7	12	19	35	43	14	8	1	-
Male	3	1	6	-	3	4	11	11	3	-	-	-
Female	2	6	6	7	9	15	24	32	11	8	1	-
White, both sexes	5	6	12	7	12	19	33	42	13	8	1	-
Male	3	1	6	-	3	4	10	11	3	-	-	-
Female	2	5	6	7	9	15	23	31	10	8	1	-
Black, both sexes	-	1	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Rhode Island												
All races, both sexes	2	1	-	1	-	3	7	4	-	-	-	-
Male	1	-	-	-	-	-	3	2	-	-	-	-
Female	1	1	-	1	-	3	4	2	-	-	-	-
White, both sexes	1	1	-	1	-	3	7	4	-	-	-	-
Male	1	-	-	-	-	-	3	2	-	-	-	-
Female	-	1	-	1	-	3	4	2	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	2	1	7	3	8	9	2	9	4	1	1	-
Male	-	1	5	2	2	4	1	2	1	-	-	-
Female	2	-	2	1	6	5	1	7	3	1	1	-
White, both sexes	1	1	6	2	8	7	1	6	4	1	1	-
Male	-	1	4	2	2	3	-	2	1	-	-	-
Female	1	-	2	-	6	4	1	4	3	1	1	-
Black, both sexes	1	-	1	1	-	2	1	3	-	-	-	-
Male	-	-	1	-	-	1	1	-	-	-	-	-
Female	1	-	-	1	-	1	-	3	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
South Dakota												
All races, both sexes	-	-	2	1	1	-	5	3	1	1	-	-
Male	-	-	1	1	-	-	1	2	-	-	-	-
Female	-	-	1	-	1	-	4	1	1	1	-	-
White, both sexes	-	-	2	1	1	-	5	3	1	1	-	-
Male	-	-	1	1	-	-	1	2	-	-	-	-
Female	-	-	1	-	1	-	4	1	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	3	3	2	5	9	16	9	14	5	3	-	-
Male	1	2	1	1	2	6	-	3	-	1	-	-
Female	2	1	1	4	7	10	9	11	5	2	-	-
White, both sexes	2	2	1	5	9	15	8	11	2	3	-	-
Male	-	1	-	1	2	6	-	2	-	1	-	-
Female	2	1	1	4	7	9	8	9	2	2	-	-
Black, both sexes	1	1	1	-	-	1	1	3	3	-	-	-
Male	1	1	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	1	2	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Texas												
All races, both sexes	5	11	11	10	18	15	25	29	8	4	2	-
Male	1	3	2	3	4	2	6	8	1	-	-	-
Female	4	8	9	7	14	13	19	21	7	4	2	-
White, both sexes	5	6	9	6	16	14	23	28	7	4	1	-
Male	1	1	1	3	4	2	5	8	1	-	-	-
Female	4	5	8	3	12	12	18	20	6	4	1	-
Black, both sexes	-	3	1	4	1	1	1	1	1	-	1	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	2	-	4	1	1	1	1	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	1	-	1	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	1	1	-	1	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	1	2	4	5	3	3	5	1	-	-	-
Male	-	1	1	1	1	2	-	4	-	-	-	-
Female	-	-	1	3	4	1	3	1	1	-	-	-
White, both sexes	-	1	2	4	5	3	3	5	1	-	-	-
Male	-	1	1	1	1	2	-	4	-	-	-	-
Female	-	-	1	3	4	1	3	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Vermont												
All races, both sexes	-	-	1	-	-	5	5	3	1	-	-	-
Male	-	-	1	-	-	2	1	-	-	-	-	-
Female	-	-	-	-	-	3	4	3	1	-	-	-
White, both sexes	-	-	1	-	-	5	5	3	1	-	-	-
Male	-	-	1	-	-	2	1	-	-	-	-	-
Female	-	-	-	-	-	3	4	3	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	3	3	6	3	7	6	17	9	1	4	1	-
Male	2	2	2	-	1	1	3	3	-	1	-	-
Female	1	1	4	3	6	5	14	6	1	3	1	-
White, both sexes	1	1	3	2	7	4	17	9	-	4	1	-
Male	1	1	2	-	1	1	3	3	-	1	-	-
Female	-	-	1	2	6	3	14	6	-	3	1	-
Black, both sexes	2	1	3	1	-	2	-	-	1	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	3	1	-	2	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Washington												
All races, both sexes	1	3	5	7	12	13	21	16	10	8	-	-
Male	-	1	2	3	3	4	6	6	4	2	-	-
Female	1	2	3	4	9	9	15	10	6	6	-	-
White, both sexes	-	3	5	5	12	13	21	16	10	8	-	-
Male	-	1	2	3	3	4	6	6	4	2	-	-
Female	-	2	3	2	9	9	15	10	6	6	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	4	1	1	3	7	1	1	2	-	-	-
Male	-	3	1	-	2	5	-	-	-	-	-	-
Female	-	1	-	1	1	2	1	1	2	-	-	-
White, both sexes	-	4	1	1	3	7	1	1	2	-	-	-
Male	-	3	1	-	2	5	-	-	-	-	-	-
Female	-	1	-	1	1	2	1	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Wisconsin												
All races, both sexes	2	2	3	5	5	15	18	13	9	4	-	-
Male	2	1	1	2	4	6	6	1	3	1	-	-
Female	-	1	2	3	1	9	12	12	6	3	-	-
White, both sexes	2	2	3	4	5	14	18	13	9	4	-	-
Male	2	1	1	2	4	6	6	1	3	1	-	-
Female	-	1	2	2	1	8	12	12	6	3	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	-	2	-	-	1	4	1	2	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	2	-	-	1	4	1	1	-	-	-
White, both sexes	1	-	2	-	-	1	4	1	2	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	2	-	-	1	4	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
United States												
All races, both sexes	1,572	1	-	-	1	1	2	5	7	13	26	52
Male	497	-	-	-	-	1	1	1	4	6	9	19
Female	1,075	1	-	-	1	-	1	4	3	7	17	33
White, both sexes	1,397	1	-	-	-	1	2	4	7	10	19	37
Male	440	-	-	-	-	1	1	-	4	4	6	12
Female	957	1	-	-	-	-	1	4	3	6	13	25
Black, both sexes	126	-	-	-	1	-	-	-	-	2	6	13
Male	41	-	-	-	-	-	-	-	-	1	2	7
Female	85	-	-	-	1	-	-	-	-	1	4	6
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	1	1	-
Male	4	-	-	-	-	-	-	-	-	1	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	41	-	-	-	-	-	-	1	-	-	-	2
Male	12	-	-	-	-	-	-	1	-	-	-	-
Female	29	-	-	-	-	-	-	-	-	-	-	2
Alabama												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Alaska												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	20	1	-	-	-	-	-	-	1	-	-	2
Male	9	-	-	-	-	-	-	-	1	-	-	-
Female	11	1	-	-	-	-	-	-	-	-	-	2
White, both sexes	20	1	-	-	-	-	-	-	1	-	-	2
Male	9	-	-	-	-	-	-	-	1	-	-	-
Female	11	1	-	-	-	-	-	-	-	-	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Arkansas												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	190	-	-	-	-	-	1	1	1	1	2	6
Male	71	-	-	-	-	-	-	-	-	-	1	3
Female	119	-	-	-	-	-	1	1	1	1	1	3
White, both sexes	160	-	-	-	-	-	1	1	1	1	2	5
Male	60	-	-	-	-	-	-	-	-	-	1	3
Female	100	-	-	-	-	-	1	1	1	1	1	2
Black, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Colorado												
All races, both sexes	41	-	-	-	-	-	-	-	1	-	-	1
Male	15	-	-	-	-	-	-	-	1	-	-	-
Female	26	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	40	-	-	-	-	-	-	-	1	-	-	1
Male	15	-	-	-	-	-	-	-	1	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	22	-	-	-	-	-	-	-	-	-	-	2
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	21	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Delaware												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Florida												
All races, both sexes	93	-	-	-	-	-	-	-	-	3	3	4
Male	42	-	-	-	-	-	-	-	-	3	1	2
Female	51	-	-	-	-	-	-	-	-	-	2	2
White, both sexes	85	-	-	-	-	-	-	-	-	2	2	4
Male	38	-	-	-	-	-	-	-	-	2	1	2
Female	47	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	8	-	-	-	-	-	-	-	-	1	1	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	30	-	-	-	-	-	-	-	1	1	-	-
Male	7	-	-	-	-	-	-	-	1	-	-	-
Female	23	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	23	-	-	-	-	-	-	-	1	-	-	-
Male	6	-	-	-	-	-	-	-	1	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Hawaii												
All races, both sexes	7	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Illinois												
All races, both sexes	65	-	-	-	-	-	-	1	-	1	1	2
Male	19	-	-	-	-	-	-	-	-	-	-	1
Female	46	-	-	-	-	-	-	1	-	1	1	1
White, both sexes	57	-	-	-	-	-	-	1	-	1	1	1
Male	18	-	-	-	-	-	-	-	-	-	-	1
Female	39	-	-	-	-	-	-	1	-	1	1	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	32	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	31	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Iowa												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	21	-	-	-	-	1	-	-	-	-	-	1
Male	7	-	-	-	-	1	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	20	-	-	-	-	1	-	-	-	-	-	1
Male	6	-	-	-	-	1	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Kentucky												
All races, both sexes	29	-	-	-	-	-	-	-	-	2	-	-
Male	10	-	-	-	-	-	-	-	-	1	-	-
Female	19	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	29	-	-	-	-	-	-	-	-	2	-	-
Male	10	-	-	-	-	-	-	-	-	1	-	-
Female	19	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	19	-	-	-	-	-	-	-	1	1	-	1
Male	6	-	-	-	-	-	-	-	-	1	-	1
Female	13	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	1	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	11	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Maine												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	27	-	-	-	1	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	1	-	-	-	-	-	-	1
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	8	-	-	-	1	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Massachusetts												
All races, both sexes	29	-	-	-	-	-	-	-	-	-	-	2
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	26	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	61	-	-	-	-	-	-	-	-	1	1	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	50	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	52	-	-	-	-	-	-	-	-	1	1	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	44	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Minnesota												
All races, both sexes	35	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	32	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	2
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Missouri												
All races, both sexes	28	-	-	-	-	-	-	-	-	-	-	2
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	21	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	23	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Nebraska												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
New Hampshire												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	55	-	-	-	-	-	-	-	1	-	1	-
Male	20	-	-	-	-	-	-	-	1	-	1	-
Female	35	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	52	-	-	-	-	-	-	-	1	-	1	-
Male	19	-	-	-	-	-	-	-	1	-	1	-
Female	33	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
New Mexico												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	81	-	-	-	-	-	1	-	-	-	4	5
Male	26	-	-	-	-	-	1	-	-	-	1	1
Female	55	-	-	-	-	-	-	-	-	-	3	4
White, both sexes	71	-	-	-	-	-	1	-	-	-	2	3
Male	22	-	-	-	-	-	1	-	-	-	-	1
Female	49	-	-	-	-	-	-	-	-	-	2	2
Black, both sexes	10	-	-	-	-	-	-	-	-	-	2	2
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
North Carolina												
All races, both sexes	37	-	-	-	-	-	-	1	-	-	3	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	26	-	-	-	-	-	-	1	-	-	3	-
White, both sexes	27	-	-	-	-	-	-	1	-	-	2	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	1	-	-	2	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	8	-	-	-	-	-	-	-	-	-	1	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Ohio												
All races, both sexes	83	-	-	-	-	-	-	1	-	1	3	1
Male	15	-	-	-	-	-	-	-	-	-	2	-
Female	68	-	-	-	-	-	-	1	-	1	1	1
White, both sexes	78	-	-	-	-	-	-	1	-	1	3	1
Male	14	-	-	-	-	-	-	-	-	-	2	-
Female	64	-	-	-	-	-	-	1	-	1	1	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	1	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	12	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Oregon												
All races, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	70	-	-	-	-	-	-	-	1	-	-	1
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	54	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	68	-	-	-	-	-	-	-	1	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	53	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Rhode Island												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	26	-	-	-	-	-	-	-	-	-	2	2
Male	12	-	-	-	-	-	-	-	-	-	1	1
Female	14	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	24	-	-	-	-	-	-	-	-	-	2	1
Male	10	-	-	-	-	-	-	-	-	-	1	-
Female	14	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
South Dakota												
All races, both sexes	9	-	-	-	-	-	-	-	-	1	1	-
Male	4	-	-	-	-	-	-	-	-	1	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	1	1	-
Male	2	-	-	-	-	-	-	-	-	1	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	32	-	-	-	-	-	-	-	-	-	-	2
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	24	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	28	-	-	-	-	-	-	-	-	-	-	2
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	21	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Texas												
All races, both sexes	89	-	-	-	-	-	-	-	-	1	2	4
Male	17	-	-	-	-	-	-	-	-	-	1	2
Female	72	-	-	-	-	-	-	-	-	1	1	2
White, both sexes	74	-	-	-	-	-	-	-	-	1	-	3
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	62	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	10	-	-	-	-	-	-	-	-	-	2	1
Male	3	-	-	-	-	-	-	-	-	-	1	1
Female	7	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Vermont												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	37	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	31	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Washington												
All races, both sexes	42	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	40	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	26	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Wisconsin												
All races, both sexes	39	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	38	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
United States												
All races, both sexes	50	81	111	114	167	228	305	250	122	32	4	-
Male	21	29	46	36	59	80	82	71	24	8	-	-
Female	29	52	65	78	108	148	223	179	98	24	4	-
White, both sexes	39	55	90	91	148	215	286	239	118	32	3	-
Male	20	20	39	32	54	75	72	70	22	8	-	-
Female	19	35	51	59	94	140	214	169	96	24	3	-
Black, both sexes	11	18	18	14	11	11	11	5	4	-	1	-
Male	1	7	6	2	2	5	6	-	2	-	-	-
Female	10	11	12	12	9	6	5	5	2	-	1	-
American Indian 1/, both sexes	-	-	2	2	2	-	-	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	1	2	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	8	1	7	6	2	8	6	-	-	-	-
Male	-	2	-	2	2	-	4	1	-	-	-	-
Female	-	6	1	5	4	2	4	5	-	-	-	-
Alabama												
All races, both sexes	1	-	1	3	4	1	4	2	1	-	-	-
Male	-	-	-	1	1	-	1	-	1	-	-	-
Female	1	-	1	2	3	1	3	2	-	-	-	-
White, both sexes	-	-	1	3	4	1	4	2	1	-	-	-
Male	-	-	-	1	1	-	1	-	1	-	-	-
Female	-	-	1	2	3	1	3	2	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Alaska												
All races, both sexes	-	2	-	1	1	-	-	-	1	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	2	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	1	-	1	1	-	-	-	1	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	2	2	1	5	4	1	1	-	-	-
Male	-	-	1	1	-	3	1	1	1	-	-	-
Female	-	-	1	1	1	2	3	-	-	-	-	-
White, both sexes	-	-	2	2	1	5	4	1	1	-	-	-
Male	-	-	1	1	-	3	1	1	1	-	-	-
Female	-	-	1	1	1	2	3	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Arkansas												
All races, both sexes	1	1	1	3	2	2	5	3	1	-	-	-
Male	1	-	-	1	2	1	1	1	-	-	-	-
Female	-	1	1	2	-	1	4	2	1	-	-	-
White, both sexes	1	1	1	3	2	2	5	2	1	-	-	-
Male	1	-	-	1	2	1	1	1	-	-	-	-
Female	-	1	1	2	-	1	4	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	9	9	13	15	22	22	38	27	19	3	1	-
Male	5	5	7	7	9	4	18	8	4	-	-	-
Female	4	4	6	8	13	18	20	19	15	3	1	-
White, both sexes	8	7	10	12	13	21	29	26	19	3	1	-
Male	5	4	6	5	7	4	13	8	4	-	-	-
Female	3	3	4	7	6	17	16	18	15	3	1	-
Black, both sexes	1	1	3	-	5	-	3	-	-	-	-	-
Male	-	1	1	-	1	-	2	-	-	-	-	-
Female	1	-	2	-	4	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	3	4	1	6	1	-	-	-	-
Male	-	-	-	2	1	-	3	-	-	-	-	-
Female	-	1	-	1	3	1	3	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Colorado												
All races, both sexes	2	1	1	1	5	7	6	14	2	-	-	-
Male	-	-	-	-	1	4	2	6	1	-	-	-
Female	2	1	1	1	4	3	4	8	1	-	-	-
White, both sexes	2	1	1	-	5	7	6	14	2	-	-	-
Male	-	-	-	-	1	4	2	6	1	-	-	-
Female	2	1	1	-	4	3	4	8	1	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	1	1	-	2	8	4	4	-	-	-	-
Male	-	1	-	-	-	2	1	2	-	-	-	-
Female	-	-	1	-	2	6	3	2	-	-	-	-
White, both sexes	-	1	1	-	2	8	4	4	-	-	-	-
Male	-	1	-	-	-	2	1	2	-	-	-	-
Female	-	-	1	-	2	6	3	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Delaware												
All races, both sexes	-	1	-	1	2	1	1	4	-	-	-	-
Male	-	-	-	-	1	1	-	1	-	-	-	-
Female	-	1	-	1	1	-	1	3	-	-	-	-
White, both sexes	-	1	-	1	2	1	1	4	-	-	-	-
Male	-	-	-	-	1	1	-	1	-	-	-	-
Female	-	1	-	1	1	-	1	3	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	-	2	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	2	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Florida												
All races, both sexes	3	5	7	4	10	18	12	11	12	1	-	-
Male	3	2	4	1	4	9	3	6	3	1	-	-
Female	-	3	3	3	6	9	9	5	9	-	-	-
White, both sexes	3	1	7	4	9	18	12	10	12	1	-	-
Male	3	-	4	1	3	9	3	6	3	1	-	-
Female	-	1	3	3	6	9	9	4	9	-	-	-
Black, both sexes	-	4	-	-	1	-	-	1	-	-	-	-
Male	-	2	-	-	1	-	-	-	-	-	-	-
Female	-	2	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	1	3	2	4	2	4	4	5	1	2	-	-
Male	-	-	1	1	-	-	2	1	-	1	-	-
Female	1	3	1	3	2	4	2	4	1	1	-	-
White, both sexes	-	1	1	4	2	3	4	4	1	2	-	-
Male	-	-	-	1	-	-	2	1	-	1	-	-
Female	-	1	1	3	2	3	2	3	1	1	-	-
Black, both sexes	1	1	1	-	-	1	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Hawaii												
All races, both sexes	-	-	-	2	-	-	3	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	3	1	-	-	-	-
White, both sexes	-	-	-	1	-	-	3	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	3	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Idaho												
All races, both sexes	1	1	1	-	-	2	1	1	1	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	1	1	1	-	-	2	-	-	1	-	-	-
White, both sexes	1	-	1	-	-	2	1	1	1	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	1	-	1	-	-	2	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Illinois												
All races, both sexes	1	3	5	4	3	16	12	10	5	-	1	-
Male	1	-	1	2	1	6	5	1	1	-	-	-
Female	-	3	4	2	2	10	7	9	4	-	1	-
White, both sexes	1	2	4	4	3	14	10	9	5	-	1	-
Male	1	-	1	2	1	5	5	1	1	-	-	-
Female	-	2	3	2	2	9	5	8	4	-	1	-
Black, both sexes	-	1	1	-	-	2	2	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	1	-	-	1	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Indiana												
All races, both sexes	1	1	5	3	4	2	7	4	5	-	-	-
Male	-	-	2	1	1	1	2	2	1	-	-	-
Female	1	1	3	2	3	1	5	2	4	-	-	-
White, both sexes	1	-	5	3	4	2	7	4	5	-	-	-
Male	-	-	2	1	1	1	2	2	1	-	-	-
Female	1	-	3	2	3	1	5	2	4	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Iowa												
All races, both sexes	-	1	-	1	1	3	4	3	2	-	-	-
Male	-	-	-	-	-	3	1	1	-	-	-	-
Female	-	1	-	1	1	-	3	2	2	-	-	-
White, both sexes	-	1	-	1	1	3	4	3	2	-	-	-
Male	-	-	-	-	-	3	1	1	-	-	-	-
Female	-	1	-	1	1	-	3	2	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	2	1	2	2	3	1	3	3	2	-	-	-
Male	1	-	1	-	-	1	-	2	-	-	-	-
Female	1	1	1	2	3	-	3	1	2	-	-	-
White, both sexes	2	1	2	2	3	1	3	2	2	-	-	-
Male	1	-	1	-	-	1	-	1	-	-	-	-
Female	1	1	1	2	3	-	3	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Kentucky												
All races, both sexes	1	1	5	5	3	3	7	2	-	-	-	-
Male	-	-	3	1	1	-	3	1	-	-	-	-
Female	1	1	2	4	2	3	4	1	-	-	-	-
White, both sexes	1	1	5	5	3	3	7	2	-	-	-	-
Male	-	-	3	1	1	-	3	1	-	-	-	-
Female	1	1	2	4	2	3	4	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	4	-	2	2	2	3	1	2	-	-	-
Male	-	2	-	-	1	-	1	-	-	-	-	-
Female	-	2	-	2	1	2	2	1	2	-	-	-
White, both sexes	-	2	-	-	2	2	3	1	2	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	2	-	-	1	2	2	1	2	-	-	-
Black, both sexes	-	2	-	2	-	-	-	-	-	-	-	-
Male	-	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	2	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Maine												
All races, both sexes	1	1	1	-	-	2	-	1	1	-	-	-
Male	-	-	1	-	-	1	-	-	1	-	-	-
Female	1	1	-	-	-	1	-	1	-	-	-	-
White, both sexes	1	1	1	-	-	2	-	1	1	-	-	-
Male	-	-	1	-	-	1	-	-	1	-	-	-
Female	1	1	-	-	-	1	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	4	3	1	1	6	4	4	1	-	-	-
Male	-	3	2	1	-	3	1	-	-	-	-	-
Female	1	1	1	-	1	3	3	4	1	-	-	-
White, both sexes	-	2	1	1	1	3	4	4	1	-	-	-
Male	-	1	1	1	-	2	1	-	-	-	-	-
Female	-	1	-	-	1	1	3	4	1	-	-	-
Black, both sexes	1	1	2	-	-	3	-	-	-	-	-	-
Male	-	1	1	-	-	1	-	-	-	-	-	-
Female	1	-	1	-	-	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Massachusetts												
All races, both sexes	-	1	-	3	1	3	12	4	3	-	-	-
Male	-	1	-	1	-	3	3	2	2	-	-	-
Female	-	-	-	2	1	-	9	2	1	-	-	-
White, both sexes	-	1	-	1	1	3	12	4	3	-	-	-
Male	-	1	-	-	-	3	3	2	2	-	-	-
Female	-	-	-	1	1	-	9	2	1	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1	3	6	4	12	3	10	13	7	-	-	-
Male	-	-	-	1	5	1	2	1	1	-	-	-
Female	1	3	6	3	7	2	8	12	6	-	-	-
White, both sexes	-	2	5	3	10	3	9	12	6	-	-	-
Male	-	-	-	1	4	1	1	1	-	-	-	-
Female	-	2	5	2	6	2	8	11	6	-	-	-
Black, both sexes	1	-	1	-	1	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	1	-	1	-	1	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Minnesota												
All races, both sexes	1	1	2	3	3	4	7	9	5	-	-	-
Male	1	-	-	1	1	2	3	2	1	-	-	-
Female	-	1	2	2	2	2	4	7	4	-	-	-
White, both sexes	1	-	-	3	3	4	7	9	5	-	-	-
Male	1	-	-	1	1	2	3	2	1	-	-	-
Female	-	-	-	2	2	2	4	7	4	-	-	-
Black, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1	1	-	2	1	1	2	2	1	-	-	-
Male	-	-	-	1	-	1	1	-	-	-	-	-
Female	1	1	-	1	1	-	1	2	1	-	-	-
White, both sexes	1	-	-	2	1	1	2	2	1	-	-	-
Male	-	-	-	1	-	1	1	-	-	-	-	-
Female	1	-	-	1	1	-	1	2	1	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Missouri												
All races, both sexes	1	1	1	-	1	7	4	5	5	1	-	-
Male	-	-	-	-	1	2	-	1	2	-	-	-
Female	1	1	1	-	-	5	4	4	3	1	-	-
White, both sexes	-	1	1	-	1	6	4	4	4	1	-	-
Male	-	-	-	-	1	2	-	1	1	-	-	-
Female	-	1	1	-	-	4	4	3	3	1	-	-
Black, both sexes	1	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	1	-	3	2	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	1	-	2	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	3	2	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	2	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Nebraska												
All races, both sexes	-	1	1	1	1	1	4	-	2	-	-	-
Male	-	-	-	1	1	-	1	-	1	-	-	-
Female	-	1	1	-	-	1	3	-	1	-	-	-
White, both sexes	-	-	1	1	1	1	3	-	2	-	-	-
Male	-	-	-	1	1	-	1	-	1	-	-	-
Female	-	-	1	-	-	1	2	-	1	-	-	-
Black, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1	-	-	-	1	1	-	-	1	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
White, both sexes	1	-	-	-	1	1	-	-	1	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
New Hampshire												
All races, both sexes	-	-	1	-	1	1	1	2	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	1	2	-	-	-	-
White, both sexes	-	-	1	-	1	1	1	2	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	1	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	3	7	2	6	13	13	3	6	-	-	-
Male	-	2	1	1	4	5	4	1	-	-	-	-
Female	-	1	6	1	2	8	9	2	6	-	-	-
White, both sexes	-	3	6	2	6	11	13	3	6	-	-	-
Male	-	2	1	1	4	4	4	1	-	-	-	-
Female	-	1	5	1	2	7	9	2	6	-	-	-
Black, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
New Mexico												
All races, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	2	1	5	2	6	14	17	18	3	3	-	-
Male	-	-	4	-	3	5	5	5	-	1	-	-
Female	2	1	1	2	3	9	12	13	3	2	-	-
White, both sexes	1	-	4	1	6	13	16	18	3	3	-	-
Male	-	-	3	-	3	4	4	5	-	1	-	-
Female	1	-	1	1	3	9	12	13	3	2	-	-
Black, both sexes	1	1	1	1	-	1	1	-	-	-	-	-
Male	-	-	1	-	-	1	1	-	-	-	-	-
Female	1	1	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
North Carolina												
All races, both sexes	1	-	4	5	7	3	7	2	2	1	-	-
Male	-	-	2	1	1	1	2	2	1	-	-	-
Female	1	-	2	4	6	2	5	-	1	1	-	-
White, both sexes	-	-	1	3	6	3	6	2	2	1	-	-
Male	-	-	1	1	1	1	2	2	1	-	-	-
Female	-	-	-	2	5	2	4	-	1	1	-	-
Black, both sexes	1	-	3	1	1	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	2	1	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	1	2	1	1	3	-	-	-	-	-
Male	-	-	1	-	1	-	1	-	-	-	-	-
Female	-	-	-	2	-	1	2	-	-	-	-	-
White, both sexes	-	-	-	2	-	1	3	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	2	-	1	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Ohio												
All races, both sexes	-	5	3	5	6	11	17	19	9	2	-	-
Male	-	2	2	-	-	4	1	3	-	1	-	-
Female	-	3	1	5	6	7	16	16	9	1	-	-
White, both sexes	-	4	3	5	4	10	17	18	9	2	-	-
Male	-	2	2	-	-	3	1	3	-	1	-	-
Female	-	2	1	5	4	7	16	15	9	1	-	-
Black, both sexes	-	1	-	-	2	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	2	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Oklahoma												
All races, both sexes	-	1	-	3	3	2	2	-	1	-	-	-
Male	-	1	-	2	1	1	-	-	-	-	-	-
Female	-	-	-	1	2	1	2	-	1	-	-	-
White, both sexes	-	1	-	1	3	2	2	-	1	-	-	-
Male	-	1	-	1	1	1	-	-	-	-	-	-
Female	-	-	-	-	2	1	2	-	1	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Oregon												
All races, both sexes	1	1	1	1	2	2	4	9	1	3	-	-
Male	1	1	1	1	1	-	1	2	1	-	-	-
Female	-	-	-	-	1	2	3	7	-	3	-	-
White, both sexes	1	1	1	1	2	2	4	9	1	3	-	-
Male	1	1	1	1	1	-	1	2	1	-	-	-
Female	-	-	-	-	1	2	3	7	-	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	5	4	6	3	3	9	14	17	3	4	-	-
Male	3	1	2	-	-	2	4	4	-	-	-	-
Female	2	3	4	3	3	7	10	13	3	4	-	-
White, both sexes	5	4	6	3	3	9	12	17	3	4	-	-
Male	3	1	2	-	-	2	3	4	-	-	-	-
Female	2	3	4	3	3	7	9	13	3	4	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Rhode Island												
All races, both sexes	2	-	-	1	-	3	3	2	-	-	-	-
Male	1	-	-	-	-	-	2	1	-	-	-	-
Female	1	-	-	1	-	3	1	1	-	-	-	-
White, both sexes	1	-	-	1	-	3	3	2	-	-	-	-
Male	1	-	-	-	-	-	2	1	-	-	-	-
Female	-	-	-	1	-	3	1	1	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	-	4	1	6	4	1	4	-	-	1	-
Male	-	-	3	1	2	2	1	1	-	-	-	-
Female	1	-	1	-	4	2	-	3	-	-	1	-
White, both sexes	1	-	4	1	6	4	-	4	-	-	1	-
Male	-	-	3	1	2	2	-	1	-	-	-	-
Female	1	-	1	-	4	2	-	3	-	-	1	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
South Dakota												
All races, both sexes	-	-	1	-	-	-	4	2	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	1	-	-	-	3	1	-	-	-	-
White, both sexes	-	-	1	-	-	-	4	2	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	1	-	-	-	3	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	1	2	3	3	8	4	6	1	2	-	-
Male	-	-	1	1	2	3	-	-	-	-	-	-
Female	-	1	1	2	1	5	4	6	1	2	-	-
White, both sexes	-	1	1	3	3	7	4	5	-	2	-	-
Male	-	-	-	1	2	3	-	-	-	-	-	-
Female	-	1	1	2	1	4	4	5	-	2	-	-
Black, both sexes	-	-	1	-	-	1	-	1	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Texas												
All races, both sexes	4	9	6	6	13	9	16	11	4	3	1	-
Male	1	3	1	2	3	-	1	3	-	-	-	-
Female	3	6	5	4	10	9	15	8	4	3	1	-
White, both sexes	4	5	5	3	11	9	15	11	4	3	-	-
Male	1	1	1	2	3	-	-	3	-	-	-	-
Female	3	4	4	1	8	9	15	8	4	3	-	-
Black, both sexes	-	2	-	3	1	-	-	-	-	-	1	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	3	1	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	1	-	1	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	1	1	-	1	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	1	2	3	2	2	-	-	-	-	-
Male	-	-	1	1	1	1	-	-	-	-	-	-
Female	-	-	-	1	2	1	2	-	-	-	-	-
White, both sexes	-	-	1	2	3	2	2	-	-	-	-	-
Male	-	-	1	1	1	1	-	-	-	-	-	-
Female	-	-	-	1	2	1	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Vermont												
All races, both sexes	-	-	1	-	-	1	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
White, both sexes	-	-	1	-	-	1	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	2	1	3	2	6	1	15	4	1	2	-	-
Male	1	1	-	-	1	1	3	2	-	1	-	-
Female	1	-	3	2	5	-	12	2	1	1	-	-
White, both sexes	1	1	-	1	6	1	15	4	-	2	-	-
Male	1	1	-	-	1	1	3	2	-	1	-	-
Female	-	-	-	1	5	-	12	2	-	1	-	-
Black, both sexes	1	-	3	1	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	3	1	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Washington												
All races, both sexes	-	2	3	4	8	4	7	7	4	3	-	-
Male	-	-	2	1	3	1	1	4	-	2	-	-
Female	-	2	1	3	5	3	6	3	4	1	-	-
White, both sexes	-	2	3	2	8	4	7	7	4	3	-	-
Male	-	-	2	1	3	1	1	4	-	2	-	-
Female	-	2	1	1	5	3	6	3	4	1	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	4	-	-	2	2	1	-	1	-	-	-
Male	-	3	-	-	1	1	-	-	-	-	-	-
Female	-	1	-	-	1	1	1	-	1	-	-	-
White, both sexes	-	4	-	-	2	2	1	-	1	-	-	-
Male	-	3	-	-	1	1	-	-	-	-	-	-
Female	-	1	-	-	1	1	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Wisconsin												
All races, both sexes	-	1	2	3	2	11	6	8	4	2	-	-
Male	-	1	1	1	2	5	1	1	1	1	-	-
Female	-	-	1	2	-	6	5	7	3	1	-	-
White, both sexes	-	1	2	2	2	11	6	8	4	2	-	-
Male	-	1	1	1	2	5	1	1	1	1	-	-
Female	-	-	1	1	-	6	5	7	3	1	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	-	-	-	-	1	2	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	1	2	-	-	-	-	-
White, both sexes	1	-	-	-	-	1	2	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	1	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
United States												
All races, both sexes	102	-	-	-	-	-	-	1	1	-	-	3
Male	34	-	-	-	-	-	-	-	1	-	-	1
Female	68	-	-	-	-	-	-	1	-	-	-	2
White, both sexes	99	-	-	-	-	-	-	-	1	-	-	2
Male	33	-	-	-	-	-	-	-	1	-	-	1
Female	66	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	1	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	1	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Colorado												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Florida												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Illinois												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Iowa												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Maine												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Massachusetts												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Minnesota												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Missouri												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Nebraska												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
New Hampshire												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
North Carolina												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Ohio												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Oregon												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
South Dakota												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	5	-	-	-	-	-	-	1	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	1	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	1	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Texas												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Vermont												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Washington												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Wisconsin												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
United States												
All races, both sexes	2	5	3	3	8	12	18	28	8	6	4	-
Male	1	3	1	2	1	8	5	7	2	2	-	-
Female	1	2	2	1	7	4	13	21	6	4	4	-
White, both sexes	2	5	3	3	8	12	18	28	8	5	4	-
Male	1	3	1	2	1	8	5	7	2	1	-	-
Female	1	2	2	1	7	4	13	21	6	4	4	-
Black, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1	1	1	1	1	-	1	3	2	-	2	-
Male	1	1	1	1	-	-	1	1	1	-	-	-
Female	-	-	-	-	1	-	-	2	1	-	2	-
White, both sexes	1	1	1	1	1	-	1	3	2	-	2	-
Male	1	1	1	1	-	-	1	1	1	-	-	-
Female	-	-	-	-	1	-	-	2	1	-	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Colorado												
All races, both sexes	-	-	-	-	-	1	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	1	-	-	-
White, both sexes	-	-	-	-	-	1	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Florida												
All races, both sexes	-	1	-	1	2	-	1	1	1	-	-	-
Male	-	1	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	1	1	-	-	1	1	-	-	-
White, both sexes	-	1	-	1	2	-	1	1	1	-	-	-
Male	-	1	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	1	1	-	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Illinois												
All races, both sexes	-	-	-	1	-	-	1	3	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	3	-	-	-	-
White, both sexes	-	-	-	1	-	-	1	3	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	3	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Iowa												
All races, both sexes	-	-	-	-	-	-	1	3	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	2	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	3	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	1	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Massachusetts												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Minnesota												
All races, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Missouri												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
New Hampshire												
All races, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1	-	-	-	-	-	-	3	-	2	1	-
Male	-	-	-	-	-	-	-	1	-	1	-	-
Female	1	-	-	-	-	-	-	2	-	1	1	-
White, both sexes	1	-	-	-	-	-	-	3	-	1	1	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	2	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
North Carolina												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Ohio												
All races, both sexes	-	-	-	-	-	2	1	1	1	1	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	1	1	-	-
White, both sexes	-	-	-	-	-	2	1	1	1	1	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Oregon												
All races, both sexes	-	-	-	-	1	1	1	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	1	-	-	1	-	-
White, both sexes	-	-	-	-	1	1	1	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	1	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	1	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	1	-	-	-	-
White, both sexes	-	-	-	-	1	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Texas												
All races, both sexes	-	1	-	-	-	2	-	2	-	-	-	-
Male	-	-	-	-	-	2	-	1	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	1	-	-	-	2	-	2	-	-	-	-
Male	-	-	-	-	-	2	-	1	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Vermont												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	1	1	1	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	1	-
White, both sexes	-	-	-	-	-	1	1	1	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Washington												
All races, both sexes	-	-	-	-	-	-	1	2	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	1	-	-	-
White, both sexes	-	-	-	-	-	-	1	2	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
United States												
All races, both sexes	651	1	-	-	-	1	-	2	1	4	7	6
Male	238	-	-	-	-	1	-	1	1	4	2	3
Female	413	1	-	-	-	-	-	1	-	-	5	3
White, both sexes	606	-	-	-	-	1	-	1	1	3	4	6
Male	222	-	-	-	-	1	-	1	1	3	2	3
Female	384	-	-	-	-	-	-	-	-	-	2	3
Black, both sexes	31	-	-	-	-	-	-	-	-	1	2	-
Male	13	-	-	-	-	-	-	-	-	1	-	-
Female	18	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	1	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	1	-	-	1	-
Alabama												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	7	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Arkansas												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	93	-	-	-	-	1	-	1	1	-	1	-
Male	38	-	-	-	-	1	-	-	1	-	-	-
Female	55	-	-	-	-	-	-	1	-	-	1	-
White, both sexes	80	-	-	-	-	1	-	-	1	-	-	-
Male	33	-	-	-	-	1	-	-	1	-	-	-
Female	47	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	1	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	1	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Colorado												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Delaware												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Florida												
All races, both sexes	54	-	-	-	-	-	-	-	-	1	1	-
Male	22	-	-	-	-	-	-	-	-	1	-	-
Female	32	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	52	-	-	-	-	-	-	-	-	1	-	-
Male	22	-	-	-	-	-	-	-	-	1	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Hawaii												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Illinois												
All races, both sexes	15	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Iowa												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Kentucky												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Maine												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Massachusetts												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Minnesota												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	2	-
Male	8	-	-	-	-	-	-	-	-	-	2	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	23	-	-	-	-	-	-	-	-	-	2	-
Male	8	-	-	-	-	-	-	-	-	-	2	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Missouri												
All races, both sexes	22	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	21	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Nebraska												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
New Hampshire												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
New Mexico												
All races, both sexes	5	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	27	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
North Carolina												
All races, both sexes	12	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Ohio												
All races, both sexes	32	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	30	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Oregon												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	38	-	-	-	-	-	-	-	-	-	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	37	-	-	-	-	-	-	-	-	-	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Rhode Island												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
South Dakota												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Texas												
All races, both sexes	33	-	-	-	-	-	-	1	-	-	1	2
Male	12	-	-	-	-	-	-	1	-	-	-	1
Female	21	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	29	-	-	-	-	-	-	1	-	-	1	2
Male	11	-	-	-	-	-	-	1	-	-	-	1
Female	18	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Vermont												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Washington												
All races, both sexes	27	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	26	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Wisconsin												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
United States												
All races, both sexes	17	21	29	31	64	78	124	139	93	27	6	-
Male	8	11	14	12	23	30	48	48	29	3	-	-
Female	9	10	15	19	41	48	76	91	64	24	6	-
White, both sexes	12	15	24	31	60	74	121	128	92	27	6	-
Male	7	8	12	12	22	27	46	45	29	3	-	-
Female	5	7	12	19	38	47	75	83	63	24	6	-
Black, both sexes	3	4	4	-	2	3	2	9	1	-	-	-
Male	1	1	2	-	1	2	2	3	-	-	-	-
Female	2	3	2	-	1	1	-	6	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	1	-	2	1	1	2	-	-	-	-
Male	-	2	-	-	-	1	-	-	-	-	-	-
Female	2	-	1	-	2	-	1	2	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	2	1	3	1	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	1	1	2	1	-	-	-	-
White, both sexes	-	-	-	-	2	1	3	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	1	1	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1	-	-	1	-	-	2	2	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	2	1	-	-	-	-
White, both sexes	1	-	-	1	-	-	2	2	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	2	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Arkansas												
All races, both sexes	-	-	1	-	-	-	1	1	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	1	2	-	-	-
White, both sexes	-	-	1	-	-	-	1	1	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1	7	8	4	9	14	13	15	16	1	1	-
Male	-	4	7	1	2	5	7	6	3	1	-	-
Female	1	3	1	3	7	9	6	9	13	-	1	-
White, both sexes	1	4	7	4	7	14	12	11	16	1	1	-
Male	-	1	7	1	2	5	7	4	3	1	-	-
Female	1	3	-	3	5	9	5	7	13	-	1	-
Black, both sexes	-	1	-	-	1	-	-	2	-	-	-	-
Male	-	1	-	-	-	-	-	2	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	1	-	1	-	1	2	-	-	-	-
Male	-	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	1	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Colorado												
All races, both sexes	-	-	-	2	2	1	2	1	1	-	-	-
Male	-	-	-	-	2	-	-	-	-	-	-	-
Female	-	-	-	2	-	1	2	1	1	-	-	-
White, both sexes	-	-	-	2	2	1	2	1	1	-	-	-
Male	-	-	-	-	2	-	-	-	-	-	-	-
Female	-	-	-	2	-	1	2	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	1	2	3	2	5	1	1	1	-
Male	-	-	-	1	1	2	1	3	1	-	-	-
Female	-	-	-	-	1	1	1	2	-	1	1	-
White, both sexes	-	-	-	1	1	3	2	5	1	1	1	-
Male	-	-	-	1	1	2	1	3	1	-	-	-
Female	-	-	-	-	-	1	1	2	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Florida												
All races, both sexes	1	3	2	1	4	4	13	14	7	3	-	-
Male	1	2	-	1	2	1	2	5	6	1	-	-
Female	-	1	2	-	2	3	11	9	1	2	-	-
White, both sexes	1	2	2	1	4	4	13	14	7	3	-	-
Male	1	2	-	1	2	1	2	5	6	1	-	-
Female	-	-	2	-	2	3	11	9	1	2	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	2	2	2	1	1	-	1	-
Male	-	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	-	-	1	1	1	1	1	-	1	-
White, both sexes	-	-	-	-	1	2	2	1	1	-	1	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	1	1	1	1	1	-	1	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Hawaii												
All races, both sexes	1	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	1	-	1	3	1	-	-	-
Male	-	-	-	-	-	-	1	2	-	-	-	-
Female	-	-	-	-	1	-	-	1	1	-	-	-
White, both sexes	-	-	-	-	1	-	1	3	1	-	-	-
Male	-	-	-	-	-	-	1	2	-	-	-	-
Female	-	-	-	-	1	-	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Illinois												
All races, both sexes	-	-	1	2	2	2	4	2	1	-	-	-
Male	-	-	-	2	1	-	-	1	-	-	-	-
Female	-	-	1	-	1	2	4	1	1	-	-	-
White, both sexes	-	-	1	2	2	2	4	2	1	-	-	-
Male	-	-	-	2	1	-	-	1	-	-	-	-
Female	-	-	1	-	1	2	4	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	1	1	5	3	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	1	4	3	-	-	-
White, both sexes	-	-	-	-	-	1	1	5	3	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	1	4	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Iowa												
All races, both sexes	-	-	-	1	3	2	1	2	2	1	-	-
Male	-	-	-	-	2	-	1	-	1	-	-	-
Female	-	-	-	1	1	2	-	2	1	1	-	-
White, both sexes	-	-	-	1	3	2	1	2	2	1	-	-
Male	-	-	-	-	2	-	1	-	1	-	-	-
Female	-	-	-	1	1	2	-	2	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	-	-	-	2	-	3	-	1	2	-	-
Male	-	-	-	-	1	-	1	-	1	-	-	-
Female	1	-	-	-	1	-	2	-	-	2	-	-
White, both sexes	1	-	-	-	2	-	3	-	1	2	-	-
Male	-	-	-	-	1	-	1	-	1	-	-	-
Female	1	-	-	-	1	-	2	-	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Kentucky												
All races, both sexes	-	1	-	-	-	1	1	2	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	2	-	-	-	-
White, both sexes	-	1	-	-	-	1	1	2	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	1	-	2	-	1	2	1	-	-	-	-
Male	1	-	-	1	-	1	1	-	-	-	-	-
Female	-	1	-	1	-	-	1	1	-	-	-	-
White, both sexes	-	1	-	2	-	1	2	1	-	-	-	-
Male	-	-	-	1	-	1	1	-	-	-	-	-
Female	-	1	-	1	-	-	1	1	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Maine												
All races, both sexes	1	-	-	-	1	-	1	3	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	1	-	1	2	-	-	-	-
White, both sexes	1	-	-	-	1	-	1	3	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	1	-	1	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	1	1	2	2	-	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	1	1	1	1	-	-	-
White, both sexes	-	-	-	-	-	1	1	2	2	-	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	1	1	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	4	2	5	1	-	-
Male	-	-	-	-	-	-	3	2	2	-	-	-
Female	-	-	-	-	-	-	1	-	3	1	-	-
White, both sexes	-	-	-	-	-	-	4	2	5	1	-	-
Male	-	-	-	-	-	-	3	2	2	-	-	-
Female	-	-	-	-	-	-	1	-	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1	1	1	-	2	3	3	2	3	1	-	-
Male	1	1	-	-	1	1	2	2	1	-	-	-
Female	-	-	1	-	1	2	1	-	2	1	-	-
White, both sexes	1	1	1	-	2	3	3	2	3	1	-	-
Male	1	1	-	-	1	1	2	2	1	-	-	-
Female	-	-	1	-	1	2	1	-	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Minnesota												
All races, both sexes	1	-	1	2	1	3	2	5	5	1	-	-
Male	-	-	1	1	-	1	-	3	-	-	-	-
Female	1	-	-	1	1	2	2	2	5	1	-	-
White, both sexes	1	-	1	2	1	3	2	5	5	1	-	-
Male	-	-	1	1	-	1	-	3	-	-	-	-
Female	1	-	-	1	1	2	2	2	5	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	1	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Missouri												
All races, both sexes	-	-	-	1	3	6	4	5	1	-	1	-
Male	-	-	-	-	-	2	2	-	-	-	-	-
Female	-	-	-	1	3	4	2	5	1	-	1	-
White, both sexes	-	-	-	1	3	5	4	5	1	-	1	-
Male	-	-	-	-	-	1	2	-	-	-	-	-
Female	-	-	-	1	3	4	2	5	1	-	1	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Nebraska												
All races, both sexes	1	-	1	-	1	-	-	-	1	-	-	-
Male	1	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	1	-	1	-	1	-	-	-	1	-	-	-
Male	1	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	2	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	2	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
New Hampshire												
All races, both sexes	-	-	-	1	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	1	-	-	-
White, both sexes	-	-	-	1	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	1	-	-	3	3	2	4	2	-	-
Male	-	-	-	-	-	-	1	-	2	-	-	-
Female	-	-	1	-	-	3	2	2	2	2	-	-
White, both sexes	-	-	-	-	-	3	3	2	4	2	-	-
Male	-	-	-	-	-	-	1	-	2	-	-	-
Female	-	-	-	-	-	3	2	2	2	2	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
New Mexico												
All races, both sexes	-	-	-	1	-	-	2	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	1	1	-	-	-	-
White, both sexes	-	-	-	1	-	-	2	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1	-	2	1	-	4	8	5	4	1	1	-
Male	1	-	1	-	-	2	5	1	2	-	-	-
Female	-	-	1	1	-	2	3	4	2	1	1	-
White, both sexes	1	-	1	1	-	4	6	5	4	1	1	-
Male	1	-	-	-	-	2	3	1	2	-	-	-
Female	-	-	1	1	-	2	3	4	2	1	1	-
Black, both sexes	-	-	1	-	-	-	2	-	-	-	-	-
Male	-	-	1	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
North Carolina												
All races, both sexes	1	-	-	1	2	1	2	2	2	-	-	-
Male	-	-	-	1	1	1	-	-	-	-	-	-
Female	1	-	-	-	1	-	2	2	2	-	-	-
White, both sexes	-	-	-	1	2	1	2	1	2	-	-	-
Male	-	-	-	1	1	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	2	1	2	-	-	-
Black, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Ohio												
All races, both sexes	1	2	3	-	3	1	8	9	4	1	-	-
Male	-	2	-	-	-	1	2	3	1	-	-	-
Female	1	-	3	-	3	-	6	6	3	1	-	-
White, both sexes	-	2	2	-	3	1	8	9	4	1	-	-
Male	-	2	-	-	-	1	2	3	1	-	-	-
Female	-	-	2	-	3	-	6	6	3	1	-	-
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1	1	-	1	4	-	-	2	1	-	-	-
Male	1	-	-	-	2	-	-	2	-	-	-	-
Female	-	1	-	1	2	-	-	-	1	-	-	-
White, both sexes	1	1	-	1	4	-	-	2	1	-	-	-
Male	1	-	-	-	2	-	-	2	-	-	-	-
Female	-	1	-	1	2	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Oregon												
All races, both sexes	-	-	-	1	1	2	1	5	3	1	-	-
Male	-	-	-	-	1	1	1	1	2	-	-	-
Female	-	-	-	1	-	1	-	4	1	1	-	-
White, both sexes	-	-	-	1	1	2	1	5	3	1	-	-
Male	-	-	-	-	1	1	1	1	2	-	-	-
Female	-	-	-	1	-	1	-	4	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	1	2	-	3	4	8	9	6	4	-	-
Male	-	-	1	-	1	2	4	4	2	-	-	-
Female	-	1	1	-	2	2	4	5	4	4	-	-
White, both sexes	-	-	2	-	3	4	8	9	6	4	-	-
Male	-	-	1	-	1	2	4	4	2	-	-	-
Female	-	-	1	-	2	2	4	5	4	4	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Rhode Island												
All races, both sexes	-	1	-	-	-	-	2	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	1	-	-	-	-	2	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	1	-	1	-	3	1	4	2	-	-	-
Male	-	1	-	1	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	2	1	3	2	-	-	-
White, both sexes	-	1	-	1	-	2	1	2	2	-	-	-
Male	-	1	-	1	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	1	1	1	2	-	-	-
Black, both sexes	-	-	-	-	-	1	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
South Dakota												
All races, both sexes	-	-	1	-	-	-	-	1	-	1	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	1	-	-	-	-	1	-	1	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	1	3	4	1	4	1	1	-	-
Male	-	-	-	-	-	2	-	2	-	1	-	-
Female	-	-	-	1	3	2	1	2	1	-	-	-
White, both sexes	-	-	-	1	3	4	1	2	1	1	-	-
Male	-	-	-	-	-	2	-	1	-	1	-	-
Female	-	-	-	1	3	2	1	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Texas												
All races, both sexes	1	1	1	2	2	1	5	11	4	-	1	-
Male	-	-	1	1	1	-	3	3	1	-	-	-
Female	1	1	-	1	1	1	2	8	3	-	1	-
White, both sexes	1	-	-	2	2	1	5	10	3	-	1	-
Male	-	-	-	1	1	-	3	3	1	-	-	-
Female	1	-	-	1	1	1	2	7	2	-	1	-
Black, both sexes	-	1	1	-	-	-	-	1	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Vermont												
All races, both sexes	-	-	-	-	-	-	2	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	1	1	-	-	-
White, both sexes	-	-	-	-	-	-	2	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Washington												
All races, both sexes	1	1	1	1	3	5	7	2	3	3	-	-
Male	-	1	-	1	-	2	2	1	2	-	-	-
Female	1	-	1	-	3	3	5	1	1	3	-	-
White, both sexes	-	1	1	1	3	5	7	2	3	3	-	-
Male	-	1	-	1	-	2	2	1	2	-	-	-
Female	-	-	1	-	3	3	5	1	1	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	1	1	-	-	-	1	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	1	-	-	-
White, both sexes	-	-	1	1	-	-	-	1	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Wisconsin												
All races, both sexes	-	-	-	1	3	1	6	2	2	-	-	-
Male	-	-	-	1	2	-	4	-	1	-	-	-
Female	-	-	-	-	1	1	2	2	1	-	-	-
White, both sexes	-	-	-	1	3	1	6	2	2	-	-	-
Male	-	-	-	1	2	-	4	-	1	-	-	-
Female	-	-	-	-	1	1	2	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
United States												
All races, both sexes	844	-	1	-	-	1	6	7	4	7	13	20
Male	251	-	1	-	-	-	3	5	1	3	3	10
Female	593	-	-	-	-	1	3	2	3	4	10	10
White, both sexes	721	-	1	-	-	1	4	3	1	6	9	13
Male	207	-	1	-	-	-	2	2	-	2	3	6
Female	514	-	-	-	-	1	2	1	1	4	6	7
Black, both sexes	84	-	-	-	-	-	2	2	1	-	3	5
Male	29	-	-	-	-	-	1	2	-	-	-	2
Female	55	-	-	-	-	-	1	-	1	-	3	3
American Indian 1/, both sexes	7	-	-	-	-	-	-	1	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	32	-	-	-	-	-	-	1	2	1	1	1
Male	13	-	-	-	-	-	-	1	1	1	-	1
Female	19	-	-	-	-	-	-	-	1	-	1	-
Alabama												
All races, both sexes	10	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	12	-	-	-	-	-	-	1	-	-	-	-
Male	5	-	-	-	-	-	-	1	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	1	-	-	-	-
Male	5	-	-	-	-	-	-	1	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Arkansas												
All races, both sexes	4	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	84	-	-	-	-	-	-	-	1	1	3	3
Male	34	-	-	-	-	-	-	-	1	-	1	2
Female	50	-	-	-	-	-	-	-	-	1	2	1
White, both sexes	66	-	-	-	-	-	-	-	-	1	3	2
Male	26	-	-	-	-	-	-	-	-	-	1	1
Female	40	-	-	-	-	-	-	-	-	1	2	1
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	1	-	-	-
Male	5	-	-	-	-	-	-	-	1	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Colorado												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Delaware												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Florida												
All races, both sexes	50	-	-	-	-	-	-	-	-	2	-	2
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	33	-	-	-	-	-	-	-	-	2	-	2
White, both sexes	42	-	-	-	-	-	-	-	-	2	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	-	-	-	-	-	2	-	1
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	12	-	-	-	-	1	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	1	-	-	1	-	-	-
White, both sexes	10	-	-	-	-	1	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	1	-	-	1	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Hawaii												
All races, both sexes	7	-	-	-	-	-	-	1	-	1	1	1
Male	4	-	-	-	-	-	-	1	-	1	-	1
Female	3	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	1	-	1	-	1
Male	4	-	-	-	-	-	-	1	-	1	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Illinois												
All races, both sexes	32	-	-	-	-	-	-	-	-	-	2	1
Male	11	-	-	-	-	-	-	-	-	-	1	1
Female	21	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	27	-	-	-	-	-	-	-	-	-	1	1
Male	9	-	-	-	-	-	-	-	-	-	1	1
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Indiana												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	1	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	15	-	-	-	-	-	-	-	-	-	1	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Iowa												
All races, both sexes	29	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	27	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Kentucky												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Maine												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	17	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Massachusetts												
All races, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	27	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	30	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	27	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Minnesota												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Missouri												
All races, both sexes	27	-	-	-	-	-	-	1	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	23	-	-	-	-	-	-	1	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Nebraska												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	2
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
New Hampshire												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
New Mexico												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	49	-	-	-	-	-	-	-	-	-	2	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	38	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	42	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	32	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	2	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
North Carolina												
All races, both sexes	32	-	-	-	-	-	1	-	-	-	-	-
Male	8	-	-	-	-	-	1	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	25	-	-	-	-	-	1	-	-	-	-	-
Male	6	-	-	-	-	-	1	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Ohio												
All races, both sexes	33	-	-	-	-	-	1	-	-	-	-	-
Male	14	-	-	-	-	-	1	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	27	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Oregon												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	56	-	-	-	-	-	-	-	-	-	1	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	45	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	54	-	-	-	-	-	-	-	-	-	1	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	43	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Rhode Island												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
South Dakota												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	28	-	-	-	-	-	3	1	-	-	2	1
Male	7	-	-	-	-	-	1	1	-	-	-	-
Female	21	-	-	-	-	-	2	-	-	-	2	1
White, both sexes	21	-	-	-	-	-	3	-	-	-	1	1
Male	4	-	-	-	-	-	1	-	-	-	-	-
Female	17	-	-	-	-	-	2	-	-	-	1	1
Black, both sexes	7	-	-	-	-	-	-	1	-	-	1	-
Male	3	-	-	-	-	-	-	1	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Texas												
All races, both sexes	27	-	1	-	-	-	-	1	-	1	-	2
Male	8	-	1	-	-	-	-	1	-	1	-	2
Female	19	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	24	-	1	-	-	-	-	1	-	1	-	2
Male	8	-	1	-	-	-	-	1	-	1	-	2
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	11	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Vermont												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	19	-	-	-	-	-	-	1	-	-	-	1
Male	6	-	-	-	-	-	-	1	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	1	-	-	-	1
Male	3	-	-	-	-	-	-	1	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Washington												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	24	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Wisconsin												
All races, both sexes	23	-	-	-	-	-	-	-	-	2	-	-
Male	6	-	-	-	-	-	-	-	-	1	-	-
Female	17	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	2	-	-
Male	6	-	-	-	-	-	-	-	-	1	-	-
Female	16	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
United States												
All races, both sexes	26	35	49	66	72	118	157	151	82	27	2	-
Male	14	13	16	24	19	33	45	33	21	7	-	-
Female	12	22	33	42	53	85	112	118	61	20	2	-
White, both sexes	14	21	42	51	62	104	145	140	78	24	2	-
Male	8	8	12	19	14	28	43	31	21	7	-	-
Female	6	13	30	32	48	76	102	109	57	17	2	-
Black, both sexes	10	9	4	11	5	9	10	8	2	3	-	-
Male	5	5	3	4	2	3	1	1	-	-	-	-
Female	5	4	1	7	3	6	9	7	2	3	-	-
American Indian 1/, both sexes	-	1	-	2	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	1	-	2	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	4	3	2	4	4	2	3	2	-	-	-
Male	1	-	1	1	2	2	1	1	-	-	-	-
Female	1	4	2	1	2	2	1	2	2	-	-	-
Alabama												
All races, both sexes	1	1	-	1	2	1	2	1	-	-	-	-
Male	1	-	-	1	2	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	2	1	-	-	-	-
White, both sexes	-	1	-	1	2	-	1	1	-	-	-	-
Male	-	-	-	1	2	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	1	-	-	-	-
Black, both sexes	1	-	-	-	-	1	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	2	-	2	-	3	1	1	2	-	-	-
Male	-	-	-	1	-	3	-	-	-	-	-	-
Female	-	2	-	1	-	-	1	1	2	-	-	-
White, both sexes	-	2	-	2	-	3	1	1	2	-	-	-
Male	-	-	-	1	-	3	-	-	-	-	-	-
Female	-	2	-	1	-	-	1	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Arkansas												
All races, both sexes	-	-	-	1	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	2	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	2	6	5	4	11	11	12	14	7	4	-	-
Male	1	3	-	2	2	7	6	5	2	2	-	-
Female	1	3	5	2	9	4	6	9	5	2	-	-
White, both sexes	-	4	5	2	8	10	9	12	7	3	-	-
Male	-	3	-	1	1	6	5	4	2	2	-	-
Female	-	1	5	1	7	4	4	8	5	1	-	-
Black, both sexes	-	-	-	2	1	-	3	-	-	1	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	1	1	-	2	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	-	-	2	1	-	2	-	-	-	-
Male	1	-	-	-	1	1	-	1	-	-	-	-
Female	1	2	-	-	1	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Colorado												
All races, both sexes	1	1	1	1	1	3	1	-	1	-	-	-
Male	1	1	1	1	-	1	-	-	1	-	-	-
Female	-	-	-	-	1	2	1	-	-	-	-	-
White, both sexes	1	1	1	1	1	2	1	-	1	-	-	-
Male	1	1	1	1	-	1	-	-	1	-	-	-
Female	-	-	-	-	1	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	1	1	1	3	-	1	3	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	1	1	2	-	1	3	-	-	-
White, both sexes	-	-	-	-	1	2	-	1	3	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	2	-	1	3	-	-	-
Black, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Delaware												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Florida												
All races, both sexes	1	2	3	2	4	6	10	9	6	3	-	-
Male	-	1	-	2	2	2	3	3	2	2	-	-
Female	1	1	3	-	2	4	7	6	4	1	-	-
White, both sexes	-	1	3	1	3	6	8	8	6	3	-	-
Male	-	-	-	1	2	2	3	2	2	2	-	-
Female	-	1	3	-	1	4	5	6	4	1	-	-
Black, both sexes	1	1	-	1	1	-	1	1	-	-	-	-
Male	-	1	-	1	-	-	-	1	-	-	-	-
Female	1	-	-	-	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Georgia												
All races, both sexes	1	2	-	2	2	1	-	2	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	2	2	1	-	2	-	-	-	-
White, both sexes	1	-	-	2	2	1	-	2	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	2	2	1	-	2	-	-	-	-
Black, both sexes	-	2	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Idaho												
All races, both sexes	-	-	1	1	2	-	2	2	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	2	-	2	2	2	-	-	-
White, both sexes	-	-	1	1	2	-	2	2	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	2	-	2	2	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Illinois												
All races, both sexes	1	1	-	3	-	8	4	10	1	1	-	-
Male	1	1	-	2	-	2	1	2	-	-	-	-
Female	-	-	-	1	-	6	3	8	1	1	-	-
White, both sexes	-	1	-	2	-	7	3	10	1	1	-	-
Male	-	1	-	2	-	1	1	2	-	-	-	-
Female	-	-	-	-	-	6	2	8	1	1	-	-
Black, both sexes	1	-	-	1	-	1	1	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	2	1	1	3	1	4	1	-	-	-
Male	-	-	-	1	-	-	1	2	1	-	-	-
Female	-	-	2	-	1	3	-	2	-	-	-	-
White, both sexes	-	-	2	1	1	3	1	4	1	-	-	-
Male	-	-	-	1	-	-	1	2	1	-	-	-
Female	-	-	2	-	1	3	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Iowa												
All races, both sexes	1	1	-	1	3	3	5	9	5	1	-	-
Male	1	-	-	1	-	1	1	2	2	1	-	-
Female	-	1	-	-	3	2	4	7	3	-	-	-
White, both sexes	1	-	-	1	2	3	5	9	5	1	-	-
Male	1	-	-	1	-	1	1	2	2	1	-	-
Female	-	-	-	-	2	2	4	7	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	1	1	-	-	-	1	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Kentucky												
All races, both sexes	-	-	-	-	2	1	2	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	2	1	2	-	1	-	-	-
White, both sexes	-	-	-	-	2	1	2	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	2	1	2	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	1	-	-	-	1	1	1	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
White, both sexes	-	-	-	-	-	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Maine												
All races, both sexes	-	-	-	1	-	1	-	1	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	1	-	1	-	1	1	-	-	-
White, both sexes	-	-	-	1	-	1	-	1	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	1	-	1	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	2	-	2	-	2	3	6	1	-	-	-
Male	-	-	-	1	-	-	-	2	-	-	-	-
Female	-	2	-	1	-	2	3	4	1	-	-	-
White, both sexes	-	1	-	1	-	2	3	5	1	-	-	-
Male	-	-	-	-	-	-	-	2	-	-	-	-
Female	-	1	-	1	-	2	3	3	1	-	-	-
Black, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Massachusetts												
All races, both sexes	-	1	1	4	1	3	8	5	4	1	-	-
Male	-	-	-	2	-	-	3	-	1	-	-	-
Female	-	1	1	2	1	3	5	5	3	1	-	-
White, both sexes	-	1	1	3	1	3	8	5	4	1	-	-
Male	-	-	-	1	-	-	3	-	1	-	-	-
Female	-	1	1	2	1	3	5	5	3	1	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	2	1	-	2	5	3	9	3	3	-	1	-
Male	1	1	-	1	3	1	3	-	1	-	-	-
Female	1	-	-	1	2	2	6	3	2	-	1	-
White, both sexes	2	1	-	2	4	2	9	2	3	-	1	-
Male	1	1	-	1	2	1	3	-	1	-	-	-
Female	1	-	-	1	2	1	6	2	2	-	1	-
Black, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Minnesota												
All races, both sexes	1	-	2	-	-	1	2	-	4	2	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	2	-	4	2	-	-
White, both sexes	1	-	2	-	-	1	2	-	4	2	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	2	-	4	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1	1	2	-	1	1	1	-	1	-	-	-
Male	1	1	-	-	-	-	-	-	1	-	-	-
Female	-	-	2	-	1	1	1	-	-	-	-	-
White, both sexes	-	-	1	-	1	1	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	1	-	1	1	1	-	-	-	-	-
Black, both sexes	1	1	1	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Missouri												
All races, both sexes	-	1	1	5	1	6	4	2	4	1	-	-
Male	-	-	1	2	1	1	1	-	-	-	-	-
Female	-	1	-	3	-	5	3	2	4	1	-	-
White, both sexes	-	1	1	5	-	4	4	2	4	1	-	-
Male	-	-	1	2	-	1	1	-	-	-	-	-
Female	-	1	-	3	-	3	3	2	4	1	-	-
Black, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	1	-	1	-	1	1	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	1	1	2	-	-	-	-
White, both sexes	-	1	-	1	-	1	1	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	1	1	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Nebraska												
All races, both sexes	-	-	1	-	-	2	2	1	-	1	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	2	2	-	-	1	-	-
White, both sexes	-	-	1	-	-	2	2	1	-	1	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	2	2	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1	-	1	1	2	-	1	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	2	-	1	-	1	-	-	-
White, both sexes	1	-	1	-	2	-	1	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	2	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
New Hampshire												
All races, both sexes	-	-	-	-	-	2	-	1	1	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	2	-	1	1	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	1	1	1	2	6	3	1	-	-	-
Male	-	-	1	-	1	1	1	1	-	-	-	-
Female	-	-	-	1	-	1	5	2	1	-	-	-
White, both sexes	-	-	-	1	1	1	6	2	1	-	-	-
Male	-	-	-	-	1	-	1	1	-	-	-	-
Female	-	-	-	1	-	1	5	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
New Mexico												
All races, both sexes	-	-	-	1	-	2	2	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	2	2	1	1	-	-	-
White, both sexes	-	-	-	-	-	2	2	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	2	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	3	1	1	4	4	9	11	9	5	-	-	-
Male	-	1	-	2	-	3	2	1	2	-	-	-
Female	3	-	1	2	4	6	9	8	3	-	-	-
White, both sexes	1	1	-	3	3	9	11	9	5	-	-	-
Male	-	1	-	1	-	3	2	1	2	-	-	-
Female	1	-	-	2	3	6	9	8	3	-	-	-
Black, both sexes	2	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
North Carolina												
All races, both sexes	-	1	3	3	5	4	6	4	1	4	-	-
Male	-	-	-	-	3	1	3	-	-	-	-	-
Female	-	1	3	3	2	3	3	4	1	4	-	-
White, both sexes	-	-	3	2	3	4	6	3	1	2	-	-
Male	-	-	-	-	1	1	3	-	-	-	-	-
Female	-	-	3	2	2	3	3	3	1	2	-	-
Black, both sexes	-	1	-	1	1	-	-	1	-	2	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	1	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	1	-	-	1	-	1	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	1	-	-	1	-	1	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Ohio												
All races, both sexes	2	1	2	3	2	4	11	3	4	-	-	-
Male	-	-	1	2	2	-	4	2	2	-	-	-
Female	2	1	1	1	-	4	7	1	2	-	-	-
White, both sexes	1	-	2	2	2	4	9	3	4	-	-	-
Male	-	-	1	2	2	-	4	2	2	-	-	-
Female	1	-	1	-	-	4	5	1	2	-	-	-
Black, both sexes	1	1	-	1	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	1	-	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	1	1	1	-	3	1	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	1	1	-	-	2	1	-	-	-
White, both sexes	-	-	-	1	1	1	-	3	1	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	1	1	-	-	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Oregon												
All races, both sexes	1	-	1	1	1	-	8	4	1	1	-	-
Male	1	-	-	-	-	-	3	-	1	1	-	-
Female	-	-	1	1	1	-	5	4	-	-	-	-
White, both sexes	1	-	1	-	1	-	8	4	1	1	-	-
Male	1	-	-	-	-	-	3	-	1	1	-	-
Female	-	-	1	-	1	-	5	4	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	2	4	4	6	6	12	16	4	-	1	-
Male	-	-	3	-	2	-	3	3	-	-	-	-
Female	-	2	1	4	4	6	9	13	4	-	1	-
White, both sexes	-	2	4	4	6	6	12	15	3	-	1	-
Male	-	-	3	-	2	-	3	3	-	-	-	-
Female	-	2	1	4	4	6	9	12	3	-	1	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	2	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	2	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	-	3	1	2	2	-	1	1	1	-	-
Male	-	-	2	-	-	1	-	-	1	-	-	-
Female	1	-	1	1	2	1	-	1	-	1	-	-
White, both sexes	-	-	2	-	2	1	-	-	1	1	-	-
Male	-	-	1	-	-	-	-	-	1	-	-	-
Female	-	-	1	-	2	1	-	-	-	1	-	-
Black, both sexes	1	-	1	1	-	1	-	1	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
South Dakota												
All races, both sexes	-	-	-	1	1	-	-	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
White, both sexes	-	-	-	1	1	-	-	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	3	2	-	1	2	4	3	3	3	-	-	-
Male	1	2	-	-	-	1	-	1	-	-	-	-
Female	2	-	-	1	2	3	3	2	3	-	-	-
White, both sexes	2	1	-	1	2	4	2	3	1	-	-	-
Male	-	1	-	-	-	1	-	1	-	-	-	-
Female	2	-	-	1	2	3	2	2	1	-	-	-
Black, both sexes	1	1	-	-	-	-	1	-	2	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Texas												
All races, both sexes	-	-	4	2	3	3	4	5	-	1	-	-
Male	-	-	-	-	-	-	2	1	-	-	-	-
Female	-	-	4	2	3	3	2	4	-	1	-	-
White, both sexes	-	-	4	1	3	2	3	5	-	1	-	-
Male	-	-	-	-	-	-	2	1	-	-	-	-
Female	-	-	4	1	3	2	1	4	-	1	-	-
Black, both sexes	-	-	-	1	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	1	2	2	-	1	3	1	-	-	-
Male	-	-	-	-	-	-	-	2	-	-	-	-
Female	-	-	1	2	2	-	1	1	1	-	-	-
White, both sexes	-	-	1	2	2	-	1	3	1	-	-	-
Male	-	-	-	-	-	-	-	2	-	-	-	-
Female	-	-	1	2	2	-	1	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Vermont												
All races, both sexes	-	-	-	-	-	3	2	2	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	2	1	2	-	-	-	-
White, both sexes	-	-	-	-	-	3	2	2	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	2	1	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1	2	3	1	1	3	1	4	-	1	-	-
Male	1	1	2	-	-	-	-	1	-	-	-	-
Female	-	1	1	1	1	3	1	3	-	1	-	-
White, both sexes	-	-	3	1	1	1	1	4	-	1	-	-
Male	-	-	2	-	-	-	-	1	-	-	-	-
Female	-	-	1	1	1	1	1	3	-	1	-	-
Black, both sexes	1	1	-	-	-	2	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Washington												
All races, both sexes	-	-	1	2	1	4	6	5	2	2	-	-
Male	-	-	-	1	-	1	3	-	2	-	-	-
Female	-	-	1	1	1	3	3	5	-	2	-	-
White, both sexes	-	-	1	2	1	4	6	5	2	2	-	-
Male	-	-	-	1	-	1	3	-	2	-	-	-
Female	-	-	1	1	1	3	3	5	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	1	3	-	-	-	-	-	-
Male	-	-	-	-	1	2	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	3	-	-	-	-	-	-
Male	-	-	-	-	1	2	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Wisconsin												
All races, both sexes	2	1	1	1	-	3	5	3	3	2	-	-
Male	2	-	-	-	-	1	1	-	1	-	-	-
Female	-	1	1	1	-	2	4	3	2	2	-	-
White, both sexes	2	1	1	1	-	2	5	3	3	2	-	-
Male	2	-	-	-	-	1	1	-	1	-	-	-
Female	-	1	1	1	-	1	4	3	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	1	-	-	-	2	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	2	-	1	-	-	-
White, both sexes	-	-	1	-	-	-	2	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	2	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
United States												
All races, both sexes	57,732	2	2	-	-	15	57	154	301	605	1,228	2,068
Male	24,984	2	2	-	-	10	41	101	199	423	836	1,397
Female	32,748	-	-	-	-	5	16	53	102	182	392	671
White, both sexes	43,585	2	2	-	-	7	23	80	130	304	688	1,232
Male	18,179	2	2	-	-	5	18	52	94	235	491	879
Female	25,406	-	-	-	-	2	5	28	36	69	197	353
Black, both sexes	12,573	-	-	-	-	8	34	66	163	285	511	788
Male	6,060	-	-	-	-	5	23	43	99	179	327	477
Female	6,513	-	-	-	-	3	11	23	64	106	184	311
American Indian 1/, both sexes	251	-	-	-	-	-	-	3	2	6	12	16
Male	129	-	-	-	-	-	-	3	2	4	7	12
Female	122	-	-	-	-	-	-	-	-	2	5	4
Asian or Pacific Islander 2/, both sexes	1,323	-	-	-	-	-	-	5	6	10	17	32
Male	616	-	-	-	-	-	-	3	4	5	11	29
Female	707	-	-	-	-	-	-	2	2	5	6	3
Alabama												
All races, both sexes	904	-	-	-	-	-	4	5	5	26	23	43
Male	391	-	-	-	-	-	3	5	2	15	17	29
Female	513	-	-	-	-	-	1	-	3	11	6	14
White, both sexes	530	-	-	-	-	-	1	1	1	7	9	20
Male	220	-	-	-	-	-	1	1	1	7	5	13
Female	310	-	-	-	-	-	-	-	-	-	4	7
Black, both sexes	372	-	-	-	-	-	3	3	4	19	14	23
Male	169	-	-	-	-	-	2	3	1	8	12	16
Female	203	-	-	-	-	-	1	-	3	11	2	7
American Indian 1/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Alaska												
All races, both sexes	78	-	-	-	-	-	-	-	1	-	-	7
Male	47	-	-	-	-	-	-	-	1	-	-	5
Female	31	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	57	-	-	-	-	-	-	-	-	-	-	6
Male	36	-	-	-	-	-	-	-	-	-	-	4
Female	21	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	1	-	-	1
Male	6	-	-	-	-	-	-	-	1	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1,088	-	-	-	-	-	1	2	3	12	19	35
Male	518	-	-	-	-	-	1	1	3	9	11	28
Female	570	-	-	-	-	-	-	1	-	3	8	7
White, both sexes	978	-	-	-	-	-	1	2	3	8	11	31
Male	462	-	-	-	-	-	1	1	3	6	8	25
Female	516	-	-	-	-	-	-	1	-	2	3	6
Black, both sexes	56	-	-	-	-	-	-	-	-	2	3	3
Male	33	-	-	-	-	-	-	-	-	2	1	3
Female	23	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	43	-	-	-	-	-	-	-	-	2	5	1
Male	18	-	-	-	-	-	-	-	-	1	2	-
Female	25	-	-	-	-	-	-	-	-	1	3	1
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Arkansas												
All races, both sexes	553	-	-	-	-	-	1	3	2	6	19	24
Male	215	-	-	-	-	-	-	1	-	5	14	17
Female	338	-	-	-	-	-	1	2	2	1	5	7
White, both sexes	429	-	-	-	-	-	1	1	-	4	10	15
Male	162	-	-	-	-	-	-	-	-	3	7	11
Female	267	-	-	-	-	-	1	1	-	1	3	4
Black, both sexes	124	-	-	-	-	-	-	2	2	2	9	9
Male	53	-	-	-	-	-	-	1	-	2	7	6
Female	71	-	-	-	-	-	-	1	2	-	2	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	7,620	-	-	-	-	1	1	18	30	66	137	235
Male	3,329	-	-	-	-	1	1	13	18	45	95	168
Female	4,291	-	-	-	-	-	-	5	12	21	42	67
White, both sexes	5,900	-	-	-	-	1	1	11	19	36	95	155
Male	2,515	-	-	-	-	1	1	8	13	26	68	111
Female	3,385	-	-	-	-	-	-	3	6	10	27	44
Black, both sexes	1,017	-	-	-	-	-	-	3	8	23	32	67
Male	488	-	-	-	-	-	-	2	3	17	21	47
Female	529	-	-	-	-	-	-	1	5	6	11	20
American Indian 1/, both sexes	36	-	-	-	-	-	-	1	-	-	1	-
Male	20	-	-	-	-	-	-	1	-	-	1	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	667	-	-	-	-	-	-	3	3	7	9	13
Male	306	-	-	-	-	-	-	2	2	2	5	10
Female	361	-	-	-	-	-	-	1	1	5	4	3

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Colorado												
All races, both sexes	522	-	-	-	-	-	-	2	1	4	7	15
Male	238	-	-	-	-	-	-	2	1	4	5	14
Female	284	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	487	-	-	-	-	-	-	1	1	4	7	13
Male	221	-	-	-	-	-	-	1	1	4	5	13
Female	266	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	30	-	-	-	-	-	-	1	-	-	-	2
Male	16	-	-	-	-	-	-	1	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	550	-	-	-	-	-	2	-	-	1	6	15
Male	210	-	-	-	-	-	1	-	-	1	5	8
Female	340	-	-	-	-	-	1	-	-	-	1	7
White, both sexes	484	-	-	-	-	-	1	-	-	1	4	12
Male	175	-	-	-	-	-	1	-	-	1	3	7
Female	309	-	-	-	-	-	-	-	-	-	1	5
Black, both sexes	64	-	-	-	-	-	1	-	-	-	2	3
Male	33	-	-	-	-	-	-	-	-	-	2	1
Female	31	-	-	-	-	-	1	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Delaware												
All races, both sexes	133	-	-	-	-	-	-	1	1	5	4	8
Male	64	-	-	-	-	-	-	-	-	4	2	5
Female	69	-	-	-	-	-	-	1	1	1	2	3
White, both sexes	97	-	-	-	-	-	-	-	-	3	3	6
Male	41	-	-	-	-	-	-	-	-	2	1	4
Female	56	-	-	-	-	-	-	-	-	1	2	2
Black, both sexes	34	-	-	-	-	-	-	1	-	1	1	2
Male	22	-	-	-	-	-	-	-	-	1	1	1
Female	12	-	-	-	-	-	-	1	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Dist. of Columbia												
All races, both sexes	222	-	-	-	-	-	-	2	4	2	11	9
Male	104	-	-	-	-	-	-	1	2	1	6	6
Female	118	-	-	-	-	-	-	1	2	1	5	3
White, both sexes	30	-	-	-	-	-	-	2	-	-	-	-
Male	8	-	-	-	-	-	-	1	-	-	-	-
Female	22	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	188	-	-	-	-	-	-	-	4	2	11	9
Male	95	-	-	-	-	-	-	-	2	1	6	6
Female	93	-	-	-	-	-	-	-	2	1	5	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Florida												
All races, both sexes	4,247	-	-	-	-	2	7	11	25	32	112	143
Male	2,006	-	-	-	-	1	5	8	20	24	76	99
Female	2,241	-	-	-	-	1	2	3	5	8	36	44
White, both sexes	3,372	-	-	-	-	1	4	10	12	17	65	84
Male	1,590	-	-	-	-	1	3	8	10	13	49	62
Female	1,782	-	-	-	-	-	1	2	2	4	16	22
Black, both sexes	840	-	-	-	-	1	3	1	13	15	45	59
Male	396	-	-	-	-	-	2	-	10	11	25	37
Female	444	-	-	-	-	1	1	1	3	4	20	22
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	32	-	-	-	-	-	-	-	-	-	1	-
Male	17	-	-	-	-	-	-	-	-	-	1	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	2,223	-	-	-	-	1	5	4	12	38	71	111
Male	1,045	-	-	-	-	-	4	4	7	27	45	75
Female	1,178	-	-	-	-	1	1	-	5	11	26	36
White, both sexes	1,159	-	-	-	-	-	2	2	4	11	27	55
Male	521	-	-	-	-	-	2	2	3	8	15	42
Female	638	-	-	-	-	-	-	-	1	3	12	13
Black, both sexes	1,051	-	-	-	-	1	3	2	8	27	44	55
Male	518	-	-	-	-	-	2	2	4	19	30	32
Female	533	-	-	-	-	1	1	-	4	8	14	23
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Hawaii												
All races, both sexes	174	-	-	-	-	-	-	1	-	-	-	3
Male	76	-	-	-	-	-	-	1	-	-	-	2
Female	98	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	47	-	-	-	-	-	-	1	-	-	-	2
Male	19	-	-	-	-	-	-	1	-	-	-	1
Female	28	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	126	-	-	-	-	-	-	-	-	-	-	1
Male	56	-	-	-	-	-	-	-	-	-	-	1
Female	70	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	190	-	-	-	-	-	-	1	1	4	4	5
Male	76	-	-	-	-	-	-	1	-	2	3	4
Female	114	-	-	-	-	-	-	-	1	2	1	1
White, both sexes	189	-	-	-	-	-	-	1	1	3	4	5
Male	75	-	-	-	-	-	-	1	-	1	3	4
Female	114	-	-	-	-	-	-	-	1	2	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Illinois												
All races, both sexes	2,358	-	-	-	-	-	4	7	19	33	55	109
Male	1,053	-	-	-	-	-	3	6	14	23	30	79
Female	1,305	-	-	-	-	-	1	1	5	10	25	30
White, both sexes	1,632	-	-	-	-	-	3	2	4	11	24	57
Male	671	-	-	-	-	-	3	1	4	8	13	47
Female	961	-	-	-	-	-	-	1	-	3	11	10
Black, both sexes	687	-	-	-	-	-	1	5	14	22	31	52
Male	363	-	-	-	-	-	-	5	9	15	17	32
Female	324	-	-	-	-	-	1	-	5	7	14	20
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	38	-	-	-	-	-	-	-	1	-	-	-
Male	18	-	-	-	-	-	-	-	1	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1,011	-	-	-	-	-	-	4	5	5	19	28
Male	406	-	-	-	-	-	-	1	3	2	13	23
Female	605	-	-	-	-	-	-	3	2	3	6	5
White, both sexes	866	-	-	-	-	-	-	2	1	4	11	17
Male	338	-	-	-	-	-	-	-	1	2	8	14
Female	528	-	-	-	-	-	-	2	-	2	3	3
Black, both sexes	141	-	-	-	-	-	-	2	4	1	8	10
Male	67	-	-	-	-	-	-	1	2	-	5	8
Female	74	-	-	-	-	-	-	1	2	1	3	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Iowa												
All races, both sexes	526	-	-	-	-	-	1	-	2	2	4	5
Male	181	-	-	-	-	-	1	-	2	2	3	4
Female	345	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	509	-	-	-	-	-	1	-	1	1	4	4
Male	172	-	-	-	-	-	1	-	1	1	3	4
Female	337	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	16	-	-	-	-	-	-	-	1	1	-	1
Male	8	-	-	-	-	-	-	-	1	1	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	331	-	-	-	-	-	-	-	-	6	5	6
Male	104	-	-	-	-	-	-	-	-	3	3	4
Female	227	-	-	-	-	-	-	-	-	3	2	2
White, both sexes	295	-	-	-	-	-	-	-	-	2	1	6
Male	93	-	-	-	-	-	-	-	-	1	1	4
Female	202	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	34	-	-	-	-	-	-	-	-	4	4	-
Male	11	-	-	-	-	-	-	-	-	2	2	-
Female	23	-	-	-	-	-	-	-	-	2	2	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Kentucky												
All races, both sexes	662	-	-	-	-	-	1	5	7	9	25	36
Male	278	-	-	-	-	-	1	2	3	5	20	16
Female	384	-	-	-	-	-	-	3	4	4	5	20
White, both sexes	580	-	-	-	-	-	-	3	4	7	22	29
Male	246	-	-	-	-	-	-	2	1	4	18	14
Female	334	-	-	-	-	-	-	1	3	3	4	15
Black, both sexes	80	-	-	-	-	-	1	2	3	2	3	7
Male	31	-	-	-	-	-	1	-	2	1	2	2
Female	49	-	-	-	-	-	-	2	1	1	1	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1,369	-	-	-	-	-	1	5	8	24	37	62
Male	617	-	-	-	-	-	1	5	4	16	26	41
Female	752	-	-	-	-	-	-	-	4	8	11	21
White, both sexes	794	-	-	-	-	-	1	-	2	9	15	29
Male	356	-	-	-	-	-	1	-	2	7	11	22
Female	438	-	-	-	-	-	-	-	-	2	4	7
Black, both sexes	570	-	-	-	-	-	-	5	6	15	22	33
Male	257	-	-	-	-	-	-	5	2	9	15	19
Female	313	-	-	-	-	-	-	-	4	6	7	14
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Maine												
All races, both sexes	145	-	-	-	-	-	-	-	-	-	3	2
Male	63	-	-	-	-	-	-	-	-	-	3	1
Female	82	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	143	-	-	-	-	-	-	-	-	-	3	1
Male	62	-	-	-	-	-	-	-	-	-	3	1
Female	81	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1,313	-	-	-	-	-	2	2	6	21	40	100
Male	656	-	-	-	-	-	1	2	6	14	26	70
Female	657	-	-	-	-	-	1	-	-	7	14	30
White, both sexes	769	-	-	-	-	-	-	1	2	10	16	48
Male	365	-	-	-	-	-	-	1	2	9	11	35
Female	404	-	-	-	-	-	-	-	-	1	5	13
Black, both sexes	520	-	-	-	-	-	2	1	4	11	24	51
Male	279	-	-	-	-	-	1	1	4	5	15	34
Female	241	-	-	-	-	-	1	-	-	6	9	17
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	23	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Massachusetts												
All races, both sexes	1,004	-	-	-	-	-	-	6	13	13	27	55
Male	433	-	-	-	-	-	-	3	9	11	24	37
Female	571	-	-	-	-	-	-	3	4	2	3	18
White, both sexes	915	-	-	-	-	-	-	3	9	10	23	45
Male	383	-	-	-	-	-	-	1	6	8	20	31
Female	532	-	-	-	-	-	-	2	3	2	3	14
Black, both sexes	67	-	-	-	-	-	-	1	4	2	4	7
Male	37	-	-	-	-	-	-	1	3	2	4	3
Female	30	-	-	-	-	-	-	-	1	-	-	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	22	-	-	-	-	-	-	2	-	1	-	3
Male	13	-	-	-	-	-	-	1	-	1	-	3
Female	9	-	-	-	-	-	-	1	-	-	-	-
Michigan												
All races, both sexes	2,269	-	-	-	-	-	-	9	13	24	42	70
Male	915	-	-	-	-	-	-	6	10	19	30	42
Female	1,354	-	-	-	-	-	-	3	3	5	12	28
White, both sexes	1,737	-	-	-	-	-	-	5	6	11	18	40
Male	659	-	-	-	-	-	-	4	6	9	10	25
Female	1,078	-	-	-	-	-	-	1	-	2	8	15
Black, both sexes	515	-	-	-	-	-	-	4	6	12	24	29
Male	246	-	-	-	-	-	-	2	3	9	20	16
Female	269	-	-	-	-	-	-	2	3	3	4	13
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	1	1	-	-
Male	6	-	-	-	-	-	-	-	1	1	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Minnesota												
All races, both sexes	623	-	-	-	-	-	1	1	-	3	4	7
Male	258	-	-	-	-	-	1	1	-	3	4	5
Female	365	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	580	-	-	-	-	-	-	-	-	2	2	6
Male	229	-	-	-	-	-	-	-	-	2	2	5
Female	351	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	30	-	-	-	-	-	1	-	-	1	2	-
Male	21	-	-	-	-	-	1	-	-	1	2	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	1	-	-	-	1
Male	4	-	-	-	-	-	-	1	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1,238	-	-	-	-	2	4	4	11	13	29	50
Male	540	-	-	-	-	2	3	3	6	9	18	31
Female	698	-	-	-	-	-	1	1	5	4	11	19
White, both sexes	653	-	-	-	-	1	-	-	4	2	5	20
Male	267	-	-	-	-	1	-	-	1	1	2	12
Female	386	-	-	-	-	-	-	-	3	1	3	8
Black, both sexes	578	-	-	-	-	1	4	4	7	11	22	30
Male	271	-	-	-	-	1	3	3	5	8	16	19
Female	307	-	-	-	-	-	1	1	2	3	6	11
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Missouri												
All races, both sexes	877	-	-	-	-	1	-	1	9	9	17	32
Male	346	-	-	-	-	1	-	1	5	7	8	16
Female	531	-	-	-	-	-	-	-	4	2	9	16
White, both sexes	636	-	-	-	-	1	-	-	2	5	7	10
Male	232	-	-	-	-	1	-	-	1	4	3	5
Female	404	-	-	-	-	-	-	-	1	1	4	5
Black, both sexes	232	-	-	-	-	-	-	1	7	4	9	22
Male	110	-	-	-	-	-	-	1	4	3	4	11
Female	122	-	-	-	-	-	-	-	3	1	5	11
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	132	-	-	-	-	-	-	-	-	1	-	5
Male	53	-	-	-	-	-	-	-	-	1	-	5
Female	79	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	131	-	-	-	-	-	-	-	-	1	-	5
Male	53	-	-	-	-	-	-	-	-	1	-	5
Female	78	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Nebraska												
All races, both sexes	334	-	-	-	-	-	-	-	-	-	11	9
Male	117	-	-	-	-	-	-	-	-	-	7	6
Female	217	-	-	-	-	-	-	-	-	-	4	3
White, both sexes	314	-	-	-	-	-	-	-	-	-	9	5
Male	107	-	-	-	-	-	-	-	-	-	7	3
Female	207	-	-	-	-	-	-	-	-	-	2	2
Black, both sexes	19	-	-	-	-	-	-	-	-	-	2	4
Male	10	-	-	-	-	-	-	-	-	-	-	3
Female	9	-	-	-	-	-	-	-	-	-	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	454	-	-	-	-	-	1	2	3	5	10	17
Male	236	-	-	-	-	-	-	2	3	3	7	9
Female	218	-	-	-	-	-	1	-	-	2	3	8
White, both sexes	384	-	-	-	-	-	-	2	3	3	4	12
Male	204	-	-	-	-	-	-	2	3	3	4	7
Female	180	-	-	-	-	-	-	-	-	-	-	5
Black, both sexes	57	-	-	-	-	-	1	-	-	2	5	5
Male	28	-	-	-	-	-	-	-	-	-	3	2
Female	29	-	-	-	-	-	1	-	-	2	2	3
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
New Hampshire												
All races, both sexes	187	-	-	-	-	-	-	-	-	1	3	8
Male	83	-	-	-	-	-	-	-	-	1	1	5
Female	104	-	-	-	-	-	-	-	-	-	2	3
White, both sexes	186	-	-	-	-	-	-	-	-	1	3	7
Male	82	-	-	-	-	-	-	-	-	1	1	4
Female	104	-	-	-	-	-	-	-	-	-	2	3
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1,418	-	-	-	-	-	1	-	5	12	13	44
Male	589	-	-	-	-	-	1	-	3	8	8	29
Female	829	-	-	-	-	-	-	-	2	4	5	15
White, both sexes	1,067	-	-	-	-	-	-	-	1	4	5	21
Male	418	-	-	-	-	-	-	-	-	4	4	15
Female	649	-	-	-	-	-	-	-	1	-	1	6
Black, both sexes	320	-	-	-	-	-	1	-	3	8	8	21
Male	159	-	-	-	-	-	1	-	2	4	4	12
Female	161	-	-	-	-	-	-	-	1	4	4	9
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	29	-	-	-	-	-	-	-	1	-	-	1
Male	10	-	-	-	-	-	-	-	1	-	-	1
Female	19	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
New Mexico												
All races, both sexes	259	-	-	-	-	-	-	-	-	4	5	3
Male	111	-	-	-	-	-	-	-	-	2	5	2
Female	148	-	-	-	-	-	-	-	-	2	-	1
White, both sexes	235	-	-	-	-	-	-	-	-	4	5	3
Male	96	-	-	-	-	-	-	-	-	2	5	2
Female	139	-	-	-	-	-	-	-	-	2	-	1
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	4,519	-	1	-	-	1	7	10	20	52	109	150
Male	2,052	-	1	-	-	-	5	7	13	32	82	114
Female	2,467	-	-	-	-	1	2	3	7	20	27	36
White, both sexes	3,159	-	1	-	-	-	2	5	7	25	54	81
Male	1,379	-	1	-	-	-	1	3	4	18	47	66
Female	1,780	-	-	-	-	-	1	2	3	7	7	15
Black, both sexes	1,241	-	-	-	-	1	5	5	13	26	54	67
Male	613	-	-	-	-	-	4	4	9	13	34	46
Female	628	-	-	-	-	1	1	1	4	13	20	21
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	117	-	-	-	-	-	-	-	-	1	1	2
Male	59	-	-	-	-	-	-	-	-	1	1	2
Female	58	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
North Carolina												
All races, both sexes	1,426	-	-	-	-	2	1	5	10	17	38	60
Male	603	-	-	-	-	2	-	2	10	10	17	37
Female	823	-	-	-	-	-	1	3	-	7	21	23
White, both sexes	819	-	-	-	-	1	-	1	5	8	15	26
Male	325	-	-	-	-	1	-	1	5	5	9	17
Female	494	-	-	-	-	-	-	-	-	3	6	9
Black, both sexes	584	-	-	-	-	1	1	4	5	9	22	33
Male	268	-	-	-	-	1	-	1	5	5	7	20
Female	316	-	-	-	-	-	1	3	-	4	15	13
American Indian 1/, both sexes	16	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	132	-	-	-	-	-	-	-	-	-	1	-
Male	40	-	-	-	-	-	-	-	-	-	-	-
Female	92	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	127	-	-	-	-	-	-	-	-	-	1	-
Male	39	-	-	-	-	-	-	-	-	-	-	-
Female	88	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Ohio												
All races, both sexes	3,145	-	-	-	-	-	3	8	12	27	51	117
Male	1,296	-	-	-	-	-	2	4	9	19	34	81
Female	1,849	-	-	-	-	-	1	4	3	8	17	36
White, both sexes	2,513	-	-	-	-	-	-	6	7	15	30	86
Male	987	-	-	-	-	-	-	3	6	11	20	62
Female	1,526	-	-	-	-	-	-	3	1	4	10	24
Black, both sexes	613	-	-	-	-	-	3	2	5	12	20	29
Male	296	-	-	-	-	-	2	1	3	8	13	17
Female	317	-	-	-	-	-	1	1	2	4	7	12
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	18	-	-	-	-	-	-	-	-	-	1	2
Male	12	-	-	-	-	-	-	-	-	-	1	2
Female	6	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	582	-	-	-	-	1	-	-	1	5	11	18
Male	221	-	-	-	-	1	-	-	-	5	5	14
Female	361	-	-	-	-	-	-	-	1	-	6	4
White, both sexes	476	-	-	-	-	-	-	-	-	5	8	13
Male	180	-	-	-	-	-	-	-	-	5	3	11
Female	296	-	-	-	-	-	-	-	-	-	5	2
Black, both sexes	71	-	-	-	-	1	-	-	1	-	2	3
Male	22	-	-	-	-	1	-	-	-	-	1	1
Female	49	-	-	-	-	-	-	-	1	-	1	2
American Indian 1/, both sexes	31	-	-	-	-	-	-	-	-	-	1	2
Male	19	-	-	-	-	-	-	-	-	-	1	2
Female	12	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Oregon												
All races, both sexes	629	-	-	-	-	-	-	-	1	6	4	15
Male	243	-	-	-	-	-	-	-	1	4	4	12
Female	386	-	-	-	-	-	-	-	-	2	-	3
White, both sexes	594	-	-	-	-	-	-	-	1	4	3	12
Male	219	-	-	-	-	-	-	-	1	2	3	9
Female	375	-	-	-	-	-	-	-	-	2	-	3
Black, both sexes	15	-	-	-	-	-	-	-	-	1	1	-
Male	10	-	-	-	-	-	-	-	-	1	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	-	1	-	2
Male	10	-	-	-	-	-	-	-	-	1	-	2
Female	3	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1,914	-	-	-	-	-	1	1	1	11	31	46
Male	785	-	-	-	-	-	1	-	-	7	23	33
Female	1,129	-	-	-	-	-	-	1	1	4	8	13
White, both sexes	1,606	-	-	-	-	-	1	1	1	3	25	27
Male	642	-	-	-	-	-	1	-	-	3	18	20
Female	964	-	-	-	-	-	-	1	1	-	7	7
Black, both sexes	298	-	-	-	-	-	-	-	-	8	6	19
Male	139	-	-	-	-	-	-	-	-	4	5	13
Female	159	-	-	-	-	-	-	-	-	4	1	6
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Rhode Island												
All races, both sexes	130	-	-	-	-	-	-	-	2	-	9	11
Male	50	-	-	-	-	-	-	-	-	-	6	8
Female	80	-	-	-	-	-	-	-	2	-	3	3
White, both sexes	122	-	-	-	-	-	-	-	2	-	8	9
Male	46	-	-	-	-	-	-	-	-	-	5	6
Female	76	-	-	-	-	-	-	-	2	-	3	3
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	1	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	838	-	-	-	-	1	-	3	6	6	8	19
Male	364	-	-	-	-	-	-	1	3	3	6	9
Female	474	-	-	-	-	1	-	2	3	3	2	10
White, both sexes	442	-	-	-	-	1	-	1	-	1	1	5
Male	178	-	-	-	-	-	-	1	-	-	1	4
Female	264	-	-	-	-	1	-	-	-	1	-	1
Black, both sexes	395	-	-	-	-	-	-	2	6	5	7	14
Male	185	-	-	-	-	-	-	-	3	3	5	5
Female	210	-	-	-	-	-	-	2	3	2	2	9
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
South Dakota												
All races, both sexes	127	-	-	-	-	-	-	-	-	-	-	-
Male	41	-	-	-	-	-	-	-	-	-	-	-
Female	86	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	122	-	-	-	-	-	-	-	-	-	-	-
Male	39	-	-	-	-	-	-	-	-	-	-	-
Female	83	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	1,511	-	-	-	-	-	1	8	16	22	33	58
Male	648	-	-	-	-	-	-	4	11	15	23	42
Female	863	-	-	-	-	-	1	4	5	7	10	16
White, both sexes	1,068	-	-	-	-	-	1	1	7	10	19	31
Male	431	-	-	-	-	-	-	1	5	8	15	21
Female	637	-	-	-	-	-	1	-	2	2	4	10
Black, both sexes	435	-	-	-	-	-	-	7	9	12	14	27
Male	212	-	-	-	-	-	-	3	6	7	8	21
Female	223	-	-	-	-	-	-	4	3	5	6	6
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Texas												
All races, both sexes	3,664	1	1	-	-	1	6	17	32	47	94	159
Male	1,575	1	1	-	-	-	5	9	21	35	67	93
Female	2,089	-	-	-	-	1	1	8	11	12	27	66
White, both sexes	2,886	1	1	-	-	1	3	12	15	32	65	106
Male	1,230	1	1	-	-	-	2	5	10	25	45	63
Female	1,656	-	-	-	-	1	1	7	5	7	20	43
Black, both sexes	723	-	-	-	-	-	3	5	17	15	29	49
Male	322	-	-	-	-	-	3	4	11	10	22	26
Female	401	-	-	-	-	-	-	1	6	5	7	23
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	3
Male	6	-	-	-	-	-	-	-	-	-	-	3
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	46	-	-	-	-	-	-	-	-	-	-	1
Male	17	-	-	-	-	-	-	-	-	-	-	1
Female	29	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	245	-	-	-	-	-	-	-	-	2	3	6
Male	99	-	-	-	-	-	-	-	-	2	3	3
Female	146	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	233	-	-	-	-	-	-	-	-	2	2	6
Male	95	-	-	-	-	-	-	-	-	2	2	3
Female	138	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Vermont												
All races, both sexes	104	-	-	-	-	-	-	-	-	-	1	3
Male	42	-	-	-	-	-	-	-	-	-	1	3
Female	62	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	103	-	-	-	-	-	-	-	-	-	1	3
Male	42	-	-	-	-	-	-	-	-	-	1	3
Female	61	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	990	1	-	-	-	1	-	4	8	19	35	43
Male	422	1	-	-	-	1	-	3	4	17	25	23
Female	568	-	-	-	-	-	-	1	4	2	10	20
White, both sexes	617	1	-	-	-	-	-	1	1	12	17	21
Male	252	1	-	-	-	-	-	1	1	12	13	14
Female	365	-	-	-	-	-	-	-	-	-	4	7
Black, both sexes	354	-	-	-	-	1	-	3	7	7	17	22
Male	163	-	-	-	-	1	-	2	3	5	11	9
Female	191	-	-	-	-	-	-	1	4	2	6	13
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	18	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Washington												
All races, both sexes	1,462	-	-	-	-	-	-	1	2	4	21	26
Male	626	-	-	-	-	-	-	1	2	3	14	18
Female	836	-	-	-	-	-	-	-	-	1	7	8
White, both sexes	1,328	-	-	-	-	-	-	1	-	3	16	15
Male	560	-	-	-	-	-	-	1	-	3	11	11
Female	768	-	-	-	-	-	-	-	-	-	5	4
Black, both sexes	62	-	-	-	-	-	-	-	2	-	3	6
Male	32	-	-	-	-	-	-	-	2	-	2	2
Female	30	-	-	-	-	-	-	-	-	-	1	4
American Indian 1/, both sexes	17	-	-	-	-	-	-	-	-	1	2	2
Male	9	-	-	-	-	-	-	-	-	-	1	2
Female	8	-	-	-	-	-	-	-	-	1	1	-
Asian or Pacific Islander 2/, both sexes	55	-	-	-	-	-	-	-	-	-	-	3
Male	25	-	-	-	-	-	-	-	-	-	-	3
Female	30	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	372	-	-	-	-	-	-	1	2	3	3	8
Male	152	-	-	-	-	-	-	1	2	2	2	4
Female	220	-	-	-	-	-	-	-	-	1	1	4
White, both sexes	352	-	-	-	-	-	-	1	2	1	3	7
Male	144	-	-	-	-	-	-	1	2	1	2	4
Female	208	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	20	-	-	-	-	-	-	-	-	2	-	1
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	12	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I15)												
Wisconsin												
All races, both sexes	871	-	-	-	-	1	1	-	2	3	13	25
Male	337	-	-	-	-	1	1	-	1	3	9	16
Female	534	-	-	-	-	-	-	-	1	-	4	9
White, both sexes	776	-	-	-	-	-	-	-	2	2	7	12
Male	283	-	-	-	-	-	-	-	1	2	6	11
Female	493	-	-	-	-	-	-	-	1	-	1	1
Black, both sexes	89	-	-	-	-	1	1	-	-	1	6	13
Male	52	-	-	-	-	1	1	-	-	1	3	5
Female	37	-	-	-	-	-	-	-	-	-	3	8
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	57	-	-	-	-	-	-	-	-	-	1	3
Male	30	-	-	-	-	-	-	-	-	-	-	2
Female	27	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	57	-	-	-	-	-	-	-	-	-	1	3
Male	30	-	-	-	-	-	-	-	-	-	-	2
Female	27	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
United States												
All races, both sexes	2,949	3,364	3,540	3,457	4,064	5,504	8,054	9,654	8,024	3,778	909	3
Male	1,956	2,198	2,262	2,015	2,043	2,590	3,094	3,015	1,987	706	105	2
Female	993	1,166	1,278	1,442	2,021	2,914	4,960	6,639	6,037	3,072	804	1
White, both sexes	1,761	2,069	2,284	2,326	2,850	4,119	6,497	8,178	6,994	3,270	768	1
Male	1,231	1,433	1,483	1,378	1,450	1,973	2,493	2,550	1,727	596	86	1
Female	530	636	801	948	1,400	2,146	4,004	5,628	5,267	2,674	682	-
Black, both sexes	1,122	1,212	1,160	1,022	1,092	1,220	1,289	1,233	822	421	123	2
Male	683	714	712	570	536	544	494	365	197	79	12	1
Female	439	498	448	452	556	676	795	868	625	342	111	1
American Indian 1/, both sexes	23	20	23	25	15	30	30	27	13	5	1	-
Male	15	12	15	15	8	14	8	11	1	1	1	-
Female	8	8	8	10	7	16	22	16	12	4	-	-
Asian or Pacific Islander 2/, both sexes	43	63	73	84	107	135	238	216	195	82	17	-
Male	27	39	52	52	49	59	99	89	62	30	6	-
Female	16	24	21	32	58	76	139	127	133	52	11	-
Alabama												
All races, both sexes	49	45	64	73	75	94	115	124	106	39	14	-
Male	31	31	40	40	51	35	34	35	19	3	1	-
Female	18	14	24	33	24	59	81	89	87	36	13	-
White, both sexes	27	21	25	34	48	56	80	85	79	27	9	-
Male	18	18	18	19	34	22	23	23	14	2	1	-
Female	9	3	7	15	14	34	57	62	65	25	8	-
Black, both sexes	22	23	39	39	27	38	35	39	27	12	5	-
Male	13	12	22	21	17	13	11	12	5	1	-	-
Female	9	11	17	18	10	25	24	27	22	11	5	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Alaska												
All races, both sexes	8	12	10	7	7	7	3	9	3	3	1	-
Male	5	8	5	6	6	4	-	2	3	1	1	-
Female	3	4	5	1	1	3	3	7	-	2	-	-
White, both sexes	5	11	8	6	7	1	2	6	3	2	-	-
Male	3	7	5	6	6	-	-	2	3	-	-	-
Female	2	4	3	-	1	1	2	4	-	2	-	-
Black, both sexes	-	1	1	-	-	2	-	2	-	-	-	-
Male	-	1	-	-	-	2	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	2	-	-	-	-
American Indian 1/, both sexes	3	-	1	1	-	3	1	1	-	-	1	-
Male	2	-	-	-	-	1	-	-	-	-	1	-
Female	1	-	1	1	-	2	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	73	73	61	73	88	105	151	195	128	57	12	-
Male	41	54	35	42	54	51	70	64	39	14	1	-
Female	32	19	26	31	34	54	81	131	89	43	11	-
White, both sexes	60	57	55	63	74	97	144	182	122	56	12	-
Male	34	41	30	36	49	47	67	60	39	14	1	-
Female	26	16	25	27	25	50	77	122	83	42	11	-
Black, both sexes	7	8	4	5	9	5	1	6	3	-	-	-
Male	6	8	3	2	4	2	-	2	-	-	-	-
Female	1	-	1	3	5	3	1	4	3	-	-	-
American Indian 1/, both sexes	6	6	1	4	4	3	3	7	-	1	-	-
Male	1	4	1	3	1	2	1	2	-	-	-	-
Female	5	2	-	1	3	1	2	5	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	2	1	1	1	-	3	-	3	-	-	-
Male	-	1	1	1	-	-	2	-	-	-	-	-
Female	-	1	-	-	1	-	1	-	3	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Arkansas												
All races, both sexes	27	39	32	33	30	58	65	92	86	28	8	-
Male	13	24	20	18	13	25	27	18	17	3	-	-
Female	14	15	12	15	17	33	38	74	69	25	8	-
White, both sexes	16	26	26	25	22	43	54	76	80	24	6	-
Male	9	17	17	13	10	22	21	15	14	3	-	-
Female	7	9	9	12	12	21	33	61	66	21	6	-
Black, both sexes	11	13	6	8	8	15	11	16	6	4	2	-
Male	4	7	3	5	3	3	6	3	3	-	-	-
Female	7	6	3	3	5	12	5	13	3	4	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	374	388	436	417	484	699	1,191	1,356	1,129	553	104	1
Male	252	242	284	235	258	350	466	457	315	117	11	1
Female	122	146	152	182	226	349	725	899	814	436	93	-
White, both sexes	251	257	308	311	351	523	955	1,128	943	470	84	1
Male	176	178	196	170	192	270	375	359	266	95	9	1
Female	75	79	112	141	159	253	580	769	677	375	75	-
Black, both sexes	100	101	91	63	81	99	113	114	76	36	10	-
Male	59	48	58	36	43	47	39	44	17	7	-	-
Female	41	53	33	27	38	52	74	70	59	29	10	-
American Indian 1/, both sexes	1	5	6	3	2	3	1	6	6	1	-	-
Male	1	3	4	3	2	1	-	3	1	-	-	-
Female	-	2	2	-	-	2	1	3	5	1	-	-
Asian or Pacific Islander 2/, both sexes	22	25	31	40	50	74	122	108	104	46	10	-
Male	16	13	26	26	21	32	52	51	31	15	2	-
Female	6	12	5	14	29	42	70	57	73	31	8	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Colorado												
All races, both sexes	22	30	39	35	29	56	64	91	83	36	8	-
Male	18	15	24	17	19	31	23	34	20	9	2	-
Female	4	15	15	18	10	25	41	57	63	27	6	-
White, both sexes	19	27	38	30	27	53	58	85	81	35	8	-
Male	16	13	24	15	17	29	21	32	19	9	2	-
Female	3	14	14	15	10	24	37	53	62	26	6	-
Black, both sexes	3	3	1	5	2	3	2	5	2	1	-	-
Male	2	2	-	2	2	2	2	1	1	-	-	-
Female	1	1	1	3	-	1	-	4	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	3	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	3	-	-	-	-	-
Connecticut												
All races, both sexes	20	31	28	25	31	62	68	97	96	53	15	-
Male	15	21	21	15	9	34	20	28	21	8	3	-
Female	5	10	7	10	22	28	48	69	75	45	12	-
White, both sexes	16	20	21	18	27	58	62	88	94	47	15	-
Male	11	14	16	10	7	31	18	26	19	8	3	-
Female	5	6	5	8	20	27	44	62	75	39	12	-
Black, both sexes	4	11	6	7	4	4	6	9	1	6	-	-
Male	4	7	4	5	2	3	2	2	1	-	-	-
Female	-	4	2	2	2	1	4	7	-	6	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Delaware												
All races, both sexes	10	7	3	10	12	11	14	22	16	6	3	-
Male	7	4	3	9	8	6	5	5	5	1	-	-
Female	3	3	-	1	4	5	9	17	11	5	3	-
White, both sexes	4	6	2	4	8	7	13	20	15	3	3	-
Male	2	3	2	3	5	4	5	4	5	1	-	-
Female	2	3	-	1	3	3	8	16	10	2	3	-
Black, both sexes	6	1	1	6	4	4	1	2	1	3	-	-
Male	5	1	1	6	3	2	-	1	-	-	-	-
Female	1	-	-	-	1	2	1	1	1	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	18	17	13	16	17	19	25	28	16	19	6	-
Male	13	7	9	11	9	8	12	9	5	5	-	-
Female	5	10	4	5	8	11	13	19	11	14	6	-
White, both sexes	1	1	-	-	1	3	3	7	4	7	1	-
Male	1	-	-	-	1	1	2	2	-	-	-	-
Female	-	1	-	-	-	2	1	5	4	7	1	-
Black, both sexes	17	16	13	16	15	16	21	20	11	12	5	-
Male	12	7	9	11	7	7	10	7	5	5	-	-
Female	5	9	4	5	8	9	11	13	6	7	5	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	1	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Florida												
All races, both sexes	217	235	230	261	350	433	602	734	538	250	65	-
Male	141	148	148	160	175	223	265	264	183	59	7	-
Female	76	87	82	101	175	210	337	470	355	191	58	-
White, both sexes	142	150	166	183	271	345	516	646	483	222	55	-
Male	100	100	110	115	135	186	228	243	166	55	6	-
Female	42	50	56	68	136	159	288	403	317	167	49	-
Black, both sexes	74	83	61	73	74	83	84	83	51	27	10	-
Male	40	46	37	42	37	34	35	19	16	4	1	-
Female	34	37	24	31	37	49	49	64	35	23	9	-
American Indian 1/, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	3	5	5	4	2	5	4	1	-	-
Male	1	1	1	3	3	2	2	2	1	-	-	-
Female	-	-	2	2	2	2	-	3	3	1	-	-
Georgia												
All races, both sexes	168	224	201	168	167	194	232	281	207	109	29	1
Male	115	146	124	104	80	91	76	77	44	24	2	-
Female	53	78	77	64	87	103	156	204	163	85	27	1
White, both sexes	62	100	94	69	91	103	135	191	137	58	18	-
Male	44	70	61	45	44	50	44	51	27	12	1	-
Female	18	30	33	24	47	53	91	140	110	46	17	-
Black, both sexes	106	124	106	96	76	89	92	89	70	51	11	1
Male	71	76	63	57	36	41	29	26	17	12	1	-
Female	35	48	43	39	40	48	63	63	53	39	10	1
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	3	-	1	5	1	-	-	-	-
Male	-	-	-	2	-	-	3	-	-	-	-	-
Female	-	-	1	1	-	1	2	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Hawaii												
All races, both sexes	4	4	5	7	13	12	27	39	38	18	3	-
Male	2	2	5	6	5	7	9	15	15	6	1	-
Female	2	2	-	1	8	5	18	24	23	12	2	-
White, both sexes	-	-	1	3	6	4	5	9	11	4	1	-
Male	-	-	1	3	2	2	1	3	4	1	-	-
Female	-	-	-	-	4	2	4	6	7	3	1	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	3	4	4	7	8	22	30	27	14	2	-
Male	2	1	4	3	3	5	8	12	11	5	1	-
Female	2	2	-	1	4	3	14	18	16	9	1	-
Idaho												
All races, both sexes	6	9	8	10	12	14	17	46	36	13	4	-
Male	4	6	8	6	5	5	6	11	11	4	-	-
Female	2	3	-	4	7	9	11	35	25	9	4	-
White, both sexes	6	9	8	10	12	14	17	46	36	13	4	-
Male	4	6	8	6	5	5	6	11	11	4	-	-
Female	2	3	-	4	7	9	11	35	25	9	4	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Illinois												
All races, both sexes	148	173	142	126	196	209	301	350	299	151	35	1
Male	98	112	85	75	107	95	128	100	70	24	3	1
Female	50	61	57	51	89	114	173	250	229	127	32	-
White, both sexes	76	89	84	75	113	141	235	291	261	134	32	-
Male	50	57	54	45	57	65	101	82	62	19	3	-
Female	26	32	30	30	56	76	134	209	199	115	29	-
Black, both sexes	71	78	56	50	81	60	59	52	35	16	3	1
Male	47	51	29	29	49	27	25	15	7	5	-	1
Female	24	27	27	21	32	33	34	37	28	11	3	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	6	2	1	2	8	7	7	3	1	-	-
Male	-	4	2	1	1	3	2	3	1	-	-	-
Female	-	2	-	-	1	5	5	4	2	1	-	-
Indiana												
All races, both sexes	43	49	46	61	56	93	154	205	169	60	14	-
Male	26	33	32	28	26	43	60	69	39	7	1	-
Female	17	16	14	33	30	50	94	136	130	53	13	-
White, both sexes	33	36	32	47	41	85	133	193	163	55	13	-
Male	23	26	22	23	19	37	54	64	37	7	1	-
Female	10	10	10	24	22	48	79	129	126	48	12	-
Black, both sexes	9	13	13	13	15	8	21	12	6	5	1	-
Male	3	7	10	5	7	6	6	5	2	-	-	-
Female	6	6	3	8	8	2	15	7	4	5	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Iowa												
All races, both sexes	14	10	14	22	24	42	91	115	114	55	11	-
Male	10	6	10	9	10	19	44	36	23	2	-	-
Female	4	4	4	13	14	23	47	79	91	53	11	-
White, both sexes	12	9	12	22	22	41	90	113	111	55	11	-
Male	8	5	9	9	9	19	43	36	22	2	-	-
Female	4	4	3	13	13	22	47	77	89	53	11	-
Black, both sexes	2	1	1	-	2	1	1	2	3	-	-	-
Male	2	1	-	-	1	-	1	-	1	-	-	-
Female	-	-	1	-	1	1	-	2	2	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	6	11	14	13	9	28	47	65	70	37	14	-
Male	5	6	10	6	4	14	18	13	12	4	2	-
Female	1	5	4	7	5	14	29	52	58	33	12	-
White, both sexes	4	10	12	12	7	23	44	60	65	36	13	-
Male	3	6	8	6	4	12	17	13	12	4	2	-
Female	1	4	4	6	3	11	27	47	53	32	11	-
Black, both sexes	2	1	2	1	2	4	3	5	5	-	1	-
Male	2	-	2	-	-	2	1	-	-	-	-	-
Female	-	1	-	1	2	2	2	5	5	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Kentucky												
All races, both sexes	35	42	40	37	41	71	85	101	80	35	12	-
Male	20	26	26	19	22	29	29	35	17	7	1	-
Female	15	16	14	18	19	42	56	66	63	28	11	-
White, both sexes	28	31	32	34	32	68	77	95	75	34	9	-
Male	18	18	21	19	19	27	28	32	17	7	1	-
Female	10	13	11	15	13	41	49	63	58	27	8	-
Black, both sexes	7	11	8	3	8	3	7	6	5	1	3	-
Male	2	8	5	-	2	2	1	3	-	-	-	-
Female	5	3	3	3	6	1	6	3	5	1	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Louisiana												
All races, both sexes	88	99	98	105	107	145	182	184	149	55	20	-
Male	57	56	55	56	55	67	83	54	28	11	2	-
Female	31	43	43	49	52	78	99	130	121	44	18	-
White, both sexes	36	52	44	52	57	88	123	132	110	35	9	-
Male	26	31	28	29	31	44	52	42	22	8	-	-
Female	10	21	16	23	26	44	71	90	88	27	9	-
Black, both sexes	51	47	54	52	49	56	59	52	38	20	11	-
Male	30	25	27	26	24	22	31	12	5	3	2	-
Female	21	22	27	26	25	34	28	40	33	17	9	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	1	1	-	-	1	-	-	-
Male	1	-	-	1	-	1	-	-	1	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Maine												
All races, both sexes	4	6	11	8	6	21	18	18	28	15	5	-
Male	2	6	8	6	3	10	6	10	6	2	-	-
Female	2	-	3	2	3	11	12	8	22	13	5	-
White, both sexes	4	6	11	8	6	21	18	17	28	15	5	-
Male	2	6	8	6	3	10	6	9	6	2	-	-
Female	2	-	3	2	3	11	12	8	22	13	5	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	97	100	110	79	108	115	142	165	137	77	12	-
Male	69	75	70	46	57	61	57	53	31	17	1	-
Female	28	25	40	33	51	54	85	112	106	60	11	-
White, both sexes	42	45	45	44	53	64	94	129	106	60	10	-
Male	31	35	31	27	30	37	40	42	20	13	1	-
Female	11	10	14	17	23	27	54	87	86	47	9	-
Black, both sexes	54	54	61	34	52	46	46	33	29	16	2	-
Male	37	39	36	18	26	23	17	10	9	4	-	-
Female	17	15	25	16	26	23	29	23	20	12	2	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	4	1	3	4	2	3	2	1	-	-
Male	1	1	3	1	1	-	-	1	2	-	-	-
Female	-	-	1	-	2	4	2	2	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Massachusetts												
All races, both sexes	55	65	49	59	54	74	122	180	145	71	16	-
Male	38	40	32	35	32	31	41	55	35	9	1	-
Female	17	25	17	24	22	43	81	125	110	62	15	-
White, both sexes	47	60	42	51	51	68	114	171	139	67	15	-
Male	32	37	29	32	30	28	36	51	34	8	-	-
Female	15	23	13	19	21	40	78	120	105	59	15	-
Black, both sexes	8	5	4	6	3	5	7	4	4	3	-	-
Male	6	3	1	3	2	3	5	1	-	-	-	-
Female	2	2	3	3	1	2	2	3	4	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	3	2	-	1	1	5	2	1	1	-
Male	-	-	2	-	-	-	-	3	1	1	1	-
Female	-	-	1	2	-	1	1	2	1	-	-	-
Michigan												
All races, both sexes	113	131	128	116	139	225	332	379	345	164	39	-
Male	69	91	84	68	51	90	110	121	87	32	5	-
Female	44	40	44	48	88	135	222	258	258	132	34	-
White, both sexes	56	78	77	72	97	175	273	324	320	150	35	-
Male	37	57	51	48	33	74	94	99	82	26	4	-
Female	19	21	26	24	64	101	179	225	238	124	31	-
Black, both sexes	56	51	50	44	39	50	55	53	24	14	4	-
Male	31	32	33	20	16	16	16	20	5	6	1	-
Female	25	19	17	24	23	34	39	33	19	8	3	-
American Indian 1/, both sexes	1	1	1	-	2	-	3	1	-	-	-	-
Male	1	1	-	-	1	-	-	1	-	-	-	-
Female	-	-	1	-	1	-	3	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	-	1	1	1	-	-	-
Male	-	1	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Minnesota												
All races, both sexes	16	32	26	24	35	59	91	119	135	57	13	-
Male	8	28	18	15	19	34	44	33	30	11	4	-
Female	8	4	8	9	16	25	47	86	105	46	9	-
White, both sexes	12	24	24	22	29	52	87	117	134	57	12	-
Male	5	22	16	13	16	31	41	31	30	11	4	-
Female	7	2	8	9	13	21	46	86	104	46	8	-
Black, both sexes	4	7	2	2	1	4	2	2	1	-	1	-
Male	3	5	2	2	-	1	2	2	-	-	-	-
Female	1	2	-	-	1	3	-	-	1	-	1	-
American Indian 1/, both sexes	-	1	-	-	2	1	-	-	-	-	-	-
Male	-	1	-	-	2	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	3	2	2	-	-	-	-	-
Male	-	-	-	-	1	2	1	-	-	-	-	-
Female	-	-	-	-	2	-	1	-	-	-	-	-
Mississippi												
All races, both sexes	63	73	99	82	121	132	167	182	136	53	17	-
Male	41	49	69	49	61	61	58	44	24	11	1	-
Female	22	24	30	33	60	71	109	138	112	42	16	-
White, both sexes	21	30	37	40	64	76	106	114	94	32	7	-
Male	14	24	26	28	35	40	36	24	16	6	1	-
Female	7	6	11	12	29	36	70	90	78	26	6	-
Black, both sexes	41	43	61	42	56	55	61	67	42	21	10	-
Male	26	25	42	21	26	21	22	20	8	5	-	-
Female	15	18	19	21	30	34	39	47	34	16	10	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	1	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Missouri												
All races, both sexes	44	43	48	50	57	93	128	156	115	66	8	-
Male	33	30	32	31	27	34	38	42	29	11	1	-
Female	11	13	16	19	30	59	90	114	86	55	7	-
White, both sexes	20	22	20	32	48	65	99	140	98	59	8	-
Male	14	17	16	21	22	24	30	40	24	9	1	-
Female	6	5	4	11	26	41	69	100	74	50	7	-
Black, both sexes	23	21	28	17	9	27	25	16	16	7	-	-
Male	18	13	16	10	5	10	6	2	5	2	-	-
Female	5	8	12	7	4	17	19	14	11	5	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	-	3	-	1	-	-	-
Male	1	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	1	-	-	-
Montana												
All races, both sexes	6	10	8	3	8	4	14	29	23	19	2	-
Male	4	6	7	3	6	-	4	8	6	3	-	-
Female	2	4	1	-	2	4	10	21	17	16	2	-
White, both sexes	6	10	8	3	8	4	13	29	23	19	2	-
Male	4	6	7	3	6	-	4	8	6	3	-	-
Female	2	4	1	-	2	4	9	21	17	16	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Nebraska												
All races, both sexes	6	6	13	11	18	36	34	55	84	37	14	-
Male	5	4	8	8	10	17	14	18	14	4	2	-
Female	1	2	5	3	8	19	20	37	70	33	12	-
White, both sexes	6	6	9	9	14	35	33	53	84	37	14	-
Male	5	4	6	7	8	17	13	17	14	4	2	-
Female	1	2	3	2	6	18	20	36	70	33	12	-
Black, both sexes	-	-	4	1	4	1	1	2	-	-	-	-
Male	-	-	2	1	2	-	1	1	-	-	-	-
Female	-	-	2	-	2	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	36	43	53	52	34	37	48	56	43	11	3	-
Male	23	27	31	34	19	23	24	21	6	4	-	-
Female	13	16	22	18	15	14	24	35	37	7	3	-
White, both sexes	29	33	46	43	30	34	40	50	42	10	3	-
Male	22	21	26	28	17	22	22	19	5	3	-	-
Female	7	12	20	15	13	12	18	31	37	7	3	-
Black, both sexes	6	8	7	8	2	2	5	4	1	1	-	-
Male	1	5	5	6	1	1	1	1	1	1	-	-
Female	5	3	2	2	1	1	4	3	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	1	2	1	2	1	-	-	-	-
Male	-	1	-	-	1	-	1	-	-	-	-	-
Female	1	-	-	1	1	1	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
New Hampshire												
All races, both sexes	11	7	6	6	10	23	21	32	35	17	7	-
Male	9	5	5	6	7	10	8	9	8	6	3	-
Female	2	2	1	-	3	13	13	23	27	11	4	-
White, both sexes	11	7	6	6	10	23	21	32	35	17	7	-
Male	9	5	5	6	7	10	8	9	8	6	3	-
Female	2	2	1	-	3	13	13	23	27	11	4	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	58	77	96	80	98	123	221	248	211	104	27	-
Male	40	54	65	49	53	49	94	72	43	19	2	-
Female	18	23	31	31	45	74	127	176	168	85	25	-
White, both sexes	35	40	61	46	63	93	177	216	185	96	24	-
Male	24	30	44	30	33	39	75	63	37	18	2	-
Female	11	10	17	16	30	54	102	153	148	78	22	-
Black, both sexes	23	35	33	31	33	29	37	24	25	6	3	-
Male	16	23	21	17	18	10	17	7	6	1	-	-
Female	7	12	12	14	15	19	20	17	19	5	3	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	2	2	2	1	7	8	1	2	-	-
Male	-	1	-	1	2	-	2	2	-	-	-	-
Female	-	1	2	1	-	1	5	6	1	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
New Mexico												
All races, both sexes	17	15	18	21	25	17	28	47	41	14	4	-
Male	14	10	12	10	13	7	9	14	11	2	-	-
Female	3	5	6	11	12	10	19	33	30	12	4	-
White, both sexes	16	14	16	17	21	15	24	43	40	13	4	-
Male	13	9	10	7	11	5	7	13	11	1	-	-
Female	3	5	6	10	10	10	17	30	29	12	4	-
Black, both sexes	1	1	1	1	3	1	3	2	1	-	-	-
Male	1	1	1	-	1	1	2	-	-	-	-	-
Female	-	-	-	1	2	-	1	2	1	-	-	-
American Indian 1/, both sexes	-	-	1	3	1	1	1	2	-	1	-	-
Male	-	-	1	3	1	1	-	1	-	1	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	241	271	328	309	344	473	590	670	588	285	70	-
Male	152	186	203	179	164	230	230	220	167	57	10	-
Female	89	85	125	130	180	243	360	450	421	228	60	-
White, both sexes	135	149	197	189	217	325	448	530	498	239	57	-
Male	89	111	118	112	103	159	166	174	148	51	8	-
Female	46	38	79	77	114	166	282	356	350	188	49	-
Black, both sexes	101	114	128	110	118	131	127	119	73	40	9	-
Male	60	67	83	61	54	65	54	40	16	3	-	-
Female	41	47	45	49	64	66	73	79	57	37	9	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	5	8	2	10	9	17	15	21	16	6	4	-
Male	3	8	1	6	7	6	10	6	3	3	2	-
Female	2	-	1	4	2	11	5	15	13	3	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
North Carolina												
All races, both sexes	88	86	102	94	109	146	192	221	166	69	20	-
Male	60	60	71	49	50	73	58	58	34	11	1	-
Female	28	26	31	45	59	73	134	163	132	58	19	-
White, both sexes	32	42	48	37	57	84	126	147	125	49	16	-
Male	27	31	34	21	23	42	33	40	24	11	1	-
Female	5	11	14	16	34	42	93	107	101	38	15	-
Black, both sexes	52	42	51	53	51	60	63	73	40	20	4	-
Male	30	27	36	27	27	30	24	18	10	-	-	-
Female	22	15	15	26	24	30	39	55	30	20	4	-
American Indian 1/, both sexes	4	1	1	3	1	2	3	-	-	-	-	-
Male	3	1	-	1	-	1	1	-	-	-	-	-
Female	1	-	1	2	1	1	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	2	1	-	-	-	1	1	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	1	1	-	-	-
North Dakota												
All races, both sexes	2	-	5	6	6	5	24	31	33	16	3	-
Male	2	-	2	2	1	3	13	8	8	1	-	-
Female	-	-	3	4	5	2	11	23	25	15	3	-
White, both sexes	1	-	4	4	6	5	24	30	33	16	3	-
Male	1	-	2	2	1	3	13	8	8	1	-	-
Female	-	-	2	2	5	2	11	22	25	15	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	2	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	2	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Ohio												
All races, both sexes	125	163	182	194	243	295	475	541	465	194	50	-
Male	88	113	122	105	118	139	170	151	111	22	8	-
Female	37	50	60	89	125	156	305	390	354	172	42	-
White, both sexes	74	109	122	146	187	222	404	470	417	174	44	-
Male	57	80	83	78	95	106	139	128	94	18	7	-
Female	17	29	39	68	92	116	265	342	323	156	37	-
Black, both sexes	51	52	58	46	54	72	66	71	47	19	6	-
Male	31	32	37	25	23	32	29	23	16	3	1	-
Female	20	20	21	21	31	40	37	48	31	16	5	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	2	2	2	1	4	-	1	1	-	-
Male	-	1	2	2	-	1	1	-	1	1	-	-
Female	-	1	-	-	2	-	3	-	-	-	-	-
Oklahoma												
All races, both sexes	20	29	27	37	47	55	96	95	82	46	12	-
Male	16	18	14	18	22	21	38	27	15	4	3	-
Female	4	11	13	19	25	34	58	68	67	42	9	-
White, both sexes	13	22	18	29	37	40	81	81	77	41	11	-
Male	10	13	11	15	19	15	32	25	14	4	3	-
Female	3	9	7	14	18	25	49	56	63	37	8	-
Black, both sexes	5	6	6	5	6	9	7	13	3	3	1	-
Male	4	4	1	1	2	2	2	2	1	-	-	-
Female	1	2	5	4	4	7	5	11	2	3	1	-
American Indian 1/, both sexes	2	1	3	3	2	6	7	1	2	1	-	-
Male	2	1	2	2	1	4	4	-	-	-	-	-
Female	-	-	1	1	1	2	3	1	2	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	1	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Oregon												
All races, both sexes	15	28	24	26	37	48	92	136	124	63	10	-
Male	11	17	19	16	18	22	34	41	31	12	1	-
Female	4	11	5	10	19	26	58	95	93	51	9	-
White, both sexes	14	23	20	24	36	45	88	134	119	61	10	-
Male	10	15	15	15	18	20	32	40	28	10	1	-
Female	4	8	5	9	18	25	56	94	91	51	9	-
Black, both sexes	1	4	2	-	-	2	2	-	2	-	-	-
Male	1	2	2	-	-	1	1	-	1	-	-	-
Female	-	2	-	-	-	1	1	-	1	-	-	-
American Indian 1/, both sexes	-	1	1	1	1	-	1	1	-	-	-	-
Male	-	-	1	1	-	-	1	-	-	-	-	-
Female	-	1	-	-	1	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	1	1	1	3	2	-	-
Male	-	-	1	-	-	1	-	1	2	2	-	-
Female	-	-	-	1	-	-	1	-	1	-	-	-
Pennsylvania												
All races, both sexes	65	86	116	92	116	187	326	367	295	133	40	-
Male	47	56	76	61	57	84	120	116	71	30	3	-
Female	18	30	40	31	59	103	206	251	224	103	37	-
White, both sexes	44	56	89	73	97	152	283	323	270	123	38	-
Male	34	40	57	48	47	71	107	101	63	29	3	-
Female	10	16	32	25	50	81	176	222	207	94	35	-
Black, both sexes	21	29	25	19	18	35	43	38	25	10	2	-
Male	13	16	18	13	9	13	13	13	8	1	-	-
Female	8	13	7	6	9	22	30	25	17	9	2	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	1	-	-	6	-	-	-	-
Male	-	-	1	-	1	-	-	2	-	-	-	-
Female	-	1	-	-	-	-	-	4	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Rhode Island												
All races, both sexes	6	7	2	6	6	9	15	30	18	7	2	-
Male	4	3	1	4	1	2	9	5	5	2	-	-
Female	2	4	1	2	5	7	6	25	13	5	2	-
White, both sexes	4	7	2	5	5	8	15	30	18	7	2	-
Male	4	3	1	4	-	2	9	5	5	2	-	-
Female	-	4	1	1	5	6	6	25	13	5	2	-
Black, both sexes	1	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	44	52	64	71	68	95	117	132	95	42	15	-
Male	23	32	40	51	36	46	42	35	26	10	1	-
Female	21	20	24	20	32	49	75	97	69	32	14	-
White, both sexes	16	19	27	32	31	42	75	90	64	26	11	-
Male	9	10	19	24	19	20	25	21	19	6	-	-
Female	7	9	8	8	12	22	50	69	45	20	11	-
Black, both sexes	28	33	37	38	37	53	42	42	31	16	4	-
Male	14	22	21	26	17	26	17	14	7	4	1	-
Female	14	11	16	12	20	27	25	28	24	12	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
South Dakota												
All races, both sexes	2	2	-	2	2	9	25	30	29	21	5	-
Male	2	-	-	1	1	5	13	9	7	3	-	-
Female	-	2	-	1	1	4	12	21	22	18	5	-
White, both sexes	1	1	-	1	2	9	24	29	29	21	5	-
Male	1	-	-	-	1	5	13	9	7	3	-	-
Female	-	1	-	1	1	4	11	20	22	18	5	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	1	-	-	1	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	88	99	82	90	119	137	209	265	183	90	11	-
Male	59	63	48	54	57	66	62	77	47	19	1	-
Female	29	36	34	36	62	71	147	188	136	71	10	-
White, both sexes	48	53	44	63	69	95	167	220	154	77	9	-
Male	37	39	25	38	33	35	49	67	40	17	1	-
Female	11	14	19	25	36	60	118	153	114	60	8	-
Black, both sexes	40	45	37	26	49	42	39	45	28	13	2	-
Male	22	24	22	15	23	31	11	10	7	2	-	-
Female	18	21	15	11	26	11	28	35	21	11	2	-
American Indian 1/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	1	-	2	-	1	-	-	-
Male	-	-	-	1	1	-	2	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Texas												
All races, both sexes	251	223	242	229	263	352	483	577	412	223	51	-
Male	158	133	140	134	110	156	192	181	93	34	12	-
Female	93	90	102	95	153	196	291	396	319	189	39	-
White, both sexes	175	145	180	172	200	284	399	503	352	197	43	-
Male	119	95	103	103	85	132	162	161	79	30	9	-
Female	56	50	77	69	115	152	237	342	273	167	34	-
Black, both sexes	73	75	58	55	60	62	72	69	51	22	8	-
Male	38	37	34	29	25	20	27	18	12	3	3	-
Female	35	38	24	26	35	42	45	51	39	19	5	-
American Indian 1/, both sexes	-	-	1	-	-	2	1	2	-	-	-	-
Male	-	-	1	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	3	3	2	3	4	11	3	9	4	-	-
Male	1	1	2	2	-	3	3	1	2	1	-	-
Female	2	2	1	-	3	1	8	2	7	3	-	-
Utah												
All races, both sexes	6	12	10	12	11	25	41	45	51	18	3	-
Male	5	7	9	7	6	10	16	18	11	2	-	-
Female	1	5	1	5	5	15	25	27	40	16	3	-
White, both sexes	5	11	9	11	11	25	40	43	48	17	3	-
Male	5	6	9	6	6	10	16	18	10	2	-	-
Female	-	5	-	5	5	15	24	25	38	15	3	-
Black, both sexes	-	1	-	1	-	-	-	-	-	1	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	-	-	2	2	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	1	-	-	-	-	2	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Vermont												
All races, both sexes	3	6	2	5	4	12	19	18	12	17	2	-
Male	2	3	1	4	2	6	7	7	2	4	-	-
Female	1	3	1	1	2	6	12	11	10	13	2	-
White, both sexes	3	5	2	5	4	12	19	18	12	17	2	-
Male	2	3	1	4	2	6	7	7	2	4	-	-
Female	1	2	1	1	2	6	12	11	10	13	2	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	56	58	57	57	64	93	131	161	134	50	18	-
Male	35	37	33	34	31	45	50	51	21	9	2	-
Female	21	21	24	23	33	48	81	110	113	41	16	-
White, both sexes	27	30	25	29	31	69	82	126	95	37	13	-
Male	16	23	13	19	15	32	30	44	15	3	-	-
Female	11	7	12	10	16	37	52	82	80	34	13	-
Black, both sexes	29	27	30	27	30	23	45	33	35	13	5	-
Male	19	14	19	14	16	12	19	6	5	6	2	-
Female	10	13	11	13	14	11	26	27	30	7	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	2	1	3	1	4	2	3	-	-	-
Male	-	-	1	1	-	1	1	1	1	-	-	-
Female	-	1	1	-	3	-	3	1	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Washington												
All races, both sexes	42	79	76	67	80	129	219	295	285	112	24	-
Male	34	63	54	33	47	56	93	104	76	26	2	-
Female	8	16	22	34	33	73	126	191	209	86	22	-
White, both sexes	34	65	67	55	72	114	193	283	276	110	24	-
Male	28	52	50	27	43	51	83	99	74	25	2	-
Female	6	13	17	28	29	63	110	184	202	85	22	-
Black, both sexes	4	11	2	9	3	9	10	2	-	1	-	-
Male	3	8	-	6	1	3	3	2	-	-	-	-
Female	1	3	2	3	2	6	7	-	-	1	-	-
American Indian 1/, both sexes	2	-	2	2	-	2	1	1	2	-	-	-
Male	2	-	2	-	-	1	-	1	-	-	-	-
Female	-	-	-	2	-	1	1	-	2	-	-	-
Asian or Pacific Islander 2/, both sexes	2	3	5	1	5	4	15	9	7	1	-	-
Male	1	3	2	-	3	1	7	2	2	1	-	-
Female	1	-	3	1	2	3	8	7	5	-	-	-
West Virginia												
All races, both sexes	15	20	22	24	30	35	47	69	60	27	6	-
Male	11	13	16	15	12	18	17	18	11	9	1	-
Female	4	7	6	9	18	17	30	51	49	18	5	-
White, both sexes	15	17	22	21	29	33	43	66	59	27	6	-
Male	11	12	16	13	11	17	16	17	11	9	1	-
Female	4	5	6	8	18	16	27	49	48	18	5	-
Black, both sexes	-	3	-	3	1	2	4	3	1	-	-	-
Male	-	1	-	2	1	1	1	1	-	-	-	-
Female	-	2	-	1	-	1	3	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I15)												
Wisconsin												
All races, both sexes	21	31	47	45	48	81	149	165	142	76	21	-
Male	17	17	26	25	31	39	46	58	34	11	2	-
Female	4	14	21	20	17	42	103	107	108	65	19	-
White, both sexes	14	22	35	38	42	70	137	161	138	75	21	-
Male	11	12	19	20	27	32	41	56	33	10	2	-
Female	3	10	16	18	15	38	96	105	105	65	19	-
Black, both sexes	7	8	12	6	5	10	11	4	3	1	-	-
Male	6	5	7	5	3	7	5	2	-	1	-	-
Female	1	3	5	1	2	3	6	2	3	-	-	-
American Indian 1/, both sexes	-	1	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Wyoming												
All races, both sexes	3	6	3	1	6	7	4	6	9	7	1	-
Male	2	5	2	1	5	5	2	5	1	-	-	-
Female	1	1	1	-	1	2	2	1	8	7	1	-
White, both sexes	3	6	3	1	6	7	4	6	9	7	1	-
Male	2	5	2	1	5	5	2	5	1	-	-	-
Female	1	1	1	-	1	2	2	1	8	7	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
United States												
All races, both sexes	14,414	-	1	-	-	5	6	10	18	53	146	238
Male	5,247	-	1	-	-	3	5	6	13	39	99	149
Female	9,167	-	-	-	-	2	1	4	5	14	47	89
White, both sexes	11,796	-	1	-	-	4	4	5	10	30	98	146
Male	4,167	-	1	-	-	2	3	4	7	27	71	101
Female	7,629	-	-	-	-	2	1	1	3	3	27	45
Black, both sexes	2,227	-	-	-	-	1	2	5	8	23	46	89
Male	906	-	-	-	-	1	2	2	6	12	27	46
Female	1,321	-	-	-	-	-	-	3	2	11	19	43
American Indian 1/, both sexes	58	-	-	-	-	-	-	-	-	-	-	2
Male	26	-	-	-	-	-	-	-	-	-	-	1
Female	32	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	333	-	-	-	-	-	-	-	-	-	2	1
Male	148	-	-	-	-	-	-	-	-	-	1	1
Female	185	-	-	-	-	-	-	-	-	-	1	-
Alabama												
All races, both sexes	311	-	-	-	-	-	1	-	1	7	8	11
Male	141	-	-	-	-	-	1	-	1	4	5	7
Female	170	-	-	-	-	-	-	-	-	3	3	4
White, both sexes	204	-	-	-	-	-	1	-	1	2	4	5
Male	92	-	-	-	-	-	1	-	1	2	2	3
Female	112	-	-	-	-	-	-	-	-	-	2	2
Black, both sexes	106	-	-	-	-	-	-	-	-	5	4	6
Male	48	-	-	-	-	-	-	-	-	2	3	4
Female	58	-	-	-	-	-	-	-	-	3	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Alaska												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	193	-	-	-	-	-	-	-	-	1	-	4
Male	67	-	-	-	-	-	-	-	-	1	-	3
Female	126	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	172	-	-	-	-	-	-	-	-	1	-	3
Male	58	-	-	-	-	-	-	-	-	1	-	3
Female	114	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Arkansas												
All races, both sexes	171	-	-	-	-	-	1	-	-	4	5	6
Male	66	-	-	-	-	-	-	-	-	4	4	4
Female	105	-	-	-	-	-	1	-	-	-	1	2
White, both sexes	138	-	-	-	-	-	1	-	-	2	4	3
Male	49	-	-	-	-	-	-	-	-	2	3	2
Female	89	-	-	-	-	-	1	-	-	-	1	1
Black, both sexes	33	-	-	-	-	-	-	-	-	2	1	3
Male	17	-	-	-	-	-	-	-	-	2	1	2
Female	16	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1,981	-	-	-	-	-	-	-	2	5	11	25
Male	724	-	-	-	-	-	-	-	2	5	7	18
Female	1,257	-	-	-	-	-	-	-	-	-	4	7
White, both sexes	1,612	-	-	-	-	-	-	-	2	4	7	21
Male	569	-	-	-	-	-	-	-	2	4	5	15
Female	1,043	-	-	-	-	-	-	-	-	-	2	6
Black, both sexes	199	-	-	-	-	-	-	-	-	1	3	4
Male	81	-	-	-	-	-	-	-	-	1	2	3
Female	118	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	163	-	-	-	-	-	-	-	-	-	1	-
Male	70	-	-	-	-	-	-	-	-	-	-	-
Female	93	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Colorado												
All races, both sexes	123	-	-	-	-	-	-	-	-	1	1	1
Male	46	-	-	-	-	-	-	-	-	1	1	1
Female	77	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	120	-	-	-	-	-	-	-	-	1	1	1
Male	44	-	-	-	-	-	-	-	-	1	1	1
Female	76	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	222	-	-	-	-	-	-	-	-	1	-	6
Male	74	-	-	-	-	-	-	-	-	1	-	3
Female	148	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	202	-	-	-	-	-	-	-	-	1	-	5
Male	65	-	-	-	-	-	-	-	-	1	-	3
Female	137	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	19	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Delaware												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	3
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	17	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	2
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Florida												
All races, both sexes	969	-	-	-	-	1	1	1	2	3	7	11
Male	380	-	-	-	-	1	1	-	2	2	5	7
Female	589	-	-	-	-	-	-	1	-	1	2	4
White, both sexes	819	-	-	-	-	1	1	-	1	2	5	6
Male	318	-	-	-	-	1	1	-	1	2	4	3
Female	501	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	143	-	-	-	-	-	-	1	1	1	2	5
Male	57	-	-	-	-	-	-	-	1	-	1	4
Female	86	-	-	-	-	-	-	1	-	1	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	645	-	-	-	-	-	2	1	2	9	12	19
Male	260	-	-	-	-	-	2	1	2	6	8	9
Female	385	-	-	-	-	-	-	-	-	3	4	10
White, both sexes	395	-	-	-	-	-	1	1	1	4	6	10
Male	157	-	-	-	-	-	1	1	1	3	5	8
Female	238	-	-	-	-	-	-	-	-	1	1	2
Black, both sexes	244	-	-	-	-	-	1	-	1	5	6	9
Male	101	-	-	-	-	-	1	-	1	3	3	1
Female	143	-	-	-	-	-	-	-	-	2	3	8
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Hawaii												
All races, both sexes	66	-	-	-	-	-	-	1	-	-	-	1
Male	30	-	-	-	-	-	-	1	-	-	-	-
Female	36	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	16	-	-	-	-	-	-	1	-	-	-	1
Male	7	-	-	-	-	-	-	1	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	50	-	-	-	-	-	-	-	-	-	-	-
Male	23	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	35	-	-	-	-	-	-	-	-	-	1	-
Male	9	-	-	-	-	-	-	-	-	-	1	-
Female	26	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	35	-	-	-	-	-	-	-	-	-	1	-
Male	9	-	-	-	-	-	-	-	-	-	1	-
Female	26	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Illinois												
All races, both sexes	515	-	-	-	-	-	-	-	1	-	5	6
Male	157	-	-	-	-	-	-	-	-	-	4	6
Female	358	-	-	-	-	-	-	-	1	-	1	-
White, both sexes	417	-	-	-	-	-	-	-	-	-	3	3
Male	126	-	-	-	-	-	-	-	-	-	3	3
Female	291	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	87	-	-	-	-	-	-	-	1	-	2	3
Male	29	-	-	-	-	-	-	-	-	-	1	3
Female	58	-	-	-	-	-	-	-	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	315	-	-	-	-	-	-	1	3	-	2	2
Male	109	-	-	-	-	-	-	1	2	-	-	1
Female	206	-	-	-	-	-	-	-	1	-	2	1
White, both sexes	291	-	-	-	-	-	-	-	1	-	2	1
Male	100	-	-	-	-	-	-	-	1	-	-	1
Female	191	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	23	-	-	-	-	-	-	1	2	-	-	1
Male	9	-	-	-	-	-	-	1	1	-	-	-
Female	14	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Iowa												
All races, both sexes	106	-	-	-	-	-	-	-	-	-	1	2
Male	33	-	-	-	-	-	-	-	-	-	-	2
Female	73	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	105	-	-	-	-	-	-	-	-	-	1	2
Male	32	-	-	-	-	-	-	-	-	-	-	2
Female	73	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	116	-	-	-	-	-	-	-	-	-	-	2
Male	34	-	-	-	-	-	-	-	-	-	-	1
Female	82	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	106	-	-	-	-	-	-	-	-	-	-	2
Male	32	-	-	-	-	-	-	-	-	-	-	1
Female	74	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Kentucky												
All races, both sexes	192	-	-	-	-	-	-	1	-	-	3	4
Male	64	-	-	-	-	-	-	1	-	-	1	2
Female	128	-	-	-	-	-	-	-	-	-	2	2
White, both sexes	169	-	-	-	-	-	-	1	-	-	3	2
Male	59	-	-	-	-	-	-	1	-	-	1	1
Female	110	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	23	-	-	-	-	-	-	-	-	-	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	18	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	234	-	-	-	-	-	-	1	-	1	5	7
Male	90	-	-	-	-	-	-	1	-	1	1	3
Female	144	-	-	-	-	-	-	-	-	-	4	4
White, both sexes	145	-	-	-	-	-	-	-	-	-	3	3
Male	50	-	-	-	-	-	-	-	-	-	1	2
Female	95	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	88	-	-	-	-	-	-	1	-	1	2	4
Male	39	-	-	-	-	-	-	1	-	1	-	1
Female	49	-	-	-	-	-	-	-	-	-	2	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Maine												
All races, both sexes	43	-	-	-	-	-	-	-	-	-	1	-
Male	13	-	-	-	-	-	-	-	-	-	1	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	43	-	-	-	-	-	-	-	-	-	1	-
Male	13	-	-	-	-	-	-	-	-	-	1	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	206	-	-	-	-	-	-	-	-	-	2	3
Male	66	-	-	-	-	-	-	-	-	-	1	2
Female	140	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	140	-	-	-	-	-	-	-	-	-	-	-
Male	39	-	-	-	-	-	-	-	-	-	-	-
Female	101	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	60	-	-	-	-	-	-	-	-	-	2	3
Male	24	-	-	-	-	-	-	-	-	-	1	2
Female	36	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Massachusetts												
All races, both sexes	329	-	-	-	-	-	-	-	2	-	2	4
Male	109	-	-	-	-	-	-	-	1	-	2	2
Female	220	-	-	-	-	-	-	-	1	-	-	2
White, both sexes	309	-	-	-	-	-	-	-	2	-	2	3
Male	103	-	-	-	-	-	-	-	1	-	2	2
Female	206	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	16	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	493	-	-	-	-	-	-	-	-	1	4	8
Male	179	-	-	-	-	-	-	-	-	1	3	6
Female	314	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	425	-	-	-	-	-	-	-	-	1	2	6
Male	152	-	-	-	-	-	-	-	-	1	1	6
Female	273	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	61	-	-	-	-	-	-	-	-	-	2	2
Male	25	-	-	-	-	-	-	-	-	-	2	-
Female	36	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Minnesota												
All races, both sexes	284	-	-	-	-	-	-	-	-	-	-	2
Male	109	-	-	-	-	-	-	-	-	-	-	2
Female	175	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	275	-	-	-	-	-	-	-	-	-	-	2
Male	106	-	-	-	-	-	-	-	-	-	-	2
Female	169	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	275	-	-	-	-	-	-	-	3	-	6	8
Male	104	-	-	-	-	-	-	-	2	-	4	6
Female	171	-	-	-	-	-	-	-	1	-	2	2
White, both sexes	153	-	-	-	-	-	-	-	1	-	1	2
Male	49	-	-	-	-	-	-	-	-	-	1	2
Female	104	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	121	-	-	-	-	-	-	-	2	-	5	6
Male	54	-	-	-	-	-	-	-	2	-	3	4
Female	67	-	-	-	-	-	-	-	-	-	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Missouri												
All races, both sexes	218	-	-	-	-	-	-	-	-	1	1	2
Male	71	-	-	-	-	-	-	-	-	-	-	2
Female	147	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	186	-	-	-	-	-	-	-	-	1	-	1
Male	59	-	-	-	-	-	-	-	-	-	-	1
Female	127	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	32	-	-	-	-	-	-	-	-	-	1	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	20	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	55	-	-	-	-	-	-	-	-	-	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	36	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	54	-	-	-	-	-	-	-	-	-	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	35	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Nebraska												
All races, both sexes	125	-	-	-	-	-	-	-	-	-	3	1
Male	35	-	-	-	-	-	-	-	-	-	2	-
Female	90	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	120	-	-	-	-	-	-	-	-	-	3	-
Male	32	-	-	-	-	-	-	-	-	-	2	-
Female	88	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	92	-	-	-	-	-	-	-	-	1	1	3
Male	39	-	-	-	-	-	-	-	-	-	1	1
Female	53	-	-	-	-	-	-	-	-	1	-	2
White, both sexes	81	-	-	-	-	-	-	-	-	-	1	2
Male	36	-	-	-	-	-	-	-	-	-	1	1
Female	45	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	8	-	-	-	-	-	-	-	-	1	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
New Hampshire												
All races, both sexes	39	-	-	-	-	-	-	-	-	-	-	1
Male	16	-	-	-	-	-	-	-	-	-	-	1
Female	23	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	39	-	-	-	-	-	-	-	-	-	-	1
Male	16	-	-	-	-	-	-	-	-	-	-	1
Female	23	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	340	-	-	-	-	-	-	-	-	1	2	9
Male	123	-	-	-	-	-	-	-	-	1	-	5
Female	217	-	-	-	-	-	-	-	-	-	2	4
White, both sexes	277	-	-	-	-	-	-	-	-	1	1	5
Male	99	-	-	-	-	-	-	-	-	1	-	3
Female	178	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	57	-	-	-	-	-	-	-	-	-	1	4
Male	22	-	-	-	-	-	-	-	-	-	-	2
Female	35	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
New Mexico												
All races, both sexes	51	-	-	-	-	-	-	-	-	-	1	-
Male	18	-	-	-	-	-	-	-	-	-	1	-
Female	33	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	45	-	-	-	-	-	-	-	-	-	1	-
Male	14	-	-	-	-	-	-	-	-	-	1	-
Female	31	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1,006	-	-	-	-	-	-	-	-	-	3	13
Male	380	-	-	-	-	-	-	-	-	-	3	10
Female	626	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	779	-	-	-	-	-	-	-	-	-	2	8
Male	290	-	-	-	-	-	-	-	-	-	2	6
Female	489	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	192	-	-	-	-	-	-	-	-	-	-	5
Male	72	-	-	-	-	-	-	-	-	-	-	4
Female	120	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	34	-	-	-	-	-	-	-	-	-	1	-
Male	18	-	-	-	-	-	-	-	-	-	1	-
Female	16	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
North Carolina												
All races, both sexes	365	-	-	-	-	1	-	1	-	3	6	8
Male	127	-	-	-	-	1	-	1	-	1	2	5
Female	238	-	-	-	-	-	-	-	-	2	4	3
White, both sexes	233	-	-	-	-	1	-	1	-	-	2	3
Male	80	-	-	-	-	1	-	1	-	-	1	2
Female	153	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	127	-	-	-	-	-	-	-	-	3	4	5
Male	46	-	-	-	-	-	-	-	-	1	1	3
Female	81	-	-	-	-	-	-	-	-	2	3	2
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	34	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	32	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Ohio												
All races, both sexes	687	-	-	-	-	-	-	-	-	1	9	8
Male	219	-	-	-	-	-	-	-	-	1	6	6
Female	468	-	-	-	-	-	-	-	-	-	3	2
White, both sexes	603	-	-	-	-	-	-	-	-	1	6	5
Male	187	-	-	-	-	-	-	-	-	1	4	3
Female	416	-	-	-	-	-	-	-	-	-	2	2
Black, both sexes	81	-	-	-	-	-	-	-	-	-	3	3
Male	31	-	-	-	-	-	-	-	-	-	2	3
Female	50	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	161	-	-	-	-	1	-	-	-	1	1	1
Male	52	-	-	-	-	1	-	-	-	1	-	1
Female	109	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	137	-	-	-	-	-	-	-	-	1	1	1
Male	41	-	-	-	-	-	-	-	-	1	-	1
Female	96	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	17	-	-	-	-	1	-	-	-	-	-	-
Male	6	-	-	-	-	1	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Oregon												
All races, both sexes	250	-	-	-	-	-	-	-	-	1	4	5
Male	105	-	-	-	-	-	-	-	-	-	4	4
Female	145	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	239	-	-	-	-	-	-	-	-	1	3	3
Male	96	-	-	-	-	-	-	-	-	-	3	2
Female	143	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	638	-	-	-	-	-	-	-	-	1	9	11
Male	240	-	-	-	-	-	-	-	-	1	7	7
Female	398	-	-	-	-	-	-	-	-	-	2	4
White, both sexes	585	-	-	-	-	-	-	-	-	1	9	10
Male	218	-	-	-	-	-	-	-	-	1	7	6
Female	367	-	-	-	-	-	-	-	-	-	2	4
Black, both sexes	50	-	-	-	-	-	-	-	-	-	-	1
Male	20	-	-	-	-	-	-	-	-	-	-	1
Female	30	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Rhode Island												
All races, both sexes	33	-	-	-	-	-	-	-	-	-	2	1
Male	10	-	-	-	-	-	-	-	-	-	1	-
Female	23	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	31	-	-	-	-	-	-	-	-	-	2	1
Male	9	-	-	-	-	-	-	-	-	-	1	-
Female	22	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	217	-	-	-	-	1	-	-	-	1	2	3
Male	95	-	-	-	-	-	-	-	-	-	2	1
Female	122	-	-	-	-	1	-	-	-	1	-	2
White, both sexes	127	-	-	-	-	1	-	-	-	-	1	-
Male	48	-	-	-	-	-	-	-	-	-	1	-
Female	79	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	90	-	-	-	-	-	-	-	-	1	1	3
Male	47	-	-	-	-	-	-	-	-	-	1	1
Female	43	-	-	-	-	-	-	-	-	1	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
South Dakota												
All races, both sexes	39	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	38	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	269	-	-	-	-	-	-	1	-	3	3	3
Male	94	-	-	-	-	-	-	-	-	2	2	1
Female	175	-	-	-	-	-	-	1	-	1	1	2
White, both sexes	209	-	-	-	-	-	-	-	-	1	2	1
Male	69	-	-	-	-	-	-	-	-	1	2	-
Female	140	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	59	-	-	-	-	-	-	1	-	2	1	2
Male	24	-	-	-	-	-	-	-	-	1	-	1
Female	35	-	-	-	-	-	-	1	-	1	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Texas												
All races, both sexes	828	-	1	-	-	1	1	1	1	3	13	16
Male	307	-	1	-	-	-	1	-	-	3	12	10
Female	521	-	-	-	-	1	-	1	1	-	1	6
White, both sexes	700	-	1	-	-	1	-	1	1	3	9	11
Male	256	-	1	-	-	-	-	-	-	3	8	8
Female	444	-	-	-	-	1	-	1	1	-	1	3
Black, both sexes	119	-	-	-	-	-	1	-	-	-	4	5
Male	49	-	-	-	-	-	1	-	-	-	4	2
Female	70	-	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	101	-	-	-	-	-	-	-	-	-	1	3
Male	36	-	-	-	-	-	-	-	-	-	1	1
Female	65	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	100	-	-	-	-	-	-	-	-	-	1	3
Male	36	-	-	-	-	-	-	-	-	-	1	1
Female	64	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Vermont												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	307	-	-	-	-	-	-	1	1	2	5	5
Male	117	-	-	-	-	-	-	-	1	2	4	2
Female	190	-	-	-	-	-	-	1	-	-	1	3
White, both sexes	216	-	-	-	-	-	-	-	-	1	4	-
Male	83	-	-	-	-	-	-	-	-	1	3	-
Female	133	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	82	-	-	-	-	-	-	1	1	1	1	5
Male	31	-	-	-	-	-	-	-	1	1	1	2
Female	51	-	-	-	-	-	-	1	-	-	-	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Washington												
All races, both sexes	239	-	-	-	-	-	-	-	-	-	-	2
Male	88	-	-	-	-	-	-	-	-	-	-	1
Female	151	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	224	-	-	-	-	-	-	-	-	-	-	2
Male	79	-	-	-	-	-	-	-	-	-	-	1
Female	145	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	159	-	-	-	-	-	-	-	-	-	2	2
Male	68	-	-	-	-	-	-	-	-	-	2	1
Female	91	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	150	-	-	-	-	-	-	-	-	-	2	2
Male	64	-	-	-	-	-	-	-	-	-	2	1
Female	86	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Wisconsin												
All races, both sexes	236	-	-	-	-	-	-	-	-	1	1	4
Male	75	-	-	-	-	-	-	-	-	1	1	3
Female	161	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	226	-	-	-	-	-	-	-	-	1	1	2
Male	70	-	-	-	-	-	-	-	-	1	1	2
Female	156	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	21	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	21	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
United States												
All races, both sexes	470	592	720	796	939	1,442	2,173	2,805	2,439	1,218	342	1
Male	297	388	451	438	458	674	738	764	520	180	24	-
Female	173	204	269	358	481	768	1,435	2,041	1,919	1,038	318	1
White, both sexes	321	402	504	590	723	1,165	1,830	2,452	2,160	1,060	291	-
Male	210	290	322	326	357	546	623	652	456	150	19	-
Female	111	112	182	264	366	619	1,207	1,800	1,704	910	272	-
Black, both sexes	138	181	192	180	182	238	267	283	217	128	46	1
Male	79	92	110	96	85	113	87	81	44	21	2	-
Female	59	89	82	84	97	125	180	202	173	107	44	1
American Indian 1/, both sexes	3	2	8	5	3	7	12	11	3	2	-	-
Male	2	2	5	1	1	5	2	7	-	-	-	-
Female	1	-	3	4	2	2	10	4	3	2	-	-
Asian or Pacific Islander 2/, both sexes	8	7	16	21	31	32	64	59	59	28	5	-
Male	6	4	14	15	15	10	26	24	20	9	3	-
Female	2	3	2	6	16	22	38	35	39	19	2	-
Alabama												
All races, both sexes	24	12	21	36	27	31	42	32	35	16	7	-
Male	17	9	14	19	20	15	10	10	8	1	-	-
Female	7	3	7	17	7	16	32	22	27	15	7	-
White, both sexes	15	8	8	22	23	21	30	21	28	12	3	-
Male	11	7	6	13	17	11	5	6	7	-	-	-
Female	4	1	2	9	6	10	25	15	21	12	3	-
Black, both sexes	9	3	13	14	4	10	12	11	7	4	4	-
Male	6	1	8	6	3	4	5	4	1	1	-	-
Female	3	2	5	8	1	6	7	7	6	3	4	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Alaska												
All races, both sexes	-	1	1	-	1	2	-	2	1	-	-	-
Male	-	-	1	-	1	1	-	-	1	-	-	-
Female	-	1	-	-	-	1	-	2	-	-	-	-
White, both sexes	-	1	1	-	1	-	-	1	1	-	-	-
Male	-	-	1	-	1	-	-	-	1	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	15	2	6	10	13	15	25	48	32	18	4	-
Male	7	2	2	6	6	8	10	12	8	1	1	-
Female	8	-	4	4	7	7	15	36	24	17	3	-
White, both sexes	10	-	6	9	12	12	24	45	29	17	4	-
Male	4	-	2	6	6	6	10	10	8	1	1	-
Female	6	-	4	3	6	6	14	35	21	16	3	-
Black, both sexes	3	1	-	1	-	-	-	-	1	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	2	1	-	-	1	3	-	3	-	1	-	-
Male	1	1	-	-	-	2	-	2	-	-	-	-
Female	1	-	-	-	1	1	-	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Arkansas												
All races, both sexes	3	15	13	11	10	16	20	28	24	12	3	-
Male	1	12	7	4	3	7	8	4	8	-	-	-
Female	2	3	6	7	7	9	12	24	16	12	3	-
White, both sexes	3	10	12	9	8	13	17	23	21	9	3	-
Male	1	8	7	2	3	6	6	3	6	-	-	-
Female	2	2	5	7	5	7	11	20	15	9	3	-
Black, both sexes	-	5	1	2	2	3	3	5	3	3	-	-
Male	-	4	-	2	-	1	2	1	2	-	-	-
Female	-	1	1	-	2	2	1	4	1	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	54	64	69	102	118	200	336	419	371	167	38	-
Male	32	44	41	58	58	94	112	126	92	30	5	-
Female	22	20	28	44	60	106	224	293	279	137	33	-
White, both sexes	37	48	51	80	88	154	279	351	316	141	33	-
Male	21	36	28	42	43	75	95	98	80	21	4	-
Female	16	12	23	38	45	79	184	253	236	120	29	-
Black, both sexes	13	13	10	11	15	26	26	38	24	11	4	-
Male	8	7	6	7	8	13	7	13	2	4	-	-
Female	5	6	4	4	7	13	19	25	22	7	4	-
American Indian 1/, both sexes	-	-	2	1	-	-	-	3	1	-	-	-
Male	-	-	1	1	-	-	-	2	-	-	-	-
Female	-	-	1	-	-	-	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	4	3	6	10	15	20	31	27	30	15	1	-
Male	3	1	6	8	7	6	10	13	10	5	1	-
Female	1	2	-	2	8	14	21	14	20	10	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Colorado												
All races, both sexes	1	4	7	6	6	13	17	27	23	12	4	-
Male	1	3	5	3	2	7	3	11	6	2	-	-
Female	-	1	2	3	4	6	14	16	17	10	4	-
White, both sexes	-	3	7	6	6	13	17	27	22	12	4	-
Male	-	2	5	3	2	7	3	11	6	2	-	-
Female	-	1	2	3	4	6	14	16	16	10	4	-
Black, both sexes	1	1	-	-	-	-	-	-	1	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	5	11	9	13	13	25	30	42	38	20	9	-
Male	4	9	7	7	3	13	8	11	6	1	1	-
Female	1	2	2	6	10	12	22	31	32	19	8	-
White, both sexes	4	7	7	11	13	25	27	37	37	19	9	-
Male	3	5	6	6	3	13	7	11	5	1	1	-
Female	1	2	1	5	10	12	20	26	32	18	8	-
Black, both sexes	1	4	2	2	-	-	3	5	-	1	-	-
Male	1	4	1	1	-	-	1	-	-	-	-	-
Female	-	-	1	1	-	-	2	5	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Delaware												
All races, both sexes	-	-	-	3	1	1	3	7	4	1	1	-
Male	-	-	-	2	1	-	1	1	1	-	-	-
Female	-	-	-	1	-	1	2	6	3	1	1	-
White, both sexes	-	-	-	2	1	-	3	6	4	-	1	-
Male	-	-	-	1	1	-	1	1	1	-	-	-
Female	-	-	-	1	-	-	2	5	3	-	1	-
Black, both sexes	-	-	-	1	-	1	-	1	-	1	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	2	1	3	2	4	2	4	4	2	-
Male	-	-	1	1	2	-	2	-	2	1	-	-
Female	-	-	1	-	1	2	2	2	2	3	2	-
White, both sexes	-	-	-	-	-	-	-	2	1	2	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	1	2	-	-
Black, both sexes	-	-	2	1	2	2	4	-	2	2	2	-
Male	-	-	1	1	1	-	2	-	2	1	-	-
Female	-	-	1	-	1	2	2	-	-	1	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Florida												
All races, both sexes	34	35	34	59	81	109	163	193	145	73	17	-
Male	22	21	22	33	34	60	64	57	33	16	-	-
Female	12	14	12	26	47	49	99	136	112	57	17	-
White, both sexes	27	27	25	43	64	88	145	174	134	64	12	-
Male	19	17	16	24	26	52	56	52	30	14	-	-
Female	8	10	9	19	38	36	89	122	104	50	12	-
Black, both sexes	7	8	8	16	15	19	17	18	11	9	5	-
Male	3	4	5	9	7	7	7	4	3	2	-	-
Female	4	4	3	7	8	12	10	14	8	7	5	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	2	1	1	1	-	-	-	-
Male	-	-	1	-	1	-	1	1	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Georgia												
All races, both sexes	43	60	56	53	47	44	66	103	79	38	10	1
Male	24	41	33	27	23	22	20	24	14	4	-	-
Female	19	19	23	26	24	22	46	79	65	34	10	1
White, both sexes	23	28	35	24	25	28	44	78	57	23	7	-
Male	15	22	21	13	11	13	12	18	11	2	-	-
Female	8	6	14	11	14	15	32	60	46	21	7	-
Black, both sexes	20	32	20	27	22	15	21	24	22	15	3	1
Male	9	19	12	13	12	9	7	6	3	2	-	-
Female	11	13	8	14	10	6	14	18	19	13	3	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	2	-	1	1	1	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	1	1	-	1	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Hawaii												
All races, both sexes	2	-	2	3	3	6	8	18	13	8	1	-
Male	1	-	2	3	-	3	3	8	6	2	1	-
Female	1	-	-	-	3	3	5	10	7	6	-	-
White, both sexes	-	-	-	2	2	2	-	5	2	1	-	-
Male	-	-	-	2	-	-	-	2	2	-	-	-
Female	-	-	-	-	2	2	-	3	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	2	1	1	4	8	13	11	7	1	-
Male	1	-	2	1	-	3	3	6	4	2	1	-
Female	1	-	-	-	1	1	5	7	7	5	-	-
Idaho												
All races, both sexes	1	1	1	3	3	-	4	9	8	3	1	-
Male	-	-	1	-	1	-	2	2	1	1	-	-
Female	1	1	-	3	2	-	2	7	7	2	1	-
White, both sexes	1	1	1	3	3	-	4	9	8	3	1	-
Male	-	-	1	-	1	-	2	2	1	1	-	-
Female	1	1	-	3	2	-	2	7	7	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Illinois												
All races, both sexes	15	21	14	16	35	51	78	107	93	58	15	-
Male	8	9	7	8	19	20	29	24	18	5	-	-
Female	7	12	7	8	16	31	49	83	75	53	15	-
White, both sexes	10	13	10	12	22	40	61	94	84	52	13	-
Male	5	7	5	6	13	16	26	20	17	5	-	-
Female	5	6	5	6	9	24	35	74	67	47	13	-
Black, both sexes	5	8	4	4	11	9	14	12	7	5	2	-
Male	3	2	2	2	5	4	2	4	1	-	-	-
Female	2	6	2	2	6	5	12	8	6	5	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	2	3	1	2	1	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	1	2	2	1	2	1	-	-
Indiana												
All races, both sexes	11	17	9	17	15	37	49	67	58	23	4	-
Male	7	12	7	7	9	16	19	18	10	-	-	-
Female	4	5	2	10	6	21	30	49	48	23	4	-
White, both sexes	10	14	8	14	15	34	46	65	56	21	4	-
Male	6	12	6	6	9	15	18	17	9	-	-	-
Female	4	2	2	8	6	19	28	48	47	21	4	-
Black, both sexes	1	3	1	2	-	3	3	2	2	2	-	-
Male	1	-	1	1	-	1	1	1	1	-	-	-
Female	-	3	-	1	-	2	2	1	1	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Iowa												
All races, both sexes	1	-	3	5	4	8	20	29	20	10	3	-
Male	1	-	3	3	1	3	11	7	2	-	-	-
Female	-	-	-	2	3	5	9	22	18	10	3	-
White, both sexes	1	-	3	5	4	8	19	29	20	10	3	-
Male	1	-	3	3	1	3	10	7	2	-	-	-
Female	-	-	-	2	3	5	9	22	18	10	3	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	4	7	6	4	9	15	18	25	14	12	-
Male	-	3	5	2	3	6	4	2	4	2	2	-
Female	-	1	2	4	1	3	11	16	21	12	10	-
White, both sexes	-	4	6	5	4	8	14	17	22	13	11	-
Male	-	3	4	2	3	5	4	2	4	2	2	-
Female	-	1	2	3	1	3	10	15	18	11	9	-
Black, both sexes	-	-	1	1	-	1	1	1	3	-	1	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	1	3	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Kentucky												
All races, both sexes	6	10	6	12	12	30	33	30	27	12	6	-
Male	3	5	3	7	4	10	10	10	4	4	-	-
Female	3	5	3	5	8	20	23	20	23	8	6	-
White, both sexes	4	8	6	11	11	29	30	26	24	11	3	-
Male	3	4	3	7	3	9	10	9	4	4	-	-
Female	1	4	3	4	8	20	20	17	20	7	3	-
Black, both sexes	2	2	-	1	1	1	3	4	3	1	3	-
Male	-	1	-	-	1	1	-	1	-	-	-	-
Female	2	1	-	1	-	-	3	3	3	1	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	14	11	15	15	20	25	34	34	37	10	5	-
Male	8	9	7	10	12	10	14	8	5	1	-	-
Female	6	2	8	5	8	15	20	26	32	9	5	-
White, both sexes	4	5	6	6	11	15	26	26	30	7	3	-
Male	2	4	3	4	7	6	10	6	4	1	-	-
Female	2	1	3	2	4	9	16	20	26	6	3	-
Black, both sexes	10	6	9	8	9	10	8	8	7	3	2	-
Male	6	5	4	5	5	4	4	2	1	-	-	-
Female	4	1	5	3	4	6	4	6	6	3	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Maine												
All races, both sexes	1	3	3	5	3	2	5	3	8	5	4	-
Male	-	3	2	3	-	1	-	1	2	-	-	-
Female	1	-	1	2	3	1	5	2	6	5	4	-
White, both sexes	1	3	3	5	3	2	5	3	8	5	4	-
Male	-	3	2	3	-	1	-	1	2	-	-	-
Female	1	-	1	2	3	1	5	2	6	5	4	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	9	7	10	9	12	22	30	40	40	19	3	-
Male	6	4	4	3	5	13	9	9	6	4	-	-
Female	3	3	6	6	7	9	21	31	34	15	3	-
White, both sexes	5	3	4	5	4	13	26	29	34	15	2	-
Male	4	3	-	1	2	7	9	6	4	3	-	-
Female	1	-	4	4	2	6	17	23	30	12	2	-
Black, both sexes	4	4	5	4	7	8	4	8	6	4	1	-
Male	2	1	3	2	3	5	-	2	2	1	-	-
Female	2	3	2	2	4	3	4	6	4	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	-	-	3	-	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Massachusetts												
All races, both sexes	9	7	13	19	17	33	51	76	61	28	7	-
Male	6	3	9	11	12	12	14	20	14	3	-	-
Female	3	4	4	8	5	21	37	56	47	25	7	-
White, both sexes	9	6	11	16	17	30	48	73	57	28	7	-
Male	6	2	9	11	12	10	13	19	13	3	-	-
Female	3	4	2	5	5	20	35	54	44	25	7	-
Black, both sexes	-	1	1	2	-	3	3	2	3	-	-	-
Male	-	1	-	-	-	2	1	1	-	-	-	-
Female	-	-	1	2	-	1	2	1	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	1	1	-	-	-	1	-	-	-	-
Michigan												
All races, both sexes	13	15	14	23	26	49	74	109	106	41	10	-
Male	8	11	8	14	8	22	26	31	33	8	-	-
Female	5	4	6	9	18	27	48	78	73	33	10	-
White, both sexes	10	10	7	19	20	42	65	100	97	37	9	-
Male	5	8	6	11	6	18	24	28	31	7	-	-
Female	5	2	1	8	14	24	41	72	66	30	9	-
Black, both sexes	3	4	7	4	5	7	5	8	9	4	1	-
Male	3	2	2	3	2	4	2	2	2	1	-	-
Female	-	2	5	1	3	3	3	6	7	3	1	-
American Indian 1/, both sexes	-	-	-	-	1	-	3	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	3	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Minnesota												
All races, both sexes	2	18	16	14	19	28	33	52	67	25	8	-
Male	-	16	11	11	10	13	14	13	13	4	2	-
Female	2	2	5	3	9	15	19	39	54	21	6	-
White, both sexes	1	16	16	13	18	27	32	52	66	25	7	-
Male	-	15	11	10	10	13	13	13	13	4	2	-
Female	1	1	5	3	8	14	19	39	53	21	5	-
Black, both sexes	1	2	-	1	1	1	1	-	1	-	1	-
Male	-	1	-	1	-	-	1	-	-	-	-	-
Female	1	1	-	-	1	1	-	-	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	12	12	25	9	24	32	45	44	30	20	5	-
Male	9	8	16	4	14	16	10	8	4	3	-	-
Female	3	4	9	5	10	16	35	36	26	17	5	-
White, both sexes	6	5	9	4	11	17	33	28	20	13	3	-
Male	4	4	5	2	8	8	7	4	3	1	-	-
Female	2	1	4	2	3	9	26	24	17	12	3	-
Black, both sexes	6	7	15	5	13	15	12	16	10	7	2	-
Male	5	4	10	2	6	8	3	4	1	2	-	-
Female	1	3	5	3	7	7	9	12	9	5	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Missouri												
All races, both sexes	7	11	9	15	14	24	38	35	34	23	4	-
Male	6	7	7	9	4	11	12	7	4	2	-	-
Female	1	4	2	6	10	13	26	28	30	21	4	-
White, both sexes	6	8	6	10	12	20	32	34	32	20	4	-
Male	5	7	5	7	3	8	11	7	4	1	-	-
Female	1	1	1	3	9	12	21	27	28	19	4	-
Black, both sexes	1	3	3	5	2	4	6	1	2	3	-	-
Male	1	-	2	2	1	3	1	-	-	1	-	-
Female	-	3	1	3	1	1	5	1	2	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	2	1	1	2	4	3	5	15	8	13	1	-
Male	2	1	1	2	3	-	2	3	2	3	-	-
Female	-	-	-	-	1	3	3	12	6	10	1	-
White, both sexes	2	1	1	2	4	3	4	15	8	13	1	-
Male	2	1	1	2	3	-	2	3	2	3	-	-
Female	-	-	-	-	1	3	2	12	6	10	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Nebraska												
All races, both sexes	-	2	3	4	6	20	10	21	35	14	6	-
Male	-	1	2	3	4	11	2	4	3	3	-	-
Female	-	1	1	1	2	9	8	17	32	11	6	-
White, both sexes	-	2	2	3	4	20	10	21	35	14	6	-
Male	-	1	1	2	3	11	2	4	3	3	-	-
Female	-	1	1	1	1	9	8	17	32	11	6	-
Black, both sexes	-	-	1	1	2	-	-	-	-	-	-	-
Male	-	-	1	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	5	3	8	6	7	12	14	13	14	4	1	-
Male	2	2	4	3	4	6	7	4	2	3	-	-
Female	3	1	4	3	3	6	7	9	12	1	1	-
White, both sexes	4	3	8	5	6	10	13	11	14	3	1	-
Male	2	2	4	2	4	6	7	3	2	2	-	-
Female	2	1	4	3	2	4	6	8	12	1	1	-
Black, both sexes	1	-	-	1	-	1	1	1	-	1	-	-
Male	-	-	-	1	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
New Hampshire												
All races, both sexes	-	2	3	2	2	4	6	7	8	3	1	-
Male	-	2	3	2	1	1	2	1	1	2	-	-
Female	-	-	-	-	1	3	4	6	7	1	1	-
White, both sexes	-	2	3	2	2	4	6	7	8	3	1	-
Male	-	2	3	2	1	1	2	1	1	2	-	-
Female	-	-	-	-	1	3	4	6	7	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	7	15	28	22	23	37	36	65	50	31	14	-
Male	4	10	19	13	10	11	14	20	11	4	1	-
Female	3	5	9	9	13	26	22	45	39	27	13	-
White, both sexes	7	9	17	18	14	29	28	60	49	27	12	-
Male	4	7	12	12	6	8	11	19	11	4	1	-
Female	3	2	5	6	8	21	17	41	38	23	11	-
Black, both sexes	-	5	11	3	9	8	6	3	1	4	2	-
Male	-	3	7	-	4	3	2	1	-	-	-	-
Female	-	2	4	3	5	5	4	2	1	4	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	-	2	2	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	1	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
New Mexico												
All races, both sexes	1	3	1	1	8	2	5	12	13	3	1	-
Male	1	1	1	-	5	1	1	3	3	1	-	-
Female	-	2	-	1	3	1	4	9	10	2	1	-
White, both sexes	1	3	-	1	5	1	4	12	13	3	1	-
Male	1	1	-	-	3	-	1	3	3	1	-	-
Female	-	2	-	1	2	1	3	9	10	2	1	-
Black, both sexes	-	-	-	-	2	1	-	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	1	-	1	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	32	33	58	49	74	116	158	188	172	89	21	-
Male	16	22	34	27	43	56	59	58	37	14	1	-
Female	16	11	24	22	31	60	99	130	135	75	20	-
White, both sexes	19	21	40	31	54	87	127	155	147	71	17	-
Male	10	18	24	15	34	43	46	48	33	11	-	-
Female	9	3	16	16	20	44	81	107	114	60	17	-
Black, both sexes	12	11	17	15	18	27	26	27	19	14	1	-
Male	5	3	9	10	7	13	9	8	3	1	-	-
Female	7	8	8	5	11	14	17	19	16	13	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	3	2	2	5	6	5	4	3	-
Male	1	1	1	2	2	-	4	2	1	2	1	-
Female	-	-	-	1	-	2	1	4	4	2	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
North Carolina												
All races, both sexes	15	19	19	16	20	42	55	70	57	21	12	-
Male	8	13	11	8	7	22	9	24	11	4	-	-
Female	7	6	8	8	13	20	46	46	46	17	12	-
White, both sexes	9	8	10	9	11	27	38	50	44	10	10	-
Male	6	6	7	4	3	14	6	18	7	4	-	-
Female	3	2	3	5	8	13	32	32	37	6	10	-
Black, both sexes	6	10	8	6	9	15	15	20	13	11	2	-
Male	2	6	4	4	4	8	3	6	4	-	-	-
Female	4	4	4	2	5	7	12	14	9	11	2	-
American Indian 1/, both sexes	-	1	1	1	-	-	2	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	1	-	-	2	1	-	7	9	7	5	1	-
Male	1	-	-	-	-	-	4	2	-	-	-	-
Female	-	-	-	2	1	-	3	7	7	5	1	-
White, both sexes	1	-	-	1	1	-	7	8	7	5	1	-
Male	1	-	-	-	-	-	4	2	-	-	-	-
Female	-	-	-	1	1	-	3	6	7	5	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Ohio												
All races, both sexes	11	19	26	33	54	71	122	130	130	59	14	-
Male	8	12	15	14	25	28	34	34	29	7	-	-
Female	3	7	11	19	29	43	88	96	101	52	14	-
White, both sexes	9	17	19	23	50	61	106	119	122	54	11	-
Male	7	11	11	10	25	26	31	30	22	6	-	-
Female	2	6	8	13	25	35	75	89	100	48	11	-
Black, both sexes	2	2	7	10	3	10	14	11	8	5	3	-
Male	1	1	4	4	-	2	2	4	7	1	-	-
Female	1	1	3	6	3	8	12	7	1	4	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Oklahoma												
All races, both sexes	2	6	5	11	6	22	29	34	23	17	2	-
Male	2	4	3	6	2	9	12	6	5	-	-	-
Female	-	2	2	5	4	13	17	28	18	17	2	-
White, both sexes	1	4	2	9	5	19	24	29	23	16	2	-
Male	1	3	1	5	2	7	10	5	5	-	-	-
Female	-	1	1	4	3	12	14	24	18	16	2	-
Black, both sexes	1	2	1	2	1	2	1	5	-	1	-	-
Male	1	1	-	1	-	1	-	1	-	-	-	-
Female	-	1	1	1	1	1	1	4	-	1	-	-
American Indian 1/, both sexes	-	-	2	-	-	1	4	-	-	-	-	-
Male	-	-	2	-	-	1	2	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Oregon												
All races, both sexes	9	13	12	14	14	26	33	52	34	30	3	-
Male	8	8	10	9	8	14	12	13	10	4	1	-
Female	1	5	2	5	6	12	21	39	24	26	2	-
White, both sexes	9	12	10	14	14	24	33	51	32	30	3	-
Male	8	8	8	9	8	13	12	12	8	4	1	-
Female	1	4	2	5	6	11	21	39	24	26	2	-
Black, both sexes	-	1	1	-	-	1	-	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	1	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	-	1	1	-	-	-
Male	-	-	1	-	-	1	-	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	15	20	46	29	44	65	105	121	108	47	17	-
Male	13	11	32	18	26	29	41	28	20	7	-	-
Female	2	9	14	11	18	36	64	93	88	40	17	-
White, both sexes	14	16	40	25	40	57	100	112	99	45	17	-
Male	12	10	28	16	22	27	39	25	18	7	-	-
Female	2	6	12	9	18	30	61	87	81	38	17	-
Black, both sexes	1	4	4	4	3	8	5	9	9	2	-	-
Male	1	1	3	2	3	2	2	3	2	-	-	-
Female	-	3	1	2	-	6	3	6	7	2	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Rhode Island												
All races, both sexes	-	1	-	2	2	4	5	9	6	1	-	-
Male	-	1	-	1	1	1	3	1	1	-	-	-
Female	-	-	-	1	1	3	2	8	5	1	-	-
White, both sexes	-	1	-	1	1	4	5	9	6	1	-	-
Male	-	1	-	1	-	1	3	1	1	-	-	-
Female	-	-	-	-	1	3	2	8	5	1	-	-
Black, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	8	11	19	11	14	22	33	36	32	16	8	-
Male	6	7	14	9	7	15	12	10	10	2	-	-
Female	2	4	5	2	7	7	21	26	22	14	8	-
White, both sexes	3	1	8	2	9	13	21	29	23	10	6	-
Male	3	-	6	2	4	8	8	7	7	2	-	-
Female	-	1	2	-	5	5	13	22	16	8	6	-
Black, both sexes	5	10	11	9	5	9	12	7	9	6	2	-
Male	3	7	8	7	3	7	4	3	3	-	-	-
Female	2	3	3	2	2	2	8	4	6	6	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
South Dakota												
All races, both sexes	2	1	-	-	1	6	8	8	6	3	4	-
Male	2	-	-	-	-	3	2	1	1	-	-	-
Female	-	1	-	-	1	3	6	7	5	3	4	-
White, both sexes	1	1	-	-	1	6	8	8	6	3	4	-
Male	1	-	-	-	-	3	2	1	1	-	-	-
Female	-	1	-	-	1	3	6	7	5	3	4	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	16	10	15	15	17	24	40	61	35	22	4	-
Male	10	5	6	6	10	9	16	15	7	5	-	-
Female	6	5	9	9	7	15	24	46	28	17	4	-
White, both sexes	9	4	11	12	9	19	35	53	31	20	2	-
Male	6	3	5	5	5	6	12	13	7	4	-	-
Female	3	1	6	7	4	13	23	40	24	16	2	-
Black, both sexes	7	6	4	3	7	5	5	8	4	2	2	-
Male	4	2	1	1	4	3	4	2	-	1	-	-
Female	3	4	3	2	3	2	1	6	4	1	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Texas												
All races, both sexes	25	35	45	54	51	68	126	168	121	78	20	-
Male	18	18	23	34	20	33	41	50	27	10	6	-
Female	7	17	22	20	31	35	85	118	94	68	14	-
White, both sexes	17	23	31	50	45	63	102	147	104	73	18	-
Male	13	14	13	32	19	32	32	43	25	8	5	-
Female	4	9	18	18	26	31	70	104	79	65	13	-
Black, both sexes	7	12	14	4	5	5	20	19	17	4	2	-
Male	4	4	10	2	1	1	8	7	2	2	1	-
Female	3	8	4	2	4	4	12	12	15	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	3	1	-	1	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	2	1	-	1	-	-
Utah												
All races, both sexes	1	3	5	4	2	12	16	21	22	9	2	-
Male	1	1	5	1	-	4	7	9	5	1	-	-
Female	-	2	-	3	2	8	9	12	17	8	2	-
White, both sexes	1	3	5	4	2	12	16	20	22	9	2	-
Male	1	1	5	1	-	4	7	9	5	1	-	-
Female	-	2	-	3	2	8	9	11	17	8	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Vermont												
All races, both sexes	-	1	1	1	-	4	5	4	2	4	1	-
Male	-	1	1	1	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	3	4	4	2	4	1	-
White, both sexes	-	1	1	1	-	4	5	4	2	4	1	-
Male	-	1	1	1	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	3	4	4	2	4	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	15	20	20	24	18	24	44	54	50	20	4	-
Male	8	14	12	14	6	14	15	17	6	1	1	-
Female	7	6	8	10	12	10	29	37	44	19	3	-
White, both sexes	10	12	10	16	10	20	29	49	35	17	3	-
Male	5	10	8	9	4	12	9	17	4	1	-	-
Female	5	2	2	7	6	8	20	32	31	16	3	-
Black, both sexes	5	8	10	7	7	4	12	5	11	3	1	-
Male	3	4	4	4	2	2	5	-	1	-	1	-
Female	2	4	6	3	5	2	7	5	10	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	3	-	3	-	-	-
Male	-	-	-	1	-	-	1	-	1	-	-	-
Female	-	-	-	-	1	-	2	-	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Washington												
All races, both sexes	6	9	15	10	14	13	41	50	49	23	7	-
Male	6	5	14	2	8	7	16	14	8	7	-	-
Female	-	4	1	8	6	6	25	36	41	16	7	-
White, both sexes	2	9	14	9	13	12	37	48	48	23	7	-
Male	2	5	13	2	8	6	14	13	8	7	-	-
Female	-	4	1	7	5	6	23	35	40	16	7	-
Black, both sexes	3	-	-	-	1	1	1	-	-	-	-	-
Male	3	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	1	-	-	-	1	-	-	-	-
Male	1	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	3	1	1	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	1	-	-	-
West Virginia												
All races, both sexes	11	15	11	12	12	11	16	30	25	11	1	-
Male	7	11	9	6	5	4	7	8	5	3	-	-
Female	4	4	2	6	7	7	9	22	20	8	1	-
White, both sexes	11	13	11	10	12	10	15	28	24	11	1	-
Male	7	10	9	5	5	4	6	7	5	3	-	-
Female	4	3	2	5	7	6	9	21	19	8	1	-
Black, both sexes	-	2	-	2	-	1	1	2	1	-	-	-
Male	-	1	-	1	-	-	1	1	-	-	-	-
Female	-	1	-	1	-	1	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Wisconsin												
All races, both sexes	2	8	14	6	13	18	30	50	46	30	13	-
Male	2	6	7	3	8	11	4	12	11	4	2	-
Female	-	2	7	3	5	7	26	38	35	26	11	-
White, both sexes	2	7	13	5	12	17	29	49	45	30	13	-
Male	2	5	7	3	7	10	4	12	10	4	2	-
Female	-	2	6	2	5	7	25	37	35	26	11	-
Black, both sexes	-	1	1	-	-	1	1	1	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	2	1	-	1	1	2	1	3	3	5	1	-
Male	1	1	-	1	-	1	-	3	-	-	-	-
Female	1	-	-	-	1	1	1	-	3	5	1	-
White, both sexes	2	1	-	1	1	2	1	3	3	5	1	-
Male	1	1	-	1	-	1	-	3	-	-	-	-
Female	1	-	-	-	1	1	1	-	3	5	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
United States												
All races, both sexes	30,780	1	1	-	-	5	39	115	223	438	934	1,556
Male	14,249	1	1	-	-	4	30	81	156	317	636	1,072
Female	16,531	-	-	-	-	1	9	34	67	121	298	484
White, both sexes	22,938	1	1	-	-	1	15	61	103	229	542	983
Male	10,209	1	1	-	-	1	14	42	76	180	387	710
Female	12,729	-	-	-	-	-	1	19	27	49	155	273
Black, both sexes	7,108	-	-	-	-	4	24	47	115	194	370	538
Male	3,669	-	-	-	-	3	16	34	75	129	234	330
Female	3,439	-	-	-	-	1	8	13	40	65	136	208
American Indian 1/, both sexes	135	-	-	-	-	-	-	3	2	6	11	13
Male	79	-	-	-	-	-	-	3	2	4	6	11
Female	56	-	-	-	-	-	-	-	-	2	5	2
Asian or Pacific Islander 2/, both sexes	599	-	-	-	-	-	-	4	3	9	11	22
Male	292	-	-	-	-	-	-	2	3	4	9	21
Female	307	-	-	-	-	-	-	2	-	5	2	1
Alabama												
All races, both sexes	392	-	-	-	-	-	3	5	2	13	13	22
Male	172	-	-	-	-	-	2	5	1	7	10	14
Female	220	-	-	-	-	-	1	-	1	6	3	8
White, both sexes	239	-	-	-	-	-	-	1	-	3	4	13
Male	92	-	-	-	-	-	-	1	-	3	2	8
Female	147	-	-	-	-	-	-	-	-	-	2	5
Black, both sexes	152	-	-	-	-	-	3	3	2	10	9	9
Male	79	-	-	-	-	-	2	3	1	4	8	6
Female	73	-	-	-	-	-	1	-	1	6	1	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Alaska												
All races, both sexes	62	-	-	-	-	-	-	-	1	-	-	6
Male	38	-	-	-	-	-	-	-	1	-	-	5
Female	24	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	45	-	-	-	-	-	-	-	-	-	-	5
Male	29	-	-	-	-	-	-	-	-	-	-	4
Female	16	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	1	-	-	1
Male	6	-	-	-	-	-	-	-	1	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	662	-	-	-	-	-	1	2	2	8	18	31
Male	341	-	-	-	-	-	1	1	2	6	10	25
Female	321	-	-	-	-	-	-	1	-	2	8	6
White, both sexes	600	-	-	-	-	-	1	2	2	5	10	28
Male	307	-	-	-	-	-	1	1	2	4	7	22
Female	293	-	-	-	-	-	-	1	-	1	3	6
Black, both sexes	35	-	-	-	-	-	-	-	-	1	3	3
Male	21	-	-	-	-	-	-	-	-	1	1	3
Female	14	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	22	-	-	-	-	-	-	-	-	2	5	-
Male	10	-	-	-	-	-	-	-	-	1	2	-
Female	12	-	-	-	-	-	-	-	-	1	3	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Arkansas												
All races, both sexes	276	-	-	-	-	-	-	3	1	1	12	16
Male	117	-	-	-	-	-	-	1	-	-	9	13
Female	159	-	-	-	-	-	-	2	1	1	3	3
White, both sexes	223	-	-	-	-	-	-	1	-	1	5	11
Male	89	-	-	-	-	-	-	-	-	-	3	9
Female	134	-	-	-	-	-	-	1	-	1	2	2
Black, both sexes	53	-	-	-	-	-	-	2	1	-	7	5
Male	28	-	-	-	-	-	-	1	-	-	6	4
Female	25	-	-	-	-	-	-	1	1	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	3,946	-	-	-	-	-	1	11	22	50	112	173
Male	1,824	-	-	-	-	-	1	8	14	32	79	126
Female	2,122	-	-	-	-	-	-	3	8	18	33	47
White, both sexes	3,073	-	-	-	-	-	1	8	14	27	83	114
Male	1,394	-	-	-	-	-	1	6	10	19	60	84
Female	1,679	-	-	-	-	-	-	2	4	8	23	30
Black, both sexes	556	-	-	-	-	-	-	-	6	16	23	51
Male	276	-	-	-	-	-	-	-	2	11	14	35
Female	280	-	-	-	-	-	-	-	4	5	9	16
American Indian 1/, both sexes	20	-	-	-	-	-	-	1	-	-	1	-
Male	12	-	-	-	-	-	-	1	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	297	-	-	-	-	-	-	2	2	7	5	8
Male	142	-	-	-	-	-	-	1	2	2	4	7
Female	155	-	-	-	-	-	-	1	-	5	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Colorado												
All races, both sexes	286	-	-	-	-	-	-	2	1	3	6	13
Male	132	-	-	-	-	-	-	2	1	3	4	12
Female	154	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	265	-	-	-	-	-	-	1	1	3	6	11
Male	124	-	-	-	-	-	-	1	1	3	4	11
Female	141	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	19	-	-	-	-	-	-	1	-	-	-	2
Male	8	-	-	-	-	-	-	1	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	230	-	-	-	-	-	2	-	-	-	6	9
Male	97	-	-	-	-	-	1	-	-	-	5	5
Female	133	-	-	-	-	-	1	-	-	-	1	4
White, both sexes	194	-	-	-	-	-	1	-	-	-	4	7
Male	76	-	-	-	-	-	1	-	-	-	3	4
Female	118	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	35	-	-	-	-	-	1	-	-	-	2	2
Male	20	-	-	-	-	-	-	-	-	-	2	1
Female	15	-	-	-	-	-	1	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Delaware												
All races, both sexes	73	-	-	-	-	-	-	1	-	5	4	4
Male	37	-	-	-	-	-	-	-	-	4	2	3
Female	36	-	-	-	-	-	-	1	-	1	2	1
White, both sexes	56	-	-	-	-	-	-	-	-	3	3	4
Male	25	-	-	-	-	-	-	-	-	2	1	3
Female	31	-	-	-	-	-	-	-	-	1	2	1
Black, both sexes	16	-	-	-	-	-	-	1	-	1	1	-
Male	11	-	-	-	-	-	-	-	-	1	1	-
Female	5	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	133	-	-	-	-	-	-	1	4	1	10	6
Male	66	-	-	-	-	-	-	1	2	1	5	5
Female	67	-	-	-	-	-	-	-	2	-	5	1
White, both sexes	17	-	-	-	-	-	-	1	-	-	-	-
Male	4	-	-	-	-	-	-	1	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	116	-	-	-	-	-	-	-	4	1	10	6
Male	62	-	-	-	-	-	-	-	2	1	5	5
Female	54	-	-	-	-	-	-	-	2	-	5	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Florida												
All races, both sexes	2,357	-	-	-	-	-	4	8	20	23	88	112
Male	1,163	-	-	-	-	-	4	8	16	18	59	82
Female	1,194	-	-	-	-	-	-	-	4	5	29	30
White, both sexes	1,883	-	-	-	-	-	2	8	10	14	54	74
Male	913	-	-	-	-	-	2	8	8	10	40	57
Female	970	-	-	-	-	-	-	-	2	4	14	17
Black, both sexes	454	-	-	-	-	-	2	-	10	9	32	38
Male	239	-	-	-	-	-	2	-	8	8	17	25
Female	215	-	-	-	-	-	-	-	2	1	15	13
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	18	-	-	-	-	-	-	-	-	-	1	-
Male	9	-	-	-	-	-	-	-	-	-	1	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	1,202	-	-	-	-	1	2	3	8	25	49	81
Male	628	-	-	-	-	-	2	3	5	18	31	58
Female	574	-	-	-	-	1	-	-	3	7	18	23
White, both sexes	607	-	-	-	-	-	1	1	3	7	17	40
Male	301	-	-	-	-	-	1	1	2	5	8	31
Female	306	-	-	-	-	-	-	-	1	2	9	9
Black, both sexes	591	-	-	-	-	1	1	2	5	18	32	41
Male	324	-	-	-	-	-	1	2	3	13	23	27
Female	267	-	-	-	-	1	-	-	2	5	9	14
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Hawaii												
All races, both sexes	60	-	-	-	-	-	-	-	-	-	-	2
Male	24	-	-	-	-	-	-	-	-	-	-	2
Female	36	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	21	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	39	-	-	-	-	-	-	-	-	-	-	1
Male	17	-	-	-	-	-	-	-	-	-	-	1
Female	22	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	108	-	-	-	-	-	-	-	1	4	3	5
Male	44	-	-	-	-	-	-	-	-	2	2	4
Female	64	-	-	-	-	-	-	-	1	2	1	1
White, both sexes	107	-	-	-	-	-	-	-	1	3	3	5
Male	43	-	-	-	-	-	-	-	-	1	2	4
Female	64	-	-	-	-	-	-	-	1	2	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Illinois												
All races, both sexes	1,396	-	-	-	-	-	4	7	17	29	47	92
Male	697	-	-	-	-	-	3	6	13	21	25	66
Female	699	-	-	-	-	-	1	1	4	8	22	26
White, both sexes	901	-	-	-	-	-	3	2	4	11	21	54
Male	416	-	-	-	-	-	3	1	4	8	10	44
Female	485	-	-	-	-	-	-	1	-	3	11	10
Black, both sexes	473	-	-	-	-	-	1	5	12	18	26	38
Male	268	-	-	-	-	-	-	5	8	13	15	22
Female	205	-	-	-	-	-	1	-	4	5	11	16
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	21	-	-	-	-	-	-	-	1	-	-	-
Male	12	-	-	-	-	-	-	-	1	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	447	-	-	-	-	-	-	2	1	3	10	24
Male	192	-	-	-	-	-	-	-	1	1	8	20
Female	255	-	-	-	-	-	-	2	-	2	2	4
White, both sexes	370	-	-	-	-	-	-	2	-	2	6	15
Male	152	-	-	-	-	-	-	-	-	1	6	12
Female	218	-	-	-	-	-	-	2	-	1	-	3
Black, both sexes	74	-	-	-	-	-	-	-	1	1	4	8
Male	39	-	-	-	-	-	-	-	1	-	2	7
Female	35	-	-	-	-	-	-	-	-	1	2	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Iowa												
All races, both sexes	229	-	-	-	-	-	1	-	1	2	3	1
Male	79	-	-	-	-	-	1	-	1	2	3	1
Female	150	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	221	-	-	-	-	-	1	-	1	1	3	1
Male	74	-	-	-	-	-	1	-	1	1	3	1
Female	147	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	148	-	-	-	-	-	-	-	-	3	4	2
Male	50	-	-	-	-	-	-	-	-	2	2	1
Female	98	-	-	-	-	-	-	-	-	1	2	1
White, both sexes	130	-	-	-	-	-	-	-	-	1	-	2
Male	44	-	-	-	-	-	-	-	-	1	-	1
Female	86	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	17	-	-	-	-	-	-	-	-	2	4	-
Male	6	-	-	-	-	-	-	-	-	1	2	-
Female	11	-	-	-	-	-	-	-	-	1	2	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Kentucky												
All races, both sexes	333	-	-	-	-	-	-	3	5	8	22	26
Male	161	-	-	-	-	-	-	1	3	5	19	12
Female	172	-	-	-	-	-	-	2	2	3	3	14
White, both sexes	301	-	-	-	-	-	-	1	3	7	19	24
Male	142	-	-	-	-	-	-	1	1	4	17	11
Female	159	-	-	-	-	-	-	-	2	3	2	13
Black, both sexes	30	-	-	-	-	-	-	2	2	1	3	2
Male	18	-	-	-	-	-	-	-	2	1	2	1
Female	12	-	-	-	-	-	-	2	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	889	-	-	-	-	-	-	4	6	19	28	43
Male	416	-	-	-	-	-	-	4	3	13	21	29
Female	473	-	-	-	-	-	-	-	3	6	7	14
White, both sexes	542	-	-	-	-	-	-	-	2	8	12	25
Male	260	-	-	-	-	-	-	-	2	7	10	19
Female	282	-	-	-	-	-	-	-	-	1	2	6
Black, both sexes	344	-	-	-	-	-	-	4	4	11	16	18
Male	154	-	-	-	-	-	-	4	1	6	11	10
Female	190	-	-	-	-	-	-	-	3	5	5	8
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Maine												
All races, both sexes	61	-	-	-	-	-	-	-	-	-	2	1
Male	32	-	-	-	-	-	-	-	-	-	2	1
Female	29	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	60	-	-	-	-	-	-	-	-	-	2	1
Male	31	-	-	-	-	-	-	-	-	-	2	1
Female	29	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	825	-	-	-	-	-	1	2	4	19	31	86
Male	455	-	-	-	-	-	1	2	4	13	21	59
Female	370	-	-	-	-	-	-	-	-	6	10	27
White, both sexes	488	-	-	-	-	-	-	1	2	9	15	45
Male	258	-	-	-	-	-	-	1	2	8	11	32
Female	230	-	-	-	-	-	-	-	-	1	4	13
Black, both sexes	324	-	-	-	-	-	1	1	2	10	16	40
Male	188	-	-	-	-	-	1	1	2	5	10	26
Female	136	-	-	-	-	-	-	-	-	5	6	14
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Massachusetts												
All races, both sexes	519	-	-	-	-	-	-	6	11	13	22	50
Male	257	-	-	-	-	-	-	3	8	11	19	34
Female	262	-	-	-	-	-	-	3	3	2	3	16
White, both sexes	470	-	-	-	-	-	-	3	7	10	19	41
Male	223	-	-	-	-	-	-	1	5	8	16	28
Female	247	-	-	-	-	-	-	2	2	2	3	13
Black, both sexes	40	-	-	-	-	-	-	1	4	2	3	6
Male	26	-	-	-	-	-	-	1	3	2	3	3
Female	14	-	-	-	-	-	-	-	1	-	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	2	-	1	-	3
Male	8	-	-	-	-	-	-	1	-	1	-	3
Female	1	-	-	-	-	-	-	1	-	-	-	-
Michigan												
All races, both sexes	1,371	-	-	-	-	-	-	6	10	21	35	55
Male	575	-	-	-	-	-	-	3	8	16	24	31
Female	796	-	-	-	-	-	-	3	2	5	11	24
White, both sexes	999	-	-	-	-	-	-	3	5	10	15	31
Male	390	-	-	-	-	-	-	2	5	8	8	17
Female	609	-	-	-	-	-	-	1	-	2	7	14
Black, both sexes	364	-	-	-	-	-	-	3	4	10	20	23
Male	178	-	-	-	-	-	-	1	2	7	16	13
Female	186	-	-	-	-	-	-	2	2	3	4	10
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	1	1	-	-
Male	5	-	-	-	-	-	-	-	1	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Minnesota												
All races, both sexes	184	-	-	-	-	-	1	1	-	2	3	5
Male	77	-	-	-	-	-	1	1	-	2	3	3
Female	107	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	165	-	-	-	-	-	-	-	-	1	2	4
Male	61	-	-	-	-	-	-	-	-	1	2	3
Female	104	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	14	-	-	-	-	-	1	-	-	1	1	-
Male	13	-	-	-	-	-	1	-	-	1	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	1	-	-	-	1
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	725	-	-	-	-	2	4	4	4	9	16	33
Male	336	-	-	-	-	2	3	3	1	7	10	21
Female	389	-	-	-	-	-	1	1	3	2	6	12
White, both sexes	400	-	-	-	-	1	-	-	2	1	2	18
Male	176	-	-	-	-	1	-	-	-	1	-	10
Female	224	-	-	-	-	-	-	-	2	-	2	8
Black, both sexes	323	-	-	-	-	1	4	4	2	8	13	15
Male	160	-	-	-	-	1	3	3	1	6	10	11
Female	163	-	-	-	-	-	1	1	1	2	3	4
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Missouri												
All races, both sexes	426	-	-	-	-	-	-	1	8	8	15	27
Male	183	-	-	-	-	-	-	1	4	7	7	13
Female	243	-	-	-	-	-	-	-	4	1	8	14
White, both sexes	273	-	-	-	-	-	-	-	2	4	7	7
Male	109	-	-	-	-	-	-	-	1	4	3	4
Female	164	-	-	-	-	-	-	-	1	-	4	3
Black, both sexes	149	-	-	-	-	-	-	1	6	4	7	20
Male	72	-	-	-	-	-	-	1	3	3	3	9
Female	77	-	-	-	-	-	-	-	3	1	4	11
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	54	-	-	-	-	-	-	-	-	-	-	5
Male	24	-	-	-	-	-	-	-	-	-	-	5
Female	30	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	54	-	-	-	-	-	-	-	-	-	-	5
Male	24	-	-	-	-	-	-	-	-	-	-	5
Female	30	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Nebraska												
All races, both sexes	127	-	-	-	-	-	-	-	-	-	7	8
Male	43	-	-	-	-	-	-	-	-	-	5	6
Female	84	-	-	-	-	-	-	-	-	-	2	2
White, both sexes	118	-	-	-	-	-	-	-	-	-	5	5
Male	39	-	-	-	-	-	-	-	-	-	5	3
Female	79	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	9	-	-	-	-	-	-	-	-	-	2	3
Male	4	-	-	-	-	-	-	-	-	-	-	3
Female	5	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	315	-	-	-	-	-	1	2	3	3	8	13
Male	168	-	-	-	-	-	-	2	3	2	5	7
Female	147	-	-	-	-	-	1	-	-	1	3	6
White, both sexes	266	-	-	-	-	-	-	2	3	2	3	10
Male	146	-	-	-	-	-	-	2	3	2	3	6
Female	120	-	-	-	-	-	-	-	-	-	-	4
Black, both sexes	41	-	-	-	-	-	1	-	-	1	4	3
Male	20	-	-	-	-	-	-	-	-	-	2	1
Female	21	-	-	-	-	-	1	-	-	1	2	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
New Hampshire												
All races, both sexes	113	-	-	-	-	-	-	-	-	1	3	7
Male	52	-	-	-	-	-	-	-	-	1	1	4
Female	61	-	-	-	-	-	-	-	-	-	2	3
White, both sexes	112	-	-	-	-	-	-	-	-	1	3	6
Male	51	-	-	-	-	-	-	-	-	1	1	3
Female	61	-	-	-	-	-	-	-	-	-	2	3
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	758	-	-	-	-	-	-	-	2	7	11	30
Male	340	-	-	-	-	-	-	-	2	5	8	23
Female	418	-	-	-	-	-	-	-	-	2	3	7
White, both sexes	577	-	-	-	-	-	-	-	-	3	4	14
Male	246	-	-	-	-	-	-	-	-	3	4	12
Female	331	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	169	-	-	-	-	-	-	-	2	4	7	14
Male	89	-	-	-	-	-	-	-	2	2	4	9
Female	80	-	-	-	-	-	-	-	-	2	3	5
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
New Mexico												
All races, both sexes	148	-	-	-	-	-	-	-	-	2	4	3
Male	63	-	-	-	-	-	-	-	-	1	4	2
Female	85	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	141	-	-	-	-	-	-	-	-	2	4	3
Male	57	-	-	-	-	-	-	-	-	1	4	2
Female	84	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	2,644	-	1	-	-	-	3	9	16	38	86	117
Male	1,263	-	1	-	-	-	2	6	11	24	65	86
Female	1,381	-	-	-	-	-	1	3	5	14	21	31
White, both sexes	1,798	-	1	-	-	-	1	5	5	19	46	66
Male	830	-	1	-	-	-	1	3	4	13	40	53
Female	968	-	-	-	-	-	-	2	1	6	6	13
Black, both sexes	789	-	-	-	-	-	2	4	11	18	40	50
Male	405	-	-	-	-	-	1	3	7	10	25	32
Female	384	-	-	-	-	-	1	1	4	8	15	18
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	56	-	-	-	-	-	-	-	-	1	-	1
Male	27	-	-	-	-	-	-	-	-	1	-	1
Female	29	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
North Carolina												
All races, both sexes	679	-	-	-	-	-	1	1	8	11	25	39
Male	320	-	-	-	-	-	-	-	8	7	12	22
Female	359	-	-	-	-	-	1	1	-	4	13	17
White, both sexes	407	-	-	-	-	-	-	-	4	6	12	19
Male	187	-	-	-	-	-	-	-	4	4	7	12
Female	220	-	-	-	-	-	-	-	-	2	5	7
Black, both sexes	260	-	-	-	-	-	1	1	4	5	12	19
Male	127	-	-	-	-	-	-	-	4	3	4	10
Female	133	-	-	-	-	-	1	1	-	2	8	9
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	55	-	-	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	38	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	54	-	-	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	37	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Ohio												
All races, both sexes	1,765	-	-	-	-	-	3	6	10	21	36	97
Male	792	-	-	-	-	-	2	4	7	16	24	68
Female	973	-	-	-	-	-	1	2	3	5	12	29
White, both sexes	1,379	-	-	-	-	-	-	4	5	12	20	73
Male	593	-	-	-	-	-	-	3	4	10	14	54
Female	786	-	-	-	-	-	-	1	1	2	6	19
Black, both sexes	376	-	-	-	-	-	3	2	5	9	15	22
Male	192	-	-	-	-	-	2	1	3	6	9	12
Female	184	-	-	-	-	-	1	1	2	3	6	10
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	1	2
Male	6	-	-	-	-	-	-	-	-	-	1	2
Female	3	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	307	-	-	-	-	-	-	-	1	4	10	16
Male	129	-	-	-	-	-	-	-	-	4	5	13
Female	178	-	-	-	-	-	-	-	1	-	5	3
White, both sexes	256	-	-	-	-	-	-	-	-	4	7	12
Male	106	-	-	-	-	-	-	-	-	4	3	10
Female	150	-	-	-	-	-	-	-	-	-	4	2
Black, both sexes	35	-	-	-	-	-	-	-	1	-	2	2
Male	13	-	-	-	-	-	-	-	-	-	1	1
Female	22	-	-	-	-	-	-	-	1	-	1	1
American Indian 1/, both sexes	14	-	-	-	-	-	-	-	-	-	1	2
Male	10	-	-	-	-	-	-	-	-	-	1	2
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Oregon												
All races, both sexes	241	-	-	-	-	-	-	-	1	4	-	8
Male	86	-	-	-	-	-	-	-	1	3	-	6
Female	155	-	-	-	-	-	-	-	-	1	-	2
White, both sexes	227	-	-	-	-	-	-	-	1	3	-	8
Male	78	-	-	-	-	-	-	-	1	2	-	6
Female	149	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	7	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	830	-	-	-	-	-	1	1	1	6	18	28
Male	358	-	-	-	-	-	1	-	-	3	14	21
Female	472	-	-	-	-	-	-	1	1	3	4	7
White, both sexes	664	-	-	-	-	-	1	1	1	1	14	16
Male	276	-	-	-	-	-	1	-	-	1	10	13
Female	388	-	-	-	-	-	-	1	1	-	4	3
Black, both sexes	163	-	-	-	-	-	-	-	-	5	4	12
Male	82	-	-	-	-	-	-	-	-	2	4	8
Female	81	-	-	-	-	-	-	-	-	3	-	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Rhode Island												
All races, both sexes	67	-	-	-	-	-	-	-	2	-	7	10
Male	28	-	-	-	-	-	-	-	-	-	5	8
Female	39	-	-	-	-	-	-	-	2	-	2	2
White, both sexes	62	-	-	-	-	-	-	-	2	-	6	8
Male	25	-	-	-	-	-	-	-	-	-	4	6
Female	37	-	-	-	-	-	-	-	2	-	2	2
Black, both sexes	3	-	-	-	-	-	-	-	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	1	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	374	-	-	-	-	-	-	-	2	3	5	13
Male	158	-	-	-	-	-	-	-	2	3	3	7
Female	216	-	-	-	-	-	-	-	-	-	2	6
White, both sexes	205	-	-	-	-	-	-	-	-	-	-	4
Male	84	-	-	-	-	-	-	-	-	-	-	3
Female	121	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	168	-	-	-	-	-	-	-	2	3	5	9
Male	73	-	-	-	-	-	-	-	2	3	3	4
Female	95	-	-	-	-	-	-	-	-	-	2	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
South Dakota												
All races, both sexes	49	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	36	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	47	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	34	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	876	-	-	-	-	-	1	6	13	15	23	38
Male	390	-	-	-	-	-	-	4	9	12	16	29
Female	486	-	-	-	-	-	1	2	4	3	7	9
White, both sexes	622	-	-	-	-	-	1	1	6	7	16	25
Male	256	-	-	-	-	-	-	1	4	6	12	18
Female	366	-	-	-	-	-	1	-	2	1	4	7
Black, both sexes	249	-	-	-	-	-	-	5	7	8	7	13
Male	131	-	-	-	-	-	-	3	5	6	4	11
Female	118	-	-	-	-	-	-	2	2	2	3	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Texas												
All races, both sexes	1,981	-	-	-	-	-	4	14	26	29	72	122
Male	917	-	-	-	-	-	4	8	18	23	49	73
Female	1,064	-	-	-	-	-	-	6	8	6	23	49
White, both sexes	1,563	-	-	-	-	-	2	10	13	21	52	82
Male	718	-	-	-	-	-	2	5	9	17	35	50
Female	845	-	-	-	-	-	-	5	4	4	17	32
Black, both sexes	385	-	-	-	-	-	2	4	13	8	20	37
Male	184	-	-	-	-	-	2	3	9	6	14	20
Female	201	-	-	-	-	-	-	1	4	2	6	17
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	3
Male	5	-	-	-	-	-	-	-	-	-	-	3
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	97	-	-	-	-	-	-	-	-	2	2	2
Male	37	-	-	-	-	-	-	-	-	2	2	1
Female	60	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	91	-	-	-	-	-	-	-	-	2	1	2
Male	35	-	-	-	-	-	-	-	-	2	1	1
Female	56	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Vermont												
All races, both sexes	59	-	-	-	-	-	-	-	-	-	1	3
Male	29	-	-	-	-	-	-	-	-	-	1	3
Female	30	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	59	-	-	-	-	-	-	-	-	-	1	3
Male	29	-	-	-	-	-	-	-	-	-	1	3
Female	30	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	445	1	-	-	-	1	-	2	4	15	25	28
Male	211	1	-	-	-	1	-	2	3	13	18	15
Female	234	-	-	-	-	-	-	-	1	2	7	13
White, both sexes	281	1	-	-	-	-	-	1	1	10	13	17
Male	124	1	-	-	-	-	-	1	1	10	10	11
Female	157	-	-	-	-	-	-	-	-	-	3	6
Black, both sexes	158	-	-	-	-	1	-	1	3	5	11	11
Male	84	-	-	-	-	1	-	1	2	3	7	4
Female	74	-	-	-	-	-	-	-	1	2	4	7
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Washington												
All races, both sexes	882	-	-	-	-	-	-	1	2	4	19	20
Male	392	-	-	-	-	-	-	1	2	3	12	15
Female	490	-	-	-	-	-	-	-	-	1	7	5
White, both sexes	799	-	-	-	-	-	-	1	-	3	16	10
Male	347	-	-	-	-	-	-	1	-	3	11	8
Female	452	-	-	-	-	-	-	-	-	-	5	2
Black, both sexes	35	-	-	-	-	-	-	-	2	-	2	5
Male	21	-	-	-	-	-	-	-	2	-	1	2
Female	14	-	-	-	-	-	-	-	-	-	1	3
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	1	1	2
Male	5	-	-	-	-	-	-	-	-	-	-	2
Female	4	-	-	-	-	-	-	-	-	1	1	-
Asian or Pacific Islander 2/, both sexes	39	-	-	-	-	-	-	-	-	-	-	3
Male	19	-	-	-	-	-	-	-	-	-	-	3
Female	20	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	131	-	-	-	-	-	-	1	1	2	1	6
Male	45	-	-	-	-	-	-	1	1	2	-	3
Female	86	-	-	-	-	-	-	-	-	-	1	3
White, both sexes	128	-	-	-	-	-	-	1	1	1	1	5
Male	44	-	-	-	-	-	-	1	1	1	-	3
Female	84	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	3	-	-	-	-	-	-	-	-	1	-	1
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Wisconsin												
All races, both sexes	418	-	-	-	-	1	1	-	2	2	11	16
Male	167	-	-	-	-	1	1	-	1	2	7	9
Female	251	-	-	-	-	-	-	-	1	-	4	7
White, both sexes	353	-	-	-	-	-	-	-	2	1	5	7
Male	130	-	-	-	-	-	-	-	1	1	4	6
Female	223	-	-	-	-	-	-	-	1	-	1	1
Black, both sexes	63	-	-	-	-	1	1	-	-	1	6	9
Male	37	-	-	-	-	1	1	-	-	1	3	3
Female	26	-	-	-	-	-	-	-	-	-	3	6
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	25	-	-	-	-	-	-	-	-	-	1	2
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	25	-	-	-	-	-	-	-	-	-	1	2
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
United States												
All races, both sexes	2,048	2,279	2,208	1,909	2,100	2,601	3,723	4,410	3,861	1,892	435	2
Male	1,401	1,535	1,474	1,186	1,080	1,248	1,425	1,315	907	326	52	2
Female	647	744	734	723	1,020	1,353	2,298	3,095	2,954	1,566	383	-
White, both sexes	1,273	1,454	1,454	1,310	1,463	1,925	2,996	3,729	3,383	1,647	367	1
Male	916	1,023	990	823	763	941	1,127	1,107	789	276	41	1
Female	357	431	464	487	700	984	1,869	2,622	2,594	1,371	326	-
Black, both sexes	732	772	695	549	583	611	624	582	395	212	60	1
Male	460	476	445	328	294	276	254	172	94	40	8	1
Female	272	296	250	221	289	335	370	410	301	172	52	-
American Indian 1/, both sexes	17	13	13	12	8	8	12	11	4	1	1	-
Male	10	8	8	9	5	3	5	3	-	1	1	-
Female	7	5	5	3	3	5	7	8	4	-	-	-
Asian or Pacific Islander 2/, both sexes	26	40	46	38	46	57	91	88	79	32	7	-
Male	15	28	31	26	18	28	39	33	24	9	2	-
Female	11	12	15	12	28	29	52	55	55	23	5	-
Alabama												
All races, both sexes	16	23	26	16	30	38	48	61	54	17	5	-
Male	10	18	18	10	24	14	13	15	8	2	1	-
Female	6	5	8	6	6	24	35	46	46	15	4	-
White, both sexes	10	11	11	6	17	24	37	45	41	11	5	-
Male	6	9	8	4	14	8	11	10	5	2	1	-
Female	4	2	3	2	3	16	26	35	36	9	4	-
Black, both sexes	6	12	15	10	13	14	11	16	13	6	-	-
Male	4	9	10	6	10	6	2	5	3	-	-	-
Female	2	3	5	4	3	8	9	11	10	6	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Alaska												
All races, both sexes	8	11	8	7	4	4	3	6	1	2	1	-
Male	5	8	3	6	4	3	-	1	1	-	1	-
Female	3	3	5	1	-	1	3	5	-	2	-	-
White, both sexes	5	10	6	6	4	-	2	4	1	2	-	-
Male	3	7	3	6	4	-	-	1	1	-	-	-
Female	2	3	3	-	-	-	2	3	-	2	-	-
Black, both sexes	-	1	1	-	-	1	-	1	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	3	-	1	1	-	2	1	1	-	-	1	-
Male	2	-	-	-	-	1	-	-	-	-	1	-
Female	1	-	1	1	-	1	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	50	62	48	36	55	63	86	98	65	32	5	-
Male	31	46	31	20	38	28	40	32	22	8	-	-
Female	19	16	17	16	17	35	46	66	43	24	5	-
White, both sexes	43	51	42	32	48	61	83	92	63	32	5	-
Male	28	36	26	17	35	28	39	31	22	8	-	-
Female	15	15	16	15	13	33	44	61	41	24	5	-
Black, both sexes	3	7	4	1	5	2	1	4	1	-	-	-
Male	3	7	3	-	2	-	-	1	-	-	-	-
Female	-	-	1	1	3	2	1	3	1	-	-	-
American Indian 1/, both sexes	4	3	1	3	1	-	1	2	-	-	-	-
Male	-	2	1	3	1	-	-	-	-	-	-	-
Female	4	1	-	-	-	-	1	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	1	-	1	-	1	-	-	-
Male	-	1	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Arkansas												
All races, both sexes	17	19	15	17	13	29	27	50	41	12	3	-
Male	9	10	10	11	7	16	13	11	5	2	-	-
Female	8	9	5	6	6	13	14	39	36	10	3	-
White, both sexes	12	13	11	15	9	24	25	43	39	12	2	-
Male	7	7	8	10	4	14	11	10	4	2	-	-
Female	5	6	3	5	5	10	14	33	35	10	2	-
Black, both sexes	5	6	4	2	4	5	2	7	2	-	1	-
Male	2	3	2	1	3	2	2	1	1	-	-	-
Female	3	3	2	1	1	3	-	6	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	262	257	274	233	242	312	527	615	510	290	54	1
Male	180	165	187	137	134	165	217	192	135	46	5	1
Female	82	92	87	96	108	147	310	423	375	244	49	-
White, both sexes	180	175	200	174	170	240	431	528	433	249	45	1
Male	131	125	137	102	100	129	176	156	112	40	5	1
Female	49	50	63	72	70	111	255	372	321	209	40	-
Black, both sexes	70	65	54	40	46	45	48	38	32	18	4	-
Male	41	30	33	23	24	22	17	14	10	-	-	-
Female	29	35	21	17	22	23	31	24	22	18	4	-
American Indian 1/, both sexes	1	4	3	2	2	-	-	3	3	-	-	-
Male	1	2	2	2	2	-	-	1	-	-	-	-
Female	-	2	1	-	-	-	-	2	3	-	-	-
Asian or Pacific Islander 2/, both sexes	11	13	17	17	24	27	48	46	42	23	5	-
Male	7	8	15	10	8	14	24	21	13	6	-	-
Female	4	5	2	7	16	13	24	25	29	17	5	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Colorado												
All races, both sexes	17	21	28	25	15	27	28	39	45	12	4	-
Male	14	9	16	13	10	16	10	9	9	2	2	-
Female	3	12	12	12	5	11	18	30	36	10	2	-
White, both sexes	15	19	27	20	15	25	26	37	44	11	4	-
Male	13	8	16	11	10	15	10	9	8	2	2	-
Female	2	11	11	9	5	10	16	28	36	9	2	-
Black, both sexes	2	2	1	5	-	2	-	2	1	1	-	-
Male	1	1	-	2	-	1	-	-	1	-	-	-
Female	1	1	1	3	-	1	-	2	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Connecticut												
All races, both sexes	13	16	16	10	11	22	19	36	46	20	4	-
Male	10	11	12	7	5	13	8	7	10	2	1	-
Female	3	5	4	3	6	9	11	29	36	18	3	-
White, both sexes	10	11	12	6	8	20	17	32	46	16	4	-
Male	7	9	9	3	3	12	7	5	10	2	1	-
Female	3	2	3	3	5	8	10	27	36	14	3	-
Black, both sexes	3	5	3	4	3	2	2	4	-	4	-	-
Male	3	2	2	4	2	1	1	2	-	-	-	-
Female	-	3	1	-	1	1	1	2	-	4	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Delaware												
All races, both sexes	9	7	3	4	5	5	5	6	9	4	2	-
Male	6	4	3	4	3	3	1	1	3	-	-	-
Female	3	3	-	-	2	2	4	5	6	4	2	-
White, both sexes	4	6	2	2	5	4	4	6	9	2	2	-
Male	2	3	2	2	3	2	1	1	3	-	-	-
Female	2	3	-	-	2	2	3	5	6	2	2	-
Black, both sexes	5	1	1	2	-	1	1	-	-	2	-	-
Male	4	1	1	2	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	13	12	9	6	10	9	12	16	10	12	2	-
Male	10	4	7	5	5	3	6	7	1	4	-	-
Female	3	8	2	1	5	6	6	9	9	8	2	-
White, both sexes	-	1	-	-	-	3	2	3	3	4	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	-	1	-	-	-	2	1	2	3	4	-	-
Black, both sexes	13	11	9	6	10	6	10	13	7	8	2	-
Male	10	4	7	5	5	2	5	6	1	4	-	-
Female	3	7	2	1	5	4	5	7	6	4	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Florida												
All races, both sexes	154	166	157	146	180	207	295	356	271	136	34	-
Male	103	111	104	95	95	101	134	115	88	28	2	-
Female	51	55	53	51	85	106	161	241	183	108	32	-
White, both sexes	104	111	120	104	141	175	253	313	246	125	29	-
Male	75	76	80	69	74	84	114	106	82	27	1	-
Female	29	35	40	35	67	91	139	207	164	98	28	-
Black, both sexes	49	53	35	38	37	31	41	40	23	11	5	-
Male	27	33	24	23	20	16	19	9	6	1	1	-
Female	22	20	11	15	17	15	22	31	17	10	4	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	2	4	2	1	1	3	2	-	-	-
Male	1	1	-	3	1	1	1	-	-	-	-	-
Female	-	-	2	1	1	-	-	3	2	-	-	-
Georgia												
All races, both sexes	103	133	116	90	88	108	112	121	93	55	14	-
Male	77	89	75	65	41	50	42	37	18	16	1	-
Female	26	44	41	25	47	58	70	84	75	39	13	-
White, both sexes	37	62	53	37	50	58	65	80	60	28	8	-
Male	27	45	36	27	25	28	25	24	9	7	-	-
Female	10	17	17	10	25	30	40	56	51	21	8	-
Black, both sexes	66	71	63	52	38	50	44	41	33	27	6	-
Male	50	44	39	37	16	22	15	13	9	9	1	-
Female	16	27	24	15	22	28	29	28	24	18	5	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	3	-	-	-	-	-
Male	-	-	-	1	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Hawaii												
All races, both sexes	1	2	1	3	4	5	9	13	15	4	1	-
Male	1	1	1	3	2	4	2	4	3	1	-	-
Female	-	1	-	-	2	1	7	9	12	3	1	-
White, both sexes	-	-	-	1	2	2	4	1	7	2	1	-
Male	-	-	-	1	1	2	1	-	1	-	-	-
Female	-	-	-	-	1	-	3	1	6	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	1	2	2	3	5	12	8	2	-	-
Male	1	1	1	2	1	2	1	4	2	1	-	-
Female	-	1	-	-	1	1	4	8	6	1	-	-
Idaho												
All races, both sexes	4	8	6	4	6	8	8	22	20	6	3	-
Male	3	6	6	4	3	3	2	5	4	-	-	-
Female	1	2	-	-	3	5	6	17	16	6	3	-
White, both sexes	4	8	6	4	6	8	8	22	20	6	3	-
Male	3	6	6	4	3	3	2	5	4	-	-	-
Female	1	2	-	-	3	5	6	17	16	6	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Illinois												
All races, both sexes	123	136	109	90	124	104	139	154	142	64	14	1
Male	84	94	66	57	66	48	62	42	30	11	2	1
Female	39	42	43	33	58	56	77	112	112	53	12	-
White, both sexes	63	71	64	51	70	67	108	120	122	57	13	-
Male	44	48	43	32	33	31	45	34	26	8	2	-
Female	19	23	21	19	37	36	63	86	96	49	11	-
Black, both sexes	59	60	43	38	54	32	28	30	20	7	1	1
Male	39	42	21	24	33	15	16	7	4	3	-	1
Female	20	18	22	14	21	17	12	23	16	4	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	5	2	1	-	5	3	4	-	-	-	-
Male	-	4	2	1	-	2	1	1	-	-	-	-
Female	-	1	-	-	-	3	2	3	-	-	-	-
Indiana												
All races, both sexes	26	26	23	25	21	33	67	82	73	22	9	-
Male	15	18	16	14	10	17	21	26	20	4	1	-
Female	11	8	7	11	11	16	46	56	53	18	8	-
White, both sexes	18	16	16	21	13	30	54	80	69	20	8	-
Male	13	11	11	11	6	14	18	25	19	4	1	-
Female	5	5	5	10	7	16	36	55	50	16	7	-
Black, both sexes	7	10	6	4	8	3	13	2	4	2	1	-
Male	2	7	5	3	4	3	3	1	1	-	-	-
Female	5	3	1	1	4	-	10	1	3	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Iowa												
All races, both sexes	10	7	7	11	14	16	29	41	56	26	4	-
Male	7	5	6	4	7	8	13	11	10	-	-	-
Female	3	2	1	7	7	8	16	30	46	26	4	-
White, both sexes	8	7	6	11	13	16	29	39	55	26	4	-
Male	5	5	5	4	6	8	13	11	10	-	-	-
Female	3	2	1	7	7	8	16	28	45	26	4	-
Black, both sexes	2	-	-	-	1	-	-	2	1	-	-	-
Male	2	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	1	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	6	6	6	5	1	12	18	32	33	19	1	-
Male	5	2	4	3	-	6	9	8	7	1	-	-
Female	1	4	2	2	1	6	9	24	26	18	1	-
White, both sexes	4	5	5	5	1	10	18	28	31	19	1	-
Male	3	2	3	3	-	6	9	8	7	1	-	-
Female	1	3	2	2	1	4	9	20	24	18	1	-
Black, both sexes	2	1	1	-	-	1	-	4	2	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	4	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Kentucky												
All races, both sexes	24	27	29	17	20	23	37	38	32	17	5	-
Male	15	19	21	10	13	11	12	10	8	2	-	-
Female	9	8	8	7	7	12	25	28	24	15	5	-
White, both sexes	21	22	24	16	17	23	34	37	31	17	5	-
Male	14	14	17	10	12	11	11	9	8	2	-	-
Female	7	8	7	6	5	12	23	28	23	15	5	-
Black, both sexes	3	5	5	1	2	-	2	1	1	-	-	-
Male	1	5	4	-	-	-	1	1	-	-	-	-
Female	2	-	1	1	2	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Louisiana												
All races, both sexes	56	76	64	67	64	96	115	116	85	36	14	-
Male	37	45	39	32	35	46	48	36	18	8	2	-
Female	19	31	25	35	29	50	67	80	67	28	12	-
White, both sexes	29	42	30	38	34	65	77	89	62	23	6	-
Male	22	27	22	19	19	35	30	29	13	6	-	-
Female	7	15	8	19	15	30	47	60	49	17	6	-
Black, both sexes	26	34	34	29	29	31	38	27	22	13	8	-
Male	14	18	17	13	16	11	18	7	4	2	2	-
Female	12	16	17	16	13	20	20	20	18	11	6	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Maine												
All races, both sexes	3	3	8	3	3	8	6	4	13	7	-	-
Male	2	3	6	3	3	3	1	3	3	2	-	-
Female	1	-	2	-	-	5	5	1	10	5	-	-
White, both sexes	3	3	8	3	3	8	6	3	13	7	-	-
Male	2	3	6	3	3	3	1	2	3	2	-	-
Female	1	-	2	-	-	5	5	1	10	5	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	76	77	82	57	57	64	68	83	71	41	6	-
Male	55	58	57	39	32	34	29	27	16	7	1	-
Female	21	19	25	18	25	30	39	56	55	34	5	-
White, both sexes	35	39	37	32	33	38	43	66	54	34	5	-
Male	26	29	29	23	18	23	18	22	10	5	1	-
Female	9	10	8	9	15	15	25	44	44	29	4	-
Black, both sexes	40	37	42	24	22	25	24	17	15	7	1	-
Male	28	28	26	15	13	11	11	5	4	2	-	-
Female	12	9	16	9	9	14	13	12	11	5	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	3	1	2	1	1	-	2	-	-	-
Male	1	1	2	1	1	-	-	-	2	-	-	-
Female	-	-	1	-	1	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Massachusetts												
All races, both sexes	43	54	34	36	27	21	40	64	56	34	8	-
Male	30	34	22	23	16	9	10	23	10	4	1	-
Female	13	20	12	13	11	12	30	41	46	30	7	-
White, both sexes	35	52	29	32	25	21	39	64	55	31	7	-
Male	24	33	19	20	14	9	9	23	10	4	-	-
Female	11	19	10	12	11	12	30	41	45	27	7	-
Black, both sexes	8	2	3	4	2	-	1	-	1	3	-	-
Male	6	1	1	3	2	-	1	-	-	-	-	-
Female	2	1	2	1	-	-	-	-	1	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	-	-	-	-	-	-	1	-
Male	-	-	2	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	91	99	99	70	88	122	172	200	181	100	22	-
Male	57	70	66	41	31	50	54	66	36	18	4	-
Female	34	29	33	29	57	72	118	134	145	82	18	-
White, both sexes	45	61	62	41	59	88	131	166	171	92	19	-
Male	32	46	41	27	19	41	43	50	35	13	3	-
Female	13	15	21	14	40	47	88	116	136	79	16	-
Black, both sexes	45	37	36	29	27	34	41	34	10	8	3	-
Male	24	23	25	14	10	9	11	16	1	5	1	-
Female	21	14	11	15	17	25	30	18	9	3	2	-
American Indian 1/, both sexes	1	1	1	-	1	-	-	-	-	-	-	-
Male	1	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Minnesota												
All races, both sexes	10	11	8	5	6	15	24	31	41	19	2	-
Male	7	11	6	2	4	11	10	5	9	1	1	-
Female	3	-	2	3	2	4	14	26	32	18	1	-
White, both sexes	7	6	7	5	4	14	23	30	41	19	2	-
Male	4	6	5	2	3	11	9	4	9	1	1	-
Female	3	-	2	3	1	3	14	26	32	18	1	-
Black, both sexes	3	4	1	-	-	1	1	1	-	-	-	-
Male	3	4	1	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	37	46	57	51	74	68	93	111	77	29	10	-
Male	25	35	41	33	36	31	38	27	15	7	1	-
Female	12	11	16	18	38	37	55	84	62	22	9	-
White, both sexes	9	25	22	28	44	39	57	74	57	17	4	-
Male	7	20	18	20	23	21	22	16	11	5	1	-
Female	2	5	4	8	21	18	35	58	46	12	3	-
Black, both sexes	28	21	35	23	30	29	36	36	20	12	6	-
Male	18	15	23	13	13	10	16	11	4	2	-	-
Female	10	6	12	10	17	19	20	25	16	10	6	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Missouri												
All races, both sexes	30	25	29	23	25	38	51	67	48	27	4	-
Male	22	20	18	15	12	12	14	15	16	6	1	-
Female	8	5	11	8	13	26	37	52	32	21	3	-
White, both sexes	14	12	11	15	21	19	41	56	37	23	4	-
Male	9	10	8	10	11	6	12	14	11	5	1	-
Female	5	2	3	5	10	13	29	42	26	18	3	-
Black, both sexes	15	13	18	8	4	18	9	11	11	4	-	-
Male	12	10	10	5	1	6	2	1	5	1	-	-
Female	3	3	8	3	3	12	7	10	6	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	4	8	6	1	3	-	4	8	10	4	1	-
Male	2	4	5	1	2	-	-	3	2	-	-	-
Female	2	4	1	-	1	-	4	5	8	4	1	-
White, both sexes	4	8	6	1	3	-	4	8	10	4	1	-
Male	2	4	5	1	2	-	-	3	2	-	-	-
Female	2	4	1	-	1	-	4	5	8	4	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Nebraska												
All races, both sexes	6	4	6	3	4	10	15	11	33	16	4	-
Male	5	3	4	3	2	4	4	1	4	1	1	-
Female	1	1	2	-	2	6	11	10	29	15	3	-
White, both sexes	6	4	5	3	3	9	14	11	33	16	4	-
Male	5	3	4	3	2	4	3	1	4	1	1	-
Female	1	1	1	-	1	5	11	10	29	15	3	-
Black, both sexes	-	-	1	-	1	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	29	36	42	42	22	21	31	32	24	4	2	-
Male	19	23	24	27	12	15	15	11	3	-	-	-
Female	10	13	18	15	10	6	16	21	21	4	2	-
White, both sexes	23	28	37	35	20	20	24	29	24	4	2	-
Male	18	17	21	23	11	14	13	10	3	-	-	-
Female	5	11	16	12	9	6	11	19	21	4	2	-
Black, both sexes	5	7	5	6	2	1	4	2	-	-	-	-
Male	1	5	3	4	1	1	1	1	-	-	-	-
Female	4	2	2	2	1	-	3	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	1	-	-	2	1	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	1	-	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
New Hampshire												
All races, both sexes	11	5	2	4	7	11	10	19	21	9	3	-
Male	9	3	2	4	5	5	4	5	6	2	1	-
Female	2	2	-	-	2	6	6	14	15	7	2	-
White, both sexes	11	5	2	4	7	11	10	19	21	9	3	-
Male	9	3	2	4	5	5	4	5	6	2	1	-
Female	2	2	-	-	2	6	6	14	15	7	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	37	47	56	39	47	59	127	115	118	53	10	-
Male	24	34	41	27	26	29	53	36	22	9	1	-
Female	13	13	15	12	21	30	74	79	96	44	9	-
White, both sexes	25	29	37	22	31	45	100	105	104	49	9	-
Male	17	22	29	17	17	24	42	32	18	8	1	-
Female	8	7	8	5	14	21	58	73	86	41	8	-
Black, both sexes	12	17	17	17	16	14	24	8	14	2	1	-
Male	7	11	12	10	9	5	11	2	4	1	-	-
Female	5	6	5	7	7	9	13	6	10	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	2	-	-	-	3	2	-	2	-	-
Male	-	1	-	-	-	-	-	2	-	-	-	-
Female	-	-	2	-	-	-	3	-	-	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
New Mexico												
All races, both sexes	11	9	16	15	10	9	14	19	23	10	3	-
Male	8	6	10	7	5	3	4	6	6	1	-	-
Female	3	3	6	8	5	6	10	13	17	9	3	-
White, both sexes	11	9	15	13	10	8	13	19	22	9	3	-
Male	8	6	9	5	5	2	3	6	6	-	-	-
Female	3	3	6	8	5	6	10	13	16	9	3	-
Black, both sexes	-	-	1	-	-	-	1	-	1	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	2	-	1	-	-	-	1	-	-
Male	-	-	-	2	-	1	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	174	203	211	195	197	264	294	332	312	151	41	-
Male	117	140	137	119	88	129	110	100	94	27	7	-
Female	57	63	74	76	109	135	184	232	218	124	34	-
White, both sexes	103	109	125	122	123	175	220	255	259	131	33	-
Male	73	80	78	78	53	84	79	75	83	26	6	-
Female	30	29	47	44	70	91	141	180	176	105	27	-
Black, both sexes	67	89	84	70	71	76	71	66	43	20	7	-
Male	42	55	58	38	33	39	30	22	9	1	-	-
Female	25	34	26	32	38	37	41	44	34	19	7	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	5	1	3	3	13	3	11	10	-	1	-
Male	2	5	-	3	2	6	1	3	2	-	1	-
Female	2	-	1	-	1	7	2	8	8	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
North Carolina												
All races, both sexes	56	50	58	46	52	57	80	88	64	38	5	-
Male	40	34	48	27	23	30	27	23	13	6	-	-
Female	16	16	10	19	29	27	53	65	51	32	5	-
White, both sexes	21	28	30	21	32	33	54	58	53	32	4	-
Male	19	20	26	15	13	18	16	17	10	6	-	-
Female	2	8	4	6	19	15	38	41	43	26	4	-
Black, both sexes	31	22	26	23	19	24	25	30	11	6	1	-
Male	18	14	21	12	10	12	10	6	3	-	-	-
Female	13	8	5	11	9	12	15	24	8	6	1	-
American Indian 1/, both sexes	4	-	-	1	1	-	1	-	-	-	-	-
Male	3	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	1	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	1	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	3	1	3	3	7	11	16	9	2	-
Male	-	-	2	1	-	2	5	-	6	1	-	-
Female	-	-	1	-	3	1	2	11	10	8	2	-
White, both sexes	-	-	2	1	3	3	7	11	16	9	2	-
Male	-	-	2	1	-	2	5	-	6	1	-	-
Female	-	-	-	-	3	1	2	11	10	8	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Ohio												
All races, both sexes	105	123	127	116	124	138	228	272	229	102	28	-
Male	75	88	91	64	64	70	84	66	53	11	5	-
Female	30	35	36	52	60	68	144	206	176	91	23	-
White, both sexes	63	82	86	95	88	101	194	234	203	93	26	-
Male	48	64	63	51	46	51	66	56	48	10	5	-
Female	15	18	23	44	42	50	128	178	155	83	21	-
Black, both sexes	42	39	40	20	35	37	32	38	26	9	2	-
Male	27	23	27	12	18	19	17	10	5	1	-	-
Female	15	16	13	8	17	18	15	28	21	8	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	1	1	1	-	1	-	-	-	-	-
Male	-	1	1	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	1	-	-	-	-	-
Oklahoma												
All races, both sexes	14	19	17	18	32	21	43	40	41	23	8	-
Male	12	12	10	11	16	7	14	14	5	4	2	-
Female	2	7	7	7	16	14	29	26	36	19	6	-
White, both sexes	10	17	14	16	26	15	36	33	38	21	7	-
Male	8	10	9	9	14	6	10	13	4	4	2	-
Female	2	7	5	7	12	9	26	20	34	17	5	-
Black, both sexes	3	1	2	-	3	6	5	6	2	1	1	-
Male	3	1	1	-	1	1	2	1	1	-	-	-
Female	-	-	1	-	2	5	3	5	1	1	1	-
American Indian 1/, both sexes	1	1	1	2	2	-	2	1	1	-	-	-
Male	1	1	-	2	1	-	2	-	-	-	-	-
Female	-	-	1	-	1	-	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Oregon												
All races, both sexes	6	9	8	6	14	9	36	46	64	25	5	-
Male	3	5	7	4	5	2	14	16	15	5	-	-
Female	3	4	1	2	9	7	22	30	49	20	5	-
White, both sexes	5	7	7	5	13	8	33	46	62	24	5	-
Male	2	4	6	4	5	1	13	16	14	4	-	-
Female	3	3	1	1	8	7	20	30	48	20	5	-
Black, both sexes	1	1	1	-	-	1	1	-	1	-	-	-
Male	1	1	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	1	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	1	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	1	1	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Pennsylvania												
All races, both sexes	35	57	53	44	41	78	130	150	111	60	16	-
Male	26	40	39	31	17	36	45	45	23	16	1	-
Female	9	17	14	13	24	42	85	105	88	44	15	-
White, both sexes	21	35	36	36	35	62	109	129	99	54	14	-
Male	16	27	26	25	15	30	39	39	18	15	1	-
Female	5	8	10	11	20	32	70	90	81	39	13	-
Black, both sexes	14	21	17	8	6	16	21	19	12	6	2	-
Male	10	13	13	6	2	6	6	6	5	1	-	-
Female	4	8	4	2	4	10	15	13	7	5	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Rhode Island												
All races, both sexes	5	4	1	3	2	1	5	11	10	4	2	-
Male	3	1	1	2	-	-	1	2	4	1	-	-
Female	2	3	-	1	2	1	4	9	6	3	2	-
White, both sexes	3	4	1	3	2	1	5	11	10	4	2	-
Male	3	1	1	2	-	-	1	2	4	1	-	-
Female	-	3	-	1	2	1	4	9	6	3	2	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	18	27	28	33	35	46	49	51	40	19	5	-
Male	7	15	17	25	21	21	12	14	5	5	1	-
Female	11	12	11	8	14	25	37	37	35	14	4	-
White, both sexes	11	13	13	17	16	19	38	34	24	13	3	-
Male	5	7	10	14	12	8	9	8	4	4	-	-
Female	6	6	3	3	4	11	29	26	20	9	3	-
Black, both sexes	7	14	15	15	19	27	11	17	16	6	2	-
Male	2	8	7	10	9	13	3	6	1	1	1	-
Female	5	6	8	5	10	14	8	11	15	5	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
South Dakota												
All races, both sexes	-	1	-	1	-	1	5	11	15	14	1	-
Male	-	-	-	-	-	1	3	5	1	3	-	-
Female	-	1	-	1	-	-	2	6	14	11	1	-
White, both sexes	-	-	-	1	-	1	5	10	15	14	1	-
Male	-	-	-	-	-	1	3	5	1	3	-	-
Female	-	-	-	1	-	-	2	5	14	11	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	61	77	52	56	64	69	102	142	98	52	7	-
Male	41	51	34	38	26	35	22	41	23	8	1	-
Female	20	26	18	18	38	34	80	101	75	44	6	-
White, both sexes	37	45	28	40	34	43	81	121	86	44	7	-
Male	29	34	18	28	12	16	16	34	19	8	1	-
Female	8	11	10	12	22	27	65	87	67	36	6	-
Black, both sexes	24	32	23	15	30	26	19	21	11	8	-	-
Male	12	17	15	9	14	19	5	7	4	-	-	-
Female	12	15	8	6	16	7	14	14	7	8	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	1	-	1	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Texas												
All races, both sexes	180	143	151	118	136	172	222	261	197	107	27	-
Male	119	92	93	70	57	81	91	80	42	13	4	-
Female	61	51	58	48	79	91	131	181	155	94	23	-
White, both sexes	138	98	117	81	107	137	189	232	169	92	23	-
Male	95	69	72	48	47	67	78	74	35	12	3	-
Female	43	29	45	33	60	70	111	158	134	80	20	-
Black, both sexes	41	42	30	35	28	33	26	28	22	12	4	-
Male	24	22	18	20	10	13	11	5	6	-	1	-
Female	17	20	12	15	18	20	15	23	16	12	3	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	3	3	2	1	2	7	-	6	3	-	-
Male	-	1	2	2	-	1	2	-	1	1	-	-
Female	1	2	1	-	1	1	5	-	5	2	-	-
Utah												
All races, both sexes	5	8	2	5	7	7	14	14	22	6	1	-
Male	4	5	1	4	5	2	2	5	4	-	-	-
Female	1	3	1	1	2	5	12	9	18	6	1	-
White, both sexes	4	7	1	5	7	7	13	14	22	5	1	-
Male	4	4	1	4	5	2	2	5	4	-	-	-
Female	-	3	-	1	2	5	11	9	18	5	1	-
Black, both sexes	-	1	-	-	-	-	-	-	-	1	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Vermont												
All races, both sexes	3	4	1	3	3	4	7	12	6	11	1	-
Male	2	2	-	3	2	2	4	6	1	3	-	-
Female	1	2	1	-	1	2	3	6	5	8	1	-
White, both sexes	3	4	1	3	3	4	7	12	6	11	1	-
Male	2	2	-	3	2	2	4	6	1	3	-	-
Female	1	2	1	-	1	2	3	6	5	8	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	34	29	18	19	27	38	55	60	57	22	10	-
Male	22	18	12	14	16	16	21	20	11	7	1	-
Female	12	11	6	5	11	22	34	40	46	15	9	-
White, both sexes	15	15	8	9	13	29	37	46	43	15	8	-
Male	9	10	4	8	7	11	13	17	9	2	-	-
Female	6	5	4	1	6	18	24	29	34	13	8	-
Black, both sexes	19	13	9	10	14	8	17	13	14	7	2	-
Male	13	8	7	6	9	4	8	3	2	5	1	-
Female	6	5	2	4	5	4	9	10	12	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	-	1	1	1	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Washington												
All races, both sexes	34	60	55	47	46	72	116	159	171	64	12	-
Male	27	49	35	25	30	30	51	55	44	13	-	-
Female	7	11	20	22	16	42	65	104	127	51	12	-
White, both sexes	31	47	47	39	41	65	106	151	166	64	12	-
Male	25	39	32	21	27	29	45	51	42	13	-	-
Female	6	8	15	18	14	36	61	100	124	51	12	-
Black, both sexes	-	10	2	6	1	2	3	2	-	-	-	-
Male	-	7	-	4	1	-	2	2	-	-	-	-
Female	-	3	2	2	-	2	1	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	1	-	2	-	-	-	-	-	-
Male	1	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	3	5	1	4	3	7	6	5	-	-	-
Male	1	3	2	-	2	-	4	2	2	-	-	-
Female	1	-	3	1	2	3	3	4	3	-	-	-
West Virginia												
All races, both sexes	4	2	5	9	9	12	18	18	28	12	3	-
Male	4	1	4	6	2	6	5	4	2	4	-	-
Female	-	1	1	3	7	6	13	14	26	8	3	-
White, both sexes	4	2	5	9	9	12	17	18	28	12	3	-
Male	4	1	4	6	2	6	5	4	2	4	-	-
Female	-	1	1	3	7	6	12	14	26	8	3	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Wisconsin												
All races, both sexes	13	19	27	28	25	40	71	63	59	33	7	-
Male	11	9	16	17	17	16	23	20	13	4	-	-
Female	2	10	11	11	8	24	48	43	46	29	7	-
White, both sexes	8	14	19	24	22	33	61	62	56	32	7	-
Male	7	7	10	13	15	12	18	20	13	3	-	-
Female	1	7	9	11	7	21	43	42	43	29	7	-
Black, both sexes	5	5	8	4	3	6	9	1	3	1	-	-
Male	4	2	6	4	2	4	5	-	-	1	-	-
Female	1	3	2	-	1	2	4	1	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Wyoming												
All races, both sexes	1	5	1	-	3	2	2	3	4	1	-	-
Male	1	4	-	-	3	1	2	2	-	-	-	-
Female	-	1	1	-	-	1	-	1	4	1	-	-
White, both sexes	1	5	1	-	3	2	2	3	4	1	-	-
Male	1	4	-	-	3	1	2	2	-	-	-	-
Female	-	1	1	-	-	1	-	1	4	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
United States												
All races, both sexes	9,551	1	-	-	-	4	8	19	38	79	106	197
Male	4,170	1	-	-	-	2	3	7	20	41	71	122
Female	5,381	-	-	-	-	2	5	12	18	38	35	75
White, both sexes	6,787	1	-	-	-	1	4	9	10	35	38	76
Male	2,939	1	-	-	-	1	1	4	7	20	27	47
Female	3,848	-	-	-	-	-	3	5	3	15	11	29
Black, both sexes	2,413	-	-	-	-	3	4	10	26	43	64	112
Male	1,073	-	-	-	-	1	2	3	12	20	42	69
Female	1,340	-	-	-	-	2	2	7	14	23	22	43
American Indian 1/, both sexes	50	-	-	-	-	-	-	-	-	-	1	1
Male	20	-	-	-	-	-	-	-	-	-	1	-
Female	30	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	301	-	-	-	-	-	-	-	2	1	3	8
Male	138	-	-	-	-	-	-	-	1	1	1	6
Female	163	-	-	-	-	-	-	-	1	-	2	2
Alabama												
All races, both sexes	164	-	-	-	-	-	-	-	2	6	2	6
Male	63	-	-	-	-	-	-	-	-	4	2	5
Female	101	-	-	-	-	-	-	-	2	2	-	1
White, both sexes	68	-	-	-	-	-	-	-	-	2	1	2
Male	29	-	-	-	-	-	-	-	-	2	1	2
Female	39	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	96	-	-	-	-	-	-	-	2	4	1	4
Male	34	-	-	-	-	-	-	-	-	2	1	3
Female	62	-	-	-	-	-	-	-	2	2	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Alaska												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	204	-	-	-	-	-	-	-	-	2	1	-
Male	95	-	-	-	-	-	-	-	-	1	1	-
Female	109	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	177	-	-	-	-	-	-	-	-	1	1	-
Male	82	-	-	-	-	-	-	-	-	-	1	-
Female	95	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	15	-	-	-	-	-	-	-	-	1	-	-
Male	9	-	-	-	-	-	-	-	-	1	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Arkansas												
All races, both sexes	86	-	-	-	-	-	-	-	1	1	2	2
Male	26	-	-	-	-	-	-	-	-	1	1	-
Female	60	-	-	-	-	-	-	-	1	-	1	2
White, both sexes	54	-	-	-	-	-	-	-	-	1	1	1
Male	19	-	-	-	-	-	-	-	-	1	1	-
Female	35	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	32	-	-	-	-	-	-	-	1	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	1	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1,212	-	-	-	-	-	-	4	5	8	7	29
Male	545	-	-	-	-	-	-	3	2	5	4	19
Female	667	-	-	-	-	-	-	1	3	3	3	10
White, both sexes	878	-	-	-	-	-	-	2	3	5	4	14
Male	393	-	-	-	-	-	-	2	1	3	2	7
Female	485	-	-	-	-	-	-	-	2	2	2	7
Black, both sexes	180	-	-	-	-	-	-	2	1	3	1	10
Male	84	-	-	-	-	-	-	1	1	2	1	9
Female	96	-	-	-	-	-	-	1	-	1	-	1
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	148	-	-	-	-	-	-	-	1	-	2	5
Male	66	-	-	-	-	-	-	-	-	-	1	3
Female	82	-	-	-	-	-	-	-	1	-	1	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Colorado												
All races, both sexes	94	-	-	-	-	-	-	-	-	-	-	-
Male	50	-	-	-	-	-	-	-	-	-	-	-
Female	44	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	83	-	-	-	-	-	-	-	-	-	-	-
Male	43	-	-	-	-	-	-	-	-	-	-	-
Female	40	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	78	-	-	-	-	-	-	-	-	-	-	-
Male	32	-	-	-	-	-	-	-	-	-	-	-
Female	46	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	70	-	-	-	-	-	-	-	-	-	-	-
Male	27	-	-	-	-	-	-	-	-	-	-	-
Female	43	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Delaware												
All races, both sexes	22	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	39	-	-	-	-	-	-	-	-	-	-	2
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	32	-	-	-	-	-	-	-	-	-	-	2
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Florida												
All races, both sexes	747	-	-	-	-	1	2	2	3	5	13	14
Male	382	-	-	-	-	-	-	-	2	4	9	7
Female	365	-	-	-	-	1	2	2	1	1	4	7
White, both sexes	545	-	-	-	-	-	1	2	1	1	6	3
Male	296	-	-	-	-	-	-	-	1	1	5	1
Female	249	-	-	-	-	-	1	2	-	-	1	2
Black, both sexes	196	-	-	-	-	1	1	-	2	4	7	11
Male	83	-	-	-	-	-	-	-	1	3	4	6
Female	113	-	-	-	-	1	1	-	1	1	3	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	289	-	-	-	-	-	-	-	2	3	8	10
Male	113	-	-	-	-	-	-	-	-	2	5	7
Female	176	-	-	-	-	-	-	-	2	1	3	3
White, both sexes	120	-	-	-	-	-	-	-	-	-	3	5
Male	48	-	-	-	-	-	-	-	-	-	2	3
Female	72	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	166	-	-	-	-	-	-	-	2	3	5	4
Male	64	-	-	-	-	-	-	-	-	2	3	3
Female	102	-	-	-	-	-	-	-	2	1	2	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Hawaii												
All races, both sexes	36	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	27	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	36	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	36	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Illinois												
All races, both sexes	341	-	-	-	-	-	-	-	1	1	2	7
Male	155	-	-	-	-	-	-	-	1	-	1	5
Female	186	-	-	-	-	-	-	-	-	1	1	2
White, both sexes	249	-	-	-	-	-	-	-	-	-	-	-
Male	106	-	-	-	-	-	-	-	-	-	-	-
Female	143	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	87	-	-	-	-	-	-	-	1	1	2	7
Male	46	-	-	-	-	-	-	-	1	-	1	5
Female	41	-	-	-	-	-	-	-	-	1	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	202	-	-	-	-	-	-	1	1	2	6	2
Male	90	-	-	-	-	-	-	-	-	1	4	2
Female	112	-	-	-	-	-	-	1	1	1	2	-
White, both sexes	166	-	-	-	-	-	-	-	-	2	3	1
Male	75	-	-	-	-	-	-	-	-	1	2	1
Female	91	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	36	-	-	-	-	-	-	1	1	-	3	1
Male	15	-	-	-	-	-	-	-	-	-	2	1
Female	21	-	-	-	-	-	-	1	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Iowa												
All races, both sexes	147	-	-	-	-	-	-	-	-	-	-	2
Male	56	-	-	-	-	-	-	-	-	-	-	1
Female	91	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	140	-	-	-	-	-	-	-	-	-	-	1
Male	54	-	-	-	-	-	-	-	-	-	-	1
Female	86	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	57	-	-	-	-	-	-	-	-	2	1	2
Male	16	-	-	-	-	-	-	-	-	-	1	2
Female	41	-	-	-	-	-	-	-	-	2	-	-
White, both sexes	50	-	-	-	-	-	-	-	-	1	1	2
Male	14	-	-	-	-	-	-	-	-	-	1	2
Female	36	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Kentucky												
All races, both sexes	102	-	-	-	-	-	1	1	-	1	-	3
Male	44	-	-	-	-	-	1	-	-	-	-	2
Female	58	-	-	-	-	-	-	1	-	1	-	1
White, both sexes	83	-	-	-	-	-	-	1	-	-	-	2
Male	37	-	-	-	-	-	-	-	-	-	-	2
Female	46	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	19	-	-	-	-	-	1	-	-	1	-	1
Male	7	-	-	-	-	-	1	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	161	-	-	-	-	-	1	-	1	3	1	5
Male	74	-	-	-	-	-	1	-	-	1	1	3
Female	87	-	-	-	-	-	-	-	1	2	-	2
White, both sexes	72	-	-	-	-	-	1	-	-	1	-	-
Male	31	-	-	-	-	-	1	-	-	-	-	-
Female	41	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	88	-	-	-	-	-	-	-	1	2	1	5
Male	42	-	-	-	-	-	-	-	-	1	1	3
Female	46	-	-	-	-	-	-	-	1	1	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Maine												
All races, both sexes	31	-	-	-	-	-	-	-	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	30	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	205	-	-	-	-	-	1	-	1	1	6	7
Male	96	-	-	-	-	-	-	-	1	1	4	5
Female	109	-	-	-	-	-	1	-	-	-	2	2
White, both sexes	105	-	-	-	-	-	-	-	-	1	-	1
Male	48	-	-	-	-	-	-	-	-	1	-	1
Female	57	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	95	-	-	-	-	-	1	-	1	-	6	6
Male	48	-	-	-	-	-	-	-	1	-	4	4
Female	47	-	-	-	-	-	1	-	-	-	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Massachusetts												
All races, both sexes	129	-	-	-	-	-	-	-	-	-	1	-
Male	55	-	-	-	-	-	-	-	-	-	1	-
Female	74	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	110	-	-	-	-	-	-	-	-	-	1	-
Male	46	-	-	-	-	-	-	-	-	-	1	-
Female	64	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	280	-	-	-	-	-	-	1	1	-	2	4
Male	113	-	-	-	-	-	-	1	1	-	2	2
Female	167	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	223	-	-	-	-	-	-	1	-	-	1	2
Male	88	-	-	-	-	-	-	1	-	-	1	1
Female	135	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	56	-	-	-	-	-	-	-	1	-	1	2
Male	24	-	-	-	-	-	-	-	1	-	1	1
Female	32	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Minnesota												
All races, both sexes	123	-	-	-	-	-	-	-	-	1	-	-
Male	59	-	-	-	-	-	-	-	-	1	-	-
Female	64	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	112	-	-	-	-	-	-	-	-	1	-	-
Male	53	-	-	-	-	-	-	-	-	1	-	-
Female	59	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	163	-	-	-	-	-	-	-	2	3	5	6
Male	65	-	-	-	-	-	-	-	2	1	2	2
Female	98	-	-	-	-	-	-	-	-	2	3	4
White, both sexes	66	-	-	-	-	-	-	-	-	1	1	-
Male	23	-	-	-	-	-	-	-	-	-	-	-
Female	43	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	95	-	-	-	-	-	-	-	2	2	3	6
Male	42	-	-	-	-	-	-	-	2	1	2	2
Female	53	-	-	-	-	-	-	-	-	1	1	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Missouri												
All races, both sexes	186	-	-	-	-	1	-	-	-	-	1	2
Male	75	-	-	-	-	1	-	-	-	-	1	1
Female	111	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	143	-	-	-	-	1	-	-	-	-	-	1
Male	53	-	-	-	-	1	-	-	-	-	-	-
Female	90	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	38	-	-	-	-	-	-	-	-	-	1	1
Male	20	-	-	-	-	-	-	-	-	-	1	1
Female	18	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	19	-	-	-	-	-	-	-	-	1	-	-
Male	9	-	-	-	-	-	-	-	-	1	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	-	-	1	-	-
Male	9	-	-	-	-	-	-	-	-	1	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Nebraska												
All races, both sexes	61	-	-	-	-	-	-	-	-	-	-	-
Male	29	-	-	-	-	-	-	-	-	-	-	-
Female	32	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	55	-	-	-	-	-	-	-	-	-	-	-
Male	26	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	37	-	-	-	-	-	-	-	-	-	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	31	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
New Hampshire												
All races, both sexes	30	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	30	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	255	-	-	-	-	-	-	-	2	2	-	4
Male	101	-	-	-	-	-	-	-	1	-	-	1
Female	154	-	-	-	-	-	-	-	1	2	-	3
White, both sexes	174	-	-	-	-	-	-	-	1	-	-	2
Male	58	-	-	-	-	-	-	-	-	-	-	-
Female	116	-	-	-	-	-	-	-	1	-	-	2
Black, both sexes	71	-	-	-	-	-	-	-	-	2	-	2
Male	38	-	-	-	-	-	-	-	-	-	-	1
Female	33	-	-	-	-	-	-	-	-	2	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
New Mexico												
All races, both sexes	48	-	-	-	-	-	-	-	-	2	-	-
Male	24	-	-	-	-	-	-	-	-	1	-	-
Female	24	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	39	-	-	-	-	-	-	-	-	2	-	-
Male	20	-	-	-	-	-	-	-	-	1	-	-
Female	19	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	639	-	-	-	-	1	2	-	-	9	13	14
Male	303	-	-	-	-	-	1	-	-	6	9	12
Female	336	-	-	-	-	1	1	-	-	3	4	2
White, both sexes	431	-	-	-	-	-	1	-	-	4	4	5
Male	194	-	-	-	-	-	-	-	-	4	3	5
Female	237	-	-	-	-	-	1	-	-	-	1	-
Black, both sexes	184	-	-	-	-	1	1	-	-	5	9	8
Male	95	-	-	-	-	-	1	-	-	2	6	6
Female	89	-	-	-	-	1	-	-	-	3	3	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	24	-	-	-	-	-	-	-	-	-	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
North Carolina												
All races, both sexes	315	-	-	-	-	1	-	3	2	2	5	12
Male	122	-	-	-	-	1	-	1	2	1	2	9
Female	193	-	-	-	-	-	-	2	-	1	3	3
White, both sexes	153	-	-	-	-	-	-	-	1	1	1	4
Male	46	-	-	-	-	-	-	-	1	-	1	3
Female	107	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	156	-	-	-	-	1	-	3	1	1	4	8
Male	73	-	-	-	-	1	-	1	1	1	1	6
Female	83	-	-	-	-	-	-	2	-	-	3	2
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	38	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	36	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Ohio												
All races, both sexes	524	-	-	-	-	-	-	1	2	4	5	8
Male	221	-	-	-	-	-	-	-	2	1	3	4
Female	303	-	-	-	-	-	-	1	-	3	2	4
White, both sexes	401	-	-	-	-	-	-	1	2	2	3	5
Male	163	-	-	-	-	-	-	-	2	-	1	3
Female	238	-	-	-	-	-	-	1	-	2	2	2
Black, both sexes	118	-	-	-	-	-	-	-	-	2	2	3
Male	54	-	-	-	-	-	-	-	-	1	2	1
Female	64	-	-	-	-	-	-	-	-	1	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	84	-	-	-	-	-	-	-	-	-	-	-
Male	28	-	-	-	-	-	-	-	-	-	-	-
Female	56	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	60	-	-	-	-	-	-	-	-	-	-	-
Male	23	-	-	-	-	-	-	-	-	-	-	-
Female	37	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Oregon												
All races, both sexes	106	-	-	-	-	-	-	-	-	1	-	1
Male	41	-	-	-	-	-	-	-	-	1	-	1
Female	65	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	98	-	-	-	-	-	-	-	-	-	-	1
Male	36	-	-	-	-	-	-	-	-	-	-	1
Female	62	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	358	-	-	-	-	-	-	-	-	3	4	5
Male	153	-	-	-	-	-	-	-	-	3	2	3
Female	205	-	-	-	-	-	-	-	-	-	2	2
White, both sexes	285	-	-	-	-	-	-	-	-	1	2	1
Male	122	-	-	-	-	-	-	-	-	1	1	1
Female	163	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	69	-	-	-	-	-	-	-	-	2	2	4
Male	29	-	-	-	-	-	-	-	-	2	1	2
Female	40	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Rhode Island												
All races, both sexes	26	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	189	-	-	-	-	-	-	3	3	1	-	3
Male	83	-	-	-	-	-	-	1	1	-	-	1
Female	106	-	-	-	-	-	-	2	2	1	-	2
White, both sexes	82	-	-	-	-	-	-	1	-	-	-	1
Male	35	-	-	-	-	-	-	1	-	-	-	1
Female	47	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	107	-	-	-	-	-	-	2	3	1	-	2
Male	48	-	-	-	-	-	-	-	1	-	-	-
Female	59	-	-	-	-	-	-	2	2	1	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
South Dakota												
All races, both sexes	30	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	259	-	-	-	-	-	-	1	2	3	6	13
Male	118	-	-	-	-	-	-	-	2	-	4	8
Female	141	-	-	-	-	-	-	1	-	3	2	5
White, both sexes	166	-	-	-	-	-	-	-	1	1	1	4
Male	79	-	-	-	-	-	-	-	1	-	1	2
Female	87	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	91	-	-	-	-	-	-	1	1	2	5	9
Male	38	-	-	-	-	-	-	-	1	-	3	6
Female	53	-	-	-	-	-	-	1	-	2	2	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Texas												
All races, both sexes	687	1	-	-	-	-	1	2	4	10	8	16
Male	278	1	-	-	-	-	-	1	3	4	6	8
Female	409	-	-	-	-	-	1	1	1	6	2	8
White, both sexes	500	1	-	-	-	-	1	1	1	5	3	9
Male	207	1	-	-	-	-	-	-	1	2	2	3
Female	293	-	-	-	-	-	1	1	-	3	1	6
Black, both sexes	176	-	-	-	-	-	-	1	3	5	5	6
Male	67	-	-	-	-	-	-	1	2	2	4	4
Female	109	-	-	-	-	-	-	-	1	3	1	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	37	-	-	-	-	-	-	-	-	-	-	1
Male	23	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	34	-	-	-	-	-	-	-	-	-	-	1
Male	21	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Vermont												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	177	-	-	-	-	-	-	-	3	1	4	9
Male	74	-	-	-	-	-	-	-	-	1	3	6
Female	103	-	-	-	-	-	-	-	3	-	1	3
White, both sexes	86	-	-	-	-	-	-	-	-	1	-	4
Male	36	-	-	-	-	-	-	-	-	1	-	3
Female	50	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	88	-	-	-	-	-	-	-	3	-	4	5
Male	37	-	-	-	-	-	-	-	-	-	3	3
Female	51	-	-	-	-	-	-	-	3	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Washington												
All races, both sexes	234	-	-	-	-	-	-	-	-	-	2	2
Male	95	-	-	-	-	-	-	-	-	-	2	1
Female	139	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	210	-	-	-	-	-	-	-	-	-	-	2
Male	87	-	-	-	-	-	-	-	-	-	-	1
Female	123	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	14	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	67	-	-	-	-	-	-	-	-	1	-	-
Male	33	-	-	-	-	-	-	-	-	-	-	-
Female	34	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	59	-	-	-	-	-	-	-	-	-	-	-
Male	30	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease and secondary hypertension (I12,I15)												
Wisconsin												
All races, both sexes	164	-	-	-	-	-	-	-	-	-	1	4
Male	69	-	-	-	-	-	-	-	-	-	1	3
Female	95	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	146	-	-	-	-	-	-	-	-	-	1	2
Male	58	-	-	-	-	-	-	-	-	-	1	2
Female	88	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	17	-	-	-	-	-	-	-	-	-	-	2
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
United States												
All races, both sexes	330	359	453	580	818	1,132	1,695	1,882	1,277	477	96	-
Male	192	189	245	297	408	520	735	722	429	146	20	-
Female	138	170	208	283	410	612	960	1,160	848	331	76	-
White, both sexes	126	159	248	337	540	799	1,319	1,543	1,073	391	78	-
Male	74	85	131	176	272	382	586	612	374	121	18	-
Female	52	74	117	161	268	417	733	931	699	270	60	-
Black, both sexes	196	183	196	218	250	286	304	281	160	63	14	-
Male	113	97	107	110	120	117	119	81	45	13	2	-
Female	83	86	89	108	130	169	185	200	115	50	12	-
American Indian 1/, both sexes	2	4	2	8	4	13	5	5	3	2	-	-
Male	2	1	2	5	2	5	1	1	-	-	-	-
Female	-	3	-	3	2	8	4	4	3	2	-	-
Asian or Pacific Islander 2/, both sexes	6	13	7	17	24	34	67	53	41	21	4	-
Male	3	6	5	6	14	16	29	28	10	12	-	-
Female	3	7	2	11	10	18	38	25	31	9	4	-
Alabama												
All races, both sexes	9	6	14	20	17	19	19	27	11	4	2	-
Male	4	2	6	11	7	3	7	10	2	-	-	-
Female	5	4	8	9	10	16	12	17	9	4	2	-
White, both sexes	2	1	4	6	8	6	10	16	7	2	1	-
Male	1	1	3	2	3	1	5	7	1	-	-	-
Female	1	-	1	4	5	5	5	9	6	2	1	-
Black, both sexes	7	5	10	14	9	13	9	11	4	2	1	-
Male	3	1	3	9	4	2	2	3	1	-	-	-
Female	4	4	7	5	5	11	7	8	3	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Alaska												
All races, both sexes	-	-	1	-	2	1	-	1	-	1	-	-
Male	-	-	1	-	1	-	-	1	-	1	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
White, both sexes	-	-	1	-	2	1	-	1	-	-	-	-
Male	-	-	1	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	5	7	6	25	20	24	36	40	30	5	3	-
Male	1	4	2	14	10	15	20	15	9	3	-	-
Female	4	3	4	11	10	9	16	25	21	2	3	-
White, both sexes	4	4	6	20	14	21	33	36	29	5	3	-
Male	-	3	2	11	8	13	18	14	9	3	-	-
Female	4	1	4	9	6	8	15	22	20	2	3	-
Black, both sexes	1	-	-	3	4	3	-	2	1	-	-	-
Male	1	-	-	2	2	2	-	1	-	-	-	-
Female	-	-	-	1	2	1	-	1	1	-	-	-
American Indian 1/, both sexes	-	2	-	1	2	-	2	2	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	1	2	-	1	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Arkansas												
All races, both sexes	6	5	2	3	6	12	14	12	14	4	2	-
Male	2	2	1	3	3	2	4	3	3	1	-	-
Female	4	3	1	-	3	10	10	9	11	3	2	-
White, both sexes	-	3	2	1	4	6	8	9	14	3	1	-
Male	-	2	1	1	3	2	2	2	3	1	-	-
Female	-	1	1	-	1	4	6	7	11	2	1	-
Black, both sexes	6	2	-	2	2	6	6	3	-	1	1	-
Male	2	-	-	2	-	-	2	1	-	-	-	-
Female	4	2	-	-	2	6	4	2	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	42	46	61	56	95	131	256	232	168	63	9	-
Male	28	24	38	22	52	61	103	97	57	30	-	-
Female	14	22	23	34	43	70	153	135	111	33	9	-
White, both sexes	26	25	37	41	72	91	197	177	129	50	5	-
Male	17	13	21	15	40	44	82	72	50	24	-	-
Female	9	12	16	26	32	47	115	105	79	26	5	-
Black, both sexes	12	15	18	8	15	22	27	27	14	4	1	-
Male	8	8	12	4	7	8	8	10	3	2	-	-
Female	4	7	6	4	8	14	19	17	11	2	1	-
American Indian 1/, both sexes	-	-	1	-	-	2	1	-	1	1	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	1	1	-	-
Asian or Pacific Islander 2/, both sexes	4	6	5	7	8	16	31	28	24	8	3	-
Male	3	3	4	3	5	8	13	15	4	4	-	-
Female	1	3	1	4	3	8	18	13	20	4	3	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Colorado												
All races, both sexes	3	3	4	3	6	14	15	21	14	11	-	-
Male	3	2	3	1	5	8	7	12	4	5	-	-
Female	-	1	1	2	1	6	8	9	10	6	-	-
White, both sexes	3	3	4	3	4	13	11	17	14	11	-	-
Male	3	2	3	1	3	7	5	10	4	5	-	-
Female	-	1	1	2	1	6	6	7	10	6	-	-
Black, both sexes	-	-	-	-	2	1	2	3	-	-	-	-
Male	-	-	-	-	2	1	2	1	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Connecticut												
All races, both sexes	2	2	3	1	7	11	13	17	10	11	1	-
Male	1	1	2	1	1	6	3	9	4	4	-	-
Female	1	1	1	-	6	5	10	8	6	7	1	-
White, both sexes	2	1	2	1	6	9	12	17	9	10	1	-
Male	1	-	1	1	1	4	3	9	3	4	-	-
Female	1	1	1	-	5	5	9	8	6	6	1	-
Black, both sexes	-	1	1	-	1	2	1	-	1	1	-	-
Male	-	1	1	-	-	2	-	-	1	-	-	-
Female	-	-	-	-	1	-	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Delaware												
All races, both sexes	-	-	-	1	4	3	5	6	2	-	-	-
Male	-	-	-	1	4	2	3	1	-	-	-	-
Female	-	-	-	-	-	1	2	5	2	-	-	-
White, both sexes	-	-	-	-	1	2	5	6	1	-	-	-
Male	-	-	-	-	1	1	3	1	-	-	-	-
Female	-	-	-	-	-	1	2	5	1	-	-	-
Black, both sexes	-	-	-	1	3	1	-	-	1	-	-	-
Male	-	-	-	1	3	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	2	2	1	6	2	5	7	6	1	3	2	-
Male	-	1	-	4	2	4	4	1	1	-	-	-
Female	2	1	1	2	-	1	3	5	-	3	2	-
White, both sexes	-	-	-	-	1	-	1	1	-	1	1	-
Male	-	-	-	-	1	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	2	2	1	6	1	5	5	4	1	2	1	-
Male	-	1	-	4	1	4	3	-	1	-	-	-
Female	2	1	1	2	-	1	2	4	-	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Florida												
All races, both sexes	23	28	29	47	78	91	121	153	96	33	8	-
Male	13	13	17	27	41	50	59	76	49	13	2	-
Female	10	15	12	20	37	41	62	77	47	20	6	-
White, both sexes	8	10	15	30	58	59	98	133	84	28	8	-
Male	5	6	10	18	32	38	50	70	45	12	2	-
Female	3	4	5	12	26	21	48	63	39	16	6	-
Black, both sexes	15	18	14	16	19	30	23	19	12	4	-	-
Male	8	7	7	9	8	11	9	5	4	1	-	-
Female	7	11	7	7	11	19	14	14	8	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	2	-	1	-	1	-	-
Male	-	-	-	-	1	1	-	1	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	1	-	-
Georgia												
All races, both sexes	17	22	20	18	26	30	48	42	26	13	4	-
Male	9	10	9	7	11	13	13	13	11	3	-	-
Female	8	12	11	11	15	17	35	29	15	10	4	-
White, both sexes	-	8	5	6	13	11	24	23	14	6	2	-
Male	-	3	3	3	6	6	6	8	6	2	-	-
Female	-	5	2	3	7	5	18	15	8	4	2	-
Black, both sexes	17	14	15	12	13	18	23	19	12	7	2	-
Male	9	7	6	4	5	7	7	5	5	1	-	-
Female	8	7	9	8	8	11	16	14	7	6	2	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Hawaii												
All races, both sexes	1	2	1	1	5	1	8	6	6	4	1	-
Male	-	1	1	-	2	-	4	3	3	2	-	-
Female	1	1	-	1	3	1	4	3	3	2	1	-
White, both sexes	-	-	1	-	2	-	1	3	1	-	-	-
Male	-	-	1	-	1	-	-	1	1	-	-	-
Female	-	-	-	-	1	-	1	2	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	1	3	1	7	3	5	4	1	-
Male	-	-	-	-	1	-	4	2	2	2	-	-
Female	1	1	-	1	2	1	3	1	3	2	1	-
Idaho												
All races, both sexes	1	-	1	3	3	6	3	10	6	3	-	-
Male	1	-	1	2	1	2	-	2	4	2	-	-
Female	-	-	-	1	2	4	3	8	2	1	-	-
White, both sexes	1	-	1	3	3	6	3	10	6	3	-	-
Male	1	-	1	2	1	2	-	2	4	2	-	-
Female	-	-	-	1	2	4	3	8	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Illinois												
All races, both sexes	9	11	15	16	30	39	61	72	49	22	6	-
Male	6	6	10	8	17	20	28	28	18	6	1	-
Female	3	5	5	8	13	19	33	44	31	16	5	-
White, both sexes	2	5	9	10	16	28	49	64	42	18	6	-
Male	1	2	5	6	8	15	24	24	16	4	1	-
Female	1	3	4	4	8	13	25	40	26	14	5	-
Black, both sexes	7	5	6	6	14	10	11	7	6	4	-	-
Male	5	4	5	2	9	4	4	3	1	2	-	-
Female	2	1	1	4	5	6	7	4	5	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	1	1	1	-	-	-
Male	-	-	-	-	-	1	-	1	1	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
Indiana												
All races, both sexes	5	5	14	12	19	19	32	45	29	10	-	-
Male	4	3	9	3	6	10	15	21	9	3	-	-
Female	1	2	5	9	13	9	17	24	20	7	-	-
White, both sexes	5	5	8	7	13	17	28	39	29	9	-	-
Male	4	3	5	2	4	8	14	19	9	3	-	-
Female	1	2	3	5	9	9	14	20	20	6	-	-
Black, both sexes	-	-	6	5	6	2	4	6	-	1	-	-
Male	-	-	4	1	2	2	1	2	-	-	-	-
Female	-	-	2	4	4	-	3	4	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Iowa												
All races, both sexes	2	2	4	5	6	14	36	34	30	11	1	-
Male	1	1	1	2	2	7	18	12	10	1	-	-
Female	1	1	3	3	4	7	18	22	20	10	1	-
White, both sexes	2	1	3	5	5	13	36	34	28	11	1	-
Male	1	-	1	2	2	7	18	12	9	1	-	-
Female	1	1	2	3	3	6	18	22	19	10	1	-
Black, both sexes	-	1	1	-	1	1	-	-	2	-	-	-
Male	-	1	-	-	-	-	-	-	1	-	-	-
Female	-	-	1	-	1	1	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	1	1	1	4	6	13	14	8	3	1	-
Male	-	1	1	1	1	2	4	2	1	-	-	-
Female	-	-	-	-	3	4	9	12	7	3	1	-
White, both sexes	-	1	1	1	2	4	11	14	8	3	1	-
Male	-	1	1	1	1	1	3	2	1	-	-	-
Female	-	-	-	-	1	3	8	12	7	3	1	-
Black, both sexes	-	-	-	-	2	2	2	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	2	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Kentucky												
All races, both sexes	4	3	4	6	5	14	9	29	16	5	1	-
Male	2	2	1	2	3	7	4	14	4	1	1	-
Female	2	1	3	4	2	7	5	15	12	4	-	-
White, both sexes	2	-	1	5	3	12	8	28	15	5	1	-
Male	1	-	-	2	3	6	4	13	4	1	1	-
Female	1	-	1	3	-	6	4	15	11	4	-	-
Black, both sexes	2	3	3	1	2	2	1	1	1	-	-	-
Male	1	2	1	-	-	1	-	1	-	-	-	-
Female	1	1	2	1	2	1	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	18	11	12	13	14	18	20	23	14	6	1	-
Male	12	2	5	9	4	9	14	8	4	1	-	-
Female	6	9	7	4	10	9	6	15	10	5	1	-
White, both sexes	3	5	5	6	9	6	11	13	9	3	-	-
Male	2	-	1	5	3	1	7	6	4	1	-	-
Female	1	5	4	1	6	5	4	7	5	2	-	-
Black, both sexes	15	6	7	7	5	11	9	10	5	3	1	-
Male	10	2	4	4	1	7	7	2	-	-	-	-
Female	5	4	3	3	4	4	2	8	5	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Maine												
All races, both sexes	-	-	-	-	-	7	4	11	7	-	1	-
Male	-	-	-	-	-	4	2	6	1	-	-	-
Female	-	-	-	-	-	3	2	5	6	-	1	-
White, both sexes	-	-	-	-	-	7	4	11	7	-	1	-
Male	-	-	-	-	-	4	2	6	1	-	-	-
Female	-	-	-	-	-	3	2	5	6	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	9	8	12	8	26	21	33	34	20	15	3	-
Male	7	6	5	1	14	9	14	16	7	6	-	-
Female	2	2	7	7	12	12	19	18	13	9	3	-
White, both sexes	1	1	3	4	13	7	22	27	13	9	3	-
Male	-	1	1	1	8	3	10	13	4	5	-	-
Female	1	-	2	3	5	4	12	14	9	4	3	-
Black, both sexes	8	7	9	4	13	11	10	7	7	5	-	-
Male	7	5	4	-	6	6	4	3	3	1	-	-
Female	1	2	5	4	7	5	6	4	4	4	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	3	1	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	3	1	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Massachusetts												
All races, both sexes	1	4	2	4	9	17	28	33	22	7	1	-
Male	-	3	1	1	4	8	15	11	10	1	-	-
Female	1	1	1	3	5	9	13	22	12	6	1	-
White, both sexes	1	2	2	3	8	14	24	27	21	6	1	-
Male	-	2	1	1	4	7	12	8	10	-	-	-
Female	1	-	1	2	4	7	12	19	11	6	1	-
Black, both sexes	-	2	-	-	1	2	3	2	-	-	-	-
Male	-	1	-	-	-	1	3	-	-	-	-	-
Female	-	1	-	-	1	1	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	1	4	1	1	-	-
Male	-	-	-	-	-	-	-	3	-	1	-	-
Female	-	-	-	1	-	1	1	1	1	-	-	-
Michigan												
All races, both sexes	5	11	14	15	18	42	63	47	41	15	1	-
Male	2	4	9	8	9	16	23	19	13	4	-	-
Female	3	7	5	7	9	26	40	28	28	11	1	-
White, both sexes	1	6	8	9	14	36	55	39	37	13	1	-
Male	-	2	4	8	7	13	20	16	11	4	-	-
Female	1	4	4	1	7	23	35	23	26	9	1	-
Black, both sexes	4	5	6	6	4	6	8	7	4	2	-	-
Male	2	2	5	-	2	3	3	2	2	-	-	-
Female	2	3	1	6	2	3	5	5	2	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Minnesota												
All races, both sexes	3	3	1	4	8	15	28	30	22	6	2	-
Male	1	1	-	2	4	9	17	11	8	4	1	-
Female	2	2	1	2	4	6	11	19	14	2	1	-
White, both sexes	3	2	1	3	5	11	26	30	22	6	2	-
Male	1	1	-	1	2	7	16	11	8	4	1	-
Female	2	1	1	2	3	4	10	19	14	2	1	-
Black, both sexes	-	1	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	2	1	-	-	-	-	-	-
Male	-	-	-	-	2	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	2	2	-	-	-	-	-
Male	-	-	-	-	-	2	1	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Mississippi												
All races, both sexes	10	12	12	16	13	22	20	16	22	2	2	-
Male	3	5	9	8	7	8	6	6	5	1	-	-
Female	7	7	3	8	6	14	14	10	17	1	2	-
White, both sexes	5	-	3	5	3	15	11	9	12	1	-	-
Male	2	-	1	3	1	7	3	4	2	-	-	-
Female	3	-	2	2	2	8	8	5	10	1	-	-
Black, both sexes	5	12	9	11	10	6	9	7	10	1	2	-
Male	1	5	8	5	6	1	3	2	3	1	-	-
Female	4	7	1	6	4	5	6	5	7	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Missouri												
All races, both sexes	5	7	6	11	17	23	35	42	24	12	-	-
Male	4	3	5	6	10	9	11	15	8	1	-	-
Female	1	4	1	5	7	14	24	27	16	11	-	-
White, both sexes	-	2	3	6	15	21	23	38	21	12	-	-
Male	-	-	3	3	8	9	6	14	8	1	-	-
Female	-	2	-	3	7	12	17	24	13	11	-	-
Black, both sexes	5	5	3	4	2	2	9	4	2	-	-	-
Male	4	3	2	3	2	-	3	1	-	-	-	-
Female	1	2	1	1	-	2	6	3	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	3	-	1	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	1	-	-	-
Montana												
All races, both sexes	-	1	1	-	1	1	5	4	4	1	-	-
Male	-	1	1	-	1	-	2	1	2	-	-	-
Female	-	-	-	-	-	1	3	3	2	1	-	-
White, both sexes	-	1	1	-	1	1	5	4	4	1	-	-
Male	-	1	1	-	1	-	2	1	2	-	-	-
Female	-	-	-	-	-	1	3	3	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Nebraska												
All races, both sexes	-	-	4	4	8	5	5	17	13	2	3	-
Male	-	-	2	2	4	2	4	9	6	-	-	-
Female	-	-	2	2	4	3	1	8	7	2	3	-
White, both sexes	-	-	2	3	7	5	5	15	13	2	3	-
Male	-	-	1	2	3	2	4	8	6	-	-	-
Female	-	-	1	1	4	3	1	7	7	2	3	-
Black, both sexes	-	-	2	-	1	-	-	2	-	-	-	-
Male	-	-	1	-	1	-	-	1	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1	4	1	2	5	4	2	10	5	3	-	-
Male	1	2	1	2	3	2	1	5	1	1	-	-
Female	-	2	-	-	2	2	1	5	4	2	-	-
White, both sexes	1	2	1	1	4	4	2	9	4	3	-	-
Male	1	2	1	1	2	2	1	5	-	1	-	-
Female	-	-	-	-	2	2	1	4	4	2	-	-
Black, both sexes	-	1	-	1	-	-	-	1	1	-	-	-
Male	-	-	-	1	-	-	-	-	1	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
New Hampshire												
All races, both sexes	-	-	-	-	-	7	5	5	6	4	3	-
Male	-	-	-	-	-	4	2	2	1	1	2	-
Female	-	-	-	-	-	3	3	3	5	3	1	-
White, both sexes	-	-	-	-	-	7	5	5	6	4	3	-
Male	-	-	-	-	-	4	2	2	1	1	2	-
Female	-	-	-	-	-	3	3	3	5	3	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	11	10	9	17	27	20	52	50	30	18	3	-
Male	9	8	5	8	16	8	24	12	4	5	-	-
Female	2	2	4	9	11	12	28	38	26	13	3	-
White, both sexes	3	2	5	6	17	13	43	40	21	18	3	-
Male	3	1	3	1	9	6	19	9	2	5	-	-
Female	-	1	2	5	8	7	24	31	19	13	3	-
Black, both sexes	8	8	4	10	8	6	7	8	8	-	-	-
Male	6	7	2	6	5	2	4	3	2	-	-	-
Female	2	1	2	4	3	4	3	5	6	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	1	2	2	1	-	-	-
Male	-	-	-	-	2	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	1	2	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
New Mexico												
All races, both sexes	5	1	1	3	6	6	6	14	4	-	-	-
Male	5	1	1	2	3	3	3	4	1	-	-	-
Female	-	-	-	1	3	3	3	10	3	-	-	-
White, both sexes	4	1	1	1	6	6	4	10	4	-	-	-
Male	4	1	1	1	3	3	2	3	1	-	-	-
Female	-	-	-	-	3	3	2	7	3	-	-	-
Black, both sexes	1	-	-	1	-	-	2	2	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	2	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	29	23	35	53	59	63	100	117	80	35	6	-
Male	15	15	19	29	27	28	46	49	32	13	2	-
Female	14	8	16	24	32	35	54	68	48	22	4	-
White, both sexes	10	11	21	31	31	43	74	93	70	27	6	-
Male	4	7	11	17	12	19	32	39	28	11	2	-
Female	6	4	10	14	19	24	42	54	42	16	4	-
Black, both sexes	19	10	14	19	24	18	20	21	9	6	-	-
Male	11	6	8	11	12	9	9	9	4	1	-	-
Female	8	4	6	8	12	9	11	12	5	5	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	3	4	2	6	3	1	2	-	-
Male	-	2	-	1	3	-	5	1	-	1	-	-
Female	-	-	-	2	1	2	1	2	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
North Carolina												
All races, both sexes	14	11	22	28	29	38	48	50	38	10	2	-
Male	9	9	10	12	15	15	19	7	8	1	1	-
Female	5	2	12	16	14	23	29	43	30	9	1	-
White, both sexes	2	4	7	7	13	20	30	31	24	7	1	-
Male	2	3	1	2	6	7	10	2	6	1	1	-
Female	-	1	6	5	7	13	20	29	18	6	-	-
Black, both sexes	12	6	15	20	16	16	18	18	13	3	1	-
Male	7	5	9	9	9	7	9	5	2	-	-	-
Female	5	1	6	11	7	9	9	13	11	3	1	-
American Indian 1/, both sexes	-	-	-	1	-	2	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	1	1	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
North Dakota												
All races, both sexes	1	-	2	3	2	2	8	10	9	1	-	-
Male	1	-	-	1	1	1	3	6	2	-	-	-
Female	-	-	2	2	1	1	5	4	7	1	-	-
White, both sexes	-	-	2	2	2	2	8	10	9	1	-	-
Male	-	-	-	1	1	1	3	6	2	-	-	-
Female	-	-	2	1	1	1	5	4	7	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Ohio												
All races, both sexes	5	16	24	37	44	68	104	105	77	19	5	-
Male	5	8	14	23	23	31	44	39	20	1	3	-
Female	-	8	10	14	21	37	60	66	57	18	2	-
White, both sexes	2	8	15	24	34	49	86	88	64	14	4	-
Male	2	3	8	16	19	24	35	33	15	-	2	-
Female	-	5	7	8	15	25	51	55	49	14	2	-
Black, both sexes	3	8	8	12	10	19	17	17	12	4	1	-
Male	3	5	5	6	4	7	9	6	4	-	1	-
Female	-	3	3	6	6	12	8	11	8	4	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	-	1	-	1	1	-	-
Male	-	-	1	1	-	-	-	-	1	1	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Oklahoma												
All races, both sexes	4	2	4	6	5	11	18	14	14	4	2	-
Male	2	-	1	-	2	5	10	4	3	-	1	-
Female	2	2	3	6	3	6	8	10	11	4	1	-
White, both sexes	2	1	2	2	2	5	16	13	12	3	2	-
Male	1	-	1	-	1	2	10	4	3	-	1	-
Female	1	1	1	2	1	3	6	9	9	3	1	-
Black, both sexes	1	1	2	3	2	1	1	1	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	1	2	3	1	1	1	1	1	-	-	-
American Indian 1/, both sexes	1	-	-	1	-	5	-	-	1	1	-	-
Male	1	-	-	-	-	3	-	-	-	-	-	-
Female	-	-	-	1	-	2	-	-	1	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Oregon												
All races, both sexes	-	4	3	3	6	10	16	34	20	6	2	-
Male	-	3	2	2	3	5	6	12	5	1	-	-
Female	-	1	1	1	3	5	10	22	15	5	2	-
White, both sexes	-	3	2	2	6	10	15	33	19	5	2	-
Male	-	3	1	1	3	5	5	12	5	-	-	-
Female	-	-	1	1	3	5	10	21	14	5	2	-
Black, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	1	-	-	-	1	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Pennsylvania												
All races, both sexes	12	7	15	15	24	40	73	74	60	20	6	-
Male	5	4	4	9	12	18	27	32	26	6	2	-
Female	7	3	11	6	12	22	46	42	34	14	4	-
White, both sexes	6	4	12	10	18	29	60	61	57	18	6	-
Male	3	3	3	5	9	13	23	27	25	6	2	-
Female	3	1	9	5	9	16	37	34	32	12	4	-
Black, both sexes	6	3	3	5	6	11	13	9	3	2	-	-
Male	2	1	1	4	3	5	4	3	1	-	-	-
Female	4	2	2	1	3	6	9	6	2	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	4	-	-	-	-
Male	-	-	-	-	-	-	-	2	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Rhode Island												
All races, both sexes	1	1	-	1	2	4	3	10	2	2	-	-
Male	1	-	-	1	-	1	3	2	-	1	-	-
Female	-	1	-	-	2	3	-	8	2	1	-	-
White, both sexes	1	1	-	1	2	3	3	10	2	2	-	-
Male	1	-	-	1	-	1	3	2	-	1	-	-
Female	-	1	-	-	2	2	-	8	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	14	11	15	20	12	22	26	37	17	4	1	-
Male	8	7	7	12	4	8	15	11	6	2	-	-
Female	6	4	8	8	8	14	11	26	11	2	1	-
White, both sexes	2	3	6	9	3	8	14	19	13	2	1	-
Male	1	1	3	5	1	3	8	6	5	-	-	-
Female	1	2	3	4	2	5	6	13	8	2	1	-
Black, both sexes	12	8	9	11	9	14	12	18	4	2	-	-
Male	7	6	4	7	3	5	7	5	1	2	-	-
Female	5	2	5	4	6	9	5	13	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
South Dakota												
All races, both sexes	-	-	-	1	-	1	11	10	4	3	-	-
Male	-	-	-	1	-	1	7	3	4	-	-	-
Female	-	-	-	-	-	-	4	7	-	3	-	-
White, both sexes	-	-	-	-	-	1	10	10	4	3	-	-
Male	-	-	-	-	-	1	7	3	4	-	-	-
Female	-	-	-	-	-	-	3	7	-	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	8	9	11	14	31	30	46	42	33	10	-	-
Male	7	6	5	9	19	13	19	12	10	4	-	-
Female	1	3	6	5	12	17	27	30	23	6	-	-
White, both sexes	2	3	4	8	22	24	35	27	26	8	-	-
Male	2	2	1	5	15	9	17	11	10	3	-	-
Female	-	1	3	3	7	15	18	16	16	5	-	-
Black, both sexes	6	5	7	6	9	6	10	15	7	2	-	-
Male	5	4	4	4	4	4	1	1	-	1	-	-
Female	1	1	3	2	5	2	9	14	7	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Texas												
All races, both sexes	33	38	34	46	65	95	104	123	74	30	3	-
Male	13	17	17	25	26	34	47	45	21	8	2	-
Female	20	21	17	21	39	61	57	78	53	22	1	-
White, both sexes	15	20	22	35	41	72	81	106	60	26	1	-
Male	7	9	13	20	15	28	40	41	16	8	1	-
Female	8	11	9	15	26	44	41	65	44	18	-	-
Black, both sexes	17	18	12	11	23	20	22	16	11	4	2	-
Male	6	8	4	5	11	4	7	4	4	-	1	-
Female	11	10	8	6	12	16	15	12	7	4	1	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	2	1	1	3	-	-	-
Male	-	-	-	-	-	2	-	-	1	-	-	-
Female	1	-	-	-	1	-	1	1	2	-	-	-
Utah												
All races, both sexes	-	1	3	2	2	6	7	7	6	2	-	-
Male	-	1	3	1	1	4	5	4	2	1	-	-
Female	-	-	-	1	1	2	2	3	4	1	-	-
White, both sexes	-	1	3	1	2	6	7	7	4	2	-	-
Male	-	1	3	-	1	4	5	4	1	1	-	-
Female	-	-	-	1	1	2	2	3	3	1	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Vermont												
All races, both sexes	-	1	-	1	1	4	6	2	2	-	-	-
Male	-	-	-	-	-	3	2	1	-	-	-	-
Female	-	1	-	1	1	1	4	1	2	-	-	-
White, both sexes	-	-	-	1	1	4	6	2	2	-	-	-
Male	-	-	-	-	-	3	2	1	-	-	-	-
Female	-	-	-	1	1	1	4	1	2	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	4	6	12	11	16	27	27	34	16	5	2	-
Male	4	3	6	5	8	15	12	8	2	1	-	-
Female	-	3	6	6	8	12	15	26	14	4	2	-
White, both sexes	2	2	3	3	6	17	15	21	8	3	1	-
Male	2	2	-	2	3	9	7	6	1	-	-	-
Female	-	-	3	1	3	8	8	15	7	3	1	-
Black, both sexes	2	4	8	8	9	10	12	12	8	2	1	-
Male	2	1	6	3	5	6	5	1	1	1	-	-
Female	-	3	2	5	4	4	7	11	7	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Washington												
All races, both sexes	2	5	5	6	14	33	48	58	40	15	4	-
Male	1	4	4	3	6	15	18	22	16	2	1	-
Female	1	1	1	3	8	18	30	36	24	13	3	-
White, both sexes	1	4	5	6	12	27	39	57	40	13	4	-
Male	1	3	4	3	5	13	17	22	16	1	1	-
Female	-	1	1	3	7	14	22	35	24	12	3	-
Black, both sexes	1	1	-	-	1	5	4	-	-	1	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	1	4	4	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	4	1	-	1	-	-
Male	-	-	-	-	1	1	1	-	-	1	-	-
Female	-	-	-	-	-	-	3	1	-	-	-	-
West Virginia												
All races, both sexes	-	3	5	3	7	10	13	16	4	4	1	-
Male	-	1	2	3	5	7	5	4	3	2	1	-
Female	-	2	3	-	2	3	8	12	1	2	-	-
White, both sexes	-	2	5	2	6	9	11	15	4	4	1	-
Male	-	1	2	2	4	6	5	4	3	2	1	-
Female	-	1	3	-	2	3	6	11	1	2	-	-
Black, both sexes	-	1	-	1	1	1	2	1	-	-	-	-
Male	-	-	-	1	1	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease and secondary hypertension (I12,I15)												
Wisconsin												
All races, both sexes	4	4	5	9	10	17	34	36	30	9	1	-
Male	2	2	2	3	6	10	13	16	8	3	-	-
Female	2	2	3	6	4	7	21	20	22	6	1	-
White, both sexes	2	1	2	7	8	15	33	35	30	9	1	-
Male	-	-	1	2	5	8	13	15	8	3	-	-
Female	2	1	1	5	3	7	20	20	22	6	1	-
Black, both sexes	2	2	3	2	2	2	1	1	-	-	-	-
Male	2	2	1	1	1	2	-	1	-	-	-	-
Female	-	-	2	1	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	2	-	2	3	1	-	1	1	-	-
Male	-	-	2	-	2	3	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	1	-	-
White, both sexes	-	-	2	-	2	3	1	-	1	1	-	-
Male	-	-	2	-	2	3	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
United States												
All races, both sexes	2,987	-	-	-	-	1	4	10	22	35	42	77
Male	1,318	-	-	-	-	1	3	7	10	26	30	54
Female	1,669	-	-	-	-	-	1	3	12	9	12	23
White, both sexes	2,064	-	-	-	-	1	-	5	7	10	10	27
Male	864	-	-	-	-	1	-	2	4	8	6	21
Female	1,200	-	-	-	-	-	-	3	3	2	4	6
Black, both sexes	825	-	-	-	-	-	4	4	14	25	31	49
Male	412	-	-	-	-	-	3	4	6	18	24	32
Female	413	-	-	-	-	-	1	-	8	7	7	17
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	90	-	-	-	-	-	-	1	1	-	1	1
Male	38	-	-	-	-	-	-	1	-	-	-	1
Female	52	-	-	-	-	-	-	-	1	-	1	-
Alabama												
All races, both sexes	37	-	-	-	-	-	-	-	-	-	-	4
Male	15	-	-	-	-	-	-	-	-	-	-	3
Female	22	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	18	-	-	-	-	-	-	-	-	-	-	4
Male	8	-	-	-	-	-	-	-	-	-	-	3
Female	10	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Alaska												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	29	-	-	-	-	-	-	-	1	1	-	-
Male	15	-	-	-	-	-	-	-	1	1	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	29	-	-	-	-	-	-	-	1	1	-	-
Male	15	-	-	-	-	-	-	-	1	1	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Arkansas												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	481	-	-	-	-	1	-	3	1	3	7	8
Male	236	-	-	-	-	1	-	2	-	3	5	5
Female	245	-	-	-	-	-	-	1	1	-	2	3
White, both sexes	337	-	-	-	-	1	-	1	-	-	1	6
Male	159	-	-	-	-	1	-	-	-	-	1	5
Female	178	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	82	-	-	-	-	-	-	1	1	3	5	2
Male	47	-	-	-	-	-	-	1	-	3	4	-
Female	35	-	-	-	-	-	-	-	1	-	1	2
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	59	-	-	-	-	-	-	1	-	-	1	-
Male	28	-	-	-	-	-	-	1	-	-	-	-
Female	31	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Colorado												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Delaware												
All races, both sexes	14	-	-	-	-	-	-	-	1	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Dist. of Columbia												
All races, both sexes	26	-	-	-	-	-	-	1	-	1	1	1
Male	12	-	-	-	-	-	-	-	-	-	1	1
Female	14	-	-	-	-	-	-	1	-	1	-	-
White, both sexes	3	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	23	-	-	-	-	-	-	-	-	1	1	1
Male	11	-	-	-	-	-	-	-	-	-	1	1
Female	12	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Florida												
All races, both sexes	174	-	-	-	-	-	-	-	-	1	4	6
Male	81	-	-	-	-	-	-	-	-	-	3	3
Female	93	-	-	-	-	-	-	-	-	1	1	3
White, both sexes	125	-	-	-	-	-	-	-	-	-	-	1
Male	63	-	-	-	-	-	-	-	-	-	-	1
Female	62	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	47	-	-	-	-	-	-	-	-	1	4	5
Male	17	-	-	-	-	-	-	-	-	-	3	2
Female	30	-	-	-	-	-	-	-	-	1	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	87	-	-	-	-	-	1	-	-	1	2	1
Male	44	-	-	-	-	-	-	-	-	1	1	1
Female	43	-	-	-	-	-	1	-	-	-	1	-
White, both sexes	37	-	-	-	-	-	-	-	-	-	1	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	50	-	-	-	-	-	1	-	-	1	1	1
Male	29	-	-	-	-	-	-	-	-	1	1	1
Female	21	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Hawaii												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	11	-	-	-	-	-	-	1	-	-	-	-
Male	8	-	-	-	-	-	-	1	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	1	-	-	-	-
Male	8	-	-	-	-	-	-	1	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Illinois												
All races, both sexes	106	-	-	-	-	-	-	-	-	3	1	4
Male	44	-	-	-	-	-	-	-	-	2	-	2
Female	62	-	-	-	-	-	-	-	-	1	1	2
White, both sexes	65	-	-	-	-	-	-	-	-	-	-	-
Male	23	-	-	-	-	-	-	-	-	-	-	-
Female	42	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	40	-	-	-	-	-	-	-	-	3	1	4
Male	20	-	-	-	-	-	-	-	-	2	-	2
Female	20	-	-	-	-	-	-	-	-	1	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	47	-	-	-	-	-	-	-	-	-	1	-
Male	15	-	-	-	-	-	-	-	-	-	1	-
Female	32	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	39	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Iowa												
All races, both sexes	44	-	-	-	-	-	-	-	1	-	-	-
Male	13	-	-	-	-	-	-	-	1	-	-	-
Female	31	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	43	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	31	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	10	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Kentucky												
All races, both sexes	35	-	-	-	-	-	-	-	2	-	-	3
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	26	-	-	-	-	-	-	-	2	-	-	3
White, both sexes	27	-	-	-	-	-	-	-	1	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	8	-	-	-	-	-	-	-	1	-	-	2
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	1	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	85	-	-	-	-	-	-	-	1	1	3	7
Male	37	-	-	-	-	-	-	-	1	1	3	6
Female	48	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	35	-	-	-	-	-	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	20	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	50	-	-	-	-	-	-	-	1	1	3	6
Male	22	-	-	-	-	-	-	-	1	1	3	5
Female	28	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Maine												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	77	-	-	-	-	-	-	-	1	1	1	4
Male	39	-	-	-	-	-	-	-	1	-	-	4
Female	38	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	36	-	-	-	-	-	-	-	-	-	1	2
Male	20	-	-	-	-	-	-	-	-	-	-	2
Female	16	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	41	-	-	-	-	-	-	-	1	1	-	2
Male	19	-	-	-	-	-	-	-	1	-	-	2
Female	22	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Massachusetts												
All races, both sexes	27	-	-	-	-	-	-	-	-	-	2	1
Male	12	-	-	-	-	-	-	-	-	-	2	1
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	26	-	-	-	-	-	-	-	-	-	1	1
Male	11	-	-	-	-	-	-	-	-	-	1	1
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	125	-	-	-	-	-	-	2	2	2	1	3
Male	48	-	-	-	-	-	-	2	1	2	1	3
Female	77	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	90	-	-	-	-	-	-	1	1	-	-	1
Male	29	-	-	-	-	-	-	1	1	-	-	1
Female	61	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	34	-	-	-	-	-	-	1	1	2	1	2
Male	19	-	-	-	-	-	-	1	-	2	1	2
Female	15	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Minnesota												
All races, both sexes	32	-	-	-	-	-	-	-	-	-	1	-
Male	13	-	-	-	-	-	-	-	-	-	1	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	75	-	-	-	-	-	-	-	2	1	2	3
Male	35	-	-	-	-	-	-	-	1	1	2	2
Female	40	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	34	-	-	-	-	-	-	-	1	-	1	-
Male	19	-	-	-	-	-	-	-	1	-	1	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	39	-	-	-	-	-	-	-	1	1	1	3
Male	15	-	-	-	-	-	-	-	-	1	1	2
Female	24	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Missouri												
All races, both sexes	47	-	-	-	-	-	-	-	1	-	-	1
Male	17	-	-	-	-	-	-	-	1	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	34	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	13	-	-	-	-	-	-	-	1	-	-	-
Male	6	-	-	-	-	-	-	-	1	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Nebraska												
All races, both sexes	21	-	-	-	-	-	-	-	-	-	1	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	21	-	-	-	-	-	-	-	-	-	1	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	10	-	-	-	-	-	-	-	-	1	1	1
Male	10	-	-	-	-	-	-	-	-	1	1	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	1	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
New Hampshire												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	65	-	-	-	-	-	1	-	1	2	-	1
Male	25	-	-	-	-	-	1	-	-	2	-	-
Female	40	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	39	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	23	-	-	-	-	-	1	-	1	2	-	1
Male	10	-	-	-	-	-	1	-	-	2	-	-
Female	13	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
New Mexico												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	230	-	-	-	-	-	2	1	4	5	7	6
Male	106	-	-	-	-	-	2	1	2	2	5	6
Female	124	-	-	-	-	-	-	-	2	3	2	-
White, both sexes	151	-	-	-	-	-	-	-	2	2	2	2
Male	65	-	-	-	-	-	-	-	-	1	2	2
Female	86	-	-	-	-	-	-	-	2	1	-	-
Black, both sexes	76	-	-	-	-	-	2	1	2	3	5	4
Male	41	-	-	-	-	-	2	1	2	1	3	4
Female	35	-	-	-	-	-	-	-	-	2	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
North Carolina												
All races, both sexes	67	-	-	-	-	-	-	-	-	1	2	1
Male	34	-	-	-	-	-	-	-	-	1	1	1
Female	33	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	26	-	-	-	-	-	-	-	-	1	-	-
Male	12	-	-	-	-	-	-	-	-	1	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	41	-	-	-	-	-	-	-	-	-	2	1
Male	22	-	-	-	-	-	-	-	-	-	1	1
Female	19	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Ohio												
All races, both sexes	169	-	-	-	-	-	-	1	-	1	1	4
Male	64	-	-	-	-	-	-	-	-	1	1	3
Female	105	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	130	-	-	-	-	-	-	1	-	-	1	3
Male	44	-	-	-	-	-	-	-	-	-	1	2
Female	86	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	38	-	-	-	-	-	-	-	-	1	-	1
Male	19	-	-	-	-	-	-	-	-	1	-	1
Female	19	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	30	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Oregon												
All races, both sexes	32	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	21	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	30	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	88	-	-	-	-	-	-	-	-	1	-	2
Male	34	-	-	-	-	-	-	-	-	-	-	2
Female	54	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	72	-	-	-	-	-	-	-	-	-	-	-
Male	26	-	-	-	-	-	-	-	-	-	-	-
Female	46	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	16	-	-	-	-	-	-	-	-	1	-	2
Male	8	-	-	-	-	-	-	-	-	-	-	2
Female	8	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Rhode Island												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	58	-	-	-	-	-	-	-	1	1	1	-
Male	28	-	-	-	-	-	-	-	-	-	1	-
Female	30	-	-	-	-	-	-	-	1	1	-	-
White, both sexes	28	-	-	-	-	-	-	-	-	1	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	30	-	-	-	-	-	-	-	1	-	1	-
Male	17	-	-	-	-	-	-	-	-	-	1	-
Female	13	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
South Dakota												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	107	-	-	-	-	-	-	-	1	1	1	4
Male	46	-	-	-	-	-	-	-	-	1	1	4
Female	61	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	71	-	-	-	-	-	-	-	-	1	-	1
Male	27	-	-	-	-	-	-	-	-	1	-	1
Female	44	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	36	-	-	-	-	-	-	-	1	-	1	3
Male	19	-	-	-	-	-	-	-	-	-	1	3
Female	17	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Texas												
All races, both sexes	168	-	-	-	-	-	-	-	1	5	1	5
Male	73	-	-	-	-	-	-	-	-	5	-	2
Female	95	-	-	-	-	-	-	-	1	-	1	3
White, both sexes	123	-	-	-	-	-	-	-	-	3	1	4
Male	49	-	-	-	-	-	-	-	-	3	-	2
Female	74	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	43	-	-	-	-	-	-	-	1	2	-	1
Male	22	-	-	-	-	-	-	-	-	2	-	-
Female	21	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Vermont												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	61	-	-	-	-	-	-	1	-	1	1	1
Male	20	-	-	-	-	-	-	1	-	1	-	-
Female	41	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	34	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	26	-	-	-	-	-	-	1	-	1	1	1
Male	11	-	-	-	-	-	-	1	-	1	-	-
Female	15	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Washington												
All races, both sexes	107	-	-	-	-	-	-	-	-	-	-	2
Male	51	-	-	-	-	-	-	-	-	-	-	1
Female	56	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	95	-	-	-	-	-	-	-	-	-	-	1
Male	47	-	-	-	-	-	-	-	-	-	-	1
Female	48	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	15	-	-	-	-	-	-	-	1	-	-	-
Male	6	-	-	-	-	-	-	-	1	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	1	-	-	-
Male	6	-	-	-	-	-	-	-	1	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Wisconsin												
All races, both sexes	53	-	-	-	-	-	-	-	-	-	-	1
Male	26	-	-	-	-	-	-	-	-	-	-	1
Female	27	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	51	-	-	-	-	-	-	-	-	-	-	1
Male	25	-	-	-	-	-	-	-	-	-	-	1
Female	26	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
United States												
All races, both sexes	101	134	159	172	207	329	463	557	447	191	36	-
Male	66	86	92	94	97	148	196	214	131	54	9	-
Female	35	48	67	78	110	181	267	343	316	137	27	-
White, both sexes	41	54	78	89	124	230	352	454	378	172	32	-
Male	31	35	40	53	58	104	157	179	108	49	8	-
Female	10	19	38	36	66	126	195	275	270	123	24	-
Black, both sexes	56	76	77	75	77	85	94	87	50	18	3	-
Male	31	49	50	36	37	38	34	31	14	5	-	-
Female	25	27	27	39	40	47	60	56	36	13	3	-
American Indian 1/, both sexes	1	1	-	-	-	2	1	-	3	-	-	-
Male	1	1	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	-	-	1	1	-	2	-	-	-
Asian or Pacific Islander 2/, both sexes	3	3	4	8	6	12	16	16	16	1	1	-
Male	3	1	2	5	2	5	5	4	8	-	1	-
Female	-	2	2	3	4	7	11	12	8	1	-	-
Alabama												
All races, both sexes	-	4	3	1	1	6	6	4	6	2	-	-
Male	-	2	2	-	-	3	4	-	1	-	-	-
Female	-	2	1	1	1	3	2	4	5	2	-	-
White, both sexes	-	1	2	-	-	5	3	3	3	2	-	-
Male	-	1	1	-	-	2	2	-	1	-	-	-
Female	-	-	1	-	-	3	1	3	2	2	-	-
Black, both sexes	-	3	1	1	1	1	3	1	3	-	-	-
Male	-	1	1	-	-	1	2	-	-	-	-	-
Female	-	2	-	1	1	-	1	1	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	3	2	1	2	-	3	4	9	1	2	-	-
Male	2	2	-	2	-	-	-	5	-	2	-	-
Female	1	-	1	-	-	3	4	4	1	-	-	-
White, both sexes	3	2	1	2	-	3	4	9	1	2	-	-
Male	2	2	-	2	-	-	-	5	-	2	-	-
Female	1	-	1	-	-	3	4	4	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Arkansas												
All races, both sexes	1	-	2	2	1	1	4	2	7	-	-	-
Male	1	-	2	-	-	-	2	-	1	-	-	-
Female	-	-	-	2	1	1	2	2	6	-	-	-
White, both sexes	1	-	1	-	1	-	4	1	6	-	-	-
Male	1	-	1	-	-	-	2	-	1	-	-	-
Female	-	-	-	-	1	-	2	1	5	-	-	-
Black, both sexes	-	-	1	2	-	1	-	1	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	2	-	1	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	16	21	32	26	29	56	72	90	80	33	3	-
Male	12	9	18	18	14	30	34	42	31	11	1	-
Female	4	12	14	8	15	26	38	48	49	22	2	-
White, both sexes	8	9	20	16	21	38	48	72	65	30	1	-
Male	7	4	10	11	9	22	22	33	24	10	-	-
Female	1	5	10	5	12	16	26	39	41	20	1	-
Black, both sexes	5	8	9	4	5	6	12	11	6	3	1	-
Male	2	3	7	2	4	4	7	7	2	1	-	-
Female	3	5	2	2	1	2	5	4	4	2	1	-
American Indian 1/, both sexes	-	1	-	-	-	1	-	-	1	-	-	-
Male	-	1	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	3	3	6	3	11	12	7	8	-	1	-
Male	3	1	1	5	1	4	5	2	4	-	1	-
Female	-	2	2	1	2	7	7	5	4	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Colorado												
All races, both sexes	1	2	-	1	2	2	4	4	1	1	-	-
Male	-	1	-	-	2	-	3	2	1	-	-	-
Female	1	1	-	1	-	2	1	2	-	1	-	-
White, both sexes	1	2	-	1	2	2	4	4	1	1	-	-
Male	-	1	-	-	2	-	3	2	1	-	-	-
Female	1	1	-	1	-	2	1	2	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	2	-	1	-	4	6	2	2	2	1	-
Male	-	-	-	-	-	2	1	1	1	1	1	-
Female	-	2	-	1	-	2	5	1	1	1	-	-
White, both sexes	-	1	-	-	-	4	6	2	2	2	1	-
Male	-	-	-	-	-	2	1	1	1	1	1	-
Female	-	1	-	-	-	2	5	1	1	1	-	-
Black, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Delaware												
All races, both sexes	1	-	-	2	2	2	1	3	1	1	-	-
Male	1	-	-	2	-	1	-	2	1	1	-	-
Female	-	-	-	-	2	1	1	1	-	-	-	-
White, both sexes	-	-	-	-	1	1	1	2	1	1	-	-
Male	-	-	-	-	-	1	-	1	1	1	-	-
Female	-	-	-	-	1	-	1	1	-	-	-	-
Black, both sexes	1	-	-	2	1	1	-	1	-	-	-	-
Male	1	-	-	2	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	3	3	1	3	2	3	2	4	1	-	-	-
Male	3	2	1	1	-	1	-	1	1	-	-	-
Female	-	1	-	2	2	2	2	3	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	2	3	1	3	2	3	2	3	1	-	-	-
Male	2	2	1	1	-	1	-	1	1	-	-	-
Female	-	1	-	2	2	2	2	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Florida												
All races, both sexes	6	6	10	9	11	26	23	32	26	8	6	-
Male	3	3	5	5	5	12	8	16	13	2	3	-
Female	3	3	5	4	6	14	15	16	13	6	3	-
White, both sexes	3	2	6	6	8	23	20	26	19	5	6	-
Male	1	1	4	4	3	12	8	15	9	2	3	-
Female	2	1	2	2	5	11	12	11	10	3	3	-
Black, both sexes	3	4	4	3	3	3	3	6	5	3	-	-
Male	2	2	1	1	2	-	-	1	3	-	-	-
Female	1	2	3	2	1	3	3	5	2	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Georgia												
All races, both sexes	5	9	9	7	6	12	6	15	9	3	1	-
Male	5	6	7	5	5	6	1	3	1	1	1	-
Female	-	3	2	2	1	6	5	12	8	2	-	-
White, both sexes	2	2	1	2	3	6	2	10	6	1	1	-
Male	2	-	1	2	2	3	1	1	1	1	1	-
Female	-	2	-	-	1	3	1	9	5	-	-	-
Black, both sexes	3	7	8	5	3	6	4	5	3	2	-	-
Male	3	6	6	3	3	3	-	2	-	-	-	-
Female	-	1	2	2	-	3	4	3	3	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Hawaii												
All races, both sexes	-	-	1	-	1	-	2	2	4	2	-	-
Male	-	-	1	-	1	-	-	-	3	1	-	-
Female	-	-	-	-	-	-	2	2	1	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	-	2	2	3	1	-	-
Male	-	-	1	-	1	-	-	-	3	-	-	-
Female	-	-	-	-	-	-	2	2	-	1	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	2	5	2	1	-	-
Male	-	-	-	-	-	-	2	2	2	1	-	-
Female	-	-	-	-	-	-	-	3	-	-	-	-
White, both sexes	-	-	-	-	-	-	2	5	2	1	-	-
Male	-	-	-	-	-	-	2	2	2	1	-	-
Female	-	-	-	-	-	-	-	3	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Illinois												
All races, both sexes	1	5	4	4	7	15	23	17	15	7	-	-
Male	-	3	2	2	5	7	9	6	4	2	-	-
Female	1	2	2	2	2	8	14	11	11	5	-	-
White, both sexes	1	-	1	2	5	6	17	13	13	7	-	-
Male	-	-	1	1	3	3	6	4	3	2	-	-
Female	1	-	-	1	2	3	11	9	10	5	-	-
Black, both sexes	-	5	3	2	2	9	6	3	2	-	-	-
Male	-	3	1	1	2	4	3	1	1	-	-	-
Female	-	2	2	1	-	5	3	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1	1	-	7	1	4	6	11	9	5	1	-
Male	-	-	-	4	1	-	5	4	-	-	-	-
Female	1	1	-	3	-	4	1	7	9	5	1	-
White, both sexes	-	1	-	5	-	4	5	9	9	5	1	-
Male	-	-	-	4	-	-	4	3	-	-	-	-
Female	-	1	-	1	-	4	1	6	9	5	1	-
Black, both sexes	1	-	-	2	1	-	1	2	-	-	-	-
Male	-	-	-	-	1	-	1	1	-	-	-	-
Female	1	-	-	2	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Iowa												
All races, both sexes	1	1	-	1	-	4	6	11	8	8	3	-
Male	1	-	-	-	-	1	2	6	1	1	-	-
Female	-	1	-	1	-	3	4	5	7	7	3	-
White, both sexes	1	1	-	1	-	4	6	11	8	8	3	-
Male	1	-	-	-	-	1	2	6	1	1	-	-
Female	-	1	-	1	-	3	4	5	7	7	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	1	-	1	1	1	4	1	-	-
Male	-	-	-	-	-	-	1	1	-	1	-	-
Female	-	-	-	1	-	1	-	-	4	-	-	-
White, both sexes	-	-	-	1	-	1	1	1	4	1	-	-
Male	-	-	-	-	-	-	1	1	-	1	-	-
Female	-	-	-	1	-	1	-	-	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Kentucky												
All races, both sexes	1	2	1	2	4	4	6	4	5	1	-	-
Male	-	-	1	-	2	1	3	1	1	-	-	-
Female	1	2	-	2	2	3	3	3	4	1	-	-
White, both sexes	1	1	1	2	1	4	5	4	5	1	-	-
Male	-	-	1	-	1	1	3	1	1	-	-	-
Female	1	1	-	2	-	3	2	3	4	1	-	-
Black, both sexes	-	1	-	-	3	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	1	-	-	2	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	1	7	10	9	6	13	11	13	3	-	-
Male	-	-	4	5	4	2	7	2	1	1	-	-
Female	-	1	3	5	5	4	6	9	12	2	-	-
White, both sexes	-	-	3	2	3	2	9	4	9	2	-	-
Male	-	-	2	1	2	2	5	1	1	-	-	-
Female	-	-	1	1	1	-	4	3	8	2	-	-
Black, both sexes	-	1	4	8	6	4	4	7	4	1	-	-
Male	-	-	2	4	2	-	2	1	-	1	-	-
Female	-	1	2	4	4	4	2	6	4	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Maine												
All races, both sexes	-	-	-	-	-	4	3	-	-	3	-	-
Male	-	-	-	-	-	2	3	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	3	-	-
White, both sexes	-	-	-	-	-	4	3	-	-	3	-	-
Male	-	-	-	-	-	2	3	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	3	8	6	5	13	8	11	8	6	2	-	-
Male	1	7	4	3	6	5	5	1	2	-	-	-
Female	2	1	2	2	7	3	6	7	4	2	-	-
White, both sexes	1	2	1	3	3	6	3	7	5	2	-	-
Male	1	2	1	2	2	4	3	1	2	-	-	-
Female	-	-	-	1	1	2	-	6	3	2	-	-
Black, both sexes	2	6	5	2	10	2	8	1	1	-	-	-
Male	-	5	3	1	4	1	2	-	-	-	-	-
Female	2	1	2	1	6	1	6	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Massachusetts												
All races, both sexes	2	-	-	-	1	3	3	7	6	2	-	-
Male	2	-	-	-	-	2	2	1	1	1	-	-
Female	-	-	-	-	1	1	1	6	5	1	-	-
White, both sexes	2	-	-	-	1	3	3	7	6	2	-	-
Male	2	-	-	-	-	2	2	1	1	1	-	-
Female	-	-	-	-	1	1	1	6	5	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	4	6	1	8	7	12	23	23	17	8	6	-
Male	2	6	1	5	3	2	7	5	5	2	1	-
Female	2	-	-	3	4	10	16	18	12	6	5	-
White, both sexes	-	1	-	3	4	9	22	19	15	8	6	-
Male	-	1	-	2	1	2	7	5	5	2	1	-
Female	-	-	-	1	3	7	15	14	10	6	5	-
Black, both sexes	4	5	1	5	3	3	1	4	1	-	-	-
Male	2	5	1	3	2	-	-	-	-	-	-	-
Female	2	-	-	2	1	3	1	4	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Minnesota												
All races, both sexes	1	-	1	1	2	1	6	6	5	7	1	-
Male	-	-	1	-	1	1	3	4	-	2	-	-
Female	1	-	-	1	1	-	3	2	5	5	1	-
White, both sexes	1	-	-	1	2	-	6	5	5	7	1	-
Male	-	-	-	-	1	-	3	3	-	2	-	-
Female	1	-	-	1	1	-	3	2	5	5	1	-
Black, both sexes	-	-	1	-	-	1	-	1	-	-	-	-
Male	-	-	1	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	4	3	5	6	10	10	9	11	7	2	-	-
Male	4	1	3	4	4	6	4	3	-	-	-	-
Female	-	2	2	2	6	4	5	8	7	2	-	-
White, both sexes	1	-	3	3	6	5	5	3	5	1	-	-
Male	1	-	2	3	3	4	4	-	-	-	-	-
Female	-	-	1	-	3	1	1	3	5	1	-	-
Black, both sexes	2	3	2	3	3	5	4	8	2	1	-	-
Male	2	1	1	1	1	2	-	3	-	-	-	-
Female	-	2	1	2	2	3	4	5	2	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Missouri												
All races, both sexes	2	-	4	1	1	8	4	12	9	4	-	-
Male	1	-	2	1	1	2	1	5	1	2	-	-
Female	1	-	2	-	-	6	3	7	8	2	-	-
White, both sexes	-	-	-	1	-	5	3	12	8	4	-	-
Male	-	-	-	1	-	1	1	5	1	2	-	-
Female	-	-	-	-	-	4	2	7	7	2	-	-
Black, both sexes	2	-	4	-	1	3	1	-	1	-	-	-
Male	1	-	2	-	1	1	-	-	-	-	-	-
Female	1	-	2	-	-	2	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	2	1	1	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	1	-	-
White, both sexes	-	-	-	-	-	-	-	2	1	1	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Nebraska												
All races, both sexes	-	-	-	-	-	1	4	6	3	5	1	-
Male	-	-	-	-	-	-	4	4	1	-	1	-
Female	-	-	-	-	-	1	-	2	2	5	-	-
White, both sexes	-	-	-	-	-	1	4	6	3	5	1	-
Male	-	-	-	-	-	-	4	4	1	-	1	-
Female	-	-	-	-	-	1	-	2	2	5	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1	-	2	2	-	-	1	1	-	-	-	-
Male	1	-	2	2	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	2	-	-	1	1	-	-	-	-
Male	1	-	-	2	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	2	-	-	-	-	-	-	-	-	-
Male	-	-	2	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
New Hampshire												
All races, both sexes	-	-	1	-	1	1	-	1	-	1	-	-
Male	-	-	-	-	1	-	-	1	-	1	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	1	-	1	1	-	1	-	1	-	-
Male	-	-	-	-	1	-	-	1	-	1	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	3	5	3	2	1	7	6	18	13	2	-	-
Male	3	2	-	1	1	1	3	4	6	1	-	-
Female	-	3	3	1	-	6	3	14	7	1	-	-
White, both sexes	-	-	2	-	1	6	6	11	11	2	-	-
Male	-	-	-	-	1	1	3	3	6	1	-	-
Female	-	-	2	-	-	5	3	8	5	1	-	-
Black, both sexes	3	5	1	1	-	1	-	5	2	-	-	-
Male	3	2	-	1	-	-	-	1	-	-	-	-
Female	-	3	1	-	-	1	-	4	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
New Mexico												
All races, both sexes	-	2	-	2	1	-	3	2	1	1	-	-
Male	-	2	-	1	-	-	1	1	1	-	-	-
Female	-	-	-	1	1	-	2	1	-	1	-	-
White, both sexes	-	1	-	2	-	-	3	2	1	1	-	-
Male	-	1	-	1	-	-	1	1	1	-	-	-
Female	-	-	-	1	-	-	2	1	-	1	-	-
Black, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	6	12	24	12	14	30	38	33	24	10	2	-
Male	4	9	13	4	6	17	15	13	4	3	-	-
Female	2	3	11	8	8	13	23	20	20	7	2	-
White, both sexes	3	8	11	5	9	20	27	27	22	10	1	-
Male	2	6	5	2	4	13	9	12	4	3	-	-
Female	1	2	6	3	5	7	18	15	18	7	1	-
Black, both sexes	3	4	13	6	5	10	10	5	2	-	1	-
Male	2	3	8	2	2	4	6	1	-	-	-	-
Female	1	1	5	4	3	6	4	4	2	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
North Carolina												
All races, both sexes	3	6	3	4	8	9	9	13	7	-	1	-
Male	3	4	2	2	5	6	3	4	2	-	-	-
Female	-	2	1	2	3	3	6	9	5	-	1	-
White, both sexes	-	2	1	-	1	4	4	8	4	-	1	-
Male	-	2	-	-	1	3	1	3	1	-	-	-
Female	-	-	1	-	-	1	3	5	3	-	1	-
Black, both sexes	3	4	2	4	7	5	5	5	3	-	-	-
Male	3	2	2	2	4	3	2	1	1	-	-	-
Female	-	2	-	2	3	2	3	4	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	2	1	1	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	1	1	-	-
White, both sexes	-	-	-	-	-	-	2	1	1	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Ohio												
All races, both sexes	4	5	5	8	21	18	21	34	29	14	3	-
Male	-	5	2	4	6	10	8	12	9	3	-	-
Female	4	-	3	4	15	8	13	22	20	11	3	-
White, both sexes	-	2	2	4	15	11	18	29	28	13	3	-
Male	-	2	1	1	5	5	7	9	9	2	-	-
Female	-	-	1	3	10	6	11	20	19	11	3	-
Black, both sexes	4	3	3	4	6	6	3	5	1	1	-	-
Male	-	3	1	3	1	4	1	3	-	1	-	-
Female	4	-	2	1	5	2	2	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	2	1	2	4	1	6	7	4	2	-	-
Male	-	2	-	1	2	-	2	3	2	-	-	-
Female	-	-	1	1	2	1	4	4	2	2	-	-
White, both sexes	-	-	-	2	4	1	5	6	4	1	-	-
Male	-	-	-	1	2	-	2	3	2	-	-	-
Female	-	-	-	1	2	1	3	3	2	1	-	-
Black, both sexes	-	2	1	-	-	-	-	1	-	1	-	-
Male	-	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Oregon												
All races, both sexes	-	2	1	3	3	3	7	4	6	2	-	-
Male	-	1	-	1	2	1	2	-	1	2	-	-
Female	-	1	1	2	1	2	5	4	5	-	-	-
White, both sexes	-	1	1	3	3	3	7	4	6	2	-	-
Male	-	-	-	1	2	1	2	-	1	2	-	-
Female	-	1	1	2	1	2	5	4	5	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	3	2	2	4	7	4	18	22	16	6	1	-
Male	3	1	1	3	2	1	7	11	2	1	-	-
Female	-	1	1	1	5	3	11	11	14	5	1	-
White, both sexes	3	1	1	2	4	4	14	21	15	6	1	-
Male	3	-	-	2	1	1	6	10	2	1	-	-
Female	-	1	1	-	3	3	8	11	13	5	1	-
Black, both sexes	-	1	1	2	3	-	4	1	1	-	-	-
Male	-	1	1	1	1	-	1	1	-	-	-	-
Female	-	-	-	1	2	-	3	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Rhode Island												
All races, both sexes	-	1	1	-	-	-	2	-	-	-	-	-
Male	-	1	-	-	-	-	2	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	1	-	-	-	2	-	-	-	-	-
Male	-	1	-	-	-	-	2	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	4	3	2	7	7	5	9	8	6	3	1	-
Male	2	3	2	5	4	2	3	-	5	1	-	-
Female	2	-	-	2	3	3	6	8	1	2	1	-
White, both sexes	-	2	-	4	3	2	2	8	4	1	1	-
Male	-	2	-	3	2	1	-	-	3	-	-	-
Female	-	-	-	1	1	1	2	8	1	1	1	-
Black, both sexes	4	1	2	3	4	3	7	-	2	2	-	-
Male	2	1	2	2	2	1	3	-	2	1	-	-
Female	2	-	-	1	2	2	4	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
South Dakota												
All races, both sexes	-	-	-	-	1	1	1	1	4	1	-	-
Male	-	-	-	-	1	-	1	-	1	-	-	-
Female	-	-	-	-	-	1	-	1	3	1	-	-
White, both sexes	-	-	-	-	1	1	1	1	4	1	-	-
Male	-	-	-	-	1	-	1	-	1	-	-	-
Female	-	-	-	-	-	1	-	1	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	3	3	4	5	7	14	21	20	17	6	-	-
Male	1	1	3	1	2	9	5	9	7	2	-	-
Female	2	2	1	4	5	5	16	11	10	4	-	-
White, both sexes	-	1	1	3	4	9	16	19	11	5	-	-
Male	-	-	1	-	1	4	4	9	4	2	-	-
Female	-	1	-	3	3	5	12	10	7	3	-	-
Black, both sexes	3	2	3	2	3	5	5	1	6	1	-	-
Male	1	1	2	1	1	5	1	-	3	-	-	-
Female	2	1	1	1	2	-	4	1	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Texas												
All races, both sexes	13	7	12	11	11	17	31	25	20	8	1	-
Male	8	6	7	5	7	8	13	6	3	3	-	-
Female	5	1	5	6	4	9	18	19	17	5	1	-
White, both sexes	5	4	10	6	7	12	27	18	19	6	1	-
Male	4	3	5	3	4	5	12	3	3	2	-	-
Female	1	1	5	3	3	7	15	15	16	4	1	-
Black, both sexes	8	3	2	5	4	4	4	6	1	2	-	-
Male	4	3	2	2	3	2	1	2	-	1	-	-
Female	4	-	-	3	1	2	3	4	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	1	-	-	4	3	1	1	-	-
Male	-	-	-	1	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	2	3	1	1	-	-
White, both sexes	-	-	-	1	-	-	4	2	-	1	-	-
Male	-	-	-	1	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	2	2	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Vermont												
All races, both sexes	-	-	-	-	-	-	1	-	2	2	-	-
Male	-	-	-	-	-	-	-	-	1	1	-	-
Female	-	-	-	-	-	-	1	-	1	1	-	-
White, both sexes	-	-	-	-	-	-	1	-	2	2	-	-
Male	-	-	-	-	-	-	-	-	1	1	-	-
Female	-	-	-	-	-	-	1	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	3	3	7	3	3	4	5	13	11	3	2	-
Male	1	2	3	1	1	-	2	6	2	-	-	-
Female	2	1	4	2	2	4	3	7	9	3	2	-
White, both sexes	-	1	4	1	2	3	1	10	9	2	1	-
Male	-	1	1	-	1	-	1	4	1	-	-	-
Female	-	-	3	1	1	3	-	6	8	2	1	-
Black, both sexes	3	2	3	2	-	1	4	3	2	1	1	-
Male	1	1	2	1	-	-	1	2	1	-	-	-
Female	2	1	1	1	-	1	3	1	1	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Washington												
All races, both sexes	-	5	1	4	6	11	14	28	25	10	1	-
Male	-	5	1	3	3	4	8	13	8	4	1	-
Female	-	-	-	1	3	7	6	15	17	6	-	-
White, both sexes	-	5	1	1	6	10	11	27	22	10	1	-
Male	-	5	1	1	3	3	7	13	8	4	1	-
Female	-	-	-	-	3	7	4	14	14	6	-	-
Black, both sexes	-	-	-	3	-	1	2	-	-	-	-	-
Male	-	-	-	2	-	1	1	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	1	-	-	-
West Virginia												
All races, both sexes	-	-	1	-	2	2	-	5	3	-	1	-
Male	-	-	1	-	-	1	-	2	1	-	-	-
Female	-	-	-	-	2	1	-	3	2	-	1	-
White, both sexes	-	-	1	-	2	2	-	5	3	-	1	-
Male	-	-	1	-	-	1	-	2	1	-	-	-
Female	-	-	-	-	2	1	-	3	2	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Wisconsin												
All races, both sexes	2	-	1	2	-	6	14	16	7	4	-	-
Male	2	-	1	2	-	2	6	10	2	-	-	-
Female	-	-	-	-	-	4	8	6	5	4	-	-
White, both sexes	2	-	1	2	-	5	14	15	7	4	-	-
Male	2	-	1	2	-	2	6	9	2	-	-	-
Female	-	-	-	-	-	3	8	6	5	4	-	-
Black, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
United States												
All races, both sexes	406,351	24	8	12	9	40	111	329	719	1,786	4,433	9,281
Male	216,050	11	3	6	6	27	80	246	563	1,339	3,393	7,051
Female	190,301	13	5	6	3	13	31	83	156	447	1,040	2,230
White, both sexes	354,481	19	4	11	7	25	73	224	519	1,352	3,455	7,377
Male	189,056	10	1	5	4	16	54	167	408	1,031	2,692	5,755
Female	165,425	9	3	6	3	9	19	57	111	321	763	1,622
Black, both sexes	42,679	3	2	1	1	15	35	87	167	358	791	1,653
Male	21,768	1	1	1	1	11	24	63	128	247	547	1,099
Female	20,911	2	1	-	-	4	11	24	39	111	244	554
American Indian 1/, both sexes	1,777	1	2	-	-	-	1	5	14	19	55	74
Male	1,048	-	1	-	-	-	1	3	10	11	38	53
Female	729	1	1	-	-	-	-	2	4	8	17	21
Asian or Pacific Islander 2/, both sexes	7,414	1	-	-	1	-	2	13	19	57	132	177
Male	4,178	-	-	-	1	-	1	13	17	50	116	144
Female	3,236	1	-	-	-	-	1	-	2	7	16	33
Alabama												
All races, both sexes	5,535	-	-	-	-	-	4	8	14	42	82	172
Male	2,964	-	-	-	-	-	2	5	12	26	56	129
Female	2,571	-	-	-	-	-	2	3	2	16	26	43
White, both sexes	4,442	-	-	-	-	-	3	5	11	34	54	136
Male	2,412	-	-	-	-	-	2	2	9	20	38	107
Female	2,030	-	-	-	-	-	1	3	2	14	16	29
Black, both sexes	1,077	-	-	-	-	-	1	3	3	8	26	36
Male	543	-	-	-	-	-	-	3	3	6	17	22
Female	534	-	-	-	-	-	1	-	-	2	9	14
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	2	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Alaska												
All races, both sexes	344	-	-	-	-	-	-	-	-	2	13	10
Male	226	-	-	-	-	-	-	-	-	-	13	9
Female	118	-	-	-	-	-	-	-	-	2	-	1
White, both sexes	272	-	-	-	-	-	-	-	-	1	12	7
Male	180	-	-	-	-	-	-	-	-	-	12	6
Female	92	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	53	-	-	-	-	-	-	-	-	-	-	2
Male	36	-	-	-	-	-	-	-	-	-	-	2
Female	17	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	1	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	1	-	-
Arizona												
All races, both sexes	7,292	-	-	-	1	1	5	1	13	30	61	144
Male	4,095	-	-	-	1	-	4	1	9	22	49	110
Female	3,197	-	-	-	-	1	1	-	4	8	12	34
White, both sexes	6,926	-	-	-	-	1	3	-	12	26	53	128
Male	3,908	-	-	-	-	-	2	-	9	19	44	100
Female	3,018	-	-	-	-	1	1	-	3	7	9	28
Black, both sexes	185	-	-	-	-	-	2	-	-	2	1	11
Male	85	-	-	-	-	-	2	-	-	1	1	5
Female	100	-	-	-	-	-	-	-	-	1	-	6
American Indian 1/, both sexes	138	-	-	-	-	-	-	1	1	-	7	3
Male	78	-	-	-	-	-	-	1	-	-	4	3
Female	60	-	-	-	-	-	-	-	1	-	3	-
Asian or Pacific Islander 2/, both sexes	43	-	-	-	1	-	-	-	-	2	-	2
Male	24	-	-	-	1	-	-	-	-	2	-	2
Female	19	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Arkansas												
All races, both sexes	4,863	-	2	-	-	1	3	3	19	36	87	173
Male	2,641	-	1	-	-	-	3	2	17	25	68	129
Female	2,222	-	1	-	-	1	-	1	2	11	19	44
White, both sexes	4,205	-	1	-	-	-	1	2	15	33	64	139
Male	2,302	-	-	-	-	-	1	2	15	24	51	107
Female	1,903	-	1	-	-	-	-	-	-	9	13	32
Black, both sexes	630	-	1	-	-	1	2	1	3	3	21	34
Male	323	-	1	-	-	-	2	-	2	1	16	22
Female	307	-	-	-	-	1	-	1	1	2	5	12
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	1	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	1	-	1	-
Asian or Pacific Islander 2/, both sexes	18	-	-	-	-	-	-	-	-	-	1	-
Male	11	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	43,215	-	1	-	-	4	10	25	59	135	326	736
Male	22,841	-	-	-	-	4	8	21	52	96	263	579
Female	20,374	-	1	-	-	-	2	4	7	39	63	157
White, both sexes	36,268	-	1	-	-	3	9	15	42	92	245	572
Male	19,156	-	-	-	-	3	7	13	38	62	198	451
Female	17,112	-	1	-	-	-	2	2	4	30	47	121
Black, both sexes	3,350	-	-	-	-	1	1	7	11	24	41	88
Male	1,723	-	-	-	-	1	1	5	8	16	27	62
Female	1,627	-	-	-	-	-	-	2	3	8	14	26
American Indian 1/, both sexes	203	-	-	-	-	-	-	-	1	2	3	10
Male	115	-	-	-	-	-	-	-	1	2	3	8
Female	88	-	-	-	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	3,394	-	-	-	-	-	-	3	5	17	37	66
Male	1,847	-	-	-	-	-	-	3	5	16	35	58
Female	1,547	-	-	-	-	-	-	-	-	1	2	8

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Colorado												
All races, both sexes	3,896	-	-	1	-	-	1	8	7	19	54	101
Male	2,223	-	-	1	-	-	1	3	6	15	42	84
Female	1,673	-	-	-	-	-	-	5	1	4	12	17
White, both sexes	3,738	-	-	1	-	-	1	5	6	19	51	96
Male	2,127	-	-	1	-	-	1	2	5	15	41	80
Female	1,611	-	-	-	-	-	-	3	1	4	10	16
Black, both sexes	117	-	-	-	-	-	-	2	-	-	1	4
Male	68	-	-	-	-	-	-	-	-	-	-	3
Female	49	-	-	-	-	-	-	2	-	-	1	1
American Indian 1/, both sexes	21	-	-	-	-	-	-	-	1	-	1	-
Male	16	-	-	-	-	-	-	-	1	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	20	-	-	-	-	-	-	1	-	-	1	1
Male	12	-	-	-	-	-	-	1	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	1	-
Connecticut												
All races, both sexes	4,336	2	-	-	-	-	1	2	4	17	49	78
Male	2,170	1	-	-	-	-	1	2	3	15	35	63
Female	2,166	1	-	-	-	-	-	-	1	2	14	15
White, both sexes	4,054	2	-	-	-	-	1	2	2	15	35	65
Male	2,005	1	-	-	-	-	1	2	1	13	24	56
Female	2,049	1	-	-	-	-	-	-	1	2	11	9
Black, both sexes	250	-	-	-	-	-	-	-	2	2	13	12
Male	143	-	-	-	-	-	-	-	2	2	10	6
Female	107	-	-	-	-	-	-	-	-	-	3	6
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	29	-	-	-	-	-	-	-	-	-	1	1
Male	20	-	-	-	-	-	-	-	-	-	1	1
Female	9	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Delaware												
All races, both sexes	1,259	-	-	-	-	-	-	1	2	7	19	42
Male	718	-	-	-	-	-	-	1	2	6	14	34
Female	541	-	-	-	-	-	-	-	-	1	5	8
White, both sexes	1,069	-	-	-	-	-	-	1	2	5	14	29
Male	611	-	-	-	-	-	-	1	2	4	11	24
Female	458	-	-	-	-	-	-	-	-	1	3	5
Black, both sexes	178	-	-	-	-	-	-	-	-	2	5	13
Male	98	-	-	-	-	-	-	-	-	2	3	10
Female	80	-	-	-	-	-	-	-	-	-	2	3
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1,018	-	-	-	-	-	-	-	3	9	11	22
Male	528	-	-	-	-	-	-	-	3	7	8	18
Female	490	-	-	-	-	-	-	-	-	2	3	4
White, both sexes	240	-	-	-	-	-	-	-	-	1	1	2
Male	146	-	-	-	-	-	-	-	-	1	-	2
Female	94	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	767	-	-	-	-	-	-	-	3	7	10	20
Male	377	-	-	-	-	-	-	-	3	5	8	16
Female	390	-	-	-	-	-	-	-	-	2	2	4
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Florida												
All races, both sexes	30,123	-	-	-	-	4	9	24	42	107	252	558
Male	16,256	-	-	-	-	3	7	20	35	83	190	430
Female	13,867	-	-	-	-	1	2	4	7	24	62	128
White, both sexes	27,298	-	-	-	-	3	5	15	28	77	195	466
Male	14,814	-	-	-	-	2	5	12	24	61	151	370
Female	12,484	-	-	-	-	1	-	3	4	16	44	96
Black, both sexes	2,637	-	-	-	-	1	4	7	13	29	49	90
Male	1,333	-	-	-	-	1	2	6	10	21	32	58
Female	1,304	-	-	-	-	-	2	1	3	8	17	32
American Indian 1/, both sexes	23	-	-	-	-	-	-	-	-	1	1	-
Male	16	-	-	-	-	-	-	-	-	1	-	-
Female	7	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	165	-	-	-	-	-	-	2	1	-	7	2
Male	93	-	-	-	-	-	-	2	1	-	7	2
Female	72	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	7,909	1	-	-	-	5	5	10	12	68	140	286
Male	4,456	1	-	-	-	4	4	8	10	50	102	206
Female	3,453	-	-	-	-	1	1	2	2	18	38	80
White, both sexes	6,007	1	-	-	-	3	2	6	8	43	92	202
Male	3,457	1	-	-	-	3	2	5	6	31	68	150
Female	2,550	-	-	-	-	-	-	1	2	12	24	52
Black, both sexes	1,839	-	-	-	-	2	3	4	4	23	48	81
Male	962	-	-	-	-	1	2	3	4	17	34	54
Female	877	-	-	-	-	1	1	1	-	6	14	27
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	58	-	-	-	-	-	-	-	-	2	-	3
Male	36	-	-	-	-	-	-	-	-	2	-	2
Female	22	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Hawaii												
All races, both sexes	1,185	-	-	-	-	-	-	3	2	8	21	26
Male	727	-	-	-	-	-	-	3	-	6	13	16
Female	458	-	-	-	-	-	-	-	2	2	8	10
White, both sexes	331	-	-	-	-	-	-	1	1	3	7	8
Male	219	-	-	-	-	-	-	1	-	3	4	7
Female	112	-	-	-	-	-	-	-	1	-	3	1
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	844	-	-	-	-	-	-	2	1	5	14	18
Male	501	-	-	-	-	-	-	2	-	3	9	9
Female	343	-	-	-	-	-	-	-	1	2	5	9
Idaho												
All races, both sexes	1,514	-	-	-	-	1	-	1	2	4	14	35
Male	901	-	-	-	-	1	-	-	2	3	8	26
Female	613	-	-	-	-	-	-	1	-	1	6	9
White, both sexes	1,490	-	-	-	-	1	-	1	1	2	12	35
Male	881	-	-	-	-	1	-	-	1	1	7	26
Female	609	-	-	-	-	-	-	1	-	1	5	9
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	2	1	-
Male	9	-	-	-	-	-	-	-	-	2	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	1	-	-	-
Male	10	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Illinois												
All races, both sexes	16,515	1	-	1	-	1	5	21	31	59	189	422
Male	8,685	-	-	1	-	1	5	15	19	52	153	304
Female	7,830	1	-	-	-	-	-	6	12	7	36	118
White, both sexes	14,039	1	-	1	-	1	4	18	24	42	136	303
Male	7,379	-	-	1	-	1	4	13	15	37	114	225
Female	6,660	1	-	-	-	-	-	5	9	5	22	78
Black, both sexes	2,277	-	-	-	-	-	-	3	7	15	50	115
Male	1,204	-	-	-	-	-	-	2	4	13	37	76
Female	1,073	-	-	-	-	-	-	1	3	2	13	39
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	188	-	-	-	-	-	1	-	-	2	3	3
Male	97	-	-	-	-	-	1	-	-	2	2	2
Female	91	-	-	-	-	-	-	-	-	-	1	1
Indiana												
All races, both sexes	8,453	2	-	1	-	-	2	7	25	60	129	250
Male	4,540	2	-	1	-	-	2	6	18	47	97	192
Female	3,913	-	-	-	-	-	-	1	7	13	32	58
White, both sexes	7,824	2	-	1	-	-	2	7	20	52	104	217
Male	4,187	2	-	1	-	-	2	6	13	42	80	168
Female	3,637	-	-	-	-	-	-	1	7	10	24	49
Black, both sexes	616	-	-	-	-	-	-	-	5	8	25	33
Male	345	-	-	-	-	-	-	-	5	5	17	24
Female	271	-	-	-	-	-	-	-	-	3	8	9
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Iowa												
All races, both sexes	5,194	1	-	1	-	-	-	3	5	11	53	98
Male	2,707	-	-	-	-	-	-	3	3	9	40	69
Female	2,487	1	-	1	-	-	-	-	2	2	13	29
White, both sexes	5,119	1	-	1	-	-	-	3	5	11	52	92
Male	2,662	-	-	-	-	-	-	3	3	9	39	65
Female	2,457	1	-	1	-	-	-	-	2	2	13	27
Black, both sexes	58	-	-	-	-	-	-	-	-	-	-	6
Male	36	-	-	-	-	-	-	-	-	-	-	4
Female	22	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	3,327	-	-	-	1	-	4	3	10	19	41	75
Male	1,804	-	-	-	-	-	3	3	6	18	28	62
Female	1,523	-	-	-	1	-	1	-	4	1	13	13
White, both sexes	3,142	-	-	-	1	-	4	2	10	13	36	71
Male	1,706	-	-	-	-	-	3	2	6	13	26	59
Female	1,436	-	-	-	1	-	1	-	4	-	10	12
Black, both sexes	144	-	-	-	-	-	-	-	-	4	1	3
Male	73	-	-	-	-	-	-	-	-	3	-	2
Female	71	-	-	-	-	-	-	-	-	1	1	1
American Indian 1/, both sexes	30	-	-	-	-	-	-	1	-	1	1	1
Male	20	-	-	-	-	-	-	1	-	1	-	1
Female	10	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	1	3	-
Male	5	-	-	-	-	-	-	-	-	1	2	-
Female	6	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Kentucky												
All races, both sexes	6,229	-	-	1	1	-	6	17	26	47	136	187
Male	3,436	-	-	-	-	-	5	12	19	36	95	151
Female	2,793	-	-	1	1	-	1	5	7	11	41	36
White, both sexes	5,830	-	-	1	1	-	3	14	23	43	123	170
Male	3,222	-	-	-	-	-	3	9	17	33	91	138
Female	2,608	-	-	1	1	-	-	5	6	10	32	32
Black, both sexes	387	-	-	-	-	-	3	3	3	4	13	16
Male	208	-	-	-	-	-	2	3	2	3	4	12
Female	179	-	-	-	-	-	1	-	1	1	9	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	5,606	1	-	-	-	3	5	12	24	40	112	221
Male	3,163	-	-	-	-	3	4	10	16	28	77	148
Female	2,443	1	-	-	-	-	1	2	8	12	35	73
White, both sexes	4,044	-	-	-	-	1	1	6	15	23	61	136
Male	2,284	-	-	-	-	1	-	6	11	18	40	95
Female	1,760	-	-	-	-	-	1	-	4	5	21	41
Black, both sexes	1,532	1	-	-	-	2	4	4	8	17	50	82
Male	862	-	-	-	-	2	4	2	4	10	36	51
Female	670	1	-	-	-	-	-	2	4	7	14	31
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	21	-	-	-	-	-	-	2	1	-	1	2
Male	14	-	-	-	-	-	-	2	1	-	1	2
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Maine												
All races, both sexes	1,765	-	-	-	-	-	1	3	1	7	27	38
Male	998	-	-	-	-	-	-	2	1	5	23	28
Female	767	-	-	-	-	-	1	1	-	2	4	10
White, both sexes	1,749	-	-	-	-	-	1	2	1	7	25	36
Male	987	-	-	-	-	-	-	2	1	5	21	27
Female	762	-	-	-	-	-	1	-	-	2	4	9
Black, both sexes	6	-	-	-	-	-	-	1	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	1	2
Male	4	-	-	-	-	-	-	-	-	-	1	1
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	7,833	-	-	1	-	-	2	7	15	34	73	164
Male	3,934	-	-	1	-	-	2	5	11	23	55	113
Female	3,899	-	-	-	-	-	-	2	4	11	18	51
White, both sexes	5,837	-	-	-	-	-	-	3	6	22	44	103
Male	2,943	-	-	-	-	-	-	3	4	14	32	77
Female	2,894	-	-	-	-	-	-	-	2	8	12	26
Black, both sexes	1,877	-	-	1	-	-	2	4	9	12	27	56
Male	921	-	-	1	-	-	2	2	7	9	21	33
Female	956	-	-	-	-	-	-	2	2	3	6	23
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	2
Male	6	-	-	-	-	-	-	-	-	-	-	2
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	110	-	-	-	-	-	-	-	-	-	2	3
Male	64	-	-	-	-	-	-	-	-	-	2	1
Female	46	-	-	-	-	-	-	-	-	-	-	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Massachusetts												
All races, both sexes	8,074	-	-	-	-	2	1	2	6	27	69	149
Male	4,163	-	-	-	-	1	-	1	4	22	54	120
Female	3,911	-	-	-	-	1	1	1	2	5	15	29
White, both sexes	7,736	-	-	-	-	2	1	2	6	25	57	137
Male	3,986	-	-	-	-	1	-	1	4	20	48	109
Female	3,750	-	-	-	-	1	1	1	2	5	9	28
Black, both sexes	253	-	-	-	-	-	-	-	-	1	9	10
Male	126	-	-	-	-	-	-	-	-	1	4	10
Female	127	-	-	-	-	-	-	-	-	-	5	-
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	76	-	-	-	-	-	-	-	-	1	3	2
Male	46	-	-	-	-	-	-	-	-	1	2	1
Female	30	-	-	-	-	-	-	-	-	-	1	1
Michigan												
All races, both sexes	16,619	1	-	-	-	1	5	7	36	74	155	383
Male	8,885	-	-	-	-	1	3	5	32	56	127	302
Female	7,734	1	-	-	-	-	2	2	4	18	28	81
White, both sexes	14,094	-	-	-	-	-	3	3	21	49	124	298
Male	7,531	-	-	-	-	-	2	2	18	41	103	236
Female	6,563	-	-	-	-	-	1	1	3	8	21	62
Black, both sexes	2,368	-	-	-	-	1	1	4	14	23	23	81
Male	1,269	-	-	-	-	1	1	3	13	14	18	62
Female	1,099	-	-	-	-	-	-	1	1	9	5	19
American Indian 1/, both sexes	85	1	-	-	-	-	-	-	-	2	6	1
Male	50	-	-	-	-	-	-	-	-	1	4	1
Female	35	1	-	-	-	-	-	-	-	1	2	-
Asian or Pacific Islander 2/, both sexes	72	-	-	-	-	-	1	-	1	-	2	3
Male	35	-	-	-	-	-	-	-	1	-	2	3
Female	37	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Minnesota												
All races, both sexes	4,298	-	-	-	-	1	-	3	8	16	42	104
Male	2,451	-	-	-	-	1	-	3	8	14	32	81
Female	1,847	-	-	-	-	-	-	-	-	2	10	23
White, both sexes	4,171	-	-	-	-	1	-	2	6	12	34	93
Male	2,373	-	-	-	-	1	-	2	6	11	26	75
Female	1,798	-	-	-	-	-	-	-	-	1	8	18
Black, both sexes	58	-	-	-	-	-	-	1	1	2	3	8
Male	36	-	-	-	-	-	-	1	1	2	3	4
Female	22	-	-	-	-	-	-	-	-	-	-	4
American Indian 1/, both sexes	46	-	-	-	-	-	-	-	1	1	2	3
Male	27	-	-	-	-	-	-	-	1	-	1	2
Female	19	-	-	-	-	-	-	-	-	1	1	1
Asian or Pacific Islander 2/, both sexes	23	-	-	-	-	-	-	-	-	1	3	-
Male	15	-	-	-	-	-	-	-	-	1	2	-
Female	8	-	-	-	-	-	-	-	-	-	1	-
Mississippi												
All races, both sexes	4,108	-	-	-	-	-	1	6	10	31	103	137
Male	2,201	-	-	-	-	-	1	4	7	21	74	96
Female	1,907	-	-	-	-	-	-	2	3	10	29	41
White, both sexes	2,874	-	-	-	-	-	-	1	5	18	59	74
Male	1,545	-	-	-	-	-	-	-	5	14	42	52
Female	1,329	-	-	-	-	-	-	1	-	4	17	22
Black, both sexes	1,219	-	-	-	-	-	1	5	5	13	44	60
Male	645	-	-	-	-	-	1	4	2	7	32	41
Female	574	-	-	-	-	-	-	1	3	6	12	19
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	2
Male	7	-	-	-	-	-	-	-	-	-	-	2
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Missouri												
All races, both sexes	9,737	-	-	-	-	-	2	8	22	38	115	259
Male	5,057	-	-	-	-	-	1	6	19	30	84	198
Female	4,680	-	-	-	-	-	1	2	3	8	31	61
White, both sexes	8,763	-	-	-	-	-	2	6	18	32	97	210
Male	4,576	-	-	-	-	-	1	4	17	24	72	167
Female	4,187	-	-	-	-	-	1	2	1	8	25	43
Black, both sexes	931	-	-	-	-	-	-	2	4	4	14	47
Male	459	-	-	-	-	-	-	2	2	4	9	29
Female	472	-	-	-	-	-	-	-	2	-	5	18
American Indian 1/, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	28	-	-	-	-	-	-	-	-	2	4	2
Male	17	-	-	-	-	-	-	-	-	2	3	2
Female	11	-	-	-	-	-	-	-	-	-	1	-
Montana												
All races, both sexes	1,059	-	-	-	-	-	-	1	-	3	14	29
Male	675	-	-	-	-	-	-	1	-	3	12	25
Female	384	-	-	-	-	-	-	-	-	-	2	4
White, both sexes	1,014	-	-	-	-	-	-	1	-	3	13	27
Male	646	-	-	-	-	-	-	1	-	3	11	24
Female	368	-	-	-	-	-	-	-	-	-	2	3
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	41	-	-	-	-	-	-	-	-	-	1	2
Male	25	-	-	-	-	-	-	-	-	-	1	1
Female	16	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Nebraska												
All races, both sexes	1,966	1	-	-	-	-	1	1	2	9	24	38
Male	1,065	-	-	-	-	-	1	1	1	5	22	29
Female	901	1	-	-	-	-	-	-	1	4	2	9
White, both sexes	1,885	1	-	-	-	-	1	1	2	6	22	35
Male	1,016	-	-	-	-	-	1	1	1	4	20	27
Female	869	1	-	-	-	-	-	-	1	2	2	8
Black, both sexes	65	-	-	-	-	-	-	-	-	2	2	3
Male	40	-	-	-	-	-	-	-	-	1	2	2
Female	25	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	2,426	-	-	-	1	-	1	6	2	10	32	58
Male	1,548	-	-	-	-	-	1	5	2	6	26	39
Female	878	-	-	-	1	-	-	1	-	4	6	19
White, both sexes	2,178	-	-	-	1	-	-	4	2	9	25	50
Male	1,397	-	-	-	-	-	-	4	2	5	20	38
Female	781	-	-	-	1	-	-	-	-	4	5	12
Black, both sexes	146	-	-	-	-	-	1	2	-	-	5	4
Male	84	-	-	-	-	-	1	1	-	-	4	1
Female	62	-	-	-	-	-	-	1	-	-	1	3
American Indian 1/, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	82	-	-	-	-	-	-	-	-	1	2	4
Male	51	-	-	-	-	-	-	-	-	1	2	-
Female	31	-	-	-	-	-	-	-	-	-	-	4

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
New Hampshire												
All races, both sexes	1,690	-	-	-	-	-	-	1	4	7	18	32
Male	892	-	-	-	-	-	-	1	1	3	17	23
Female	798	-	-	-	-	-	-	-	3	4	1	9
White, both sexes	1,681	-	-	-	-	-	-	1	4	7	18	31
Male	889	-	-	-	-	-	-	1	1	3	17	22
Female	792	-	-	-	-	-	-	-	3	4	1	9
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	12,944	2	-	-	-	-	1	4	13	26	92	240
Male	6,495	1	-	-	-	-	-	2	12	21	71	189
Female	6,449	1	-	-	-	-	1	2	1	5	21	51
White, both sexes	11,409	1	-	-	-	-	1	3	9	17	73	186
Male	5,702	1	-	-	-	-	-	1	8	13	58	152
Female	5,707	-	-	-	-	-	1	2	1	4	15	34
Black, both sexes	1,285	-	-	-	-	-	-	-	4	6	16	44
Male	638	-	-	-	-	-	-	-	4	5	10	28
Female	647	-	-	-	-	-	-	-	-	1	6	16
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	241	1	-	-	-	-	-	1	-	3	3	9
Male	153	-	-	-	-	-	-	1	-	3	3	8
Female	88	1	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
New Mexico												
All races, both sexes	2,218	-	1	-	-	-	-	3	4	7	19	53
Male	1,274	-	1	-	-	-	-	3	4	6	17	45
Female	944	-	-	-	-	-	-	-	-	1	2	8
White, both sexes	2,073	-	-	-	-	-	-	3	2	7	16	49
Male	1,186	-	-	-	-	-	-	3	2	6	14	42
Female	887	-	-	-	-	-	-	-	-	1	2	7
Black, both sexes	36	-	-	-	-	-	-	-	-	-	-	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	99	-	1	-	-	-	-	-	2	-	3	4
Male	64	-	1	-	-	-	-	-	2	-	3	3
Female	35	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	38,071	3	-	1	-	7	6	16	35	111	269	510
Male	17,924	2	-	1	-	3	4	11	30	81	201	382
Female	20,147	1	-	-	-	4	2	5	5	30	68	128
White, both sexes	31,972	3	-	1	-	3	4	7	19	84	186	352
Male	15,167	2	-	1	-	1	3	6	16	68	146	279
Female	16,805	1	-	-	-	2	1	1	3	16	40	73
Black, both sexes	5,104	-	-	-	-	4	2	9	14	22	63	137
Male	2,202	-	-	-	-	2	1	5	12	9	36	83
Female	2,902	-	-	-	-	2	1	4	2	13	27	54
American Indian 1/, both sexes	51	-	-	-	-	-	-	-	-	-	2	2
Male	29	-	-	-	-	-	-	-	-	-	2	2
Female	22	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	944	-	-	-	-	-	-	-	2	5	18	19
Male	526	-	-	-	-	-	-	-	2	4	17	18
Female	418	-	-	-	-	-	-	-	-	1	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
North Carolina												
All races, both sexes	11,233	-	-	-	-	-	1	7	34	80	176	327
Male	6,165	-	-	-	-	-	1	4	30	57	132	248
Female	5,068	-	-	-	-	-	-	3	4	23	44	79
White, both sexes	8,999	-	-	-	-	-	1	2	20	56	131	234
Male	5,008	-	-	-	-	-	1	-	18	42	104	187
Female	3,991	-	-	-	-	-	-	2	2	14	27	47
Black, both sexes	2,092	-	-	-	-	-	-	4	12	22	41	90
Male	1,064	-	-	-	-	-	-	3	10	13	25	60
Female	1,028	-	-	-	-	-	-	1	2	9	16	30
American Indian 1/, both sexes	107	-	-	-	-	-	-	1	1	-	4	2
Male	68	-	-	-	-	-	-	1	1	-	3	1
Female	39	-	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	35	-	-	-	-	-	-	-	1	2	-	1
Male	25	-	-	-	-	-	-	-	1	2	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
North Dakota												
All races, both sexes	1,040	-	-	-	-	-	1	2	1	3	10	23
Male	578	-	-	-	-	-	1	-	1	1	8	19
Female	462	-	-	-	-	-	-	2	-	2	2	4
White, both sexes	1,012	-	-	-	-	-	-	2	1	1	8	22
Male	562	-	-	-	-	-	-	-	1	1	6	18
Female	450	-	-	-	-	-	-	2	-	-	2	4
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	26	-	-	-	-	-	1	-	-	2	2	1
Male	14	-	-	-	-	-	1	-	-	-	2	1
Female	12	-	-	-	-	-	-	-	-	2	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Ohio												
All races, both sexes	18,530	1	2	-	-	1	5	11	42	90	179	447
Male	9,717	1	1	-	-	-	5	7	31	73	135	339
Female	8,813	-	1	-	-	1	-	4	11	17	44	108
White, both sexes	16,816	1	1	-	-	1	4	8	34	79	156	382
Male	8,822	1	1	-	-	-	4	5	25	64	121	300
Female	7,994	-	-	-	-	1	-	3	9	15	35	82
Black, both sexes	1,655	-	1	-	-	-	1	3	8	10	21	65
Male	854	-	-	-	-	-	1	2	6	8	13	39
Female	801	-	1	-	-	-	-	1	2	2	8	26
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	50	-	-	-	-	-	-	-	-	1	2	-
Male	36	-	-	-	-	-	-	-	-	1	1	-
Female	14	-	-	-	-	-	-	-	-	-	1	-
Oklahoma												
All races, both sexes	6,537	-	-	-	-	-	2	8	11	30	77	210
Male	3,448	-	-	-	-	-	-	6	10	20	56	150
Female	3,089	-	-	-	-	-	2	2	1	10	21	60
White, both sexes	5,792	-	-	-	-	-	2	5	8	24	67	166
Male	3,037	-	-	-	-	-	-	4	8	18	48	121
Female	2,755	-	-	-	-	-	2	1	-	6	19	45
Black, both sexes	370	-	-	-	-	-	-	2	-	3	4	29
Male	204	-	-	-	-	-	-	2	-	1	4	20
Female	166	-	-	-	-	-	-	-	-	2	-	9
American Indian 1/, both sexes	349	-	-	-	-	-	-	1	3	3	6	15
Male	199	-	-	-	-	-	-	-	2	1	4	9
Female	150	-	-	-	-	-	-	1	1	2	2	6
Asian or Pacific Islander 2/, both sexes	26	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Oregon												
All races, both sexes	3,959	-	-	-	-	-	-	-	11	14	46	80
Male	2,327	-	-	-	-	-	-	-	10	13	39	63
Female	1,632	-	-	-	-	-	-	-	1	1	7	17
White, both sexes	3,846	-	-	-	-	-	-	-	10	12	44	75
Male	2,257	-	-	-	-	-	-	-	9	11	37	61
Female	1,589	-	-	-	-	-	-	-	1	1	7	14
Black, both sexes	41	-	-	-	-	-	-	-	-	1	-	2
Male	29	-	-	-	-	-	-	-	-	1	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	28	-	-	-	-	-	-	-	-	-	1	2
Male	12	-	-	-	-	-	-	-	-	-	1	-
Female	16	-	-	-	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	44	-	-	-	-	-	-	-	1	1	1	1
Male	29	-	-	-	-	-	-	-	1	1	1	1
Female	15	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	21,297	1	1	1	-	2	6	10	23	61	160	409
Male	11,044	1	-	-	-	1	1	7	18	46	124	301
Female	10,253	-	1	1	-	1	5	3	5	15	36	108
White, both sexes	19,518	-	1	1	-	1	4	9	16	54	135	356
Male	10,157	-	-	-	-	-	1	6	12	41	103	268
Female	9,361	-	1	1	-	1	3	3	4	13	32	88
Black, both sexes	1,671	1	-	-	-	1	2	1	6	7	22	50
Male	824	1	-	-	-	1	-	1	5	5	18	31
Female	847	-	-	-	-	-	2	-	1	2	4	19
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	99	-	-	-	-	-	-	-	1	-	3	3
Male	55	-	-	-	-	-	-	-	1	-	3	2
Female	44	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Rhode Island												
All races, both sexes	2,210	-	-	-	-	-	-	-	5	11	22	24
Male	1,027	-	-	-	-	-	-	-	5	7	17	23
Female	1,183	-	-	-	-	-	-	-	-	4	5	1
White, both sexes	2,138	-	-	-	-	-	-	-	4	11	22	22
Male	989	-	-	-	-	-	-	-	4	7	17	21
Female	1,149	-	-	-	-	-	-	-	-	4	5	1
Black, both sexes	59	-	-	-	-	-	-	-	-	-	-	1
Male	30	-	-	-	-	-	-	-	-	-	-	1
Female	29	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	1	-	-	1
Male	4	-	-	-	-	-	-	-	1	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	5,238	-	-	1	-	-	2	4	8	34	89	161
Male	2,985	-	-	1	-	-	-	4	5	26	67	118
Female	2,253	-	-	-	-	-	2	-	3	8	22	43
White, both sexes	3,931	-	-	1	-	-	-	2	5	20	69	94
Male	2,292	-	-	1	-	-	-	2	4	15	51	71
Female	1,639	-	-	-	-	-	-	-	1	5	18	23
Black, both sexes	1,280	-	-	-	-	-	2	2	3	13	20	63
Male	672	-	-	-	-	-	-	2	1	10	16	43
Female	608	-	-	-	-	-	2	-	2	3	4	20
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	18	-	-	-	-	-	-	-	-	1	-	3
Male	14	-	-	-	-	-	-	-	-	1	-	3
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
South Dakota												
All races, both sexes	1,234	-	-	-	-	-	-	2	1	3	16	20
Male	695	-	-	-	-	-	-	1	-	3	14	16
Female	539	-	-	-	-	-	-	1	1	-	2	4
White, both sexes	1,165	-	-	-	-	-	-	2	1	1	9	13
Male	656	-	-	-	-	-	-	1	-	1	9	10
Female	509	-	-	-	-	-	-	1	1	-	-	3
Black, both sexes	4	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	1	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	63	-	-	-	-	-	-	-	-	2	6	6
Male	36	-	-	-	-	-	-	-	-	2	4	5
Female	27	-	-	-	-	-	-	-	-	-	2	1
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	10,312	-	-	-	-	-	-	9	25	61	153	327
Male	5,461	-	-	-	-	-	-	6	20	44	121	246
Female	4,851	-	-	-	-	-	-	3	5	17	32	81
White, both sexes	9,010	-	-	-	-	-	-	7	18	50	128	277
Male	4,783	-	-	-	-	-	-	5	13	34	100	210
Female	4,227	-	-	-	-	-	-	2	5	16	28	67
Black, both sexes	1,271	-	-	-	-	-	-	2	7	10	23	49
Male	657	-	-	-	-	-	-	1	7	9	19	35
Female	614	-	-	-	-	-	-	1	-	1	4	14
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	27	-	-	-	-	-	-	-	-	1	2	1
Male	18	-	-	-	-	-	-	-	-	1	2	1
Female	9	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Texas												
All races, both sexes	25,337	6	-	3	3	3	6	30	45	139	339	698
Male	13,954	2	-	-	3	2	5	22	27	100	257	532
Female	11,383	4	-	3	-	1	1	8	18	39	82	166
White, both sexes	22,306	5	-	3	2	3	4	23	34	105	269	575
Male	12,334	2	-	-	2	2	3	17	22	76	204	445
Female	9,972	3	-	3	-	1	1	6	12	29	65	130
Black, both sexes	2,771	1	-	-	1	-	2	6	10	30	59	113
Male	1,468	-	-	-	1	-	2	5	4	22	42	78
Female	1,303	1	-	-	-	-	-	1	6	8	17	35
American Indian 1/, both sexes	37	-	-	-	-	-	-	1	-	1	-	1
Male	24	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	223	-	-	-	-	-	-	-	1	3	11	9
Male	128	-	-	-	-	-	-	-	1	2	11	8
Female	95	-	-	-	-	-	-	-	-	1	-	1
Utah												
All races, both sexes	1,459	-	-	-	-	-	-	3	4	13	17	42
Male	875	-	-	-	-	-	-	2	4	11	12	35
Female	584	-	-	-	-	-	-	1	-	2	5	7
White, both sexes	1,415	-	-	-	-	-	-	3	4	12	15	36
Male	849	-	-	-	-	-	-	2	4	10	12	31
Female	566	-	-	-	-	-	-	1	-	2	3	5
Black, both sexes	9	-	-	-	-	-	-	-	-	-	2	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	17	-	-	-	-	-	-	-	-	-	-	3
Male	9	-	-	-	-	-	-	-	-	-	-	2
Female	8	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	18	-	-	-	-	-	-	-	-	1	-	3
Male	12	-	-	-	-	-	-	-	-	1	-	2
Female	6	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Vermont												
All races, both sexes	824	-	-	-	-	-	-	-	1	1	6	12
Male	430	-	-	-	-	-	-	-	1	-	5	10
Female	394	-	-	-	-	-	-	-	-	1	1	2
White, both sexes	820	-	-	-	-	-	-	-	1	1	6	12
Male	429	-	-	-	-	-	-	-	1	-	5	10
Female	391	-	-	-	-	-	-	-	-	1	1	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	8,150	-	-	-	-	1	3	15	23	59	105	230
Male	4,565	-	-	-	-	1	2	12	18	45	84	179
Female	3,585	-	-	-	-	-	1	3	5	14	21	51
White, both sexes	6,468	-	-	-	-	-	2	11	17	34	79	157
Male	3,652	-	-	-	-	-	2	8	13	28	64	126
Female	2,816	-	-	-	-	-	-	3	4	6	15	31
Black, both sexes	1,558	-	-	-	-	1	1	4	6	25	22	65
Male	838	-	-	-	-	1	-	4	5	17	17	45
Female	720	-	-	-	-	-	1	-	1	8	5	20
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	119	-	-	-	-	-	-	-	-	-	4	8
Male	73	-	-	-	-	-	-	-	-	-	3	8
Female	46	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Washington												
All races, both sexes	7,425	1	1	-	1	2	2	3	7	25	78	149
Male	4,109	-	-	-	1	1	1	3	4	19	60	105
Female	3,316	1	1	-	-	1	1	-	3	6	18	44
White, both sexes	6,930	1	-	-	1	1	2	1	4	21	68	132
Male	3,822	-	-	-	1	-	1	1	2	16	53	94
Female	3,108	1	-	-	-	1	1	-	2	5	15	38
Black, both sexes	173	-	-	-	-	1	-	-	1	1	7	12
Male	115	-	-	-	-	1	-	-	1	1	5	9
Female	58	-	-	-	-	-	-	-	-	-	2	3
American Indian 1/, both sexes	94	-	1	-	-	-	-	-	-	-	2	4
Male	52	-	-	-	-	-	-	-	-	-	1	1
Female	42	-	1	-	-	-	-	-	-	-	1	3
Asian or Pacific Islander 2/, both sexes	228	-	-	-	-	-	-	2	2	3	1	1
Male	120	-	-	-	-	-	-	2	1	2	1	1
Female	108	-	-	-	-	-	-	-	1	1	-	-
West Virginia												
All races, both sexes	3,497	-	-	-	-	-	1	3	9	19	53	105
Male	1,848	-	-	-	-	-	1	3	7	15	42	76
Female	1,649	-	-	-	-	-	-	-	2	4	11	29
White, both sexes	3,401	-	-	-	-	-	1	3	9	17	51	102
Male	1,795	-	-	-	-	-	1	3	7	13	41	73
Female	1,606	-	-	-	-	-	-	-	2	4	10	29
Black, both sexes	89	-	-	-	-	-	-	-	-	2	2	3
Male	50	-	-	-	-	-	-	-	-	2	1	3
Female	39	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Wisconsin												
All races, both sexes	6,856	-	-	-	1	-	-	4	10	21	57	165
Male	3,821	-	-	-	1	-	-	3	7	18	49	142
Female	3,035	-	-	-	-	-	-	1	3	3	8	23
White, both sexes	6,561	-	-	-	1	-	-	3	7	19	50	146
Male	3,661	-	-	-	1	-	-	2	5	16	44	127
Female	2,900	-	-	-	-	-	-	1	2	3	6	19
Black, both sexes	218	-	-	-	-	-	-	1	1	1	5	14
Male	112	-	-	-	-	-	-	1	1	1	3	11
Female	106	-	-	-	-	-	-	-	-	-	2	3
American Indian 1/, both sexes	50	-	-	-	-	-	-	-	1	1	1	1
Male	33	-	-	-	-	-	-	-	-	1	1	1
Female	17	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	27	-	-	-	-	-	-	-	1	-	1	4
Male	15	-	-	-	-	-	-	-	1	-	1	3
Female	12	-	-	-	-	-	-	-	-	-	-	1
Wyoming												
All races, both sexes	592	-	-	-	-	-	1	1	1	2	9	22
Male	358	-	-	-	-	-	1	1	1	1	5	20
Female	234	-	-	-	-	-	-	-	-	1	4	2
White, both sexes	579	-	-	-	-	-	1	1	-	2	8	21
Male	349	-	-	-	-	-	1	1	-	1	4	19
Female	230	-	-	-	-	-	-	-	-	1	4	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	1	-	1	-
Male	5	-	-	-	-	-	-	-	1	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
United States												
All races, both sexes	15,109	20,733	25,431	28,094	34,933	49,453	66,699	71,378	51,375	21,614	4,759	21
Male	11,475	15,446	18,030	18,881	21,896	28,400	34,293	31,157	17,665	5,263	804	15
Female	3,634	5,287	7,401	9,213	13,037	21,053	32,406	40,221	33,710	16,351	3,955	6
White, both sexes	11,889	16,539	20,929	23,208	29,366	43,130	59,766	65,173	47,404	19,787	4,210	14
Male	9,246	12,595	15,128	15,933	18,756	25,146	31,250	28,802	16,489	4,857	701	10
Female	2,643	3,944	5,801	7,275	10,610	17,984	28,516	36,371	30,915	14,930	3,509	4
Black, both sexes	2,816	3,636	3,868	4,082	4,629	5,135	5,516	4,874	3,076	1,474	454	6
Male	1,895	2,418	2,466	2,420	2,561	2,590	2,352	1,750	834	278	77	4
Female	921	1,218	1,402	1,662	2,068	2,545	3,164	3,124	2,242	1,196	377	2
American Indian 1/, both sexes	127	166	164	198	196	224	221	167	88	46	9	-
Male	100	124	108	131	116	128	110	79	25	8	2	-
Female	27	42	56	67	80	96	111	88	63	38	7	-
Asian or Pacific Islander 2/, both sexes	277	392	470	606	742	964	1,196	1,164	807	307	86	1
Male	234	309	328	397	463	536	581	526	317	120	24	1
Female	43	83	142	209	279	428	615	638	490	187	62	-
Alabama												
All races, both sexes	244	387	431	487	568	715	849	821	507	167	37	-
Male	186	285	297	298	355	397	405	311	155	39	6	-
Female	58	102	134	189	213	318	444	510	352	128	31	-
White, both sexes	173	285	320	393	450	579	712	705	421	134	27	-
Male	130	213	222	242	296	323	357	282	131	34	4	-
Female	43	72	98	151	154	256	355	423	290	100	23	-
Black, both sexes	71	102	107	91	117	136	133	114	86	33	10	-
Male	56	72	72	55	58	74	46	28	24	5	2	-
Female	15	30	35	36	59	62	87	86	62	28	8	-
American Indian 1/, both sexes	-	-	1	-	-	-	2	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	3	3	1	-	2	2	-	-	-	-
Male	-	-	2	1	1	-	2	1	-	-	-	-
Female	-	-	1	2	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Alaska												
All races, both sexes	22	35	36	34	36	47	41	42	17	9	-	-
Male	21	27	32	19	29	32	24	13	5	2	-	-
Female	1	8	4	15	7	15	17	29	12	7	-	-
White, both sexes	18	30	34	25	22	32	37	30	16	8	-	-
Male	17	23	30	15	16	24	21	9	5	2	-	-
Female	1	7	4	10	6	8	16	21	11	6	-	-
Black, both sexes	2	-	-	-	2	3	-	1	-	-	-	-
Male	2	-	-	-	2	-	-	-	-	-	-	-
Female	-	-	-	-	-	3	-	1	-	-	-	-
American Indian 1/, both sexes	2	4	1	7	11	12	4	9	-	1	-	-
Male	2	3	1	3	10	8	3	4	-	-	-	-
Female	-	1	-	4	1	4	1	5	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	2	1	-	-	2	1	-	-	-
Male	-	1	1	1	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	2	1	-	-	-
Arizona												
All races, both sexes	242	356	518	533	670	967	1,242	1,286	839	329	52	2
Male	190	261	369	367	443	584	665	602	320	88	8	2
Female	52	95	149	166	227	383	577	684	519	241	44	-
White, both sexes	223	318	491	499	635	924	1,192	1,248	810	312	49	2
Male	176	240	352	351	423	559	642	585	312	84	8	2
Female	47	78	139	148	212	365	550	663	498	228	41	-
Black, both sexes	9	19	16	17	18	22	20	22	16	8	2	-
Male	5	8	12	8	8	9	10	10	4	2	-	-
Female	4	11	4	9	10	13	10	12	12	6	2	-
American Indian 1/, both sexes	9	15	8	13	13	15	26	10	9	8	-	-
Male	8	10	3	6	10	12	12	5	3	1	-	-
Female	1	5	5	7	3	3	14	5	6	7	-	-
Asian or Pacific Islander 2/, both sexes	1	4	3	4	4	6	4	6	4	1	1	-
Male	1	3	2	2	2	4	1	2	1	1	-	-
Female	-	1	1	2	2	2	3	4	3	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Arkansas												
All races, both sexes	226	305	339	400	458	587	743	726	510	194	51	-
Male	164	217	235	263	295	342	361	292	174	44	9	-
Female	62	88	104	137	163	245	382	434	336	150	42	-
White, both sexes	167	248	299	347	399	518	661	651	457	165	38	-
Male	125	176	211	228	261	310	324	265	156	40	6	-
Female	42	72	88	119	138	208	337	386	301	125	32	-
Black, both sexes	58	55	38	53	54	63	81	74	49	27	12	-
Male	38	39	24	35	32	28	36	26	15	3	3	-
Female	20	16	14	18	22	35	45	48	34	24	9	-
American Indian 1/, both sexes	1	1	1	-	2	2	1	-	-	-	-	-
Male	1	1	-	-	1	1	1	-	-	-	-	-
Female	-	-	1	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	3	4	-	1	4	2	1	-
Male	-	1	-	-	1	3	-	1	3	1	-	-
Female	-	-	1	-	2	1	-	-	1	1	1	-
California												
All races, both sexes	1,464	2,024	2,609	2,811	3,561	5,034	7,199	8,132	5,858	2,609	617	1
Male	1,171	1,533	1,845	1,860	2,230	2,873	3,574	3,757	2,136	727	111	1
Female	293	491	764	951	1,331	2,161	3,625	4,375	3,722	1,882	506	-
White, both sexes	1,124	1,564	2,094	2,218	2,836	4,195	6,201	7,062	5,189	2,282	524	-
Male	902	1,213	1,502	1,514	1,789	2,415	3,119	3,318	1,904	615	93	-
Female	222	351	592	704	1,047	1,780	3,082	3,744	3,285	1,667	431	-
Black, both sexes	211	300	307	336	374	389	396	427	241	150	46	-
Male	151	191	201	188	228	204	169	161	66	37	7	-
Female	60	109	106	148	146	185	227	266	175	113	39	-
American Indian 1/, both sexes	15	11	20	22	28	26	22	24	12	6	1	-
Male	14	11	10	15	16	11	12	8	4	-	-	-
Female	1	-	10	7	12	15	10	16	8	6	1	-
Asian or Pacific Islander 2/, both sexes	114	149	188	235	323	424	580	619	416	171	46	1
Male	104	118	132	143	197	243	274	270	162	75	11	1
Female	10	31	56	92	126	181	306	349	254	96	35	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Colorado												
All races, both sexes	182	224	259	285	331	497	610	682	416	180	39	-
Male	148	172	186	191	223	290	313	318	165	53	12	-
Female	34	52	73	94	108	207	297	364	251	127	27	-
White, both sexes	174	206	246	271	314	472	592	661	407	177	39	-
Male	142	157	176	181	213	278	305	307	159	52	12	-
Female	32	49	70	90	101	194	287	354	248	125	27	-
Black, both sexes	5	15	8	12	14	20	11	15	8	2	-	-
Male	4	12	7	8	8	8	4	8	5	1	-	-
Female	1	3	1	4	6	12	7	7	3	1	-	-
American Indian 1/, both sexes	2	2	5	2	1	3	2	1	-	1	-	-
Male	1	2	3	2	1	3	1	1	-	-	-	-
Female	1	-	2	-	-	-	1	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	2	2	5	5	1	-	-	-
Male	1	1	-	-	1	1	3	2	1	-	-	-
Female	-	-	-	-	1	1	2	3	-	-	-	-
Connecticut												
All races, both sexes	111	159	209	205	279	516	720	929	650	327	77	1
Male	87	123	145	135	177	295	379	403	223	73	9	1
Female	24	36	64	70	102	221	341	526	427	254	68	-
White, both sexes	91	143	178	176	254	481	679	900	634	322	73	1
Male	72	109	124	116	158	274	359	395	218	73	8	1
Female	19	34	54	60	96	207	320	505	416	249	65	-
Black, both sexes	15	14	27	26	24	29	37	28	13	4	4	-
Male	11	12	17	18	18	17	19	7	3	-	1	-
Female	4	2	10	8	6	12	18	21	10	4	3	-
American Indian 1/, both sexes	1	1	-	-	-	-	1	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	1	4	3	1	6	3	1	3	1	-	-
Male	3	1	4	1	1	4	1	1	2	-	-	-
Female	1	-	-	2	-	2	2	-	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Delaware												
All races, both sexes	54	67	88	71	115	148	214	193	152	71	14	1
Male	44	51	71	51	81	91	120	85	54	10	3	-
Female	10	16	17	20	34	57	94	108	98	61	11	1
White, both sexes	41	50	71	55	92	125	196	178	141	57	11	1
Male	33	41	56	40	67	77	115	78	52	8	2	-
Female	8	9	15	15	25	48	81	100	89	49	9	1
Black, both sexes	12	17	17	15	20	21	17	13	10	13	3	-
Male	10	10	15	10	12	13	5	5	1	1	1	-
Female	2	7	2	5	8	8	12	8	9	12	2	-
American Indian 1/, both sexes	1	-	-	1	2	1	1	-	-	1	-	-
Male	1	-	-	1	1	1	-	-	-	1	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	2	1	-	-	-
Male	-	-	-	-	1	-	-	2	1	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	31	53	89	107	98	128	143	140	109	57	18	-
Male	21	34	61	67	56	73	63	58	38	17	4	-
Female	10	19	28	40	42	55	80	82	71	40	14	-
White, both sexes	3	12	11	16	17	32	36	39	38	23	9	-
Male	3	9	10	12	12	23	26	22	16	9	1	-
Female	-	3	1	4	5	9	10	17	22	14	8	-
Black, both sexes	28	41	78	90	80	93	105	101	70	32	9	-
Male	18	25	51	54	43	50	36	36	22	7	3	-
Female	10	16	27	36	37	43	69	65	48	25	6	-
American Indian 1/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	3	1	-	1	2	-	-
Male	-	-	-	-	1	-	1	-	-	1	-	-
Female	-	-	-	-	-	3	-	-	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Florida												
All races, both sexes	921	1,193	1,606	1,857	2,399	3,818	5,184	5,836	4,187	1,751	372	3
Male	694	882	1,127	1,268	1,514	2,281	2,810	2,701	1,647	487	75	2
Female	227	311	479	589	885	1,537	2,374	3,135	2,540	1,264	297	1
White, both sexes	721	1,009	1,357	1,588	2,061	3,436	4,803	5,527	3,993	1,665	347	2
Male	571	759	972	1,107	1,330	2,092	2,636	2,584	1,592	473	72	1
Female	150	250	385	481	731	1,344	2,167	2,943	2,401	1,192	275	1
Black, both sexes	187	174	236	247	314	348	359	289	183	81	25	1
Male	112	114	146	146	170	172	165	109	52	13	3	1
Female	75	60	90	101	144	176	194	180	131	68	22	-
American Indian 1/, both sexes	1	3	2	3	2	3	-	5	2	-	-	-
Male	1	2	1	3	1	2	-	4	1	-	-	-
Female	-	1	1	-	1	1	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	12	7	11	19	22	31	22	15	9	5	-	-
Male	10	7	8	12	13	15	9	4	2	1	-	-
Female	2	-	3	7	9	16	13	11	7	4	-	-
Georgia												
All races, both sexes	400	565	620	710	806	940	1,117	1,107	778	294	45	-
Male	303	426	439	479	526	531	584	451	264	65	3	-
Female	97	139	181	231	280	409	533	656	514	229	42	-
White, both sexes	258	380	424	523	603	738	907	921	636	230	30	-
Male	196	296	315	369	414	431	496	388	226	57	3	-
Female	62	84	109	154	189	307	411	533	410	173	27	-
Black, both sexes	136	181	189	182	196	193	198	183	138	64	14	-
Male	103	126	121	106	107	96	83	62	35	8	-	-
Female	33	55	68	76	89	97	115	121	103	56	14	-
American Indian 1/, both sexes	1	-	2	-	-	1	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	4	5	5	7	8	11	3	4	-	1	-
Male	4	4	2	4	5	4	5	1	3	-	-	-
Female	1	-	3	1	2	4	6	2	1	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Hawaii												
All races, both sexes	41	70	88	110	108	140	179	200	135	43	11	-
Male	28	59	66	76	74	89	122	95	56	18	6	-
Female	13	11	22	34	34	51	57	105	79	25	5	-
White, both sexes	12	18	33	34	29	41	40	61	32	6	5	-
Male	8	16	29	27	23	28	26	26	16	2	3	-
Female	4	2	4	7	6	13	14	35	16	4	2	-
Black, both sexes	2	1	1	-	-	2	-	2	-	-	-	-
Male	2	1	-	-	-	1	-	1	-	-	-	-
Female	-	-	1	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	1	-	-
Male	-	-	1	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	27	51	53	76	79	97	139	137	103	36	6	-
Male	18	42	36	49	51	60	96	68	40	15	3	-
Female	9	9	17	27	28	37	43	69	63	21	3	-
Idaho												
All races, both sexes	64	79	102	123	142	180	218	286	186	67	10	-
Male	53	62	84	84	100	102	133	154	66	22	1	-
Female	11	17	18	39	42	78	85	132	120	45	9	-
White, both sexes	62	79	100	120	140	180	214	281	186	67	9	-
Male	51	62	83	81	99	102	129	150	66	22	-	-
Female	11	17	17	39	41	78	85	131	120	45	9	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	1	-	1	1	1	-	1	1	-	-	1	-
Male	1	-	1	1	1	-	1	1	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	2	1	-	3	3	-	-	-	-
Male	1	-	-	2	-	-	3	3	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Illinois												
All races, both sexes	659	913	1,011	1,104	1,407	1,906	2,734	2,828	2,151	866	204	2
Male	493	655	693	799	880	1,053	1,395	1,223	704	203	37	-
Female	166	258	318	305	527	853	1,339	1,605	1,447	663	167	2
White, both sexes	486	673	775	849	1,122	1,619	2,421	2,564	2,003	808	188	1
Male	384	503	554	631	716	913	1,257	1,125	662	192	32	-
Female	102	170	221	218	406	706	1,164	1,439	1,341	616	156	1
Black, both sexes	167	230	220	238	266	264	280	231	123	54	13	1
Male	104	144	132	154	154	127	123	85	34	10	5	-
Female	63	86	88	84	112	137	157	146	89	44	8	1
American Indian 1/, both sexes	-	2	1	-	3	-	1	3	-	-	-	-
Male	-	-	1	-	2	-	-	1	-	-	-	-
Female	-	2	-	-	1	-	1	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	8	15	17	16	23	32	30	25	4	3	-
Male	5	8	6	14	8	13	15	12	8	1	-	-
Female	1	-	9	3	8	10	17	18	17	3	3	-
Indiana												
All races, both sexes	357	487	521	628	767	1,060	1,393	1,332	946	395	91	-
Male	272	354	373	404	497	594	702	574	314	83	8	-
Female	85	133	148	224	270	466	691	758	632	312	83	-
White, both sexes	305	435	464	583	710	980	1,304	1,266	906	377	89	-
Male	233	315	339	377	465	550	655	546	306	79	8	-
Female	72	120	125	206	245	430	649	720	600	298	81	-
Black, both sexes	49	52	56	44	54	78	87	65	40	18	2	-
Male	36	39	33	27	30	42	47	28	8	4	-	-
Female	13	13	23	17	24	36	40	37	32	14	2	-
American Indian 1/, both sexes	1	-	-	1	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	1	-	3	2	2	-	-	-	-	-
Male	2	-	1	-	2	2	-	-	-	-	-	-
Female	-	-	-	-	1	-	2	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Iowa												
All races, both sexes	168	213	244	292	407	607	844	925	846	382	94	-
Male	136	167	177	196	266	349	470	415	306	87	14	-
Female	32	46	67	96	141	258	374	510	540	295	80	-
White, both sexes	164	204	239	283	395	601	836	920	839	379	94	-
Male	133	164	175	191	257	344	465	412	301	87	14	-
Female	31	40	64	92	138	257	371	508	538	292	80	-
Black, both sexes	4	9	5	5	10	3	5	4	5	2	-	-
Male	3	3	2	2	8	3	4	3	4	-	-	-
Female	1	6	3	3	2	-	1	1	1	2	-	-
American Indian 1/, both sexes	-	-	-	2	1	1	1	1	1	1	-	-
Male	-	-	-	2	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	1	1	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	1	2	2	-	1	-	-	-
Male	-	-	-	1	1	1	1	-	1	-	-	-
Female	-	-	-	1	-	1	1	-	-	-	-	-
Kansas												
All races, both sexes	127	138	172	229	275	400	555	584	443	195	56	-
Male	95	101	131	161	171	239	313	263	155	48	7	-
Female	32	37	41	68	104	161	242	321	288	147	49	-
White, both sexes	117	124	164	208	258	369	531	563	424	191	56	-
Male	89	91	126	145	166	223	303	253	146	48	7	-
Female	28	33	38	63	92	146	228	310	278	143	49	-
Black, both sexes	10	12	5	12	14	28	17	18	16	4	-	-
Male	6	8	3	10	4	14	7	8	8	-	-	-
Female	4	4	2	2	10	14	10	10	8	4	-	-
American Indian 1/, both sexes	-	2	3	7	1	3	5	2	3	-	-	-
Male	-	2	2	5	1	2	2	2	1	-	-	-
Female	-	-	1	2	-	1	3	-	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	2	-	2	1	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	1	2	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Kentucky												
All races, both sexes	319	397	456	538	625	754	940	886	607	246	40	-
Male	240	301	319	358	380	422	477	356	193	63	9	-
Female	79	96	137	180	245	332	463	530	414	183	31	-
White, both sexes	281	354	426	497	584	715	893	848	574	241	39	-
Male	220	272	297	331	360	402	456	339	182	63	9	-
Female	61	82	129	166	224	313	437	509	392	178	30	-
Black, both sexes	36	43	28	41	39	38	43	38	33	5	1	-
Male	20	29	21	27	19	19	19	17	11	-	-	-
Female	16	14	7	14	20	19	24	21	22	5	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	2	-	2	1	4	-	-	-	-	-
Male	-	-	1	-	1	1	2	-	-	-	-	-
Female	2	-	1	-	1	-	2	-	-	-	-	-
Louisiana												
All races, both sexes	314	452	477	488	551	633	854	748	462	171	38	-
Male	231	355	328	313	336	361	431	310	159	40	13	-
Female	83	97	149	175	215	272	423	438	303	131	25	-
White, both sexes	189	284	307	340	393	485	670	594	379	138	22	-
Male	150	226	216	212	248	291	352	246	133	33	6	-
Female	39	58	91	128	145	194	318	348	246	105	16	-
Black, both sexes	124	164	168	146	154	144	181	153	82	33	15	-
Male	80	126	110	99	86	69	79	64	26	7	7	-
Female	44	38	58	47	68	75	102	89	56	26	8	-
American Indian 1/, both sexes	-	1	1	1	1	2	1	-	1	-	-	-
Male	-	-	1	1	-	1	-	-	-	-	-	-
Female	-	1	-	-	1	1	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	3	1	1	3	2	2	1	-	-	1	-
Male	1	3	1	1	2	-	-	-	-	-	-	-
Female	-	-	-	-	1	2	2	1	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Maine												
All races, both sexes	61	80	116	114	157	225	271	306	238	97	23	-
Male	53	60	91	87	108	144	137	136	93	25	5	-
Female	8	20	25	27	49	81	134	170	145	72	18	-
White, both sexes	60	79	116	113	156	222	269	306	237	96	23	-
Male	52	59	91	87	107	141	137	136	92	24	5	-
Female	8	20	25	26	49	81	132	170	145	72	18	-
Black, both sexes	1	1	-	-	-	1	-	-	-	1	-	-
Male	1	1	-	-	-	1	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	1	1	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	1	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Maryland												
All races, both sexes	315	396	499	557	724	911	1,271	1,382	962	429	91	-
Male	221	275	353	355	429	503	606	571	301	96	14	-
Female	94	121	146	202	295	408	665	811	661	333	77	-
White, both sexes	202	226	317	362	486	693	1,009	1,119	811	362	72	-
Male	149	163	231	246	302	397	495	464	268	86	12	-
Female	53	63	86	116	184	296	514	655	543	276	60	-
Black, both sexes	112	164	170	185	222	201	245	243	143	63	18	-
Male	71	107	113	102	116	100	103	94	29	9	2	-
Female	41	57	57	83	106	101	142	149	114	54	16	-
American Indian 1/, both sexes	-	-	-	-	1	2	3	1	-	-	-	-
Male	-	-	-	-	1	-	2	1	-	-	-	-
Female	-	-	-	-	-	2	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	6	12	10	15	15	14	19	8	4	1	-
Male	1	5	9	7	10	6	6	12	4	1	-	-
Female	-	1	3	3	5	9	8	7	4	3	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Massachusetts												
All races, both sexes	249	332	422	469	645	952	1,375	1,609	1,151	506	107	1
Male	201	257	304	316	413	548	702	720	368	117	14	1
Female	48	75	118	153	232	404	673	889	783	389	93	-
White, both sexes	226	313	393	432	601	906	1,336	1,569	1,127	495	107	1
Male	186	247	289	293	384	524	683	706	360	116	14	1
Female	40	66	104	139	217	382	653	863	767	379	93	-
Black, both sexes	13	13	24	25	34	36	26	33	20	9	-	-
Male	8	7	11	14	19	20	13	12	6	1	-	-
Female	5	6	13	11	15	16	13	21	14	8	-	-
American Indian 1/, both sexes	4	1	2	-	-	1	-	-	-	1	-	-
Male	2	1	2	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	1	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	6	5	3	12	10	9	13	7	4	1	-	-
Male	5	2	2	9	10	4	6	2	2	-	-	-
Female	1	3	1	3	-	5	7	5	2	1	-	-
Michigan												
All races, both sexes	699	897	1,118	1,154	1,378	2,057	2,699	2,925	2,046	804	180	-
Male	530	650	782	779	844	1,153	1,392	1,296	709	192	32	-
Female	169	247	336	375	534	904	1,307	1,629	1,337	612	148	-
White, both sexes	512	651	880	906	1,131	1,737	2,356	2,650	1,866	748	159	-
Male	411	482	626	629	715	985	1,249	1,175	648	181	28	-
Female	101	169	254	277	416	752	1,107	1,475	1,218	567	131	-
Black, both sexes	174	234	228	235	232	300	319	260	168	52	19	-
Male	109	160	148	140	118	160	135	115	58	11	3	-
Female	65	74	80	95	114	140	184	145	110	41	16	-
American Indian 1/, both sexes	9	8	7	4	7	14	10	8	6	1	1	-
Male	7	6	6	3	5	5	5	4	2	-	1	-
Female	2	2	1	1	2	9	5	4	4	1	-	-
Asian or Pacific Islander 2/, both sexes	4	4	3	9	8	6	14	7	6	3	1	-
Male	3	2	2	7	6	3	3	2	1	-	-	-
Female	1	2	1	2	2	3	11	5	5	3	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Minnesota												
All races, both sexes	160	212	238	263	318	479	650	825	668	267	44	-
Male	133	179	172	181	212	307	372	417	266	67	6	-
Female	27	33	66	82	106	172	278	408	402	200	38	-
White, both sexes	153	195	227	247	305	464	640	821	663	264	44	-
Male	129	165	166	170	202	299	368	416	264	67	6	-
Female	24	30	61	77	103	165	272	405	399	197	38	-
Black, both sexes	5	8	5	6	5	2	7	2	1	2	-	-
Male	2	7	3	4	3	-	4	1	1	-	-	-
Female	3	1	2	2	2	2	3	1	-	2	-	-
American Indian 1/, both sexes	2	6	5	8	5	9	2	1	-	1	-	-
Male	2	4	2	5	4	6	-	-	-	-	-	-
Female	-	2	3	3	1	3	2	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	3	1	2	3	4	1	1	4	-	-	-
Male	-	3	1	2	3	2	-	-	1	-	-	-
Female	-	-	-	-	-	2	1	1	3	-	-	-
Mississippi												
All races, both sexes	262	306	361	365	449	475	558	502	364	138	39	1
Male	187	223	252	245	259	267	260	177	100	22	5	1
Female	75	83	109	120	190	208	298	325	264	116	34	-
White, both sexes	145	188	237	237	312	356	424	395	294	107	22	-
Male	108	148	174	164	193	208	203	144	74	15	1	-
Female	37	40	63	73	119	148	221	251	220	92	21	-
Black, both sexes	115	118	123	125	136	118	132	106	70	31	16	1
Male	77	75	77	79	66	58	57	32	26	7	3	1
Female	38	43	46	46	70	60	75	74	44	24	13	-
American Indian 1/, both sexes	2	-	1	3	1	-	1	-	-	-	-	-
Male	2	-	1	2	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	1	-	-	1	-
Male	-	-	-	-	-	1	-	1	-	-	1	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Missouri												
All races, both sexes	397	477	609	692	843	1,188	1,556	1,705	1,167	531	127	1
Male	294	340	440	450	514	679	802	707	363	107	22	1
Female	103	137	169	242	329	509	754	998	804	424	105	-
White, both sexes	347	393	539	605	730	1,074	1,433	1,587	1,088	495	106	1
Male	258	288	388	403	445	621	751	675	344	99	18	1
Female	89	105	151	202	285	453	682	912	744	396	88	-
Black, both sexes	46	81	69	85	105	110	119	113	75	36	21	-
Male	34	51	51	47	65	56	49	30	18	8	4	-
Female	12	30	18	38	40	54	70	83	57	28	17	-
American Indian 1/, both sexes	4	1	1	2	3	1	1	1	1	-	-	-
Male	2	-	1	-	2	-	-	-	-	-	-	-
Female	2	1	-	2	1	1	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	-	5	3	3	4	3	-	-	-
Male	-	1	-	-	2	2	2	2	1	-	-	-
Female	-	1	-	-	3	1	1	2	2	-	-	-
Montana												
All races, both sexes	51	59	94	76	97	135	167	168	119	41	5	-
Male	42	49	71	56	68	89	90	97	55	16	1	-
Female	9	10	23	20	29	46	77	71	64	25	4	-
White, both sexes	48	55	88	71	92	131	160	167	114	40	4	-
Male	39	46	66	52	64	87	88	97	52	15	1	-
Female	9	9	22	19	28	44	72	70	62	25	3	-
Black, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	4	6	4	4	4	6	1	5	1	1	-
Male	2	3	5	3	3	2	1	-	3	1	-	-
Female	-	1	1	1	1	2	5	1	2	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Nebraska												
All races, both sexes	61	75	95	112	142	213	316	386	310	148	32	-
Male	51	57	77	74	82	141	193	186	103	39	3	-
Female	10	18	18	38	60	72	123	200	207	109	29	-
White, both sexes	56	67	85	105	129	203	309	376	308	148	31	-
Male	46	51	70	71	75	136	190	178	103	39	3	-
Female	10	16	15	34	54	67	119	198	205	109	28	-
Black, both sexes	4	7	7	7	9	9	3	9	2	-	1	-
Male	4	5	6	3	5	4	1	7	-	-	-	-
Female	-	2	1	4	4	5	2	2	2	-	1	-
American Indian 1/, both sexes	1	1	1	-	2	1	2	-	-	-	-	-
Male	1	1	-	-	2	1	1	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	2	-	2	1	-	-	-	-
Male	-	-	1	-	-	-	1	1	-	-	-	-
Female	-	-	1	-	2	-	1	-	-	-	-	-
Nevada												
All races, both sexes	132	189	240	260	274	330	381	296	156	52	6	-
Male	105	146	191	183	173	214	230	154	56	15	2	-
Female	27	43	49	77	101	116	151	142	100	37	4	-
White, both sexes	112	160	212	233	238	304	355	275	145	47	6	-
Male	89	123	171	168	148	197	218	146	53	13	2	-
Female	23	37	41	65	90	107	137	129	92	34	4	-
Black, both sexes	8	20	18	15	23	13	16	12	5	4	-	-
Male	5	15	11	9	16	9	6	3	1	2	-	-
Female	3	5	7	6	7	4	10	9	4	2	-	-
American Indian 1/, both sexes	3	1	2	2	2	4	4	2	-	-	-	-
Male	3	1	2	1	2	3	4	-	-	-	-	-
Female	-	-	-	1	-	1	-	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	8	8	10	11	9	6	7	6	1	-	-
Male	8	7	7	5	7	5	2	5	2	-	-	-
Female	1	1	1	5	4	4	4	2	4	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
New Hampshire												
All races, both sexes	45	68	114	99	144	191	290	313	249	94	21	-
Male	41	52	82	71	94	115	150	136	78	24	4	-
Female	4	16	32	28	50	76	140	177	171	70	17	-
White, both sexes	45	67	113	98	143	189	290	313	247	94	21	-
Male	41	51	82	70	94	115	150	136	78	24	4	-
Female	4	16	31	28	49	74	140	177	169	70	17	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	2	-	-	2	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	2	-	-	2	-	-	-
New Jersey												
All races, both sexes	321	496	574	748	1,011	1,580	2,281	2,599	1,915	841	200	-
Male	250	375	416	492	618	899	1,202	1,093	618	197	39	-
Female	71	121	158	256	393	681	1,079	1,506	1,297	644	161	-
White, both sexes	248	391	461	597	821	1,368	2,063	2,406	1,797	783	184	-
Male	193	310	341	403	504	787	1,092	1,019	592	192	36	-
Female	55	81	120	194	317	581	971	1,387	1,205	591	148	-
Black, both sexes	64	91	91	130	154	179	186	163	94	49	14	-
Male	48	52	58	71	91	96	89	59	22	3	2	-
Female	16	39	33	59	63	83	97	104	72	46	12	-
American Indian 1/, both sexes	-	-	-	-	1	1	1	1	3	1	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	3	1	-	-
Asian or Pacific Islander 2/, both sexes	9	14	22	21	35	32	31	29	21	8	2	-
Male	9	13	17	18	22	16	21	15	4	2	1	-
Female	-	1	5	3	13	16	10	14	17	6	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
New Mexico												
All races, both sexes	80	135	169	151	206	263	355	364	261	116	31	-
Male	61	107	119	103	132	155	197	176	110	31	7	-
Female	19	28	50	48	74	108	158	188	151	85	24	-
White, both sexes	74	120	155	143	189	248	331	348	248	111	29	-
Male	56	97	109	95	124	148	182	168	104	29	7	-
Female	18	23	46	48	65	100	149	180	144	82	22	-
Black, both sexes	2	4	7	1	3	3	7	3	5	-	1	-
Male	2	2	5	1	1	2	3	1	1	-	-	-
Female	-	2	2	-	2	1	4	2	4	-	1	-
American Indian 1/, both sexes	3	10	7	7	14	11	15	10	7	4	1	-
Male	2	7	5	7	7	5	11	5	4	2	-	-
Female	1	3	2	-	7	6	4	5	3	2	1	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	1	2	3	1	1	-	-
Male	1	1	-	-	-	-	1	2	1	-	-	-
Female	-	-	-	-	-	1	1	1	-	1	-	-
New York												
All races, both sexes	919	1,338	1,761	2,221	2,861	4,564	6,640	7,319	5,965	2,865	660	-
Male	673	954	1,192	1,444	1,665	2,525	3,211	2,908	1,921	607	109	-
Female	246	384	569	777	1,196	2,039	3,429	4,411	4,044	2,258	551	-
White, both sexes	662	997	1,315	1,647	2,241	3,757	5,688	6,425	5,410	2,596	575	-
Male	508	754	932	1,113	1,342	2,130	2,835	2,608	1,767	563	93	-
Female	154	243	383	534	899	1,627	2,853	3,817	3,643	2,033	482	-
Black, both sexes	229	286	389	486	529	665	794	745	440	223	67	-
Male	141	160	217	271	265	313	308	229	109	30	11	-
Female	88	126	172	215	264	352	486	516	331	193	56	-
American Indian 1/, both sexes	4	4	5	9	4	5	6	6	2	2	-	-
Male	3	1	5	4	3	4	2	3	-	-	-	-
Female	1	3	-	5	1	1	4	3	2	2	-	-
Asian or Pacific Islander 2/, both sexes	24	51	52	79	87	137	152	143	113	44	18	-
Male	21	39	38	56	55	78	66	68	45	14	5	-
Female	3	12	14	23	32	59	86	75	68	30	13	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
North Carolina												
All races, both sexes	511	693	860	910	1,097	1,386	1,760	1,723	1,126	437	103	2
Male	369	521	618	629	697	807	882	714	345	101	9	1
Female	142	172	242	281	400	579	878	1,009	781	336	94	1
White, both sexes	355	492	660	692	876	1,135	1,464	1,481	965	356	79	-
Male	257	375	494	493	565	682	763	628	310	82	7	-
Female	98	117	166	199	311	453	701	853	655	274	72	-
Black, both sexes	146	185	188	199	208	232	279	227	157	76	24	2
Male	103	134	113	122	124	112	110	80	35	17	2	1
Female	43	51	75	77	84	120	169	147	122	59	22	1
American Indian 1/, both sexes	7	11	11	14	11	13	15	12	3	2	-	-
Male	6	8	10	11	6	9	7	4	-	1	-	-
Female	1	3	1	3	5	4	8	8	3	1	-	-
Asian or Pacific Islander 2/, both sexes	3	5	1	5	2	6	2	3	1	3	-	-
Male	3	4	1	3	2	4	2	2	-	1	-	-
Female	-	1	-	2	-	2	-	1	1	2	-	-
North Dakota												
All races, both sexes	33	33	45	62	76	118	156	209	173	81	14	-
Male	28	25	33	44	57	73	88	103	72	23	2	-
Female	5	8	12	18	19	45	68	106	101	58	12	-
White, both sexes	30	30	41	61	75	115	153	208	171	80	14	-
Male	25	24	31	44	57	72	86	102	71	22	2	-
Female	5	6	10	17	18	43	67	106	100	58	12	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	3	4	1	1	3	3	1	2	-	-	-
Male	2	1	2	-	-	1	2	1	1	-	-	-
Female	-	2	2	1	1	2	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Ohio												
All races, both sexes	669	937	1,143	1,197	1,560	2,258	3,138	3,419	2,348	889	194	-
Male	502	691	827	825	971	1,271	1,592	1,460	747	211	28	-
Female	167	246	316	372	589	987	1,546	1,959	1,601	678	166	-
White, both sexes	561	781	993	1,053	1,351	2,046	2,909	3,228	2,222	828	178	-
Male	431	592	731	739	851	1,164	1,481	1,377	713	195	27	-
Female	130	189	262	314	500	882	1,428	1,851	1,509	633	151	-
Black, both sexes	105	148	142	139	203	203	221	187	123	61	14	-
Male	68	95	90	83	115	99	106	80	32	16	1	-
Female	37	53	52	56	88	104	115	107	91	45	13	-
American Indian 1/, both sexes	-	1	3	2	-	1	1	-	-	-	1	-
Male	-	-	2	1	-	1	1	-	-	-	-	-
Female	-	1	1	1	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	3	7	5	3	6	8	7	4	3	-	1	-
Male	3	4	4	2	5	7	4	3	2	-	-	-
Female	-	3	1	1	1	1	3	1	1	-	1	-
Oklahoma												
All races, both sexes	282	398	481	528	609	792	1,007	1,022	699	307	74	-
Male	213	273	312	337	372	442	512	413	255	63	14	-
Female	69	125	169	191	237	350	495	609	444	244	60	-
White, both sexes	235	327	413	459	531	685	914	957	652	282	65	-
Male	177	218	271	288	333	384	471	387	239	58	12	-
Female	58	109	142	171	198	301	443	570	413	224	53	-
Black, both sexes	22	37	29	31	37	54	49	27	24	15	7	-
Male	18	29	19	21	20	26	20	6	12	4	2	-
Female	4	8	10	10	17	28	29	21	12	11	5	-
American Indian 1/, both sexes	24	31	34	36	36	50	41	35	23	9	2	-
Male	17	25	20	27	18	31	21	19	4	1	-	-
Female	7	6	14	9	18	19	20	16	19	8	2	-
Asian or Pacific Islander 2/, both sexes	1	3	5	2	5	3	3	3	-	1	-	-
Male	1	1	2	1	1	1	-	1	-	-	-	-
Female	-	2	3	1	4	2	3	2	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Oregon												
All races, both sexes	142	223	258	283	357	462	616	691	542	198	35	1
Male	113	162	189	215	237	291	380	333	223	52	6	1
Female	29	61	69	68	120	171	236	358	319	146	29	-
White, both sexes	131	210	250	272	346	448	598	683	538	193	35	1
Male	105	152	182	207	230	284	370	330	221	51	6	1
Female	26	58	68	65	116	164	228	353	317	142	29	-
Black, both sexes	4	9	1	5	3	6	5	1	2	2	-	-
Male	3	7	1	4	3	5	3	-	1	-	-	-
Female	1	2	-	1	-	1	2	1	1	2	-	-
American Indian 1/, both sexes	5	1	1	4	4	2	4	3	-	1	-	-
Male	3	1	1	2	1	-	1	2	-	-	-	-
Female	2	-	-	2	3	2	3	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	2	3	6	2	4	6	9	4	2	2	-	-
Male	2	2	5	2	3	2	6	1	1	1	-	-
Female	-	1	1	-	1	4	3	3	1	1	-	-
Pennsylvania												
All races, both sexes	608	945	1,191	1,279	1,688	2,693	3,754	4,125	2,963	1,137	237	3
Male	476	725	854	858	1,071	1,535	1,922	1,792	979	289	42	2
Female	132	220	337	421	617	1,158	1,832	2,333	1,984	848	195	1
White, both sexes	497	794	1,026	1,123	1,494	2,447	3,521	3,925	2,821	1,072	219	2
Male	397	626	758	760	956	1,422	1,822	1,722	951	272	39	1
Female	100	168	268	363	538	1,025	1,699	2,203	1,870	800	180	1
Black, both sexes	105	142	162	144	183	233	212	182	136	63	18	1
Male	74	93	96	89	107	106	93	60	24	16	3	1
Female	31	49	66	55	76	127	119	122	112	47	15	-
American Indian 1/, both sexes	-	2	-	1	1	1	2	2	-	-	-	-
Male	-	2	-	1	1	1	2	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	7	3	11	10	12	19	16	6	2	-	-
Male	5	4	-	8	7	6	5	9	4	1	-	-
Female	1	3	3	3	3	6	14	7	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Rhode Island												
All races, both sexes	55	93	87	105	139	232	360	468	393	182	34	-
Male	47	73	60	75	84	131	160	184	117	43	1	-
Female	8	20	27	30	55	101	200	284	276	139	33	-
White, both sexes	51	86	76	101	128	221	352	458	391	181	34	-
Male	43	68	52	73	78	127	158	181	117	42	1	-
Female	8	18	24	28	50	94	194	277	274	139	33	-
Black, both sexes	2	5	9	3	9	9	8	10	2	1	-	-
Male	2	5	7	1	5	3	2	3	-	1	-	-
Female	-	-	2	2	4	6	6	7	2	-	-	-
American Indian 1/, both sexes	1	1	2	-	1	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	1	1	2	-	-	-	-	-	-
Male	1	-	-	1	1	1	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
South Carolina												
All races, both sexes	259	346	466	490	559	688	805	689	455	157	24	1
Male	188	256	347	335	344	400	412	301	144	32	4	1
Female	71	90	119	155	215	288	393	388	311	125	20	-
White, both sexes	169	235	329	356	403	529	646	553	384	119	16	1
Male	130	173	251	252	256	326	343	260	127	26	3	1
Female	39	62	78	104	147	203	303	293	257	93	13	-
Black, both sexes	86	109	135	128	153	157	157	136	70	38	8	-
Male	54	82	95	78	85	73	68	41	17	6	1	-
Female	32	27	40	50	68	84	89	95	53	32	7	-
American Indian 1/, both sexes	2	2	2	1	1	-	-	-	-	-	-	-
Male	2	1	1	1	1	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	5	2	2	2	-	1	-	-	-
Male	2	-	-	4	2	1	1	-	-	-	-	-
Female	-	-	-	1	-	1	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
South Dakota												
All races, both sexes	35	55	53	73	94	132	201	226	210	89	24	-
Male	31	46	38	52	60	85	120	114	86	26	3	-
Female	4	9	15	21	34	47	81	112	124	63	21	-
White, both sexes	31	48	48	68	87	127	192	220	207	88	23	-
Male	29	41	35	49	57	82	115	112	86	26	3	-
Female	2	7	13	19	30	45	77	108	121	62	20	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	1	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	4	6	4	5	7	5	8	6	3	1	-	-
Male	2	5	2	3	3	3	5	2	-	-	-	-
Female	2	1	2	2	4	2	3	4	3	1	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
Tennessee												
All races, both sexes	508	656	757	845	1,013	1,290	1,562	1,596	1,011	413	86	-
Male	355	478	529	553	599	747	751	625	292	84	11	-
Female	153	178	228	292	414	543	811	971	719	329	75	-
White, both sexes	419	538	642	731	862	1,147	1,394	1,440	917	366	74	-
Male	299	394	458	485	525	665	676	566	267	76	10	-
Female	120	144	184	246	337	482	718	874	650	290	64	-
Black, both sexes	86	117	113	109	147	137	166	153	93	47	12	-
Male	55	84	70	64	72	77	73	57	25	8	1	-
Female	31	33	43	45	75	60	93	96	68	39	11	-
American Indian 1/, both sexes	-	-	-	1	1	1	1	-	-	-	-	-
Male	-	-	-	1	-	1	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	1	2	4	3	5	1	3	1	-	-	-
Male	1	-	1	3	2	4	1	2	-	-	-	-
Female	2	1	1	1	1	1	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Texas												
All races, both sexes	1,160	1,612	1,981	1,988	2,365	3,219	3,991	3,916	2,576	1,052	205	-
Male	857	1,183	1,373	1,323	1,483	1,826	2,053	1,745	862	266	33	-
Female	303	429	608	665	882	1,393	1,938	2,171	1,714	786	172	-
White, both sexes	934	1,345	1,685	1,682	2,050	2,864	3,611	3,629	2,373	936	174	-
Male	697	1,003	1,173	1,153	1,302	1,646	1,884	1,639	802	237	25	-
Female	237	342	512	529	748	1,218	1,727	1,990	1,571	699	149	-
Black, both sexes	212	251	267	269	290	319	355	257	190	111	28	-
Male	150	169	179	149	164	166	159	94	53	26	5	-
Female	62	82	88	120	126	153	196	163	137	85	23	-
American Indian 1/, both sexes	2	3	2	12	6	2	1	4	2	-	-	-
Male	2	3	1	8	5	1	1	2	-	-	-	-
Female	-	-	1	4	1	1	-	2	2	-	-	-
Asian or Pacific Islander 2/, both sexes	12	13	27	25	19	34	24	26	11	5	3	-
Male	8	8	20	13	12	13	9	10	7	3	3	-
Female	4	5	7	12	7	21	15	16	4	2	-	-
Utah												
All races, both sexes	51	97	97	98	123	159	272	254	148	75	6	-
Male	37	79	74	67	88	90	148	135	69	23	1	-
Female	14	18	23	31	35	69	124	119	79	52	5	-
White, both sexes	48	90	89	96	120	156	268	251	147	74	6	-
Male	35	73	68	67	87	88	147	132	69	23	1	-
Female	13	17	21	29	33	68	121	119	78	51	5	-
Black, both sexes	1	-	3	-	-	1	1	1	-	-	-	-
Male	1	-	2	-	-	1	-	1	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	1	2	1	2	2	2	2	1	1	-	-	-
Male	1	2	1	-	-	1	1	1	-	-	-	-
Female	-	-	-	2	2	1	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	5	4	-	1	-	1	1	-	1	-	-
Male	-	4	3	-	1	-	-	1	-	-	-	-
Female	1	1	1	-	-	-	1	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Vermont												
All races, both sexes	32	37	51	55	57	104	145	149	113	49	12	-
Male	25	26	40	32	37	68	69	65	38	13	1	-
Female	7	11	11	23	20	36	76	84	75	36	11	-
White, both sexes	32	37	51	54	57	104	142	149	113	49	12	-
Male	25	26	40	32	37	68	68	65	38	13	1	-
Female	7	11	11	22	20	36	74	84	75	36	11	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Virginia												
All races, both sexes	383	517	543	656	759	1,043	1,330	1,261	828	324	70	-
Male	284	402	392	438	496	607	667	550	286	88	14	-
Female	99	115	151	218	263	436	663	711	542	236	56	-
White, both sexes	275	374	396	489	609	815	1,118	1,064	709	274	45	-
Male	213	302	290	340	413	474	579	470	244	77	9	-
Female	62	72	106	149	196	341	539	594	465	197	36	-
Black, both sexes	103	135	139	156	136	210	193	180	108	49	25	-
Male	68	93	95	90	76	123	80	72	36	11	5	-
Female	35	42	44	66	60	87	113	108	72	38	20	-
American Indian 1/, both sexes	1	1	-	-	-	1	2	-	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	7	8	11	14	17	17	17	11	1	-	-
Male	3	6	7	8	7	10	7	8	6	-	-	-
Female	1	1	1	3	7	7	10	9	5	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Washington												
All races, both sexes	261	334	458	453	644	917	1,178	1,373	1,031	414	92	1
Male	211	262	336	332	420	534	626	660	402	111	20	1
Female	50	72	122	121	224	383	552	713	629	303	72	-
White, both sexes	235	300	417	410	592	841	1,099	1,320	991	403	90	1
Male	189	232	305	301	386	496	591	639	385	109	20	1
Female	46	68	112	109	206	345	508	681	606	294	70	-
Black, both sexes	10	11	16	17	22	28	22	10	12	1	2	-
Male	10	10	12	12	17	14	11	5	6	1	-	-
Female	-	1	4	5	5	14	11	5	6	-	2	-
American Indian 1/, both sexes	7	11	11	11	11	13	8	11	1	3	-	-
Male	5	9	9	6	4	7	4	5	1	-	-	-
Female	2	2	2	5	7	6	4	6	-	3	-	-
Asian or Pacific Islander 2/, both sexes	9	12	14	15	19	35	49	32	27	7	-	-
Male	7	11	10	13	13	17	20	11	10	1	-	-
Female	2	1	4	2	6	18	29	21	17	6	-	-
West Virginia												
All races, both sexes	151	177	218	293	317	434	612	550	384	144	27	-
Male	115	134	151	188	204	223	333	217	115	21	3	-
Female	36	43	67	105	113	211	279	333	269	123	24	-
White, both sexes	145	175	210	283	307	421	606	533	372	140	26	-
Male	110	132	143	180	198	215	331	213	112	20	3	-
Female	35	43	67	103	109	206	275	320	260	120	23	-
Black, both sexes	6	2	8	6	10	13	5	15	12	4	1	-
Male	5	2	8	5	6	8	2	4	3	1	-	-
Female	1	-	-	1	4	5	3	11	9	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	4	-	-	1	2	-	-	-	-
Male	-	-	-	3	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Wisconsin												
All races, both sexes	239	351	420	435	568	814	1,118	1,238	956	373	86	-
Male	182	286	306	319	392	488	638	550	339	89	12	-
Female	57	65	114	116	176	326	480	688	617	284	74	-
White, both sexes	210	310	395	407	535	785	1,089	1,211	942	367	84	-
Male	161	262	290	303	374	473	627	540	336	88	12	-
Female	49	48	105	104	161	312	462	671	606	279	72	-
Black, both sexes	27	29	18	19	26	21	18	20	11	6	1	-
Male	19	14	11	12	15	11	5	6	1	1	-	-
Female	8	15	7	7	11	10	13	14	10	5	1	-
American Indian 1/, both sexes	2	10	4	7	3	6	8	4	1	-	1	-
Male	2	8	3	4	2	3	4	3	1	-	-	-
Female	-	2	1	3	1	3	4	1	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	2	3	2	4	2	3	3	2	-	-	-
Male	-	2	2	-	1	1	2	1	1	-	-	-
Female	-	-	1	2	3	1	1	2	1	-	-	-
Wyoming												
All races, both sexes	32	52	37	51	56	72	105	87	49	11	4	-
Male	23	35	31	34	40	45	55	41	19	4	2	-
Female	9	17	6	17	16	27	50	46	30	7	2	-
White, both sexes	32	49	37	50	55	70	102	87	49	11	4	-
Male	23	33	31	33	39	44	54	41	19	4	2	-
Female	9	16	6	17	16	26	48	46	30	7	2	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	3	-	1	-	1	2	-	-	-	-	-
Male	-	2	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
United States												
All races, both sexes	132,968	10	4	5	6	16	38	125	275	684	1,718	3,609
Male	71,712	5	-	-	5	10	23	91	206	495	1,273	2,727
Female	61,256	5	4	5	1	6	15	34	69	189	445	882
White, both sexes	115,900	9	2	5	5	10	26	87	197	540	1,366	2,952
Male	63,011	5	-	-	4	6	16	64	149	401	1,029	2,291
Female	52,889	4	2	5	1	4	10	23	48	139	337	661
Black, both sexes	14,097	1	1	-	1	6	12	30	66	121	294	564
Male	6,997	-	-	-	1	4	7	20	49	80	196	366
Female	7,100	1	1	-	-	2	5	10	17	41	98	198
American Indian 1/, both sexes	591	-	1	-	-	-	-	3	5	9	15	25
Male	359	-	-	-	-	-	-	2	3	4	10	17
Female	232	-	1	-	-	-	-	1	2	5	5	8
Asian or Pacific Islander 2/, both sexes	2,380	-	-	-	-	-	-	5	7	14	43	68
Male	1,345	-	-	-	-	-	-	5	5	10	38	53
Female	1,035	-	-	-	-	-	-	-	2	4	5	15
Alabama												
All races, both sexes	2,587	-	-	-	-	-	4	6	10	25	48	100
Male	1,418	-	-	-	-	-	2	3	9	15	32	72
Female	1,169	-	-	-	-	-	2	3	1	10	16	28
White, both sexes	2,038	-	-	-	-	-	3	5	7	21	35	81
Male	1,148	-	-	-	-	-	2	2	6	12	25	63
Female	890	-	-	-	-	-	1	3	1	9	10	18
Black, both sexes	540	-	-	-	-	-	1	1	3	4	13	19
Male	267	-	-	-	-	-	-	1	3	3	7	9
Female	273	-	-	-	-	-	1	-	-	1	6	10
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Alaska												
All races, both sexes	59	-	-	-	-	-	-	-	-	1	3	2
Male	39	-	-	-	-	-	-	-	-	-	3	1
Female	20	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	43	-	-	-	-	-	-	-	-	1	2	2
Male	26	-	-	-	-	-	-	-	-	-	2	1
Female	17	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1,699	-	-	-	-	1	1	1	3	7	14	31
Male	908	-	-	-	-	-	-	1	1	2	12	25
Female	791	-	-	-	-	1	1	-	2	5	2	6
White, both sexes	1,601	-	-	-	-	1	1	-	2	6	11	28
Male	861	-	-	-	-	-	-	-	1	1	9	23
Female	740	-	-	-	-	1	1	-	1	5	2	5
Black, both sexes	39	-	-	-	-	-	-	-	-	-	1	2
Male	14	-	-	-	-	-	-	-	-	-	1	1
Female	25	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	43	-	-	-	-	-	-	1	1	-	2	1
Male	25	-	-	-	-	-	-	1	-	-	2	1
Female	18	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	-	-	1	-	-
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Arkansas												
All races, both sexes	2,975	-	1	-	-	1	3	2	10	26	63	139
Male	1,667	-	-	-	-	-	3	2	9	20	47	103
Female	1,308	-	1	-	-	1	-	-	1	6	16	36
White, both sexes	2,552	-	1	-	-	-	1	2	7	24	49	114
Male	1,447	-	-	-	-	-	1	2	7	20	38	89
Female	1,105	-	1	-	-	-	-	-	-	4	11	25
Black, both sexes	405	-	-	-	-	1	2	-	2	2	12	25
Male	209	-	-	-	-	-	2	-	2	-	8	14
Female	196	-	-	-	-	1	-	-	-	2	4	11
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	1	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	1	-	1	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	11,963	-	1	-	-	1	4	8	12	37	90	218
Male	6,320	-	-	-	-	1	2	6	11	24	66	168
Female	5,643	-	1	-	-	-	2	2	1	13	24	50
White, both sexes	9,853	-	1	-	-	-	3	4	8	29	73	172
Male	5,249	-	-	-	-	-	1	3	7	19	53	133
Female	4,604	-	1	-	-	-	2	1	1	10	20	39
Black, both sexes	935	-	-	-	-	1	1	2	3	5	8	18
Male	430	-	-	-	-	1	1	1	3	2	4	12
Female	505	-	-	-	-	-	-	1	-	3	4	6
American Indian 1/, both sexes	56	-	-	-	-	-	-	-	-	1	-	1
Male	35	-	-	-	-	-	-	-	-	1	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1,119	-	-	-	-	-	-	2	1	2	9	27
Male	606	-	-	-	-	-	-	2	1	2	9	23
Female	513	-	-	-	-	-	-	-	-	-	-	4

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Colorado												
All races, both sexes	1,208	-	-	-	-	-	-	3	1	5	13	28
Male	686	-	-	-	-	-	-	1	-	4	8	26
Female	522	-	-	-	-	-	-	2	1	1	5	2
White, both sexes	1,148	-	-	-	-	-	-	1	1	5	11	27
Male	649	-	-	-	-	-	-	1	-	4	8	25
Female	499	-	-	-	-	-	-	-	1	1	3	2
Black, both sexes	41	-	-	-	-	-	-	2	-	-	1	1
Male	25	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	2	-	-	1	-
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
Connecticut												
All races, both sexes	1,203	1	-	-	-	-	1	-	-	7	16	28
Male	577	-	-	-	-	-	1	-	-	6	9	25
Female	626	1	-	-	-	-	-	-	-	1	7	3
White, both sexes	1,116	1	-	-	-	-	1	-	-	7	10	26
Male	526	-	-	-	-	-	1	-	-	6	3	24
Female	590	1	-	-	-	-	-	-	-	1	7	2
Black, both sexes	75	-	-	-	-	-	-	-	-	-	6	1
Male	43	-	-	-	-	-	-	-	-	-	6	-
Female	32	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Delaware												
All races, both sexes	314	-	-	-	-	-	-	-	1	-	6	9
Male	187	-	-	-	-	-	-	-	1	-	4	7
Female	127	-	-	-	-	-	-	-	-	-	2	2
White, both sexes	265	-	-	-	-	-	-	-	1	-	3	7
Male	161	-	-	-	-	-	-	-	1	-	3	6
Female	104	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	47	-	-	-	-	-	-	-	-	-	3	2
Male	25	-	-	-	-	-	-	-	-	-	1	1
Female	22	-	-	-	-	-	-	-	-	-	2	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	194	-	-	-	-	-	-	-	2	2	1	4
Male	92	-	-	-	-	-	-	-	2	2	-	3
Female	102	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	48	-	-	-	-	-	-	-	-	-	-	-
Male	28	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	144	-	-	-	-	-	-	-	2	2	1	4
Male	63	-	-	-	-	-	-	-	2	2	-	3
Female	81	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Florida												
All races, both sexes	7,966	-	-	-	-	2	2	8	11	34	60	160
Male	4,226	-	-	-	-	2	1	6	10	24	43	121
Female	3,740	-	-	-	-	-	1	2	1	10	17	39
White, both sexes	7,157	-	-	-	-	2	1	5	9	26	45	132
Male	3,833	-	-	-	-	2	1	4	8	19	32	101
Female	3,324	-	-	-	-	-	-	1	1	7	13	31
Black, both sexes	754	-	-	-	-	-	1	3	2	7	13	27
Male	359	-	-	-	-	-	-	2	2	4	9	19
Female	395	-	-	-	-	-	1	1	-	3	4	8
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	51	-	-	-	-	-	-	-	-	-	2	1
Male	31	-	-	-	-	-	-	-	-	-	2	1
Female	20	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	2,928	1	-	-	-	1	1	5	3	23	51	115
Male	1,603	1	-	-	-	-	-	4	3	17	33	78
Female	1,325	-	-	-	-	1	1	1	-	6	18	37
White, both sexes	2,203	1	-	-	-	-	-	2	2	12	37	84
Male	1,236	1	-	-	-	-	-	2	2	8	25	60
Female	967	-	-	-	-	-	-	-	-	4	12	24
Black, both sexes	701	-	-	-	-	1	1	3	1	10	14	31
Male	353	-	-	-	-	-	-	2	1	8	8	18
Female	348	-	-	-	-	1	1	1	-	2	6	13
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	22	-	-	-	-	-	-	-	-	1	-	-
Male	14	-	-	-	-	-	-	-	-	1	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Hawaii												
All races, both sexes	393	-	-	-	-	-	-	1	1	5	7	9
Male	223	-	-	-	-	-	-	1	-	3	4	5
Female	170	-	-	-	-	-	-	-	1	2	3	4
White, both sexes	104	-	-	-	-	-	-	-	-	1	2	2
Male	62	-	-	-	-	-	-	-	-	1	1	2
Female	42	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	286	-	-	-	-	-	-	1	1	4	5	7
Male	159	-	-	-	-	-	-	1	-	2	3	3
Female	127	-	-	-	-	-	-	-	1	2	2	4
Idaho												
All races, both sexes	762	-	-	-	-	1	-	1	1	3	8	21
Male	494	-	-	-	-	1	-	-	1	2	5	17
Female	268	-	-	-	-	-	-	1	-	1	3	4
White, both sexes	745	-	-	-	-	1	-	1	1	2	6	21
Male	479	-	-	-	-	1	-	-	1	1	4	17
Female	266	-	-	-	-	-	-	1	-	1	2	4
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	1	1	-
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Illinois												
All races, both sexes	6,136	-	-	-	-	-	-	5	12	26	85	187
Male	3,264	-	-	-	-	-	-	5	8	24	68	138
Female	2,872	-	-	-	-	-	-	-	4	2	17	49
White, both sexes	5,220	-	-	-	-	-	-	5	8	18	64	143
Male	2,820	-	-	-	-	-	-	5	5	17	55	108
Female	2,400	-	-	-	-	-	-	-	3	1	9	35
Black, both sexes	840	-	-	-	-	-	-	-	4	7	20	42
Male	407	-	-	-	-	-	-	-	3	6	13	29
Female	433	-	-	-	-	-	-	-	1	1	7	13
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	73	-	-	-	-	-	-	-	-	1	1	2
Male	34	-	-	-	-	-	-	-	-	1	-	1
Female	39	-	-	-	-	-	-	-	-	-	1	1
Indiana												
All races, both sexes	3,279	1	-	-	-	-	-	2	17	32	67	126
Male	1,850	1	-	-	-	-	-	1	11	29	48	94
Female	1,429	-	-	-	-	-	-	1	6	3	19	32
White, both sexes	3,071	1	-	-	-	-	-	2	14	29	56	110
Male	1,723	1	-	-	-	-	-	1	8	26	40	81
Female	1,348	-	-	-	-	-	-	1	6	3	16	29
Black, both sexes	203	-	-	-	-	-	-	-	3	3	11	16
Male	122	-	-	-	-	-	-	-	3	3	8	13
Female	81	-	-	-	-	-	-	-	-	-	3	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Iowa												
All races, both sexes	1,668	-	-	1	-	-	-	1	3	4	25	42
Male	899	-	-	-	-	-	-	1	2	3	17	33
Female	769	-	-	1	-	-	-	-	1	1	8	9
White, both sexes	1,646	-	-	1	-	-	-	1	3	4	24	41
Male	883	-	-	-	-	-	-	1	2	3	16	32
Female	763	-	-	1	-	-	-	-	1	1	8	9
Black, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1,073	-	-	-	-	-	-	3	1	5	10	32
Male	581	-	-	-	-	-	-	3	-	5	8	26
Female	492	-	-	-	-	-	-	-	1	-	2	6
White, both sexes	1,019	-	-	-	-	-	-	2	1	3	9	31
Male	554	-	-	-	-	-	-	2	-	3	8	25
Female	465	-	-	-	-	-	-	-	1	-	1	6
Black, both sexes	41	-	-	-	-	-	-	-	-	2	1	1
Male	20	-	-	-	-	-	-	-	-	2	-	1
Female	21	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	9	-	-	-	-	-	-	1	-	-	-	-
Male	6	-	-	-	-	-	-	1	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Kentucky												
All races, both sexes	3,144	-	-	1	1	-	3	11	14	28	77	119
Male	1,817	-	-	-	-	-	2	8	11	21	53	92
Female	1,327	-	-	1	1	-	1	3	3	7	24	27
White, both sexes	2,908	-	-	1	1	-	1	9	12	26	71	106
Male	1,685	-	-	-	-	-	1	6	10	19	52	82
Female	1,223	-	-	1	1	-	-	3	2	7	19	24
Black, both sexes	229	-	-	-	-	-	2	2	2	2	6	13
Male	128	-	-	-	-	-	1	2	1	2	1	10
Female	101	-	-	-	-	-	1	-	1	-	5	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	2,426	-	-	-	-	2	2	5	18	21	66	118
Male	1,436	-	-	-	-	2	2	3	12	15	45	84
Female	990	-	-	-	-	-	-	2	6	6	21	34
White, both sexes	1,646	-	-	-	-	1	-	2	9	10	34	68
Male	968	-	-	-	-	1	-	2	7	7	23	50
Female	678	-	-	-	-	-	-	-	2	3	11	18
Black, both sexes	761	-	-	-	-	1	2	3	8	11	31	49
Male	457	-	-	-	-	1	2	1	4	8	21	33
Female	304	-	-	-	-	-	-	2	4	3	10	16
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	1	-	1	1
Male	9	-	-	-	-	-	-	-	1	-	1	1
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Maine												
All races, both sexes	638	-	-	-	-	-	-	2	-	3	6	16
Male	358	-	-	-	-	-	-	1	-	2	5	11
Female	280	-	-	-	-	-	-	1	-	1	1	5
White, both sexes	629	-	-	-	-	-	-	1	-	3	5	15
Male	352	-	-	-	-	-	-	1	-	2	4	11
Female	277	-	-	-	-	-	-	-	-	1	1	4
Black, both sexes	4	-	-	-	-	-	-	1	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	2,402	-	-	-	-	-	-	3	7	15	24	64
Male	1,189	-	-	-	-	-	-	2	5	12	19	40
Female	1,213	-	-	-	-	-	-	1	2	3	5	24
White, both sexes	1,772	-	-	-	-	-	-	1	4	8	12	43
Male	887	-	-	-	-	-	-	1	3	7	9	29
Female	885	-	-	-	-	-	-	-	1	1	3	14
Black, both sexes	583	-	-	-	-	-	-	2	3	7	10	19
Male	273	-	-	-	-	-	-	1	2	5	8	10
Female	310	-	-	-	-	-	-	1	1	2	2	9
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	44	-	-	-	-	-	-	-	-	-	2	1
Male	27	-	-	-	-	-	-	-	-	-	2	-
Female	17	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Massachusetts												
All races, both sexes	2,432	-	-	-	-	1	1	-	-	8	22	35
Male	1,198	-	-	-	-	1	-	-	-	5	18	28
Female	1,234	-	-	-	-	-	1	-	-	3	4	7
White, both sexes	2,322	-	-	-	-	1	1	-	-	8	19	33
Male	1,143	-	-	-	-	1	-	-	-	5	16	26
Female	1,179	-	-	-	-	-	1	-	-	3	3	7
Black, both sexes	77	-	-	-	-	-	-	-	-	-	2	2
Male	34	-	-	-	-	-	-	-	-	-	1	2
Female	43	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	31	-	-	-	-	-	-	-	-	-	1	-
Male	20	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	4,644	-	-	-	-	1	3	1	16	21	49	115
Male	2,449	-	-	-	-	1	2	1	14	17	41	91
Female	2,195	-	-	-	-	-	1	-	2	4	8	24
White, both sexes	4,063	-	-	-	-	-	3	1	11	16	42	94
Male	2,166	-	-	-	-	-	2	1	10	14	34	74
Female	1,897	-	-	-	-	-	1	-	1	2	8	20
Black, both sexes	518	-	-	-	-	1	-	-	4	4	5	19
Male	246	-	-	-	-	1	-	-	3	3	5	15
Female	272	-	-	-	-	-	-	-	1	1	-	4
American Indian 1/, both sexes	33	-	-	-	-	-	-	-	-	1	2	1
Male	19	-	-	-	-	-	-	-	-	-	2	1
Female	14	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	30	-	-	-	-	-	-	-	1	-	-	1
Male	18	-	-	-	-	-	-	-	1	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Minnesota												
All races, both sexes	1,246	-	-	-	-	-	-	1	1	4	14	29
Male	696	-	-	-	-	-	-	1	1	3	10	22
Female	550	-	-	-	-	-	-	-	-	1	4	7
White, both sexes	1,213	-	-	-	-	-	-	1	1	2	10	25
Male	676	-	-	-	-	-	-	1	1	2	8	20
Female	537	-	-	-	-	-	-	-	-	-	2	5
Black, both sexes	13	-	-	-	-	-	-	-	-	1	1	1
Male	9	-	-	-	-	-	-	-	-	1	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	1	1	3
Male	7	-	-	-	-	-	-	-	-	-	-	2
Female	6	-	-	-	-	-	-	-	-	1	1	1
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	2	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
Mississippi												
All races, both sexes	1,842	-	-	-	-	-	-	2	5	15	56	75
Male	1,061	-	-	-	-	-	-	1	4	11	44	55
Female	781	-	-	-	-	-	-	1	1	4	12	20
White, both sexes	1,225	-	-	-	-	-	-	1	3	6	32	42
Male	720	-	-	-	-	-	-	-	3	6	26	30
Female	505	-	-	-	-	-	-	1	-	-	6	12
Black, both sexes	612	-	-	-	-	-	-	1	2	9	24	32
Male	339	-	-	-	-	-	-	1	1	5	18	24
Female	273	-	-	-	-	-	-	-	1	4	6	8
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Missouri												
All races, both sexes	4,169	-	-	-	-	-	2	6	15	22	47	126
Male	2,226	-	-	-	-	-	1	4	13	17	35	97
Female	1,943	-	-	-	-	-	1	2	2	5	12	29
White, both sexes	3,784	-	-	-	-	-	2	4	12	22	39	109
Male	2,038	-	-	-	-	-	1	2	12	17	29	85
Female	1,746	-	-	-	-	-	1	2	-	5	10	24
Black, both sexes	366	-	-	-	-	-	-	2	3	-	6	15
Male	178	-	-	-	-	-	-	2	1	-	4	10
Female	188	-	-	-	-	-	-	-	2	-	2	5
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	-	2	2
Male	10	-	-	-	-	-	-	-	-	-	2	2
Female	4	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	290	-	-	-	-	-	-	1	-	2	3	9
Male	182	-	-	-	-	-	-	1	-	2	3	8
Female	108	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	275	-	-	-	-	-	-	1	-	2	2	9
Male	172	-	-	-	-	-	-	1	-	2	2	8
Female	103	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Nebraska												
All races, both sexes	601	-	-	-	-	-	-	-	1	6	11	17
Male	340	-	-	-	-	-	-	-	-	3	10	12
Female	261	-	-	-	-	-	-	-	1	3	1	5
White, both sexes	577	-	-	-	-	-	-	-	1	5	11	16
Male	326	-	-	-	-	-	-	-	-	3	10	12
Female	251	-	-	-	-	-	-	-	1	2	1	4
Black, both sexes	19	-	-	-	-	-	-	-	-	1	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	522	-	-	-	-	-	-	4	1	2	11	17
Male	331	-	-	-	-	-	-	3	1	1	9	12
Female	191	-	-	-	-	-	-	1	-	1	2	5
White, both sexes	475	-	-	-	-	-	-	2	1	2	9	14
Male	305	-	-	-	-	-	-	2	1	1	8	11
Female	170	-	-	-	-	-	-	-	-	1	1	3
Black, both sexes	29	-	-	-	-	-	-	2	-	-	2	1
Male	15	-	-	-	-	-	-	1	-	-	1	1
Female	14	-	-	-	-	-	-	1	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	-	-	-	-	-	-	2
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
New Hampshire												
All races, both sexes	431	-	-	-	-	-	-	-	2	4	7	7
Male	247	-	-	-	-	-	-	-	1	1	7	6
Female	184	-	-	-	-	-	-	-	1	3	-	1
White, both sexes	427	-	-	-	-	-	-	-	2	4	7	7
Male	246	-	-	-	-	-	-	-	1	1	7	6
Female	181	-	-	-	-	-	-	-	1	3	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	3,887	-	-	-	-	-	1	2	2	9	33	84
Male	1,955	-	-	-	-	-	-	2	2	6	26	63
Female	1,932	-	-	-	-	-	1	-	-	3	7	21
White, both sexes	3,339	-	-	-	-	-	1	1	-	7	27	62
Male	1,680	-	-	-	-	-	-	1	-	4	22	48
Female	1,659	-	-	-	-	-	1	-	-	3	5	14
Black, both sexes	459	-	-	-	-	-	-	-	2	1	5	17
Male	213	-	-	-	-	-	-	-	2	1	3	11
Female	246	-	-	-	-	-	-	-	-	-	2	6
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	85	-	-	-	-	-	-	1	-	1	1	4
Male	60	-	-	-	-	-	-	1	-	1	1	3
Female	25	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
New Mexico												
All races, both sexes	555	-	-	-	-	-	-	-	3	3	4	16
Male	334	-	-	-	-	-	-	-	3	2	3	14
Female	221	-	-	-	-	-	-	-	-	1	1	2
White, both sexes	518	-	-	-	-	-	-	-	2	3	3	13
Male	309	-	-	-	-	-	-	-	2	2	2	12
Female	209	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	27	-	-	-	-	-	-	-	1	-	1	3
Male	20	-	-	-	-	-	-	-	1	-	1	2
Female	7	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	8,691	1	-	-	-	3	1	4	11	41	94	160
Male	4,276	1	-	-	-	2	1	4	11	29	64	124
Female	4,415	-	-	-	-	1	-	-	-	12	30	36
White, both sexes	7,537	1	-	-	-	2	1	1	8	33	70	120
Male	3,761	1	-	-	-	1	1	1	8	26	49	97
Female	3,776	-	-	-	-	1	-	-	-	7	21	23
Black, both sexes	967	-	-	-	-	1	-	3	3	6	19	35
Male	415	-	-	-	-	1	-	3	3	2	10	22
Female	552	-	-	-	-	-	-	-	-	4	9	13
American Indian 1/, both sexes	23	-	-	-	-	-	-	-	-	-	1	1
Male	12	-	-	-	-	-	-	-	-	-	1	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	164	-	-	-	-	-	-	-	-	2	4	4
Male	88	-	-	-	-	-	-	-	-	1	4	4
Female	76	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
North Carolina												
All races, both sexes	3,666	-	-	-	-	-	-	1	11	24	51	95
Male	1,974	-	-	-	-	-	-	1	9	13	37	72
Female	1,692	-	-	-	-	-	-	-	2	11	14	23
White, both sexes	2,942	-	-	-	-	-	-	-	9	17	40	74
Male	1,622	-	-	-	-	-	-	-	7	10	30	58
Female	1,320	-	-	-	-	-	-	-	2	7	10	16
Black, both sexes	682	-	-	-	-	-	-	1	2	7	11	21
Male	330	-	-	-	-	-	-	1	2	3	7	14
Female	352	-	-	-	-	-	-	-	-	4	4	7
American Indian 1/, both sexes	33	-	-	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	408	-	-	-	-	-	-	-	-	2	4	10
Male	246	-	-	-	-	-	-	-	-	-	3	9
Female	162	-	-	-	-	-	-	-	-	2	1	1
White, both sexes	395	-	-	-	-	-	-	-	-	-	4	9
Male	238	-	-	-	-	-	-	-	-	-	3	8
Female	157	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	-	2	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	2	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Ohio												
All races, both sexes	5,649	-	1	-	-	-	2	2	13	28	74	153
Male	2,990	-	-	-	-	-	2	1	8	22	56	111
Female	2,659	-	1	-	-	-	-	1	5	6	18	42
White, both sexes	5,175	-	-	-	-	-	2	2	10	26	67	141
Male	2,757	-	-	-	-	-	2	1	6	22	51	106
Female	2,418	-	-	-	-	-	-	1	4	4	16	35
Black, both sexes	457	-	1	-	-	-	-	-	3	2	6	12
Male	218	-	-	-	-	-	-	-	2	-	4	5
Female	239	-	1	-	-	-	-	-	1	2	2	7
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	-	1	-
Male	14	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1,592	-	-	-	-	-	-	4	4	15	23	60
Male	871	-	-	-	-	-	-	3	4	10	17	46
Female	721	-	-	-	-	-	-	1	-	5	6	14
White, both sexes	1,404	-	-	-	-	-	-	4	3	13	21	49
Male	766	-	-	-	-	-	-	3	3	10	15	41
Female	638	-	-	-	-	-	-	1	-	3	6	8
Black, both sexes	82	-	-	-	-	-	-	-	-	1	2	9
Male	41	-	-	-	-	-	-	-	-	-	2	4
Female	41	-	-	-	-	-	-	-	-	1	-	5
American Indian 1/, both sexes	100	-	-	-	-	-	-	-	1	1	-	2
Male	64	-	-	-	-	-	-	-	1	-	-	1
Female	36	-	-	-	-	-	-	-	-	1	-	1
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Oregon												
All races, both sexes	1,295	-	-	-	-	-	-	-	1	2	17	23
Male	743	-	-	-	-	-	-	-	1	2	15	15
Female	552	-	-	-	-	-	-	-	-	-	2	8
White, both sexes	1,263	-	-	-	-	-	-	-	1	2	17	21
Male	722	-	-	-	-	-	-	-	1	2	15	15
Female	541	-	-	-	-	-	-	-	-	-	2	6
Black, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	7,494	-	-	1	-	1	1	2	7	24	73	187
Male	4,032	-	-	-	-	-	-	1	6	18	55	138
Female	3,462	-	-	1	-	1	1	1	1	6	18	49
White, both sexes	6,951	-	-	1	-	1	1	2	6	22	65	170
Male	3,778	-	-	-	-	-	-	1	5	16	47	128
Female	3,173	-	-	1	-	1	1	1	1	6	18	42
Black, both sexes	499	-	-	-	-	-	-	-	-	2	7	15
Male	230	-	-	-	-	-	-	-	-	2	7	9
Female	269	-	-	-	-	-	-	-	-	-	-	6
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	40	-	-	-	-	-	-	-	1	-	1	2
Male	20	-	-	-	-	-	-	-	1	-	1	1
Female	20	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Rhode Island												
All races, both sexes	693	-	-	-	-	-	-	-	2	1	9	4
Male	344	-	-	-	-	-	-	-	2	-	6	4
Female	349	-	-	-	-	-	-	-	-	1	3	-
White, both sexes	673	-	-	-	-	-	-	-	1	1	9	4
Male	333	-	-	-	-	-	-	-	1	-	6	4
Female	340	-	-	-	-	-	-	-	-	1	3	-
Black, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	2,289	-	-	-	-	-	1	1	4	13	40	88
Male	1,325	-	-	-	-	-	-	1	2	10	30	64
Female	964	-	-	-	-	-	1	-	2	3	10	24
White, both sexes	1,682	-	-	-	-	-	-	1	3	9	32	48
Male	1,008	-	-	-	-	-	-	1	2	8	24	36
Female	674	-	-	-	-	-	-	-	1	1	8	12
Black, both sexes	598	-	-	-	-	-	1	-	1	4	8	39
Male	309	-	-	-	-	-	-	-	-	2	6	27
Female	289	-	-	-	-	-	1	-	1	2	2	12
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
South Dakota												
All races, both sexes	545	-	-	-	-	-	-	2	-	1	7	13
Male	317	-	-	-	-	-	-	1	-	1	6	11
Female	228	-	-	-	-	-	-	1	-	-	1	2
White, both sexes	512	-	-	-	-	-	-	2	-	-	4	9
Male	295	-	-	-	-	-	-	1	-	-	4	7
Female	217	-	-	-	-	-	-	1	-	-	-	2
Black, both sexes	3	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	1	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	29	-	-	-	-	-	-	-	-	1	2	3
Male	19	-	-	-	-	-	-	-	-	1	1	3
Female	10	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	4,247	-	-	-	-	-	-	2	12	25	72	142
Male	2,294	-	-	-	-	-	-	1	11	18	58	115
Female	1,953	-	-	-	-	-	-	1	1	7	14	27
White, both sexes	3,728	-	-	-	-	-	-	2	8	20	63	127
Male	2,041	-	-	-	-	-	-	1	7	13	50	105
Female	1,687	-	-	-	-	-	-	1	1	7	13	22
Black, both sexes	505	-	-	-	-	-	-	-	4	5	8	15
Male	245	-	-	-	-	-	-	-	4	5	7	10
Female	260	-	-	-	-	-	-	-	-	-	1	5
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Texas												
All races, both sexes	10,691	5	-	2	3	-	3	11	23	55	145	287
Male	5,834	2	-	-	3	-	2	7	11	38	105	213
Female	4,857	3	-	2	-	-	1	4	12	17	40	74
White, both sexes	9,436	4	-	2	2	-	2	9	14	43	116	246
Male	5,200	2	-	-	2	-	1	6	6	31	85	187
Female	4,236	2	-	2	-	-	1	3	8	12	31	59
Black, both sexes	1,133	1	-	-	1	-	1	1	8	10	25	35
Male	558	-	-	-	1	-	1	1	4	6	16	21
Female	575	1	-	-	-	-	-	-	4	4	9	14
American Indian 1/, both sexes	18	-	-	-	-	-	-	1	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	104	-	-	-	-	-	-	-	1	2	4	5
Male	63	-	-	-	-	-	-	-	1	1	4	4
Female	41	-	-	-	-	-	-	-	-	1	-	1
Utah												
All races, both sexes	516	-	-	-	-	-	-	1	1	5	6	20
Male	310	-	-	-	-	-	-	1	1	4	3	17
Female	206	-	-	-	-	-	-	-	-	1	3	3
White, both sexes	497	-	-	-	-	-	-	1	1	5	5	18
Male	299	-	-	-	-	-	-	1	1	4	3	16
Female	198	-	-	-	-	-	-	-	-	1	2	2
Black, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Vermont												
All races, both sexes	210	-	-	-	-	-	-	-	-	-	-	5
Male	100	-	-	-	-	-	-	-	-	-	-	5
Female	110	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	209	-	-	-	-	-	-	-	-	-	-	5
Male	100	-	-	-	-	-	-	-	-	-	-	5
Female	109	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	2,920	-	-	-	-	-	1	4	7	18	32	82
Male	1,642	-	-	-	-	-	1	3	4	14	27	60
Female	1,278	-	-	-	-	-	-	1	3	4	5	22
White, both sexes	2,333	-	-	-	-	-	1	4	6	13	26	63
Male	1,329	-	-	-	-	-	1	3	3	10	22	47
Female	1,004	-	-	-	-	-	-	1	3	3	4	16
Black, both sexes	548	-	-	-	-	-	-	-	1	5	5	16
Male	289	-	-	-	-	-	-	-	1	4	4	10
Female	259	-	-	-	-	-	-	-	-	1	1	6
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	38	-	-	-	-	-	-	-	-	-	1	3
Male	23	-	-	-	-	-	-	-	-	-	1	3
Female	15	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Washington												
All races, both sexes	2,261	1	1	-	1	1	1	2	3	10	34	57
Male	1,253	-	-	-	1	-	1	2	-	7	24	40
Female	1,008	1	1	-	-	1	-	-	3	3	10	17
White, both sexes	2,101	1	-	-	1	1	1	1	2	10	29	53
Male	1,160	-	-	-	1	-	1	1	-	7	21	37
Female	941	1	-	-	-	1	-	-	2	3	8	16
Black, both sexes	58	-	-	-	-	-	-	-	-	-	3	4
Male	35	-	-	-	-	-	-	-	-	-	2	3
Female	23	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	40	-	1	-	-	-	-	-	-	-	2	-
Male	23	-	-	-	-	-	-	-	-	-	1	-
Female	17	-	1	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	62	-	-	-	-	-	-	1	1	-	-	-
Male	35	-	-	-	-	-	-	1	-	-	-	-
Female	27	-	-	-	-	-	-	-	1	-	-	-
West Virginia												
All races, both sexes	1,086	-	-	-	-	-	-	2	-	4	9	29
Male	562	-	-	-	-	-	-	2	-	2	8	20
Female	524	-	-	-	-	-	-	-	-	2	1	9
White, both sexes	1,061	-	-	-	-	-	-	2	-	4	8	29
Male	546	-	-	-	-	-	-	2	-	2	7	20
Female	515	-	-	-	-	-	-	-	-	2	1	9
Black, both sexes	22	-	-	-	-	-	-	-	-	-	1	-
Male	14	-	-	-	-	-	-	-	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Wisconsin												
All races, both sexes	2,787	-	-	-	1	-	-	3	3	10	25	79
Male	1,567	-	-	-	1	-	-	2	1	8	21	71
Female	1,220	-	-	-	-	-	-	1	2	2	4	8
White, both sexes	2,712	-	-	-	1	-	-	2	3	9	24	73
Male	1,523	-	-	-	1	-	-	1	1	7	20	66
Female	1,189	-	-	-	-	-	-	1	2	2	4	7
Black, both sexes	47	-	-	-	-	-	-	1	-	1	-	3
Male	24	-	-	-	-	-	-	1	-	1	-	2
Female	23	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	18	-	-	-	-	-	-	-	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	1	2
Male	7	-	-	-	-	-	-	-	-	-	1	2
Female	3	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	292	-	-	-	-	-	-	-	-	1	6	17
Male	179	-	-	-	-	-	-	-	-	1	3	15
Female	113	-	-	-	-	-	-	-	-	-	3	2
White, both sexes	287	-	-	-	-	-	-	-	-	1	6	16
Male	175	-	-	-	-	-	-	-	-	1	3	14
Female	112	-	-	-	-	-	-	-	-	-	3	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
United States												
All races, both sexes	5,858	8,085	9,750	10,713	12,728	16,712	20,917	21,094	14,065	5,398	1,154	4
Male	4,402	5,920	6,824	7,037	7,787	9,279	10,419	8,905	4,786	1,333	182	3
Female	1,456	2,165	2,926	3,676	4,941	7,433	10,498	12,189	9,279	4,065	972	1
White, both sexes	4,742	6,656	8,224	9,024	10,821	14,646	18,664	19,173	12,869	4,884	996	2
Male	3,661	4,995	5,880	6,073	6,720	8,250	9,467	8,205	4,431	1,204	158	2
Female	1,081	1,661	2,344	2,951	4,101	6,396	9,197	10,968	8,438	3,680	838	-
Black, both sexes	962	1,249	1,305	1,412	1,588	1,673	1,804	1,531	933	409	133	2
Male	619	790	791	786	867	807	726	531	250	88	18	1
Female	343	459	514	626	721	866	1,078	1,000	683	321	115	1
American Indian 1/, both sexes	52	70	58	60	71	64	69	54	25	8	2	-
Male	38	53	39	39	40	37	38	28	9	1	1	-
Female	14	17	19	21	31	27	31	26	16	7	1	-
Asian or Pacific Islander 2/, both sexes	102	110	163	217	248	329	380	336	238	97	23	-
Male	84	82	114	139	160	185	188	141	96	40	5	-
Female	18	28	49	78	88	144	192	195	142	57	18	-
Alabama												
All races, both sexes	158	216	247	259	285	350	311	328	169	57	14	-
Male	114	163	167	154	182	183	144	112	51	14	1	-
Female	44	53	80	105	103	167	167	216	118	43	13	-
White, both sexes	121	161	177	214	224	283	256	264	132	45	9	-
Male	87	123	122	131	147	151	126	98	41	11	1	-
Female	34	38	55	83	77	132	130	166	91	34	8	-
Black, both sexes	37	55	67	43	61	67	53	62	37	12	5	-
Male	27	40	43	23	35	32	18	13	10	3	-	-
Female	10	15	24	20	26	35	35	49	27	9	5	-
American Indian 1/, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	3	2	-	-	-	2	-	-	-	-
Male	-	-	2	-	-	-	-	1	-	-	-	-
Female	-	-	1	2	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Alaska												
All races, both sexes	4	4	3	8	9	7	8	8	1	1	-	-
Male	4	4	3	5	9	6	2	2	-	-	-	-
Female	-	-	-	3	-	1	6	6	1	1	-	-
White, both sexes	2	3	3	5	4	5	8	6	1	1	-	-
Male	2	3	3	3	4	4	2	2	-	-	-	-
Female	-	-	-	2	-	1	6	4	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	1	5	2	-	2	-	-	-	-
Male	2	-	-	1	5	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	2	-	-	-	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	48	84	110	127	173	245	309	283	173	77	11	1
Male	33	59	67	80	101	140	172	129	60	23	2	1
Female	15	25	43	47	72	105	137	154	113	54	9	-
White, both sexes	46	71	107	118	163	235	291	275	164	72	9	1
Male	32	52	67	76	96	133	164	125	58	21	2	1
Female	14	19	40	42	67	102	127	150	106	51	7	-
Black, both sexes	1	5	1	4	5	5	6	2	4	1	2	-
Male	1	1	-	2	1	3	2	1	-	1	-	-
Female	-	4	1	2	4	2	4	1	4	-	2	-
American Indian 1/, both sexes	1	6	1	3	2	3	11	3	4	4	-	-
Male	-	4	-	2	2	3	5	2	2	1	-	-
Female	1	2	1	1	-	-	6	1	2	3	-	-
Asian or Pacific Islander 2/, both sexes	-	2	1	2	3	2	1	3	1	-	-	-
Male	-	2	-	-	2	1	1	1	-	-	-	-
Female	-	-	1	2	1	1	-	2	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Arkansas												
All races, both sexes	181	229	242	263	281	358	424	374	258	95	25	-
Male	133	158	167	178	179	203	199	148	88	25	5	-
Female	48	71	75	85	102	155	225	226	170	70	20	-
White, both sexes	138	191	216	226	246	313	371	328	231	80	14	-
Male	104	130	153	151	158	183	173	132	80	23	3	-
Female	34	61	63	75	88	130	198	196	151	57	11	-
Black, both sexes	43	36	24	37	32	40	52	46	26	15	10	-
Male	29	26	14	27	19	16	25	16	7	2	2	-
Female	14	10	10	10	13	24	27	30	19	13	8	-
American Indian 1/, both sexes	-	1	1	-	2	2	1	-	-	-	-	-
Male	-	1	-	-	1	1	1	-	-	-	-	-
Female	-	-	1	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	1	3	-	-	1	-	1	-
Male	-	1	-	-	1	3	-	-	1	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	1	-
California												
All races, both sexes	415	577	748	895	1,074	1,515	2,027	2,125	1,479	604	133	-
Male	327	421	515	579	660	839	1,031	949	532	174	15	-
Female	88	156	233	316	414	676	996	1,176	947	430	118	-
White, both sexes	333	449	592	696	838	1,240	1,705	1,809	1,280	511	110	-
Male	259	345	420	472	513	697	883	831	459	141	13	-
Female	74	104	172	224	325	543	822	978	821	370	97	-
Black, both sexes	34	82	85	103	110	112	116	133	66	42	14	-
Male	24	40	49	52	68	51	49	43	19	10	1	-
Female	10	42	36	51	42	61	67	90	47	32	13	-
American Indian 1/, both sexes	7	4	4	8	11	5	6	5	4	-	-	-
Male	6	4	1	4	9	1	5	3	1	-	-	-
Female	1	-	3	4	2	4	1	2	3	-	-	-
Asian or Pacific Islander 2/, both sexes	41	42	67	88	115	158	200	178	129	51	9	-
Male	38	32	45	51	70	90	94	72	53	23	1	-
Female	3	10	22	37	45	68	106	106	76	28	8	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Colorado												
All races, both sexes	69	76	97	97	114	160	202	195	101	38	9	-
Male	54	59	68	67	74	91	94	89	39	8	4	-
Female	15	17	29	30	40	69	108	106	62	30	5	-
White, both sexes	65	69	92	90	110	152	193	188	98	37	9	-
Male	51	53	65	62	71	86	90	84	37	8	4	-
Female	14	16	27	28	39	66	103	104	61	29	5	-
Black, both sexes	3	6	2	5	3	6	4	5	3	-	-	-
Male	2	5	2	3	2	3	1	4	2	-	-	-
Female	1	1	-	2	1	3	3	1	1	-	-	-
American Indian 1/, both sexes	1	-	3	2	1	2	1	1	-	1	-	-
Male	1	-	1	2	1	2	-	1	-	-	-	-
Female	-	-	2	-	-	-	1	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	4	1	-	-	-	-
Male	-	1	-	-	-	-	3	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Connecticut												
All races, both sexes	35	44	68	74	86	151	180	253	165	76	17	1
Male	31	31	42	49	51	78	89	97	53	12	2	1
Female	4	13	26	25	35	73	91	156	112	64	15	-
White, both sexes	29	40	61	66	75	136	171	241	161	74	16	1
Male	25	28	37	43	42	69	86	94	53	12	2	1
Female	4	12	24	23	33	67	85	147	108	62	14	-
Black, both sexes	3	4	7	7	11	11	8	12	3	1	1	-
Male	3	3	5	5	9	6	3	3	-	-	-	-
Female	-	1	2	2	2	5	5	9	3	1	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	1	-	4	1	-	1	1	-	-
Male	2	-	-	1	-	3	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Delaware												
All races, both sexes	18	21	26	23	33	33	50	55	27	8	4	-
Male	15	17	25	18	20	19	27	22	11	1	-	-
Female	3	4	1	5	13	14	23	33	16	7	4	-
White, both sexes	16	17	17	18	28	29	44	52	23	7	3	-
Male	13	15	16	15	17	17	25	22	10	1	-	-
Female	3	2	1	3	11	12	19	30	13	6	3	-
Black, both sexes	2	4	9	5	4	4	6	3	3	1	1	-
Male	2	2	9	3	3	2	2	-	-	-	-	-
Female	-	2	-	2	1	2	4	3	3	1	1	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	9	9	14	18	17	20	22	34	27	12	3	-
Male	7	6	9	9	8	11	7	14	10	4	-	-
Female	2	3	5	9	9	9	15	20	17	8	3	-
White, both sexes	1	2	-	4	3	6	6	14	8	3	1	-
Male	1	2	-	3	2	4	4	7	4	1	-	-
Female	-	-	-	1	1	2	2	7	4	2	1	-
Black, both sexes	8	7	14	14	14	14	15	20	18	9	2	-
Male	6	4	9	6	6	7	2	7	6	3	-	-
Female	2	3	5	8	8	7	13	13	12	6	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Florida												
All races, both sexes	262	361	485	555	724	1,058	1,353	1,466	1,001	362	62	-
Male	182	256	330	365	435	587	715	644	380	114	11	-
Female	80	105	155	190	289	471	638	822	621	248	51	-
White, both sexes	208	300	408	487	622	947	1,242	1,382	944	340	57	-
Male	151	219	287	331	380	535	665	610	370	107	11	-
Female	57	81	121	156	242	412	577	772	574	233	46	-
Black, both sexes	49	56	73	63	95	103	103	77	55	22	5	-
Male	27	33	39	31	50	48	46	32	10	7	-	-
Female	22	23	34	32	45	55	57	45	45	15	5	-
American Indian 1/, both sexes	-	1	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	4	4	5	7	7	8	6	2	-	-	-
Male	4	4	4	3	5	3	4	1	-	-	-	-
Female	1	-	-	2	2	4	4	5	2	-	-	-
Georgia												
All races, both sexes	160	252	258	306	317	361	389	338	241	92	14	-
Male	111	182	173	200	186	198	185	125	86	21	-	-
Female	49	70	85	106	131	163	204	213	155	71	14	-
White, both sexes	98	176	179	225	232	284	312	279	198	73	9	-
Male	65	135	124	154	141	161	156	107	76	19	-	-
Female	33	41	55	71	91	123	156	172	122	54	9	-
Black, both sexes	58	76	79	78	82	72	72	58	41	19	5	-
Male	44	47	49	43	44	34	27	18	8	2	-	-
Female	14	29	30	35	38	38	45	40	33	17	5	-
American Indian 1/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	3	3	5	4	1	2	-	-	-
Male	2	-	-	3	1	3	2	-	2	-	-	-
Female	1	-	-	-	2	2	2	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Hawaii												
All races, both sexes	10	15	35	37	37	50	53	70	41	17	5	-
Male	7	9	30	22	26	28	37	25	18	7	1	-
Female	3	6	5	15	11	22	16	45	23	10	4	-
White, both sexes	1	2	13	13	13	12	12	24	7	1	1	-
Male	-	1	12	10	10	8	7	5	5	-	-	-
Female	1	1	1	3	3	4	5	19	2	1	1	-
Black, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	13	21	24	24	38	41	45	34	16	4	-
Male	6	8	17	12	16	20	30	20	13	7	1	-
Female	2	5	4	12	8	18	11	25	21	9	3	-
Idaho												
All races, both sexes	47	54	75	82	82	81	98	124	55	26	3	-
Male	41	43	62	58	60	43	61	68	22	9	1	-
Female	6	11	13	24	22	38	37	56	33	17	2	-
White, both sexes	46	54	73	79	81	81	95	121	55	26	2	-
Male	40	43	61	55	59	43	58	65	22	9	-	-
Female	6	11	12	24	22	38	37	56	33	17	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	1	1	-	1	1	-	-	1	-
Male	1	-	1	1	1	-	1	1	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	2	-	-	2	2	-	-	-	-
Male	-	-	-	2	-	-	2	2	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Illinois												
All races, both sexes	267	384	430	468	595	710	975	972	708	242	69	1
Male	187	264	284	329	368	386	476	430	233	53	11	-
Female	80	120	146	139	227	324	499	542	475	189	58	1
White, both sexes	213	303	339	360	495	608	863	866	653	220	62	-
Male	160	221	238	260	312	332	433	394	225	46	9	-
Female	53	82	101	100	183	276	430	472	428	174	53	-
Black, both sexes	54	79	83	99	91	97	102	91	46	19	5	1
Male	27	41	42	63	50	51	37	31	6	6	2	-
Female	27	38	41	36	41	46	65	60	40	13	3	1
American Indian 1/, both sexes	-	-	1	-	2	-	-	-	-	-	-	-
Male	-	-	1	-	2	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	7	9	7	5	10	15	9	3	2	-
Male	-	2	3	6	4	3	6	5	2	1	-	-
Female	-	-	4	3	3	2	4	10	7	2	2	-
Indiana												
All races, both sexes	192	245	246	282	314	406	501	452	278	104	14	-
Male	149	176	184	187	198	223	239	189	95	25	1	-
Female	43	69	62	95	116	183	262	263	183	79	13	-
White, both sexes	171	224	226	265	293	379	476	438	271	102	14	-
Male	132	163	173	176	184	207	227	185	93	25	1	-
Female	39	61	53	89	109	172	249	253	178	77	13	-
Black, both sexes	18	21	20	17	20	26	25	14	7	2	-	-
Male	14	13	11	11	13	15	12	4	2	-	-	-
Female	4	8	9	6	7	11	13	10	5	2	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	1	1	-	-	-	-	-	-
Male	2	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Iowa												
All races, both sexes	70	86	108	113	130	207	270	252	219	110	27	-
Male	56	73	80	70	88	117	153	98	84	21	3	-
Female	14	13	28	43	42	90	117	154	135	89	24	-
White, both sexes	68	83	107	111	129	204	267	251	217	108	27	-
Male	54	71	80	69	87	115	150	97	82	21	3	-
Female	14	12	27	42	42	89	117	154	135	87	24	-
Black, both sexes	2	3	1	1	1	1	3	1	2	1	-	-
Male	2	2	-	1	1	1	3	1	2	-	-	-
Female	-	1	1	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
Kansas												
All races, both sexes	37	51	67	82	103	141	167	177	137	47	13	-
Male	28	38	51	56	61	80	99	67	46	11	2	-
Female	9	13	16	26	42	61	68	110	91	36	11	-
White, both sexes	34	49	65	78	96	133	158	168	132	47	13	-
Male	27	36	50	54	58	76	95	64	43	11	2	-
Female	7	13	15	24	38	57	63	104	89	36	11	-
Black, both sexes	3	2	2	1	6	6	4	8	5	-	-	-
Male	1	2	1	-	2	3	2	3	3	-	-	-
Female	2	-	1	1	4	3	2	5	2	-	-	-
American Indian 1/, both sexes	-	-	-	2	1	2	3	-	-	-	-	-
Male	-	-	-	2	1	1	1	-	-	-	-	-
Female	-	-	-	-	-	1	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	2	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Kentucky												
All races, both sexes	206	246	294	289	351	359	417	391	225	100	12	-
Male	166	184	202	190	214	203	211	150	78	29	3	-
Female	40	62	92	99	137	156	206	241	147	71	9	-
White, both sexes	177	218	278	263	327	337	390	371	211	97	12	-
Male	150	166	191	174	204	189	195	142	72	29	3	-
Female	27	52	87	89	123	148	195	229	139	68	9	-
Black, both sexes	27	28	15	26	23	21	25	20	14	3	-	-
Male	16	18	11	16	9	13	14	8	6	-	-	-
Female	11	10	4	10	14	8	11	12	8	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	1	-	1	1	2	-	-	-	-	-
Male	-	-	-	-	1	1	2	-	-	-	-	-
Female	2	-	1	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	172	244	269	247	242	273	304	238	150	44	11	-
Male	129	193	177	159	156	146	143	96	60	8	6	-
Female	43	51	92	88	86	127	161	142	90	36	5	-
White, both sexes	100	148	167	164	168	201	233	181	119	35	6	-
Male	82	117	110	100	112	116	113	71	47	7	3	-
Female	18	31	57	64	56	85	120	110	72	28	3	-
Black, both sexes	71	93	100	81	72	69	70	57	30	9	4	-
Male	46	74	65	57	43	30	30	25	13	1	3	-
Female	25	19	35	24	29	39	40	32	17	8	1	-
American Indian 1/, both sexes	-	1	1	1	1	1	-	-	1	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	1	1	1	2	1	-	-	-	1	-
Male	1	2	1	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	1	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Maine												
All races, both sexes	23	35	56	55	59	73	103	105	76	20	6	-
Male	21	24	45	40	37	45	48	41	31	6	1	-
Female	2	11	11	15	22	28	55	64	45	14	5	-
White, both sexes	22	35	56	55	58	72	102	105	75	19	6	-
Male	20	24	45	40	36	44	48	41	30	5	1	-
Female	2	11	11	15	22	28	54	64	45	14	5	-
Black, both sexes	1	-	-	-	-	1	-	-	-	1	-	-
Male	1	-	-	-	-	1	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	97	132	164	211	239	300	377	397	261	95	16	-
Male	61	90	113	127	131	145	169	157	92	23	3	-
Female	36	42	51	84	108	155	208	240	169	72	13	-
White, both sexes	60	82	103	141	152	234	305	315	221	79	12	-
Male	44	57	74	90	86	116	142	125	82	19	3	-
Female	16	25	29	51	66	118	163	190	139	60	9	-
Black, both sexes	37	50	54	64	82	61	66	73	37	14	4	-
Male	17	33	34	32	41	28	24	26	9	3	-	-
Female	20	17	20	32	41	33	42	47	28	11	4	-
American Indian 1/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	7	6	5	4	5	9	3	2	-	-
Male	-	-	5	5	4	1	2	6	1	1	-	-
Female	-	-	2	1	1	3	3	3	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Massachusetts												
All races, both sexes	78	112	125	161	196	306	403	496	300	161	27	-
Male	63	84	84	101	113	173	188	215	86	36	3	-
Female	15	28	41	60	83	133	215	281	214	125	24	-
White, both sexes	68	108	119	148	183	286	391	483	291	156	27	-
Male	57	81	82	94	106	162	183	209	83	35	3	-
Female	11	27	37	54	77	124	208	274	208	121	24	-
Black, both sexes	4	2	5	8	11	15	7	10	7	4	-	-
Male	2	1	1	3	5	8	3	5	2	1	-	-
Female	2	1	4	5	6	7	4	5	5	3	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	2	1	5	2	5	5	3	2	1	-	-
Male	3	2	1	4	2	3	2	1	1	-	-	-
Female	1	-	-	1	-	2	3	2	1	1	-	-
Michigan												
All races, both sexes	199	251	327	336	418	598	766	777	542	181	43	-
Male	149	175	216	224	246	316	387	323	191	49	6	-
Female	50	76	111	112	172	282	379	454	351	132	37	-
White, both sexes	155	206	281	282	369	532	672	707	489	167	36	-
Male	123	142	194	193	225	281	351	296	173	47	6	-
Female	32	64	87	89	144	251	321	411	316	120	30	-
Black, both sexes	35	37	43	47	45	58	87	64	48	14	7	-
Male	19	28	20	25	18	31	35	24	17	2	-	-
Female	16	9	23	22	27	27	52	40	31	12	7	-
American Indian 1/, both sexes	6	5	2	1	3	5	3	2	2	-	-	-
Male	4	4	1	1	2	2	1	1	-	-	-	-
Female	2	1	1	-	1	3	2	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes	3	3	1	6	1	3	4	4	3	-	-	-
Male	3	1	1	5	1	2	-	2	1	-	-	-
Female	-	2	-	1	-	1	4	2	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Minnesota												
All races, both sexes	44	67	76	105	113	137	177	219	180	68	11	-
Male	40	53	56	64	69	88	92	105	70	21	1	-
Female	4	14	20	41	44	49	85	114	110	47	10	-
White, both sexes	43	63	74	100	111	135	177	217	176	67	11	-
Male	40	50	55	60	67	86	92	104	68	21	1	-
Female	3	13	19	40	44	49	85	113	108	46	10	-
Black, both sexes	1	2	1	2	1	-	-	1	1	1	-	-
Male	-	2	-	2	1	-	-	1	1	-	-	-
Female	1	-	1	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	2	1	2	-	2	-	1	-	-	-	-
Male	-	1	1	1	-	2	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	-	-	3	-	-	-
Male	-	-	-	1	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-
Mississippi												
All races, both sexes	149	157	211	204	212	194	211	177	118	47	8	1
Male	104	113	150	133	134	107	101	59	34	8	2	1
Female	45	44	61	71	78	87	110	118	84	39	6	-
White, both sexes	81	104	139	134	137	144	149	125	91	34	3	-
Male	61	78	104	90	94	78	79	45	22	4	-	-
Female	20	26	35	44	43	66	70	80	69	30	3	-
Black, both sexes	68	53	71	69	74	50	61	52	27	13	5	1
Male	43	35	45	43	40	29	22	14	12	4	2	1
Female	25	18	26	26	34	21	39	38	15	9	3	-
American Indian 1/, both sexes	-	-	1	1	1	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Missouri												
All races, both sexes	194	238	323	353	399	539	655	636	421	159	34	-
Male	142	158	231	227	240	297	323	267	131	36	7	-
Female	52	80	92	126	159	242	332	369	290	123	27	-
White, both sexes	174	205	292	319	350	484	609	598	385	150	30	-
Male	130	143	207	207	212	269	308	254	122	34	6	-
Female	44	62	85	112	138	215	301	344	263	116	24	-
Black, both sexes	19	30	31	33	46	54	43	36	35	9	4	-
Male	12	14	24	20	27	27	14	12	8	2	1	-
Female	7	16	7	13	19	27	29	24	27	7	3	-
American Indian 1/, both sexes	1	1	-	1	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	1	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	-	2	1	2	2	1	-	-	-
Male	-	1	-	-	1	1	1	1	1	-	-	-
Female	-	1	-	-	1	-	1	1	-	-	-	-
Montana												
All races, both sexes	15	18	28	25	30	37	37	42	33	10	-	-
Male	13	13	24	18	21	21	19	20	15	4	-	-
Female	2	5	4	7	9	16	18	22	18	6	-	-
White, both sexes	14	18	24	22	28	37	34	41	33	10	-	-
Male	12	13	21	16	19	21	18	20	15	4	-	-
Female	2	5	3	6	9	16	16	21	18	6	-	-
Black, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	4	2	1	-	2	1	-	-	-	-
Male	-	-	3	1	1	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	2	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Nebraska												
All races, both sexes	30	38	43	46	54	72	91	87	68	29	8	-
Male	24	33	36	26	28	49	52	38	22	5	2	-
Female	6	5	7	20	26	23	39	49	46	24	6	-
White, both sexes	28	35	40	45	50	69	88	84	68	29	8	-
Male	22	30	33	26	26	48	51	36	22	5	2	-
Female	6	5	7	19	24	21	37	48	46	24	6	-
Black, both sexes	1	2	3	1	3	3	1	3	-	-	-	-
Male	1	2	3	-	2	1	-	2	-	-	-	-
Female	-	-	-	1	1	2	1	1	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	-	1	-	-	-	-	-
Male	1	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Nevada												
All races, both sexes	29	35	52	50	58	71	82	68	31	9	2	-
Male	24	27	44	35	35	49	48	34	7	1	1	-
Female	5	8	8	15	23	22	34	34	24	8	1	-
White, both sexes	23	31	50	46	53	64	76	65	29	8	2	-
Male	18	25	42	32	32	44	47	33	7	1	1	-
Female	5	6	8	14	21	20	29	32	22	7	1	-
Black, both sexes	4	3	1	2	3	3	4	2	1	1	-	-
Male	4	1	1	1	2	2	1	-	-	-	-	-
Female	-	2	-	1	1	1	3	2	1	1	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	1	1	2	4	2	1	1	-	-	-
Male	2	1	1	1	1	3	-	1	-	-	-	-
Female	-	-	-	-	1	1	2	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
New Hampshire												
All races, both sexes	12	22	30	29	36	59	80	72	55	11	5	-
Male	12	19	26	19	23	33	44	34	16	5	1	-
Female	-	3	4	10	13	26	36	38	39	6	4	-
White, both sexes	12	21	30	29	36	58	80	72	53	11	5	-
Male	12	18	26	19	23	33	44	34	16	5	1	-
Female	-	3	4	10	13	25	36	38	37	6	4	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	-	-	2	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	2	-	-	-
New Jersey												
All races, both sexes	117	193	209	237	367	499	668	758	487	170	51	-
Male	88	141	141	149	221	272	344	295	153	42	10	-
Female	29	52	68	88	146	227	324	463	334	128	41	-
White, both sexes	91	154	169	184	289	423	588	696	444	157	46	-
Male	67	120	116	120	177	233	306	272	146	40	8	-
Female	24	34	53	64	112	190	282	424	298	117	38	-
Black, both sexes	23	35	32	45	60	68	67	52	37	11	4	-
Male	18	17	18	21	33	34	29	18	6	1	1	-
Female	5	18	14	24	27	34	38	34	31	10	3	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	3	4	8	8	17	8	12	10	5	2	1	-
Male	3	4	7	8	10	5	9	5	1	1	1	-
Female	-	-	1	-	7	3	3	5	4	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
New Mexico												
All races, both sexes	30	47	42	42	55	61	96	85	46	21	4	-
Male	24	36	27	29	41	39	50	42	17	7	-	-
Female	6	11	15	13	14	22	46	43	29	14	4	-
White, both sexes	29	42	40	39	51	59	88	83	44	19	3	-
Male	23	32	25	26	39	38	44	41	16	7	-	-
Female	6	10	15	13	12	21	44	42	28	12	3	-
Black, both sexes	-	2	1	-	-	-	2	-	1	-	-	-
Male	-	1	1	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	3	1	3	4	2	4	2	1	1	1	-
Male	-	3	1	3	2	1	4	1	1	-	-	-
Female	-	-	-	-	2	1	-	1	-	1	1	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	2	-	-	1	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	1	-	-
New York												
All races, both sexes	276	383	477	573	729	1,039	1,480	1,638	1,163	505	113	-
Male	209	270	324	367	413	571	728	656	373	114	15	-
Female	67	113	153	206	316	468	752	982	790	391	98	-
White, both sexes	225	308	397	474	610	894	1,301	1,463	1,063	468	98	-
Male	180	228	279	319	351	496	662	602	343	105	12	-
Female	45	80	118	155	259	398	639	861	720	363	86	-
Black, both sexes	43	63	68	86	99	114	159	147	82	28	11	-
Male	24	36	37	40	50	55	60	40	24	6	2	-
Female	19	27	31	46	49	59	99	107	58	22	9	-
American Indian 1/, both sexes	2	2	3	4	1	4	1	1	2	1	-	-
Male	1	-	3	2	-	4	-	-	-	-	-	-
Female	1	2	-	2	1	-	1	1	2	1	-	-
Asian or Pacific Islander 2/, both sexes	6	10	9	9	19	27	19	27	16	8	4	-
Male	4	6	5	6	12	16	6	14	6	3	1	-
Female	2	4	4	3	7	11	13	13	10	5	3	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
North Carolina												
All races, both sexes	160	251	284	336	377	463	607	537	316	127	26	-
Male	117	186	194	228	225	260	275	228	101	26	2	-
Female	43	65	90	108	152	203	332	309	215	101	24	-
White, both sexes	111	184	227	253	298	382	503	450	269	103	22	-
Male	82	137	161	184	177	224	242	197	91	20	2	-
Female	29	47	66	69	121	158	261	253	178	83	20	-
Black, both sexes	44	61	52	77	74	76	100	83	47	22	4	-
Male	30	46	29	40	46	34	32	31	10	5	-	-
Female	14	15	23	37	28	42	68	52	37	17	4	-
American Indian 1/, both sexes	4	5	4	4	4	4	4	4	-	-	-	-
Male	4	3	3	3	1	2	1	-	-	-	-	-
Female	-	2	1	1	3	2	3	4	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	2	1	1	-	-	-	2	-	-
Male	1	-	1	1	1	-	-	-	-	1	-	-
Female	-	1	-	1	-	1	-	-	-	1	-	-
North Dakota												
All races, both sexes	9	21	23	24	32	48	74	86	53	22	-	-
Male	9	15	17	17	25	31	45	43	23	9	-	-
Female	-	6	6	7	7	17	29	43	30	13	-	-
White, both sexes	8	19	20	24	32	47	73	86	52	21	-	-
Male	8	14	15	17	25	31	44	43	22	8	-	-
Female	-	5	5	7	7	16	29	43	30	13	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	2	3	-	-	1	1	-	1	-	-	-
Male	1	1	2	-	-	-	1	-	1	-	-	-
Female	-	1	1	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Ohio												
All races, both sexes	244	347	404	439	522	712	911	952	603	198	44	-
Male	182	257	295	291	315	388	435	400	173	46	8	-
Female	62	90	109	148	207	324	476	552	430	152	36	-
White, both sexes	210	310	359	395	464	644	842	905	567	189	42	-
Male	158	238	269	268	286	354	400	380	163	45	8	-
Female	52	72	90	127	178	290	442	525	404	144	34	-
Black, both sexes	33	36	43	41	55	65	67	46	36	9	2	-
Male	23	18	25	21	26	31	33	19	10	1	-	-
Female	10	18	18	20	29	34	34	27	26	8	2	-
American Indian 1/, both sexes	-	-	2	1	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	2	3	3	2	1	-	-	-	-
Male	1	1	-	2	3	3	2	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	70	95	135	154	164	194	234	229	135	60	16	-
Male	55	62	86	97	95	117	114	95	54	15	1	-
Female	15	33	49	57	69	77	120	134	81	45	15	-
White, both sexes	57	78	118	136	145	169	208	209	125	54	15	-
Male	46	48	76	84	88	103	101	85	49	13	1	-
Female	11	30	42	52	57	66	107	124	76	41	14	-
Black, both sexes	5	5	7	7	8	11	12	4	4	6	1	-
Male	4	3	4	5	4	5	5	1	2	2	-	-
Female	1	2	3	2	4	6	7	3	2	4	1	-
American Indian 1/, both sexes	8	12	9	11	9	14	13	14	6	-	-	-
Male	5	11	6	8	3	9	8	9	3	-	-	-
Female	3	1	3	3	6	5	5	5	3	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	2	-	1	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	2	-	1	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Oregon												
All races, both sexes	48	72	92	122	129	155	198	217	154	53	12	-
Male	35	49	67	90	79	95	114	101	65	12	3	-
Female	13	23	25	32	50	60	84	116	89	41	9	-
White, both sexes	42	69	88	119	125	151	194	216	153	53	12	-
Male	31	47	63	87	78	92	112	100	64	12	3	-
Female	11	22	25	32	47	59	82	116	89	41	9	-
Black, both sexes	1	3	1	2	1	3	1	-	1	-	-	-
Male	1	2	1	2	1	3	-	-	1	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	3	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	2	-	-	-	3	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	3	1	-	1	3	-	-	-	-	-
Male	1	-	3	1	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Pennsylvania												
All races, both sexes	284	428	513	555	680	992	1,266	1,276	857	294	53	-
Male	217	333	384	367	417	565	622	547	286	67	9	-
Female	67	95	129	188	263	427	644	729	571	227	44	-
White, both sexes	249	380	469	505	618	907	1,192	1,225	817	273	48	-
Male	194	304	359	341	380	531	595	528	279	61	9	-
Female	55	76	110	164	238	376	597	697	538	212	39	-
Black, both sexes	32	43	43	47	56	81	66	45	37	20	5	-
Male	21	25	25	24	33	32	25	16	5	6	-	-
Female	11	18	18	23	23	49	41	29	32	14	5	-
American Indian 1/, both sexes	-	2	-	-	-	-	1	1	-	-	-	-
Male	-	2	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	3	1	3	6	4	7	5	3	1	-	-
Male	2	2	-	2	4	2	1	2	2	-	-	-
Female	1	1	1	1	2	2	6	3	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Rhode Island												
All races, both sexes	18	30	34	47	62	86	118	143	96	35	8	-
Male	14	21	21	34	39	46	59	53	35	10	-	-
Female	4	9	13	13	23	40	59	90	61	25	8	-
White, both sexes	18	26	30	45	59	84	115	142	96	35	8	-
Male	14	19	17	33	37	46	58	53	35	10	-	-
Female	4	7	13	12	22	38	57	89	61	25	8	-
Black, both sexes	-	2	3	1	3	2	3	1	-	-	-	-
Male	-	2	3	-	2	-	1	-	-	-	-	-
Female	-	-	-	1	1	2	2	1	-	-	-	-
American Indian 1/, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	133	181	236	257	275	300	309	246	150	48	7	-
Male	95	131	175	171	167	159	155	105	51	9	-	-
Female	38	50	61	86	108	141	154	141	99	39	7	-
White, both sexes	88	127	167	182	206	227	239	191	127	30	5	-
Male	66	90	132	130	124	131	120	91	46	7	-	-
Female	22	37	35	52	82	96	119	100	81	23	5	-
Black, both sexes	43	53	67	73	69	72	70	55	23	18	2	-
Male	27	40	42	39	43	27	35	14	5	2	-	-
Female	16	13	25	34	26	45	35	41	18	16	2	-
American Indian 1/, both sexes	1	1	2	-	-	-	-	-	-	-	-	-
Male	1	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	2	-	1	-	-	-	-	-	-
Male	1	-	-	2	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
South Dakota												
All races, both sexes	18	32	29	36	54	69	85	74	88	28	9	-
Male	17	28	22	26	31	43	55	31	35	8	2	-
Female	1	4	7	10	23	26	30	43	53	20	7	-
White, both sexes	17	28	25	34	51	65	81	72	87	28	9	-
Male	16	25	19	24	30	41	52	31	35	8	2	-
Female	1	3	6	10	21	24	29	41	52	20	7	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	4	3	2	3	4	3	2	1	-	-	-
Male	1	3	2	2	1	2	3	-	-	-	-	-
Female	-	1	1	-	2	2	-	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Tennessee												
All races, both sexes	219	303	347	374	469	537	583	593	396	139	34	-
Male	155	224	248	245	280	304	273	216	116	25	5	-
Female	64	79	99	129	189	233	310	377	280	114	29	-
White, both sexes	191	254	301	326	406	480	514	532	357	120	27	-
Male	139	190	219	220	250	275	248	195	103	22	4	-
Female	52	64	82	106	156	205	266	337	254	98	23	-
Black, both sexes	27	48	45	46	61	52	69	60	39	19	7	-
Male	15	34	28	24	30	25	25	21	13	3	1	-
Female	12	14	17	22	31	27	44	39	26	16	6	-
American Indian 1/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	2	1	4	-	1	-	-	-	-
Male	1	-	1	1	-	3	-	-	-	-	-	-
Female	-	1	-	1	1	1	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Texas												
All races, both sexes	541	758	890	906	1,071	1,398	1,664	1,507	936	404	82	-
Male	409	530	586	576	644	783	823	679	314	92	17	-
Female	132	228	304	330	427	615	841	828	622	312	65	-
White, both sexes	457	642	784	781	930	1,247	1,488	1,393	856	354	66	-
Male	351	460	516	511	567	708	751	634	290	79	13	-
Female	106	182	268	270	363	539	737	759	566	275	53	-
Black, both sexes	79	107	91	108	127	133	162	105	77	47	15	-
Male	55	63	59	57	64	68	66	41	21	11	3	-
Female	24	44	32	51	63	65	96	64	56	36	12	-
American Indian 1/, both sexes	1	2	1	4	4	1	1	2	-	-	-	-
Male	1	2	1	1	4	-	1	2	-	-	-	-
Female	-	-	-	3	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	7	14	13	10	17	13	7	3	3	1	-
Male	2	5	10	7	9	7	5	2	3	2	1	-
Female	2	2	4	6	1	10	8	5	-	1	-	-
Utah												
All races, both sexes	17	34	43	45	48	60	95	75	44	19	3	-
Male	12	29	34	35	32	31	49	34	22	5	1	-
Female	5	5	9	10	16	29	46	41	22	14	2	-
White, both sexes	16	32	39	45	46	58	94	73	43	18	3	-
Male	12	27	31	35	31	30	48	32	22	5	1	-
Female	4	5	8	10	15	28	46	41	21	13	2	-
Black, both sexes	-	-	2	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	1	2	1	-	1	-	-	-
Male	-	1	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	2	-	1	-	-	1	-	1	-	-
Male	-	1	2	-	1	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Vermont												
All races, both sexes	5	7	10	24	13	28	40	37	28	11	2	-
Male	4	6	10	14	8	15	16	14	7	1	-	-
Female	1	1	-	10	5	13	24	23	21	10	2	-
White, both sexes	5	7	10	24	13	28	39	37	28	11	2	-
Male	4	6	10	14	8	15	16	14	7	1	-	-
Female	1	1	-	10	5	13	23	23	21	10	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Virginia												
All races, both sexes	151	211	233	258	286	387	453	414	266	98	19	-
Male	112	168	175	168	186	212	212	179	89	29	3	-
Female	39	43	58	90	100	175	241	235	177	69	16	-
White, both sexes	115	163	185	194	221	308	363	338	238	82	13	-
Male	91	137	140	127	153	173	172	142	79	26	3	-
Female	24	26	45	67	68	135	191	196	159	56	10	-
Black, both sexes	36	47	46	58	61	73	82	72	25	15	6	-
Male	21	30	33	36	31	36	37	35	8	3	-	-
Female	15	17	13	22	30	37	45	37	17	12	6	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	6	4	6	8	4	3	1	-	-
Male	-	-	2	5	2	3	3	2	2	-	-	-
Female	-	-	-	1	2	3	5	2	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Washington												
All races, both sexes	105	123	161	161	206	302	332	374	261	99	26	-
Male	85	100	122	118	136	164	163	165	92	29	4	-
Female	20	23	39	43	70	138	169	209	169	70	22	-
White, both sexes	92	110	146	149	191	276	314	359	244	96	25	-
Male	75	88	112	108	128	150	153	160	85	29	4	-
Female	17	22	34	41	63	126	161	199	159	67	21	-
Black, both sexes	3	4	6	4	7	14	5	2	5	-	1	-
Male	3	4	3	4	4	7	3	-	2	-	-	-
Female	-	-	3	-	3	7	2	2	3	-	1	-
American Indian 1/, both sexes	4	6	6	3	5	3	1	8	1	-	-	-
Male	3	5	5	1	1	2	-	4	1	-	-	-
Female	1	1	1	2	4	1	1	4	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	3	3	5	3	9	12	5	11	3	-	-
Male	4	3	2	5	3	5	7	1	4	-	-	-
Female	2	-	1	-	-	4	5	4	7	3	-	-
West Virginia												
All races, both sexes	48	64	87	97	112	145	198	149	96	37	9	-
Male	33	47	59	61	73	67	96	57	30	7	-	-
Female	15	17	28	36	39	78	102	92	66	30	9	-
White, both sexes	46	63	85	91	109	140	196	148	95	37	8	-
Male	32	46	57	57	71	63	95	57	30	7	-	-
Female	14	17	28	34	38	77	101	91	65	30	8	-
Black, both sexes	2	1	2	3	3	5	2	1	1	-	1	-
Male	1	1	2	2	2	4	1	-	-	-	-	-
Female	1	-	-	1	1	1	1	1	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	3	-	-	-	-	-	-	-	-
Male	-	-	-	2	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Wisconsin												
All races, both sexes	115	170	220	203	264	334	424	454	332	121	29	-
Male	88	142	154	148	186	193	218	201	102	24	7	-
Female	27	28	66	55	78	141	206	253	230	97	22	-
White, both sexes	108	161	213	198	252	327	416	446	330	120	29	-
Male	83	136	150	145	177	189	215	199	102	24	7	-
Female	25	25	63	53	75	138	201	247	228	96	22	-
Black, both sexes	6	3	4	3	9	4	4	6	2	1	-	-
Male	4	1	2	1	7	3	1	1	-	-	-	-
Female	2	2	2	2	2	1	3	5	2	1	-	-
American Indian 1/, both sexes	1	5	2	2	1	1	3	2	-	-	-	-
Male	1	4	1	2	1	-	2	1	-	-	-	-
Female	-	1	1	-	-	1	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	2	2	1	-	-	-	-	-
Male	-	1	1	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	1	1	1	-	-	-	-	-
Wyoming												
All races, both sexes	20	32	24	23	32	32	40	39	18	7	1	-
Male	15	20	22	17	21	20	18	17	7	3	-	-
Female	5	12	2	6	11	12	22	22	11	4	1	-
White, both sexes	20	31	24	23	31	30	40	39	18	7	1	-
Male	15	19	22	17	20	19	18	17	7	3	-	-
Female	5	12	2	6	11	11	22	22	11	4	1	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
United States												
All races, both sexes	4,092	2	-	-	-	1	2	6	11	32	77	148
Male	2,159	1	-	-	-	1	1	5	9	21	55	100
Female	1,933	1	-	-	-	-	1	1	2	11	22	48
White, both sexes	3,441	2	-	-	-	-	1	4	4	19	59	120
Male	1,792	1	-	-	-	-	1	3	4	16	43	83
Female	1,649	1	-	-	-	-	-	1	-	3	16	37
Black, both sexes	539	-	-	-	-	1	1	2	5	11	13	22
Male	303	-	-	-	-	1	-	2	3	5	10	14
Female	236	-	-	-	-	-	1	-	2	6	3	8
American Indian 1/, both sexes	61	-	-	-	-	-	-	-	1	2	4	4
Male	34	-	-	-	-	-	-	-	1	-	1	1
Female	27	-	-	-	-	-	-	-	-	2	3	3
Asian or Pacific Islander 2/, both sexes	51	-	-	-	-	-	-	-	1	-	1	2
Male	30	-	-	-	-	-	-	-	1	-	1	2
Female	21	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	27	-	-	-	-	-	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	20	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Alaska												
All races, both sexes	26	-	-	-	-	-	-	-	-	-	1	2
Male	20	-	-	-	-	-	-	-	-	-	1	2
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	1	1
Male	14	-	-	-	-	-	-	-	-	-	1	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	31	-	-	-	-	-	-	-	-	-	-	2
Male	21	-	-	-	-	-	-	-	-	-	-	2
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	29	-	-	-	-	-	-	-	-	-	-	2
Male	21	-	-	-	-	-	-	-	-	-	-	2
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Arkansas												
All races, both sexes	15	-	-	-	-	-	-	-	1	1	1	1
Male	9	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	1	1	-	1
White, both sexes	9	-	-	-	-	-	-	-	-	1	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	6	-	-	-	-	-	-	-	1	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	171	-	-	-	-	-	-	-	-	-	4	7
Male	103	-	-	-	-	-	-	-	-	-	4	2
Female	68	-	-	-	-	-	-	-	-	-	-	5
White, both sexes	151	-	-	-	-	-	-	-	-	-	4	6
Male	93	-	-	-	-	-	-	-	-	-	4	2
Female	58	-	-	-	-	-	-	-	-	-	-	4
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Colorado												
All races, both sexes	38	-	-	-	-	-	-	-	-	-	2	1
Male	19	-	-	-	-	-	-	-	-	-	1	1
Female	19	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	37	-	-	-	-	-	-	-	-	-	2	1
Male	18	-	-	-	-	-	-	-	-	-	1	1
Female	19	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	27	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	27	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Delaware												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	2
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Florida												
All races, both sexes	124	-	-	-	-	1	1	1	-	-	1	1
Male	64	-	-	-	-	1	1	1	-	-	-	1
Female	60	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	114	-	-	-	-	-	1	1	-	-	1	1
Male	58	-	-	-	-	-	1	1	-	-	-	1
Female	56	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	10	-	-	-	-	1	-	-	-	-	-	-
Male	6	-	-	-	-	1	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	75	-	-	-	-	-	-	-	1	1	-	1
Male	41	-	-	-	-	-	-	-	1	1	-	1
Female	34	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	52	-	-	-	-	-	-	-	-	1	-	-
Male	28	-	-	-	-	-	-	-	-	1	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	22	-	-	-	-	-	-	-	1	-	-	1
Male	12	-	-	-	-	-	-	-	1	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Hawaii												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Illinois												
All races, both sexes	116	-	-	-	-	-	-	-	-	-	-	1
Male	60	-	-	-	-	-	-	-	-	-	-	-
Female	56	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	103	-	-	-	-	-	-	-	-	-	-	1
Male	53	-	-	-	-	-	-	-	-	-	-	-
Female	50	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	55	-	-	-	-	-	-	-	-	1	1	2
Male	29	-	-	-	-	-	-	-	-	1	-	2
Female	26	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	49	-	-	-	-	-	-	-	-	1	1	2
Male	28	-	-	-	-	-	-	-	-	1	-	2
Female	21	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Iowa												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Kentucky												
All races, both sexes	20	-	-	-	-	-	-	-	-	1	2	1
Male	13	-	-	-	-	-	-	-	-	1	2	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	18	-	-	-	-	-	-	-	-	1	2	1
Male	11	-	-	-	-	-	-	-	-	1	2	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	75	-	-	-	-	-	-	-	-	-	-	7
Male	37	-	-	-	-	-	-	-	-	-	-	3
Female	38	-	-	-	-	-	-	-	-	-	-	4
White, both sexes	48	-	-	-	-	-	-	-	-	-	-	3
Male	23	-	-	-	-	-	-	-	-	-	-	2
Female	25	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	26	-	-	-	-	-	-	-	-	-	-	3
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Maine												
All races, both sexes	29	-	-	-	-	-	-	-	-	-	1	1
Male	19	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	29	-	-	-	-	-	-	-	-	-	1	1
Male	19	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	55	-	-	-	-	-	-	-	-	-	-	1
Male	30	-	-	-	-	-	-	-	-	-	-	1
Female	25	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	37	-	-	-	-	-	-	-	-	-	-	1
Male	22	-	-	-	-	-	-	-	-	-	-	1
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Massachusetts												
All races, both sexes	63	-	-	-	-	-	-	-	1	-	1	3
Male	34	-	-	-	-	-	-	-	1	-	-	2
Female	29	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	59	-	-	-	-	-	-	-	1	-	1	2
Male	31	-	-	-	-	-	-	-	1	-	-	1
Female	28	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	69	-	-	-	-	-	-	-	-	2	1	3
Male	38	-	-	-	-	-	-	-	-	1	1	-
Female	31	-	-	-	-	-	-	-	-	1	-	3
White, both sexes	53	-	-	-	-	-	-	-	-	1	1	2
Male	29	-	-	-	-	-	-	-	-	1	1	-
Female	24	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	16	-	-	-	-	-	-	-	-	1	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Minnesota												
All races, both sexes	39	-	-	-	-	-	-	-	-	-	-	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	38	-	-	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	21	-	-	-	-	-	-	-	-	1	-	2
Male	11	-	-	-	-	-	-	-	-	-	-	2
Female	10	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	1	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Missouri												
All races, both sexes	68	-	-	-	-	-	-	-	-	-	-	4
Male	25	-	-	-	-	-	-	-	-	-	-	4
Female	43	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	60	-	-	-	-	-	-	-	-	-	-	4
Male	21	-	-	-	-	-	-	-	-	-	-	4
Female	39	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Nebraska												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
New Hampshire												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	106	-	-	-	-	-	-	-	-	1	1	-
Male	40	-	-	-	-	-	-	-	-	1	1	-
Female	66	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	98	-	-	-	-	-	-	-	-	1	1	-
Male	36	-	-	-	-	-	-	-	-	1	1	-
Female	62	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
New Mexico												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	168	1	-	-	-	-	-	1	-	-	1	4
Male	71	-	-	-	-	-	-	1	-	-	-	2
Female	97	1	-	-	-	-	-	-	-	-	1	2
White, both sexes	152	1	-	-	-	-	-	1	-	-	1	3
Male	62	-	-	-	-	-	-	1	-	-	-	1
Female	90	1	-	-	-	-	-	-	-	-	1	2
Black, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
North Carolina												
All races, both sexes	78	-	-	-	-	-	-	-	1	1	-	2
Male	51	-	-	-	-	-	-	-	1	1	-	2
Female	27	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	65	-	-	-	-	-	-	-	-	-	-	2
Male	44	-	-	-	-	-	-	-	-	-	-	2
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	11	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Ohio												
All races, both sexes	150	1	-	-	-	-	-	-	1	2	2	10
Male	90	1	-	-	-	-	-	-	1	2	1	6
Female	60	-	-	-	-	-	-	-	-	-	1	4
White, both sexes	145	1	-	-	-	-	-	-	1	2	2	10
Male	86	1	-	-	-	-	-	-	1	2	1	6
Female	59	-	-	-	-	-	-	-	-	-	1	4
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	998	-	-	-	-	-	-	1	1	3	22	46
Male	523	-	-	-	-	-	-	1	1	-	16	31
Female	475	-	-	-	-	-	-	-	-	3	6	15
White, both sexes	893	-	-	-	-	-	-	-	-	1	20	43
Male	461	-	-	-	-	-	-	-	-	-	16	30
Female	432	-	-	-	-	-	-	-	-	1	4	13
Black, both sexes	57	-	-	-	-	-	-	1	-	1	-	2
Male	34	-	-	-	-	-	-	1	-	-	-	1
Female	23	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	42	-	-	-	-	-	-	-	1	1	2	1
Male	25	-	-	-	-	-	-	-	1	-	-	-
Female	17	-	-	-	-	-	-	-	-	1	2	1
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Oregon												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	174	-	-	-	-	-	-	-	-	2	-	6
Male	75	-	-	-	-	-	-	-	-	2	-	4
Female	99	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	165	-	-	-	-	-	-	-	-	2	-	6
Male	74	-	-	-	-	-	-	-	-	2	-	4
Female	91	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Rhode Island												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	189	-	-	-	-	-	-	2	2	1	7	7
Male	108	-	-	-	-	-	-	2	1	1	5	5
Female	81	-	-	-	-	-	-	-	1	-	2	2
White, both sexes	102	-	-	-	-	-	-	1	-	-	3	5
Male	58	-	-	-	-	-	-	1	-	-	2	4
Female	44	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	84	-	-	-	-	-	-	1	2	1	4	1
Male	49	-	-	-	-	-	-	1	1	1	3	-
Female	35	-	-	-	-	-	-	-	1	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
South Dakota												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	2	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	2	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	54	-	-	-	-	-	-	-	-	-	1	1
Male	29	-	-	-	-	-	-	-	-	-	1	1
Female	25	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	43	-	-	-	-	-	-	-	-	-	-	1
Male	23	-	-	-	-	-	-	-	-	-	-	1
Female	20	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	10	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Texas												
All races, both sexes	182	-	-	-	-	-	-	-	-	2	5	1
Male	85	-	-	-	-	-	-	-	-	-	2	1
Female	97	-	-	-	-	-	-	-	-	2	3	-
White, both sexes	154	-	-	-	-	-	-	-	-	1	2	1
Male	71	-	-	-	-	-	-	-	-	-	-	1
Female	83	-	-	-	-	-	-	-	-	1	2	-
Black, both sexes	24	-	-	-	-	-	-	-	-	-	3	-
Male	13	-	-	-	-	-	-	-	-	-	2	-
Female	11	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Vermont												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	557	-	-	-	-	-	1	1	3	12	19	25
Male	318	-	-	-	-	-	-	-	3	9	15	18
Female	239	-	-	-	-	-	1	1	-	3	4	7
White, both sexes	387	-	-	-	-	-	-	1	2	6	15	14
Male	216	-	-	-	-	-	-	-	2	6	12	10
Female	171	-	-	-	-	-	-	1	-	-	3	4
Black, both sexes	163	-	-	-	-	-	1	-	1	6	4	10
Male	97	-	-	-	-	-	-	-	1	3	3	7
Female	66	-	-	-	-	-	1	-	-	3	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Washington												
All races, both sexes	30	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	22	-	-	-	-	-	-	-	-	1	-	-
Male	13	-	-	-	-	-	-	-	-	1	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	1	-	-
Male	13	-	-	-	-	-	-	-	-	1	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Wisconsin												
All races, both sexes	40	-	-	-	-	-	-	-	-	-	-	1
Male	21	-	-	-	-	-	-	-	-	-	-	1
Female	19	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	38	-	-	-	-	-	-	-	-	-	-	1
Male	19	-	-	-	-	-	-	-	-	-	-	1
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
United States												
All races, both sexes	228	320	359	361	379	472	549	542	396	164	43	-
Male	168	240	247	245	234	238	262	197	99	25	11	-
Female	60	80	112	116	145	234	287	345	297	139	32	-
White, both sexes	166	247	290	292	312	398	493	488	363	146	37	-
Male	123	187	197	202	194	198	239	179	93	21	8	-
Female	43	60	93	90	118	200	254	309	270	125	29	-
Black, both sexes	53	62	55	56	57	65	49	44	26	11	6	-
Male	38	44	40	34	35	35	18	14	4	3	3	-
Female	15	18	15	22	22	30	31	30	22	8	3	-
American Indian 1/, both sexes	5	8	6	7	5	4	3	4	3	5	-	-
Male	5	7	4	3	3	4	2	3	-	-	-	-
Female	-	1	2	4	2	-	1	1	3	5	-	-
Asian or Pacific Islander 2/, both sexes	4	3	8	6	5	5	4	6	4	2	-	-
Male	2	2	6	6	2	1	3	1	2	1	-	-
Female	2	1	2	-	3	4	1	5	2	1	-	-
Alabama												
All races, both sexes	2	2	2	2	4	2	6	5	-	1	-	-
Male	1	1	-	1	2	2	5	2	-	-	-	-
Female	1	1	2	1	2	-	1	3	-	1	-	-
White, both sexes	1	2	2	2	2	1	6	2	-	1	-	-
Male	1	1	-	1	1	1	5	-	-	-	-	-
Female	-	1	2	1	1	-	1	2	-	1	-	-
Black, both sexes	1	-	-	-	2	1	-	3	-	-	-	-
Male	-	-	-	-	1	1	-	2	-	-	-	-
Female	1	-	-	-	1	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Alaska												
All races, both sexes	1	4	7	3	-	4	1	3	-	-	-	-
Male	1	3	6	-	-	4	1	2	-	-	-	-
Female	-	1	1	3	-	-	-	1	-	-	-	-
White, both sexes	1	3	5	2	-	3	1	2	-	-	-	-
Male	1	2	4	-	-	3	1	1	-	-	-	-
Female	-	1	1	2	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	1	1	-	1	-	1	-	-	-	-
Male	-	1	1	-	-	1	-	1	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	2	2	3	4	1	6	4	4	3	-	-
Male	-	2	1	2	4	-	5	2	3	-	-	-
Female	-	-	1	1	-	1	1	2	1	3	-	-
White, both sexes	-	2	2	3	4	1	6	4	4	1	-	-
Male	-	2	1	2	4	-	5	2	3	-	-	-
Female	-	-	1	1	-	1	1	2	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	2	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	2	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Arkansas												
All races, both sexes	2	-	-	2	2	-	4	1	-	-	-	-
Male	1	-	-	2	2	-	3	-	-	-	-	-
Female	1	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	-	-	2	2	-	2	1	-	-	-	-
Male	-	-	-	2	2	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	2	-	-	-	-	-	2	-	-	-	-	-
Male	1	-	-	-	-	-	2	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	7	10	16	22	14	17	19	21	22	9	3	-
Male	6	9	11	18	9	13	13	8	9	1	-	-
Female	1	1	5	4	5	4	6	13	13	8	3	-
White, both sexes	4	10	14	18	13	14	18	20	19	8	3	-
Male	4	9	10	15	8	11	13	8	8	1	-	-
Female	-	1	4	3	5	3	5	12	11	7	3	-
Black, both sexes	1	-	-	1	-	2	1	-	-	-	-	-
Male	1	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	2	2	1	-	-	1	3	1	-	-
Male	1	-	1	2	1	-	-	-	1	-	-	-
Female	1	-	1	-	-	-	-	1	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Colorado												
All races, both sexes	2	2	-	6	3	4	8	4	3	3	-	-
Male	2	1	-	4	1	-	3	2	3	1	-	-
Female	-	1	-	2	2	4	5	2	-	2	-	-
White, both sexes	2	2	-	6	3	4	7	4	3	3	-	-
Male	2	1	-	4	1	-	2	2	3	1	-	-
Female	-	1	-	2	2	4	5	2	-	2	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	1	3	-	1	2	3	9	5	3	-	-
Male	-	1	2	-	1	1	-	3	3	-	-	-
Female	-	-	1	-	-	1	3	6	2	3	-	-
White, both sexes	-	1	3	-	1	2	3	9	5	3	-	-
Male	-	1	2	-	1	1	-	3	3	-	-	-
Female	-	-	1	-	-	1	3	6	2	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Delaware												
All races, both sexes	1	-	1	1	-	-	2	-	1	-	-	-
Male	1	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	1	-	-	-
White, both sexes	1	-	1	1	-	-	2	-	1	-	-	-
Male	1	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Florida												
All races, both sexes	7	4	11	9	11	20	18	20	14	5	-	-
Male	3	3	6	8	8	11	7	8	6	-	-	-
Female	4	1	5	1	3	9	11	12	8	5	-	-
White, both sexes	6	2	9	7	11	19	17	20	14	5	-	-
Male	3	2	5	7	8	10	6	8	6	-	-	-
Female	3	-	4	-	3	9	11	12	8	5	-	-
Black, both sexes	1	2	2	2	-	1	1	-	-	-	-	-
Male	-	1	1	1	-	1	1	-	-	-	-	-
Female	1	1	1	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	8	5	5	5	11	7	14	5	11	1	-	-
Male	6	4	1	3	8	2	8	2	3	1	-	-
Female	2	1	4	2	3	5	6	3	8	-	-	-
White, both sexes	5	3	2	4	11	4	10	4	8	-	-	-
Male	3	2	1	2	8	1	6	1	3	-	-	-
Female	2	1	1	2	3	3	4	3	5	-	-	-
Black, both sexes	3	2	3	1	-	3	3	1	3	1	-	-
Male	3	2	-	1	-	1	1	1	-	1	-	-
Female	-	-	3	-	-	2	2	-	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Hawaii												
All races, both sexes	-	-	-	1	-	-	1	2	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	1	2	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Idaho												
All races, both sexes	1	-	-	2	2	-	-	2	3	1	-	-
Male	-	-	-	2	2	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	2	3	1	-	-
White, both sexes	1	-	-	2	2	-	-	2	3	1	-	-
Male	-	-	-	2	2	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	2	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Illinois												
All races, both sexes	5	6	11	9	9	17	16	20	14	5	3	-
Male	5	4	6	8	5	9	7	9	4	1	2	-
Female	-	2	5	1	4	8	9	11	10	4	1	-
White, both sexes	4	3	11	8	6	15	15	18	14	5	3	-
Male	4	2	6	7	4	8	7	8	4	1	2	-
Female	-	1	5	1	2	7	8	10	10	4	1	-
Black, both sexes	1	3	-	1	2	2	1	1	-	-	-	-
Male	1	2	-	1	1	1	-	1	-	-	-	-
Female	-	1	-	-	1	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
Indiana												
All races, both sexes	4	1	5	1	4	4	15	7	8	1	1	-
Male	2	1	5	1	4	2	7	3	1	-	-	-
Female	2	-	-	-	-	2	8	4	7	1	1	-
White, both sexes	3	1	5	1	4	4	12	7	7	-	1	-
Male	2	1	5	1	4	2	6	3	1	-	-	-
Female	1	-	-	-	-	2	6	4	6	-	1	-
Black, both sexes	1	-	-	-	-	-	3	-	1	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	2	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Iowa												
All races, both sexes	1	3	1	1	2	2	8	2	3	-	-	-
Male	1	2	1	1	-	2	4	-	1	-	-	-
Female	-	1	-	-	2	-	4	2	2	-	-	-
White, both sexes	1	3	1	1	2	2	8	2	3	-	-	-
Male	1	2	1	1	-	2	4	-	1	-	-	-
Female	-	1	-	-	2	-	4	2	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	2	-	2	-	3	3	4	4	-	-	-
Male	1	1	-	2	-	2	1	1	-	-	-	-
Female	-	1	-	-	-	1	2	3	4	-	-	-
White, both sexes	1	2	-	2	-	3	3	4	3	-	-	-
Male	1	1	-	2	-	2	1	1	-	-	-	-
Female	-	1	-	-	-	1	2	3	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Kentucky												
All races, both sexes	-	2	1	-	3	2	3	3	1	1	-	-
Male	-	2	1	-	3	-	1	1	1	-	-	-
Female	-	-	-	-	-	2	2	2	-	1	-	-
White, both sexes	-	1	1	-	2	2	3	3	1	1	-	-
Male	-	1	1	-	2	-	1	1	1	-	-	-
Female	-	-	-	-	-	2	2	2	-	1	-	-
Black, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	2	9	8	9	7	5	11	11	3	2	1	-
Male	2	7	6	5	4	3	4	3	-	-	-	-
Female	-	2	2	4	3	2	7	8	3	2	1	-
White, both sexes	-	4	6	6	3	3	8	9	3	2	1	-
Male	-	4	5	3	2	1	4	2	-	-	-	-
Female	-	-	1	3	1	2	4	7	3	2	1	-
Black, both sexes	2	5	2	3	4	2	3	2	-	-	-	-
Male	2	3	1	2	2	2	-	1	-	-	-	-
Female	-	2	1	1	2	-	3	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Maine												
All races, both sexes	3	4	4	2	1	4	3	2	1	3	-	-
Male	3	4	4	2	-	2	-	-	-	3	-	-
Female	-	-	-	-	1	2	3	2	1	-	-	-
White, both sexes	3	4	4	2	1	4	3	2	1	3	-	-
Male	3	4	4	2	-	2	-	-	-	3	-	-
Female	-	-	-	-	1	2	3	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	3	2	6	9	7	9	6	10	-	2	-	-
Male	3	2	4	6	4	3	2	5	-	-	-	-
Female	-	-	2	3	3	6	4	5	-	2	-	-
White, both sexes	2	2	2	5	4	6	4	9	-	2	-	-
Male	2	2	2	4	2	2	2	5	-	-	-	-
Female	-	-	-	1	2	4	2	4	-	2	-	-
Black, both sexes	1	-	4	4	3	2	2	1	-	-	-	-
Male	1	-	2	2	2	1	-	-	-	-	-	-
Female	-	-	2	2	1	1	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Massachusetts												
All races, both sexes	2	1	4	4	2	12	9	13	8	3	-	-
Male	2	1	4	3	2	8	4	5	2	-	-	-
Female	-	-	-	1	-	4	5	8	6	3	-	-
White, both sexes	1	1	3	4	2	12	8	13	8	3	-	-
Male	1	1	3	3	2	8	4	5	2	-	-	-
Female	-	-	-	1	-	4	4	8	6	3	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1	3	6	6	10	11	9	7	9	-	1	-
Male	1	3	4	3	7	8	5	3	2	-	-	-
Female	-	-	2	3	3	3	4	4	7	-	1	-
White, both sexes	1	2	4	4	7	8	9	6	7	-	1	-
Male	1	2	3	2	4	6	5	2	2	-	-	-
Female	-	-	1	2	3	2	4	4	5	-	1	-
Black, both sexes	-	1	2	2	3	3	-	1	2	-	-	-
Male	-	1	1	1	3	2	-	1	-	-	-	-
Female	-	-	1	1	-	1	-	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Minnesota												
All races, both sexes	1	2	2	4	1	8	8	5	5	2	1	-
Male	-	2	1	3	1	3	4	3	-	-	1	-
Female	1	-	1	1	-	5	4	2	5	2	-	-
White, both sexes	1	2	2	3	1	8	8	5	5	2	1	-
Male	-	2	1	2	1	3	4	3	-	-	1	-
Female	1	-	1	1	-	5	4	2	5	2	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	2	-	4	2	3	1	3	3	-	-	-
Male	-	2	-	3	-	1	-	2	1	-	-	-
Female	-	-	-	1	2	2	1	1	2	-	-	-
White, both sexes	-	1	-	1	1	3	1	1	3	-	-	-
Male	-	1	-	1	-	1	-	1	1	-	-	-
Female	-	-	-	-	1	2	1	-	2	-	-	-
Black, both sexes	-	1	-	3	1	-	-	2	-	-	-	-
Male	-	1	-	2	-	-	-	1	-	-	-	-
Female	-	-	-	1	1	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Missouri												
All races, both sexes	-	2	6	7	5	4	9	14	12	3	2	-
Male	-	2	3	5	2	1	2	5	-	-	1	-
Female	-	-	3	2	3	3	7	9	12	3	1	-
White, both sexes	-	2	4	5	5	2	7	14	12	3	2	-
Male	-	2	1	4	2	1	1	5	-	-	1	-
Female	-	-	3	1	3	1	6	9	12	3	1	-
Black, both sexes	-	-	2	2	-	2	2	-	-	-	-	-
Male	-	-	2	1	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	2	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	1	-	-	1	3	1	1	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	3	-	1	-	-	-
White, both sexes	-	-	1	-	-	1	3	1	1	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	3	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Nebraska												
All races, both sexes	-	-	2	2	-	2	4	2	2	1	-	-
Male	-	-	1	1	-	-	3	1	-	1	-	-
Female	-	-	1	1	-	2	1	1	2	-	-	-
White, both sexes	-	-	2	1	-	2	4	2	2	1	-	-
Male	-	-	1	1	-	-	3	1	-	1	-	-
Female	-	-	1	-	-	2	1	1	2	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	2	1	2	4	2	3	1	-	-	-
Male	-	-	2	1	2	3	-	3	-	-	-	-
Female	-	-	-	-	-	1	2	-	1	-	-	-
White, both sexes	-	-	2	1	2	4	2	2	1	-	-	-
Male	-	-	2	1	2	3	-	2	-	-	-	-
Female	-	-	-	-	-	1	2	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
New Hampshire												
All races, both sexes	1	1	1	1	2	-	2	-	1	-	-	-
Male	1	1	1	1	1	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	1	-	-	-
White, both sexes	1	1	1	1	2	-	2	-	1	-	-	-
Male	1	1	1	1	1	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	3	2	2	5	10	15	18	14	20	13	2	-
Male	1	1	2	3	3	7	10	4	5	2	-	-
Female	2	1	-	2	7	8	8	10	15	11	2	-
White, both sexes	3	1	1	5	9	14	16	12	20	13	2	-
Male	1	-	1	3	2	7	9	4	5	2	-	-
Female	2	1	-	2	7	7	7	8	15	11	2	-
Black, both sexes	-	1	-	-	1	1	1	2	-	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
New Mexico												
All races, both sexes	-	3	1	-	1	1	2	2	2	1	-	-
Male	-	2	1	-	1	1	1	1	1	-	-	-
Female	-	1	-	-	-	-	1	1	1	1	-	-
White, both sexes	-	3	1	-	1	1	2	2	2	1	-	-
Male	-	2	1	-	1	1	1	1	1	-	-	-
Female	-	1	-	-	-	-	1	1	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	3	6	12	8	12	20	24	30	30	13	3	-
Male	3	4	9	5	5	11	15	9	6	1	-	-
Female	-	2	3	3	7	9	9	21	24	12	3	-
White, both sexes	3	6	10	6	10	17	23	29	27	12	3	-
Male	3	4	7	3	4	10	14	9	5	1	-	-
Female	-	2	3	3	6	7	9	20	22	11	3	-
Black, both sexes	-	-	2	1	2	3	1	-	2	1	-	-
Male	-	-	2	1	1	1	1	-	-	-	-	-
Female	-	-	-	-	1	2	-	-	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	1	1	-	-	-
Male	-	-	-	1	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
North Carolina												
All races, both sexes	5	6	4	10	10	10	12	9	6	2	-	-
Male	4	5	3	9	7	8	4	5	2	-	-	-
Female	1	1	1	1	3	2	8	4	4	2	-	-
White, both sexes	4	4	4	9	9	7	11	9	5	1	-	-
Male	3	4	3	8	7	6	4	5	2	-	-	-
Female	1	-	1	1	2	1	7	4	3	1	-	-
Black, both sexes	1	2	-	1	1	2	1	-	1	1	-	-
Male	1	1	-	1	-	1	-	-	-	-	-	-
Female	-	1	-	-	1	1	1	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	1	1	-	2	-	-	1	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	1	-	-	-
White, both sexes	-	-	1	1	-	2	-	-	1	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Ohio												
All races, both sexes	11	16	14	7	11	22	19	17	15	2	-	-
Male	11	12	8	6	8	9	13	8	4	-	-	-
Female	-	4	6	1	3	13	6	9	11	2	-	-
White, both sexes	11	15	14	6	11	22	17	16	15	2	-	-
Male	11	12	8	5	8	9	11	7	4	-	-	-
Female	-	3	6	1	3	13	6	9	11	2	-	-
Black, both sexes	-	1	-	1	-	-	2	1	-	-	-	-
Male	-	-	-	1	-	-	2	1	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	64	110	99	88	104	95	106	128	76	43	12	-
Male	49	76	69	53	69	46	47	39	16	6	4	-
Female	15	34	30	35	35	49	59	89	60	37	8	-
White, both sexes	53	89	87	81	93	85	96	122	72	40	11	-
Male	38	61	60	49	62	40	43	37	15	6	4	-
Female	15	28	27	32	31	45	53	85	57	34	7	-
Black, both sexes	6	12	6	2	6	7	8	3	2	-	1	-
Male	6	8	5	1	4	4	3	-	1	-	-	-
Female	-	4	1	1	2	3	5	3	1	-	1	-
American Indian 1/, both sexes	5	7	4	4	5	2	2	3	2	3	-	-
Male	5	6	3	2	3	2	1	2	-	-	-	-
Female	-	1	1	2	2	-	1	1	2	3	-	-
Asian or Pacific Islander 2/, both sexes	-	2	2	1	-	1	-	-	-	-	-	-
Male	-	1	1	1	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Oregon												
All races, both sexes	1	1	1	4	4	4	3	2	3	-	-	-
Male	-	1	1	4	3	2	1	-	1	-	-	-
Female	1	-	-	-	1	2	2	2	2	-	-	-
White, both sexes	1	1	1	4	4	3	3	2	3	-	-	-
Male	-	1	1	4	3	2	1	-	1	-	-	-
Female	1	-	-	-	1	1	2	2	2	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	4	11	17	19	8	20	32	24	20	8	3	-
Male	3	6	14	11	3	10	10	5	6	1	-	-
Female	1	5	3	8	5	10	22	19	14	7	3	-
White, both sexes	4	10	17	18	7	19	30	23	19	8	2	-
Male	3	6	14	11	3	10	9	5	6	1	-	-
Female	1	4	3	7	4	9	21	18	13	7	2	-
Black, both sexes	-	1	-	1	1	1	2	1	1	-	1	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	1	1	1	1	1	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Rhode Island												
All races, both sexes	-	2	-	1	1	-	1	2	-	2	-	-
Male	-	2	-	1	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	2	-	-
White, both sexes	-	2	-	1	1	-	1	2	-	2	-	-
Male	-	2	-	1	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	14	23	28	21	20	25	14	16	9	-	-	-
Male	9	18	17	10	13	14	4	8	1	-	-	-
Female	5	5	11	11	7	11	10	8	8	-	-	-
White, both sexes	3	15	16	11	9	14	10	10	5	-	-	-
Male	3	11	7	4	7	8	4	6	1	-	-	-
Female	-	4	9	7	2	6	6	4	4	-	-	-
Black, both sexes	11	8	12	10	11	10	3	6	4	-	-	-
Male	6	7	10	6	6	6	-	2	-	-	-	-
Female	5	1	2	4	5	4	3	4	4	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
South Dakota												
All races, both sexes	-	-	1	-	-	1	1	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	1	-	-	1	1	-	-	-	-	-
White, both sexes	-	-	1	-	-	1	1	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	1	-	-	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	5	1	4	8	5	5	10	8	3	2	1	-
Male	2	-	3	6	3	2	6	4	-	1	-	-
Female	3	1	1	2	2	3	4	4	3	1	1	-
White, both sexes	3	1	3	6	5	5	9	6	3	1	-	-
Male	1	-	2	4	3	2	6	4	-	-	-	-
Female	2	1	1	2	2	3	3	2	3	1	-	-
Black, both sexes	2	-	1	1	-	-	1	2	-	1	1	-
Male	1	-	1	1	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	1	2	-	-	1	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Texas												
All races, both sexes	7	7	12	18	29	30	24	27	14	4	2	-
Male	5	6	7	12	12	10	13	10	6	1	-	-
Female	2	1	5	6	17	20	11	17	8	3	2	-
White, both sexes	4	6	11	15	22	26	24	25	12	3	2	-
Male	3	5	7	11	8	9	13	10	4	-	-	-
Female	1	1	4	4	14	17	11	15	8	3	2	-
Black, both sexes	3	1	1	3	6	3	-	2	2	-	-	-
Male	2	1	-	1	4	1	-	-	2	-	-	-
Female	1	-	1	2	2	2	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	2	-	-	2	3	3	4	1	-	-
Male	-	-	2	-	-	-	2	1	1	-	-	-
Female	-	-	-	-	-	2	1	2	3	1	-	-
White, both sexes	-	-	2	-	-	2	3	3	4	1	-	-
Male	-	-	2	-	-	-	2	1	1	-	-	-
Female	-	-	-	-	-	2	1	2	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Vermont												
All races, both sexes	-	-	1	-	-	-	2	-	1	-	-	-
Male	-	-	1	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	2	-	1	-	-	-
Male	-	-	1	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	53	57	47	43	45	58	70	60	39	16	8	-
Male	36	47	35	30	25	30	35	19	8	5	3	-
Female	17	10	12	13	20	28	35	41	31	11	5	-
White, both sexes	35	35	28	28	30	40	62	45	31	10	5	-
Male	23	31	19	21	15	18	32	16	7	4	-	-
Female	12	4	9	7	15	22	30	29	24	6	5	-
Black, both sexes	17	21	18	14	13	18	8	15	8	6	3	-
Male	13	15	15	8	9	12	3	3	1	1	3	-
Female	4	6	3	6	4	6	5	12	7	5	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	1	2	-	-	-	-	-	-	-
Male	-	1	1	1	1	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Washington												
All races, both sexes	1	2	3	2	1	6	2	4	5	4	-	-
Male	1	1	1	2	1	3	2	2	-	-	-	-
Female	-	1	2	-	-	3	-	2	5	4	-	-
White, both sexes	1	2	2	2	1	6	1	4	5	4	-	-
Male	1	1	1	2	1	3	1	2	-	-	-	-
Female	-	1	1	-	-	3	-	2	5	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	1	1	2	3	5	-	3	2	2	2	-	-
Male	1	1	-	3	5	-	2	-	-	-	-	-
Female	-	-	2	-	-	-	1	2	2	2	-	-
White, both sexes	1	1	2	3	5	-	3	2	2	2	-	-
Male	1	1	-	3	5	-	2	-	-	-	-	-
Female	-	-	2	-	-	-	1	2	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Wisconsin												
All races, both sexes	1	1	1	5	4	6	5	8	6	2	-	-
Male	1	-	1	3	3	4	3	4	1	-	-	-
Female	-	1	-	2	1	2	2	4	5	2	-	-
White, both sexes	1	1	1	4	4	6	4	8	6	2	-	-
Male	1	-	1	2	3	4	2	4	1	-	-	-
Female	-	1	-	2	1	2	2	4	5	2	-	-
Black, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	1	-	-	-	1	3	1	-	-	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	-	1	-	-	-	-	2	-	-	-	-	-
White, both sexes	-	1	-	-	-	1	3	1	-	-	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	-	1	-	-	-	-	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
United States												
All races, both sexes	269,291	12	4	7	3	23	71	198	433	1,070	2,638	5,524
Male	142,179	5	3	6	1	16	56	150	348	823	2,065	4,224
Female	127,112	7	1	1	2	7	15	48	85	247	573	1,300
White, both sexes	235,140	8	2	6	2	15	46	133	318	793	2,030	4,305
Male	124,253	4	1	5	-	10	37	100	255	614	1,620	3,381
Female	110,887	4	1	1	2	5	9	33	63	179	410	924
Black, both sexes	28,043	2	1	1	-	8	22	55	96	226	484	1,067
Male	14,468	1	1	1	-	6	17	41	76	162	341	719
Female	13,575	1	-	-	-	2	5	14	20	64	143	348
American Indian 1/, both sexes	1,125	1	1	-	-	-	1	2	8	8	36	45
Male	655	-	1	-	-	-	1	1	6	7	27	35
Female	470	1	-	-	-	-	-	1	2	1	9	10
Asian or Pacific Islander 2/, both sexes	4,983	1	-	-	1	-	2	8	11	43	88	107
Male	2,803	-	-	-	1	-	1	8	11	40	77	89
Female	2,180	1	-	-	-	-	1	-	-	3	11	18
Alabama												
All races, both sexes	2,921	-	-	-	-	-	-	2	4	17	34	71
Male	1,531	-	-	-	-	-	-	2	3	11	24	56
Female	1,390	-	-	-	-	-	-	-	1	6	10	15
White, both sexes	2,384	-	-	-	-	-	-	-	4	13	19	54
Male	1,253	-	-	-	-	-	-	-	3	8	13	43
Female	1,131	-	-	-	-	-	-	-	1	5	6	11
Black, both sexes	530	-	-	-	-	-	-	2	-	4	13	17
Male	272	-	-	-	-	-	-	2	-	3	10	13
Female	258	-	-	-	-	-	-	-	-	1	3	4
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	2	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Alaska												
All races, both sexes	259	-	-	-	-	-	-	-	-	1	9	6
Male	167	-	-	-	-	-	-	-	-	-	9	6
Female	92	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	210	-	-	-	-	-	-	-	-	-	9	4
Male	140	-	-	-	-	-	-	-	-	-	9	4
Female	70	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	35	-	-	-	-	-	-	-	-	-	-	1
Male	21	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	1	-	-
Arizona												
All races, both sexes	5,562	-	-	-	1	-	4	-	10	23	47	111
Male	3,166	-	-	-	1	-	4	-	8	20	37	83
Female	2,396	-	-	-	-	-	-	-	2	3	10	28
White, both sexes	5,296	-	-	-	-	-	2	-	10	20	42	98
Male	3,026	-	-	-	-	-	2	-	8	18	35	75
Female	2,270	-	-	-	-	-	-	-	2	2	7	23
Black, both sexes	146	-	-	-	-	-	2	-	-	2	-	9
Male	71	-	-	-	-	-	2	-	-	1	-	4
Female	75	-	-	-	-	-	-	-	-	1	-	5
American Indian 1/, both sexes	93	-	-	-	-	-	-	-	-	-	5	2
Male	53	-	-	-	-	-	-	-	-	-	2	2
Female	40	-	-	-	-	-	-	-	-	-	3	-
Asian or Pacific Islander 2/, both sexes	27	-	-	-	1	-	-	-	-	1	-	2
Male	16	-	-	-	1	-	-	-	-	1	-	2
Female	11	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Arkansas												
All races, both sexes	1,873	-	1	-	-	-	-	1	8	9	23	33
Male	965	-	1	-	-	-	-	-	8	5	20	26
Female	908	-	-	-	-	-	-	1	-	4	3	7
White, both sexes	1,644	-	-	-	-	-	-	-	8	8	15	24
Male	850	-	-	-	-	-	-	-	8	4	13	18
Female	794	-	-	-	-	-	-	-	-	4	2	6
Black, both sexes	219	-	1	-	-	-	-	1	-	1	8	9
Male	110	-	1	-	-	-	-	-	-	1	7	8
Female	109	-	-	-	-	-	-	1	-	-	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	31,081	-	-	-	-	3	6	17	47	98	232	511
Male	16,418	-	-	-	-	3	6	15	41	72	193	409
Female	14,663	-	-	-	-	-	-	2	6	26	39	102
White, both sexes	26,264	-	-	-	-	3	6	11	34	63	168	394
Male	13,814	-	-	-	-	3	6	10	31	43	141	316
Female	12,450	-	-	-	-	-	-	1	3	20	27	78
Black, both sexes	2,410	-	-	-	-	-	-	5	8	19	33	70
Male	1,290	-	-	-	-	-	-	4	5	14	23	50
Female	1,120	-	-	-	-	-	-	1	3	5	10	20
American Indian 1/, both sexes	144	-	-	-	-	-	-	-	1	1	3	8
Male	79	-	-	-	-	-	-	-	1	1	3	8
Female	65	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2,263	-	-	-	-	-	-	1	4	15	28	39
Male	1,235	-	-	-	-	-	-	1	4	14	26	35
Female	1,028	-	-	-	-	-	-	-	-	1	2	4

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Colorado												
All races, both sexes	2,650	-	-	1	-	-	1	5	6	14	39	72
Male	1,518	-	-	1	-	-	1	2	6	11	33	57
Female	1,132	-	-	-	-	-	-	3	-	3	6	15
White, both sexes	2,553	-	-	1	-	-	1	4	5	14	38	68
Male	1,460	-	-	1	-	-	1	1	5	11	32	54
Female	1,093	-	-	-	-	-	-	3	-	3	6	14
Black, both sexes	75	-	-	-	-	-	-	-	-	-	-	3
Male	42	-	-	-	-	-	-	-	-	-	-	2
Female	33	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	1	-	1	-
Male	8	-	-	-	-	-	-	-	1	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	1	-	-	-	1
Male	8	-	-	-	-	-	-	1	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	3,106	1	-	-	-	-	-	2	4	10	33	50
Male	1,582	1	-	-	-	-	-	2	3	9	26	38
Female	1,524	-	-	-	-	-	-	-	1	1	7	12
White, both sexes	2,911	1	-	-	-	-	-	2	2	8	25	39
Male	1,468	1	-	-	-	-	-	2	1	7	21	32
Female	1,443	-	-	-	-	-	-	-	1	1	4	7
Black, both sexes	175	-	-	-	-	-	-	-	2	2	7	11
Male	100	-	-	-	-	-	-	-	2	2	4	6
Female	75	-	-	-	-	-	-	-	-	-	3	5
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	18	-	-	-	-	-	-	-	-	-	1	-
Male	13	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Delaware												
All races, both sexes	939	-	-	-	-	-	-	1	1	7	13	33
Male	528	-	-	-	-	-	-	1	1	6	10	27
Female	411	-	-	-	-	-	-	-	-	1	3	6
White, both sexes	798	-	-	-	-	-	-	1	1	5	11	22
Male	447	-	-	-	-	-	-	1	1	4	8	18
Female	351	-	-	-	-	-	-	-	-	1	3	4
Black, both sexes	131	-	-	-	-	-	-	-	-	2	2	11
Male	73	-	-	-	-	-	-	-	-	2	2	9
Female	58	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	819	-	-	-	-	-	-	-	1	7	10	16
Male	433	-	-	-	-	-	-	-	1	5	8	13
Female	386	-	-	-	-	-	-	-	-	2	2	3
White, both sexes	191	-	-	-	-	-	-	-	-	1	1	1
Male	117	-	-	-	-	-	-	-	-	1	-	1
Female	74	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	619	-	-	-	-	-	-	-	1	5	9	15
Male	312	-	-	-	-	-	-	-	1	3	8	12
Female	307	-	-	-	-	-	-	-	-	2	1	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Florida												
All races, both sexes	22,033	-	-	-	-	1	6	15	31	73	191	397
Male	11,966	-	-	-	-	-	5	13	25	59	147	308
Female	10,067	-	-	-	-	1	1	2	6	14	44	89
White, both sexes	20,027	-	-	-	-	1	3	9	19	51	149	333
Male	10,923	-	-	-	-	-	3	7	16	42	119	268
Female	9,104	-	-	-	-	1	-	2	3	9	30	65
Black, both sexes	1,873	-	-	-	-	-	3	4	11	22	36	63
Male	968	-	-	-	-	-	2	4	8	17	23	39
Female	905	-	-	-	-	-	1	-	3	5	13	24
American Indian 1/, both sexes	19	-	-	-	-	-	-	-	-	-	1	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	114	-	-	-	-	-	-	2	1	-	5	1
Male	62	-	-	-	-	-	-	2	1	-	5	1
Female	52	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	4,906	-	-	-	-	4	4	5	8	44	89	170
Male	2,812	-	-	-	-	4	4	4	6	32	69	127
Female	2,094	-	-	-	-	-	-	1	2	12	20	43
White, both sexes	3,752	-	-	-	-	3	2	4	6	30	55	118
Male	2,193	-	-	-	-	3	2	3	4	22	43	90
Female	1,559	-	-	-	-	-	-	1	2	8	12	28
Black, both sexes	1,116	-	-	-	-	1	2	1	2	13	34	49
Male	597	-	-	-	-	1	2	1	2	9	26	35
Female	519	-	-	-	-	-	-	-	-	4	8	14
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	35	-	-	-	-	-	-	-	-	1	-	3
Male	21	-	-	-	-	-	-	-	-	1	-	2
Female	14	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Hawaii												
All races, both sexes	787	-	-	-	-	-	-	2	1	3	13	17
Male	501	-	-	-	-	-	-	2	-	3	8	11
Female	286	-	-	-	-	-	-	-	1	-	5	6
White, both sexes	227	-	-	-	-	-	-	1	1	2	5	6
Male	157	-	-	-	-	-	-	1	-	2	3	5
Female	70	-	-	-	-	-	-	-	1	-	2	1
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	553	-	-	-	-	-	-	1	-	1	8	11
Male	339	-	-	-	-	-	-	1	-	1	5	6
Female	214	-	-	-	-	-	-	-	-	-	3	5
Idaho												
All races, both sexes	741	-	-	-	-	-	-	-	1	1	6	14
Male	403	-	-	-	-	-	-	-	1	1	3	9
Female	338	-	-	-	-	-	-	-	-	-	3	5
White, both sexes	734	-	-	-	-	-	-	-	-	-	6	14
Male	398	-	-	-	-	-	-	-	-	-	3	9
Female	336	-	-	-	-	-	-	-	-	-	3	5
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Illinois												
All races, both sexes	10,263	1	-	1	-	1	5	16	19	33	104	234
Male	5,361	-	-	1	-	1	5	10	11	28	85	166
Female	4,902	1	-	-	-	-	-	6	8	5	19	68
White, both sexes	8,716	1	-	1	-	1	4	13	16	24	72	159
Male	4,506	-	-	1	-	1	4	8	10	20	59	117
Female	4,210	1	-	-	-	-	-	5	6	4	13	42
Black, both sexes	1,426	-	-	-	-	-	-	3	3	8	30	73
Male	790	-	-	-	-	-	-	2	1	7	24	47
Female	636	-	-	-	-	-	-	1	2	1	6	26
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	113	-	-	-	-	-	1	-	-	1	2	1
Male	63	-	-	-	-	-	1	-	-	1	2	1
Female	50	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	5,119	1	-	1	-	-	2	5	8	27	61	122
Male	2,661	1	-	1	-	-	2	5	7	17	49	96
Female	2,458	-	-	-	-	-	-	-	1	10	12	26
White, both sexes	4,704	1	-	1	-	-	2	5	6	22	47	105
Male	2,436	1	-	1	-	-	2	5	5	15	40	85
Female	2,268	-	-	-	-	-	-	-	1	7	7	20
Black, both sexes	407	-	-	-	-	-	-	-	2	5	14	17
Male	222	-	-	-	-	-	-	-	2	2	9	11
Female	185	-	-	-	-	-	-	-	-	3	5	6
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Iowa												
All races, both sexes	3,503	1	-	-	-	-	-	2	2	7	28	56
Male	1,796	-	-	-	-	-	-	2	1	6	23	36
Female	1,707	1	-	-	-	-	-	-	1	1	5	20
White, both sexes	3,450	1	-	-	-	-	-	2	2	7	28	51
Male	1,767	-	-	-	-	-	-	2	1	6	23	33
Female	1,683	1	-	-	-	-	-	-	1	1	5	18
Black, both sexes	41	-	-	-	-	-	-	-	-	-	-	5
Male	22	-	-	-	-	-	-	-	-	-	-	3
Female	19	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	2,234	-	-	-	1	-	4	-	9	14	31	42
Male	1,214	-	-	-	-	-	3	-	6	13	20	35
Female	1,020	-	-	-	1	-	1	-	3	1	11	7
White, both sexes	2,104	-	-	-	1	-	4	-	9	10	27	39
Male	1,143	-	-	-	-	-	3	-	6	10	18	33
Female	961	-	-	-	1	-	1	-	3	-	9	6
Black, both sexes	103	-	-	-	-	-	-	-	-	2	-	2
Male	53	-	-	-	-	-	-	-	-	1	-	1
Female	50	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	20	-	-	-	-	-	-	-	-	1	1	1
Male	14	-	-	-	-	-	-	-	-	1	-	1
Female	6	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	1	3	-
Male	4	-	-	-	-	-	-	-	-	1	2	-
Female	3	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Kentucky												
All races, both sexes	3,065	-	-	-	-	-	3	6	12	18	57	67
Male	1,606	-	-	-	-	-	3	4	8	14	40	58
Female	1,459	-	-	-	-	-	-	2	4	4	17	9
White, both sexes	2,904	-	-	-	-	-	2	5	11	16	50	63
Male	1,526	-	-	-	-	-	2	3	7	13	37	55
Female	1,378	-	-	-	-	-	-	2	4	3	13	8
Black, both sexes	156	-	-	-	-	-	1	1	1	2	7	3
Male	78	-	-	-	-	-	1	1	1	1	3	2
Female	78	-	-	-	-	-	-	-	-	1	4	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	3,105	1	-	-	-	1	3	7	6	19	46	96
Male	1,690	-	-	-	-	1	2	7	4	13	32	61
Female	1,415	1	-	-	-	-	1	-	2	6	14	35
White, both sexes	2,350	-	-	-	-	-	1	4	6	13	27	65
Male	1,293	-	-	-	-	-	-	4	4	11	17	43
Female	1,057	-	-	-	-	-	1	-	2	2	10	22
Black, both sexes	745	1	-	-	-	1	2	1	-	6	19	30
Male	391	-	-	-	-	1	2	1	-	2	15	17
Female	354	1	-	-	-	-	-	-	-	4	4	13
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	2	-	-	-	1
Male	5	-	-	-	-	-	-	2	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Maine												
All races, both sexes	1,098	-	-	-	-	-	1	1	1	4	20	21
Male	621	-	-	-	-	-	-	1	1	3	17	17
Female	477	-	-	-	-	-	1	-	-	1	3	4
White, both sexes	1,091	-	-	-	-	-	1	1	1	4	19	20
Male	616	-	-	-	-	-	-	1	1	3	16	16
Female	475	-	-	-	-	-	1	-	-	1	3	4
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	5,376	-	-	1	-	-	2	4	8	19	49	99
Male	2,715	-	-	1	-	-	2	3	6	11	36	72
Female	2,661	-	-	-	-	-	-	1	2	8	13	27
White, both sexes	4,028	-	-	-	-	-	-	2	2	14	32	59
Male	2,034	-	-	-	-	-	-	2	1	7	23	47
Female	1,994	-	-	-	-	-	-	-	1	7	9	12
Black, both sexes	1,277	-	-	1	-	-	2	2	6	5	17	37
Male	640	-	-	1	-	-	2	1	5	4	13	23
Female	637	-	-	-	-	-	-	1	1	1	4	14
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	65	-	-	-	-	-	-	-	-	-	-	2
Male	37	-	-	-	-	-	-	-	-	-	-	1
Female	28	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Massachusetts												
All races, both sexes	5,579	-	-	-	-	1	-	2	5	19	46	111
Male	2,931	-	-	-	-	-	-	1	3	17	36	90
Female	2,648	-	-	-	-	1	-	1	2	2	10	21
White, both sexes	5,355	-	-	-	-	1	-	2	5	17	37	102
Male	2,812	-	-	-	-	-	-	1	3	15	32	82
Female	2,543	-	-	-	-	1	-	1	2	2	5	20
Black, both sexes	174	-	-	-	-	-	-	-	-	1	7	7
Male	91	-	-	-	-	-	-	-	-	1	3	7
Female	83	-	-	-	-	-	-	-	-	-	4	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	43	-	-	-	-	-	-	-	-	1	2	2
Male	24	-	-	-	-	-	-	-	-	1	1	1
Female	19	-	-	-	-	-	-	-	-	-	1	1
Michigan												
All races, both sexes	11,906	1	-	-	-	-	2	6	20	51	105	265
Male	6,398	-	-	-	-	-	1	4	18	38	85	211
Female	5,508	1	-	-	-	-	1	2	2	13	20	54
White, both sexes	9,978	-	-	-	-	-	-	2	10	32	81	202
Male	5,336	-	-	-	-	-	-	1	8	26	68	162
Female	4,642	-	-	-	-	-	-	1	2	6	13	40
Black, both sexes	1,834	-	-	-	-	-	1	4	10	18	18	61
Male	1,014	-	-	-	-	-	1	3	10	11	13	47
Female	820	-	-	-	-	-	-	1	-	7	5	14
American Indian 1/, both sexes	52	1	-	-	-	-	-	-	-	1	4	-
Male	31	-	-	-	-	-	-	-	-	1	2	-
Female	21	1	-	-	-	-	-	-	-	-	2	-
Asian or Pacific Islander 2/, both sexes	42	-	-	-	-	-	1	-	-	-	2	2
Male	17	-	-	-	-	-	-	-	-	-	2	2
Female	25	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Minnesota												
All races, both sexes	3,013	-	-	-	-	1	-	2	7	12	28	75
Male	1,737	-	-	-	-	1	-	2	7	11	22	59
Female	1,276	-	-	-	-	-	-	-	-	1	6	16
White, both sexes	2,920	-	-	-	-	1	-	1	5	10	24	68
Male	1,680	-	-	-	-	1	-	1	5	9	18	55
Female	1,240	-	-	-	-	-	-	-	-	1	6	13
Black, both sexes	44	-	-	-	-	-	-	1	1	1	2	7
Male	26	-	-	-	-	-	-	1	1	1	2	4
Female	18	-	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes	33	-	-	-	-	-	-	-	1	-	1	-
Male	20	-	-	-	-	-	-	-	1	-	1	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	-	-	1	1	-
Male	11	-	-	-	-	-	-	-	-	1	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	2,245	-	-	-	-	-	1	4	5	15	47	60
Male	1,129	-	-	-	-	-	1	3	3	10	30	39
Female	1,116	-	-	-	-	-	-	1	2	5	17	21
White, both sexes	1,637	-	-	-	-	-	-	-	2	12	27	31
Male	819	-	-	-	-	-	-	-	2	8	16	21
Female	818	-	-	-	-	-	-	-	-	4	11	10
Black, both sexes	598	-	-	-	-	-	1	4	3	3	20	27
Male	301	-	-	-	-	-	1	3	1	2	14	16
Female	297	-	-	-	-	-	-	1	2	1	6	11
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Missouri												
All races, both sexes	5,500	-	-	-	-	-	-	2	7	16	68	129
Male	2,806	-	-	-	-	-	-	2	6	13	49	97
Female	2,694	-	-	-	-	-	-	-	1	3	19	32
White, both sexes	4,919	-	-	-	-	-	-	2	6	10	58	97
Male	2,517	-	-	-	-	-	-	2	5	7	43	78
Female	2,402	-	-	-	-	-	-	-	1	3	15	19
Black, both sexes	557	-	-	-	-	-	-	-	1	4	8	32
Male	277	-	-	-	-	-	-	-	1	4	5	19
Female	280	-	-	-	-	-	-	-	-	-	3	13
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	2	2	-
Male	7	-	-	-	-	-	-	-	-	2	1	-
Female	7	-	-	-	-	-	-	-	-	-	1	-
Montana												
All races, both sexes	762	-	-	-	-	-	-	-	-	1	11	20
Male	491	-	-	-	-	-	-	-	-	1	9	17
Female	271	-	-	-	-	-	-	-	-	-	2	3
White, both sexes	732	-	-	-	-	-	-	-	-	1	11	18
Male	472	-	-	-	-	-	-	-	-	1	9	16
Female	260	-	-	-	-	-	-	-	-	-	2	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	30	-	-	-	-	-	-	-	-	-	-	2
Male	19	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Nebraska												
All races, both sexes	1,350	1	-	-	-	-	1	1	1	3	13	21
Male	718	-	-	-	-	-	1	1	1	2	12	17
Female	632	1	-	-	-	-	-	-	-	1	1	4
White, both sexes	1,294	1	-	-	-	-	1	1	1	1	11	19
Male	683	-	-	-	-	-	1	1	1	1	10	15
Female	611	1	-	-	-	-	-	-	-	-	1	4
Black, both sexes	45	-	-	-	-	-	-	-	-	1	2	2
Male	29	-	-	-	-	-	-	-	-	1	2	2
Female	16	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1,889	-	-	-	1	-	1	2	1	8	21	41
Male	1,206	-	-	-	-	-	1	2	1	5	17	27
Female	683	-	-	-	1	-	-	-	-	3	4	14
White, both sexes	1,689	-	-	-	1	-	-	2	1	7	16	36
Male	1,082	-	-	-	-	-	-	2	1	4	12	27
Female	607	-	-	-	1	-	-	-	-	3	4	9
Black, both sexes	117	-	-	-	-	-	1	-	-	-	3	3
Male	69	-	-	-	-	-	1	-	-	-	3	-
Female	48	-	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	64	-	-	-	-	-	-	-	-	1	2	2
Male	40	-	-	-	-	-	-	-	-	1	2	-
Female	24	-	-	-	-	-	-	-	-	-	-	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
New Hampshire												
All races, both sexes	1,249	-	-	-	-	-	-	1	2	3	11	24
Male	638	-	-	-	-	-	-	1	-	2	10	16
Female	611	-	-	-	-	-	-	-	2	1	1	8
White, both sexes	1,244	-	-	-	-	-	-	1	2	3	11	23
Male	636	-	-	-	-	-	-	1	-	2	10	15
Female	608	-	-	-	-	-	-	-	2	1	1	8
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	8,951	2	-	-	-	-	-	2	11	16	58	156
Male	4,500	1	-	-	-	-	-	-	10	14	44	126
Female	4,451	1	-	-	-	-	-	2	1	2	14	30
White, both sexes	7,972	1	-	-	-	-	-	2	9	9	45	124
Male	3,986	1	-	-	-	-	-	-	8	8	35	104
Female	3,986	-	-	-	-	-	-	2	1	1	10	20
Black, both sexes	820	-	-	-	-	-	-	-	2	5	11	27
Male	423	-	-	-	-	-	-	-	2	4	7	17
Female	397	-	-	-	-	-	-	-	-	1	4	10
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	154	1	-	-	-	-	-	-	-	2	2	5
Male	91	-	-	-	-	-	-	-	-	2	2	5
Female	63	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
New Mexico												
All races, both sexes	1,650	-	1	-	-	-	-	3	1	4	15	37
Male	932	-	1	-	-	-	-	3	1	4	14	31
Female	718	-	-	-	-	-	-	-	-	-	1	6
White, both sexes	1,542	-	-	-	-	-	-	3	-	4	13	36
Male	869	-	-	-	-	-	-	3	-	4	12	30
Female	673	-	-	-	-	-	-	-	-	-	1	6
Black, both sexes	30	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	72	-	1	-	-	-	-	-	1	-	2	1
Male	44	-	1	-	-	-	-	-	1	-	2	1
Female	28	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	29,212	1	-	1	-	4	5	11	24	70	174	346
Male	13,577	1	-	1	-	1	3	6	19	52	137	256
Female	15,635	-	-	-	-	3	2	5	5	18	37	90
White, both sexes	24,283	1	-	1	-	1	3	5	11	51	115	229
Male	11,344	1	-	1	-	-	2	4	8	42	97	181
Female	12,939	-	-	-	-	1	1	1	3	9	18	48
Black, both sexes	4,124	-	-	-	-	3	2	6	11	16	44	101
Male	1,780	-	-	-	-	1	1	2	9	7	26	60
Female	2,344	-	-	-	-	2	1	4	2	9	18	41
American Indian 1/, both sexes	28	-	-	-	-	-	-	-	-	-	1	1
Male	17	-	-	-	-	-	-	-	-	-	1	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	777	-	-	-	-	-	-	-	2	3	14	15
Male	436	-	-	-	-	-	-	-	2	3	13	14
Female	341	-	-	-	-	-	-	-	-	-	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
North Carolina												
All races, both sexes	7,489	-	-	-	-	-	1	6	22	55	125	230
Male	4,140	-	-	-	-	-	1	3	20	43	95	174
Female	3,349	-	-	-	-	-	-	3	2	12	30	56
White, both sexes	5,992	-	-	-	-	-	1	2	11	39	91	158
Male	3,342	-	-	-	-	-	1	-	11	32	74	127
Female	2,650	-	-	-	-	-	-	2	-	7	17	31
Black, both sexes	1,399	-	-	-	-	-	-	3	10	14	30	69
Male	729	-	-	-	-	-	-	2	8	9	18	46
Female	670	-	-	-	-	-	-	1	2	5	12	23
American Indian 1/, both sexes	74	-	-	-	-	-	-	1	1	-	4	2
Male	51	-	-	-	-	-	-	1	1	-	3	1
Female	23	-	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	24	-	-	-	-	-	-	-	-	2	-	1
Male	18	-	-	-	-	-	-	-	-	2	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
North Dakota												
All races, both sexes	627	-	-	-	-	-	1	2	1	1	6	13
Male	330	-	-	-	-	-	1	-	1	1	5	10
Female	297	-	-	-	-	-	-	2	-	-	1	3
White, both sexes	612	-	-	-	-	-	-	2	1	1	4	13
Male	322	-	-	-	-	-	-	-	1	1	3	10
Female	290	-	-	-	-	-	-	2	-	-	1	3
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	14	-	-	-	-	-	1	-	-	-	2	-
Male	7	-	-	-	-	-	1	-	-	-	2	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Ohio												
All races, both sexes	12,731	-	1	-	-	1	3	9	28	60	103	284
Male	6,637	-	1	-	-	-	3	6	22	49	78	222
Female	6,094	-	-	-	-	1	-	3	6	11	25	62
White, both sexes	11,496	-	1	-	-	1	2	6	23	51	87	231
Male	5,979	-	1	-	-	-	2	4	18	40	69	188
Female	5,517	-	-	-	-	1	-	2	5	11	18	43
Black, both sexes	1,193	-	-	-	-	-	1	3	5	8	15	53
Male	632	-	-	-	-	-	1	2	4	8	9	34
Female	561	-	-	-	-	-	-	1	1	-	6	19
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	36	-	-	-	-	-	-	-	-	1	1	-
Male	22	-	-	-	-	-	-	-	-	1	-	-
Female	14	-	-	-	-	-	-	-	-	-	1	-
Oklahoma												
All races, both sexes	3,947	-	-	-	-	-	2	3	6	12	32	104
Male	2,054	-	-	-	-	-	-	2	5	10	23	73
Female	1,893	-	-	-	-	-	2	1	1	2	9	31
White, both sexes	3,495	-	-	-	-	-	2	1	5	10	26	74
Male	1,810	-	-	-	-	-	-	1	5	8	17	50
Female	1,685	-	-	-	-	-	2	-	-	2	9	24
Black, both sexes	231	-	-	-	-	-	-	1	-	1	2	18
Male	129	-	-	-	-	-	-	1	-	1	2	15
Female	102	-	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes	207	-	-	-	-	-	-	1	1	1	4	12
Male	110	-	-	-	-	-	-	-	-	1	4	8
Female	97	-	-	-	-	-	-	1	1	-	-	4
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Oregon												
All races, both sexes	2,641	-	-	-	-	-	-	-	10	12	29	57
Male	1,571	-	-	-	-	-	-	-	9	11	24	48
Female	1,070	-	-	-	-	-	-	-	1	1	5	9
White, both sexes	2,561	-	-	-	-	-	-	-	9	10	27	54
Male	1,522	-	-	-	-	-	-	-	8	9	22	46
Female	1,039	-	-	-	-	-	-	-	1	1	5	8
Black, both sexes	27	-	-	-	-	-	-	-	-	1	-	2
Male	18	-	-	-	-	-	-	-	-	1	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	18	-	-	-	-	-	-	-	-	-	1	-
Male	9	-	-	-	-	-	-	-	-	-	1	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	35	-	-	-	-	-	-	-	1	1	1	1
Male	22	-	-	-	-	-	-	-	1	1	1	1
Female	13	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	13,629	1	1	-	-	1	5	8	16	35	87	216
Male	6,937	1	-	-	-	1	1	6	12	26	69	159
Female	6,692	-	1	-	-	-	4	2	4	9	18	57
White, both sexes	12,402	-	1	-	-	-	3	7	10	30	70	180
Male	6,305	-	-	-	-	-	1	5	7	23	56	136
Female	6,097	-	1	-	-	-	2	2	3	7	14	44
Black, both sexes	1,163	1	-	-	-	1	2	1	6	5	15	35
Male	593	1	-	-	-	1	-	1	5	3	11	22
Female	570	-	-	-	-	-	2	-	1	2	4	13
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	59	-	-	-	-	-	-	-	-	-	2	1
Male	35	-	-	-	-	-	-	-	-	-	2	1
Female	24	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Rhode Island												
All races, both sexes	1,507	-	-	-	-	-	-	-	3	10	12	20
Male	677	-	-	-	-	-	-	-	3	7	10	19
Female	830	-	-	-	-	-	-	-	-	3	2	1
White, both sexes	1,455	-	-	-	-	-	-	-	3	10	12	18
Male	650	-	-	-	-	-	-	-	3	7	10	17
Female	805	-	-	-	-	-	-	-	-	3	2	1
Black, both sexes	44	-	-	-	-	-	-	-	-	-	-	1
Male	22	-	-	-	-	-	-	-	-	-	-	1
Female	22	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	2,760	-	-	1	-	-	1	1	2	20	42	66
Male	1,552	-	-	1	-	-	-	1	2	15	32	49
Female	1,208	-	-	-	-	-	1	-	-	5	10	17
White, both sexes	2,147	-	-	1	-	-	-	-	2	11	34	41
Male	1,226	-	-	1	-	-	-	-	2	7	25	31
Female	921	-	-	-	-	-	-	-	-	4	9	10
Black, both sexes	598	-	-	-	-	-	1	1	-	8	8	23
Male	314	-	-	-	-	-	-	1	-	7	7	16
Female	284	-	-	-	-	-	1	-	-	1	1	7
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	1	-	1
Male	8	-	-	-	-	-	-	-	-	1	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
South Dakota												
All races, both sexes	682	-	-	-	-	-	-	-	1	2	7	7
Male	375	-	-	-	-	-	-	-	-	2	7	5
Female	307	-	-	-	-	-	-	-	1	-	-	2
White, both sexes	648	-	-	-	-	-	-	-	1	1	5	4
Male	359	-	-	-	-	-	-	-	-	1	5	3
Female	289	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	32	-	-	-	-	-	-	-	-	1	2	3
Male	16	-	-	-	-	-	-	-	-	1	2	2
Female	16	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	6,011	-	-	-	-	-	-	7	13	36	80	184
Male	3,138	-	-	-	-	-	-	5	9	26	62	130
Female	2,873	-	-	-	-	-	-	2	4	10	18	54
White, both sexes	5,239	-	-	-	-	-	-	5	10	30	65	149
Male	2,719	-	-	-	-	-	-	4	6	21	50	104
Female	2,520	-	-	-	-	-	-	1	4	9	15	45
Black, both sexes	756	-	-	-	-	-	-	2	3	5	14	34
Male	407	-	-	-	-	-	-	1	3	4	11	25
Female	349	-	-	-	-	-	-	1	-	1	3	9
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	-	1	1	1
Male	11	-	-	-	-	-	-	-	-	1	1	1
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Texas												
All races, both sexes	14,464	1	-	1	-	3	3	19	22	82	189	410
Male	8,035	-	-	-	-	2	3	15	16	62	150	318
Female	6,429	1	-	1	-	1	-	4	6	20	39	92
White, both sexes	12,716	1	-	1	-	3	2	14	20	61	151	328
Male	7,063	-	-	-	-	2	2	11	16	45	119	257
Female	5,653	1	-	1	-	1	-	3	4	16	32	71
Black, both sexes	1,614	-	-	-	-	-	1	5	2	20	31	78
Male	897	-	-	-	-	-	1	4	-	16	24	57
Female	717	-	-	-	-	-	-	1	2	4	7	21
American Indian 1/, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	116	-	-	-	-	-	-	-	-	1	7	4
Male	64	-	-	-	-	-	-	-	-	1	7	4
Female	52	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	928	-	-	-	-	-	-	2	3	8	11	22
Male	559	-	-	-	-	-	-	1	3	7	9	18
Female	369	-	-	-	-	-	-	1	-	1	2	4
White, both sexes	903	-	-	-	-	-	-	2	3	7	10	18
Male	544	-	-	-	-	-	-	1	3	6	9	15
Female	359	-	-	-	-	-	-	1	-	1	1	3
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	2
Male	6	-	-	-	-	-	-	-	-	-	-	2
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	1	-	2
Male	6	-	-	-	-	-	-	-	-	1	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Vermont												
All races, both sexes	610	-	-	-	-	-	-	-	1	1	6	7
Male	327	-	-	-	-	-	-	-	1	-	5	5
Female	283	-	-	-	-	-	-	-	-	1	1	2
White, both sexes	607	-	-	-	-	-	-	-	1	1	6	7
Male	326	-	-	-	-	-	-	-	1	-	5	5
Female	281	-	-	-	-	-	-	-	-	1	1	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	4,673	-	-	-	-	1	1	10	13	29	54	123
Male	2,605	-	-	-	-	1	1	9	11	22	42	101
Female	2,068	-	-	-	-	-	-	1	2	7	12	22
White, both sexes	3,748	-	-	-	-	-	1	6	9	15	38	80
Male	2,107	-	-	-	-	-	1	5	8	12	30	69
Female	1,641	-	-	-	-	-	-	1	1	3	8	11
Black, both sexes	847	-	-	-	-	1	-	4	4	14	13	39
Male	452	-	-	-	-	1	-	4	3	10	10	28
Female	395	-	-	-	-	-	-	-	1	4	3	11
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	74	-	-	-	-	-	-	-	-	-	3	4
Male	45	-	-	-	-	-	-	-	-	-	2	4
Female	29	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Washington												
All races, both sexes	5,134	-	-	-	-	1	1	1	4	15	44	92
Male	2,843	-	-	-	-	1	-	1	4	12	36	65
Female	2,291	-	-	-	-	-	1	-	-	3	8	27
White, both sexes	4,801	-	-	-	-	-	1	-	2	11	39	79
Male	2,650	-	-	-	-	-	-	-	2	9	32	57
Female	2,151	-	-	-	-	-	1	-	-	2	7	22
Black, both sexes	115	-	-	-	-	1	-	-	1	1	4	8
Male	80	-	-	-	-	1	-	-	1	1	3	6
Female	35	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	52	-	-	-	-	-	-	-	-	-	-	4
Male	28	-	-	-	-	-	-	-	-	-	-	1
Female	24	-	-	-	-	-	-	-	-	-	-	3
Asian or Pacific Islander 2/, both sexes	166	-	-	-	-	-	-	1	1	3	1	1
Male	85	-	-	-	-	-	-	1	1	2	1	1
Female	81	-	-	-	-	-	-	-	-	1	-	-
West Virginia												
All races, both sexes	2,389	-	-	-	-	-	1	1	9	14	44	76
Male	1,273	-	-	-	-	-	1	1	7	12	34	56
Female	1,116	-	-	-	-	-	-	-	2	2	10	20
White, both sexes	2,318	-	-	-	-	-	1	1	9	12	43	73
Male	1,236	-	-	-	-	-	1	1	7	10	34	53
Female	1,082	-	-	-	-	-	-	-	2	2	9	20
Black, both sexes	67	-	-	-	-	-	-	-	-	2	1	3
Male	36	-	-	-	-	-	-	-	-	2	-	3
Female	31	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Wisconsin												
All races, both sexes	4,029	-	-	-	-	-	-	1	7	11	32	85
Male	2,233	-	-	-	-	-	-	1	6	10	28	70
Female	1,796	-	-	-	-	-	-	-	1	1	4	15
White, both sexes	3,811	-	-	-	-	-	-	1	4	10	26	72
Male	2,119	-	-	-	-	-	-	1	4	9	24	60
Female	1,692	-	-	-	-	-	-	-	-	1	2	12
Black, both sexes	169	-	-	-	-	-	-	-	1	-	5	11
Male	86	-	-	-	-	-	-	-	1	-	3	9
Female	83	-	-	-	-	-	-	-	-	-	2	2
American Indian 1/, both sexes	32	-	-	-	-	-	-	-	1	1	1	-
Male	20	-	-	-	-	-	-	-	-	1	1	-
Female	12	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	-	-	-	1	-	-	2
Male	8	-	-	-	-	-	-	-	1	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	1
Wyoming												
All races, both sexes	294	-	-	-	-	-	1	1	1	1	3	5
Male	176	-	-	-	-	-	1	1	1	-	2	5
Female	118	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	286	-	-	-	-	-	1	1	-	1	2	5
Male	171	-	-	-	-	-	1	1	-	-	1	5
Female	115	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	1	-	1	-
Male	4	-	-	-	-	-	-	-	1	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
United States												
All races, both sexes	9,023	12,328	15,322	17,020	21,826	32,269	45,233	49,742	36,914	16,052	3,562	17
Male	6,905	9,286	10,959	11,599	13,875	18,883	23,612	22,055	12,780	3,905	611	12
Female	2,118	3,042	4,363	5,421	7,951	13,386	21,621	27,687	24,134	12,147	2,951	5
White, both sexes	6,981	9,636	12,415	13,892	18,233	28,086	40,609	45,512	34,172	14,757	3,177	12
Male	5,462	7,413	9,051	9,658	11,842	16,698	21,544	20,418	11,965	3,632	535	8
Female	1,519	2,223	3,364	4,234	6,391	11,388	19,065	25,094	22,207	11,125	2,642	4
Black, both sexes	1,801	2,325	2,508	2,614	2,984	3,397	3,663	3,299	2,117	1,054	315	4
Male	1,238	1,584	1,635	1,600	1,659	1,748	1,608	1,205	580	187	56	3
Female	563	741	873	1,014	1,325	1,649	2,055	2,094	1,537	867	259	1
American Indian 1/, both sexes	70	88	100	131	120	156	149	109	60	33	7	-
Male	57	64	65	89	73	87	70	48	16	7	1	-
Female	13	24	35	42	47	69	79	61	44	26	6	-
Asian or Pacific Islander 2/, both sexes	171	279	299	383	489	630	812	822	565	208	63	1
Male	148	225	208	252	301	350	390	384	219	79	19	1
Female	23	54	91	131	188	280	422	438	346	129	44	-
Alabama												
All races, both sexes	84	169	182	226	279	363	532	488	338	109	23	-
Male	71	121	130	143	171	212	256	197	104	25	5	-
Female	13	48	52	83	108	151	276	291	234	84	18	-
White, both sexes	51	122	141	177	224	295	450	439	289	88	18	-
Male	42	89	100	110	148	171	226	184	90	23	3	-
Female	9	33	41	67	76	124	224	255	199	65	15	-
Black, both sexes	33	47	40	48	54	68	80	49	49	21	5	-
Male	29	32	29	32	22	41	28	13	14	2	2	-
Female	4	15	11	16	32	27	52	36	35	19	3	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	2	-	-	-	-	-
Male	-	-	-	1	1	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Alaska												
All races, both sexes	17	27	26	23	27	36	32	31	16	8	-	-
Male	16	20	23	14	20	22	21	9	5	2	-	-
Female	1	7	3	9	7	14	11	22	11	6	-	-
White, both sexes	15	24	26	18	18	24	28	22	15	7	-	-
Male	14	18	23	12	12	17	18	6	5	2	-	-
Female	1	6	3	6	6	7	10	16	10	5	-	-
Black, both sexes	2	-	-	-	2	3	-	1	-	-	-	-
Male	2	-	-	-	2	-	-	-	-	-	-	-
Female	-	-	-	-	-	3	-	1	-	-	-	-
American Indian 1/, both sexes	-	3	-	5	6	9	4	6	-	1	-	-
Male	-	2	-	2	5	5	3	3	-	-	-	-
Female	-	1	-	3	1	4	1	3	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	2	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	1	-	-	-
Arizona												
All races, both sexes	194	270	406	403	493	721	927	999	662	249	41	1
Male	157	200	301	285	338	444	488	471	257	65	6	1
Female	37	70	105	118	155	277	439	528	405	184	35	-
White, both sexes	177	245	382	378	468	688	895	969	642	239	40	1
Male	144	186	284	273	323	426	473	458	251	63	6	1
Female	33	59	98	105	145	262	422	511	391	176	34	-
Black, both sexes	8	14	15	13	13	17	14	20	12	7	-	-
Male	4	7	12	6	7	6	8	9	4	1	-	-
Female	4	7	3	7	6	11	6	11	8	6	-	-
American Indian 1/, both sexes	8	9	7	10	11	12	15	7	5	2	-	-
Male	8	6	3	4	8	9	7	3	1	-	-	-
Female	-	3	4	6	3	3	8	4	4	2	-	-
Asian or Pacific Islander 2/, both sexes	1	2	2	2	1	4	3	3	3	1	1	-
Male	1	1	2	2	-	3	-	1	1	1	-	-
Female	-	1	-	-	1	1	3	2	2	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Arkansas												
All races, both sexes	43	76	97	135	175	229	315	351	252	99	26	-
Male	30	59	68	83	114	139	159	144	86	19	4	-
Female	13	17	29	52	61	90	156	207	166	80	22	-
White, both sexes	29	57	83	119	151	205	288	322	226	85	24	-
Male	21	46	58	75	101	127	150	133	76	17	3	-
Female	8	11	25	44	50	78	138	189	150	68	21	-
Black, both sexes	13	19	14	16	22	23	27	28	23	12	2	-
Male	8	13	10	8	13	12	9	10	8	1	1	-
Female	5	6	4	8	9	11	18	18	15	11	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	1	-	1	3	2	-	-
Male	-	-	-	-	-	-	-	1	2	1	-	-
Female	-	-	-	-	2	1	-	-	1	1	-	-
California												
All races, both sexes	1,042	1,437	1,845	1,894	2,473	3,502	5,153	5,986	4,357	1,996	481	1
Male	838	1,103	1,319	1,263	1,561	2,021	2,530	2,800	1,595	552	96	1
Female	204	334	526	631	912	1,481	2,623	3,186	2,762	1,444	385	-
White, both sexes	787	1,105	1,488	1,504	1,985	2,941	4,478	5,233	3,890	1,763	411	-
Male	639	859	1,072	1,027	1,268	1,707	2,223	2,479	1,437	473	80	-
Female	148	246	416	477	717	1,234	2,255	2,754	2,453	1,290	331	-
Black, both sexes	176	218	222	232	264	275	279	294	175	108	32	-
Male	126	151	152	135	160	152	120	118	47	27	6	-
Female	50	67	70	97	104	123	159	176	128	81	26	-
American Indian 1/, both sexes	8	7	16	13	17	20	16	19	8	6	1	-
Male	8	7	9	11	7	9	7	5	3	-	-	-
Female	-	-	7	2	10	11	9	14	5	6	1	-
Asian or Pacific Islander 2/, both sexes	71	107	119	145	207	266	380	440	284	119	37	1
Male	65	86	86	90	126	153	180	198	108	52	10	1
Female	6	21	33	55	81	113	200	242	176	67	27	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Colorado												
All races, both sexes	111	146	162	182	214	333	400	483	312	139	30	-
Male	92	112	118	120	148	199	216	227	123	44	8	-
Female	19	34	44	62	66	134	184	256	189	95	22	-
White, both sexes	107	135	154	175	201	316	392	469	306	137	30	-
Male	89	103	111	115	141	192	213	221	119	43	8	-
Female	18	32	43	60	60	124	179	248	187	94	22	-
Black, both sexes	2	9	6	7	11	14	6	10	5	2	-	-
Male	2	7	5	5	6	5	2	4	3	1	-	-
Female	-	2	1	2	5	9	4	6	2	1	-	-
American Indian 1/, both sexes	1	2	2	-	-	1	1	-	-	-	-	-
Male	-	2	2	-	-	1	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	2	2	1	4	1	-	-	-
Male	1	-	-	-	1	1	-	2	1	-	-	-
Female	-	-	-	-	1	1	1	2	-	-	-	-
Connecticut												
All races, both sexes	76	114	138	131	192	363	537	667	480	248	60	-
Male	56	91	101	86	125	216	290	303	167	61	7	-
Female	20	23	37	45	67	147	247	364	313	187	53	-
White, both sexes	62	102	114	110	178	343	505	650	468	245	57	-
Male	47	80	85	73	115	204	273	298	162	61	6	-
Female	15	22	29	37	63	139	232	352	306	184	51	-
Black, both sexes	12	10	20	19	13	18	29	16	10	3	3	-
Male	8	9	12	13	9	11	16	4	3	-	1	-
Female	4	1	8	6	4	7	13	12	7	3	2	-
American Indian 1/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	4	2	1	2	2	1	2	-	-	-
Male	1	1	4	-	1	1	1	1	2	-	-	-
Female	1	-	-	2	-	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Delaware												
All races, both sexes	35	46	61	47	82	115	162	138	124	63	10	1
Male	28	34	45	33	61	72	92	63	43	9	3	-
Female	7	12	16	14	21	43	70	75	81	54	7	1
White, both sexes	24	33	53	36	64	96	150	126	117	50	8	1
Male	19	26	39	25	50	60	89	56	42	7	2	-
Female	5	7	14	11	14	36	61	70	75	43	6	1
Black, both sexes	10	13	8	10	16	17	11	10	7	12	2	-
Male	8	8	6	7	9	11	3	5	1	1	1	-
Female	2	5	2	3	7	6	8	5	6	11	1	-
American Indian 1/, both sexes	1	-	-	1	1	1	1	-	-	1	-	-
Male	1	-	-	1	1	1	-	-	-	1	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	2	-	-	-	-
Male	-	-	-	-	1	-	-	2	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	22	44	75	89	81	107	120	105	82	45	15	-
Male	14	28	52	58	48	62	56	43	28	13	4	-
Female	8	16	23	31	33	45	64	62	54	32	11	-
White, both sexes	2	10	11	12	14	26	30	25	30	20	8	-
Male	2	7	10	9	10	19	22	15	12	8	1	-
Female	-	3	1	3	4	7	8	10	18	12	7	-
Black, both sexes	20	34	64	76	66	78	89	80	52	23	7	-
Male	12	21	42	48	37	43	34	28	16	4	3	-
Female	8	13	22	28	29	35	55	52	36	19	4	-
American Indian 1/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	3	-	-	-	2	-	-
Male	-	-	-	-	1	-	-	-	-	1	-	-
Female	-	-	-	-	-	3	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Florida												
All races, both sexes	652	828	1,110	1,293	1,664	2,740	3,813	4,350	3,172	1,384	310	3
Male	509	623	791	895	1,071	1,683	2,088	2,049	1,261	373	64	2
Female	143	205	319	398	593	1,057	1,725	2,301	1,911	1,011	246	1
White, both sexes	507	707	940	1,094	1,428	2,470	3,544	4,125	3,035	1,320	290	2
Male	417	538	680	769	942	1,547	1,965	1,966	1,216	366	61	1
Female	90	169	260	325	486	923	1,579	2,159	1,819	954	229	1
Black, both sexes	137	116	161	182	219	244	255	212	128	59	20	1
Male	85	80	106	114	120	123	118	77	42	6	3	1
Female	52	36	55	68	99	121	137	135	86	53	17	-
American Indian 1/, both sexes	1	2	2	3	2	2	-	4	2	-	-	-
Male	1	2	1	3	1	1	-	3	1	-	-	-
Female	-	-	1	-	1	1	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	7	3	7	14	15	24	14	9	7	5	-	-
Male	6	3	4	9	8	12	5	3	2	1	-	-
Female	1	-	3	5	7	12	9	6	5	4	-	-
Georgia												
All races, both sexes	232	308	357	399	478	572	714	764	526	201	31	-
Male	186	240	265	276	332	331	391	324	175	43	3	-
Female	46	68	92	123	146	241	323	440	351	158	28	-
White, both sexes	155	201	243	294	360	450	585	638	430	157	21	-
Male	128	159	190	213	265	269	334	280	147	38	3	-
Female	27	42	53	81	95	181	251	358	283	119	18	-
Black, both sexes	75	103	107	103	114	118	123	124	94	44	9	-
Male	56	77	72	62	63	61	55	43	27	5	-	-
Female	19	26	35	41	51	57	68	81	67	39	9	-
American Indian 1/, both sexes	-	-	2	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	4	5	2	4	3	6	2	2	-	1	-
Male	2	4	2	1	4	1	2	1	1	-	-	-
Female	-	-	3	1	-	2	4	1	1	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Hawaii												
All races, both sexes	31	55	53	72	71	90	125	128	94	26	6	-
Male	21	50	36	53	48	61	84	70	38	11	5	-
Female	10	5	17	19	23	29	41	58	56	15	1	-
White, both sexes	11	16	20	21	16	29	28	37	25	5	4	-
Male	8	15	17	17	13	20	19	21	11	2	3	-
Female	3	1	3	4	3	9	9	16	14	3	1	-
Black, both sexes	1	1	1	-	-	2	-	1	-	-	-	-
Male	1	1	-	-	-	1	-	1	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	19	38	32	51	55	59	97	90	69	20	2	-
Male	12	34	19	36	35	40	65	48	27	8	2	-
Female	7	4	13	15	20	19	32	42	42	12	-	-
Idaho												
All races, both sexes	16	25	27	39	58	99	120	160	128	40	7	-
Male	12	19	22	24	38	59	72	86	44	13	-	-
Female	4	6	5	15	20	40	48	74	84	27	7	-
White, both sexes	15	25	27	39	57	99	119	158	128	40	7	-
Male	11	19	22	24	38	59	71	85	44	13	-	-
Female	4	6	5	15	19	40	48	73	84	27	7	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	1	1	-	-	-	-
Male	1	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Illinois												
All races, both sexes	387	523	570	627	803	1,179	1,743	1,836	1,429	619	132	1
Male	301	387	403	462	507	658	912	784	467	149	24	-
Female	86	136	167	165	296	521	831	1,052	962	470	108	1
White, both sexes	269	367	425	481	621	996	1,543	1,680	1,336	583	123	1
Male	220	280	310	364	400	573	817	723	433	145	21	-
Female	49	87	115	117	221	423	726	957	903	438	102	1
Black, both sexes	112	148	137	138	173	165	177	139	77	35	8	-
Male	76	101	90	90	103	75	86	53	28	4	3	-
Female	36	47	47	48	70	90	91	86	49	31	5	-
American Indian 1/, both sexes	-	2	-	-	1	-	1	3	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	2	-	-	1	-	1	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	6	8	8	8	18	22	14	16	1	1	-
Male	5	6	3	8	4	10	9	7	6	-	-	-
Female	1	-	5	-	4	8	13	7	10	1	1	-
Indiana												
All races, both sexes	161	241	270	345	449	650	877	873	660	290	76	-
Male	121	177	184	216	295	369	456	382	218	58	7	-
Female	40	64	86	129	154	281	421	491	442	232	69	-
White, both sexes	131	210	233	317	413	597	816	821	628	275	74	-
Male	99	151	161	200	277	341	422	358	212	54	7	-
Female	32	59	72	117	136	256	394	463	416	221	67	-
Black, both sexes	30	31	36	27	34	52	59	51	32	15	2	-
Male	22	26	22	16	17	27	34	24	6	4	-	-
Female	8	5	14	11	17	25	25	27	26	11	2	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	2	1	2	-	-	-	-	-
Male	-	-	1	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	2	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Iowa												
All races, both sexes	97	124	135	178	275	398	566	671	624	272	67	-
Male	79	92	96	125	178	230	313	317	221	66	11	-
Female	18	32	39	53	97	168	253	354	403	206	56	-
White, both sexes	95	118	131	171	264	395	561	667	619	271	67	-
Male	78	91	94	121	170	227	311	315	218	66	11	-
Female	17	27	37	50	94	168	250	352	401	205	56	-
Black, both sexes	2	6	4	4	9	2	2	3	3	1	-	-
Male	1	1	2	1	7	2	1	2	2	-	-	-
Female	1	5	2	3	2	-	1	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	2	1	1	1	1	1	-	-	-
Male	-	-	-	2	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	2	-	1	-	-	-
Male	-	-	-	1	1	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Kansas												
All races, both sexes	89	85	105	145	172	256	385	403	302	148	43	-
Male	66	62	80	103	110	157	213	195	109	37	5	-
Female	23	23	25	42	62	99	172	208	193	111	38	-
White, both sexes	82	73	99	128	162	233	370	391	289	144	43	-
Male	61	54	76	89	108	145	207	188	103	37	5	-
Female	21	19	23	39	54	88	163	203	186	107	38	-
Black, both sexes	7	10	3	11	8	22	13	10	11	4	-	-
Male	5	6	2	10	2	11	5	5	5	-	-	-
Female	2	4	1	1	6	11	8	5	6	4	-	-
American Indian 1/, both sexes	-	2	3	5	-	1	2	2	2	-	-	-
Male	-	2	2	3	-	1	1	2	1	-	-	-
Female	-	-	1	2	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	2	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Kentucky												
All races, both sexes	113	149	161	249	271	393	520	492	381	145	28	-
Male	74	115	116	168	163	219	265	205	114	34	6	-
Female	39	34	45	81	108	174	255	287	267	111	22	-
White, both sexes	104	135	147	234	255	376	500	474	362	143	27	-
Male	70	105	105	157	154	213	260	196	109	34	6	-
Female	34	30	42	77	101	163	240	278	253	109	21	-
Black, both sexes	9	14	13	15	15	17	18	18	19	2	1	-
Male	4	10	10	11	9	6	5	9	5	-	-	-
Female	5	4	3	4	6	11	13	9	14	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	-	2	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	2	-	-	-	-	-
Louisiana												
All races, both sexes	140	199	200	232	302	355	539	499	309	125	26	-
Male	100	155	145	149	176	212	284	211	99	32	7	-
Female	40	44	55	83	126	143	255	288	210	93	19	-
White, both sexes	89	132	134	170	222	281	429	404	257	101	15	-
Male	68	105	101	109	134	174	235	173	86	26	3	-
Female	21	27	33	61	88	107	194	231	171	75	12	-
Black, both sexes	51	66	66	62	78	73	108	94	52	24	11	-
Male	32	49	44	40	41	37	49	38	13	6	4	-
Female	19	17	22	22	37	36	59	56	39	18	7	-
American Indian 1/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	2	-	1	1	-	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Maine												
All races, both sexes	35	41	56	57	97	148	165	199	161	74	17	-
Male	29	32	42	45	71	97	89	95	62	16	4	-
Female	6	9	14	12	26	51	76	104	99	58	13	-
White, both sexes	35	40	56	56	97	146	164	199	161	74	17	-
Male	29	31	42	45	71	95	89	95	62	16	4	-
Female	6	9	14	11	26	51	75	104	99	58	13	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Maryland												
All races, both sexes	215	262	329	337	478	602	888	975	701	332	75	-
Male	157	183	236	222	294	355	435	409	209	73	11	-
Female	58	79	93	115	184	247	453	566	492	259	64	-
White, both sexes	140	142	212	216	330	453	700	795	590	281	60	-
Male	103	104	155	152	214	279	351	334	186	67	9	-
Female	37	38	57	64	116	174	349	461	404	214	51	-
Black, both sexes	74	114	112	117	137	138	177	169	106	49	14	-
Male	53	74	77	68	73	71	79	68	20	6	2	-
Female	21	40	35	49	64	67	98	101	86	43	12	-
American Indian 1/, both sexes	-	-	-	-	1	1	2	1	-	-	-	-
Male	-	-	-	-	1	-	1	1	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	6	5	4	10	10	9	10	5	2	1	-
Male	1	5	4	2	6	5	4	6	3	-	-	-
Female	-	1	1	2	4	5	5	4	2	2	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Massachusetts												
All races, both sexes	169	219	293	304	447	634	963	1,100	843	342	80	1
Male	136	172	216	212	298	367	510	500	280	81	11	1
Female	33	47	77	92	149	267	453	600	563	261	69	-
White, both sexes	157	204	271	280	416	608	937	1,073	828	336	80	1
Male	128	165	204	196	276	354	496	492	275	81	11	1
Female	29	39	67	84	140	254	441	581	553	255	69	-
Black, both sexes	9	11	19	17	23	21	18	23	13	5	-	-
Male	6	6	10	11	14	12	10	7	4	-	-	-
Female	3	5	9	6	9	9	8	16	9	5	-	-
American Indian 1/, both sexes	2	1	2	-	-	1	-	-	-	1	-	-
Male	1	1	2	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	3	1	7	8	4	8	4	2	-	-	-
Male	1	-	-	5	8	1	4	1	1	-	-	-
Female	-	3	1	2	-	3	4	3	1	-	-	-
Michigan												
All races, both sexes	499	643	785	812	950	1,448	1,924	2,141	1,495	623	136	-
Male	380	472	562	552	591	829	1,000	970	516	143	26	-
Female	119	171	223	260	359	619	924	1,171	979	480	110	-
White, both sexes	356	443	595	620	755	1,197	1,675	1,937	1,370	581	122	-
Male	287	338	429	434	486	698	893	877	473	134	22	-
Female	69	105	166	186	269	499	782	1,060	897	447	100	-
Black, both sexes	139	196	183	186	184	239	232	195	118	38	12	-
Male	90	131	127	114	97	127	100	90	41	9	3	-
Female	49	65	56	72	87	112	132	105	77	29	9	-
American Indian 1/, both sexes	3	3	5	3	4	9	7	6	4	1	1	-
Male	3	2	5	2	3	3	4	3	2	-	1	-
Female	-	1	-	1	1	6	3	3	2	1	-	-
Asian or Pacific Islander 2/, both sexes	1	1	2	3	7	3	10	3	3	3	1	-
Male	-	1	1	2	5	1	3	-	-	-	-	-
Female	1	-	1	1	2	2	7	3	3	3	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Minnesota												
All races, both sexes	115	143	160	154	204	334	465	601	483	197	32	-
Male	93	124	115	114	142	216	276	309	196	46	4	-
Female	22	19	45	40	62	118	189	292	287	151	28	-
White, both sexes	109	130	151	144	193	321	455	599	482	195	32	-
Male	89	113	110	108	134	210	272	309	196	46	4	-
Female	20	17	41	36	59	111	183	290	286	149	28	-
Black, both sexes	4	6	4	3	4	2	7	1	-	1	-	-
Male	2	5	3	1	2	-	4	-	-	-	-	-
Female	2	1	1	2	2	2	3	1	-	1	-	-
American Indian 1/, both sexes	2	4	4	6	5	7	2	-	-	1	-	-
Male	2	3	1	4	4	4	-	-	-	-	-	-
Female	-	1	3	2	1	3	2	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	3	1	1	2	4	1	1	1	-	-	-
Male	-	3	1	1	2	2	-	-	-	-	-	-
Female	-	-	-	-	-	2	1	1	1	-	-	-
Mississippi												
All races, both sexes	113	147	150	157	235	278	346	322	243	91	31	-
Male	83	108	102	109	125	159	159	116	65	14	3	-
Female	30	39	48	48	110	119	187	206	178	77	28	-
White, both sexes	64	83	98	102	174	209	274	269	200	73	19	-
Male	47	69	70	73	99	129	124	98	51	11	1	-
Female	17	14	28	29	75	80	150	171	149	62	18	-
Black, both sexes	47	64	52	53	61	68	71	52	43	18	11	-
Male	34	39	32	34	26	29	35	17	14	3	1	-
Female	13	25	20	19	35	39	36	35	29	15	10	-
American Indian 1/, both sexes	2	-	-	2	-	-	1	-	-	-	-	-
Male	2	-	-	2	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	-	-	1	-
Male	-	-	-	-	-	1	-	1	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Missouri												
All races, both sexes	203	237	280	332	439	645	892	1,055	734	369	91	1
Male	152	180	206	218	272	381	477	435	232	71	14	1
Female	51	57	74	114	167	264	415	620	502	298	77	-
White, both sexes	173	186	243	281	375	588	817	975	691	342	74	1
Male	128	143	180	192	231	351	442	416	222	65	11	1
Female	45	43	63	89	144	237	375	559	469	277	63	-
Black, both sexes	27	51	36	50	59	54	74	77	40	27	17	-
Male	22	37	25	26	38	29	34	18	10	6	3	-
Female	5	14	11	24	21	25	40	59	30	21	14	-
American Indian 1/, both sexes	3	-	1	1	2	1	-	1	1	-	-	-
Male	2	-	1	-	2	-	-	-	-	-	-	-
Female	1	-	-	1	-	1	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	3	2	1	2	2	-	-	-
Male	-	-	-	-	1	1	1	1	-	-	-	-
Female	-	-	-	-	2	1	-	1	2	-	-	-
Montana												
All races, both sexes	36	41	65	51	67	97	127	125	85	31	5	-
Male	29	36	46	38	47	68	71	76	40	12	1	-
Female	7	5	19	13	20	29	56	49	45	19	4	-
White, both sexes	34	37	63	49	64	93	123	125	80	30	4	-
Male	27	33	44	36	45	66	70	76	37	11	1	-
Female	7	4	19	13	19	27	53	49	43	19	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	4	2	2	3	4	4	-	5	1	1	-
Male	2	3	2	2	2	2	1	-	3	1	-	-
Female	-	1	-	-	1	2	3	-	2	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Nebraska												
All races, both sexes	31	37	50	64	88	139	221	297	240	118	24	-
Male	27	24	40	47	54	92	138	147	81	33	1	-
Female	4	13	10	17	34	47	83	150	159	85	23	-
White, both sexes	28	32	43	59	79	132	217	290	238	118	23	-
Male	24	21	36	44	49	88	136	141	81	33	1	-
Female	4	11	7	15	30	44	81	149	157	85	22	-
Black, both sexes	3	5	4	5	6	6	2	6	2	-	1	-
Male	3	3	3	3	3	3	1	5	-	-	-	-
Female	-	2	1	2	3	3	1	1	2	-	1	-
American Indian 1/, both sexes	-	-	1	-	2	1	1	-	-	-	-	-
Male	-	-	-	-	2	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	1	-	1	1	-	-	-	-
Male	-	-	1	-	-	-	1	1	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-
Nevada												
All races, both sexes	103	154	186	209	214	255	297	225	124	43	4	-
Male	81	119	145	147	136	162	182	117	49	14	1	-
Female	22	35	41	62	78	93	115	108	75	29	3	-
White, both sexes	89	129	160	186	183	236	277	208	115	39	4	-
Male	71	98	127	135	114	150	171	111	46	12	1	-
Female	18	31	33	51	69	86	106	97	69	27	3	-
Black, both sexes	4	17	17	13	20	10	12	10	4	3	-	-
Male	1	14	10	8	14	7	5	3	1	2	-	-
Female	3	3	7	5	6	3	7	7	3	1	-	-
American Indian 1/, both sexes	3	1	2	1	2	4	4	2	-	-	-	-
Male	3	1	2	-	2	3	4	-	-	-	-	-
Female	-	-	-	1	-	1	-	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	7	7	9	9	5	4	5	5	1	-	-
Male	6	6	6	4	6	2	2	3	2	-	-	-
Female	1	1	1	5	3	3	2	2	3	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
New Hampshire												
All races, both sexes	32	45	83	69	106	132	208	241	193	83	16	-
Male	28	32	55	51	70	82	105	102	62	19	3	-
Female	4	13	28	18	36	50	103	139	131	64	13	-
White, both sexes	32	45	82	68	105	131	208	241	193	83	16	-
Male	28	32	55	50	70	82	105	102	62	19	3	-
Female	4	13	27	18	35	49	103	139	131	64	13	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
New Jersey												
All races, both sexes	201	301	363	506	634	1,066	1,595	1,827	1,408	658	147	-
Male	161	233	273	340	394	620	848	794	460	153	29	-
Female	40	68	90	166	240	446	747	1,033	948	505	118	-
White, both sexes	154	236	291	408	523	931	1,459	1,698	1,333	613	136	-
Male	125	190	224	280	325	547	777	743	441	150	28	-
Female	29	46	67	128	198	384	682	955	892	463	108	-
Black, both sexes	41	55	59	85	93	110	118	109	57	38	10	-
Male	30	34	40	50	57	62	60	41	16	2	1	-
Female	11	21	19	35	36	48	58	68	41	36	9	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	1	2	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	2	1	-	-
Asian or Pacific Islander 2/, both sexes	6	10	13	13	18	24	18	19	16	6	1	-
Male	6	9	9	10	12	11	11	10	3	1	-	-
Female	-	1	4	3	6	13	7	9	13	5	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
New Mexico												
All races, both sexes	50	85	126	109	150	201	257	277	213	94	27	-
Male	37	69	91	74	90	115	146	133	92	24	7	-
Female	13	16	35	35	60	86	111	144	121	70	20	-
White, both sexes	45	75	114	104	137	188	241	263	202	91	26	-
Male	33	63	83	69	84	109	137	126	87	22	7	-
Female	12	12	31	35	53	79	104	137	115	69	19	-
Black, both sexes	2	2	6	1	3	3	5	3	4	-	1	-
Male	2	1	4	1	1	2	2	1	1	-	-	-
Female	-	1	2	-	2	1	3	2	3	-	1	-
American Indian 1/, both sexes	3	7	6	4	10	9	11	8	6	3	-	-
Male	2	4	4	4	5	4	7	4	3	2	-	-
Female	1	3	2	-	5	5	4	4	3	1	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	-	3	1	-	-	-
Male	-	1	-	-	-	-	-	2	1	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
New York												
All races, both sexes	640	949	1,272	1,640	2,120	3,505	5,136	5,651	4,772	2,347	544	-
Male	461	680	859	1,072	1,247	1,943	2,468	2,243	1,542	492	94	-
Female	179	269	413	568	873	1,562	2,668	3,408	3,230	1,855	450	-
White, both sexes	434	683	908	1,167	1,621	2,846	4,364	4,933	4,320	2,116	474	-
Male	325	522	646	791	987	1,624	2,159	1,997	1,419	457	81	-
Female	109	161	262	376	634	1,222	2,205	2,936	2,901	1,659	393	-
Black, both sexes	186	223	319	399	428	548	634	598	356	194	56	-
Male	117	124	178	230	214	257	247	189	85	24	9	-
Female	69	99	141	169	214	291	387	409	271	170	47	-
American Indian 1/, both sexes	2	2	2	5	3	1	5	5	-	1	-	-
Male	2	1	2	2	3	-	2	3	-	-	-	-
Female	-	1	-	3	-	1	3	2	-	1	-	-
Asian or Pacific Islander 2/, both sexes	18	41	43	69	68	110	133	115	96	36	14	-
Male	17	33	33	49	43	62	60	54	38	11	4	-
Female	1	8	10	20	25	48	73	61	58	25	10	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
North Carolina												
All races, both sexes	346	436	572	564	710	913	1,141	1,177	804	308	77	2
Male	248	330	421	392	465	539	603	481	242	75	7	1
Female	98	106	151	172	245	374	538	696	562	233	70	1
White, both sexes	240	304	429	430	569	746	950	1,022	691	252	57	-
Male	172	234	330	301	381	452	517	426	217	62	5	-
Female	68	70	99	129	188	294	433	596	474	190	52	-
Black, both sexes	101	122	136	121	133	154	178	144	109	53	20	2
Male	72	87	84	81	78	77	78	49	25	12	2	1
Female	29	35	52	40	55	77	100	95	84	41	18	1
American Indian 1/, both sexes	3	6	7	10	7	9	11	8	3	2	-	-
Male	2	5	7	8	5	7	6	4	-	1	-	-
Female	1	1	-	2	2	2	5	4	3	1	-	-
Asian or Pacific Islander 2/, both sexes	2	4	-	3	1	4	2	3	1	1	-	-
Male	2	4	-	2	1	3	2	2	-	-	-	-
Female	-	-	-	1	-	1	-	1	1	1	-	-
North Dakota												
All races, both sexes	24	12	21	37	44	68	82	123	119	59	14	-
Male	19	10	15	26	32	42	43	60	49	14	2	-
Female	5	2	6	11	12	26	39	63	70	45	12	-
White, both sexes	22	11	20	36	43	66	80	122	118	59	14	-
Male	17	10	15	26	32	41	42	59	49	14	2	-
Female	5	1	5	10	11	25	38	63	69	45	12	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	1	1	1	2	2	1	1	-	-	-
Male	1	-	-	-	-	1	1	1	-	-	-	-
Female	-	1	1	1	1	1	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Ohio												
All races, both sexes	414	574	725	751	1,027	1,524	2,208	2,450	1,730	689	150	-
Male	309	422	524	528	648	874	1,144	1,052	570	165	20	-
Female	105	152	201	223	379	650	1,064	1,398	1,160	524	130	-
White, both sexes	340	456	620	652	876	1,380	2,050	2,307	1,640	637	136	-
Male	262	342	454	466	557	801	1,070	990	546	150	19	-
Female	78	114	166	186	319	579	980	1,317	1,094	487	117	-
Black, both sexes	72	111	99	97	148	138	152	140	87	52	12	-
Male	45	77	65	61	89	68	71	60	22	15	1	-
Female	27	34	34	36	59	70	81	80	65	37	11	-
American Indian 1/, both sexes	-	1	1	1	-	1	1	-	-	-	1	-
Male	-	-	1	1	-	1	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	2	6	5	1	3	5	5	3	3	-	1	-
Male	2	3	4	-	2	4	2	2	2	-	-	-
Female	-	3	1	1	1	1	3	1	1	-	1	-
Oklahoma												
All races, both sexes	148	193	247	286	341	503	667	665	488	204	46	-
Male	109	135	157	187	208	279	351	279	185	42	9	-
Female	39	58	90	99	133	224	316	386	303	162	37	-
White, both sexes	125	160	208	242	293	431	610	626	455	188	39	-
Male	93	109	135	155	183	241	327	265	175	39	7	-
Female	32	51	73	87	110	190	283	361	280	149	32	-
Black, both sexes	11	20	16	22	23	36	29	20	18	9	5	-
Male	8	18	10	15	12	17	12	5	9	2	2	-
Female	3	2	6	7	11	19	17	15	9	7	3	-
American Indian 1/, both sexes	11	12	21	21	22	34	26	18	15	6	2	-
Male	7	8	11	17	12	20	12	8	1	1	-	-
Female	4	4	10	4	10	14	14	10	14	5	2	-
Asian or Pacific Islander 2/, both sexes	1	1	2	1	3	2	2	1	-	1	-	-
Male	1	-	1	-	1	1	-	1	-	-	-	-
Female	-	1	1	1	2	1	2	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Oregon												
All races, both sexes	93	150	165	157	224	303	415	472	385	145	23	1
Male	78	112	121	121	155	194	265	232	157	40	3	1
Female	15	38	44	36	69	109	150	240	228	105	20	-
White, both sexes	88	140	161	149	217	294	401	465	382	140	23	1
Male	74	104	118	116	149	190	257	230	156	39	3	1
Female	14	36	43	33	68	104	144	235	226	101	20	-
Black, both sexes	3	6	-	3	2	2	4	1	1	2	-	-
Male	2	5	-	2	2	2	3	-	-	-	-	-
Female	1	1	-	1	-	-	1	1	1	2	-	-
American Indian 1/, both sexes	1	1	1	4	1	2	4	2	-	1	-	-
Male	1	1	1	2	1	-	1	1	-	-	-	-
Female	-	-	-	2	-	2	3	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	3	3	1	4	5	6	4	2	2	-	-
Male	1	2	2	1	3	2	4	1	1	1	-	-
Female	-	1	1	-	1	3	2	3	1	1	-	-
Pennsylvania												
All races, both sexes	320	506	661	705	1,000	1,681	2,456	2,825	2,086	835	181	3
Male	256	386	456	480	651	960	1,290	1,240	687	221	33	2
Female	64	120	205	225	349	721	1,166	1,585	1,399	614	148	1
White, both sexes	244	404	540	600	869	1,521	2,299	2,677	1,985	791	169	2
Male	200	316	385	408	573	881	1,218	1,189	666	210	30	1
Female	44	88	155	192	296	640	1,081	1,488	1,319	581	139	1
Black, both sexes	73	98	119	96	126	151	144	136	98	43	12	1
Male	53	68	71	65	74	74	67	44	19	10	3	1
Female	20	30	48	31	52	77	77	92	79	33	9	-
American Indian 1/, both sexes	-	-	-	1	1	1	1	1	-	-	-	-
Male	-	-	-	1	1	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	4	2	8	4	8	12	11	3	1	-	-
Male	3	2	-	6	3	4	4	7	2	1	-	-
Female	-	2	2	2	1	4	8	4	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Rhode Island												
All races, both sexes	37	61	53	57	76	146	241	323	297	145	26	-
Male	33	50	39	40	44	85	101	130	82	33	1	-
Female	4	11	14	17	32	61	140	193	215	112	25	-
White, both sexes	33	58	46	55	68	137	236	314	295	144	26	-
Male	29	47	35	39	40	81	100	127	82	32	1	-
Female	4	11	11	16	28	56	136	187	213	112	25	-
Black, both sexes	2	3	6	2	6	7	5	9	2	1	-	-
Male	2	3	4	1	3	3	1	3	-	1	-	-
Female	-	-	2	1	3	4	4	6	2	-	-	-
American Indian 1/, both sexes	1	-	1	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	2	-	-	-	-	-	-
Male	1	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
South Carolina												
All races, both sexes	112	142	202	212	264	363	482	427	296	109	17	1
Male	84	107	155	154	164	227	253	188	92	23	4	1
Female	28	35	47	58	100	136	229	239	204	86	13	-
White, both sexes	78	93	146	163	188	288	397	352	252	89	11	1
Male	61	72	112	118	125	187	219	163	80	19	3	1
Female	17	21	34	45	63	101	178	189	172	70	8	-
Black, both sexes	32	48	56	45	73	75	84	75	43	20	6	-
Male	21	35	43	33	36	40	33	25	12	4	1	-
Female	11	13	13	12	37	35	51	50	31	16	5	-
American Indian 1/, both sexes	1	1	-	1	1	-	-	-	-	-	-	-
Male	1	-	-	1	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	3	2	-	1	-	1	-	-	-
Male	1	-	-	2	2	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
South Dakota												
All races, both sexes	17	23	23	37	40	62	115	151	121	61	15	-
Male	14	18	16	26	29	42	65	82	50	18	1	-
Female	3	5	7	11	11	20	50	69	71	43	14	-
White, both sexes	14	20	22	34	36	61	110	147	119	60	14	-
Male	13	16	16	25	27	41	63	80	50	18	1	-
Female	1	4	6	9	9	20	47	67	69	42	13	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	3	2	1	3	4	1	5	4	2	1	-	-
Male	1	2	-	1	2	1	2	2	-	-	-	-
Female	2	-	1	2	2	-	3	2	2	1	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	284	352	406	463	539	748	969	995	612	272	51	-
Male	198	254	278	302	316	441	472	405	176	58	6	-
Female	86	98	128	161	223	307	497	590	436	214	45	-
White, both sexes	225	283	338	399	451	662	871	902	557	245	47	-
Male	159	204	237	261	272	388	422	367	164	54	6	-
Female	66	79	101	138	179	274	449	535	393	191	41	-
Black, both sexes	57	69	67	62	86	85	96	91	54	27	4	-
Male	39	50	41	39	42	52	48	36	12	4	-	-
Female	18	19	26	23	44	33	48	55	42	23	4	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	1	2	2	1	1	2	1	-	-	-
Male	-	-	-	2	2	1	1	2	-	-	-	-
Female	2	-	1	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Texas												
All races, both sexes	612	847	1,079	1,064	1,265	1,791	2,303	2,382	1,626	644	121	-
Male	443	647	780	735	827	1,033	1,217	1,056	542	173	16	-
Female	169	200	299	329	438	758	1,086	1,326	1,084	471	105	-
White, both sexes	473	697	890	886	1,098	1,591	2,099	2,211	1,505	579	106	-
Male	343	538	650	631	727	929	1,120	995	508	158	12	-
Female	130	159	240	255	371	662	979	1,216	997	421	94	-
Black, both sexes	130	143	175	158	157	183	193	150	111	64	13	-
Male	93	105	120	91	96	97	93	53	30	15	2	-
Female	37	38	55	67	61	86	100	97	81	49	11	-
American Indian 1/, both sexes	1	1	1	8	2	1	-	2	2	-	-	-
Male	1	1	-	7	1	1	-	-	-	-	-	-
Female	-	-	1	1	1	-	-	2	2	-	-	-
Asian or Pacific Islander 2/, both sexes	8	6	13	12	8	16	11	19	8	1	2	-
Male	6	3	10	6	3	6	4	8	4	-	2	-
Female	2	3	3	6	5	10	7	11	4	1	-	-
Utah												
All races, both sexes	34	63	52	53	75	97	174	176	100	55	3	-
Male	25	50	38	32	56	59	97	100	46	18	-	-
Female	9	13	14	21	19	38	77	76	54	37	3	-
White, both sexes	32	58	48	51	74	96	171	175	100	55	3	-
Male	23	46	35	32	56	58	97	99	46	18	-	-
Female	9	12	13	19	18	38	74	76	54	37	3	-
Black, both sexes	1	-	1	-	-	1	1	-	-	-	-	-
Male	1	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	1	1	1	2	1	-	1	1	-	-	-	-
Male	1	1	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	2	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	4	2	-	-	-	1	-	-	-	-	-
Male	-	3	1	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Vermont												
All races, both sexes	27	30	40	31	44	76	103	112	84	38	10	-
Male	21	20	29	18	29	53	52	51	30	12	1	-
Female	6	10	11	13	15	23	51	61	54	26	9	-
White, both sexes	27	30	40	30	44	76	101	112	84	38	10	-
Male	21	20	29	18	29	53	51	51	30	12	1	-
Female	6	10	11	12	15	23	50	61	54	26	9	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	179	249	263	355	428	598	807	787	523	210	43	-
Male	136	187	182	240	285	365	420	352	189	54	8	-
Female	43	62	81	115	143	233	387	435	334	156	35	-
White, both sexes	125	176	183	267	358	467	693	681	440	182	27	-
Male	99	134	131	192	245	283	375	312	158	47	6	-
Female	26	42	52	75	113	184	318	369	282	135	21	-
Black, both sexes	50	67	75	84	62	119	103	93	75	28	16	-
Male	34	48	47	46	36	75	40	34	27	7	2	-
Female	16	19	28	38	26	44	63	59	48	21	14	-
American Indian 1/, both sexes	1	-	-	-	-	1	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	6	5	4	8	11	9	13	8	-	-	-
Male	3	5	4	2	4	7	4	6	4	-	-	-
Female	-	1	1	2	4	4	5	7	4	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Washington												
All races, both sexes	155	209	294	290	437	609	844	995	765	311	66	1
Male	125	161	213	212	283	367	461	493	310	82	16	1
Female	30	48	81	78	154	242	383	502	455	229	50	-
White, both sexes	142	188	269	259	400	559	784	957	742	303	65	1
Male	113	143	192	191	257	343	437	477	300	80	16	1
Female	29	45	77	68	143	216	347	480	442	223	49	-
Black, both sexes	7	7	10	13	15	14	17	8	7	1	1	-
Male	7	6	9	8	13	7	8	5	4	1	-	-
Female	-	1	1	5	2	7	9	3	3	-	1	-
American Indian 1/, both sexes	3	5	4	8	6	10	6	3	-	3	-	-
Male	2	4	4	5	3	5	3	1	-	-	-	-
Female	1	1	-	3	3	5	3	2	-	3	-	-
Asian or Pacific Islander 2/, both sexes	3	9	11	10	16	26	37	27	16	4	-	-
Male	3	8	8	8	10	12	13	10	6	1	-	-
Female	-	1	3	2	6	14	24	17	10	3	-	-
West Virginia												
All races, both sexes	102	112	129	193	200	289	411	399	286	105	18	-
Male	81	86	92	124	126	156	235	160	85	14	3	-
Female	21	26	37	69	74	133	176	239	201	91	15	-
White, both sexes	98	111	123	189	193	281	407	383	275	101	18	-
Male	77	85	86	120	122	152	234	156	82	13	3	-
Female	21	26	37	69	71	129	173	227	193	88	15	-
Black, both sexes	4	1	6	3	7	8	3	14	11	4	-	-
Male	4	1	6	3	4	4	1	4	3	1	-	-
Female	-	-	-	-	3	4	2	10	8	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	1	2	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Wisconsin												
All races, both sexes	123	180	199	227	300	474	689	776	618	250	57	-
Male	93	144	151	168	203	291	417	345	236	65	5	-
Female	30	36	48	59	97	183	272	431	382	185	52	-
White, both sexes	101	148	181	205	279	452	669	757	606	245	55	-
Male	77	126	139	156	194	280	410	337	233	64	5	-
Female	24	22	42	49	85	172	259	420	373	181	50	-
Black, both sexes	21	26	14	15	17	17	13	14	9	5	1	-
Male	15	13	9	10	8	8	3	5	1	1	-	-
Female	6	13	5	5	9	9	10	9	8	4	1	-
American Indian 1/, both sexes	1	5	2	5	2	5	5	2	1	-	1	-
Male	1	4	2	2	1	3	2	2	1	-	-	-
Female	-	1	-	3	1	2	3	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	1	2	2	2	-	2	3	2	-	-	-
Male	-	1	1	-	-	-	2	1	1	-	-	-
Female	-	-	1	2	2	-	-	2	1	-	-	-
Wyoming												
All races, both sexes	12	19	13	28	24	39	62	47	31	4	3	-
Male	8	15	9	17	19	24	36	23	12	1	2	-
Female	4	4	4	11	5	15	26	24	19	3	1	-
White, both sexes	12	17	13	27	24	39	59	47	31	4	3	-
Male	8	14	9	16	19	24	35	23	12	1	2	-
Female	4	3	4	11	5	15	24	24	19	3	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	2	-	1	-	-	2	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
United States												
All races, both sexes	59,051	1	-	-	-	3	14	70	141	396	999	2,191
Male	33,401	1	-	-	-	2	10	57	119	310	799	1,713
Female	25,650	-	-	-	-	1	4	13	22	86	200	478
White, both sexes	48,660	1	-	-	-	2	8	44	100	287	762	1,677
Male	27,365	1	-	-	-	1	6	34	83	229	622	1,340
Female	21,295	-	-	-	-	1	2	10	17	58	140	337
Black, both sexes	8,921	-	-	-	-	1	5	21	33	91	188	455
Male	5,155	-	-	-	-	1	4	18	29	66	137	325
Female	3,766	-	-	-	-	-	1	3	4	25	51	130
American Indian 1/, both sexes	325	-	-	-	-	-	-	-	4	2	11	20
Male	200	-	-	-	-	-	-	-	3	2	8	14
Female	125	-	-	-	-	-	-	-	1	-	3	6
Asian or Pacific Islander 2/, both sexes	1,145	-	-	-	-	-	1	5	4	16	38	39
Male	681	-	-	-	-	-	-	5	4	13	32	34
Female	464	-	-	-	-	-	1	-	-	3	6	5
Alabama												
All races, both sexes	585	-	-	-	-	-	-	1	-	5	11	28
Male	298	-	-	-	-	-	-	1	-	3	8	23
Female	287	-	-	-	-	-	-	-	-	2	3	5
White, both sexes	466	-	-	-	-	-	-	-	-	5	5	20
Male	242	-	-	-	-	-	-	-	-	3	4	18
Female	224	-	-	-	-	-	-	-	-	2	1	2
Black, both sexes	118	-	-	-	-	-	-	1	-	-	6	8
Male	55	-	-	-	-	-	-	1	-	-	4	5
Female	63	-	-	-	-	-	-	-	-	-	2	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Alaska												
All races, both sexes	109	-	-	-	-	-	-	-	-	1	6	5
Male	83	-	-	-	-	-	-	-	-	-	6	5
Female	26	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	91	-	-	-	-	-	-	-	-	-	6	4
Male	72	-	-	-	-	-	-	-	-	-	6	4
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
Arizona												
All races, both sexes	1,772	-	-	-	-	-	-	-	6	11	17	52
Male	1,076	-	-	-	-	-	-	-	5	9	12	39
Female	696	-	-	-	-	-	-	-	1	2	5	13
White, both sexes	1,660	-	-	-	-	-	-	-	6	8	16	43
Male	1,013	-	-	-	-	-	-	-	5	7	12	34
Female	647	-	-	-	-	-	-	-	1	1	4	9
Black, both sexes	66	-	-	-	-	-	-	-	-	2	-	7
Male	32	-	-	-	-	-	-	-	-	1	-	3
Female	34	-	-	-	-	-	-	-	-	1	-	4
American Indian 1/, both sexes	34	-	-	-	-	-	-	-	-	-	1	-
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	1	-	2
Male	9	-	-	-	-	-	-	-	-	1	-	2
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Arkansas												
All races, both sexes	251	-	-	-	-	-	-	-	5	6	12	14
Male	144	-	-	-	-	-	-	-	5	3	10	12
Female	107	-	-	-	-	-	-	-	-	3	2	2
White, both sexes	206	-	-	-	-	-	-	-	5	5	7	11
Male	110	-	-	-	-	-	-	-	5	2	6	9
Female	96	-	-	-	-	-	-	-	-	3	1	2
Black, both sexes	41	-	-	-	-	-	-	-	-	1	5	3
Male	33	-	-	-	-	-	-	-	-	1	4	3
Female	8	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	7,243	-	-	-	-	-	1	7	9	24	76	181
Male	4,177	-	-	-	-	-	1	6	9	17	66	155
Female	3,066	-	-	-	-	-	-	1	-	7	10	26
White, both sexes	5,876	-	-	-	-	-	1	3	7	16	54	140
Male	3,375	-	-	-	-	-	1	3	7	10	48	119
Female	2,501	-	-	-	-	-	-	-	-	6	6	21
Black, both sexes	829	-	-	-	-	-	-	3	1	5	12	26
Male	496	-	-	-	-	-	-	2	1	5	8	23
Female	333	-	-	-	-	-	-	1	-	-	4	3
American Indian 1/, both sexes	36	-	-	-	-	-	-	-	-	-	1	3
Male	24	-	-	-	-	-	-	-	-	-	1	3
Female	12	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	502	-	-	-	-	-	-	1	1	3	9	12
Male	282	-	-	-	-	-	-	1	1	2	9	10
Female	220	-	-	-	-	-	-	-	-	1	-	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Colorado												
All races, both sexes	624	-	-	-	-	-	1	3	3	5	19	24
Male	372	-	-	-	-	-	1	1	3	4	17	19
Female	252	-	-	-	-	-	-	2	-	1	2	5
White, both sexes	590	-	-	-	-	-	1	2	2	5	18	23
Male	350	-	-	-	-	-	1	-	2	4	16	18
Female	240	-	-	-	-	-	-	2	-	1	2	5
Black, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	1	-	1	-
Male	2	-	-	-	-	-	-	-	1	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	1	-	-	-	1
Male	3	-	-	-	-	-	-	1	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	623	-	-	-	-	-	-	-	-	2	7	24
Male	361	-	-	-	-	-	-	-	-	2	6	18
Female	262	-	-	-	-	-	-	-	-	-	1	6
White, both sexes	564	-	-	-	-	-	-	-	-	1	5	18
Male	318	-	-	-	-	-	-	-	-	1	5	15
Female	246	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes	55	-	-	-	-	-	-	-	-	1	2	6
Male	40	-	-	-	-	-	-	-	-	1	1	3
Female	15	-	-	-	-	-	-	-	-	-	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Delaware												
All races, both sexes	315	-	-	-	-	-	-	1	1	5	8	20
Male	191	-	-	-	-	-	-	1	1	4	6	18
Female	124	-	-	-	-	-	-	-	-	1	2	2
White, both sexes	254	-	-	-	-	-	-	1	1	4	8	12
Male	148	-	-	-	-	-	-	1	1	3	6	11
Female	106	-	-	-	-	-	-	-	-	1	2	1
Black, both sexes	58	-	-	-	-	-	-	-	-	1	-	8
Male	40	-	-	-	-	-	-	-	-	1	-	7
Female	18	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	545	-	-	-	-	-	-	-	1	6	7	11
Male	290	-	-	-	-	-	-	-	1	4	6	8
Female	255	-	-	-	-	-	-	-	-	2	1	3
White, both sexes	135	-	-	-	-	-	-	-	-	1	-	1
Male	82	-	-	-	-	-	-	-	-	1	-	1
Female	53	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	403	-	-	-	-	-	-	-	1	4	7	10
Male	205	-	-	-	-	-	-	-	1	2	6	7
Female	198	-	-	-	-	-	-	-	-	2	1	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Florida												
All races, both sexes	4,523	-	-	-	-	-	-	2	6	23	66	125
Male	2,487	-	-	-	-	-	-	2	5	21	53	99
Female	2,036	-	-	-	-	-	-	-	1	2	13	26
White, both sexes	4,119	-	-	-	-	-	-	2	5	16	51	110
Male	2,260	-	-	-	-	-	-	2	4	15	43	90
Female	1,859	-	-	-	-	-	-	-	1	1	8	20
Black, both sexes	370	-	-	-	-	-	-	-	-	7	12	15
Male	208	-	-	-	-	-	-	-	-	6	7	9
Female	162	-	-	-	-	-	-	-	-	1	5	6
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	29	-	-	-	-	-	-	-	1	-	3	-
Male	17	-	-	-	-	-	-	-	1	-	3	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	966	-	-	-	-	-	-	1	2	14	26	52
Male	577	-	-	-	-	-	-	-	1	11	22	42
Female	389	-	-	-	-	-	-	1	1	3	4	10
White, both sexes	697	-	-	-	-	-	-	1	1	7	15	41
Male	418	-	-	-	-	-	-	-	-	6	13	33
Female	279	-	-	-	-	-	-	1	1	1	2	8
Black, both sexes	260	-	-	-	-	-	-	-	1	6	11	10
Male	153	-	-	-	-	-	-	-	1	4	9	8
Female	107	-	-	-	-	-	-	-	-	2	2	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	1	-	1
Male	6	-	-	-	-	-	-	-	-	1	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Hawaii												
All races, both sexes	186	-	-	-	-	-	-	1	-	2	5	4
Male	120	-	-	-	-	-	-	1	-	2	3	3
Female	66	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	58	-	-	-	-	-	-	1	-	1	1	1
Male	46	-	-	-	-	-	-	1	-	1	1	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	125	-	-	-	-	-	-	-	-	1	4	3
Male	73	-	-	-	-	-	-	-	-	1	2	2
Female	52	-	-	-	-	-	-	-	-	-	2	1
Idaho												
All races, both sexes	68	-	-	-	-	-	-	-	-	-	3	-
Male	31	-	-	-	-	-	-	-	-	-	2	-
Female	37	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	67	-	-	-	-	-	-	-	-	-	3	-
Male	31	-	-	-	-	-	-	-	-	-	2	-
Female	36	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Illinois												
All races, both sexes	1,793	-	-	-	-	-	-	-	1	6	16	87
Male	1,092	-	-	-	-	-	-	-	1	5	14	60
Female	701	-	-	-	-	-	-	-	-	1	2	27
White, both sexes	1,309	-	-	-	-	-	-	-	1	5	9	48
Male	781	-	-	-	-	-	-	-	1	4	7	36
Female	528	-	-	-	-	-	-	-	-	1	2	12
Black, both sexes	458	-	-	-	-	-	-	-	-	1	6	38
Male	295	-	-	-	-	-	-	-	-	1	6	23
Female	163	-	-	-	-	-	-	-	-	-	-	15
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	24	-	-	-	-	-	-	-	-	-	1	1
Male	16	-	-	-	-	-	-	-	-	-	1	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	534	-	-	-	-	-	1	1	2	2	7	24
Male	279	-	-	-	-	-	1	1	2	2	4	19
Female	255	-	-	-	-	-	-	-	-	-	3	5
White, both sexes	475	-	-	-	-	-	1	1	1	2	5	24
Male	248	-	-	-	-	-	1	1	1	2	3	19
Female	227	-	-	-	-	-	-	-	-	-	2	5
Black, both sexes	57	-	-	-	-	-	-	-	1	-	2	-
Male	31	-	-	-	-	-	-	-	1	-	1	-
Female	26	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Iowa												
All races, both sexes	900	-	-	-	-	-	-	1	2	1	13	25
Male	443	-	-	-	-	-	-	1	1	1	11	15
Female	457	-	-	-	-	-	-	-	1	-	2	10
White, both sexes	878	-	-	-	-	-	-	1	2	1	13	22
Male	431	-	-	-	-	-	-	1	1	1	11	14
Female	447	-	-	-	-	-	-	-	1	-	2	8
Black, both sexes	15	-	-	-	-	-	-	-	-	-	-	3
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	281	-	-	-	-	-	1	-	1	3	6	5
Male	155	-	-	-	-	-	-	-	-	3	3	5
Female	126	-	-	-	-	-	1	-	1	-	3	-
White, both sexes	262	-	-	-	-	-	1	-	1	1	4	5
Male	142	-	-	-	-	-	-	-	-	1	2	5
Female	120	-	-	-	-	-	1	-	1	-	2	-
Black, both sexes	14	-	-	-	-	-	-	-	-	1	-	-
Male	10	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	2	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Kentucky												
All races, both sexes	505	-	-	-	-	-	-	3	5	4	19	27
Male	283	-	-	-	-	-	-	2	2	4	12	25
Female	222	-	-	-	-	-	-	1	3	-	7	2
White, both sexes	475	-	-	-	-	-	-	2	5	4	18	26
Male	265	-	-	-	-	-	-	1	2	4	12	24
Female	210	-	-	-	-	-	-	1	3	-	6	2
Black, both sexes	28	-	-	-	-	-	-	1	-	-	1	-
Male	17	-	-	-	-	-	-	1	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	998	-	-	-	-	-	2	3	2	5	21	39
Male	521	-	-	-	-	-	2	3	1	4	15	28
Female	477	-	-	-	-	-	-	-	1	1	6	11
White, both sexes	734	-	-	-	-	-	-	1	2	5	11	25
Male	375	-	-	-	-	-	-	1	1	4	7	18
Female	359	-	-	-	-	-	-	-	1	1	4	7
Black, both sexes	261	-	-	-	-	-	2	-	-	-	10	13
Male	143	-	-	-	-	-	2	-	-	-	8	9
Female	118	-	-	-	-	-	-	-	-	-	2	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	2	-	-	-	1
Male	3	-	-	-	-	-	-	2	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Maine												
All races, both sexes	166	-	-	-	-	-	-	1	1	1	8	4
Male	84	-	-	-	-	-	-	1	1	1	7	4
Female	82	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	165	-	-	-	-	-	-	1	1	1	8	4
Male	84	-	-	-	-	-	-	1	1	1	7	4
Female	81	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	2,383	-	-	-	-	-	2	4	5	12	34	59
Male	1,191	-	-	-	-	-	2	3	4	6	25	47
Female	1,192	-	-	-	-	-	-	1	1	6	9	12
White, both sexes	1,681	-	-	-	-	-	-	2	2	8	23	36
Male	822	-	-	-	-	-	-	2	1	3	17	32
Female	859	-	-	-	-	-	-	-	1	5	6	4
Black, both sexes	677	-	-	-	-	-	2	2	3	4	11	20
Male	354	-	-	-	-	-	2	1	3	3	8	13
Female	323	-	-	-	-	-	-	1	-	1	3	7
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	22	-	-	-	-	-	-	-	-	-	-	2
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Massachusetts												
All races, both sexes	1,307	-	-	-	-	-	-	2	1	9	25	52
Male	787	-	-	-	-	-	-	1	1	9	20	38
Female	520	-	-	-	-	-	-	1	-	-	5	14
White, both sexes	1,209	-	-	-	-	-	-	2	1	7	19	48
Male	726	-	-	-	-	-	-	1	1	7	17	34
Female	483	-	-	-	-	-	-	1	-	-	2	14
Black, both sexes	78	-	-	-	-	-	-	-	-	1	6	4
Male	48	-	-	-	-	-	-	-	-	1	3	4
Female	30	-	-	-	-	-	-	-	-	-	3	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	-	1	-	-
Male	10	-	-	-	-	-	-	-	-	1	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	3,812	-	-	-	-	-	1	5	7	33	57	157
Male	2,286	-	-	-	-	-	-	4	7	24	48	125
Female	1,526	-	-	-	-	-	1	1	-	9	9	32
White, both sexes	2,627	-	-	-	-	-	-	2	2	21	41	110
Male	1,590	-	-	-	-	-	-	1	2	16	35	88
Female	1,037	-	-	-	-	-	-	1	-	5	6	22
Black, both sexes	1,153	-	-	-	-	-	-	3	5	12	13	47
Male	679	-	-	-	-	-	-	3	5	8	12	37
Female	474	-	-	-	-	-	-	-	-	4	1	10
American Indian 1/, both sexes	16	-	-	-	-	-	-	-	-	-	2	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	2	-
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	1	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Minnesota												
All races, both sexes	333	-	-	-	-	-	-	-	1	-	1	9
Male	195	-	-	-	-	-	-	-	1	-	1	8
Female	138	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	322	-	-	-	-	-	-	-	1	-	1	8
Male	190	-	-	-	-	-	-	-	1	-	1	7
Female	132	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	555	-	-	-	-	-	-	-	-	-	5	3
Male	217	-	-	-	-	-	-	-	-	-	3	3
Female	338	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	433	-	-	-	-	-	-	-	-	-	3	2
Male	163	-	-	-	-	-	-	-	-	-	2	2
Female	270	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	121	-	-	-	-	-	-	-	-	-	2	1
Male	53	-	-	-	-	-	-	-	-	-	1	1
Female	68	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Missouri												
All races, both sexes	748	-	-	-	-	-	-	-	2	1	19	48
Male	399	-	-	-	-	-	-	-	2	1	13	36
Female	349	-	-	-	-	-	-	-	-	-	6	12
White, both sexes	663	-	-	-	-	-	-	-	2	1	16	38
Male	350	-	-	-	-	-	-	-	2	1	11	31
Female	313	-	-	-	-	-	-	-	-	-	5	7
Black, both sexes	80	-	-	-	-	-	-	-	-	-	2	10
Male	47	-	-	-	-	-	-	-	-	-	2	5
Female	33	-	-	-	-	-	-	-	-	-	-	5
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Montana												
All races, both sexes	222	-	-	-	-	-	-	-	-	-	-	8
Male	151	-	-	-	-	-	-	-	-	-	-	7
Female	71	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	216	-	-	-	-	-	-	-	-	-	-	6
Male	147	-	-	-	-	-	-	-	-	-	-	6
Female	69	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Nebraska												
All races, both sexes	117	-	-	-	-	-	-	-	-	-	2	1
Male	53	-	-	-	-	-	-	-	-	-	2	1
Female	64	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	116	-	-	-	-	-	-	-	-	-	2	1
Male	52	-	-	-	-	-	-	-	-	-	2	1
Female	64	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	978	-	-	-	-	-	-	-	1	2	12	23
Male	642	-	-	-	-	-	-	-	1	2	11	16
Female	336	-	-	-	-	-	-	-	-	-	1	7
White, both sexes	860	-	-	-	-	-	-	-	1	1	9	21
Male	559	-	-	-	-	-	-	-	1	1	8	16
Female	301	-	-	-	-	-	-	-	-	-	1	5
Black, both sexes	67	-	-	-	-	-	-	-	-	-	2	1
Male	47	-	-	-	-	-	-	-	-	-	2	-
Female	20	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	39	-	-	-	-	-	-	-	-	1	1	1
Male	26	-	-	-	-	-	-	-	-	1	1	-
Female	13	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
New Hampshire												
All races, both sexes	360	-	-	-	-	-	-	1	-	2	8	14
Male	209	-	-	-	-	-	-	1	-	2	8	9
Female	151	-	-	-	-	-	-	-	-	-	-	5
White, both sexes	359	-	-	-	-	-	-	1	-	2	8	14
Male	209	-	-	-	-	-	-	1	-	2	8	9
Female	150	-	-	-	-	-	-	-	-	-	-	5
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1,846	1	-	-	-	-	-	-	5	5	24	63
Male	1,024	1	-	-	-	-	-	-	5	4	13	52
Female	822	-	-	-	-	-	-	-	-	1	11	11
White, both sexes	1,552	1	-	-	-	-	-	-	3	3	18	49
Male	855	1	-	-	-	-	-	-	3	3	11	43
Female	697	-	-	-	-	-	-	-	-	-	7	6
Black, both sexes	263	-	-	-	-	-	-	-	2	1	5	13
Male	150	-	-	-	-	-	-	-	2	-	1	8
Female	113	-	-	-	-	-	-	-	-	1	4	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	31	-	-	-	-	-	-	-	-	1	1	1
Male	19	-	-	-	-	-	-	-	-	1	1	1
Female	12	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
New Mexico												
All races, both sexes	637	-	-	-	-	-	-	3	1	1	12	26
Male	396	-	-	-	-	-	-	3	1	1	11	22
Female	241	-	-	-	-	-	-	-	-	-	1	4
White, both sexes	582	-	-	-	-	-	-	3	-	1	10	25
Male	362	-	-	-	-	-	-	3	-	1	9	21
Female	220	-	-	-	-	-	-	-	-	-	1	4
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	42	-	-	-	-	-	-	-	1	-	2	1
Male	25	-	-	-	-	-	-	-	1	-	2	1
Female	17	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	3,851	-	-	-	-	-	-	3	11	30	82	128
Male	2,182	-	-	-	-	-	-	2	9	25	67	97
Female	1,669	-	-	-	-	-	-	1	2	5	15	31
White, both sexes	2,779	-	-	-	-	-	-	1	3	21	51	76
Male	1,579	-	-	-	-	-	-	1	2	19	48	62
Female	1,200	-	-	-	-	-	-	-	1	2	3	14
Black, both sexes	958	-	-	-	-	-	-	2	8	7	26	44
Male	516	-	-	-	-	-	-	1	7	4	14	27
Female	442	-	-	-	-	-	-	1	1	3	12	17
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	110	-	-	-	-	-	-	-	-	2	5	8
Male	84	-	-	-	-	-	-	-	-	2	5	8
Female	26	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
North Carolina												
All races, both sexes	1,170	-	-	-	-	-	-	3	5	19	35	57
Male	627	-	-	-	-	-	-	2	5	16	29	46
Female	543	-	-	-	-	-	-	1	-	3	6	11
White, both sexes	927	-	-	-	-	-	-	1	3	12	29	40
Male	497	-	-	-	-	-	-	-	3	11	24	32
Female	430	-	-	-	-	-	-	1	-	1	5	8
Black, both sexes	231	-	-	-	-	-	-	2	2	7	6	17
Male	121	-	-	-	-	-	-	2	2	5	5	14
Female	110	-	-	-	-	-	-	-	-	2	1	3
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	100	-	-	-	-	-	-	-	-	-	3	3
Male	54	-	-	-	-	-	-	-	-	-	3	2
Female	46	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	97	-	-	-	-	-	-	-	-	-	2	3
Male	53	-	-	-	-	-	-	-	-	-	2	2
Female	44	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Ohio												
All races, both sexes	2,122	-	-	-	-	-	-	3	8	22	45	113
Male	1,189	-	-	-	-	-	-	2	7	19	34	91
Female	933	-	-	-	-	-	-	1	1	3	11	22
White, both sexes	1,837	-	-	-	-	-	-	2	8	19	40	88
Male	1,019	-	-	-	-	-	-	1	7	16	31	72
Female	818	-	-	-	-	-	-	1	1	3	9	16
Black, both sexes	277	-	-	-	-	-	-	1	-	3	4	25
Male	168	-	-	-	-	-	-	1	-	3	3	19
Female	109	-	-	-	-	-	-	-	-	-	1	6
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
Oklahoma												
All races, both sexes	1,052	-	-	-	-	-	-	1	4	7	14	49
Male	585	-	-	-	-	-	-	1	3	5	10	36
Female	467	-	-	-	-	-	-	-	1	2	4	13
White, both sexes	916	-	-	-	-	-	-	1	3	6	11	35
Male	504	-	-	-	-	-	-	1	3	4	7	25
Female	412	-	-	-	-	-	-	-	-	2	4	10
Black, both sexes	65	-	-	-	-	-	-	-	-	-	-	5
Male	41	-	-	-	-	-	-	-	-	-	-	5
Female	24	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	65	-	-	-	-	-	-	-	1	1	3	9
Male	38	-	-	-	-	-	-	-	-	1	3	6
Female	27	-	-	-	-	-	-	-	1	-	-	3
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Oregon												
All races, both sexes	251	-	-	-	-	-	-	-	1	1	4	4
Male	131	-	-	-	-	-	-	-	1	1	2	2
Female	120	-	-	-	-	-	-	-	-	-	2	2
White, both sexes	242	-	-	-	-	-	-	-	1	1	4	4
Male	125	-	-	-	-	-	-	-	1	1	2	2
Female	117	-	-	-	-	-	-	-	-	-	2	2
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	3,263	-	-	-	-	-	2	4	5	13	35	103
Male	1,708	-	-	-	-	-	1	4	2	9	27	80
Female	1,555	-	-	-	-	-	1	-	3	4	8	23
White, both sexes	2,722	-	-	-	-	-	1	3	3	11	30	83
Male	1,389	-	-	-	-	-	1	3	1	9	23	66
Female	1,333	-	-	-	-	-	-	-	2	2	7	17
Black, both sexes	519	-	-	-	-	-	1	1	2	2	5	20
Male	307	-	-	-	-	-	-	1	1	-	4	14
Female	212	-	-	-	-	-	1	-	1	2	1	6
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Rhode Island												
All races, both sexes	520	-	-	-	-	-	-	-	1	7	9	11
Male	267	-	-	-	-	-	-	-	1	5	7	11
Female	253	-	-	-	-	-	-	-	-	2	2	-
White, both sexes	496	-	-	-	-	-	-	-	1	7	9	10
Male	252	-	-	-	-	-	-	-	1	5	7	10
Female	244	-	-	-	-	-	-	-	-	2	2	-
Black, both sexes	21	-	-	-	-	-	-	-	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	390	-	-	-	-	-	-	-	1	3	12	11
Male	228	-	-	-	-	-	-	-	1	3	11	10
Female	162	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	300	-	-	-	-	-	-	-	1	2	10	6
Male	174	-	-	-	-	-	-	-	1	2	9	6
Female	126	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	88	-	-	-	-	-	-	-	-	1	2	5
Male	52	-	-	-	-	-	-	-	-	1	2	4
Female	36	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
South Dakota												
All races, both sexes	84	-	-	-	-	-	-	-	-	-	1	-
Male	39	-	-	-	-	-	-	-	-	-	1	-
Female	45	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	82	-	-	-	-	-	-	-	-	-	1	-
Male	38	-	-	-	-	-	-	-	-	-	1	-
Female	44	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	1,929	-	-	-	-	-	-	2	4	25	34	101
Male	1,065	-	-	-	-	-	-	2	2	18	27	73
Female	864	-	-	-	-	-	-	-	2	7	7	28
White, both sexes	1,628	-	-	-	-	-	-	2	3	21	25	81
Male	874	-	-	-	-	-	-	2	1	15	18	57
Female	754	-	-	-	-	-	-	-	2	6	7	24
Black, both sexes	295	-	-	-	-	-	-	-	1	3	8	20
Male	186	-	-	-	-	-	-	-	1	2	8	16
Female	109	-	-	-	-	-	-	-	-	1	-	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	1	1	-
Male	5	-	-	-	-	-	-	-	-	1	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Texas												
All races, both sexes	3,817	-	-	-	-	2	-	9	13	42	98	225
Male	2,373	-	-	-	-	1	-	7	12	33	78	170
Female	1,444	-	-	-	-	1	-	2	1	9	20	55
White, both sexes	3,205	-	-	-	-	2	-	6	12	28	79	179
Male	1,983	-	-	-	-	1	-	4	12	21	61	138
Female	1,222	-	-	-	-	1	-	2	-	7	18	41
Black, both sexes	574	-	-	-	-	-	-	3	1	14	15	45
Male	363	-	-	-	-	-	-	3	-	12	13	31
Female	211	-	-	-	-	-	-	-	1	2	2	14
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	35	-	-	-	-	-	-	-	-	-	4	1
Male	24	-	-	-	-	-	-	-	-	-	4	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	83	-	-	-	-	-	-	-	-	-	-	1
Male	37	-	-	-	-	-	-	-	-	-	-	1
Female	46	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	79	-	-	-	-	-	-	-	-	-	-	1
Male	36	-	-	-	-	-	-	-	-	-	-	1
Female	43	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Vermont												
All races, both sexes	274	-	-	-	-	-	-	-	1	1	6	6
Male	153	-	-	-	-	-	-	-	1	-	5	4
Female	121	-	-	-	-	-	-	-	-	1	1	2
White, both sexes	273	-	-	-	-	-	-	-	1	1	6	6
Male	153	-	-	-	-	-	-	-	1	-	5	4
Female	120	-	-	-	-	-	-	-	-	1	1	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1,039	-	-	-	-	1	-	4	9	12	29	59
Male	567	-	-	-	-	1	-	4	7	8	24	52
Female	472	-	-	-	-	-	-	-	2	4	5	7
White, both sexes	798	-	-	-	-	-	-	2	6	6	23	39
Male	437	-	-	-	-	-	-	2	5	5	20	35
Female	361	-	-	-	-	-	-	-	1	1	3	4
Black, both sexes	219	-	-	-	-	1	-	2	3	6	3	18
Male	115	-	-	-	-	1	-	2	2	3	2	15
Female	104	-	-	-	-	-	-	-	1	3	1	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	21	-	-	-	-	-	-	-	-	-	3	2
Male	14	-	-	-	-	-	-	-	-	-	2	2
Female	7	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Washington												
All races, both sexes	1,423	-	-	-	-	-	1	1	3	12	21	61
Male	799	-	-	-	-	-	-	1	3	9	19	41
Female	624	-	-	-	-	-	1	-	-	3	2	20
White, both sexes	1,300	-	-	-	-	-	1	-	1	10	19	52
Male	726	-	-	-	-	-	-	-	1	8	17	35
Female	574	-	-	-	-	-	1	-	-	2	2	17
Black, both sexes	47	-	-	-	-	-	-	-	1	-	1	5
Male	34	-	-	-	-	-	-	-	1	-	1	4
Female	13	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	19	-	-	-	-	-	-	-	-	-	-	3
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	57	-	-	-	-	-	-	1	1	2	1	1
Male	28	-	-	-	-	-	-	1	1	1	1	1
Female	29	-	-	-	-	-	-	-	-	1	-	-
West Virginia												
All races, both sexes	426	-	-	-	-	-	1	-	1	3	7	10
Male	207	-	-	-	-	-	1	-	1	3	6	7
Female	219	-	-	-	-	-	-	-	-	-	1	3
White, both sexes	411	-	-	-	-	-	1	-	1	2	7	10
Male	198	-	-	-	-	-	1	-	1	2	6	7
Female	213	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	14	-	-	-	-	-	-	-	-	1	-	-
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Wisconsin												
All races, both sexes	936	-	-	-	-	-	-	-	3	8	12	33
Male	552	-	-	-	-	-	-	-	3	7	11	27
Female	384	-	-	-	-	-	-	-	-	1	1	6
White, both sexes	833	-	-	-	-	-	-	-	1	8	9	26
Male	489	-	-	-	-	-	-	-	1	7	9	21
Female	344	-	-	-	-	-	-	-	-	1	-	5
Black, both sexes	87	-	-	-	-	-	-	-	1	-	3	6
Male	55	-	-	-	-	-	-	-	1	-	2	5
Female	32	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	1	-	-	1
Male	4	-	-	-	-	-	-	-	1	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	36	-	-	-	-	-	1	-	1	-	-	2
Male	24	-	-	-	-	-	1	-	1	-	-	2
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	34	-	-	-	-	-	1	-	-	-	-	2
Male	23	-	-	-	-	-	1	-	-	-	-	2
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
United States												
All races, both sexes	3,719	4,721	5,334	4,782	5,102	6,298	7,923	8,007	6,021	2,690	629	10
Male	2,905	3,657	3,926	3,305	3,244	3,589	3,963	3,298	1,806	595	95	7
Female	814	1,064	1,408	1,477	1,858	2,709	3,960	4,709	4,215	2,095	534	3
White, both sexes	2,815	3,590	4,180	3,700	4,037	5,186	6,811	7,103	5,411	2,394	543	9
Male	2,250	2,835	3,122	2,618	2,602	2,988	3,430	2,954	1,634	534	76	6
Female	565	755	1,058	1,082	1,435	2,198	3,381	4,149	3,777	1,860	467	3
Black, both sexes	799	995	1,017	947	930	933	920	742	512	255	76	1
Male	562	712	701	601	565	505	447	278	137	50	16	1
Female	237	283	316	346	365	428	473	464	375	205	60	-
American Indian 1/, both sexes	31	29	32	38	38	45	36	23	8	8	-	-
Male	26	22	22	27	24	23	18	8	2	1	-	-
Female	5	7	10	11	14	22	18	15	6	7	-	-
Asian or Pacific Islander 2/, both sexes	74	107	105	97	97	134	156	139	90	33	10	-
Male	67	88	81	59	53	73	68	58	33	10	3	-
Female	7	19	24	38	44	61	88	81	57	23	7	-
Alabama												
All races, both sexes	22	63	33	42	55	56	93	88	53	30	5	-
Male	18	48	26	28	32	24	34	27	17	9	-	-
Female	4	15	7	14	23	32	59	61	36	21	5	-
White, both sexes	16	50	26	31	42	41	75	79	47	25	4	-
Male	12	40	20	21	27	17	31	26	15	8	-	-
Female	4	10	6	10	15	24	44	53	32	17	4	-
Black, both sexes	6	13	7	10	13	15	18	9	6	5	1	-
Male	6	8	6	6	5	7	3	1	2	1	-	-
Female	-	5	1	4	8	8	15	8	4	4	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Alaska												
All races, both sexes	12	18	17	12	9	13	5	5	5	1	-	-
Male	11	14	17	7	9	8	2	1	3	-	-	-
Female	1	4	-	5	-	5	3	4	2	1	-	-
White, both sexes	10	17	17	8	6	8	5	4	5	1	-	-
Male	9	13	17	5	6	6	2	1	3	-	-	-
Female	1	4	-	3	-	2	3	3	2	1	-	-
Black, both sexes	2	-	-	-	-	1	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	4	3	4	-	-	-	-	-	-
Male	-	1	-	2	3	2	-	-	-	-	-	-
Female	-	-	-	2	-	2	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Arizona												
All races, both sexes	100	134	207	191	172	220	243	216	129	68	5	1
Male	78	114	157	133	112	132	133	89	42	20	-	1
Female	22	20	50	58	60	88	110	127	87	48	5	-
White, both sexes	91	122	196	179	161	207	235	203	122	65	5	1
Male	71	106	149	127	106	123	126	85	41	20	-	1
Female	20	16	47	52	55	84	109	118	81	45	5	-
Black, both sexes	4	7	7	6	7	7	4	8	5	2	-	-
Male	2	4	5	2	4	3	4	3	1	-	-	-
Female	2	3	2	4	3	4	-	5	4	2	-	-
American Indian 1/, both sexes	4	5	3	4	3	4	4	4	1	1	-	-
Male	4	4	2	2	2	4	3	1	-	-	-	-
Female	-	1	1	2	1	-	1	3	1	1	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	2	1	2	-	1	1	-	-	-
Male	1	-	1	2	-	2	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Arkansas												
All races, both sexes	15	20	21	18	19	24	28	28	25	15	1	-
Male	11	18	17	10	10	13	14	10	9	2	-	-
Female	4	2	4	8	9	11	14	18	16	13	1	-
White, both sexes	9	16	17	13	17	20	25	25	21	14	1	-
Male	6	14	13	8	9	10	13	7	6	2	-	-
Female	3	2	4	5	8	10	12	18	15	12	1	-
Black, both sexes	5	4	4	5	1	4	3	3	3	-	-	-
Male	4	4	4	2	1	3	1	3	3	-	-	-
Female	1	-	-	3	-	1	2	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	1	-	-
California												
All races, both sexes	503	621	752	634	643	765	946	939	727	325	90	-
Male	428	508	565	442	406	435	428	382	231	79	19	-
Female	75	113	187	192	237	330	518	557	496	246	71	-
White, both sexes	378	481	600	503	514	617	802	790	618	278	74	-
Male	324	395	451	359	328	350	360	337	200	67	16	-
Female	54	86	149	144	186	267	442	453	418	211	58	-
Black, both sexes	89	98	98	93	85	82	73	75	55	26	8	-
Male	70	75	73	61	54	50	37	21	11	5	-	-
Female	19	23	25	32	31	32	36	54	44	21	8	-
American Indian 1/, both sexes	4	1	6	5	4	3	3	3	1	2	-	-
Male	4	1	5	4	1	2	1	1	1	-	-	-
Female	-	-	1	1	3	1	2	2	-	2	-	-
Asian or Pacific Islander 2/, both sexes	32	41	48	33	40	63	68	71	53	19	8	-
Male	30	37	36	18	23	33	30	23	19	7	3	-
Female	2	4	12	15	17	30	38	48	34	12	5	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Colorado												
All races, both sexes	42	59	61	60	44	63	81	81	44	27	7	-
Male	37	45	43	43	28	39	38	33	14	7	-	-
Female	5	14	18	17	16	24	43	48	30	20	7	-
White, both sexes	40	54	58	56	38	60	79	79	43	25	7	-
Male	35	41	40	40	25	38	38	32	14	6	-	-
Female	5	13	18	16	13	22	41	47	29	19	7	-
Black, both sexes	1	5	3	4	6	3	1	2	1	2	-	-
Male	1	4	3	3	3	1	-	1	-	1	-	-
Female	-	1	-	1	3	2	1	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Connecticut												
All races, both sexes	37	47	60	40	43	59	79	95	71	45	14	-
Male	31	40	43	26	27	36	47	45	25	13	2	-
Female	6	7	17	14	16	23	32	50	46	32	12	-
White, both sexes	29	40	51	31	40	56	72	94	69	45	13	-
Male	26	33	36	19	24	33	43	44	25	13	1	-
Female	3	7	15	12	16	23	29	50	44	32	12	-
Black, both sexes	6	6	8	9	3	3	7	1	2	-	1	-
Male	4	6	6	7	3	3	4	1	-	-	1	-
Female	2	-	2	2	-	-	3	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	1	-	-	-	-	-	-	-	-	-
Male	1	1	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Delaware												
All races, both sexes	20	32	36	19	31	30	31	30	28	18	4	1
Male	16	25	26	13	23	20	13	14	7	3	1	-
Female	4	7	10	6	8	10	18	16	21	15	3	1
White, both sexes	12	24	30	13	26	25	28	25	26	14	4	1
Male	9	19	21	8	21	15	12	11	7	2	1	-
Female	3	5	9	5	5	10	16	14	19	12	3	1
Black, both sexes	8	8	6	5	5	5	3	4	2	3	-	-
Male	7	6	5	4	2	5	1	2	-	-	-	-
Female	1	2	1	1	3	-	2	2	2	3	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	1	-	-
Male	-	-	-	1	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	11	32	64	69	57	66	78	59	50	25	9	-
Male	7	22	45	46	33	36	35	22	18	4	3	-
Female	4	10	19	23	24	30	43	37	32	21	6	-
White, both sexes	1	8	11	11	11	18	20	18	19	11	5	-
Male	1	5	10	8	8	12	14	10	9	3	-	-
Female	-	3	1	3	3	6	6	8	10	8	5	-
Black, both sexes	10	24	53	57	45	46	57	41	31	13	4	-
Male	6	17	35	37	24	24	21	12	9	1	3	-
Female	4	7	18	20	21	22	36	29	22	12	1	-
American Indian 1/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	2	-	-	-	1	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Florida												
All races, both sexes	192	259	304	320	334	470	683	787	589	300	61	2
Male	153	198	227	219	211	283	362	350	215	74	14	1
Female	39	61	77	101	123	187	321	437	374	226	47	1
White, both sexes	153	234	264	277	281	423	634	761	565	282	59	2
Male	127	178	197	189	182	259	338	343	207	71	14	1
Female	26	56	67	88	99	164	296	418	358	211	45	1
Black, both sexes	36	24	37	39	47	41	45	26	23	16	2	-
Male	23	19	28	28	26	22	23	7	8	2	-	-
Female	13	5	9	11	21	19	22	19	15	14	2	-
American Indian 1/, both sexes	-	-	1	1	1	1	-	-	1	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	3	1	2	3	5	5	4	-	-	2	-	-
Male	3	1	2	1	2	2	1	-	-	1	-	-
Female	-	-	-	2	3	3	3	-	-	1	-	-
Georgia												
All races, both sexes	72	74	98	87	91	83	107	125	88	38	8	-
Male	57	57	77	64	67	43	53	48	30	4	1	-
Female	15	17	21	23	24	40	54	77	58	34	7	-
White, both sexes	49	46	67	59	72	58	84	95	72	24	6	-
Male	41	35	52	44	56	33	40	36	25	3	1	-
Female	8	11	15	15	16	25	44	59	47	21	5	-
Black, both sexes	23	26	30	27	19	24	21	30	16	14	2	-
Male	16	20	24	20	11	10	12	12	5	1	-	-
Female	7	6	6	7	8	14	9	18	11	13	2	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	1	1	-	-	2	-	-	-	-	-
Male	-	2	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Hawaii												
All races, both sexes	8	21	21	30	17	21	20	15	16	5	-	-
Male	6	19	13	21	12	12	15	7	4	2	-	-
Female	2	2	8	9	5	9	5	8	12	3	-	-
White, both sexes	3	8	8	11	7	9	1	-	5	2	-	-
Male	3	8	7	10	5	6	1	-	1	1	-	-
Female	-	-	1	1	2	3	-	-	4	1	-	-
Black, both sexes	-	1	1	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	12	12	19	10	11	19	15	11	3	-	-
Male	3	10	6	11	7	6	14	7	3	1	-	-
Female	2	2	6	8	3	5	5	8	8	2	-	-
Idaho												
All races, both sexes	5	2	4	5	1	9	9	13	10	6	1	-
Male	2	1	3	5	-	4	5	3	4	2	-	-
Female	3	1	1	-	1	5	4	10	6	4	1	-
White, both sexes	5	2	4	5	1	9	9	12	10	6	1	-
Male	2	1	3	5	-	4	5	3	4	2	-	-
Female	3	1	1	-	1	5	4	9	6	4	1	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Illinois												
All races, both sexes	180	221	187	157	178	154	230	179	118	62	16	1
Male	139	158	134	124	116	85	124	81	36	12	3	-
Female	41	63	53	33	62	69	106	98	82	50	13	1
White, both sexes	119	138	122	109	119	120	180	159	108	58	13	1
Male	99	101	94	86	76	66	99	68	32	11	1	-
Female	20	37	28	23	43	54	81	91	76	47	12	1
Black, both sexes	57	79	63	48	58	30	45	17	9	4	3	-
Male	36	55	38	38	40	17	24	10	4	1	2	-
Female	21	24	25	10	18	13	21	7	5	3	1	-
American Indian 1/, both sexes	-	2	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	2	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	2	2	-	1	4	5	3	1	-	-	-
Male	4	2	2	-	-	2	1	3	-	-	-	-
Female	-	-	-	-	1	2	4	-	1	-	-	-
Indiana												
All races, both sexes	31	30	40	42	34	69	75	64	71	31	10	-
Male	20	23	31	25	23	40	38	31	16	3	-	-
Female	11	7	9	17	11	29	37	33	55	28	10	-
White, both sexes	25	24	32	36	30	62	70	58	65	30	9	-
Male	17	18	26	23	20	35	35	29	15	3	-	-
Female	8	6	6	13	10	27	35	29	50	27	9	-
Black, both sexes	6	6	8	5	3	7	5	6	6	1	1	-
Male	3	5	5	2	3	5	3	2	1	-	-	-
Female	3	1	3	3	-	2	2	4	5	1	1	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Iowa												
All races, both sexes	34	53	53	61	64	101	129	147	139	65	12	-
Male	28	38	36	38	41	65	63	56	34	13	2	-
Female	6	15	17	23	23	36	66	91	105	52	10	-
White, both sexes	34	52	51	57	58	98	128	147	137	65	12	-
Male	28	38	35	36	37	62	63	56	33	13	2	-
Female	6	14	16	21	21	36	65	91	104	52	10	-
Black, both sexes	-	1	2	2	4	2	-	-	1	-	-	-
Male	-	-	1	-	3	2	-	-	1	-	-	-
Female	-	1	1	2	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	1	1	1	-	1	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	18	14	19	23	26	28	46	42	28	14	7	-
Male	16	8	14	13	14	19	26	24	6	2	2	-
Female	2	6	5	10	12	9	20	18	22	12	5	-
White, both sexes	17	13	17	20	24	27	44	42	26	13	7	-
Male	15	7	12	11	14	18	24	24	5	2	2	-
Female	2	6	5	9	10	9	20	18	21	11	5	-
Black, both sexes	1	1	1	2	2	1	2	-	2	1	-	-
Male	1	1	1	2	-	1	2	-	1	-	-	-
Female	-	-	-	-	2	-	-	-	1	1	-	-
American Indian 1/, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Kentucky												
All races, both sexes	26	32	33	45	44	49	64	84	47	19	4	-
Male	18	27	23	33	30	30	33	32	10	2	-	-
Female	8	5	10	12	14	19	31	52	37	17	4	-
White, both sexes	24	28	30	43	40	46	61	82	43	19	4	-
Male	17	24	20	31	28	29	32	31	8	2	-	-
Female	7	4	10	12	12	17	29	51	35	17	4	-
Black, both sexes	2	4	3	2	3	3	3	2	4	-	-	-
Male	1	3	3	2	2	1	1	1	2	-	-	-
Female	1	1	-	-	1	2	2	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	59	71	66	81	100	94	145	153	97	47	13	-
Male	39	53	54	52	57	51	70	58	23	9	2	-
Female	20	18	12	29	43	43	75	95	74	38	11	-
White, both sexes	38	44	48	61	74	67	109	119	84	40	6	-
Male	27	35	41	37	42	37	54	42	21	8	-	-
Female	11	9	7	24	32	30	55	77	63	32	6	-
Black, both sexes	21	27	18	20	26	27	36	34	13	7	7	-
Male	12	18	13	15	15	14	16	16	2	1	2	-
Female	9	9	5	5	11	13	20	18	11	6	5	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Maine												
All races, both sexes	10	7	9	7	15	22	20	23	23	12	3	-
Male	8	5	6	2	11	12	9	11	3	3	-	-
Female	2	2	3	5	4	10	11	12	20	9	3	-
White, both sexes	10	7	9	6	15	22	20	23	23	12	3	-
Male	8	5	6	2	11	12	9	11	3	3	-	-
Female	2	2	3	4	4	10	11	12	20	9	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	147	152	176	151	197	231	357	381	297	142	36	-
Male	106	111	124	95	115	131	174	154	63	28	3	-
Female	41	41	52	56	82	100	183	227	234	114	33	-
White, both sexes	94	86	110	89	122	166	266	300	235	114	28	-
Male	67	64	79	60	72	97	129	122	52	24	1	-
Female	27	22	31	29	50	69	137	178	183	90	27	-
Black, both sexes	53	61	66	61	74	63	86	75	61	27	8	-
Male	39	43	45	34	43	33	43	28	10	4	2	-
Female	14	18	21	27	31	30	43	47	51	23	6	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	5	-	1	1	2	4	5	1	1	-	-
Male	-	4	-	1	-	1	2	3	1	-	-	-
Female	-	1	-	-	1	1	2	2	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Massachusetts												
All races, both sexes	92	104	116	110	139	148	172	165	122	41	8	1
Male	77	85	89	80	92	85	93	67	41	8	-	1
Female	15	19	27	30	47	63	79	98	81	33	8	-
White, both sexes	82	96	104	101	123	137	163	160	117	40	8	1
Male	70	82	80	74	79	80	89	64	39	8	-	1
Female	12	14	24	27	44	57	74	96	78	32	8	-
Black, both sexes	7	5	11	7	12	8	8	4	4	1	-	-
Male	5	2	8	5	9	4	3	3	1	-	-	-
Female	2	3	3	2	3	4	5	1	3	1	-	-
American Indian 1/, both sexes	2	1	1	-	-	1	-	-	-	-	-	-
Male	1	1	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	-	2	4	2	1	1	1	-	-	-
Male	1	-	-	1	4	1	1	-	1	-	-	-
Female	-	2	-	1	-	1	-	1	-	-	-	-
Michigan												
All races, both sexes	307	393	438	358	337	440	479	414	269	93	24	-
Male	229	287	316	247	212	247	242	184	88	21	5	-
Female	78	106	122	111	125	193	237	230	181	72	19	-
White, both sexes	202	239	290	230	224	290	344	333	208	74	17	-
Male	161	184	214	171	149	167	176	144	63	16	3	-
Female	41	55	76	59	75	123	168	189	145	58	14	-
Black, both sexes	102	154	143	126	112	147	127	79	59	18	6	-
Male	66	103	98	74	62	78	61	40	25	5	2	-
Female	36	51	45	52	50	69	66	39	34	13	4	-
American Indian 1/, both sexes	2	-	3	2	1	3	3	-	-	-	-	-
Male	2	-	3	2	1	2	2	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	2	-	-	-	5	2	2	1	1	-
Male	-	-	1	-	-	-	3	-	-	-	-	-
Female	1	-	1	-	-	-	2	2	2	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Minnesota												
All races, both sexes	13	15	27	15	12	43	63	57	54	20	3	-
Male	12	13	19	12	9	30	36	24	23	6	1	-
Female	1	2	8	3	3	13	27	33	31	14	2	-
White, both sexes	11	15	25	15	12	40	62	57	54	18	3	-
Male	10	13	18	12	9	29	36	24	23	6	1	-
Female	1	2	7	3	3	11	26	33	31	12	2	-
Black, both sexes	1	-	1	-	-	-	1	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	3	-	-	-	1	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	16	23	32	26	42	62	82	99	104	46	15	-
Male	10	17	19	18	23	29	33	32	18	10	2	-
Female	6	6	13	8	19	33	49	67	86	36	13	-
White, both sexes	12	14	22	19	33	47	65	82	87	39	8	-
Male	8	10	12	14	19	21	25	27	14	8	1	-
Female	4	4	10	5	14	26	40	55	73	31	7	-
Black, both sexes	4	9	10	7	9	14	17	17	17	7	7	-
Male	2	7	7	4	4	7	8	5	4	2	1	-
Female	2	2	3	3	5	7	9	12	13	5	6	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Missouri												
All races, both sexes	44	52	58	56	61	79	102	105	77	33	10	1
Male	28	44	44	36	30	47	53	39	19	3	3	1
Female	16	8	14	20	31	32	49	66	58	30	7	-
White, both sexes	38	41	54	49	51	70	92	95	73	33	9	1
Male	24	33	42	33	21	42	48	38	18	3	2	1
Female	14	8	12	16	30	28	44	57	55	30	7	-
Black, both sexes	5	11	4	7	9	8	10	9	4	-	1	-
Male	4	11	2	3	8	4	5	1	1	-	1	-
Female	1	-	2	4	1	4	5	8	3	-	-	-
American Indian 1/, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Montana												
All races, both sexes	15	18	29	25	27	31	34	22	9	4	-	-
Male	11	15	21	16	17	25	17	15	4	3	-	-
Female	4	3	8	9	10	6	17	7	5	1	-	-
White, both sexes	15	17	28	25	27	31	32	22	9	4	-	-
Male	11	14	20	16	17	25	16	15	4	3	-	-
Female	4	3	8	9	10	6	16	7	5	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	1	-	-	-	2	-	-	-	-	-
Male	-	1	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Nebraska												
All races, both sexes	2	6	7	1	5	8	21	31	17	14	2	-
Male	1	5	6	1	3	6	11	8	9	-	-	-
Female	1	1	1	-	2	2	10	23	8	14	2	-
White, both sexes	2	6	7	1	5	8	21	30	17	14	2	-
Male	1	5	6	1	3	6	11	7	9	-	-	-
Female	1	1	1	-	2	2	10	23	8	14	2	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	73	117	127	121	123	118	117	83	45	15	1	-
Male	56	92	100	88	76	71	67	40	19	3	-	-
Female	17	25	27	33	47	47	50	43	26	12	1	-
White, both sexes	64	97	111	107	104	107	108	74	42	13	1	-
Male	50	74	89	79	62	63	61	35	17	3	-	-
Female	14	23	22	28	42	44	47	39	25	10	1	-
Black, both sexes	3	12	12	9	11	6	4	6	-	1	-	-
Male	1	11	7	6	9	5	3	3	-	-	-	-
Female	2	1	5	3	2	1	1	3	-	1	-	-
American Indian 1/, both sexes	2	1	1	1	2	3	2	-	-	-	-	-
Male	2	1	1	-	2	2	2	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	7	3	4	6	2	3	3	3	1	-	-
Male	3	6	3	3	3	1	1	2	2	-	-	-
Female	1	1	-	1	3	1	2	1	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
New Hampshire												
All races, both sexes	22	25	37	31	30	36	44	49	43	14	4	-
Male	20	17	30	21	21	21	19	25	10	3	2	-
Female	2	8	7	10	9	15	25	24	33	11	2	-
White, both sexes	22	25	36	31	30	36	44	49	43	14	4	-
Male	20	17	30	21	21	21	19	25	10	3	2	-
Female	2	8	6	10	9	15	25	24	33	11	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	88	123	143	139	168	217	260	272	234	84	20	-
Male	76	103	114	97	111	114	134	112	65	21	2	-
Female	12	20	29	42	57	103	126	160	169	63	18	-
White, both sexes	68	96	118	103	138	183	221	242	214	76	19	-
Male	61	84	95	74	90	94	114	102	59	19	2	-
Female	7	12	23	29	48	89	107	140	155	57	17	-
Black, both sexes	18	24	22	34	26	30	36	27	17	7	1	-
Male	13	17	16	21	18	18	19	10	5	2	-	-
Female	5	7	6	13	8	12	17	17	12	5	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	3	3	2	4	4	3	3	3	1	-	-
Male	2	2	3	2	3	2	1	-	1	-	-	-
Female	-	1	-	-	1	2	2	3	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
New Mexico												
All races, both sexes	35	59	72	54	61	82	70	74	51	28	8	-
Male	27	49	51	41	37	45	42	36	22	5	3	-
Female	8	10	21	13	24	37	28	38	29	23	5	-
White, both sexes	31	54	67	51	52	74	65	67	47	27	8	-
Male	24	46	47	38	33	42	38	32	20	5	3	-
Female	7	8	20	13	19	32	27	35	27	22	5	-
Black, both sexes	2	1	1	-	1	1	1	1	1	-	-	-
Male	2	-	1	-	-	1	1	1	-	-	-	-
Female	-	1	-	-	1	-	-	-	1	-	-	-
American Indian 1/, both sexes	2	3	4	3	8	6	4	5	2	1	-	-
Male	1	2	3	3	4	2	3	2	1	-	-	-
Female	1	1	1	-	4	4	1	3	1	1	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	-	1	1	-	-	-
Male	-	1	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
New York												
All races, both sexes	228	293	362	362	366	495	518	435	354	156	28	-
Male	166	225	256	239	242	274	249	190	106	33	2	-
Female	62	68	106	123	124	221	269	245	248	123	26	-
White, both sexes	142	198	223	219	253	358	416	354	308	135	21	-
Male	106	163	163	153	174	198	200	162	96	31	1	-
Female	36	35	60	66	79	160	216	192	212	104	20	-
Black, both sexes	75	79	126	130	109	121	90	75	42	18	6	-
Male	49	49	80	75	67	66	43	24	8	1	1	-
Female	26	30	46	55	42	55	47	51	34	17	5	-
American Indian 1/, both sexes	1	1	-	1	-	1	-	-	-	-	-	-
Male	1	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	15	13	12	4	15	12	6	4	3	1	-
Male	10	12	13	10	1	10	6	4	2	1	-	-
Female	-	3	-	2	3	5	6	2	2	2	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
North Carolina												
All races, both sexes	70	77	82	64	81	114	157	180	147	60	18	1
Male	51	61	62	43	52	61	83	66	35	13	1	1
Female	19	16	20	21	29	53	74	114	112	47	17	-
White, both sexes	44	54	61	49	62	94	131	160	128	47	12	-
Male	32	42	51	33	42	54	73	60	31	9	-	-
Female	12	12	10	16	20	40	58	100	97	38	12	-
Black, both sexes	24	21	21	14	18	19	24	18	18	13	6	1
Male	17	17	11	9	9	7	8	5	4	4	1	1
Female	7	4	10	5	9	12	16	13	14	9	5	-
American Indian 1/, both sexes	1	1	-	-	1	1	1	2	1	-	-	-
Male	1	1	-	-	1	-	1	1	-	-	-	-
Female	-	-	-	-	-	1	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	1	-	-	1	-	-	-	-	-
Male	1	1	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	3	6	6	10	10	8	11	15	16	8	1	-
Male	3	5	3	5	6	8	4	8	6	1	-	-
Female	-	1	3	5	4	-	7	7	10	7	1	-
White, both sexes	3	6	5	10	10	8	10	15	16	8	1	-
Male	3	5	3	5	6	8	4	8	6	1	-	-
Female	-	1	2	5	4	-	6	7	10	7	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Ohio												
All races, both sexes	143	192	175	143	188	212	276	288	223	70	21	-
Male	107	140	127	100	116	112	131	122	61	17	3	-
Female	36	52	48	43	72	100	145	166	162	53	18	-
White, both sexes	115	142	148	122	154	182	255	271	209	63	19	-
Male	90	106	109	88	95	100	120	111	56	14	3	-
Female	25	36	39	34	59	82	135	160	153	49	16	-
Black, both sexes	28	46	26	21	33	30	20	17	14	7	2	-
Male	17	33	18	12	20	12	11	11	5	3	-	-
Female	11	13	8	9	13	18	9	6	9	4	2	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	3	1	-	1	-	1	-	-	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	2	1	-	-	-	1	-	-	-	-	-
Oklahoma												
All races, both sexes	77	82	85	86	93	114	150	143	85	46	16	-
Male	56	60	54	60	56	64	81	59	26	10	4	-
Female	21	22	31	26	37	50	69	84	59	36	12	-
White, both sexes	63	69	75	72	80	93	136	132	83	42	15	-
Male	46	50	48	49	49	52	76	55	26	9	4	-
Female	17	19	27	23	31	41	60	77	57	33	11	-
Black, both sexes	8	8	6	9	5	9	6	4	1	3	1	-
Male	6	6	5	7	3	4	3	1	-	1	-	-
Female	2	2	1	2	2	5	3	3	1	2	1	-
American Indian 1/, both sexes	6	5	4	4	7	11	6	6	1	1	-	-
Male	4	4	1	4	4	7	2	2	-	-	-	-
Female	2	1	3	-	3	4	4	4	1	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	1	2	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	1	1	-	2	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Oregon												
All races, both sexes	14	11	8	15	27	29	38	46	37	15	1	-
Male	12	10	5	11	17	18	28	13	7	4	-	-
Female	2	1	3	4	10	11	10	33	30	11	1	-
White, both sexes	12	10	8	15	26	27	36	45	37	15	1	-
Male	10	9	5	11	16	17	27	13	7	4	-	-
Female	2	1	3	4	10	10	9	32	30	11	1	-
Black, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	2	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	1	1	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Pennsylvania												
All races, both sexes	156	229	264	191	258	387	479	511	389	193	43	1
Male	127	170	189	137	170	198	241	219	90	37	6	1
Female	29	59	75	54	88	189	238	292	299	156	37	-
White, both sexes	109	165	190	137	196	325	425	457	367	178	41	1
Male	92	126	139	100	127	166	213	195	86	35	6	1
Female	17	39	51	37	69	159	212	262	281	143	35	-
Black, both sexes	44	63	73	51	61	59	49	50	21	15	2	-
Male	32	43	50	36	42	30	26	22	4	2	-	-
Female	12	20	23	15	19	29	23	28	17	13	2	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	1	1	3	-	3	4	4	1	-	-	-
Male	3	1	-	1	-	2	1	2	-	-	-	-
Female	-	-	1	2	-	1	3	2	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Rhode Island												
All races, both sexes	25	36	25	26	23	49	80	92	80	52	4	-
Male	22	31	17	20	12	32	43	34	18	13	1	-
Female	3	5	8	6	11	17	37	58	62	39	3	-
White, both sexes	22	34	21	25	21	46	77	88	80	51	4	-
Male	19	29	15	19	11	30	42	33	18	12	1	-
Female	3	5	6	6	10	16	35	55	62	39	3	-
Black, both sexes	2	2	3	1	2	2	3	4	-	1	-	-
Male	2	2	2	1	1	1	1	1	-	1	-	-
Female	-	-	1	-	1	1	2	3	-	-	-	-
American Indian 1/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	23	33	37	32	35	35	64	55	27	18	4	-
Male	21	23	25	29	19	16	33	25	8	4	-	-
Female	2	10	12	3	16	19	31	30	19	14	4	-
White, both sexes	19	21	26	24	27	26	50	45	26	15	2	-
Male	17	15	17	21	14	13	26	23	7	3	-	-
Female	2	6	9	3	13	13	24	22	19	12	2	-
Black, both sexes	3	12	11	7	8	9	14	10	1	3	2	-
Male	3	8	8	7	5	3	7	2	1	1	-	-
Female	-	4	3	-	3	6	7	8	-	2	2	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
South Dakota												
All races, both sexes	1	3	1	4	4	6	16	24	11	7	6	-
Male	1	3	1	2	3	4	7	13	2	2	-	-
Female	-	-	-	2	1	2	9	11	9	5	6	-
White, both sexes	1	3	1	3	4	6	16	23	11	7	6	-
Male	1	3	1	1	3	4	7	13	2	2	-	-
Female	-	-	-	2	1	2	9	10	9	5	6	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	143	170	154	154	166	202	241	257	183	77	16	-
Male	112	131	106	96	99	115	110	92	61	19	2	-
Female	31	39	48	58	67	87	131	165	122	58	14	-
White, both sexes	109	134	126	132	132	173	212	231	164	67	16	-
Male	83	100	87	83	79	96	95	81	56	19	2	-
Female	26	34	39	49	53	77	117	150	108	48	14	-
Black, both sexes	34	36	28	22	32	29	29	25	18	10	-	-
Male	29	31	19	13	18	19	15	10	5	-	-	-
Female	5	5	9	9	14	10	14	15	13	10	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	-	-	1	1	-	-	-
Male	-	-	-	-	2	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Texas												
All races, both sexes	320	388	449	369	325	338	408	390	292	120	29	-
Male	242	305	345	269	219	200	219	151	93	29	-	-
Female	78	83	104	100	106	138	189	239	199	91	29	-
White, both sexes	250	315	368	299	270	290	363	357	259	102	26	-
Male	189	252	288	223	183	171	191	141	86	22	-	-
Female	61	63	80	76	87	119	172	216	173	80	26	-
Black, both sexes	68	71	74	64	51	43	41	30	33	18	3	-
Male	51	52	52	41	34	26	27	7	7	7	-	-
Female	17	19	22	23	17	17	14	23	26	11	3	-
American Indian 1/, both sexes	-	-	-	3	-	-	-	-	-	-	-	-
Male	-	-	-	3	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	7	3	4	5	4	3	-	-	-	-
Male	2	1	5	2	2	3	1	3	-	-	-	-
Female	-	1	2	1	2	2	3	-	-	-	-	-
Utah												
All races, both sexes	3	6	4	8	6	8	13	20	10	4	-	-
Male	3	3	3	7	4	4	4	5	3	-	-	-
Female	-	3	1	1	2	4	9	15	7	4	-	-
White, both sexes	3	5	3	8	5	8	12	20	10	4	-	-
Male	3	3	2	7	4	4	4	5	3	-	-	-
Female	-	2	1	1	1	4	8	15	7	4	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Vermont												
All races, both sexes	19	19	29	18	24	23	44	37	35	10	2	-
Male	17	14	21	9	13	14	25	12	14	3	1	-
Female	2	5	8	9	11	9	19	25	21	7	1	-
White, both sexes	19	19	29	17	24	23	44	37	35	10	2	-
Male	17	14	21	9	13	14	25	12	14	3	1	-
Female	2	5	8	8	11	9	19	25	21	7	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	62	77	61	77	84	109	135	156	113	42	9	-
Male	47	59	44	43	54	65	61	54	35	8	1	-
Female	15	18	17	34	30	44	74	102	78	34	8	-
White, both sexes	44	54	46	47	68	84	113	129	94	35	8	-
Male	38	43	33	28	46	50	52	45	29	5	1	-
Female	6	11	13	19	22	34	61	84	65	30	7	-
Black, both sexes	18	22	12	29	14	22	21	22	18	7	1	-
Male	9	15	9	14	7	13	8	7	5	3	-	-
Female	9	7	3	15	7	9	13	15	13	4	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	3	1	2	3	-	5	1	-	-	-
Male	-	1	2	1	1	2	-	2	1	-	-	-
Female	-	-	1	-	1	1	-	3	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Washington												
All races, both sexes	91	99	117	106	117	138	185	217	157	77	19	1
Male	72	80	83	79	73	79	83	97	53	22	4	1
Female	19	19	34	27	44	59	102	120	104	55	15	-
White, both sexes	83	87	105	96	102	130	164	206	150	75	18	1
Male	64	69	72	74	62	78	79	91	50	21	4	1
Female	19	18	33	22	40	52	85	115	100	54	14	-
Black, both sexes	4	4	6	5	8	2	5	2	2	1	1	-
Male	4	4	5	3	7	-	1	2	1	1	-	-
Female	-	-	1	2	1	2	4	-	1	-	1	-
American Indian 1/, both sexes	2	3	2	2	3	1	1	1	-	1	-	-
Male	2	3	2	1	2	-	-	-	-	-	-	-
Female	-	-	-	1	1	1	1	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	2	5	4	3	4	5	15	8	5	-	-	-
Male	2	4	4	1	2	1	3	4	2	-	-	-
Female	-	1	-	2	2	4	12	4	3	-	-	-
West Virginia												
All races, both sexes	25	25	32	22	35	42	72	70	59	18	4	-
Male	21	18	22	13	23	24	31	20	17	-	-	-
Female	4	7	10	9	12	18	41	50	42	18	4	-
White, both sexes	23	25	29	21	33	42	72	67	58	16	4	-
Male	19	18	19	12	22	24	31	20	16	-	-	-
Female	4	7	10	9	11	18	41	47	42	16	4	-
Black, both sexes	2	-	3	-	2	-	-	3	1	2	-	-
Male	2	-	3	-	1	-	-	-	1	-	-	-
Female	-	-	-	-	1	-	-	3	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Wisconsin												
All races, both sexes	62	75	91	61	78	95	116	144	122	29	7	-
Male	47	57	69	45	56	60	64	58	42	6	-	-
Female	15	18	22	16	22	35	52	86	80	23	7	-
White, both sexes	47	52	81	50	70	86	109	138	121	28	7	-
Male	35	43	63	38	54	54	61	55	42	6	-	-
Female	12	9	18	12	16	32	48	83	79	22	7	-
Black, both sexes	15	20	9	9	6	9	3	4	1	1	-	-
Male	12	12	5	7	2	6	1	2	-	-	-	-
Female	3	8	4	2	4	3	2	2	1	1	-	-
American Indian 1/, both sexes	-	3	1	2	-	-	3	-	-	-	-	-
Male	-	2	1	-	-	-	1	-	-	-	-	-
Female	-	1	-	2	-	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	-	1	2	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	2	-	-	1	-	-	-	-
Wyoming												
All races, both sexes	3	3	5	4	3	3	7	2	1	1	-	-
Male	2	3	4	2	2	3	3	-	1	-	-	-
Female	1	-	1	2	1	-	4	2	-	1	-	-
White, both sexes	3	3	5	4	3	3	6	2	1	1	-	-
Male	2	3	4	2	2	3	3	-	1	-	-	-
Female	1	-	1	2	1	-	3	2	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
United States												
All races, both sexes	210,240	11	4	7	3	20	57	128	292	674	1,639	3,333
Male	108,778	4	3	6	1	14	46	93	229	513	1,266	2,511
Female	101,462	7	1	1	2	6	11	35	63	161	373	822
White, both sexes	186,480	7	2	6	2	13	38	89	218	506	1,268	2,628
Male	96,888	3	1	5	-	9	31	66	172	385	998	2,041
Female	89,592	4	1	1	2	4	7	23	46	121	270	587
Black, both sexes	19,122	2	1	1	-	7	17	34	63	135	296	612
Male	9,313	1	1	1	-	5	13	23	47	96	204	394
Female	9,809	1	-	-	-	2	4	11	16	39	92	218
American Indian 1/, both sexes	800	1	1	-	-	-	1	2	4	6	25	25
Male	455	-	1	-	-	-	1	1	3	5	19	21
Female	345	1	-	-	-	-	-	1	1	1	6	4
Asian or Pacific Islander 2/, both sexes	3,838	1	-	-	1	-	1	3	7	27	50	68
Male	2,122	-	-	-	1	-	1	3	7	27	45	55
Female	1,716	1	-	-	-	-	-	-	-	-	5	13
Alabama												
All races, both sexes	2,336	-	-	-	-	-	-	1	4	12	23	43
Male	1,233	-	-	-	-	-	-	1	3	8	16	33
Female	1,103	-	-	-	-	-	-	-	1	4	7	10
White, both sexes	1,918	-	-	-	-	-	-	-	4	8	14	34
Male	1,011	-	-	-	-	-	-	-	3	5	9	25
Female	907	-	-	-	-	-	-	-	1	3	5	9
Black, both sexes	412	-	-	-	-	-	-	1	-	4	7	9
Male	217	-	-	-	-	-	-	1	-	3	6	8
Female	195	-	-	-	-	-	-	-	-	1	1	1
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	2	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Alaska												
All races, both sexes	150	-	-	-	-	-	-	-	-	-	3	1
Male	84	-	-	-	-	-	-	-	-	-	3	1
Female	66	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	119	-	-	-	-	-	-	-	-	-	3	-
Male	68	-	-	-	-	-	-	-	-	-	3	-
Female	51	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	3,790	-	-	-	1	-	4	-	4	12	30	59
Male	2,090	-	-	-	1	-	4	-	3	11	25	44
Female	1,700	-	-	-	-	-	-	-	1	1	5	15
White, both sexes	3,636	-	-	-	-	-	2	-	4	12	26	55
Male	2,013	-	-	-	-	-	2	-	3	11	23	41
Female	1,623	-	-	-	-	-	-	-	1	1	3	14
Black, both sexes	80	-	-	-	-	-	2	-	-	-	-	2
Male	39	-	-	-	-	-	2	-	-	-	-	1
Female	41	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	59	-	-	-	-	-	-	-	-	-	4	2
Male	31	-	-	-	-	-	-	-	-	-	2	2
Female	28	-	-	-	-	-	-	-	-	-	2	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	1	-	-	-	-	-	-	-
Male	7	-	-	-	1	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Arkansas												
All races, both sexes	1,622	-	1	-	-	-	-	1	3	3	11	19
Male	821	-	1	-	-	-	-	-	3	2	10	14
Female	801	-	-	-	-	-	-	1	-	1	1	5
White, both sexes	1,438	-	-	-	-	-	-	-	3	3	8	13
Male	740	-	-	-	-	-	-	-	3	2	7	9
Female	698	-	-	-	-	-	-	-	-	1	1	4
Black, both sexes	178	-	1	-	-	-	-	1	-	-	3	6
Male	77	-	1	-	-	-	-	-	-	-	3	5
Female	101	-	-	-	-	-	-	1	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	23,838	-	-	-	-	3	5	10	38	74	156	330
Male	12,241	-	-	-	-	3	5	9	32	55	127	254
Female	11,597	-	-	-	-	-	-	1	6	19	29	76
White, both sexes	20,388	-	-	-	-	3	5	8	27	47	114	254
Male	10,439	-	-	-	-	3	5	7	24	33	93	197
Female	9,949	-	-	-	-	-	-	1	3	14	21	57
Black, both sexes	1,581	-	-	-	-	-	-	2	7	14	21	44
Male	794	-	-	-	-	-	-	2	4	9	15	27
Female	787	-	-	-	-	-	-	-	3	5	6	17
American Indian 1/, both sexes	108	-	-	-	-	-	-	-	1	1	2	5
Male	55	-	-	-	-	-	-	-	1	1	2	5
Female	53	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1,761	-	-	-	-	-	-	-	3	12	19	27
Male	953	-	-	-	-	-	-	-	3	12	17	25
Female	808	-	-	-	-	-	-	-	-	-	2	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Colorado												
All races, both sexes	2,026	-	-	1	-	-	-	2	3	9	20	48
Male	1,146	-	-	1	-	-	-	1	3	7	16	38
Female	880	-	-	-	-	-	-	1	-	2	4	10
White, both sexes	1,963	-	-	1	-	-	-	2	3	9	20	45
Male	1,110	-	-	1	-	-	-	1	3	7	16	36
Female	853	-	-	-	-	-	-	1	-	2	4	9
Black, both sexes	47	-	-	-	-	-	-	-	-	-	-	3
Male	25	-	-	-	-	-	-	-	-	-	-	2
Female	22	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	2,483	1	-	-	-	-	-	2	4	8	26	26
Male	1,221	1	-	-	-	-	-	2	3	7	20	20
Female	1,262	-	-	-	-	-	-	-	1	1	6	6
White, both sexes	2,347	1	-	-	-	-	-	2	2	7	20	21
Male	1,150	1	-	-	-	-	-	2	1	6	16	17
Female	1,197	-	-	-	-	-	-	-	1	1	4	4
Black, both sexes	120	-	-	-	-	-	-	-	2	1	5	5
Male	60	-	-	-	-	-	-	-	2	1	3	3
Female	60	-	-	-	-	-	-	-	-	-	2	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	-	1	-
Male	10	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Delaware												
All races, both sexes	624	-	-	-	-	-	-	-	-	2	5	13
Male	337	-	-	-	-	-	-	-	-	2	4	9
Female	287	-	-	-	-	-	-	-	-	-	1	4
White, both sexes	544	-	-	-	-	-	-	-	-	1	3	10
Male	299	-	-	-	-	-	-	-	-	1	2	7
Female	245	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	73	-	-	-	-	-	-	-	-	1	2	3
Male	33	-	-	-	-	-	-	-	-	1	2	2
Female	40	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	274	-	-	-	-	-	-	-	-	1	3	5
Male	143	-	-	-	-	-	-	-	-	1	2	5
Female	131	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	56	-	-	-	-	-	-	-	-	-	1	-
Male	35	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	216	-	-	-	-	-	-	-	-	1	2	5
Male	107	-	-	-	-	-	-	-	-	1	2	5
Female	109	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Florida												
All races, both sexes	17,510	-	-	-	-	1	6	13	25	50	125	272
Male	9,479	-	-	-	-	-	5	11	20	38	94	209
Female	8,031	-	-	-	-	1	1	2	5	12	31	63
White, both sexes	15,908	-	-	-	-	1	3	7	14	35	98	223
Male	8,663	-	-	-	-	-	3	5	12	27	76	178
Female	7,245	-	-	-	-	1	-	2	2	8	22	45
Black, both sexes	1,503	-	-	-	-	-	3	4	11	15	24	48
Male	760	-	-	-	-	-	2	4	8	11	16	30
Female	743	-	-	-	-	-	1	-	3	4	8	18
American Indian 1/, both sexes	14	-	-	-	-	-	-	-	-	-	1	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	85	-	-	-	-	-	-	2	-	-	2	1
Male	45	-	-	-	-	-	-	2	-	-	2	1
Female	40	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	3,940	-	-	-	-	4	4	4	6	30	63	118
Male	2,235	-	-	-	-	4	4	4	5	21	47	85
Female	1,705	-	-	-	-	-	-	-	1	9	16	33
White, both sexes	3,055	-	-	-	-	3	2	3	5	23	40	77
Male	1,775	-	-	-	-	3	2	3	4	16	30	57
Female	1,280	-	-	-	-	-	-	-	1	7	10	20
Black, both sexes	856	-	-	-	-	1	2	1	1	7	23	39
Male	444	-	-	-	-	1	2	1	1	5	17	27
Female	412	-	-	-	-	-	-	-	-	2	6	12
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	27	-	-	-	-	-	-	-	-	-	-	2
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Hawaii												
All races, both sexes	601	-	-	-	-	-	-	1	1	1	8	13
Male	381	-	-	-	-	-	-	1	-	1	5	8
Female	220	-	-	-	-	-	-	-	1	-	3	5
White, both sexes	169	-	-	-	-	-	-	-	1	1	4	5
Male	111	-	-	-	-	-	-	-	-	1	2	4
Female	58	-	-	-	-	-	-	-	1	-	2	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	428	-	-	-	-	-	-	1	-	-	4	8
Male	266	-	-	-	-	-	-	1	-	-	3	4
Female	162	-	-	-	-	-	-	-	-	-	1	4
Idaho												
All races, both sexes	673	-	-	-	-	-	-	-	1	1	3	14
Male	372	-	-	-	-	-	-	-	1	1	1	9
Female	301	-	-	-	-	-	-	-	-	-	2	5
White, both sexes	667	-	-	-	-	-	-	-	-	-	3	14
Male	367	-	-	-	-	-	-	-	-	-	1	9
Female	300	-	-	-	-	-	-	-	-	-	2	5
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Illinois												
All races, both sexes	8,470	1	-	1	-	1	5	16	18	27	88	147
Male	4,269	-	-	1	-	1	5	10	10	23	71	106
Female	4,201	1	-	-	-	-	-	6	8	4	17	41
White, both sexes	7,407	1	-	1	-	1	4	13	15	19	63	111
Male	3,725	-	-	1	-	1	4	8	9	16	52	81
Female	3,682	1	-	-	-	-	-	5	6	3	11	30
Black, both sexes	968	-	-	-	-	-	-	3	3	7	24	35
Male	495	-	-	-	-	-	-	2	1	6	18	24
Female	473	-	-	-	-	-	-	1	2	1	6	11
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	89	-	-	-	-	-	1	-	-	1	1	-
Male	47	-	-	-	-	-	1	-	-	1	1	-
Female	42	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	4,585	1	-	1	-	-	1	4	6	25	54	98
Male	2,382	1	-	1	-	-	1	4	5	15	45	77
Female	2,203	-	-	-	-	-	-	-	1	10	9	21
White, both sexes	4,229	1	-	1	-	-	1	4	5	20	42	81
Male	2,188	1	-	1	-	-	1	4	4	13	37	66
Female	2,041	-	-	-	-	-	-	-	1	7	5	15
Black, both sexes	350	-	-	-	-	-	-	-	1	5	12	17
Male	191	-	-	-	-	-	-	-	1	2	8	11
Female	159	-	-	-	-	-	-	-	-	3	4	6
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Iowa												
All races, both sexes	2,603	1	-	-	-	-	-	1	-	6	15	31
Male	1,353	-	-	-	-	-	-	1	-	5	12	21
Female	1,250	1	-	-	-	-	-	-	-	1	3	10
White, both sexes	2,572	1	-	-	-	-	-	1	-	6	15	29
Male	1,336	-	-	-	-	-	-	1	-	5	12	19
Female	1,236	1	-	-	-	-	-	-	-	1	3	10
Black, both sexes	26	-	-	-	-	-	-	-	-	-	-	2
Male	14	-	-	-	-	-	-	-	-	-	-	2
Female	12	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1,953	-	-	-	1	-	3	-	8	11	25	37
Male	1,059	-	-	-	-	-	3	-	6	10	17	30
Female	894	-	-	-	1	-	-	-	2	1	8	7
White, both sexes	1,842	-	-	-	1	-	3	-	8	9	23	34
Male	1,001	-	-	-	-	-	3	-	6	9	16	28
Female	841	-	-	-	1	-	-	-	2	-	7	6
Black, both sexes	89	-	-	-	-	-	-	-	-	1	-	2
Male	43	-	-	-	-	-	-	-	-	-	-	1
Female	46	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	17	-	-	-	-	-	-	-	-	-	1	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	1	1	-
Male	3	-	-	-	-	-	-	-	-	1	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Kentucky												
All races, both sexes	2,560	-	-	-	-	-	3	3	7	14	38	40
Male	1,323	-	-	-	-	-	3	2	6	10	28	33
Female	1,237	-	-	-	-	-	-	1	1	4	10	7
White, both sexes	2,429	-	-	-	-	-	2	3	6	12	32	37
Male	1,261	-	-	-	-	-	2	2	5	9	25	31
Female	1,168	-	-	-	-	-	-	1	1	3	7	6
Black, both sexes	128	-	-	-	-	-	1	-	1	2	6	3
Male	61	-	-	-	-	-	1	-	1	1	3	2
Female	67	-	-	-	-	-	-	-	-	1	3	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	2,107	1	-	-	-	1	1	4	4	14	25	57
Male	1,169	-	-	-	-	1	-	4	3	9	17	33
Female	938	1	-	-	-	-	1	-	1	5	8	24
White, both sexes	1,616	-	-	-	-	-	1	3	4	8	16	40
Male	918	-	-	-	-	-	-	3	3	7	10	25
Female	698	-	-	-	-	-	1	-	1	1	6	15
Black, both sexes	484	1	-	-	-	1	-	1	-	6	9	17
Male	248	-	-	-	-	1	-	1	-	2	7	8
Female	236	1	-	-	-	-	-	-	-	4	2	9
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Maine												
All races, both sexes	932	-	-	-	-	-	1	-	-	3	12	17
Male	537	-	-	-	-	-	-	-	-	2	10	13
Female	395	-	-	-	-	-	1	-	-	1	2	4
White, both sexes	926	-	-	-	-	-	1	-	-	3	11	16
Male	532	-	-	-	-	-	-	-	-	2	9	12
Female	394	-	-	-	-	-	1	-	-	1	2	4
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	2,993	-	-	1	-	-	-	-	3	7	15	40
Male	1,524	-	-	1	-	-	-	-	2	5	11	25
Female	1,469	-	-	-	-	-	-	-	1	2	4	15
White, both sexes	2,347	-	-	-	-	-	-	-	-	6	9	23
Male	1,212	-	-	-	-	-	-	-	-	4	6	15
Female	1,135	-	-	-	-	-	-	-	-	2	3	8
Black, both sexes	600	-	-	1	-	-	-	-	3	1	6	17
Male	286	-	-	1	-	-	-	-	2	1	5	10
Female	314	-	-	-	-	-	-	-	1	-	1	7
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	43	-	-	-	-	-	-	-	-	-	-	-
Male	24	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Massachusetts												
All races, both sexes	4,272	-	-	-	-	1	-	-	4	10	21	59
Male	2,144	-	-	-	-	-	-	-	2	8	16	52
Female	2,128	-	-	-	-	1	-	-	2	2	5	7
White, both sexes	4,146	-	-	-	-	1	-	-	4	10	18	54
Male	2,086	-	-	-	-	-	-	-	2	8	15	48
Female	2,060	-	-	-	-	1	-	-	2	2	3	6
Black, both sexes	96	-	-	-	-	-	-	-	-	-	1	3
Male	43	-	-	-	-	-	-	-	-	-	-	3
Female	53	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	28	-	-	-	-	-	-	-	-	-	2	2
Male	14	-	-	-	-	-	-	-	-	-	1	1
Female	14	-	-	-	-	-	-	-	-	-	1	1
Michigan												
All races, both sexes	8,094	1	-	-	-	-	1	1	13	18	48	108
Male	4,112	-	-	-	-	-	1	-	11	14	37	86
Female	3,982	1	-	-	-	-	-	1	2	4	11	22
White, both sexes	7,351	-	-	-	-	-	-	-	8	11	40	92
Male	3,746	-	-	-	-	-	-	-	6	10	33	74
Female	3,605	-	-	-	-	-	-	-	2	1	7	18
Black, both sexes	681	-	-	-	-	-	1	1	5	6	5	14
Male	335	-	-	-	-	-	1	-	5	3	1	10
Female	346	-	-	-	-	-	-	1	-	3	4	4
American Indian 1/, both sexes	36	1	-	-	-	-	-	-	-	1	2	-
Male	19	-	-	-	-	-	-	-	-	1	2	-
Female	17	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	26	-	-	-	-	-	-	-	-	-	1	2
Male	12	-	-	-	-	-	-	-	-	-	1	2
Female	14	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Minnesota												
All races, both sexes	2,680	-	-	-	-	1	-	2	6	12	27	66
Male	1,542	-	-	-	-	1	-	2	6	11	21	51
Female	1,138	-	-	-	-	-	-	-	-	1	6	15
White, both sexes	2,598	-	-	-	-	1	-	1	4	10	23	60
Male	1,490	-	-	-	-	1	-	1	4	9	17	48
Female	1,108	-	-	-	-	-	-	-	-	1	6	12
Black, both sexes	39	-	-	-	-	-	-	1	1	1	2	6
Male	24	-	-	-	-	-	-	1	1	1	2	3
Female	15	-	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes	28	-	-	-	-	-	-	-	1	-	1	-
Male	18	-	-	-	-	-	-	-	1	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	-	1	1	-
Male	10	-	-	-	-	-	-	-	-	1	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1,690	-	-	-	-	-	1	4	5	15	42	57
Male	912	-	-	-	-	-	1	3	3	10	27	36
Female	778	-	-	-	-	-	-	1	2	5	15	21
White, both sexes	1,204	-	-	-	-	-	-	-	2	12	24	29
Male	656	-	-	-	-	-	-	-	2	8	14	19
Female	548	-	-	-	-	-	-	-	-	4	10	10
Black, both sexes	477	-	-	-	-	-	1	4	3	3	18	26
Male	248	-	-	-	-	-	1	3	1	2	13	15
Female	229	-	-	-	-	-	-	1	2	1	5	11
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Missouri												
All races, both sexes	4,752	-	-	-	-	-	-	2	5	15	49	81
Male	2,407	-	-	-	-	-	-	2	4	12	36	61
Female	2,345	-	-	-	-	-	-	-	1	3	13	20
White, both sexes	4,256	-	-	-	-	-	-	2	4	9	42	59
Male	2,167	-	-	-	-	-	-	2	3	6	32	47
Female	2,089	-	-	-	-	-	-	-	1	3	10	12
Black, both sexes	477	-	-	-	-	-	-	-	1	4	6	22
Male	230	-	-	-	-	-	-	-	1	4	3	14
Female	247	-	-	-	-	-	-	-	-	-	3	8
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	2	1	-
Male	6	-	-	-	-	-	-	-	-	2	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	540	-	-	-	-	-	-	-	-	1	11	12
Male	340	-	-	-	-	-	-	-	-	1	9	10
Female	200	-	-	-	-	-	-	-	-	-	2	2
White, both sexes	516	-	-	-	-	-	-	-	-	1	11	12
Male	325	-	-	-	-	-	-	-	-	1	9	10
Female	191	-	-	-	-	-	-	-	-	-	2	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Nebraska												
All races, both sexes	1,233	1	-	-	-	-	1	1	1	3	11	20
Male	665	-	-	-	-	-	1	1	1	2	10	16
Female	568	1	-	-	-	-	-	-	-	1	1	4
White, both sexes	1,178	1	-	-	-	-	1	1	1	1	9	18
Male	631	-	-	-	-	-	1	1	1	1	8	14
Female	547	1	-	-	-	-	-	-	-	-	1	4
Black, both sexes	44	-	-	-	-	-	-	-	-	1	2	2
Male	28	-	-	-	-	-	-	-	-	1	2	2
Female	16	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	911	-	-	-	1	-	1	2	-	6	9	18
Male	564	-	-	-	-	-	1	2	-	3	6	11
Female	347	-	-	-	1	-	-	-	-	3	3	7
White, both sexes	829	-	-	-	1	-	-	2	-	6	7	15
Male	523	-	-	-	-	-	-	2	-	3	4	11
Female	306	-	-	-	1	-	-	-	-	3	3	4
Black, both sexes	50	-	-	-	-	-	1	-	-	-	1	2
Male	22	-	-	-	-	-	1	-	-	-	1	-
Female	28	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	25	-	-	-	-	-	-	-	-	-	1	1
Male	14	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
New Hampshire												
All races, both sexes	889	-	-	-	-	-	-	-	2	1	3	10
Male	429	-	-	-	-	-	-	-	-	-	2	7
Female	460	-	-	-	-	-	-	-	2	1	1	3
White, both sexes	885	-	-	-	-	-	-	-	2	1	3	9
Male	427	-	-	-	-	-	-	-	-	-	2	6
Female	458	-	-	-	-	-	-	-	2	1	1	3
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	7,105	1	-	-	-	-	-	2	6	11	34	93
Male	3,476	-	-	-	-	-	-	-	5	10	31	74
Female	3,629	1	-	-	-	-	-	2	1	1	3	19
White, both sexes	6,420	-	-	-	-	-	-	2	6	6	27	75
Male	3,131	-	-	-	-	-	-	-	5	5	24	61
Female	3,289	-	-	-	-	-	-	2	1	1	3	14
Black, both sexes	557	-	-	-	-	-	-	-	-	4	6	14
Male	273	-	-	-	-	-	-	-	-	4	6	9
Female	284	-	-	-	-	-	-	-	-	-	-	5
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	123	1	-	-	-	-	-	-	-	1	1	4
Male	72	-	-	-	-	-	-	-	-	1	1	4
Female	51	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
New Mexico												
All races, both sexes	1,013	-	1	-	-	-	-	-	-	3	3	11
Male	536	-	1	-	-	-	-	-	-	3	3	9
Female	477	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	960	-	-	-	-	-	-	-	-	3	3	11
Male	507	-	-	-	-	-	-	-	-	3	3	9
Female	453	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	30	-	1	-	-	-	-	-	-	-	-	-
Male	19	-	1	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	25,361	1	-	1	-	4	5	8	13	40	92	218
Male	11,395	1	-	1	-	1	3	4	10	27	70	159
Female	13,966	-	-	-	-	3	2	4	3	13	22	59
White, both sexes	21,504	1	-	1	-	1	3	4	8	30	64	153
Male	9,765	1	-	1	-	-	2	3	6	23	49	119
Female	11,739	-	-	-	-	1	1	1	2	7	15	34
Black, both sexes	3,166	-	-	-	-	3	2	4	3	9	18	57
Male	1,264	-	-	-	-	1	1	1	2	3	12	33
Female	1,902	-	-	-	-	2	1	3	1	6	6	24
American Indian 1/, both sexes	24	-	-	-	-	-	-	-	-	-	1	1
Male	14	-	-	-	-	-	-	-	-	-	1	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	667	-	-	-	-	-	-	-	2	1	9	7
Male	352	-	-	-	-	-	-	-	2	1	8	6
Female	315	-	-	-	-	-	-	-	-	-	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
North Carolina												
All races, both sexes	6,319	-	-	-	-	-	1	3	17	36	90	173
Male	3,513	-	-	-	-	-	1	1	15	27	66	128
Female	2,806	-	-	-	-	-	-	2	2	9	24	45
White, both sexes	5,065	-	-	-	-	-	1	1	8	27	62	118
Male	2,845	-	-	-	-	-	1	-	8	21	50	95
Female	2,220	-	-	-	-	-	-	1	-	6	12	23
Black, both sexes	1,168	-	-	-	-	-	-	1	8	7	24	52
Male	608	-	-	-	-	-	-	-	6	4	13	32
Female	560	-	-	-	-	-	-	1	2	3	11	20
American Indian 1/, both sexes	66	-	-	-	-	-	-	1	1	-	4	2
Male	46	-	-	-	-	-	-	1	1	-	3	1
Female	20	-	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	20	-	-	-	-	-	-	-	-	2	-	1
Male	14	-	-	-	-	-	-	-	-	2	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
North Dakota												
All races, both sexes	527	-	-	-	-	-	1	2	1	1	3	10
Male	276	-	-	-	-	-	1	-	1	1	2	8
Female	251	-	-	-	-	-	-	2	-	-	1	2
White, both sexes	515	-	-	-	-	-	-	2	1	1	2	10
Male	269	-	-	-	-	-	-	-	1	1	1	8
Female	246	-	-	-	-	-	-	2	-	-	1	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	11	-	-	-	-	-	1	-	-	-	1	-
Male	6	-	-	-	-	-	1	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Ohio												
All races, both sexes	10,609	-	1	-	-	1	3	6	20	38	58	171
Male	5,448	-	1	-	-	-	3	4	15	30	44	131
Female	5,161	-	-	-	-	1	-	2	5	8	14	40
White, both sexes	9,659	-	1	-	-	1	2	4	15	32	47	143
Male	4,960	-	1	-	-	-	2	3	11	24	38	116
Female	4,699	-	-	-	-	1	-	1	4	8	9	27
Black, both sexes	916	-	-	-	-	-	1	2	5	5	11	28
Male	464	-	-	-	-	-	1	1	4	5	6	15
Female	452	-	-	-	-	-	-	1	1	-	5	13
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	29	-	-	-	-	-	-	-	-	1	-	-
Male	20	-	-	-	-	-	-	-	-	1	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2,895	-	-	-	-	-	2	2	2	5	18	55
Male	1,469	-	-	-	-	-	-	1	2	5	13	37
Female	1,426	-	-	-	-	-	2	1	-	-	5	18
White, both sexes	2,579	-	-	-	-	-	2	-	2	4	15	39
Male	1,306	-	-	-	-	-	-	-	2	4	10	25
Female	1,273	-	-	-	-	-	2	-	-	-	5	14
Black, both sexes	166	-	-	-	-	-	-	1	-	1	2	13
Male	88	-	-	-	-	-	-	1	-	1	2	10
Female	78	-	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes	142	-	-	-	-	-	-	1	-	-	1	3
Male	72	-	-	-	-	-	-	-	-	-	1	2
Female	70	-	-	-	-	-	-	1	-	-	-	1
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Oregon												
All races, both sexes	2,390	-	-	-	-	-	-	-	9	11	25	53
Male	1,440	-	-	-	-	-	-	-	8	10	22	46
Female	950	-	-	-	-	-	-	-	1	1	3	7
White, both sexes	2,319	-	-	-	-	-	-	-	8	9	23	50
Male	1,397	-	-	-	-	-	-	-	7	8	20	44
Female	922	-	-	-	-	-	-	-	1	1	3	6
Black, both sexes	24	-	-	-	-	-	-	-	-	1	-	2
Male	15	-	-	-	-	-	-	-	-	1	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	16	-	-	-	-	-	-	-	-	-	1	-
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	31	-	-	-	-	-	-	-	1	1	1	1
Male	20	-	-	-	-	-	-	-	1	1	1	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	10,366	1	1	-	-	1	3	4	11	22	52	113
Male	5,229	1	-	-	-	1	-	2	10	17	42	79
Female	5,137	-	1	-	-	-	3	2	1	5	10	34
White, both sexes	9,680	-	1	-	-	-	2	4	7	19	40	97
Male	4,916	-	-	-	-	-	-	2	6	14	33	70
Female	4,764	-	1	-	-	-	2	2	1	5	7	27
Black, both sexes	644	1	-	-	-	1	1	-	4	3	10	15
Male	286	1	-	-	-	1	-	-	4	3	7	8
Female	358	-	-	-	-	-	1	-	-	-	3	7
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	39	-	-	-	-	-	-	-	-	-	2	1
Male	25	-	-	-	-	-	-	-	-	-	2	1
Female	14	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Rhode Island												
All races, both sexes	987	-	-	-	-	-	-	-	2	3	3	9
Male	410	-	-	-	-	-	-	-	2	2	3	8
Female	577	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	959	-	-	-	-	-	-	-	2	3	3	8
Male	398	-	-	-	-	-	-	-	2	2	3	7
Female	561	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	2,370	-	-	1	-	-	1	1	1	17	30	55
Male	1,324	-	-	1	-	-	-	1	1	12	21	39
Female	1,046	-	-	-	-	-	1	-	-	5	9	16
White, both sexes	1,847	-	-	1	-	-	-	-	1	9	24	35
Male	1,052	-	-	1	-	-	-	-	1	5	16	25
Female	795	-	-	-	-	-	-	-	-	4	8	10
Black, both sexes	510	-	-	-	-	-	1	1	-	7	6	18
Male	262	-	-	-	-	-	-	1	-	6	5	12
Female	248	-	-	-	-	-	1	-	-	1	1	6
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	1	-	1
Male	7	-	-	-	-	-	-	-	-	1	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
South Dakota												
All races, both sexes	598	-	-	-	-	-	-	-	1	2	6	7
Male	336	-	-	-	-	-	-	-	-	2	6	5
Female	262	-	-	-	-	-	-	-	1	-	-	2
White, both sexes	566	-	-	-	-	-	-	-	1	1	4	4
Male	321	-	-	-	-	-	-	-	-	1	4	3
Female	245	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	30	-	-	-	-	-	-	-	-	1	2	3
Male	15	-	-	-	-	-	-	-	-	1	2	2
Female	15	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	4,082	-	-	-	-	-	-	5	9	11	46	83
Male	2,073	-	-	-	-	-	-	3	7	8	35	57
Female	2,009	-	-	-	-	-	-	2	2	3	11	26
White, both sexes	3,611	-	-	-	-	-	-	3	7	9	40	68
Male	1,845	-	-	-	-	-	-	2	5	6	32	47
Female	1,766	-	-	-	-	-	-	1	2	3	8	21
Black, both sexes	461	-	-	-	-	-	-	2	2	2	6	14
Male	221	-	-	-	-	-	-	1	2	2	3	9
Female	240	-	-	-	-	-	-	1	-	-	3	5
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Texas												
All races, both sexes	10,647	1	-	1	-	1	3	10	9	40	91	185
Male	5,662	-	-	-	-	1	3	8	4	29	72	148
Female	4,985	1	-	1	-	-	-	2	5	11	19	37
White, both sexes	9,511	1	-	1	-	1	2	8	8	33	72	149
Male	5,080	-	-	-	-	1	2	7	4	24	58	119
Female	4,431	1	-	1	-	-	-	1	4	9	14	30
Black, both sexes	1,040	-	-	-	-	-	1	2	1	6	16	33
Male	534	-	-	-	-	-	1	1	-	4	11	26
Female	506	-	-	-	-	-	-	1	1	2	5	7
American Indian 1/, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	81	-	-	-	-	-	-	-	-	1	3	3
Male	40	-	-	-	-	-	-	-	-	1	3	3
Female	41	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	845	-	-	-	-	-	-	2	3	8	11	21
Male	522	-	-	-	-	-	-	1	3	7	9	17
Female	323	-	-	-	-	-	-	1	-	1	2	4
White, both sexes	824	-	-	-	-	-	-	2	3	7	10	17
Male	508	-	-	-	-	-	-	1	3	6	9	14
Female	316	-	-	-	-	-	-	1	-	1	1	3
Black, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	2
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	1	-	2
Male	6	-	-	-	-	-	-	-	-	1	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Vermont												
All races, both sexes	336	-	-	-	-	-	-	-	-	-	-	1
Male	174	-	-	-	-	-	-	-	-	-	-	1
Female	162	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	334	-	-	-	-	-	-	-	-	-	-	1
Male	173	-	-	-	-	-	-	-	-	-	-	1
Female	161	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	3,634	-	-	-	-	-	1	6	4	17	25	64
Male	2,038	-	-	-	-	-	1	5	4	14	18	49
Female	1,596	-	-	-	-	-	-	1	-	3	7	15
White, both sexes	2,950	-	-	-	-	-	1	4	3	9	15	41
Male	1,670	-	-	-	-	-	1	3	3	7	10	34
Female	1,280	-	-	-	-	-	-	1	-	2	5	7
Black, both sexes	628	-	-	-	-	-	-	2	1	8	10	21
Male	337	-	-	-	-	-	-	2	1	7	8	13
Female	291	-	-	-	-	-	-	-	-	1	2	8
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	53	-	-	-	-	-	-	-	-	-	-	2
Male	31	-	-	-	-	-	-	-	-	-	-	2
Female	22	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Washington												
All races, both sexes	3,711	-	-	-	-	1	-	-	1	3	23	31
Male	2,044	-	-	-	-	1	-	-	1	3	17	24
Female	1,667	-	-	-	-	-	-	-	-	-	6	7
White, both sexes	3,501	-	-	-	-	-	-	-	1	1	20	27
Male	1,924	-	-	-	-	-	-	-	1	1	15	22
Female	1,577	-	-	-	-	-	-	-	-	-	5	5
Black, both sexes	68	-	-	-	-	1	-	-	-	1	3	3
Male	46	-	-	-	-	1	-	-	-	1	2	2
Female	22	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	33	-	-	-	-	-	-	-	-	-	-	1
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	109	-	-	-	-	-	-	-	-	1	-	-
Male	57	-	-	-	-	-	-	-	-	1	-	-
Female	52	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	1,963	-	-	-	-	-	-	1	8	11	37	66
Male	1,066	-	-	-	-	-	-	1	6	9	28	49
Female	897	-	-	-	-	-	-	-	2	2	9	17
White, both sexes	1,907	-	-	-	-	-	-	1	8	10	36	63
Male	1,038	-	-	-	-	-	-	1	6	8	28	46
Female	869	-	-	-	-	-	-	-	2	2	8	17
Black, both sexes	53	-	-	-	-	-	-	-	-	1	1	3
Male	28	-	-	-	-	-	-	-	-	1	-	3
Female	25	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Wisconsin												
All races, both sexes	3,093	-	-	-	-	-	-	1	4	3	20	52
Male	1,681	-	-	-	-	-	-	1	3	3	17	43
Female	1,412	-	-	-	-	-	-	-	1	-	3	9
White, both sexes	2,978	-	-	-	-	-	-	1	3	2	17	46
Male	1,630	-	-	-	-	-	-	1	3	2	15	39
Female	1,348	-	-	-	-	-	-	-	-	-	2	7
Black, both sexes	82	-	-	-	-	-	-	-	-	-	2	5
Male	31	-	-	-	-	-	-	-	-	-	1	4
Female	51	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	23	-	-	-	-	-	-	-	1	1	1	-
Male	16	-	-	-	-	-	-	-	-	1	1	-
Female	7	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
Wyoming												
All races, both sexes	258	-	-	-	-	-	-	1	-	1	3	3
Male	152	-	-	-	-	-	-	1	-	-	2	3
Female	106	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	252	-	-	-	-	-	-	1	-	1	2	3
Male	148	-	-	-	-	-	-	1	-	-	1	3
Female	104	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
United States												
All races, both sexes	5,304	7,607	9,988	12,238	16,724	25,971	37,310	41,735	30,893	13,362	2,933	7
Male	4,000	5,629	7,033	8,294	10,631	15,294	19,649	18,757	10,974	3,310	516	5
Female	1,304	1,978	2,955	3,944	6,093	10,677	17,661	22,978	19,919	10,052	2,417	2
White, both sexes	4,166	6,046	8,235	10,192	14,196	22,900	33,798	38,409	28,761	12,363	2,634	3
Male	3,212	4,578	5,929	7,040	9,240	13,710	18,114	17,464	10,331	3,098	459	2
Female	954	1,468	2,306	3,152	4,956	9,190	15,684	20,945	18,430	9,265	2,175	1
Black, both sexes	1,002	1,330	1,491	1,667	2,054	2,464	2,743	2,557	1,605	799	239	3
Male	676	872	934	999	1,094	1,243	1,161	927	443	137	40	2
Female	326	458	557	668	960	1,221	1,582	1,630	1,162	662	199	1
American Indian 1/, both sexes	39	59	68	93	82	111	113	86	52	25	7	-
Male	31	42	43	62	49	64	52	40	14	6	1	-
Female	8	17	25	31	33	47	61	46	38	19	6	-
Asian or Pacific Islander 2/, both sexes	97	172	194	286	392	496	656	683	475	175	53	1
Male	81	137	127	193	248	277	322	326	186	69	16	1
Female	16	35	67	93	144	219	334	357	289	106	37	-
Alabama												
All races, both sexes	62	106	149	184	224	307	439	400	285	79	18	-
Male	53	73	104	115	139	188	222	170	87	16	5	-
Female	9	33	45	69	85	119	217	230	198	63	13	-
White, both sexes	35	72	115	146	182	254	375	360	242	63	14	-
Male	30	49	80	89	121	154	195	158	75	15	3	-
Female	5	23	35	57	61	100	180	202	167	48	11	-
Black, both sexes	27	34	33	38	41	53	62	40	43	16	4	-
Male	23	24	23	26	17	34	25	12	12	1	2	-
Female	4	10	10	12	24	19	37	28	31	15	2	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	2	-	-	-	-	-
Male	-	-	-	-	1	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Alaska												
All races, both sexes	5	9	9	11	18	23	27	26	11	7	-	-
Male	5	6	6	7	11	14	19	8	2	2	-	-
Female	-	3	3	4	7	9	8	18	9	5	-	-
White, both sexes	5	7	9	10	12	16	23	18	10	6	-	-
Male	5	5	6	7	6	11	16	5	2	2	-	-
Female	-	2	3	3	6	5	7	13	8	4	-	-
Black, both sexes	-	-	-	-	2	2	-	1	-	-	-	-
Male	-	-	-	-	2	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	1	-	-	-	-
American Indian 1/, both sexes	-	2	-	1	3	5	4	6	-	1	-	-
Male	-	1	-	-	2	3	3	3	-	-	-	-
Female	-	1	-	1	1	2	1	3	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	1	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Arizona												
All races, both sexes	94	136	199	212	321	501	684	783	533	181	36	-
Male	79	86	144	152	226	312	355	382	215	45	6	-
Female	15	50	55	60	95	189	329	401	318	136	30	-
White, both sexes	86	123	186	199	307	481	660	766	520	174	35	-
Male	73	80	135	146	217	303	347	373	210	43	6	-
Female	13	43	51	53	90	178	313	393	310	131	29	-
Black, both sexes	4	7	8	7	6	10	10	12	7	5	-	-
Male	2	3	7	4	3	3	4	6	3	1	-	-
Female	2	4	1	3	3	7	6	6	4	4	-	-
American Indian 1/, both sexes	4	4	4	6	8	8	11	3	4	1	-	-
Male	4	2	1	2	6	5	4	2	1	-	-	-
Female	-	2	3	4	2	3	7	1	3	1	-	-
Asian or Pacific Islander 2/, both sexes	-	2	1	-	-	2	3	2	2	1	1	-
Male	-	1	1	-	-	1	-	1	1	1	-	-
Female	-	1	-	-	-	1	3	1	1	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Arkansas												
All races, both sexes	28	56	76	117	156	205	287	323	227	84	25	-
Male	19	41	51	73	104	126	145	134	77	17	4	-
Female	9	15	25	44	52	79	142	189	150	67	21	-
White, both sexes	20	41	66	106	134	185	263	297	205	71	23	-
Male	15	32	45	67	92	117	137	126	70	15	3	-
Female	5	9	21	39	42	68	126	171	135	56	20	-
Black, both sexes	8	15	10	11	21	19	24	25	20	12	2	-
Male	4	9	6	6	12	9	8	7	5	1	1	-
Female	4	6	4	5	9	10	16	18	15	11	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	1	2	1	-	-
Male	-	-	-	-	-	-	-	1	2	1	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
California												
All races, both sexes	539	816	1,093	1,260	1,830	2,737	4,207	5,047	3,630	1,671	391	1
Male	410	595	754	821	1,155	1,586	2,102	2,418	1,364	473	77	1
Female	129	221	339	439	675	1,151	2,105	2,629	2,266	1,198	314	-
White, both sexes	409	624	888	1,001	1,471	2,324	3,676	4,443	3,272	1,485	337	-
Male	315	464	621	668	940	1,357	1,863	2,142	1,237	406	64	-
Female	94	160	267	333	531	967	1,813	2,301	2,035	1,079	273	-
Black, both sexes	87	120	124	139	179	193	206	219	120	82	24	-
Male	56	76	79	74	106	102	83	97	36	22	6	-
Female	31	44	45	65	73	91	123	122	84	60	18	-
American Indian 1/, both sexes	4	6	10	8	13	17	13	16	7	4	1	-
Male	4	6	4	7	6	7	6	4	2	-	-	-
Female	-	-	6	1	7	10	7	12	5	4	1	-
Asian or Pacific Islander 2/, both sexes	39	66	71	112	167	203	312	369	231	100	29	1
Male	35	49	50	72	103	120	150	175	89	45	7	1
Female	4	17	21	40	64	83	162	194	142	55	22	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Colorado												
All races, both sexes	69	87	101	122	170	270	319	402	268	112	23	-
Male	55	67	75	77	120	160	178	194	109	37	8	-
Female	14	20	26	45	50	110	141	208	159	75	15	-
White, both sexes	67	81	96	119	163	256	313	390	263	112	23	-
Male	54	62	71	75	116	154	175	189	105	37	8	-
Female	13	19	25	44	47	102	138	201	158	75	15	-
Black, both sexes	1	4	3	3	5	11	5	8	4	-	-	-
Male	1	3	2	2	3	4	2	3	3	-	-	-
Female	-	1	1	1	2	7	3	5	1	-	-	-
American Indian 1/, both sexes	1	2	2	-	-	1	1	-	-	-	-	-
Male	-	2	2	-	-	1	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	2	-	4	1	-	-	-
Male	-	-	-	-	1	1	-	2	1	-	-	-
Female	-	-	-	-	1	1	-	2	-	-	-	-
Connecticut												
All races, both sexes	39	67	78	91	149	304	458	572	409	203	46	-
Male	25	51	58	60	98	180	243	258	142	48	5	-
Female	14	16	20	31	51	124	215	314	267	155	41	-
White, both sexes	33	62	63	79	138	287	433	556	399	200	44	-
Male	21	47	49	54	91	171	230	254	137	48	5	-
Female	12	15	14	25	47	116	203	302	262	152	39	-
Black, both sexes	6	4	12	10	10	15	22	15	8	3	2	-
Male	4	3	6	6	6	8	12	3	3	-	-	-
Female	2	1	6	4	4	7	10	12	5	3	2	-
American Indian 1/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	3	2	1	2	2	1	2	-	-	-
Male	-	-	3	-	1	1	1	1	2	-	-	-
Female	-	-	-	2	-	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Delaware												
All races, both sexes	15	14	25	28	51	85	131	108	96	45	6	-
Male	12	9	19	20	38	52	79	49	36	6	2	-
Female	3	5	6	8	13	33	52	59	60	39	4	-
White, both sexes	12	9	23	23	38	71	122	101	91	36	4	-
Male	10	7	18	17	29	45	77	45	35	5	1	-
Female	2	2	5	6	9	26	45	56	56	31	3	-
Black, both sexes	2	5	2	5	11	12	8	6	5	9	2	-
Male	1	2	1	3	7	6	2	3	1	1	1	-
Female	1	3	1	2	4	6	6	3	4	8	1	-
American Indian 1/, both sexes	1	-	-	-	1	1	1	-	-	-	-	-
Male	1	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	11	12	11	20	24	41	42	46	32	20	6	-
Male	7	6	7	12	15	26	21	21	10	9	1	-
Female	4	6	4	8	9	15	21	25	22	11	5	-
White, both sexes	1	2	-	1	3	8	10	7	11	9	3	-
Male	1	2	-	1	2	7	8	5	3	5	1	-
Female	-	-	-	-	1	1	2	2	8	4	2	-
Black, both sexes	10	10	11	19	21	32	32	39	21	10	3	-
Male	6	4	7	11	13	19	13	16	7	3	-	-
Female	4	6	4	8	8	13	19	23	14	7	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Florida												
All races, both sexes	460	569	806	973	1,330	2,270	3,130	3,563	2,583	1,084	249	1
Male	356	425	564	676	860	1,400	1,726	1,699	1,046	299	50	1
Female	104	144	242	297	470	870	1,404	1,864	1,537	785	199	-
White, both sexes	354	473	676	817	1,147	2,047	2,910	3,364	2,470	1,038	231	-
Male	290	360	483	580	760	1,288	1,627	1,623	1,009	295	47	-
Female	64	113	193	237	387	759	1,283	1,741	1,461	743	184	-
Black, both sexes	101	92	124	143	172	203	210	186	105	43	18	1
Male	62	61	78	86	94	101	95	70	34	4	3	1
Female	39	31	46	57	78	102	115	116	71	39	15	-
American Indian 1/, both sexes	1	2	1	2	1	1	-	4	1	-	-	-
Male	1	2	1	2	-	1	-	3	1	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	2	5	11	10	19	10	9	7	3	-	-
Male	3	2	2	8	6	10	4	3	2	-	-	-
Female	1	-	3	3	4	9	6	6	5	3	-	-
Georgia												
All races, both sexes	160	234	259	312	387	489	607	639	438	163	23	-
Male	129	183	188	212	265	288	338	276	145	39	2	-
Female	31	51	71	100	122	201	269	363	293	124	21	-
White, both sexes	106	155	176	235	288	392	501	543	358	133	15	-
Male	87	124	138	169	209	236	294	244	122	35	2	-
Female	19	31	38	66	79	156	207	299	236	98	13	-
Black, both sexes	52	77	77	76	95	94	102	94	78	30	7	-
Male	40	57	48	42	52	51	43	31	22	4	-	-
Female	12	20	29	34	43	43	59	63	56	26	7	-
American Indian 1/, both sexes	-	-	2	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	4	1	4	3	4	2	2	-	1	-
Male	2	2	1	1	4	1	1	1	1	-	-	-
Female	-	-	3	-	-	2	3	1	1	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Hawaii												
All races, both sexes	23	34	32	42	54	69	105	113	78	21	6	-
Male	15	31	23	32	36	49	69	63	34	9	5	-
Female	8	3	9	10	18	20	36	50	44	12	1	-
White, both sexes	8	8	12	10	9	20	27	37	20	3	4	-
Male	5	7	10	7	8	14	18	21	10	1	3	-
Female	3	1	2	3	1	6	9	16	10	2	1	-
Black, both sexes	1	-	-	-	-	1	-	1	-	-	-	-
Male	1	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	26	20	32	45	48	78	75	58	17	2	-
Male	9	24	13	25	28	34	51	41	24	7	2	-
Female	5	2	7	7	17	14	27	34	34	10	-	-
Idaho												
All races, both sexes	11	23	23	34	57	90	111	147	118	34	6	-
Male	10	18	19	19	38	55	67	83	40	11	-	-
Female	1	5	4	15	19	35	44	64	78	23	6	-
White, both sexes	10	23	23	34	56	90	110	146	118	34	6	-
Male	9	18	19	19	38	55	66	82	40	11	-	-
Female	1	5	4	15	18	35	44	64	78	23	6	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	1	1	-	-	-	-
Male	1	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Illinois												
All races, both sexes	207	302	383	470	625	1,025	1,513	1,657	1,311	557	116	-
Male	162	229	269	338	391	573	788	703	431	137	21	-
Female	45	73	114	132	234	452	725	954	880	420	95	-
White, both sexes	150	229	303	372	502	876	1,363	1,521	1,228	525	110	-
Male	121	179	216	278	324	507	718	655	401	134	20	-
Female	29	50	87	94	178	369	645	866	827	391	90	-
Black, both sexes	55	69	74	90	115	135	132	122	68	31	5	-
Male	40	46	52	52	63	58	62	43	24	3	1	-
Female	15	23	22	38	52	77	70	79	44	28	4	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	3	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	1	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	4	6	8	7	14	17	11	15	1	1	-
Male	1	4	1	8	4	8	8	4	6	-	-	-
Female	1	-	5	-	3	6	9	7	9	1	1	-
Indiana												
All races, both sexes	130	211	230	303	415	581	802	809	589	259	66	-
Male	101	154	153	191	272	329	418	351	202	55	7	-
Female	29	57	77	112	143	252	384	458	387	204	59	-
White, both sexes	106	186	201	281	383	535	746	763	563	245	65	-
Male	82	133	135	177	257	306	387	329	197	51	7	-
Female	24	53	66	104	126	229	359	434	366	194	58	-
Black, both sexes	24	25	28	22	31	45	54	45	26	14	1	-
Male	19	21	17	14	14	22	31	22	5	4	-	-
Female	5	4	11	8	17	23	23	23	21	10	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	1	2	-	-	-	-	-
Male	-	-	1	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Iowa												
All races, both sexes	63	71	82	117	211	297	437	524	485	207	55	-
Male	51	54	60	87	137	165	250	261	187	53	9	-
Female	12	17	22	30	74	132	187	263	298	154	46	-
White, both sexes	61	66	80	114	206	297	433	520	482	206	55	-
Male	50	53	59	85	133	165	248	259	185	53	9	-
Female	11	13	21	29	73	132	185	261	297	153	46	-
Black, both sexes	2	5	2	2	5	-	2	3	2	1	-	-
Male	1	1	1	1	4	-	1	2	1	-	-	-
Female	1	4	1	1	1	-	1	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Kansas												
All races, both sexes	71	71	86	122	146	228	339	361	274	134	36	-
Male	50	54	66	90	96	138	187	171	103	35	3	-
Female	21	17	20	32	50	90	152	190	171	99	33	-
White, both sexes	65	60	82	108	138	206	326	349	263	131	36	-
Male	46	47	64	78	94	127	183	164	98	35	3	-
Female	19	13	18	30	44	79	143	185	165	96	33	-
Black, both sexes	6	9	2	9	6	21	11	10	9	3	-	-
Male	4	5	1	8	2	10	3	5	4	-	-	-
Female	2	4	1	1	4	11	8	5	5	3	-	-
American Indian 1/, both sexes	-	2	2	4	-	1	2	2	2	-	-	-
Male	-	2	1	3	-	1	1	2	1	-	-	-
Female	-	-	1	1	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	2	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Kentucky												
All races, both sexes	87	117	128	204	227	344	456	408	334	126	24	-
Male	56	88	93	135	133	189	232	173	104	32	6	-
Female	31	29	35	69	94	155	224	235	230	94	18	-
White, both sexes	80	107	117	191	215	330	439	392	319	124	23	-
Male	53	81	85	126	126	184	228	165	101	32	6	-
Female	27	26	32	65	89	146	211	227	218	92	17	-
Black, both sexes	7	10	10	13	12	14	15	16	15	2	1	-
Male	3	7	7	9	7	5	4	8	3	-	-	-
Female	4	3	3	4	5	9	11	8	12	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	2	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Louisiana												
All races, both sexes	81	128	134	151	202	261	394	346	212	78	13	-
Male	61	102	91	97	119	161	214	153	76	23	5	-
Female	20	26	43	54	83	100	180	193	136	55	8	-
White, both sexes	51	88	86	109	148	214	320	285	173	61	9	-
Male	41	70	60	72	92	137	181	131	65	18	3	-
Female	10	18	26	37	56	77	139	154	108	43	6	-
Black, both sexes	30	39	48	42	52	46	72	60	39	17	4	-
Male	20	31	31	25	26	23	33	22	11	5	2	-
Female	10	8	17	17	26	23	39	38	28	12	2	-
American Indian 1/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	2	-	1	1	-	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Maine												
All races, both sexes	25	34	47	50	82	126	145	176	138	62	14	-
Male	21	27	36	43	60	85	80	84	59	13	4	-
Female	4	7	11	7	22	41	65	92	79	49	10	-
White, both sexes	25	33	47	50	82	124	144	176	138	62	14	-
Male	21	26	36	43	60	83	80	84	59	13	4	-
Female	4	7	11	7	22	41	64	92	79	49	10	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Maryland												
All races, both sexes	68	110	153	186	281	371	531	594	404	190	39	-
Male	51	72	112	127	179	224	261	255	146	45	8	-
Female	17	38	41	59	102	147	270	339	258	145	31	-
White, both sexes	46	56	102	127	208	287	434	495	355	167	32	-
Male	36	40	76	92	142	182	222	212	134	43	8	-
Female	10	16	26	35	66	105	212	283	221	124	24	-
Black, both sexes	21	53	46	56	63	75	91	94	45	22	6	-
Male	14	31	32	34	30	38	36	40	10	2	-	-
Female	7	22	14	22	33	37	55	54	35	20	6	-
American Indian 1/, both sexes	-	-	-	-	1	1	1	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	5	3	9	8	5	5	4	1	1	-
Male	1	1	4	1	6	4	2	3	2	-	-	-
Female	-	-	1	2	3	4	3	2	2	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Massachusetts												
All races, both sexes	77	115	177	194	308	486	791	935	721	301	72	-
Male	59	87	127	132	206	282	417	433	239	73	11	-
Female	18	28	50	62	102	204	374	502	482	228	61	-
White, both sexes	75	108	167	179	293	471	774	913	711	296	72	-
Male	58	83	124	122	197	274	407	428	236	73	11	-
Female	17	25	43	57	96	197	367	485	475	223	61	-
Black, both sexes	2	6	8	10	11	13	10	19	9	4	-	-
Male	1	4	2	6	5	8	7	4	3	-	-	-
Female	1	2	6	4	6	5	3	15	6	4	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	1	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	5	4	2	7	3	1	-	-	-
Male	-	-	-	4	4	-	3	1	-	-	-	-
Female	-	1	1	1	-	2	4	2	1	-	-	-
Michigan												
All races, both sexes	192	250	347	454	613	1,008	1,445	1,727	1,226	530	112	-
Male	151	185	246	305	379	582	758	786	428	122	21	-
Female	41	65	101	149	234	426	687	941	798	408	91	-
White, both sexes	154	204	305	390	531	907	1,331	1,604	1,162	507	105	-
Male	126	154	215	263	337	531	717	733	410	118	19	-
Female	28	50	90	127	194	376	614	871	752	389	86	-
Black, both sexes	37	42	40	60	72	92	105	116	59	20	6	-
Male	24	28	29	40	35	49	39	50	16	4	1	-
Female	13	14	11	20	37	43	66	66	43	16	5	-
American Indian 1/, both sexes	1	3	2	1	3	6	4	6	4	1	1	-
Male	1	2	2	-	2	1	2	3	2	-	1	-
Female	-	1	-	1	1	5	2	3	2	1	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	3	7	3	5	1	1	2	-	-
Male	-	1	-	2	5	1	-	-	-	-	-	-
Female	-	-	-	1	2	2	5	1	1	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Minnesota												
All races, both sexes	102	128	133	139	192	291	402	544	429	177	29	-
Male	81	111	96	102	133	186	240	285	173	40	3	-
Female	21	17	37	37	59	105	162	259	256	137	26	-
White, both sexes	98	115	126	129	181	281	393	542	428	177	29	-
Male	79	100	92	96	125	181	236	285	173	40	3	-
Female	19	15	34	33	56	100	157	257	255	137	26	-
Black, both sexes	3	6	3	3	4	2	6	1	-	-	-	-
Male	1	5	3	1	2	-	4	-	-	-	-	-
Female	2	1	-	2	2	2	2	1	-	-	-	-
American Indian 1/, both sexes	1	4	4	6	5	4	2	-	-	-	-	-
Male	1	3	1	4	4	3	-	-	-	-	-	-
Female	-	1	3	2	1	1	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	3	-	1	2	4	1	1	1	-	-	-
Male	-	3	-	1	2	2	-	-	-	-	-	-
Female	-	-	-	-	-	2	1	1	1	-	-	-
Mississippi												
All races, both sexes	97	124	118	131	193	216	264	223	139	45	16	-
Male	73	91	83	91	102	130	126	84	47	4	1	-
Female	24	33	35	40	91	86	138	139	92	41	15	-
White, both sexes	52	69	76	83	141	162	209	187	113	34	11	-
Male	39	59	58	59	80	108	99	71	37	3	-	-
Female	13	10	18	24	61	54	110	116	76	31	11	-
Black, both sexes	43	55	42	46	52	54	54	35	26	11	4	-
Male	32	32	25	30	22	22	27	12	10	1	-	-
Female	11	23	17	16	30	32	27	23	16	10	4	-
American Indian 1/, both sexes	2	-	-	2	-	-	1	-	-	-	-	-
Male	2	-	-	2	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	1	-
Male	-	-	-	-	-	-	-	1	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Missouri												
All races, both sexes	159	185	222	276	378	566	790	950	657	336	81	-
Male	124	136	162	182	242	334	424	396	213	68	11	-
Female	35	49	60	94	136	232	366	554	444	268	70	-
White, both sexes	135	145	189	232	324	518	725	880	618	309	65	-
Male	104	110	138	159	210	309	394	378	204	62	9	-
Female	31	35	51	73	114	209	331	502	414	247	56	-
Black, both sexes	22	40	32	43	50	46	64	68	36	27	16	-
Male	18	26	23	23	30	25	29	17	9	6	2	-
Female	4	14	9	20	20	21	35	51	27	21	14	-
American Indian 1/, both sexes	2	-	1	1	1	1	-	1	1	-	-	-
Male	2	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	3	1	1	1	2	-	-	-
Male	-	-	-	-	1	-	1	1	-	-	-	-
Female	-	-	-	-	2	1	-	-	2	-	-	-
Montana												
All races, both sexes	21	23	36	26	40	66	93	103	76	27	5	-
Male	18	21	25	22	30	43	54	61	36	9	1	-
Female	3	2	11	4	10	23	39	42	40	18	4	-
White, both sexes	19	20	35	24	37	62	91	103	71	26	4	-
Male	16	19	24	20	28	41	54	61	33	8	1	-
Female	3	1	11	4	9	21	37	42	38	18	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	3	1	2	3	4	2	-	5	1	1	-
Male	2	2	1	2	2	2	-	-	3	1	-	-
Female	-	1	-	-	1	2	2	-	2	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Nebraska												
All races, both sexes	29	31	43	63	83	131	200	266	223	104	22	-
Male	26	19	34	46	51	86	127	139	72	33	1	-
Female	3	12	9	17	32	45	73	127	151	71	21	-
White, both sexes	26	26	36	58	74	124	196	260	221	104	21	-
Male	23	16	30	43	46	82	125	134	72	33	1	-
Female	3	10	6	15	28	42	71	126	149	71	20	-
Black, both sexes	3	5	4	5	6	6	2	5	2	-	1	-
Male	3	3	3	3	3	3	1	4	-	-	-	-
Female	-	2	1	2	3	3	1	1	2	-	1	-
American Indian 1/, both sexes	-	-	1	-	2	1	1	-	-	-	-	-
Male	-	-	-	-	2	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	1	-	1	1	-	-	-	-
Male	-	-	1	-	-	-	1	1	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-
Nevada												
All races, both sexes	30	37	59	88	91	137	180	142	79	28	3	-
Male	25	27	45	59	60	91	115	77	30	11	1	-
Female	5	10	14	29	31	46	65	65	49	17	2	-
White, both sexes	25	32	49	79	79	129	169	134	73	26	3	-
Male	21	24	38	56	52	87	110	76	29	9	1	-
Female	4	8	11	23	27	42	59	58	44	17	2	-
Black, both sexes	1	5	5	4	9	4	8	4	4	2	-	-
Male	-	3	3	2	5	2	2	-	1	2	-	-
Female	1	2	2	2	4	2	6	4	3	-	-	-
American Indian 1/, both sexes	1	-	1	-	-	1	2	2	-	-	-	-
Male	1	-	1	-	-	1	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	4	5	3	3	1	2	2	-	-	-
Male	3	-	3	1	3	1	1	1	-	-	-	-
Female	-	-	1	4	-	2	-	1	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
New Hampshire												
All races, both sexes	10	20	46	38	76	96	164	192	150	69	12	-
Male	8	15	25	30	49	61	86	77	52	16	1	-
Female	2	5	21	8	27	35	78	115	98	53	11	-
White, both sexes	10	20	46	37	75	95	164	192	150	69	12	-
Male	8	15	25	29	49	61	86	77	52	16	1	-
Female	2	5	21	8	26	34	78	115	98	53	11	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
New Jersey												
All races, both sexes	113	178	220	367	466	849	1,335	1,555	1,174	574	127	-
Male	85	130	159	243	283	506	714	682	395	132	27	-
Female	28	48	61	124	183	343	621	873	779	442	100	-
White, both sexes	86	140	173	305	385	748	1,238	1,456	1,119	537	117	-
Male	64	106	129	206	235	453	663	641	382	131	26	-
Female	22	34	44	99	150	295	575	815	737	406	91	-
Black, both sexes	23	31	37	51	67	80	82	82	40	31	9	-
Male	17	17	24	29	39	44	41	31	11	-	1	-
Female	6	14	13	22	28	36	41	51	29	31	8	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	1	2	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	2	1	-	-
Asian or Pacific Islander 2/, both sexes	4	7	10	11	14	20	15	16	13	5	1	-
Male	4	7	6	8	9	9	10	10	2	1	-	-
Female	-	-	4	3	5	11	5	6	11	4	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
New Mexico												
All races, both sexes	15	26	54	55	89	119	187	203	162	66	19	-
Male	10	20	40	33	53	70	104	97	70	19	4	-
Female	5	6	14	22	36	49	83	106	92	47	15	-
White, both sexes	14	21	47	53	85	114	176	196	155	64	18	-
Male	9	17	36	31	51	67	99	94	67	17	4	-
Female	5	4	11	22	34	47	77	102	88	47	14	-
Black, both sexes	-	1	5	1	2	2	4	2	3	-	1	-
Male	-	1	3	1	1	1	1	-	1	-	-	-
Female	-	-	2	-	1	1	3	2	2	-	1	-
American Indian 1/, both sexes	1	4	2	1	2	3	7	3	4	2	-	-
Male	1	2	1	1	1	2	4	2	2	2	-	-
Female	-	2	1	-	1	1	3	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
New York												
All races, both sexes	412	656	910	1,278	1,754	3,010	4,618	5,216	4,418	2,191	516	-
Male	295	455	603	833	1,005	1,669	2,219	2,053	1,436	459	92	-
Female	117	201	307	445	749	1,341	2,399	3,163	2,982	1,732	424	-
White, both sexes	292	485	685	948	1,368	2,488	3,948	4,579	4,012	1,981	453	-
Male	219	359	483	638	813	1,426	1,959	1,835	1,323	426	80	-
Female	73	126	202	310	555	1,062	1,989	2,744	2,689	1,555	373	-
Black, both sexes	111	144	193	269	319	427	544	523	314	176	50	-
Male	68	75	98	155	147	191	204	165	77	23	8	-
Female	43	69	95	114	172	236	340	358	237	153	42	-
American Indian 1/, both sexes	1	1	2	4	3	-	5	5	-	1	-	-
Male	1	-	2	1	3	-	2	3	-	-	-	-
Female	-	1	-	3	-	-	3	2	-	1	-	-
Asian or Pacific Islander 2/, both sexes	8	26	30	57	64	95	121	109	92	33	13	-
Male	7	21	20	39	42	52	54	50	36	10	4	-
Female	1	5	10	18	22	43	67	59	56	23	9	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
North Carolina												
All races, both sexes	276	359	490	500	629	799	984	997	657	248	59	1
Male	197	269	359	349	413	478	520	415	207	62	6	-
Female	79	90	131	151	216	321	464	582	450	186	53	1
White, both sexes	196	250	368	381	507	652	819	862	563	205	45	-
Male	140	192	279	268	339	398	444	366	186	53	5	-
Female	56	58	89	113	168	254	375	496	377	152	40	-
Black, both sexes	77	101	115	107	115	135	154	126	91	40	14	1
Male	55	70	73	72	69	70	70	44	21	8	1	-
Female	22	31	42	35	46	65	84	82	70	32	13	1
American Indian 1/, both sexes	2	5	7	10	6	8	10	6	2	2	-	-
Male	1	4	7	8	4	7	5	3	-	1	-	-
Female	1	1	-	2	2	1	5	3	2	1	-	-
Asian or Pacific Islander 2/, both sexes	1	3	-	2	1	4	1	3	1	1	-	-
Male	1	3	-	1	1	3	1	2	-	-	-	-
Female	-	-	-	1	-	1	-	1	1	1	-	-
North Dakota												
All races, both sexes	21	6	15	27	34	60	71	108	103	51	13	-
Male	16	5	12	21	26	34	39	52	43	13	2	-
Female	5	1	3	6	8	26	32	56	60	38	11	-
White, both sexes	19	5	15	26	33	58	70	107	102	51	13	-
Male	14	5	12	21	26	33	38	51	43	13	2	-
Female	5	-	3	5	7	25	32	56	59	38	11	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	1	1	2	1	1	1	-	-	-
Male	1	-	-	-	-	1	1	1	-	-	-	-
Female	-	1	-	1	1	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Ohio												
All races, both sexes	271	382	550	608	839	1,312	1,932	2,162	1,507	619	129	-
Male	202	282	397	428	532	762	1,013	930	509	148	17	-
Female	69	100	153	180	307	550	919	1,232	998	471	112	-
White, both sexes	225	314	472	530	722	1,198	1,795	2,036	1,431	574	117	-
Male	172	236	345	378	462	701	950	879	490	136	16	-
Female	53	78	127	152	260	497	845	1,157	941	438	101	-
Black, both sexes	44	65	73	76	115	108	132	123	73	45	10	-
Male	28	44	47	49	69	56	60	49	17	12	1	-
Female	16	21	26	27	46	52	72	74	56	33	9	-
American Indian 1/, both sexes	-	-	1	1	-	1	1	-	-	-	1	-
Male	-	-	1	1	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	2	3	4	1	2	5	4	3	3	-	1	-
Male	2	2	4	-	1	4	2	2	2	-	-	-
Female	-	1	-	1	1	1	2	1	1	-	1	-
Oklahoma												
All races, both sexes	71	111	162	200	248	389	517	522	403	158	30	-
Male	53	75	103	127	152	215	270	220	159	32	5	-
Female	18	36	59	73	96	174	247	302	244	126	25	-
White, both sexes	62	91	133	170	213	338	474	494	372	146	24	-
Male	47	59	87	106	134	189	251	210	149	30	3	-
Female	15	32	46	64	79	149	223	284	223	116	21	-
Black, both sexes	3	12	10	13	18	27	23	16	17	6	4	-
Male	2	12	5	8	9	13	9	4	9	1	2	-
Female	1	-	5	5	9	14	14	12	8	5	2	-
American Indian 1/, both sexes	5	7	17	17	15	23	20	12	14	5	2	-
Male	3	4	10	13	8	13	10	6	1	1	-	-
Female	2	3	7	4	7	10	10	6	13	4	2	-
Asian or Pacific Islander 2/, both sexes	1	1	2	-	2	1	-	-	-	1	-	-
Male	1	-	1	-	1	-	-	-	-	-	-	-
Female	-	1	1	-	1	1	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Oregon												
All races, both sexes	79	139	157	142	197	274	377	426	348	130	22	1
Male	66	102	116	110	138	176	237	219	150	36	3	1
Female	13	37	41	32	59	98	140	207	198	94	19	-
White, both sexes	76	130	153	134	191	267	365	420	345	125	22	1
Male	64	95	113	105	133	173	230	217	149	35	3	1
Female	12	35	40	29	58	94	135	203	196	90	19	-
Black, both sexes	1	6	-	3	2	2	3	1	1	2	-	-
Male	-	5	-	2	2	2	2	-	-	-	-	-
Female	1	1	-	1	-	-	1	1	1	2	-	-
American Indian 1/, both sexes	1	1	1	4	-	1	4	2	-	1	-	-
Male	1	1	1	2	-	-	1	1	-	-	-	-
Female	-	-	-	2	-	1	3	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	2	3	1	4	4	5	3	2	2	-	-
Male	1	1	2	1	3	1	4	1	1	1	-	-
Female	-	1	1	-	1	3	1	2	1	1	-	-
Pennsylvania												
All races, both sexes	164	277	397	514	742	1,294	1,977	2,314	1,697	642	138	2
Male	129	216	267	343	481	762	1,049	1,021	597	184	27	1
Female	35	61	130	171	261	532	928	1,293	1,100	458	111	1
White, both sexes	135	239	350	463	673	1,196	1,874	2,220	1,618	613	128	1
Male	108	190	246	308	446	715	1,005	994	580	175	24	-
Female	27	49	104	155	227	481	869	1,226	1,038	438	104	1
Black, both sexes	29	35	46	45	65	92	95	86	77	28	10	1
Male	21	25	21	29	32	44	41	22	15	8	3	1
Female	8	10	25	16	33	48	54	64	62	20	7	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	1	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	3	1	5	4	5	8	7	2	1	-	-
Male	-	1	-	5	3	2	3	5	2	1	-	-
Female	-	2	1	-	1	3	5	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Rhode Island												
All races, both sexes	12	25	28	31	53	97	161	231	217	93	22	-
Male	11	19	22	20	32	53	58	96	64	20	-	-
Female	1	6	6	11	21	44	103	135	153	73	22	-
White, both sexes	11	24	25	30	47	91	159	226	215	93	22	-
Male	10	18	20	20	29	51	58	94	64	20	-	-
Female	1	6	5	10	18	40	101	132	151	73	22	-
Black, both sexes	-	1	3	1	4	5	2	5	2	-	-	-
Male	-	1	2	-	2	2	-	2	-	-	-	-
Female	-	-	1	1	2	3	2	3	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	1	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
South Carolina												
All races, both sexes	89	109	165	180	229	328	418	372	269	91	13	1
Male	63	84	130	125	145	211	220	163	84	19	4	1
Female	26	25	35	55	84	117	198	209	185	72	9	-
White, both sexes	59	72	120	139	161	262	347	307	226	74	9	1
Male	44	57	95	97	111	174	193	140	73	16	3	1
Female	15	15	25	42	50	88	154	167	153	58	6	-
Black, both sexes	29	36	45	38	65	66	70	65	42	17	4	-
Male	18	27	35	26	31	37	26	23	11	3	1	-
Female	11	9	10	12	34	29	44	42	31	14	3	-
American Indian 1/, both sexes	-	1	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	2	2	-	1	-	1	-	-	-
Male	1	-	-	1	2	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
South Dakota												
All races, both sexes	16	20	22	33	36	56	99	127	110	54	9	-
Male	13	15	15	24	26	38	58	69	48	16	1	-
Female	3	5	7	9	10	18	41	58	62	38	8	-
White, both sexes	13	17	21	31	32	55	94	124	108	53	8	-
Male	12	13	15	24	24	37	56	67	48	16	1	-
Female	1	4	6	7	8	18	38	57	60	37	7	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	3	2	1	2	4	1	5	3	2	1	-	-
Male	1	2	-	-	2	1	2	2	-	-	-	-
Female	2	-	1	2	2	-	3	1	2	1	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	141	182	252	309	373	546	728	738	429	195	35	-
Male	86	123	172	206	217	326	362	313	115	39	4	-
Female	55	59	80	103	156	220	366	425	314	156	31	-
White, both sexes	116	149	212	267	319	489	659	671	393	178	31	-
Male	76	104	150	178	193	292	327	286	108	35	4	-
Female	40	45	62	89	126	197	332	385	285	143	27	-
Black, both sexes	23	33	39	40	54	56	67	66	36	17	4	-
Male	10	19	22	26	24	33	33	26	7	4	-	-
Female	13	14	17	14	30	23	34	40	29	13	4	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	1	2	-	1	1	1	-	-	-	-
Male	-	-	-	2	-	1	1	1	-	-	-	-
Female	2	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Texas												
All races, both sexes	292	459	630	695	940	1,453	1,895	1,992	1,334	524	92	-
Male	201	342	435	466	608	833	998	905	449	144	16	-
Female	91	117	195	229	332	620	897	1,087	885	380	76	-
White, both sexes	223	382	522	587	828	1,301	1,736	1,854	1,246	477	80	-
Male	154	286	362	408	544	758	929	854	422	136	12	-
Female	69	96	160	179	284	543	807	1,000	824	341	68	-
Black, both sexes	62	72	101	94	106	140	152	120	78	46	10	-
Male	42	53	68	50	62	71	66	46	23	8	2	-
Female	20	19	33	44	44	69	86	74	55	38	8	-
American Indian 1/, both sexes	1	1	1	5	2	1	-	2	2	-	-	-
Male	1	1	-	4	1	1	-	-	-	-	-	-
Female	-	-	1	1	1	-	-	2	2	-	-	-
Asian or Pacific Islander 2/, both sexes	6	4	6	9	4	11	7	16	8	1	2	-
Male	4	2	5	4	1	3	3	5	4	-	2	-
Female	2	2	1	5	3	8	4	11	4	1	-	-
Utah												
All races, both sexes	31	57	48	45	69	89	161	156	90	51	3	-
Male	22	47	35	25	52	55	93	95	43	18	-	-
Female	9	10	13	20	17	34	68	61	47	33	3	-
White, both sexes	29	53	45	43	69	88	159	155	90	51	3	-
Male	20	43	33	25	52	54	93	94	43	18	-	-
Female	9	10	12	18	17	34	66	61	47	33	3	-
Black, both sexes	1	-	1	-	-	1	-	-	-	-	-	-
Male	1	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	2	-	-	1	1	-	-	-	-
Male	1	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	2	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	3	2	-	-	-	1	-	-	-	-	-
Male	-	3	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Vermont												
All races, both sexes	8	11	11	13	20	53	59	75	49	28	8	-
Male	4	6	8	9	16	39	27	39	16	9	-	-
Female	4	5	3	4	4	14	32	36	33	19	8	-
White, both sexes	8	11	11	13	20	53	57	75	49	28	8	-
Male	4	6	8	9	16	39	26	39	16	9	-	-
Female	4	5	3	4	4	14	31	36	33	19	8	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	117	172	202	278	344	489	672	631	410	168	34	-
Male	89	128	138	197	231	300	359	298	154	46	7	-
Female	28	44	64	81	113	189	313	333	256	122	27	-
White, both sexes	81	122	137	220	290	383	580	552	346	147	19	-
Male	61	91	98	164	199	233	323	267	129	42	5	-
Female	20	31	39	56	91	150	257	285	217	105	14	-
Black, both sexes	32	45	63	55	48	97	82	71	57	21	15	-
Male	25	33	38	32	29	62	32	27	22	4	2	-
Female	7	12	25	23	19	35	50	44	35	17	13	-
American Indian 1/, both sexes	1	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	5	2	3	6	8	9	8	7	-	-	-
Male	3	4	2	1	3	5	4	4	3	-	-	-
Female	-	1	-	2	3	3	5	4	4	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Washington												
All races, both sexes	64	110	177	184	320	471	659	778	608	234	47	-
Male	53	81	130	133	210	288	378	396	257	60	12	-
Female	11	29	47	51	110	183	281	382	351	174	35	-
White, both sexes	59	101	164	163	298	429	620	751	592	228	47	-
Male	49	74	120	117	195	265	358	386	250	59	12	-
Female	10	27	44	46	103	164	262	365	342	169	35	-
Black, both sexes	3	3	4	8	7	12	12	6	5	-	-	-
Male	3	2	4	5	6	7	7	3	3	-	-	-
Female	-	1	-	3	1	5	5	3	2	-	-	-
American Indian 1/, both sexes	1	2	2	6	3	9	5	2	-	2	-	-
Male	-	1	2	4	1	5	3	1	-	-	-	-
Female	1	1	-	2	2	4	2	1	-	2	-	-
Asian or Pacific Islander 2/, both sexes	1	4	7	7	12	21	22	19	11	4	-	-
Male	1	4	4	7	8	11	10	6	4	1	-	-
Female	-	-	3	-	4	10	12	13	7	3	-	-
West Virginia												
All races, both sexes	77	87	97	171	165	247	339	329	227	87	14	-
Male	60	68	70	111	103	132	204	140	68	14	3	-
Female	17	19	27	60	62	115	135	189	159	73	11	-
White, both sexes	75	86	94	168	160	239	335	316	217	85	14	-
Male	58	67	67	108	100	128	203	136	66	13	3	-
Female	17	19	27	60	60	111	132	180	151	72	11	-
Black, both sexes	2	1	3	3	5	8	3	11	10	2	-	-
Male	2	1	3	3	3	4	1	4	2	1	-	-
Female	-	-	-	-	2	4	2	7	8	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Wisconsin												
All races, both sexes	61	105	108	166	222	379	573	632	496	221	50	-
Male	46	87	82	123	147	231	353	287	194	59	5	-
Female	15	18	26	43	75	148	220	345	302	162	45	-
White, both sexes	54	96	100	155	209	366	560	619	485	217	48	-
Male	42	83	76	118	140	226	349	282	191	58	5	-
Female	12	13	24	37	69	140	211	337	294	159	43	-
Black, both sexes	6	6	5	6	11	8	10	10	8	4	1	-
Male	3	1	4	3	6	2	2	3	1	1	-	-
Female	3	5	1	3	5	6	8	7	7	3	1	-
American Indian 1/, both sexes	1	2	1	3	2	5	2	2	1	-	1	-
Male	1	2	1	2	1	3	1	2	1	-	-	-
Female	-	-	-	1	1	2	1	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	1	2	2	-	-	1	1	2	-	-	-
Male	-	1	1	-	-	-	1	-	1	-	-	-
Female	-	-	1	2	-	-	-	1	1	-	-	-
Wyoming												
All races, both sexes	9	16	8	24	21	36	55	45	30	3	3	-
Male	6	12	5	15	17	21	33	23	11	1	2	-
Female	3	4	3	9	4	15	22	22	19	2	1	-
White, both sexes	9	14	8	23	21	36	53	45	30	3	3	-
Male	6	11	5	14	17	21	32	23	11	1	2	-
Female	3	3	3	9	4	15	21	22	19	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	2	-	1	-	-	1	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
United States												
All races, both sexes	172,748	397	163	98	121	296	576	800	977	1,615	2,486	3,870
Male	77,178	225	77	57	64	196	353	506	594	1,009	1,568	2,414
Female	95,570	172	86	41	57	100	223	294	383	606	918	1,456
White, both sexes	149,300	233	106	68	80	194	388	505	631	1,060	1,697	2,719
Male	65,872	132	44	37	44	123	231	327	398	669	1,065	1,749
Female	83,428	101	62	31	36	71	157	178	233	391	632	970
Black, both sexes	20,352	136	41	22	38	94	169	260	299	501	708	1,062
Male	9,736	78	27	15	18	67	110	154	166	302	446	604
Female	10,616	58	14	7	20	27	59	106	133	199	262	458
American Indian 1/, both sexes	711	8	5	1	-	4	7	12	23	14	30	46
Male	382	5	4	1	-	3	1	9	16	9	20	34
Female	329	3	1	-	-	1	6	3	7	5	10	12
Asian or Pacific Islander 2/, both sexes	2,385	20	11	7	3	4	12	23	24	40	51	43
Male	1,188	10	2	4	2	3	11	16	14	29	37	27
Female	1,197	10	9	3	1	1	1	7	10	11	14	16
Alabama												
All races, both sexes	5,926	10	4	4	2	4	21	24	31	55	123	185
Male	2,747	4	4	3	-	2	15	11	18	23	75	124
Female	3,179	6	-	1	2	2	6	13	13	32	48	61
White, both sexes	4,437	5	-	2	1	3	13	11	16	27	82	107
Male	2,036	2	-	2	-	2	11	6	11	14	49	66
Female	2,401	3	-	-	1	1	2	5	5	13	33	41
Black, both sexes	1,472	5	4	2	1	1	8	13	15	28	41	77
Male	703	2	4	1	-	-	4	5	7	9	26	57
Female	769	3	-	1	1	1	4	8	8	19	15	20
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Alaska												
All races, both sexes	200	-	1	1	-	2	-	1	5	-	8	7
Male	97	-	1	-	-	1	-	-	3	-	5	5
Female	103	-	-	1	-	1	-	1	2	-	3	2
White, both sexes	151	-	-	1	-	2	-	-	1	-	5	6
Male	69	-	-	-	-	1	-	-	1	-	3	4
Female	82	-	-	1	-	1	-	-	-	-	2	2
Black, both sexes	3	-	-	-	-	-	-	-	-	-	2	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	41	-	1	-	-	-	-	1	4	-	1	-
Male	24	-	1	-	-	-	-	-	2	-	1	-
Female	17	-	-	-	-	-	-	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2,279	9	2	2	1	7	11	19	16	17	46	58
Male	1,104	3	1	1	1	5	7	12	11	12	28	40
Female	1,175	6	1	1	-	2	4	7	5	5	18	18
White, both sexes	2,140	6	2	1	1	5	8	18	11	17	37	53
Male	1,038	2	1	1	1	4	7	12	7	12	23	37
Female	1,102	4	1	-	-	1	1	6	4	5	14	16
Black, both sexes	61	1	-	1	-	1	1	-	1	-	4	3
Male	29	-	-	-	-	-	-	-	1	-	2	1
Female	32	1	-	1	-	1	1	-	-	-	2	2
American Indian 1/, both sexes	66	2	-	-	-	1	2	1	4	-	4	1
Male	33	1	-	-	-	1	-	-	3	-	2	1
Female	33	1	-	-	-	-	2	1	1	-	2	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	1	1
Female	8	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Arkansas												
All races, both sexes	2,026	6	4	-	3	5	7	11	9	14	27	48
Male	846	4	2	-	3	4	5	7	6	11	18	28
Female	1,180	2	2	-	-	1	2	4	3	3	9	20
White, both sexes	1,745	3	2	-	2	3	5	4	5	11	16	39
Male	721	1	1	-	2	2	3	3	3	10	10	23
Female	1,024	2	1	-	-	1	2	1	2	1	6	16
Black, both sexes	268	3	2	-	1	2	1	7	4	3	10	9
Male	121	3	1	-	1	2	1	4	3	1	8	5
Female	147	-	1	-	-	-	-	3	1	2	2	4
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	1	-	-	-	-	-
Male	3	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	13,666	37	20	9	11	27	52	88	104	152	239	371
Male	6,317	23	8	6	7	20	37	60	69	105	170	239
Female	7,349	14	12	3	4	7	15	28	35	47	69	132
White, both sexes	11,862	27	16	7	10	18	40	56	76	116	171	274
Male	5,387	15	7	4	7	13	27	41	50	77	120	180
Female	6,475	12	9	3	3	5	13	15	26	39	51	94
Black, both sexes	1,002	5	3	-	1	7	9	20	20	22	46	75
Male	504	4	1	-	-	5	7	9	13	18	34	46
Female	498	1	2	-	1	2	2	11	7	4	12	29
American Indian 1/, both sexes	72	-	-	-	-	-	-	2	1	-	6	7
Male	45	-	-	-	-	-	-	2	1	-	5	4
Female	27	-	-	-	-	-	-	-	-	-	1	3
Asian or Pacific Islander 2/, both sexes	730	5	1	2	-	2	3	10	7	14	16	15
Male	381	4	-	2	-	2	3	8	5	10	11	9
Female	349	1	1	-	-	-	-	2	2	4	5	6

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Colorado												
All races, both sexes	1,841	6	4	3	-	4	7	6	8	20	32	42
Male	810	3	3	2	-	2	4	5	3	15	18	31
Female	1,031	3	1	1	-	2	3	1	5	5	14	11
White, both sexes	1,762	5	4	3	-	3	5	6	7	19	27	38
Male	772	3	3	2	-	1	2	5	2	15	17	28
Female	990	2	1	1	-	2	3	1	5	4	10	10
Black, both sexes	56	1	-	-	-	-	2	-	1	1	4	3
Male	27	-	-	-	-	-	2	-	1	-	1	2
Female	29	1	-	-	-	-	-	-	-	1	3	1
American Indian 1/, both sexes	8	-	-	-	-	1	-	-	-	-	-	1
Male	6	-	-	-	-	1	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1	-
Connecticut												
All races, both sexes	2,654	2	3	2	-	6	6	3	18	20	23	54
Male	1,201	2	1	1	-	5	3	-	9	15	18	33
Female	1,453	-	2	1	-	1	3	3	9	5	5	21
White, both sexes	2,458	2	1	2	-	2	4	3	9	13	18	46
Male	1,109	2	-	1	-	2	2	-	6	9	14	29
Female	1,349	-	1	1	-	-	2	3	3	4	4	17
Black, both sexes	184	-	1	-	-	3	2	-	9	6	5	8
Male	86	-	1	-	-	3	1	-	3	5	4	4
Female	98	-	-	-	-	-	1	-	6	1	1	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	1	-	-	1	-	-	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	1	-	-
Female	5	-	1	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Delaware												
All races, both sexes	555	-	2	-	-	1	4	1	2	12	12	12
Male	243	-	-	-	-	1	2	-	-	6	8	6
Female	312	-	2	-	-	-	2	1	2	6	4	6
White, both sexes	469	-	1	-	-	1	2	1	1	11	8	7
Male	201	-	-	-	-	1	-	-	-	6	5	5
Female	268	-	1	-	-	-	2	1	1	5	3	2
Black, both sexes	83	-	1	-	-	-	2	-	1	1	4	5
Male	40	-	-	-	-	-	2	-	-	-	3	1
Female	43	-	1	-	-	-	-	-	1	1	1	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	251	-	-	1	-	-	3	1	-	4	2	9
Male	103	-	-	1	-	-	1	-	-	3	2	5
Female	148	-	-	-	-	-	2	1	-	1	-	4
White, both sexes	63	-	-	-	-	-	1	-	-	-	1	2
Male	26	-	-	-	-	-	-	-	-	-	1	1
Female	37	-	-	-	-	-	1	-	-	-	-	1
Black, both sexes	188	-	-	1	-	-	2	1	-	4	1	7
Male	77	-	-	1	-	-	1	-	-	3	1	4
Female	111	-	-	-	-	-	1	1	-	1	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Florida												
All races, both sexes	9,392	32	4	6	10	17	29	25	48	89	109	182
Male	4,657	15	3	3	6	13	19	10	31	67	68	108
Female	4,735	17	1	3	4	4	10	15	17	22	41	74
White, both sexes	8,335	10	3	5	3	10	20	13	25	51	80	120
Male	4,124	5	2	3	1	8	11	5	17	43	48	76
Female	4,211	5	1	2	2	2	9	8	8	8	32	44
Black, both sexes	999	20	1	1	6	7	9	12	21	36	26	59
Male	496	9	1	-	4	5	8	5	13	23	19	29
Female	503	11	-	1	2	2	1	7	8	13	7	30
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	1	-	-	2
Male	7	-	-	-	-	-	-	-	1	-	-	2
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	50	2	-	-	1	-	-	-	1	2	3	1
Male	30	1	-	-	1	-	-	-	-	1	1	1
Female	20	1	-	-	-	-	-	-	1	1	2	-
Georgia												
All races, both sexes	6,931	26	10	6	5	14	28	33	40	87	131	174
Male	3,191	14	2	5	3	9	18	20	22	56	75	108
Female	3,740	12	8	1	2	5	10	13	18	31	56	66
White, both sexes	5,066	8	4	3	2	9	12	16	18	52	64	89
Male	2,313	6	-	2	1	5	4	11	10	34	32	59
Female	2,753	2	4	1	1	4	8	5	8	18	32	30
Black, both sexes	1,822	18	3	3	3	5	15	17	22	34	65	85
Male	864	8	2	3	2	4	13	9	12	21	42	49
Female	958	10	1	-	1	1	2	8	10	13	23	36
American Indian 1/, both sexes	7	-	1	-	-	-	-	-	-	-	2	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	1	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	36	-	2	-	-	-	1	-	-	1	-	-
Male	12	-	-	-	-	-	1	-	-	1	-	-
Female	24	-	2	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Hawaii												
All races, both sexes	953	3	1	1	-	1	2	1	9	11	15	24
Male	519	1	1	1	-	1	1	1	3	6	11	15
Female	434	2	-	-	-	-	1	-	6	5	4	9
White, both sexes	245	1	1	-	-	-	1	-	-	1	5	4
Male	139	-	1	-	-	-	-	-	-	1	4	3
Female	106	1	-	-	-	-	1	-	-	-	1	1
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	699	2	-	1	-	1	1	1	9	10	10	19
Male	375	1	-	1	-	1	1	1	3	5	7	11
Female	324	1	-	-	-	-	-	-	6	5	3	8
Idaho												
All races, both sexes	775	1	1	-	-	-	2	2	2	5	10	20
Male	353	1	-	-	-	-	1	2	1	2	4	12
Female	422	-	1	-	-	-	1	-	1	3	6	8
White, both sexes	759	1	1	-	-	-	2	2	2	5	9	20
Male	342	1	-	-	-	-	1	2	1	2	3	12
Female	417	-	1	-	-	-	1	-	1	3	6	8
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Illinois												
All races, both sexes	7,679	19	9	4	4	12	21	47	42	85	107	175
Male	3,320	11	4	2	1	10	14	33	24	48	65	117
Female	4,359	8	5	2	3	2	7	14	18	37	42	58
White, both sexes	6,555	11	5	3	4	8	18	30	27	47	59	108
Male	2,749	7	3	1	1	7	12	20	17	32	37	75
Female	3,806	4	2	2	3	1	6	10	10	15	22	33
Black, both sexes	1,062	7	3	1	-	4	2	15	15	37	46	66
Male	542	4	1	1	-	3	1	13	7	16	26	41
Female	520	3	2	-	-	1	1	2	8	21	20	25
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	1	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	56	1	1	-	-	-	1	2	-	-	1	1
Male	26	-	-	-	-	-	1	-	-	-	1	1
Female	30	1	1	-	-	-	-	2	-	-	-	-
Indiana												
All races, both sexes	4,676	11	4	-	5	8	16	22	23	40	80	112
Male	2,064	8	-	-	2	5	12	9	13	20	51	75
Female	2,612	3	4	-	3	3	4	13	10	20	29	37
White, both sexes	4,329	7	4	-	5	7	11	14	18	29	64	102
Male	1,891	5	-	-	2	5	10	5	11	15	40	68
Female	2,438	2	4	-	3	2	1	9	7	14	24	34
Black, both sexes	340	4	-	-	-	1	5	8	5	10	15	10
Male	169	3	-	-	-	-	2	4	2	4	10	7
Female	171	1	-	-	-	1	3	4	3	6	5	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	1	1	-
Male	4	-	-	-	-	-	-	-	-	1	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Iowa												
All races, both sexes	1,353	3	1	-	-	2	4	7	9	18	11	24
Male	576	1	1	-	-	2	2	5	5	11	9	16
Female	777	2	-	-	-	-	2	2	4	7	2	8
White, both sexes	1,332	2	1	-	-	2	4	4	9	17	11	22
Male	566	1	1	-	-	2	2	3	5	10	9	14
Female	766	1	-	-	-	-	2	1	4	7	2	8
Black, both sexes	19	1	-	-	-	-	-	3	-	-	-	2
Male	9	-	-	-	-	-	-	2	-	-	-	2
Female	10	1	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	2,228	5	1	-	-	3	8	4	10	12	24	47
Male	909	3	-	-	-	1	4	3	6	7	12	31
Female	1,319	2	1	-	-	2	4	1	4	5	12	16
White, both sexes	2,086	3	1	-	-	2	7	3	8	10	23	38
Male	844	1	-	-	-	1	4	3	4	6	12	26
Female	1,242	2	1	-	-	1	3	-	4	4	11	12
Black, both sexes	111	1	-	-	-	-	1	1	2	2	1	5
Male	54	1	-	-	-	-	-	-	2	1	-	2
Female	57	-	-	-	-	-	1	1	-	1	1	3
American Indian 1/, both sexes	22	1	-	-	-	1	-	-	-	-	-	4
Male	10	1	-	-	-	-	-	-	-	-	-	3
Female	12	-	-	-	-	1	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Kentucky												
All races, both sexes	3,277	4	2	1	3	2	9	9	13	29	52	71
Male	1,405	1	2	-	1	-	4	5	6	11	29	37
Female	1,872	3	-	1	2	2	5	4	7	18	23	34
White, both sexes	3,054	4	2	1	3	2	6	9	10	25	42	57
Male	1,303	1	2	-	1	-	2	5	5	10	25	30
Female	1,751	3	-	1	2	2	4	4	5	15	17	27
Black, both sexes	218	-	-	-	-	-	3	-	3	4	10	14
Male	98	-	-	-	-	-	2	-	1	1	4	7
Female	120	-	-	-	-	-	1	-	2	3	6	7
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	3,336	8	4	1	6	8	13	9	20	41	43	73
Male	1,442	4	4	1	3	6	6	3	10	25	27	39
Female	1,894	4	-	-	3	2	7	6	10	16	16	34
White, both sexes	2,428	3	-	-	3	5	5	4	7	24	21	38
Male	1,027	2	-	-	2	4	3	1	2	15	13	18
Female	1,401	1	-	-	1	1	2	3	5	9	8	20
Black, both sexes	893	5	3	1	3	3	8	4	13	17	22	35
Male	409	2	3	1	1	2	3	2	8	10	14	21
Female	484	3	-	-	2	1	5	2	5	7	8	14
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	1	-	-	-	-	1	-	-	-	-
Male	5	-	1	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Maine												
All races, both sexes	995	2	-	1	2	2	4	-	4	5	5	15
Male	463	1	-	-	1	-	2	-	2	4	3	10
Female	532	1	-	1	1	2	2	-	2	1	2	5
White, both sexes	988	2	-	1	2	2	4	-	4	5	4	13
Male	457	1	-	-	1	-	2	-	2	4	3	8
Female	531	1	-	1	1	2	2	-	2	1	1	5
Black, both sexes	3	-	-	-	-	-	-	-	-	-	1	2
Male	2	-	-	-	-	-	-	-	-	-	-	2
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	2,526	5	2	2	4	8	11	20	21	43	59	61
Male	1,149	2	1	2	-	7	9	11	12	28	43	33
Female	1,377	3	1	-	4	1	2	9	9	15	16	28
White, both sexes	1,838	2	1	1	3	1	6	11	8	20	31	31
Male	818	1	-	1	-	1	5	5	6	13	24	18
Female	1,020	1	1	-	3	-	1	6	2	7	7	13
Black, both sexes	643	2	1	1	1	7	5	8	13	23	26	30
Male	310	1	1	1	-	6	4	5	6	15	17	15
Female	333	1	-	-	1	1	1	3	7	8	9	15
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	43	1	-	-	-	-	-	1	-	-	2	-
Male	21	-	-	-	-	-	-	1	-	-	2	-
Female	22	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Massachusetts												
All races, both sexes	4,017	5	1	1	4	10	11	18	12	18	42	45
Male	1,667	2	-	-	2	6	7	11	11	10	29	31
Female	2,350	3	1	1	2	4	4	7	1	8	13	14
White, both sexes	3,812	4	1	1	3	8	10	15	11	15	34	39
Male	1,556	2	-	-	1	5	6	9	10	8	21	27
Female	2,256	2	1	1	2	3	4	6	1	7	13	12
Black, both sexes	166	-	-	-	1	2	1	2	1	1	7	6
Male	90	-	-	-	1	1	1	1	1	-	7	4
Female	76	-	-	-	-	1	-	1	-	1	-	2
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	35	1	-	-	-	-	-	1	-	2	1	-
Male	19	-	-	-	-	-	-	1	-	2	1	-
Female	16	1	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	5,925	13	2	1	3	11	18	32	41	65	78	147
Male	2,575	7	2	-	2	7	9	24	22	42	51	91
Female	3,350	6	-	1	1	4	9	8	19	23	27	56
White, both sexes	5,098	10	1	1	2	7	9	18	20	33	45	100
Male	2,169	5	1	-	2	3	4	15	10	24	32	65
Female	2,929	5	-	1	-	4	5	3	10	9	13	35
Black, both sexes	784	3	-	-	1	4	9	14	21	31	30	47
Male	388	2	-	-	-	4	5	9	12	18	16	26
Female	396	1	-	-	1	-	4	5	9	13	14	21
American Indian 1/, both sexes	16	-	-	-	-	-	-	-	-	1	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	12	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	27	-	1	-	-	-	-	-	-	-	2	-
Male	14	-	1	-	-	-	-	-	-	-	2	-
Female	13	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Minnesota												
All races, both sexes	2,891	7	5	2	-	3	6	12	13	13	31	41
Male	1,263	7	2	2	-	-	2	8	7	4	22	21
Female	1,628	-	3	-	-	3	4	4	6	9	9	20
White, both sexes	2,792	5	5	1	-	3	6	5	9	9	25	34
Male	1,211	5	2	1	-	-	2	4	5	2	17	17
Female	1,581	-	3	-	-	3	4	1	4	7	8	17
Black, both sexes	62	-	-	-	-	-	-	6	2	3	5	4
Male	36	-	-	-	-	-	-	3	1	2	4	3
Female	26	-	-	-	-	-	-	3	1	1	1	1
American Indian 1/, both sexes	24	1	-	-	-	-	-	-	2	1	1	3
Male	10	1	-	-	-	-	-	-	1	-	1	1
Female	14	-	-	-	-	-	-	-	1	1	-	2
Asian or Pacific Islander 2/, both sexes	13	1	-	1	-	-	-	1	-	-	-	-
Male	6	1	-	1	-	-	-	1	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	3,104	8	1	1	3	6	13	13	11	29	47	83
Male	1,343	5	1	1	2	1	8	8	5	17	26	52
Female	1,761	3	-	-	1	5	5	5	6	12	21	31
White, both sexes	2,059	3	-	1	1	2	6	2	5	10	18	37
Male	898	2	-	1	1	-	4	2	3	5	10	24
Female	1,161	1	-	-	-	2	2	-	2	5	8	13
Black, both sexes	1,034	5	1	-	2	4	7	11	6	19	28	46
Male	439	3	1	-	1	1	4	6	2	12	15	28
Female	595	2	-	-	1	3	3	5	4	7	13	18
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Missouri												
All races, both sexes	4,050	5	3	5	2	7	12	15	27	38	53	93
Male	1,776	2	2	3	1	7	8	10	16	25	29	69
Female	2,274	3	1	2	1	-	4	5	11	13	24	24
White, both sexes	3,711	2	3	4	1	5	11	12	19	26	44	72
Male	1,611	1	2	2	1	5	7	9	12	16	23	56
Female	2,100	1	1	2	-	-	4	3	7	10	21	16
Black, both sexes	326	2	-	1	1	2	1	3	7	12	9	21
Male	160	-	-	1	-	2	1	1	4	9	6	13
Female	166	2	-	-	1	-	-	2	3	3	3	8
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	8	1	-	-	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	739	-	1	-	1	2	2	4	1	1	6	12
Male	337	-	1	-	1	2	-	1	-	1	4	5
Female	402	-	-	-	-	-	2	3	1	-	2	7
White, both sexes	717	-	-	-	1	2	1	4	-	1	6	10
Male	324	-	-	-	1	2	-	1	-	1	4	3
Female	393	-	-	-	-	-	1	3	-	-	2	7
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	21	-	1	-	-	-	1	-	1	-	-	2
Male	13	-	1	-	-	-	-	-	-	-	-	2
Female	8	-	-	-	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Nebraska												
All races, both sexes	1,382	1	1	-	1	-	1	1	3	12	18	20
Male	578	-	-	-	-	-	-	1	1	10	11	12
Female	804	1	1	-	1	-	1	-	2	2	7	8
White, both sexes	1,331	1	1	-	1	-	1	-	3	11	15	18
Male	559	-	-	-	-	-	-	-	1	9	9	11
Female	772	1	1	-	1	-	1	-	2	2	6	7
Black, both sexes	42	-	-	-	-	-	-	1	-	1	3	1
Male	15	-	-	-	-	-	-	1	-	1	2	1
Female	27	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1,823	3	2	2	3	3	10	12	13	27	40	55
Male	1,005	2	-	2	2	1	7	7	9	21	30	37
Female	818	1	2	-	1	2	3	5	4	6	10	18
White, both sexes	1,562	1	-	2	2	3	6	6	10	17	32	43
Male	866	-	-	2	2	1	5	3	7	14	23	30
Female	696	1	-	-	-	2	1	3	3	3	9	13
Black, both sexes	186	2	1	-	-	-	4	5	1	10	5	9
Male	106	2	-	-	-	-	2	3	-	7	4	5
Female	80	-	1	-	-	-	2	2	1	3	1	4
American Indian 1/, both sexes	17	-	-	-	-	-	-	1	1	-	2	1
Male	10	-	-	-	-	-	-	1	1	-	2	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	58	-	1	-	1	-	-	-	1	-	1	2
Male	23	-	-	-	-	-	-	-	1	-	1	2
Female	35	-	1	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
New Hampshire												
All races, both sexes	687	-	-	-	1	1	1	4	-	-	3	8
Male	298	-	-	-	-	1	-	2	-	-	1	4
Female	389	-	-	-	1	-	1	2	-	-	2	4
White, both sexes	678	-	-	-	1	1	1	4	-	-	3	7
Male	296	-	-	-	-	1	-	2	-	-	1	4
Female	382	-	-	-	1	-	1	2	-	-	2	3
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	4,976	11	1	2	2	6	12	16	20	42	64	98
Male	2,194	8	-	1	1	4	4	11	18	24	32	58
Female	2,782	3	1	1	1	2	8	5	2	18	32	40
White, both sexes	4,346	5	1	-	1	1	8	7	13	27	39	62
Male	1,897	4	-	-	1	1	3	5	12	15	24	40
Female	2,449	1	1	-	-	-	5	2	1	12	15	22
Black, both sexes	560	6	-	1	1	5	4	7	6	14	24	35
Male	263	4	-	1	-	3	1	5	5	8	8	18
Female	297	2	-	-	1	2	3	2	1	6	16	17
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	67	-	-	1	-	-	-	2	1	1	1	1
Male	33	-	-	-	-	-	-	1	1	1	-	-
Female	34	-	-	1	-	-	-	1	-	-	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
New Mexico												
All races, both sexes	910	2	-	-	-	4	1	5	4	11	7	17
Male	411	2	-	-	-	4	-	4	1	6	6	15
Female	499	-	-	-	-	-	1	1	3	5	1	2
White, both sexes	858	2	-	-	-	3	1	3	3	11	6	14
Male	377	2	-	-	-	3	-	3	-	6	5	12
Female	481	-	-	-	-	-	1	-	3	5	1	2
Black, both sexes	14	-	-	-	-	1	-	-	-	-	1	-
Male	9	-	-	-	-	1	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	34	-	-	-	-	-	-	2	1	-	-	3
Male	23	-	-	-	-	-	-	1	1	-	-	3
Female	11	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	8,418	25	9	6	9	14	32	31	39	68	109	159
Male	3,673	15	4	2	3	9	22	23	33	51	71	105
Female	4,745	10	5	4	6	5	10	8	6	17	38	54
White, both sexes	7,562	15	3	4	4	8	20	17	30	43	77	113
Male	3,266	9	-	1	1	5	14	12	27	31	50	80
Female	4,296	6	3	3	3	3	6	5	3	12	27	33
Black, both sexes	762	9	5	2	5	6	10	14	8	22	31	43
Male	362	5	4	1	2	4	6	11	5	18	20	22
Female	400	4	1	1	3	2	4	3	3	4	11	21
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	1	-	3
Male	7	-	-	-	-	-	-	-	-	1	-	3
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	83	1	1	-	-	-	2	-	1	2	1	-
Male	38	1	-	-	-	-	2	-	1	1	1	-
Female	45	-	1	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
North Carolina												
All races, both sexes	5,333	21	12	7	5	6	16	24	33	67	70	135
Male	2,361	14	6	4	3	4	5	17	21	43	40	77
Female	2,972	7	6	3	2	2	11	7	12	24	30	58
White, both sexes	4,131	12	7	6	2	1	8	10	18	31	40	66
Male	1,817	7	2	3	1	-	2	7	13	21	20	41
Female	2,314	5	5	3	1	1	6	3	5	10	20	25
Black, both sexes	1,144	8	4	1	3	5	6	13	14	34	29	65
Male	519	7	3	1	2	4	2	9	7	21	19	33
Female	625	1	1	-	1	1	4	4	7	13	10	32
American Indian 1/, both sexes	40	1	1	-	-	-	2	-	-	2	1	4
Male	19	-	1	-	-	-	1	-	-	1	1	3
Female	21	1	-	-	-	-	1	-	-	1	-	1
Asian or Pacific Islander 2/, both sexes	18	-	-	-	-	-	-	1	1	-	-	-
Male	6	-	-	-	-	-	-	1	1	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	299	3	1	-	-	-	1	4	3	2	3	7
Male	130	2	-	-	-	-	-	3	1	2	2	4
Female	169	1	1	-	-	-	1	1	2	-	1	3
White, both sexes	283	3	1	-	-	-	-	3	3	1	2	5
Male	119	2	-	-	-	-	-	2	1	1	1	3
Female	164	1	1	-	-	-	-	1	2	-	1	2
Black, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	15	-	-	-	-	-	1	-	-	1	1	2
Male	10	-	-	-	-	-	-	-	-	1	1	1
Female	5	-	-	-	-	-	1	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Ohio												
All races, both sexes	6,137	8	10	4	3	7	23	27	44	56	86	144
Male	2,626	5	6	2	2	4	15	18	24	37	49	88
Female	3,511	3	4	2	1	3	8	9	20	19	37	56
White, both sexes	5,525	3	8	4	1	5	17	18	35	42	65	113
Male	2,336	2	4	2	-	2	10	12	20	26	37	72
Female	3,189	1	4	2	1	3	7	6	15	16	28	41
Black, both sexes	598	5	2	-	1	2	6	8	9	13	21	31
Male	283	3	2	-	1	2	5	5	4	10	12	16
Female	315	2	-	-	-	-	1	3	5	3	9	15
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	1	-	-	1	-	1	-	-
Male	5	-	-	-	1	-	-	1	-	1	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2,695	5	3	2	2	3	2	17	16	23	42	63
Male	1,180	2	1	2	2	1	2	12	14	15	23	42
Female	1,515	3	2	-	-	2	-	5	2	8	19	21
White, both sexes	2,371	3	1	1	2	3	1	13	10	17	33	39
Male	1,031	-	-	1	2	1	1	8	8	11	18	27
Female	1,340	3	1	-	-	2	-	5	2	6	15	12
Black, both sexes	193	-	1	-	-	-	1	-	3	2	5	17
Male	82	-	-	-	-	-	1	-	3	1	3	10
Female	111	-	1	-	-	-	-	-	-	1	2	7
American Indian 1/, both sexes	123	2	1	1	-	-	-	4	3	4	3	7
Male	63	2	1	1	-	-	-	4	3	3	1	5
Female	60	-	-	-	-	-	-	-	-	1	2	2
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Oregon												
All races, both sexes	2,359	5	1	1	1	5	6	9	12	14	14	33
Male	1,007	3	-	1	1	3	3	8	9	6	9	22
Female	1,352	2	1	-	-	2	3	1	3	8	5	11
White, both sexes	2,306	4	1	1	1	4	6	6	10	11	14	32
Male	979	3	-	1	1	2	3	5	7	4	9	21
Female	1,327	1	1	-	-	2	3	1	3	7	5	11
Black, both sexes	14	-	-	-	-	1	-	2	1	3	-	-
Male	10	-	-	-	-	1	-	2	1	2	-	-
Female	4	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	26	1	-	-	-	-	-	1	1	-	-	1
Male	9	-	-	-	-	-	-	1	1	-	-	1
Female	17	1	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	10,479	7	4	5	2	12	33	39	54	65	104	195
Male	4,511	2	-	1	2	8	22	22	34	35	73	114
Female	5,968	5	4	4	-	4	11	17	20	30	31	81
White, both sexes	9,601	7	3	2	2	7	23	26	34	41	75	148
Male	4,084	2	-	-	2	5	16	14	21	20	53	91
Female	5,517	5	3	2	-	2	7	12	13	21	22	57
Black, both sexes	838	-	-	3	-	5	9	13	20	23	28	47
Male	408	-	-	1	-	3	5	8	13	14	19	23
Female	430	-	-	2	-	2	4	5	7	9	9	24
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	39	-	1	-	-	-	1	-	-	1	1	-
Male	19	-	-	-	-	-	1	-	-	1	1	-
Female	20	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Rhode Island												
All races, both sexes	452	2	1	-	-	1	-	1	2	-	4	5
Male	183	1	1	-	-	-	-	1	2	-	3	5
Female	269	1	-	-	-	1	-	-	-	-	1	-
White, both sexes	438	1	1	-	-	-	-	1	2	-	3	5
Male	178	-	1	-	-	-	-	1	2	-	2	5
Female	260	1	-	-	-	-	-	-	-	-	1	-
Black, both sexes	10	-	-	-	-	1	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	3,270	10	1	-	5	5	12	22	30	43	76	85
Male	1,543	7	-	-	5	4	5	13	10	30	49	45
Female	1,727	3	1	-	-	1	7	9	20	13	27	40
White, both sexes	2,275	5	1	-	3	2	7	11	12	18	43	43
Male	1,052	4	-	-	3	1	2	5	3	13	31	24
Female	1,223	1	1	-	-	1	5	6	9	5	12	19
Black, both sexes	984	5	-	-	2	3	5	11	18	25	33	42
Male	486	3	-	-	2	3	3	8	7	17	18	21
Female	498	2	-	-	-	-	2	3	11	8	15	21
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
South Dakota												
All races, both sexes	325	-	2	-	-	2	-	2	4	2	2	6
Male	138	-	2	-	-	1	-	2	2	1	1	3
Female	187	-	-	-	-	1	-	-	2	1	1	3
White, both sexes	308	-	2	-	-	1	-	1	3	2	1	5
Male	129	-	2	-	-	-	-	1	1	1	1	2
Female	179	-	-	-	-	1	-	-	2	1	-	3
Black, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	15	-	-	-	-	1	-	-	1	-	1	1
Male	8	-	-	-	-	1	-	-	1	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	2,905	8	1	2	1	6	18	17	25	34	71	92
Male	1,260	6	-	-	-	5	13	12	14	22	35	60
Female	1,645	2	1	2	1	1	5	5	11	12	36	32
White, both sexes	2,412	4	1	1	1	5	12	15	18	21	48	65
Male	1,043	2	-	-	-	5	10	10	11	14	22	40
Female	1,369	2	1	1	1	-	2	5	7	7	26	25
Black, both sexes	484	4	-	1	-	1	6	2	7	13	23	27
Male	214	4	-	-	-	-	3	2	3	8	13	20
Female	270	-	-	1	-	1	3	-	4	5	10	7
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Texas												
All races, both sexes	11,269	32	13	6	6	27	52	72	62	131	178	269
Male	5,126	15	6	3	2	15	33	45	40	82	117	166
Female	6,143	17	7	3	4	12	19	27	22	49	61	103
White, both sexes	9,738	25	11	3	4	18	36	49	45	97	129	209
Male	4,387	13	6	1	2	9	21	33	29	62	82	136
Female	5,351	12	5	2	2	9	15	16	16	35	47	73
Black, both sexes	1,411	6	2	2	2	9	15	22	17	33	46	58
Male	683	2	-	2	-	6	12	12	11	19	32	28
Female	728	4	2	-	2	3	3	10	6	14	14	30
American Indian 1/, both sexes	19	-	-	-	-	-	1	-	-	-	-	2
Male	11	-	-	-	-	-	-	-	-	-	-	2
Female	8	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	101	1	-	1	-	-	-	1	-	1	3	-
Male	45	-	-	-	-	-	-	-	-	1	3	-
Female	56	1	-	1	-	-	-	1	-	-	-	-
Utah												
All races, both sexes	1,389	3	2	-	1	1	3	7	9	11	16	23
Male	618	2	-	-	1	1	2	4	5	6	8	12
Female	771	1	2	-	-	-	1	3	4	5	8	11
White, both sexes	1,353	3	2	-	1	1	3	7	8	10	13	21
Male	599	2	-	-	1	1	2	4	4	5	6	11
Female	754	1	2	-	-	-	1	3	4	5	7	10
Black, both sexes	8	-	-	-	-	-	-	-	-	-	2	1
Male	5	-	-	-	-	-	-	-	-	-	1	1
Female	3	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	1	1	1	-
Male	7	-	-	-	-	-	-	-	1	1	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	20	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Vermont												
All races, both sexes	263	1	-	-	-	-	1	2	2	-	1	1
Male	118	-	-	-	-	-	-	1	2	-	-	1
Female	145	1	-	-	-	-	1	1	-	-	1	-
White, both sexes	263	1	-	-	-	-	1	2	2	-	1	1
Male	118	-	-	-	-	-	-	1	2	-	-	1
Female	145	1	-	-	-	-	1	1	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	5,029	10	2	3	5	7	21	26	26	38	77	120
Male	2,259	8	2	2	2	4	12	17	17	19	48	71
Female	2,770	2	-	1	3	3	9	9	9	19	29	49
White, both sexes	3,934	5	1	2	2	5	11	15	14	27	37	77
Male	1,728	5	1	2	1	2	5	11	10	13	22	50
Female	2,206	-	-	-	1	3	6	4	4	14	15	27
Black, both sexes	1,032	4	1	-	3	2	8	11	11	10	36	43
Male	499	3	1	-	1	2	6	6	7	5	24	21
Female	533	1	-	-	2	-	2	5	4	5	12	22
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	57	1	-	1	-	-	2	-	1	1	4	-
Male	29	-	-	-	-	-	1	-	-	1	2	-
Female	28	1	-	1	-	-	1	-	1	-	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Washington												
All races, both sexes	2,526	4	2	2	2	4	3	16	17	18	32	57
Male	1,087	1	1	2	-	2	3	12	9	9	20	34
Female	1,439	3	1	-	2	2	-	4	8	9	12	23
White, both sexes	2,388	2	1	2	2	4	3	14	14	15	25	50
Male	1,026	1	1	2	-	2	3	11	8	6	16	30
Female	1,362	1	-	-	2	2	-	3	6	9	9	20
Black, both sexes	57	-	-	-	-	-	-	2	1	1	5	7
Male	29	-	-	-	-	-	-	1	-	1	3	4
Female	28	-	-	-	-	-	-	1	1	-	2	3
American Indian 1/, both sexes	23	1	-	-	-	-	-	-	2	1	1	-
Male	11	-	-	-	-	-	-	-	1	1	-	-
Female	12	1	-	-	-	-	-	-	1	-	1	-
Asian or Pacific Islander 2/, both sexes	58	1	1	-	-	-	-	-	-	1	1	-
Male	21	-	-	-	-	-	-	-	-	1	1	-
Female	37	1	1	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	1,544	2	2	1	1	4	3	6	5	9	11	27
Male	672	-	2	1	1	3	1	3	2	5	8	17
Female	872	2	-	-	-	1	2	3	3	4	3	10
White, both sexes	1,493	1	1	1	1	4	3	6	5	7	10	25
Male	648	-	1	1	1	3	1	3	2	4	7	16
Female	845	1	-	-	-	1	2	3	3	3	3	9
Black, both sexes	49	1	1	-	-	-	-	-	-	1	1	2
Male	22	-	1	-	-	-	-	-	-	-	1	1
Female	27	1	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Wisconsin												
All races, both sexes	3,705	6	1	1	2	6	5	12	13	17	36	63
Male	1,580	5	-	-	1	5	4	8	10	10	28	39
Female	2,125	1	1	1	1	1	1	4	3	7	8	24
White, both sexes	3,532	3	-	1	2	6	3	10	11	15	30	49
Male	1,495	2	-	-	1	5	2	7	8	8	23	28
Female	2,037	1	-	1	1	1	1	3	3	7	7	21
Black, both sexes	146	3	1	-	-	-	2	2	1	2	6	13
Male	73	3	-	-	-	-	2	1	1	2	5	10
Female	73	-	1	-	-	-	-	1	-	-	1	3
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	1	-	-	-
Male	7	-	-	-	-	-	-	-	1	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	328	1	-	-	-	-	-	1	2	2	7	3
Male	143	1	-	-	-	-	-	1	1	1	4	2
Female	185	-	-	-	-	-	-	-	1	1	3	1
White, both sexes	321	1	-	-	-	-	-	-	2	2	6	3
Male	141	1	-	-	-	-	-	-	1	1	4	2
Female	180	-	-	-	-	-	-	-	1	1	2	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	1	-	-	1	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
United States												
All races, both sexes	5,215	6,503	7,743	9,297	12,348	18,764	28,170	33,259	26,410	11,220	2,416	4
Male	3,300	4,104	4,658	5,348	6,772	9,397	12,564	12,713	8,242	2,618	395	4
Female	1,915	2,399	3,085	3,949	5,576	9,367	15,606	20,546	18,168	8,602	2,021	-
White, both sexes	3,701	4,780	6,039	7,364	10,218	16,352	25,300	30,729	24,623	10,364	2,145	4
Male	2,390	3,056	3,676	4,299	5,657	8,284	11,360	11,847	7,725	2,416	339	4
Female	1,311	1,724	2,363	3,065	4,561	8,068	13,940	18,882	16,898	7,948	1,806	-
Black, both sexes	1,408	1,583	1,525	1,712	1,892	2,072	2,376	2,032	1,461	718	243	-
Male	837	957	877	931	977	934	981	663	400	147	45	-
Female	571	626	648	781	915	1,138	1,395	1,369	1,061	571	198	-
American Indian 1/, both sexes	35	54	54	70	52	79	81	69	43	16	8	-
Male	23	30	32	38	31	44	36	24	14	4	4	-
Female	12	24	22	32	21	35	45	45	29	12	4	-
Asian or Pacific Islander 2/, both sexes	71	86	125	151	186	261	413	429	283	122	20	-
Male	50	61	73	80	107	135	187	179	103	51	7	-
Female	21	25	52	71	79	126	226	250	180	71	13	-
Alabama												
All races, both sexes	234	328	324	383	523	701	925	988	705	274	78	-
Male	149	205	205	224	300	344	406	360	208	57	10	-
Female	85	123	119	159	223	357	519	628	497	217	68	-
White, both sexes	136	221	215	272	374	534	735	833	580	221	49	-
Male	84	131	142	167	226	269	324	309	170	45	6	-
Female	52	90	73	105	148	265	411	524	410	176	43	-
Black, both sexes	96	106	108	109	147	163	189	154	124	53	28	-
Male	64	74	62	57	73	71	82	51	38	12	4	-
Female	32	32	46	52	74	92	107	103	86	41	24	-
American Indian 1/, both sexes	-	1	1	1	-	2	-	1	-	-	1	-
Male	-	-	1	-	-	2	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	1	-	-	1	-
Asian or Pacific Islander 2/, both sexes	2	-	-	1	2	2	1	-	1	-	-	-
Male	1	-	-	-	1	2	-	-	-	-	-	-
Female	1	-	-	1	1	-	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Alaska												
All races, both sexes	8	14	4	15	21	26	30	25	23	8	1	-
Male	4	9	3	7	10	17	13	12	6	1	-	-
Female	4	5	1	8	11	9	17	13	17	7	1	-
White, both sexes	4	10	4	8	15	20	23	23	22	6	1	-
Male	2	5	3	4	8	12	9	10	6	1	-	-
Female	2	5	1	4	7	8	14	13	16	5	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	3	-	7	5	6	6	2	1	2	-	-
Male	1	3	-	3	2	5	4	2	-	-	-	-
Female	1	-	-	4	3	1	2	-	1	2	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	1	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	1	-	-	-	-	-
Arizona												
All races, both sexes	72	90	116	116	178	262	342	435	323	132	25	-
Male	51	65	67	68	101	143	159	185	109	31	4	-
Female	21	25	49	48	77	119	183	250	214	101	21	-
White, both sexes	60	76	103	110	168	249	325	424	316	126	24	-
Male	42	56	61	64	97	137	154	179	108	29	4	-
Female	18	20	42	46	71	112	171	245	208	97	20	-
Black, both sexes	8	9	2	4	6	6	5	3	3	2	1	-
Male	5	5	1	2	3	3	2	2	1	1	-	-
Female	3	4	1	2	3	3	3	1	2	1	1	-
American Indian 1/, both sexes	4	4	8	2	3	6	9	7	4	4	-	-
Male	4	3	4	2	1	3	3	4	-	1	-	-
Female	-	1	4	-	2	3	6	3	4	3	-	-
Asian or Pacific Islander 2/, both sexes	-	1	3	-	1	1	3	1	-	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	2	-	1	1	3	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Arkansas												
All races, both sexes	55	83	111	113	160	230	333	345	281	151	30	-
Male	29	50	54	61	93	116	129	110	82	28	6	-
Female	26	33	57	52	67	114	204	235	199	123	24	-
White, both sexes	38	53	88	89	147	201	305	314	256	141	23	-
Male	20	35	45	48	86	101	119	103	74	27	5	-
Female	18	18	43	41	61	100	186	211	182	114	18	-
Black, both sexes	17	29	20	24	13	28	26	28	25	9	7	-
Male	9	15	8	13	7	14	9	7	8	1	1	-
Female	8	14	12	11	6	14	17	21	17	8	6	-
American Indian 1/, both sexes	-	1	2	-	-	1	1	3	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	2	-	-	-	1	3	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	1	-	-	1	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
California												
All races, both sexes	417	484	498	591	866	1,358	2,190	2,714	2,281	954	203	-
Male	260	306	308	350	470	682	999	1,111	786	268	33	-
Female	157	178	190	241	396	676	1,191	1,603	1,495	686	170	-
White, both sexes	317	372	387	467	721	1,188	1,962	2,463	2,101	882	191	-
Male	206	238	237	279	392	599	895	1,000	721	247	32	-
Female	111	134	150	188	329	589	1,067	1,463	1,380	635	159	-
Black, both sexes	75	90	70	75	84	89	96	98	80	31	6	-
Male	39	52	41	46	42	36	39	42	25	4	1	-
Female	36	38	29	29	42	53	57	56	55	27	5	-
American Indian 1/, both sexes	3	4	5	8	5	7	10	6	6	1	1	-
Male	2	3	4	5	4	5	3	3	3	1	-	-
Female	1	1	1	3	1	2	7	3	3	-	1	-
Asian or Pacific Islander 2/, both sexes	22	18	36	41	56	74	122	147	94	40	5	-
Male	13	13	26	20	32	42	62	66	37	16	-	-
Female	9	5	10	21	24	32	60	81	57	24	5	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Colorado												
All races, both sexes	57	61	77	78	122	177	286	385	316	125	25	-
Male	38	34	49	46	71	84	124	149	97	27	5	-
Female	19	27	28	32	51	93	162	236	219	98	20	-
White, both sexes	52	56	71	74	113	170	276	378	308	122	25	-
Male	34	32	44	44	65	81	122	148	92	27	5	-
Female	18	24	27	30	48	89	154	230	216	95	20	-
Black, both sexes	4	5	4	3	7	4	7	4	4	2	-	-
Male	3	2	4	1	5	1	1	1	3	-	-	-
Female	1	3	-	2	2	3	6	3	1	2	-	-
American Indian 1/, both sexes	-	-	2	1	-	1	-	-	2	-	-	-
Male	-	-	1	1	-	1	-	-	1	-	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	2	2	3	3	2	1	-	-
Male	1	-	-	-	1	1	1	-	1	-	-	-
Female	-	-	-	-	1	1	2	3	1	1	-	-
Connecticut												
All races, both sexes	72	79	103	123	169	270	412	581	457	207	44	-
Male	52	56	62	75	104	141	199	215	147	54	9	-
Female	20	23	41	48	65	129	213	366	310	153	35	-
White, both sexes	59	66	88	100	147	251	400	561	442	201	43	-
Male	45	49	54	62	93	130	195	209	145	53	9	-
Female	14	17	34	38	54	121	205	352	297	148	34	-
Black, both sexes	12	13	15	21	22	18	11	18	14	5	1	-
Male	7	7	8	11	11	11	3	5	1	1	-	-
Female	5	6	7	10	11	7	8	13	13	4	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	2	-	1	1	2	1	-	-	-
Male	-	-	-	2	-	-	1	1	1	-	-	-
Female	1	-	-	-	-	1	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Delaware												
All races, both sexes	22	23	20	38	40	70	91	97	76	27	4	1
Male	14	16	12	24	24	26	32	38	25	7	1	1
Female	8	7	8	14	16	44	59	59	51	20	3	-
White, both sexes	16	14	12	30	34	63	78	91	69	26	3	1
Male	11	10	7	17	21	24	27	36	23	7	-	1
Female	5	4	5	13	13	39	51	55	46	19	3	-
Black, both sexes	6	8	8	8	6	7	12	5	7	1	1	-
Male	3	5	5	7	3	2	5	1	2	-	1	-
Female	3	3	3	1	3	5	7	4	5	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	10	12	16	17	20	12	41	39	40	20	4	-
Male	7	6	10	8	10	7	13	14	10	4	2	-
Female	3	6	6	9	10	5	28	25	30	16	2	-
White, both sexes	1	1	3	1	2	1	10	13	20	5	2	-
Male	1	-	1	1	1	1	3	6	8	1	1	-
Female	-	1	2	-	1	-	7	7	12	4	1	-
Black, both sexes	9	11	13	16	18	11	31	26	20	15	2	-
Male	6	6	9	7	9	6	10	8	2	3	1	-
Female	3	5	4	9	9	5	21	18	18	12	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Florida												
All races, both sexes	282	349	389	528	668	1,104	1,601	1,850	1,388	583	99	-
Male	170	225	238	337	376	613	817	810	529	166	33	-
Female	112	124	151	191	292	491	784	1,040	859	417	66	-
White, both sexes	184	268	311	437	581	996	1,480	1,758	1,333	560	87	-
Male	115	183	199	277	326	567	762	783	507	158	28	-
Female	69	85	112	160	255	429	718	975	826	402	59	-
Black, both sexes	97	78	75	85	84	101	111	86	53	20	11	-
Male	55	39	37	55	47	42	50	25	21	5	4	-
Female	42	39	38	30	37	59	61	61	32	15	7	-
American Indian 1/, both sexes	-	1	1	1	-	-	1	-	-	-	1	-
Male	-	1	1	1	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	2	5	3	7	9	6	2	3	-	-
Male	-	2	1	4	3	4	5	2	1	3	-	-
Female	1	-	1	1	-	3	4	4	1	-	-	-
Georgia												
All races, both sexes	298	375	445	506	578	744	1,057	1,104	838	353	79	-
Male	184	253	280	271	308	374	468	399	239	66	17	-
Female	114	122	165	235	270	370	589	705	599	287	62	-
White, both sexes	179	238	286	335	390	570	846	912	707	273	53	-
Male	115	159	180	192	228	285	389	335	205	48	13	-
Female	64	79	106	143	162	285	457	577	502	225	40	-
Black, both sexes	118	136	156	170	184	165	204	188	127	78	26	-
Male	68	94	99	79	78	85	77	63	34	18	4	-
Female	50	42	57	91	106	80	127	125	93	60	22	-
American Indian 1/, both sexes	-	-	-	-	1	1	1	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	3	1	3	8	6	3	4	2	-	-
Male	1	-	1	-	1	4	2	1	-	-	-	-
Female	-	1	2	1	2	4	4	2	4	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Hawaii												
All races, both sexes	36	40	43	64	50	99	149	199	134	56	15	-
Male	31	29	25	38	33	55	78	101	58	23	7	-
Female	5	11	18	26	17	44	71	98	76	33	8	-
White, both sexes	15	18	13	16	8	20	31	47	38	21	5	-
Male	13	12	9	11	4	8	18	27	21	5	2	-
Female	2	6	4	5	4	12	13	20	17	16	3	-
Black, both sexes	-	1	-	-	2	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	2	-	-	-	-
Female	-	1	-	-	2	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	1	-	1	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	21	21	30	47	40	79	117	150	95	35	10	-
Male	18	17	16	26	29	47	59	72	37	18	5	-
Female	3	4	14	21	11	32	58	78	58	17	5	-
Idaho												
All races, both sexes	12	31	30	45	49	71	127	171	122	63	11	-
Male	6	22	18	27	25	39	63	64	48	17	1	-
Female	6	9	12	18	24	32	64	107	74	46	10	-
White, both sexes	11	29	30	44	48	68	123	169	121	63	11	-
Male	6	21	18	27	25	37	59	62	47	17	1	-
Female	5	8	12	17	23	31	64	107	74	46	10	-
Black, both sexes	1	1	-	-	-	1	-	1	-	-	-	-
Male	-	1	-	-	-	1	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	1	1	2	1	-	1	-	-	-
Male	-	-	-	-	-	1	1	-	1	-	-	-
Female	-	1	-	1	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	3	1	-	-	-	-
Male	-	-	-	-	-	-	3	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Illinois												
All races, both sexes	209	272	310	405	513	800	1,223	1,461	1,238	591	132	-
Male	123	174	180	240	272	377	529	572	383	125	16	-
Female	86	98	130	165	241	423	694	889	855	466	116	-
White, both sexes	147	194	231	300	399	677	1,089	1,363	1,165	554	116	-
Male	87	124	125	177	214	315	466	534	361	121	13	-
Female	60	70	106	123	185	362	623	829	804	433	103	-
Black, both sexes	62	76	74	103	105	116	120	93	67	35	15	-
Male	36	49	50	63	53	59	56	37	20	4	2	-
Female	26	27	24	40	52	57	64	56	47	31	13	-
American Indian 1/, both sexes	-	1	-	1	-	-	1	-	-	-	1	-
Male	-	-	-	-	-	-	1	-	-	-	1	-
Female	-	1	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	5	1	9	7	13	5	6	2	-	-
Male	-	1	5	-	5	3	6	1	2	-	-	-
Female	-	-	-	1	4	4	7	4	4	2	-	-
Indiana												
All races, both sexes	140	200	260	268	347	511	710	846	692	311	70	-
Male	89	125	154	152	188	243	322	333	189	61	13	-
Female	51	75	106	116	159	268	388	513	503	250	57	-
White, both sexes	113	169	232	235	315	481	661	822	668	303	69	-
Male	71	105	138	133	169	232	303	324	184	58	13	-
Female	42	64	94	102	146	249	358	498	484	245	56	-
Black, both sexes	27	30	28	33	31	29	49	24	23	7	1	-
Male	18	19	16	19	18	11	19	9	5	3	-	-
Female	9	11	12	14	13	18	30	15	18	4	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	1	-	-	1	1	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Iowa												
All races, both sexes	24	53	49	51	81	135	206	255	267	130	23	-
Male	13	31	26	32	35	73	91	103	87	32	1	-
Female	11	22	23	19	46	62	115	152	180	98	22	-
White, both sexes	22	51	46	50	79	132	205	255	267	130	23	-
Male	13	30	26	32	34	71	90	103	87	32	1	-
Female	9	21	20	18	45	61	115	152	180	98	22	-
Black, both sexes	2	2	2	1	2	3	1	-	-	-	-	-
Male	-	1	-	-	1	2	1	-	-	-	-	-
Female	2	1	2	1	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	65	57	79	76	138	220	346	449	427	200	57	-
Male	43	37	47	39	83	90	143	172	138	45	5	-
Female	22	20	32	37	55	130	203	277	289	155	52	-
White, both sexes	55	47	69	63	127	204	330	434	415	194	53	-
Male	36	30	42	32	77	85	137	167	136	41	4	-
Female	19	17	27	31	50	119	193	267	279	153	49	-
Black, both sexes	8	8	9	10	8	12	15	11	9	4	4	-
Male	6	6	5	6	4	3	6	5	2	4	1	-
Female	2	2	4	4	4	9	9	6	7	-	3	-
American Indian 1/, both sexes	1	1	1	3	2	3	-	3	-	2	-	-
Male	1	-	-	1	2	2	-	-	-	-	-	-
Female	-	1	1	2	-	1	-	3	-	2	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	1	1	1	1	3	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	1	1	1	3	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Kentucky												
All races, both sexes	106	158	156	228	260	412	560	562	433	172	35	-
Male	76	96	85	133	141	201	234	184	121	33	5	-
Female	30	62	71	95	119	211	326	378	312	139	30	-
White, both sexes	95	139	148	206	241	385	528	543	411	166	31	-
Male	71	83	81	122	127	192	222	176	111	32	5	-
Female	24	56	67	84	114	193	306	367	300	134	26	-
Black, both sexes	11	17	8	20	19	27	32	19	21	6	4	-
Male	5	11	4	10	14	9	12	8	9	1	-	-
Female	6	6	4	10	5	18	20	11	12	5	4	-
American Indian 1/, both sexes	-	2	-	1	-	-	-	-	1	-	-	-
Male	-	2	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	123	139	171	215	267	369	600	589	433	166	38	-
Male	77	82	89	138	139	183	237	214	113	35	7	-
Female	46	57	82	77	128	186	363	375	320	131	31	-
White, both sexes	58	82	102	130	181	271	468	503	351	139	33	-
Male	37	48	58	81	95	146	182	185	100	29	6	-
Female	21	34	44	49	86	125	286	318	251	110	27	-
Black, both sexes	65	56	68	84	84	98	129	84	80	26	5	-
Male	40	34	31	56	44	37	53	27	13	6	1	-
Female	25	22	37	28	40	61	76	57	67	20	4	-
American Indian 1/, both sexes	-	1	1	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	2	-	3	1	2	1	-	-
Male	-	-	-	1	-	-	2	1	-	-	-	-
Female	-	-	-	-	2	-	1	-	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Maine												
All races, both sexes	23	31	52	51	76	126	157	208	147	68	16	-
Male	17	18	35	28	47	68	74	80	49	22	2	-
Female	6	13	17	23	29	58	83	128	98	46	14	-
White, both sexes	23	31	51	51	75	126	156	208	147	67	16	-
Male	17	18	34	28	46	68	73	80	49	21	2	-
Female	6	13	17	23	29	58	83	128	98	46	14	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	1	-	1	-	-	-	-	-
Male	-	-	1	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	83	93	120	153	175	259	407	484	335	154	27	-
Male	55	51	79	83	90	127	169	186	117	37	7	-
Female	28	42	41	70	85	132	238	298	218	117	20	-
White, both sexes	34	56	69	87	115	184	324	408	301	122	23	-
Male	25	31	45	45	57	99	138	162	108	28	6	-
Female	9	25	24	42	58	85	186	246	193	94	17	-
Black, both sexes	48	36	48	62	57	66	77	67	31	30	4	-
Male	29	19	32	35	32	23	30	21	8	9	1	-
Female	19	17	16	27	25	43	47	46	23	21	3	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	2	4	3	9	6	9	2	2	-	-
Male	1	1	2	3	1	5	1	3	1	-	-	-
Female	-	-	-	1	2	4	5	6	1	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Massachusetts												
All races, both sexes	74	90	119	162	222	402	717	893	762	320	89	-
Male	42	62	66	102	138	216	302	317	228	71	14	-
Female	32	28	53	60	84	186	415	576	534	249	75	-
White, both sexes	66	78	98	143	202	384	688	868	751	308	85	-
Male	37	52	53	89	127	206	289	309	224	68	13	-
Female	29	26	45	54	75	178	399	559	527	240	72	-
Black, both sexes	7	10	18	15	17	14	22	21	8	9	4	-
Male	4	8	12	10	10	7	12	5	4	1	1	-
Female	3	2	6	5	7	7	10	16	4	8	3	-
American Indian 1/, both sexes	-	-	-	2	-	1	1	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	3	2	3	3	6	4	3	3	-	-
Male	1	2	1	2	1	2	1	3	-	2	-	-
Female	-	-	2	-	2	1	5	1	3	1	-	-
Michigan												
All races, both sexes	203	191	229	278	392	617	950	1,180	975	409	90	-
Male	140	121	138	162	189	312	421	435	295	95	10	-
Female	63	70	91	116	203	305	529	745	680	314	80	-
White, both sexes	131	125	175	223	325	540	840	1,112	920	383	78	-
Male	90	85	109	131	159	272	379	412	275	88	8	-
Female	41	40	66	92	166	268	461	700	645	295	70	-
Black, both sexes	71	65	50	52	64	75	101	62	51	23	10	-
Male	49	35	27	30	29	39	40	22	17	7	1	-
Female	22	30	23	22	35	36	61	40	34	16	9	-
American Indian 1/, both sexes	1	-	1	-	2	1	4	3	1	-	1	-
Male	1	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	1	-	1	-	4	3	1	-	1	-
Asian or Pacific Islander 2/, both sexes	-	1	3	3	1	1	5	3	3	3	1	-
Male	-	1	2	1	-	-	2	1	3	-	1	-
Female	-	-	1	2	1	1	3	2	-	3	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Minnesota												
All races, both sexes	58	51	101	101	150	292	431	657	592	271	54	-
Male	41	32	59	61	90	151	223	271	187	67	6	-
Female	17	19	42	40	60	141	208	386	405	204	48	-
White, both sexes	49	45	96	92	146	281	420	651	587	270	53	-
Male	35	29	57	55	88	146	217	271	185	67	6	-
Female	14	16	39	37	58	135	203	380	402	203	47	-
Black, both sexes	5	4	3	5	2	10	5	3	4	-	1	-
Male	3	3	1	4	2	5	3	-	2	-	-	-
Female	2	1	2	1	-	5	2	3	2	-	1	-
American Indian 1/, both sexes	2	2	2	4	1	-	3	1	1	-	-	-
Male	1	-	1	2	-	-	2	-	-	-	-	-
Female	1	2	1	2	1	-	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	1	1	3	2	-	1	-	-
Male	2	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	1	2	2	-	1	-	-
Mississippi												
All races, both sexes	114	127	170	204	272	354	504	513	380	201	49	1
Male	69	85	94	108	142	159	218	188	95	44	14	1
Female	45	42	76	96	130	195	286	325	285	157	35	-
White, both sexes	53	63	93	120	177	239	386	390	289	139	24	1
Male	31	42	53	66	99	108	181	151	78	29	7	1
Female	22	21	40	54	78	131	205	239	211	110	17	-
Black, both sexes	61	64	76	82	93	113	117	122	91	61	25	-
Male	38	43	41	41	41	49	37	37	17	15	7	-
Female	23	21	35	41	52	64	80	85	74	46	18	-
American Indian 1/, both sexes	-	-	1	2	1	1	-	1	-	1	-	-
Male	-	-	-	1	1	1	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	1	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Missouri												
All races, both sexes	111	121	193	221	299	457	634	735	671	285	63	-
Male	69	81	102	127	160	234	288	257	207	71	8	-
Female	42	40	91	94	139	223	346	478	464	214	55	-
White, both sexes	89	94	168	202	266	417	601	697	645	273	60	-
Male	54	67	88	115	141	219	276	247	196	67	7	-
Female	35	27	80	87	125	198	325	450	449	206	53	-
Black, both sexes	21	26	25	18	32	38	32	36	24	12	3	-
Male	14	14	14	11	18	14	12	10	11	4	1	-
Female	7	12	11	7	14	24	20	26	13	8	2	-
American Indian 1/, both sexes	1	-	-	-	1	-	-	1	1	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	2	1	1	1	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	1	-	-	-	1	1	1	1	-	-	-
Montana												
All races, both sexes	11	17	21	38	47	73	116	162	155	60	9	-
Male	10	10	12	23	28	37	57	69	56	17	3	-
Female	1	7	9	15	19	36	59	93	99	43	6	-
White, both sexes	11	13	20	34	45	69	114	162	155	60	9	-
Male	10	8	11	20	26	36	56	69	56	17	3	-
Female	1	5	9	14	19	33	58	93	99	43	6	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	4	1	4	2	4	1	-	-	-	-	-
Male	-	2	1	3	2	1	1	-	-	-	-	-
Female	-	2	-	1	-	3	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Nebraska												
All races, both sexes	36	41	46	76	85	120	212	265	283	132	28	-
Male	24	28	26	58	48	67	90	93	84	22	3	-
Female	12	13	20	18	37	53	122	172	199	110	25	-
White, both sexes	32	37	43	71	82	115	206	261	275	130	28	-
Male	21	27	25	56	46	64	90	92	83	22	3	-
Female	11	10	18	15	36	51	116	169	192	108	25	-
Black, both sexes	3	4	2	3	2	5	6	1	8	2	-	-
Male	2	1	1	1	1	3	-	-	1	-	-	-
Female	1	3	1	2	1	2	6	1	7	2	-	-
American Indian 1/, both sexes	1	-	1	2	-	-	-	1	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	2	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Nevada												
All races, both sexes	71	108	118	166	171	236	315	262	154	40	11	1
Male	52	69	77	106	101	138	171	112	48	11	1	1
Female	19	39	41	60	70	98	144	150	106	29	10	-
White, both sexes	49	79	101	137	142	204	290	245	147	36	9	1
Male	37	51	67	90	85	125	159	108	45	10	1	1
Female	12	28	34	47	57	79	131	137	102	26	8	-
Black, both sexes	17	21	13	20	22	25	17	6	4	3	1	-
Male	12	13	8	12	12	13	10	1	2	-	-	-
Female	5	8	5	8	10	12	7	5	2	3	1	-
American Indian 1/, both sexes	2	2	1	1	1	1	1	1	2	-	-	-
Male	1	2	1	1	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	1	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes	3	6	3	8	6	6	7	10	1	1	1	-
Male	2	3	1	3	3	-	2	3	1	1	-	-
Female	1	3	2	5	3	6	5	7	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
New Hampshire												
All races, both sexes	17	16	39	27	49	71	116	153	114	48	19	-
Male	14	12	25	17	29	33	45	70	33	9	3	-
Female	3	4	14	10	20	38	71	83	81	39	16	-
White, both sexes	16	16	38	26	49	67	116	153	113	48	19	-
Male	14	12	24	16	29	33	45	70	33	9	3	-
Female	2	4	14	10	20	34	71	83	80	39	16	-
Black, both sexes	1	-	1	1	-	1	-	-	1	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	3	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	3	-	-	-	-	-	-
New Jersey												
All races, both sexes	109	144	180	220	340	501	876	1,110	810	343	69	-
Male	65	84	112	121	198	248	408	434	268	80	15	-
Female	44	60	68	99	142	253	468	676	542	263	54	-
White, both sexes	82	105	143	157	271	423	803	1,041	773	321	63	-
Male	53	66	91	85	161	212	370	409	257	73	15	-
Female	29	39	52	72	110	211	433	632	516	248	48	-
Black, both sexes	26	36	32	55	64	71	62	61	30	15	5	-
Male	11	16	19	30	33	33	33	22	9	4	-	-
Female	15	20	13	25	31	38	29	39	21	11	5	-
American Indian 1/, both sexes	-	1	-	2	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	5	6	5	7	11	8	7	7	1	-
Male	1	2	2	5	4	3	5	3	2	3	-	-
Female	-	-	3	1	1	4	6	5	5	4	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
New Mexico												
All races, both sexes	21	39	44	40	56	108	165	183	137	53	13	-
Male	13	22	19	23	33	54	75	65	47	17	5	-
Female	8	17	25	17	23	54	90	118	90	36	8	-
White, both sexes	20	34	42	38	51	100	155	179	134	51	11	-
Male	12	20	17	22	31	47	69	62	45	17	4	-
Female	8	14	25	16	20	53	86	117	89	34	7	-
Black, both sexes	-	1	-	-	2	2	5	-	-	2	-	-
Male	-	-	-	-	2	2	3	-	-	-	-	-
Female	-	1	-	-	-	-	2	-	-	2	-	-
American Indian 1/, both sexes	1	3	2	2	3	5	4	3	3	-	2	-
Male	1	2	2	1	-	5	2	2	2	-	1	-
Female	-	1	-	1	3	-	2	1	1	-	1	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
New York												
All races, both sexes	227	263	311	362	554	852	1,401	1,738	1,447	614	148	-
Male	152	158	199	199	315	420	632	643	461	139	17	-
Female	75	105	112	163	239	432	769	1,095	986	475	131	-
White, both sexes	172	205	252	294	477	758	1,312	1,643	1,385	593	137	-
Male	121	127	161	162	268	379	595	622	448	136	17	-
Female	51	78	91	132	209	379	717	1,021	937	457	120	-
Black, both sexes	49	52	55	63	64	82	72	83	59	17	11	-
Male	27	28	37	34	38	37	31	17	13	2	-	-
Female	22	24	18	29	26	45	41	66	46	15	11	-
American Indian 1/, both sexes	1	1	-	1	1	1	1	1	-	-	-	-
Male	-	1	-	-	1	-	-	1	-	-	-	-
Female	1	-	-	1	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	5	4	4	12	11	16	11	3	4	-	-
Male	4	2	1	3	8	4	6	3	-	1	-	-
Female	1	3	3	1	4	7	10	8	3	3	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
North Carolina												
All races, both sexes	170	210	282	316	443	612	910	943	703	281	67	-
Male	106	125	178	166	249	299	386	354	203	55	6	-
Female	64	85	104	150	194	313	524	589	500	226	61	-
White, both sexes	103	128	197	212	322	488	758	813	629	226	54	-
Male	70	83	125	119	184	241	331	319	181	44	3	-
Female	33	45	72	93	138	247	427	494	448	182	51	-
Black, both sexes	67	80	80	99	116	118	144	121	70	55	12	-
Male	36	41	50	45	63	55	53	33	21	11	3	-
Female	31	39	30	54	53	63	91	88	49	44	9	-
American Indian 1/, both sexes	-	1	2	4	4	3	5	7	3	-	-	-
Male	-	-	2	2	2	1	2	2	1	-	-	-
Female	-	1	-	2	2	2	3	5	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	3	1	1	3	3	2	1	-	1	-
Male	-	1	1	-	-	2	-	-	-	-	-	-
Female	-	-	2	1	1	1	3	2	1	-	1	-
North Dakota												
All races, both sexes	9	10	11	10	12	22	30	74	59	30	8	-
Male	7	7	6	3	7	17	16	30	18	5	-	-
Female	2	3	5	7	5	5	14	44	41	25	8	-
White, both sexes	8	7	8	10	12	21	30	72	59	30	8	-
Male	6	4	5	3	7	16	16	29	18	5	-	-
Female	2	3	3	7	5	5	14	43	41	25	8	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	3	3	-	-	1	-	2	-	-	-	-
Male	1	3	1	-	-	1	-	1	-	-	-	-
Female	-	-	2	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Ohio												
All races, both sexes	205	239	280	302	457	665	1,050	1,183	939	344	61	-
Male	112	159	169	158	210	318	452	426	284	80	8	-
Female	93	80	111	144	247	347	598	757	655	264	53	-
White, both sexes	157	195	227	252	398	613	979	1,121	887	327	58	-
Male	90	136	142	135	183	292	413	405	269	77	7	-
Female	67	59	85	117	215	321	566	716	618	250	51	-
Black, both sexes	46	44	53	49	58	52	68	60	50	17	3	-
Male	20	23	27	23	27	26	37	21	15	3	1	-
Female	26	21	26	26	31	26	31	39	35	14	2	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	2	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	3	2	-	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	1	1	-	1	2	-	-	-	-
Oklahoma												
All races, both sexes	85	97	144	167	215	319	446	492	365	153	34	-
Male	44	55	87	87	100	152	196	194	115	27	7	-
Female	41	42	57	80	115	167	250	298	250	126	27	-
White, both sexes	61	72	119	142	187	283	409	457	342	144	32	-
Male	35	45	70	76	89	137	180	185	108	23	6	-
Female	26	27	49	66	98	146	229	272	234	121	26	-
Black, both sexes	18	13	13	14	19	20	19	25	16	6	1	-
Male	7	7	8	6	7	10	7	6	4	2	-	-
Female	11	6	5	8	12	10	12	19	12	4	1	-
American Indian 1/, both sexes	6	11	10	9	9	16	17	10	7	2	1	-
Male	2	3	8	5	4	5	8	3	3	1	1	-
Female	4	8	2	4	5	11	9	7	4	1	-	-
Asian or Pacific Islander 2/, both sexes	-	1	2	2	-	-	1	-	-	1	-	-
Male	-	-	1	-	-	-	1	-	-	1	-	-
Female	-	1	1	2	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Oregon												
All races, both sexes	55	65	62	102	133	221	403	519	463	190	45	-
Male	40	34	31	60	70	107	179	195	160	58	8	-
Female	15	31	31	42	63	114	224	324	303	132	37	-
White, both sexes	53	65	60	99	128	220	390	509	459	188	45	-
Male	38	34	29	58	67	106	173	194	159	57	8	-
Female	15	31	31	41	61	114	217	315	300	131	37	-
Black, both sexes	1	-	-	-	2	-	1	2	1	-	-	-
Male	1	-	-	-	2	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	1	-	-	-
American Indian 1/, both sexes	1	-	2	3	2	1	2	2	-	-	-	-
Male	1	-	2	2	1	1	2	-	-	-	-	-
Female	-	-	-	1	1	-	-	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	10	6	3	2	-	-
Male	-	-	-	-	-	-	3	1	1	1	-	-
Female	-	-	-	-	1	-	7	5	2	1	-	-
Pennsylvania												
All races, both sexes	258	357	428	508	657	1,145	1,797	2,324	1,704	645	136	-
Male	166	226	272	283	369	567	771	869	507	151	17	-
Female	92	131	156	225	288	578	1,026	1,455	1,197	494	119	-
White, both sexes	194	284	376	445	594	1,059	1,680	2,227	1,628	617	129	-
Male	127	179	239	252	329	533	718	834	487	146	16	-
Female	67	105	137	193	265	526	962	1,393	1,141	471	113	-
Black, both sexes	63	71	50	61	57	83	113	91	67	27	7	-
Male	38	45	32	30	37	33	53	32	17	4	1	-
Female	25	26	18	31	20	50	60	59	50	23	6	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	2	2	6	3	4	5	9	1	-	-
Male	1	2	1	1	3	1	-	3	3	1	-	-
Female	-	-	1	1	3	2	4	2	6	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Rhode Island												
All races, both sexes	7	10	20	22	29	50	80	100	71	42	5	-
Male	5	6	14	14	19	16	31	32	20	13	-	-
Female	2	4	6	8	10	34	49	68	51	29	5	-
White, both sexes	6	9	20	20	28	50	77	98	71	41	5	-
Male	5	6	14	13	18	16	31	32	20	12	-	-
Female	1	3	6	7	10	34	46	66	51	29	5	-
Black, both sexes	1	1	-	2	1	-	2	1	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	1	1	-	1	-	-	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
South Carolina												
All races, both sexes	149	199	232	246	273	391	475	501	347	139	28	1
Male	98	140	135	149	149	188	210	179	93	29	4	1
Female	51	59	97	97	124	203	265	322	254	110	24	-
White, both sexes	73	99	136	150	187	289	375	413	276	112	19	1
Male	45	71	82	92	99	144	169	159	74	26	4	1
Female	28	28	54	58	88	145	206	254	202	86	15	-
Black, both sexes	76	97	95	96	86	101	98	86	69	27	9	-
Male	53	67	53	57	50	43	40	20	18	3	-	-
Female	23	30	42	39	36	58	58	66	51	24	9	-
American Indian 1/, both sexes	-	2	1	-	-	-	1	-	1	-	-	-
Male	-	2	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	1	2	1	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	1	-	-	-	-	1	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
South Dakota												
All races, both sexes	8	7	9	15	29	32	44	60	62	32	7	-
Male	3	3	5	10	21	21	15	16	20	11	1	-
Female	5	4	4	5	8	11	29	44	42	21	6	-
White, both sexes	7	7	7	14	27	31	43	58	60	32	7	-
Male	3	3	5	9	20	20	15	15	19	11	1	-
Female	4	4	2	5	7	11	28	43	41	21	6	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	1	2	1	1	2	2	-	-	-
Male	-	-	-	1	1	1	-	1	1	-	-	-
Female	1	-	1	-	1	-	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	97	128	158	197	225	351	470	502	338	138	26	-
Male	60	87	89	103	129	172	195	147	86	22	3	-
Female	37	41	69	94	96	179	275	355	252	116	23	-
White, both sexes	62	87	127	156	186	312	396	446	309	119	21	-
Male	41	61	72	84	116	155	170	129	80	20	1	-
Female	21	26	55	72	70	157	226	317	229	99	20	-
Black, both sexes	35	41	31	40	39	37	70	56	29	17	5	-
Male	19	26	17	19	13	16	24	18	6	1	2	-
Female	16	15	14	21	26	21	46	38	23	16	3	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	4	-	-	2	-	-
Male	-	-	-	-	-	-	1	-	-	1	-	-
Female	-	-	-	1	-	1	3	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Texas												
All races, both sexes	390	480	558	686	908	1,364	1,751	2,000	1,489	659	136	-
Male	237	290	342	379	506	681	779	770	449	139	30	-
Female	153	190	216	307	402	683	972	1,230	1,040	520	106	-
White, both sexes	290	351	440	567	757	1,213	1,576	1,834	1,375	592	117	-
Male	179	210	276	317	423	609	703	707	419	126	24	-
Female	111	141	164	250	334	604	873	1,127	956	466	93	-
Black, both sexes	95	120	114	113	140	135	155	149	96	63	19	-
Male	56	73	65	59	77	64	64	58	25	12	6	-
Female	39	47	49	54	63	71	91	91	71	51	13	-
American Indian 1/, both sexes	2	2	-	2	1	3	3	2	1	-	-	-
Male	1	2	-	1	1	2	1	1	-	-	-	-
Female	1	-	-	1	-	1	2	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	3	7	4	4	10	13	17	15	17	4	-	-
Male	1	5	1	2	5	6	11	4	5	1	-	-
Female	2	2	3	2	5	7	6	11	12	3	-	-
Utah												
All races, both sexes	30	42	43	82	95	160	272	266	218	98	7	-
Male	20	21	27	52	39	83	120	97	75	40	3	-
Female	10	21	16	30	56	77	152	169	143	58	4	-
White, both sexes	28	38	42	78	93	158	266	263	213	98	7	-
Male	18	20	26	50	38	82	117	96	73	40	3	-
Female	10	18	16	28	55	76	149	167	140	58	4	-
Black, both sexes	1	1	-	1	-	-	1	-	1	-	-	-
Male	1	-	-	1	-	-	-	-	1	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	1	2	-	1	-	-	1	-	-	-	-	-
Male	1	1	-	1	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	2	2	2	4	3	4	-	-	-
Male	-	-	1	-	1	1	2	1	1	-	-	-
Female	-	1	-	2	1	1	2	2	3	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Vermont												
All races, both sexes	4	8	11	15	15	30	44	55	46	23	4	-
Male	2	6	7	9	10	11	20	26	15	6	2	-
Female	2	2	4	6	5	19	24	29	31	17	2	-
White, both sexes	4	8	11	15	15	30	44	55	46	23	4	-
Male	2	6	7	9	10	11	20	26	15	6	2	-
Female	2	2	4	6	5	19	24	29	31	17	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	151	223	247	288	394	577	817	952	691	292	62	-
Male	98	133	149	162	215	293	365	359	210	66	7	-
Female	53	90	98	126	179	284	452	593	481	226	55	-
White, both sexes	94	128	155	204	282	447	682	831	605	257	53	-
Male	57	73	101	119	155	230	305	318	184	59	5	-
Female	37	55	54	85	127	217	377	513	421	198	48	-
Black, both sexes	56	94	87	81	104	124	126	114	78	31	8	-
Male	40	60	43	42	55	60	56	39	21	6	1	-
Female	16	34	44	39	49	64	70	75	57	25	7	-
American Indian 1/, both sexes	-	-	-	-	1	3	-	-	-	2	-	-
Male	-	-	-	-	1	1	-	-	-	1	-	-
Female	-	-	-	-	-	2	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	1	5	3	7	3	9	7	8	2	1	-
Male	1	-	5	1	4	2	4	2	5	-	1	-
Female	-	1	-	2	3	1	5	5	3	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Washington												
All races, both sexes	74	78	119	112	162	216	383	514	471	205	35	-
Male	43	49	61	69	93	104	161	212	147	48	7	-
Female	31	29	58	43	69	112	222	302	324	157	28	-
White, both sexes	64	72	107	104	151	201	366	494	459	204	34	-
Male	37	44	56	63	87	99	156	204	145	48	7	-
Female	27	28	51	41	64	102	210	290	314	156	27	-
Black, both sexes	6	2	6	2	8	6	3	4	2	1	1	-
Male	3	2	3	2	4	3	1	2	-	-	-	-
Female	3	-	3	-	4	3	2	2	2	1	1	-
American Indian 1/, both sexes	2	-	2	-	1	4	3	5	1	-	-	-
Male	1	-	1	-	1	1	2	2	1	-	-	-
Female	1	-	1	-	-	3	1	3	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	4	4	6	2	5	11	11	9	-	-	-
Male	2	3	1	4	1	1	2	4	1	-	-	-
Female	-	1	3	2	1	4	9	7	8	-	-	-
West Virginia												
All races, both sexes	36	53	81	103	117	195	271	272	233	90	22	-
Male	25	33	57	60	58	97	119	104	62	11	3	-
Female	11	20	24	43	59	98	152	168	171	79	19	-
White, both sexes	35	48	77	100	115	189	264	264	229	87	21	-
Male	24	29	54	58	57	94	115	103	61	11	3	-
Female	11	19	23	42	58	95	149	161	168	76	18	-
Black, both sexes	1	5	3	3	2	6	7	8	4	3	1	-
Male	1	4	2	2	1	3	4	1	1	-	-	-
Female	-	1	1	1	1	3	3	7	3	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Wisconsin												
All races, both sexes	76	106	98	152	218	344	644	800	718	313	74	-
Male	50	68	64	87	123	183	292	319	211	67	6	-
Female	26	38	34	65	95	161	352	481	507	246	68	-
White, both sexes	62	86	87	139	196	329	626	787	708	309	73	-
Male	42	58	59	83	110	177	287	314	208	67	6	-
Female	20	28	28	56	86	152	339	473	500	242	67	-
Black, both sexes	14	19	10	9	18	10	15	9	9	2	1	-
Male	8	9	4	3	10	3	5	4	3	-	-	-
Female	6	10	6	6	8	7	10	5	6	2	1	-
American Indian 1/, both sexes	-	-	-	3	2	3	1	-	1	1	-	-
Male	-	-	-	1	2	1	-	-	-	-	-	-
Female	-	-	-	2	-	2	1	-	1	1	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	1	2	2	2	4	-	1	-	-
Male	-	1	1	-	1	2	-	1	-	-	-	-
Female	-	-	-	1	1	-	2	3	-	1	-	-
Wyoming												
All races, both sexes	11	11	16	15	28	31	53	64	57	25	1	-
Male	5	8	10	9	14	16	28	18	17	8	-	-
Female	6	3	6	6	14	15	25	46	40	17	1	-
White, both sexes	11	11	15	15	27	31	53	63	56	24	1	-
Male	5	8	9	9	14	16	28	18	17	8	-	-
Female	6	3	6	6	13	15	25	45	39	16	1	-
Black, both sexes	-	-	-	-	1	-	-	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	1	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
United States												
All races, both sexes	12,552	100	26	4	13	40	68	146	149	266	355	534
Male	4,868	61	8	3	3	13	21	53	59	123	153	231
Female	7,684	39	18	1	10	27	47	93	90	143	202	303
White, both sexes	10,284	61	17	4	8	32	50	89	91	177	249	367
Male	3,992	43	2	3	3	11	15	33	35	85	103	164
Female	6,292	18	15	1	5	21	35	56	56	92	146	203
Black, both sexes	2,057	30	5	-	4	6	17	50	50	79	100	161
Male	791	13	4	-	-	1	5	17	21	34	47	65
Female	1,266	17	1	-	4	5	12	33	29	45	53	96
American Indian 1/, both sexes	57	4	1	-	-	2	-	5	5	2	3	2
Male	24	3	1	-	-	1	-	2	2	-	-	1
Female	33	1	-	-	-	1	-	3	3	2	3	1
Asian or Pacific Islander 2/, both sexes	154	5	3	-	1	-	1	2	3	8	3	4
Male	61	2	1	-	-	-	1	1	1	4	3	1
Female	93	3	2	-	1	-	-	1	2	4	-	3
Alabama												
All races, both sexes	238	2	1	-	-	-	1	5	2	8	8	6
Male	104	-	1	-	-	-	1	-	1	2	5	4
Female	134	2	-	-	-	-	-	5	1	6	3	2
White, both sexes	153	-	-	-	-	-	-	-	1	2	2	3
Male	61	-	-	-	-	-	-	-	-	-	1	1
Female	92	-	-	-	-	-	-	-	1	2	1	2
Black, both sexes	83	2	1	-	-	-	1	5	1	6	6	3
Male	43	-	1	-	-	-	1	-	1	2	4	3
Female	40	2	-	-	-	-	-	5	-	4	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Alaska												
All races, both sexes	24	-	1	-	-	1	-	1	2	-	1	1
Male	9	-	1	-	-	1	-	-	-	-	-	1
Female	15	-	-	-	-	-	-	1	2	-	1	-
White, both sexes	18	-	-	-	-	1	-	-	-	-	1	1
Male	7	-	-	-	-	1	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	1	-	-	-	-	1	2	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	181	3	-	-	-	1	-	5	4	2	13	2
Male	77	1	-	-	-	1	-	2	2	1	7	2
Female	104	2	-	-	-	-	-	3	2	1	6	-
White, both sexes	154	1	-	-	-	-	-	4	1	2	10	2
Male	64	-	-	-	-	-	-	2	-	1	6	2
Female	90	1	-	-	-	-	-	2	1	1	4	-
Black, both sexes	13	-	-	-	-	-	-	-	-	-	2	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	11	2	-	-	-	1	-	1	3	-	1	-
Male	5	1	-	-	-	1	-	-	2	-	-	-
Female	6	1	-	-	-	-	-	1	1	-	1	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Arkansas												
All races, both sexes	198	2	1	-	-	-	-	3	3	1	5	6
Male	79	2	-	-	-	-	-	1	2	1	4	2
Female	119	-	1	-	-	-	-	2	1	-	1	4
White, both sexes	170	-	1	-	-	-	-	2	2	1	3	6
Male	65	-	-	-	-	-	-	1	1	1	3	2
Female	105	-	1	-	-	-	-	1	1	-	-	4
Black, both sexes	28	2	-	-	-	-	-	1	1	-	2	-
Male	14	2	-	-	-	-	-	-	1	-	1	-
Female	14	-	-	-	-	-	-	1	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	800	5	7	1	2	6	2	10	18	19	23	40
Male	303	4	2	1	1	2	1	4	5	9	10	12
Female	497	1	5	-	1	4	1	6	13	10	13	28
White, both sexes	657	4	6	1	2	6	1	6	13	13	20	27
Male	244	3	1	1	1	2	1	3	3	6	8	6
Female	413	1	5	-	1	4	-	3	10	7	12	21
Black, both sexes	97	-	1	-	-	-	1	4	3	1	3	13
Male	40	-	1	-	-	-	-	1	1	1	2	6
Female	57	-	-	-	-	-	1	3	2	-	1	7
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	43	1	-	-	-	-	-	-	2	5	-	-
Male	16	1	-	-	-	-	-	-	1	2	-	-
Female	27	-	-	-	-	-	-	-	1	3	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Colorado												
All races, both sexes	206	1	1	-	-	-	1	-	1	2	4	5
Male	93	1	-	-	-	-	-	-	-	1	-	3
Female	113	-	1	-	-	-	1	-	1	1	4	2
White, both sexes	192	1	1	-	-	-	1	-	1	2	3	5
Male	87	1	-	-	-	-	-	-	-	1	-	3
Female	105	-	1	-	-	-	1	-	1	1	3	2
Black, both sexes	13	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	147	1	-	-	-	-	-	-	5	3	2	3
Male	51	1	-	-	-	-	-	-	2	1	-	1
Female	96	-	-	-	-	-	-	-	3	2	2	2
White, both sexes	128	1	-	-	-	-	-	-	3	3	1	2
Male	45	1	-	-	-	-	-	-	2	1	-	-
Female	83	-	-	-	-	-	-	-	1	2	1	2
Black, both sexes	17	-	-	-	-	-	-	-	2	-	1	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	2	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Delaware												
All races, both sexes	50	-	1	-	-	-	-	-	-	2	1	-
Male	22	-	-	-	-	-	-	-	-	1	-	-
Female	28	-	1	-	-	-	-	-	-	1	1	-
White, both sexes	40	-	1	-	-	-	-	-	-	2	1	-
Male	17	-	-	-	-	-	-	-	-	1	-	-
Female	23	-	1	-	-	-	-	-	-	1	1	-
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	24	-	-	-	-	-	2	-	-	-	-	2
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	2	-	-	-	-	2
White, both sexes	4	-	-	-	-	-	1	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	1	-	-	-	-	1
Black, both sexes	20	-	-	-	-	-	1	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	1	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Florida												
All races, both sexes	637	8	-	1	-	1	4	7	4	7	13	27
Male	250	2	-	-	-	1	-	-	2	4	5	10
Female	387	6	-	1	-	-	4	7	2	3	8	17
White, both sexes	519	4	-	1	-	1	3	5	1	4	9	15
Male	218	2	-	-	-	1	-	-	1	2	3	6
Female	301	2	-	1	-	-	3	5	-	2	6	9
Black, both sexes	115	3	-	-	-	-	1	2	3	3	4	12
Male	30	-	-	-	-	-	-	-	1	2	2	4
Female	85	3	-	-	-	-	1	2	2	1	2	8
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	1	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	430	3	1	-	-	2	4	9	6	10	15	25
Male	171	1	-	-	-	-	-	6	5	2	7	9
Female	259	2	1	-	-	2	4	3	1	8	8	16
White, both sexes	285	-	-	-	-	1	3	4	3	6	5	9
Male	116	-	-	-	-	-	-	3	3	2	3	3
Female	169	-	-	-	-	1	3	1	-	4	2	6
Black, both sexes	143	3	1	-	-	1	1	5	3	4	9	16
Male	55	1	-	-	-	-	-	3	2	-	4	6
Female	88	2	1	-	-	1	1	2	1	4	5	10
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Hawaii												
All races, both sexes	37	1	1	-	-	-	-	-	1	1	1	3
Male	17	1	1	-	-	-	-	-	-	-	1	1
Female	20	-	-	-	-	-	-	-	1	1	-	2
White, both sexes	15	-	1	-	-	-	-	-	-	-	-	1
Male	8	-	1	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	22	1	-	-	-	-	-	-	1	1	1	2
Male	9	1	-	-	-	-	-	-	-	-	1	-
Female	13	-	-	-	-	-	-	-	1	1	-	2
Idaho												
All races, both sexes	81	-	-	-	-	-	-	1	-	3	2	8
Male	29	-	-	-	-	-	-	1	-	1	1	1
Female	52	-	-	-	-	-	-	-	-	2	1	7
White, both sexes	79	-	-	-	-	-	-	1	-	3	2	8
Male	28	-	-	-	-	-	-	1	-	1	1	1
Female	51	-	-	-	-	-	-	-	-	2	1	7
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Illinois												
All races, both sexes	594	3	1	-	-	2	3	9	2	13	19	24
Male	219	-	-	-	-	1	2	4	-	7	5	15
Female	375	3	1	-	-	1	1	5	2	6	14	9
White, both sexes	470	2	-	-	-	1	2	6	1	8	12	17
Male	170	-	-	-	-	1	1	3	-	5	3	10
Female	300	2	-	-	-	-	1	3	1	3	9	7
Black, both sexes	115	-	-	-	-	1	-	2	1	5	7	7
Male	47	-	-	-	-	-	-	1	-	2	2	5
Female	68	-	-	-	-	1	-	1	1	3	5	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	1	1	-	-	-	1	1	-	-	-	-
Male	2	-	-	-	-	-	1	-	-	-	-	-
Female	7	1	1	-	-	-	-	1	-	-	-	-
Indiana												
All races, both sexes	405	2	2	-	2	1	1	4	5	11	15	20
Male	172	2	-	-	1	-	1	1	3	6	12	11
Female	233	-	2	-	1	1	-	3	2	5	3	9
White, both sexes	364	2	2	-	2	1	1	2	5	7	12	19
Male	153	2	-	-	1	-	1	-	3	3	10	11
Female	211	-	2	-	1	1	-	2	2	4	2	8
Black, both sexes	41	-	-	-	-	-	-	2	-	4	3	1
Male	19	-	-	-	-	-	-	1	-	3	2	-
Female	22	-	-	-	-	-	-	1	-	1	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Iowa												
All races, both sexes	152	1	-	-	-	-	2	-	-	3	2	4
Male	51	1	-	-	-	-	-	-	-	1	2	2
Female	101	-	-	-	-	-	2	-	-	2	-	2
White, both sexes	148	1	-	-	-	-	2	-	-	3	2	4
Male	50	1	-	-	-	-	-	-	-	1	2	2
Female	98	-	-	-	-	-	2	-	-	2	-	2
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	170	2	-	-	-	3	2	1	1	4	6	8
Male	66	1	-	-	-	1	1	-	-	1	3	5
Female	104	1	-	-	-	2	1	1	1	3	3	3
White, both sexes	162	1	-	-	-	2	2	-	1	3	6	8
Male	65	-	-	-	-	1	1	-	-	1	3	5
Female	97	1	-	-	-	1	1	-	1	2	3	3
Black, both sexes	5	-	-	-	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	1	-	1	-	-
American Indian 1/, both sexes	2	1	-	-	-	1	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Kentucky												
All races, both sexes	222	-	-	-	-	1	-	1	1	2	9	10
Male	94	-	-	-	-	-	-	-	-	1	2	4
Female	128	-	-	-	-	1	-	1	1	1	7	6
White, both sexes	206	-	-	-	-	1	-	1	1	2	6	9
Male	85	-	-	-	-	-	-	-	-	1	1	3
Female	121	-	-	-	-	1	-	1	1	1	5	6
Black, both sexes	16	-	-	-	-	-	-	-	-	-	3	1
Male	9	-	-	-	-	-	-	-	-	-	1	1
Female	7	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	244	3	-	-	2	1	-	2	4	10	4	14
Male	92	1	-	-	1	-	-	-	1	3	4	6
Female	152	2	-	-	1	1	-	2	3	7	-	8
White, both sexes	151	2	-	-	1	1	-	2	2	3	2	5
Male	56	1	-	-	1	-	-	-	-	-	2	2
Female	95	1	-	-	-	1	-	2	2	3	-	3
Black, both sexes	92	1	-	-	1	-	-	-	2	7	2	9
Male	35	-	-	-	-	-	-	-	1	3	2	4
Female	57	1	-	-	1	-	-	-	1	4	-	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Maine												
All races, both sexes	75	-	-	-	-	-	1	-	1	1	1	6
Male	31	-	-	-	-	-	-	-	-	1	1	4
Female	44	-	-	-	-	-	1	-	1	-	-	2
White, both sexes	75	-	-	-	-	-	1	-	1	1	1	6
Male	31	-	-	-	-	-	-	-	-	1	1	4
Female	44	-	-	-	-	-	1	-	1	-	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	242	2	-	-	-	-	-	5	2	6	7	14
Male	88	1	-	-	-	-	-	3	1	4	3	11
Female	154	1	-	-	-	-	-	2	1	2	4	3
White, both sexes	155	1	-	-	-	-	-	2	1	3	2	7
Male	60	1	-	-	-	-	-	1	1	2	1	6
Female	95	-	-	-	-	-	-	1	-	1	1	1
Black, both sexes	80	1	-	-	-	-	-	2	1	3	5	7
Male	24	-	-	-	-	-	-	1	-	2	2	5
Female	56	1	-	-	-	-	-	1	1	1	3	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	1	-	-	-	-
Male	4	-	-	-	-	-	-	1	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Massachusetts												
All races, both sexes	232	1	-	-	1	-	-	2	-	3	2	5
Male	84	-	-	-	-	-	-	1	-	2	-	1
Female	148	1	-	-	1	-	-	1	-	1	2	4
White, both sexes	211	-	-	-	1	-	-	2	-	2	2	4
Male	73	-	-	-	-	-	-	1	-	1	-	1
Female	138	-	-	-	1	-	-	1	-	1	2	3
Black, both sexes	16	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	1	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	3	1	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	398	3	1	-	-	-	3	6	5	7	7	13
Male	148	1	1	-	-	-	-	4	3	4	2	4
Female	250	2	-	-	-	-	3	2	2	3	5	9
White, both sexes	307	1	-	-	-	-	2	5	-	4	3	10
Male	117	-	-	-	-	-	-	3	-	3	1	3
Female	190	1	-	-	-	-	2	2	-	1	2	7
Black, both sexes	86	2	-	-	-	-	1	1	5	2	4	3
Male	30	1	-	-	-	-	-	1	3	1	1	1
Female	56	1	-	-	-	-	1	-	2	1	3	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	3	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Minnesota												
All races, both sexes	246	1	-	-	-	1	3	1	4	5	8	7
Male	89	1	-	-	-	-	-	-	2	1	3	1
Female	157	-	-	-	-	1	3	1	2	4	5	6
White, both sexes	236	1	-	-	-	1	3	1	3	5	8	6
Male	86	1	-	-	-	-	-	-	2	1	3	1
Female	150	-	-	-	-	1	3	1	1	4	5	5
Black, both sexes	8	-	-	-	-	-	-	-	1	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	173	3	-	-	-	3	4	1	1	3	6	7
Male	64	2	-	-	-	-	1	-	-	2	2	5
Female	109	1	-	-	-	3	3	1	1	1	4	2
White, both sexes	100	1	-	-	-	2	2	-	-	1	3	3
Male	37	1	-	-	-	-	1	-	-	-	1	3
Female	63	-	-	-	-	2	1	-	-	1	2	-
Black, both sexes	73	2	-	-	-	1	2	1	1	2	3	4
Male	27	1	-	-	-	-	-	-	-	2	1	2
Female	46	1	-	-	-	1	2	1	1	-	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Missouri												
All races, both sexes	303	-	-	1	-	1	1	3	5	10	11	18
Male	122	-	-	1	-	1	1	1	2	2	1	13
Female	181	-	-	-	-	-	-	2	3	8	10	5
White, both sexes	255	-	-	1	-	-	1	2	3	7	9	15
Male	102	-	-	1	-	-	1	1	1	2	-	10
Female	153	-	-	-	-	-	-	1	2	5	9	5
Black, both sexes	46	-	-	-	-	1	-	1	2	3	2	3
Male	20	-	-	-	-	1	-	-	1	-	1	3
Female	26	-	-	-	-	-	-	1	1	3	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	41	-	-	-	-	-	-	1	-	1	1	5
Male	17	-	-	-	-	-	-	-	-	1	-	2
Female	24	-	-	-	-	-	-	1	-	-	1	3
White, both sexes	41	-	-	-	-	-	-	1	-	1	1	5
Male	17	-	-	-	-	-	-	-	-	1	-	2
Female	24	-	-	-	-	-	-	1	-	-	1	3
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Nebraska												
All races, both sexes	113	-	-	-	-	-	-	1	2	3	3	1
Male	41	-	-	-	-	-	-	1	-	3	2	1
Female	72	-	-	-	-	-	-	-	2	-	1	-
White, both sexes	107	-	-	-	-	-	-	-	2	3	3	1
Male	38	-	-	-	-	-	-	-	-	3	2	1
Female	69	-	-	-	-	-	-	-	2	-	1	-
Black, both sexes	5	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	85	1	-	-	1	-	-	4	1	5	4	3
Male	37	1	-	-	-	-	-	1	-	3	2	-
Female	48	-	-	-	1	-	-	3	1	2	2	3
White, both sexes	61	-	-	-	-	-	-	2	1	3	2	2
Male	24	-	-	-	-	-	-	-	-	2	1	-
Female	37	-	-	-	-	-	-	2	1	1	1	2
Black, both sexes	20	1	-	-	-	-	-	2	-	2	1	1
Male	10	1	-	-	-	-	-	1	-	1	-	-
Female	10	-	-	-	-	-	-	1	-	1	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	1	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
New Hampshire												
All races, both sexes	55	-	-	-	-	-	-	-	-	-	1	-
Male	23	-	-	-	-	-	-	-	-	-	1	-
Female	32	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	53	-	-	-	-	-	-	-	-	-	1	-
Male	22	-	-	-	-	-	-	-	-	-	1	-
Female	31	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	359	4	-	-	-	1	3	4	3	9	16	15
Male	136	4	-	-	-	-	-	1	2	5	8	5
Female	223	-	-	-	-	1	3	3	1	4	8	10
White, both sexes	285	2	-	-	-	-	2	1	2	6	10	7
Male	101	2	-	-	-	-	-	-	2	4	5	2
Female	184	-	-	-	-	-	2	1	-	2	5	5
Black, both sexes	66	2	-	-	-	1	1	3	1	3	6	8
Male	30	2	-	-	-	-	-	1	-	1	3	3
Female	36	-	-	-	-	1	1	2	1	2	3	5
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
New Mexico												
All races, both sexes	84	1	-	-	-	-	-	2	-	4	1	3
Male	34	1	-	-	-	-	-	1	-	2	-	3
Female	50	-	-	-	-	-	-	1	-	2	1	-
White, both sexes	79	1	-	-	-	-	-	1	-	4	1	2
Male	32	1	-	-	-	-	-	1	-	2	-	2
Female	47	-	-	-	-	-	-	-	-	2	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	1	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	544	13	1	-	-	1	2	3	2	13	12	16
Male	218	8	-	-	-	1	1	3	2	10	5	7
Female	326	5	1	-	-	-	1	-	-	3	7	9
White, both sexes	456	8	-	-	-	1	2	1	1	10	10	10
Male	191	6	-	-	-	1	1	1	1	8	4	4
Female	265	2	-	-	-	-	1	-	-	2	6	6
Black, both sexes	79	5	-	-	-	-	-	2	1	3	2	6
Male	25	2	-	-	-	-	-	2	1	2	1	3
Female	54	3	-	-	-	-	-	-	-	1	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	1	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
North Carolina												
All races, both sexes	477	5	3	-	1	-	4	5	8	15	8	23
Male	180	4	2	-	-	-	1	2	2	6	3	7
Female	297	1	1	-	1	-	3	3	6	9	5	16
White, both sexes	330	3	1	-	-	-	3	2	3	5	3	8
Male	127	3	-	-	-	-	-	1	2	2	-	4
Female	203	-	1	-	-	-	3	1	1	3	3	4
Black, both sexes	143	2	2	-	1	-	1	3	5	9	5	15
Male	53	1	2	-	-	-	1	1	-	4	3	3
Female	90	1	-	-	1	-	-	2	5	5	2	12
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	29	1	-	-	-	-	-	1	-	-	1	1
Male	12	1	-	-	-	-	-	1	-	-	-	1
Female	17	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	29	1	-	-	-	-	-	1	-	-	1	1
Male	12	1	-	-	-	-	-	1	-	-	-	1
Female	17	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Ohio												
All races, both sexes	455	1	-	-	-	2	1	5	10	9	14	20
Male	150	-	-	-	-	-	-	1	2	6	6	6
Female	305	1	-	-	-	2	1	4	8	3	8	14
White, both sexes	390	-	-	-	-	2	1	3	8	5	11	16
Male	132	-	-	-	-	-	-	-	2	3	4	6
Female	258	-	-	-	-	2	1	3	6	2	7	10
Black, both sexes	63	1	-	-	-	-	-	2	2	3	3	4
Male	16	-	-	-	-	-	-	1	-	2	2	-
Female	47	1	-	-	-	-	-	1	2	1	1	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	161	1	-	-	-	1	-	6	1	4	5	8
Male	57	1	-	-	-	-	-	2	-	1	2	3
Female	104	-	-	-	-	1	-	4	1	3	3	5
White, both sexes	138	-	-	-	-	1	-	5	1	3	4	5
Male	49	-	-	-	-	-	-	1	-	1	2	3
Female	89	-	-	-	-	1	-	4	1	2	2	2
Black, both sexes	11	-	-	-	-	-	-	-	-	1	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	1	-	2
American Indian 1/, both sexes	12	1	-	-	-	-	-	1	-	-	1	1
Male	4	1	-	-	-	-	-	1	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Oregon												
All races, both sexes	167	2	-	-	-	1	3	1	-	2	2	8
Male	66	1	-	-	-	-	2	1	-	-	1	3
Female	101	1	-	-	-	1	1	-	-	2	1	5
White, both sexes	163	2	-	-	-	1	3	-	-	2	2	7
Male	63	1	-	-	-	-	2	-	-	-	1	2
Female	100	1	-	-	-	1	1	-	-	2	1	5
Black, both sexes	3	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	678	2	2	-	-	-	4	8	11	11	23	31
Male	270	1	-	-	-	-	2	2	6	6	11	15
Female	408	1	2	-	-	-	2	6	5	5	12	16
White, both sexes	587	2	2	-	-	-	3	5	8	8	16	23
Male	228	1	-	-	-	-	2	2	3	5	7	12
Female	359	1	2	-	-	-	1	3	5	3	9	11
Black, both sexes	90	-	-	-	-	-	1	3	3	3	7	8
Male	42	-	-	-	-	-	-	-	3	1	4	3
Female	48	-	-	-	-	-	1	3	-	2	3	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Rhode Island												
All races, both sexes	50	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	38	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	49	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	37	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	254	3	1	-	-	-	1	4	7	8	7	13
Male	94	3	-	-	-	-	-	-	3	4	3	5
Female	160	-	1	-	-	-	1	4	4	4	4	8
White, both sexes	167	1	1	-	-	-	-	3	2	1	3	7
Male	56	1	-	-	-	-	-	-	-	1	2	2
Female	111	-	1	-	-	-	-	3	2	-	1	5
Black, both sexes	84	2	-	-	-	-	1	1	5	7	4	6
Male	36	2	-	-	-	-	-	-	3	3	1	3
Female	48	-	-	-	-	-	1	1	2	4	3	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
South Dakota												
All races, both sexes	32	-	-	-	-	-	-	-	-	1	-	2
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	19	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	32	-	-	-	-	-	-	-	-	1	-	2
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	19	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	315	2	-	-	-	1	6	2	7	6	10	17
Male	124	1	-	-	-	1	4	-	3	3	3	5
Female	191	1	-	-	-	-	2	2	4	3	7	12
White, both sexes	252	2	-	-	-	1	3	2	5	4	8	10
Male	101	1	-	-	-	1	2	-	2	3	3	2
Female	151	1	-	-	-	-	1	2	3	1	5	8
Black, both sexes	62	-	-	-	-	-	3	-	2	2	2	7
Male	22	-	-	-	-	-	2	-	1	-	-	3
Female	40	-	-	-	-	-	1	-	1	2	2	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Texas												
All races, both sexes	952	7	1	-	2	2	6	8	9	24	27	37
Male	374	5	-	-	-	-	2	1	3	10	14	17
Female	578	2	1	-	2	2	4	7	6	14	13	20
White, both sexes	764	7	1	-	2	1	4	5	6	20	21	30
Male	304	5	-	-	-	-	1	-	2	9	9	15
Female	460	2	1	-	2	1	3	5	4	11	12	15
Black, both sexes	177	-	-	-	-	1	2	3	3	4	5	7
Male	67	-	-	-	-	-	1	1	1	1	4	2
Female	110	-	-	-	-	1	1	2	2	3	1	5
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	121	-	-	-	-	1	2	2	2	2	4	5
Male	43	-	-	-	-	1	1	1	-	-	1	1
Female	78	-	-	-	-	-	1	1	2	2	3	4
White, both sexes	116	-	-	-	-	1	2	2	2	2	3	4
Male	42	-	-	-	-	1	1	1	-	-	1	1
Female	74	-	-	-	-	-	1	1	2	2	2	3
Black, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Vermont												
All races, both sexes	28	1	-	-	-	-	-	-	-	-	1	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	16	1	-	-	-	-	-	-	-	-	1	-
White, both sexes	28	1	-	-	-	-	-	-	-	-	1	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	16	1	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	343	5	-	-	2	2	1	3	3	4	8	15
Male	135	4	-	-	-	-	-	2	1	2	1	5
Female	208	1	-	-	2	2	1	1	2	2	7	10
White, both sexes	253	4	-	-	-	2	1	2	2	3	6	6
Male	101	4	-	-	-	-	-	2	1	1	-	4
Female	152	-	-	-	-	2	1	-	1	2	6	2
Black, both sexes	89	1	-	-	2	-	-	1	1	1	2	9
Male	33	-	-	-	-	-	-	-	-	1	1	1
Female	56	1	-	-	2	-	-	1	1	-	1	8
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Washington												
All races, both sexes	250	1	-	1	-	2	-	4	2	3	10	8
Male	89	1	-	1	-	1	-	2	-	-	4	3
Female	161	-	-	-	-	1	-	2	2	3	6	5
White, both sexes	233	1	-	1	-	2	-	3	2	3	7	6
Male	85	1	-	1	-	1	-	2	-	-	3	2
Female	148	-	-	-	-	1	-	1	2	3	4	4
Black, both sexes	11	-	-	-	-	-	-	1	-	-	3	2
Male	3	-	-	-	-	-	-	-	-	-	1	1
Female	8	-	-	-	-	-	-	1	-	-	2	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	109	1	-	-	-	2	1	1	1	2	2	9
Male	50	-	-	-	-	1	-	-	1	2	2	4
Female	59	1	-	-	-	1	1	1	-	-	-	5
White, both sexes	103	1	-	-	-	2	1	1	1	2	1	9
Male	46	-	-	-	-	1	-	-	1	2	1	4
Female	57	1	-	-	-	1	1	1	-	-	-	5
Black, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Wisconsin												
All races, both sexes	305	1	-	-	-	-	-	4	2	4	8	14
Male	134	1	-	-	-	-	-	2	2	1	4	8
Female	171	-	-	-	-	-	-	2	-	3	4	6
White, both sexes	280	1	-	-	-	-	-	4	1	4	7	11
Male	124	1	-	-	-	-	-	2	1	1	4	6
Female	156	-	-	-	-	-	-	2	-	3	3	5
Black, both sexes	21	-	-	-	-	-	-	-	1	-	1	3
Male	10	-	-	-	-	-	-	-	1	-	-	2
Female	11	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	36	1	-	-	-	-	-	1	1	-	2	1
Male	15	1	-	-	-	-	-	1	1	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	34	1	-	-	-	-	-	-	1	-	2	1
Male	13	1	-	-	-	-	-	-	1	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
United States												
All races, both sexes	674	795	919	1,037	1,215	1,509	1,839	1,590	940	291	42	-
Male	322	387	422	448	517	615	645	474	239	62	9	-
Female	352	408	497	589	698	894	1,194	1,116	701	229	33	-
White, both sexes	499	607	753	821	1,000	1,273	1,611	1,427	852	267	29	-
Male	238	307	352	356	437	535	574	419	214	59	4	-
Female	261	300	401	465	563	738	1,037	1,008	638	208	25	-
Black, both sexes	162	170	155	197	195	212	206	143	80	22	13	-
Male	76	71	65	82	72	73	66	48	23	3	5	-
Female	86	99	90	115	123	139	140	95	57	19	8	-
American Indian 1/, both sexes	6	5	3	4	3	7	2	2	1	-	-	-
Male	4	2	1	2	1	2	1	1	-	-	-	-
Female	2	3	2	2	2	5	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	7	13	8	15	17	17	20	18	7	2	-	-
Male	4	7	4	8	7	5	4	6	2	-	-	-
Female	3	6	4	7	10	12	16	12	5	2	-	-
Alabama												
All races, both sexes	13	15	15	21	29	35	35	28	12	2	-	-
Male	7	10	9	12	13	13	11	12	3	-	-	-
Female	6	5	6	9	16	22	24	16	9	2	-	-
White, both sexes	8	11	13	14	18	24	27	21	7	2	-	-
Male	3	6	8	9	8	9	7	8	1	-	-	-
Female	5	5	5	5	10	15	20	13	6	2	-	-
Black, both sexes	5	4	2	6	10	11	8	7	5	-	-	-
Male	4	4	1	3	5	4	4	4	2	-	-	-
Female	1	-	1	3	5	7	4	3	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Alaska												
All races, both sexes	2	2	-	-	4	4	2	-	3	-	-	-
Male	-	-	-	-	4	2	-	-	-	-	-	-
Female	2	2	-	-	-	2	2	-	3	-	-	-
White, both sexes	2	2	-	-	4	3	1	-	3	-	-	-
Male	-	-	-	-	4	1	-	-	-	-	-	-
Female	2	2	-	-	-	2	1	-	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Arizona												
All races, both sexes	6	11	14	17	24	27	19	15	13	5	-	-
Male	3	8	6	8	7	12	5	5	4	3	-	-
Female	3	3	8	9	17	15	14	10	9	2	-	-
White, both sexes	2	7	11	17	22	24	18	15	13	5	-	-
Male	1	5	5	8	6	11	5	5	4	3	-	-
Female	1	2	6	9	16	13	13	10	9	2	-	-
Black, both sexes	3	3	1	-	1	2	1	-	-	-	-	-
Male	1	2	1	-	1	1	-	-	-	-	-	-
Female	2	1	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	1	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Arkansas												
All races, both sexes	12	12	14	23	22	24	29	20	16	3	2	-
Male	4	8	5	9	9	10	10	5	6	1	-	-
Female	8	4	9	14	13	14	19	15	10	2	2	-
White, both sexes	10	11	12	16	22	19	25	20	15	3	2	-
Male	3	7	5	5	9	9	8	5	5	1	-	-
Female	7	4	7	11	13	10	17	15	10	2	2	-
Black, both sexes	2	1	2	7	-	5	4	-	1	-	-	-
Male	1	1	-	4	-	1	2	-	1	-	-	-
Female	1	-	2	3	-	4	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	55	55	51	51	69	91	112	95	63	24	1	-
Male	22	23	20	23	29	37	44	33	16	4	1	-
Female	33	32	31	28	40	54	68	62	47	20	-	-
White, both sexes	45	42	39	35	56	80	98	85	56	22	-	-
Male	18	18	14	14	24	33	40	30	14	4	-	-
Female	27	24	25	21	32	47	58	55	42	18	-	-
Black, both sexes	8	11	9	8	7	9	7	5	5	1	1	-
Male	4	5	4	4	2	3	2	1	2	-	1	-
Female	4	6	5	4	5	6	5	4	3	1	-	-
American Indian 1/, both sexes	-	-	-	2	1	-	-	-	-	-	-	-
Male	-	-	-	2	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	3	6	5	2	7	5	2	1	-	-
Male	-	-	2	3	2	1	2	2	-	-	-	-
Female	2	2	1	3	3	1	5	3	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Colorado												
All races, both sexes	15	14	14	17	21	34	29	26	17	3	1	-
Male	10	4	8	9	12	17	11	10	6	1	-	-
Female	5	10	6	8	9	17	18	16	11	2	1	-
White, both sexes	14	12	14	16	17	31	28	26	16	3	1	-
Male	9	4	8	9	9	16	11	10	5	1	-	-
Female	5	8	6	7	8	15	17	16	11	2	1	-
Black, both sexes	1	2	-	1	4	2	1	-	1	-	-	-
Male	1	-	-	-	3	-	-	-	1	-	-	-
Female	-	2	-	1	1	2	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	3	8	9	12	11	24	22	23	10	10	1	-
Male	2	5	3	6	5	5	8	7	2	2	1	-
Female	1	3	6	6	6	19	14	16	8	8	-	-
White, both sexes	2	5	7	9	10	21	22	23	9	9	1	-
Male	1	4	2	4	5	5	8	7	2	2	1	-
Female	1	1	5	5	5	16	14	16	7	7	-	-
Black, both sexes	1	3	2	2	1	2	-	-	1	1	-	-
Male	1	1	1	1	-	-	-	-	-	-	-	-
Female	-	2	1	1	1	2	-	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Delaware												
All races, both sexes	5	3	4	1	7	3	5	8	6	3	1	-
Male	3	2	-	1	5	1	3	2	2	1	1	-
Female	2	1	4	-	2	2	2	6	4	2	-	-
White, both sexes	3	3	1	1	5	2	4	8	6	3	-	-
Male	2	2	-	1	4	-	2	2	2	1	-	-
Female	1	1	1	-	1	2	2	6	4	2	-	-
Black, both sexes	2	-	3	-	2	1	1	-	-	-	1	-
Male	1	-	-	-	1	1	1	-	-	-	1	-
Female	1	-	3	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	3	3	2	1	5	5	-	-	-	-
Male	1	-	2	1	-	-	1	2	-	-	-	-
Female	-	-	1	2	2	1	4	3	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	3	3	2	1	5	4	-	-	-	-
Male	-	-	2	1	-	-	1	1	-	-	-	-
Female	-	-	1	2	2	1	4	3	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Florida												
All races, both sexes	28	36	31	49	70	95	109	100	31	14	2	-
Male	7	17	16	26	26	38	48	32	14	1	1	-
Female	21	19	15	23	44	57	61	68	17	13	1	-
White, both sexes	18	24	25	40	62	80	96	90	26	13	2	-
Male	4	12	14	19	26	37	46	30	13	1	1	-
Female	14	12	11	21	36	43	50	60	13	12	1	-
Black, both sexes	10	11	6	8	8	15	13	10	5	1	-	-
Male	3	4	2	6	-	1	2	2	1	-	-	-
Female	7	7	4	2	8	14	11	8	4	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	30	33	33	35	37	53	58	44	26	4	2	-
Male	15	15	17	16	15	26	21	11	4	1	-	-
Female	15	18	16	19	22	27	37	33	22	3	2	-
White, both sexes	16	24	22	26	27	35	45	35	21	3	-	-
Male	7	14	12	13	12	14	17	8	4	1	-	-
Female	9	10	10	13	15	21	28	27	17	2	-	-
Black, both sexes	14	8	11	9	10	18	13	9	5	1	2	-
Male	8	1	5	3	3	12	4	3	-	-	-	-
Female	6	7	6	6	7	6	9	6	5	1	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Hawaii												
All races, both sexes	1	1	4	2	4	4	5	4	3	1	-	-
Male	1	1	3	1	3	1	2	1	-	-	-	-
Female	-	-	1	1	1	3	3	3	3	1	-	-
White, both sexes	1	-	4	2	-	-	2	1	2	1	-	-
Male	1	-	3	1	-	-	1	-	-	-	-	-
Female	-	-	1	1	-	-	1	1	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	4	4	3	3	1	-	-	-
Male	-	1	-	-	3	1	1	1	-	-	-	-
Female	-	-	-	-	1	3	2	2	1	-	-	-
Idaho												
All races, both sexes	2	4	8	8	7	7	8	11	8	3	1	-
Male	-	4	1	3	2	4	4	2	4	1	-	-
Female	2	-	7	5	5	3	4	9	4	2	1	-
White, both sexes	2	4	8	8	6	7	7	11	8	3	1	-
Male	-	4	1	3	2	4	3	2	4	1	-	-
Female	2	-	7	5	4	3	4	9	4	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Illinois												
All races, both sexes	30	29	52	50	55	71	92	67	53	16	3	-
Male	13	14	23	21	26	24	28	24	8	3	1	-
Female	17	15	29	29	29	47	64	43	45	13	2	-
White, both sexes	24	17	44	37	42	57	75	56	51	16	2	-
Male	11	8	20	18	20	21	19	19	8	3	-	-
Female	13	9	24	19	22	36	56	37	43	13	2	-
Black, both sexes	6	12	7	13	13	13	15	10	2	-	1	-
Male	2	6	2	3	6	3	9	5	-	-	1	-
Female	4	6	5	10	7	10	6	5	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	2	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	2	1	-	-	-	-
Indiana												
All races, both sexes	23	35	46	39	36	46	57	34	19	6	1	-
Male	13	15	19	19	18	19	15	9	5	3	-	-
Female	10	20	27	20	18	27	42	25	14	3	1	-
White, both sexes	20	28	44	36	31	43	54	32	18	5	-	-
Male	10	13	19	17	14	18	15	9	4	3	-	-
Female	10	15	25	19	17	25	39	23	14	2	-	-
Black, both sexes	3	7	2	3	5	3	3	2	1	1	1	-
Male	3	2	-	2	4	1	-	-	1	-	-	-
Female	-	5	2	1	1	2	3	2	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Iowa												
All races, both sexes	8	15	6	12	16	17	29	16	14	7	-	-
Male	2	6	1	6	3	4	9	9	3	2	-	-
Female	6	9	5	6	13	13	20	7	11	5	-	-
White, both sexes	7	13	5	12	16	17	29	16	14	7	-	-
Male	2	5	1	6	3	4	9	9	3	2	-	-
Female	5	8	4	6	13	13	20	7	11	5	-	-
Black, both sexes	1	2	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	15	11	14	8	12	27	17	15	18	5	1	-
Male	10	5	8	3	5	8	4	6	5	-	-	-
Female	5	6	6	5	7	19	13	9	13	5	1	-
White, both sexes	15	11	14	7	12	24	17	15	18	5	1	-
Male	10	5	8	3	5	8	4	6	5	-	-	-
Female	5	6	6	4	7	16	13	9	13	5	1	-
Black, both sexes	-	-	-	1	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Kentucky												
All races, both sexes	13	19	20	18	27	28	25	32	12	3	1	-
Male	9	9	7	12	16	11	11	6	5	1	-	-
Female	4	10	13	6	11	17	14	26	7	2	1	-
White, both sexes	11	17	19	17	24	27	23	32	12	3	1	-
Male	7	8	7	11	14	11	10	6	5	1	-	-
Female	4	9	12	6	10	16	13	26	7	2	1	-
Black, both sexes	2	2	1	1	3	1	2	-	-	-	-	-
Male	2	1	-	1	2	-	1	-	-	-	-	-
Female	-	1	1	-	1	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	24	17	28	19	22	18	43	19	12	2	-	-
Male	14	9	9	8	8	5	13	6	4	-	-	-
Female	10	8	19	11	14	13	30	13	8	2	-	-
White, both sexes	14	12	16	9	14	9	32	15	11	1	-	-
Male	9	6	7	3	6	3	7	5	4	-	-	-
Female	5	6	9	6	8	6	25	10	7	1	-	-
Black, both sexes	10	5	12	10	8	9	11	3	1	1	-	-
Male	5	3	2	5	2	2	6	-	-	-	-	-
Female	5	2	10	5	6	7	5	3	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Maine												
All races, both sexes	1	4	2	8	6	20	6	12	4	2	-	-
Male	-	1	1	4	2	12	-	4	1	-	-	-
Female	1	3	1	4	4	8	6	8	3	2	-	-
White, both sexes	1	4	2	8	6	20	6	12	4	2	-	-
Male	-	1	1	4	2	12	-	4	1	-	-	-
Female	1	3	1	4	4	8	6	8	3	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	21	17	16	22	18	21	38	31	17	5	-	-
Male	11	10	8	8	3	8	5	8	3	1	-	-
Female	10	7	8	14	15	13	33	23	14	4	-	-
White, both sexes	12	10	13	13	12	13	23	26	14	3	-	-
Male	7	6	7	4	3	6	4	8	3	-	-	-
Female	5	4	6	9	9	7	19	18	11	3	-	-
Black, both sexes	8	6	3	8	6	7	14	4	3	2	-	-
Male	3	3	1	3	-	2	1	-	-	1	-	-
Female	5	3	2	5	6	5	13	4	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	1	-	1	1	1	-	-	-	-
Male	1	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Massachusetts												
All races, both sexes	8	16	13	16	15	29	54	40	22	3	2	-
Male	2	7	6	10	8	12	18	11	6	-	-	-
Female	6	9	7	6	7	17	36	29	16	3	2	-
White, both sexes	8	14	11	12	12	27	51	39	21	3	2	-
Male	2	6	5	7	6	12	17	10	5	-	-	-
Female	6	8	6	5	6	15	34	29	16	3	2	-
Black, both sexes	-	1	2	4	2	1	3	1	1	-	-	-
Male	-	-	1	3	2	-	1	1	1	-	-	-
Female	-	1	1	1	-	1	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	1	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Michigan												
All races, both sexes	23	18	29	37	41	57	51	56	27	12	2	-
Male	13	12	15	13	14	22	13	16	7	4	-	-
Female	10	6	14	24	27	35	38	40	20	8	2	-
White, both sexes	17	14	20	25	32	49	42	47	25	9	2	-
Male	11	10	10	12	11	18	12	13	6	4	-	-
Female	6	4	10	13	21	31	30	34	19	5	2	-
Black, both sexes	6	4	7	12	9	8	9	8	2	3	-	-
Male	2	2	5	1	3	4	1	3	1	-	-	-
Female	4	2	2	11	6	4	8	5	1	3	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Minnesota												
All races, both sexes	11	4	20	15	22	28	38	46	27	5	-	-
Male	6	-	10	6	11	10	15	14	9	-	-	-
Female	5	4	10	9	11	18	23	32	18	5	-	-
White, both sexes	8	4	19	14	22	27	37	45	27	5	-	-
Male	4	-	10	5	11	10	15	14	9	-	-	-
Female	4	4	9	9	11	17	22	31	18	5	-	-
Black, both sexes	1	-	1	1	-	1	1	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	14	4	18	13	22	26	19	15	13	1	-	-
Male	8	2	8	3	5	11	6	5	4	-	-	-
Female	6	2	10	10	17	15	13	10	9	1	-	-
White, both sexes	6	3	9	7	14	14	16	10	9	-	-	-
Male	2	2	4	2	3	7	5	3	3	-	-	-
Female	4	1	5	5	11	7	11	7	6	-	-	-
Black, both sexes	8	1	9	6	8	12	3	5	4	1	-	-
Male	6	-	4	1	2	4	1	2	1	-	-	-
Female	2	1	5	5	6	8	2	3	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Missouri												
All races, both sexes	21	17	32	28	30	32	36	26	19	11	1	-
Male	9	12	18	15	13	11	11	6	2	3	-	-
Female	12	5	14	13	17	21	25	20	17	8	1	-
White, both sexes	16	11	27	25	27	28	32	24	15	11	1	-
Male	6	10	15	14	11	10	10	6	1	3	-	-
Female	10	1	12	11	16	18	22	18	14	8	1	-
Black, both sexes	5	5	5	3	3	4	4	2	3	-	-	-
Male	3	2	3	1	2	1	1	-	1	-	-	-
Female	2	3	2	2	1	3	3	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	2	2	4	4	5	3	7	5	-	-	-
Male	1	-	-	1	3	4	2	2	1	-	-	-
Female	-	2	2	3	1	1	1	5	4	-	-	-
White, both sexes	1	2	2	4	4	5	3	7	5	-	-	-
Male	1	-	-	1	3	4	2	2	1	-	-	-
Female	-	2	2	3	1	1	1	5	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Nebraska												
All races, both sexes	3	4	5	4	10	12	22	17	22	3	1	-
Male	2	2	2	3	2	5	4	6	7	1	-	-
Female	1	2	3	1	8	7	18	11	15	2	1	-
White, both sexes	3	4	5	4	9	11	21	16	21	3	1	-
Male	2	2	2	3	2	5	4	5	6	1	-	-
Female	1	2	3	1	7	6	17	11	15	2	1	-
Black, both sexes	-	-	-	-	1	1	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	4	8	4	8	5	12	14	3	8	-	-	-
Male	4	3	3	2	2	8	5	2	1	-	-	-
Female	-	5	1	6	3	4	9	1	7	-	-	-
White, both sexes	2	6	3	7	3	7	12	3	8	-	-	-
Male	2	2	2	2	1	4	5	2	1	-	-	-
Female	-	4	1	5	2	3	7	1	7	-	-	-
Black, both sexes	1	1	1	1	2	5	2	-	-	-	-	-
Male	1	-	1	-	1	4	-	-	-	-	-	-
Female	-	1	-	1	1	1	2	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
New Hampshire												
All races, both sexes	4	1	7	2	8	6	12	5	6	2	1	-
Male	1	-	3	1	7	1	3	1	4	1	-	-
Female	3	1	4	1	1	5	9	4	2	1	1	-
White, both sexes	3	1	6	2	8	6	12	5	6	2	1	-
Male	1	-	2	1	7	1	3	1	4	1	-	-
Female	2	1	4	1	1	5	9	4	2	1	1	-
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	16	20	27	36	30	40	51	47	23	13	1	-
Male	8	11	14	10	14	16	19	12	5	2	-	-
Female	8	9	13	26	16	24	32	35	18	11	1	-
White, both sexes	13	15	23	25	23	36	46	42	20	12	-	-
Male	6	8	10	4	12	12	18	11	3	2	-	-
Female	7	7	13	21	11	24	28	31	17	10	-	-
Black, both sexes	3	3	3	9	7	3	4	5	2	1	1	-
Male	2	2	3	5	2	3	1	1	1	-	-	-
Female	1	1	-	4	5	-	3	4	1	1	1	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	2	-	1	1	-	1	-	-	-
Male	-	1	1	1	-	1	-	-	1	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
New Mexico												
All races, both sexes	2	7	10	4	10	14	8	12	3	3	-	-
Male	2	4	2	-	4	6	2	3	2	2	-	-
Female	-	3	8	4	6	8	6	9	1	1	-	-
White, both sexes	2	7	10	4	9	14	7	11	3	3	-	-
Male	2	4	2	-	4	6	2	2	2	2	-	-
Female	-	3	8	4	5	8	5	9	1	1	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	31	37	22	47	52	69	86	77	41	17	2	-
Male	16	14	11	19	24	25	35	24	9	4	-	-
Female	15	23	11	28	28	44	51	53	32	13	2	-
White, both sexes	22	31	18	35	43	61	79	70	38	15	1	-
Male	15	13	10	14	22	23	35	20	9	4	-	-
Female	7	18	8	21	21	38	44	50	29	11	1	-
Black, both sexes	8	6	4	12	7	6	5	6	3	2	1	-
Male	1	1	1	5	1	1	-	4	-	-	-	-
Female	7	5	3	7	6	5	5	2	3	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	2	2	2	1	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	1	-	-	-	1	1	2	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
North Carolina												
All races, both sexes	22	30	37	40	46	50	77	59	34	10	-	-
Male	8	14	19	17	21	22	28	16	5	3	-	-
Female	14	16	18	23	25	28	49	43	29	7	-	-
White, both sexes	15	14	24	32	30	37	65	47	30	8	-	-
Male	5	9	12	15	15	18	23	12	4	2	-	-
Female	10	5	12	17	15	19	42	35	26	6	-	-
Black, both sexes	7	16	13	7	15	12	12	12	4	2	-	-
Male	3	5	7	2	6	4	5	4	1	1	-	-
Female	4	11	6	5	9	8	7	8	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	1	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	1	2	2	2	5	3	6	2	2	-	-
Male	-	-	-	1	1	2	2	3	-	-	-	-
Female	-	1	2	1	1	3	1	3	2	2	-	-
White, both sexes	-	1	2	2	2	5	3	6	2	2	-	-
Male	-	-	-	1	1	2	2	3	-	-	-	-
Female	-	1	2	1	1	3	1	3	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Ohio												
All races, both sexes	28	28	31	33	55	44	64	65	37	8	-	-
Male	10	15	11	13	15	15	24	19	6	1	-	-
Female	18	13	20	20	40	29	40	46	31	7	-	-
White, both sexes	22	22	26	28	49	38	62	56	34	7	-	-
Male	8	13	11	12	14	14	22	16	6	1	-	-
Female	14	9	15	16	35	24	40	40	28	6	-	-
Black, both sexes	5	6	5	5	6	6	2	9	3	1	-	-
Male	1	2	-	1	1	1	2	3	-	-	-	-
Female	4	4	5	4	5	5	-	6	3	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	7	12	21	17	14	23	19	12	7	2	1	-
Male	3	4	8	9	6	7	5	4	2	-	-	-
Female	4	8	13	8	8	16	14	8	5	2	1	-
White, both sexes	5	9	20	17	13	17	17	11	7	2	1	-
Male	2	4	7	9	6	5	4	3	2	-	-	-
Female	3	5	13	8	7	12	13	8	5	2	1	-
Black, both sexes	-	1	1	-	1	3	1	1	-	-	-	-
Male	-	-	1	-	-	2	-	1	-	-	-	-
Female	-	1	-	-	1	1	1	-	-	-	-	-
American Indian 1/, both sexes	2	2	-	-	-	3	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	1	2	-	-	-	3	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Oregon												
All races, both sexes	9	14	8	15	16	14	28	22	14	7	1	-
Male	4	7	3	7	7	4	12	8	4	2	-	-
Female	5	7	5	8	9	10	16	14	10	5	1	-
White, both sexes	9	14	8	15	15	14	28	21	14	7	1	-
Male	4	7	3	7	6	4	12	8	4	2	-	-
Female	5	7	5	8	9	10	16	13	10	5	1	-
Black, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	28	44	43	54	60	78	113	98	57	9	2	-
Male	17	22	19	23	34	31	43	23	13	2	-	-
Female	11	22	24	31	26	47	70	75	44	7	2	-
White, both sexes	24	36	39	45	51	70	98	92	55	8	2	-
Male	15	17	17	17	29	28	37	22	13	1	-	-
Female	9	19	22	28	22	42	61	70	42	7	2	-
Black, both sexes	4	8	4	9	9	8	15	6	1	1	-	-
Male	2	5	2	6	5	3	6	1	-	1	-	-
Female	2	3	2	3	4	5	9	5	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Rhode Island												
All races, both sexes	2	6	1	4	6	8	7	8	4	3	-	-
Male	1	3	1	-	3	1	2	-	-	-	-	-
Female	1	3	-	4	3	7	5	8	4	3	-	-
White, both sexes	1	6	1	4	6	8	7	8	4	3	-	-
Male	1	3	1	-	3	1	2	-	-	-	-	-
Female	-	3	-	4	3	7	5	8	4	3	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	14	19	23	27	13	32	27	32	15	5	3	-
Male	6	13	6	11	3	13	8	12	3	1	-	-
Female	8	6	17	16	10	19	19	20	12	4	3	-
White, both sexes	6	12	14	17	9	27	22	26	10	5	1	-
Male	2	9	5	6	1	10	7	8	1	1	-	-
Female	4	3	9	11	8	17	15	18	9	4	1	-
Black, both sexes	8	6	9	10	4	5	5	5	4	-	2	-
Male	4	3	1	5	2	3	1	4	1	-	-	-
Female	4	3	8	5	2	2	4	1	3	-	2	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
South Dakota												
All races, both sexes	-	2	4	1	4	5	6	4	2	-	1	-
Male	-	-	3	-	3	5	1	-	-	-	-	-
Female	-	2	1	1	1	-	5	4	2	-	1	-
White, both sexes	-	2	4	1	4	5	6	4	2	-	1	-
Male	-	-	3	-	3	5	1	-	-	-	-	-
Female	-	2	1	1	1	-	5	4	2	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	10	24	23	29	29	39	44	45	13	6	2	-
Male	7	13	11	7	15	18	15	10	4	3	1	-
Female	3	11	12	22	14	21	29	35	9	3	1	-
White, both sexes	6	18	21	22	24	34	37	38	11	6	-	-
Male	5	10	10	6	13	16	13	8	3	3	-	-
Female	1	8	11	16	11	18	24	30	8	3	-	-
Black, both sexes	4	6	2	7	5	4	7	7	2	-	2	-
Male	2	3	1	1	2	1	2	2	1	-	1	-
Female	2	3	1	6	3	3	5	5	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Texas												
All races, both sexes	55	58	75	81	112	99	124	122	76	23	4	-
Male	22	24	39	35	41	47	47	41	20	3	3	-
Female	33	34	36	46	71	52	77	81	56	20	1	-
White, both sexes	38	43	58	63	89	77	106	111	60	19	3	-
Male	18	21	28	28	31	37	42	38	15	3	2	-
Female	20	22	30	35	58	40	64	73	45	16	1	-
Black, both sexes	16	13	17	17	21	21	18	9	16	3	1	-
Male	4	3	11	7	9	10	5	2	5	-	1	-
Female	12	10	6	10	12	11	13	7	11	3	-	-
American Indian 1/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	-	2	1	-	2	-	1	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	2	-	-	1	1	-	1	-	1	-	-
Utah												
All races, both sexes	7	10	10	6	8	15	9	23	13	2	-	-
Male	3	5	5	2	2	8	2	7	3	1	-	-
Female	4	5	5	4	6	7	7	16	10	1	-	-
White, both sexes	7	8	10	6	8	15	8	23	13	2	-	-
Male	3	4	5	2	2	8	2	7	3	1	-	-
Female	4	4	5	4	6	7	6	16	10	1	-	-
Black, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Vermont												
All races, both sexes	-	1	2	4	2	4	5	4	3	1	-	-
Male	-	-	2	2	1	2	1	1	2	1	-	-
Female	-	1	-	2	1	2	4	3	1	-	-	-
White, both sexes	-	1	2	4	2	4	5	4	3	1	-	-
Male	-	-	2	2	1	2	1	1	2	1	-	-
Female	-	1	-	2	1	2	4	3	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	17	29	18	35	39	38	53	38	22	10	1	-
Male	10	14	11	16	16	15	18	11	7	2	-	-
Female	7	15	7	19	23	23	35	27	15	8	1	-
White, both sexes	12	20	12	23	28	31	43	28	19	10	1	-
Male	6	8	9	9	13	14	15	7	6	2	-	-
Female	6	12	3	14	15	17	28	21	13	8	1	-
Black, both sexes	5	9	6	11	11	7	10	10	3	-	-	-
Male	4	6	2	6	3	1	3	4	1	-	-	-
Female	1	3	4	5	8	6	7	6	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Washington												
All races, both sexes	14	12	23	20	20	29	29	35	27	10	-	-
Male	4	5	12	8	8	12	12	9	7	-	-	-
Female	10	7	11	12	12	17	17	26	20	10	-	-
White, both sexes	12	12	23	19	19	27	28	32	26	10	-	-
Male	3	5	12	8	7	12	12	9	7	-	-	-
Female	9	7	11	11	12	15	16	23	19	10	-	-
Black, both sexes	1	-	-	-	1	2	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	2	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	-	1	1	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	1	1	-	-	-
West Virginia												
All races, both sexes	4	5	8	13	12	10	19	13	6	-	-	-
Male	2	2	6	7	4	8	8	3	-	-	-	-
Female	2	3	2	6	8	2	11	10	6	-	-	-
White, both sexes	4	4	8	12	11	10	18	13	5	-	-	-
Male	2	1	6	6	4	8	7	3	-	-	-	-
Female	2	3	2	6	7	2	11	10	5	-	-	-
Black, both sexes	-	1	-	1	1	-	1	-	1	-	-	-
Male	-	1	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Wisconsin												
All races, both sexes	9	19	15	26	25	32	68	41	32	5	-	-
Male	6	12	7	10	17	15	28	10	10	1	-	-
Female	3	7	8	16	8	17	40	31	22	4	-	-
White, both sexes	7	14	14	23	24	29	63	41	32	5	-	-
Male	4	10	7	10	17	15	25	10	10	1	-	-
Female	3	4	7	13	7	14	38	31	22	4	-	-
Black, both sexes	2	5	1	2	1	2	3	-	-	-	-	-
Male	2	2	-	-	-	-	3	-	-	-	-	-
Female	-	3	1	2	1	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Wyoming												
All races, both sexes	2	2	2	1	4	4	5	7	3	-	-	-
Male	-	1	1	1	2	2	3	1	1	-	-	-
Female	2	1	1	-	2	2	2	6	2	-	-	-
White, both sexes	2	2	1	1	4	4	5	7	3	-	-	-
Male	-	1	-	1	2	2	3	1	1	-	-	-
Female	2	1	1	-	2	2	2	6	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
United States												
All races, both sexes	7,097	4	2	1	2	22	47	103	99	185	261	344
Male	3,006	2	-	1	-	5	17	37	46	95	107	159
Female	4,091	2	2	-	2	17	30	66	53	90	154	185
White, both sexes	5,834	3	2	1	-	16	32	64	66	128	188	241
Male	2,449	2	-	1	-	4	11	25	30	65	75	114
Female	3,385	1	2	-	-	12	21	39	36	63	113	127
Black, both sexes	1,182	1	-	-	2	4	14	35	31	51	69	101
Male	519	-	-	-	-	-	5	9	14	26	30	45
Female	663	1	-	-	2	4	9	26	17	25	39	56
American Indian 1/, both sexes	27	-	-	-	-	2	-	3	2	2	2	-
Male	11	-	-	-	-	1	-	2	2	-	-	-
Female	16	-	-	-	-	1	-	1	-	2	2	-
Asian or Pacific Islander 2/, both sexes	54	-	-	-	-	-	1	1	-	4	2	2
Male	27	-	-	-	-	-	1	1	-	4	2	-
Female	27	-	-	-	-	-	-	-	-	-	-	2
Alabama												
All races, both sexes	171	-	-	-	-	-	1	4	2	7	7	4
Male	75	-	-	-	-	-	1	-	1	1	4	2
Female	96	-	-	-	-	-	-	4	1	6	3	2
White, both sexes	115	-	-	-	-	-	-	-	1	2	2	3
Male	47	-	-	-	-	-	-	-	-	-	1	1
Female	68	-	-	-	-	-	-	-	1	2	1	2
Black, both sexes	55	-	-	-	-	-	1	4	1	5	5	1
Male	28	-	-	-	-	-	1	-	1	1	3	1
Female	27	-	-	-	-	-	-	4	-	4	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Alaska												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	97	-	-	-	-	1	-	2	3	1	6	2
Male	46	-	-	-	-	1	-	1	2	-	3	2
Female	51	-	-	-	-	-	-	1	1	1	3	-
White, both sexes	83	-	-	-	-	-	-	1	1	1	5	2
Male	38	-	-	-	-	-	-	1	-	-	3	2
Female	45	-	-	-	-	-	-	-	1	1	2	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	5	-	-	-	-	1	-	1	2	-	-	-
Male	3	-	-	-	-	1	-	-	2	-	-	-
Female	2	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Arkansas												
All races, both sexes	125	-	1	-	-	-	-	1	3	1	4	5
Male	52	-	-	-	-	-	-	-	2	1	3	2
Female	73	-	1	-	-	-	-	1	1	-	1	3
White, both sexes	107	-	1	-	-	-	-	1	2	1	3	5
Male	43	-	-	-	-	-	-	-	1	1	3	2
Female	64	-	1	-	-	-	-	1	1	-	-	3
Black, both sexes	18	-	-	-	-	-	-	-	1	-	1	-
Male	9	-	-	-	-	-	-	-	1	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	322	-	-	-	-	2	1	8	5	11	13	14
Male	124	-	-	-	-	-	1	4	2	7	3	6
Female	198	-	-	-	-	2	-	4	3	4	10	8
White, both sexes	270	-	-	-	-	2	1	4	4	8	11	7
Male	102	-	-	-	-	-	1	3	2	4	2	2
Female	168	-	-	-	-	2	-	1	2	4	9	5
Black, both sexes	42	-	-	-	-	-	-	4	1	1	2	7
Male	17	-	-	-	-	-	-	1	-	1	1	4
Female	25	-	-	-	-	-	-	3	1	-	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	2	-	-
Male	5	-	-	-	-	-	-	-	-	2	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Colorado												
All races, both sexes	117	-	-	-	-	-	1	-	-	2	2	4
Male	63	-	-	-	-	-	-	-	-	1	-	2
Female	54	-	-	-	-	-	1	-	-	1	2	2
White, both sexes	111	-	-	-	-	-	1	-	-	2	1	4
Male	60	-	-	-	-	-	-	-	-	1	-	2
Female	51	-	-	-	-	-	1	-	-	1	1	2
Black, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	82	-	-	-	-	-	-	-	3	2	2	1
Male	27	-	-	-	-	-	-	-	1	1	-	-
Female	55	-	-	-	-	-	-	-	2	1	2	1
White, both sexes	69	-	-	-	-	-	-	-	2	2	1	1
Male	23	-	-	-	-	-	-	-	1	1	-	-
Female	46	-	-	-	-	-	-	-	1	1	1	1
Black, both sexes	11	-	-	-	-	-	-	-	1	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Delaware												
All races, both sexes	31	-	-	-	-	-	-	-	-	2	1	-
Male	16	-	-	-	-	-	-	-	-	1	-	-
Female	15	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	25	-	-	-	-	-	-	-	-	2	1	-
Male	12	-	-	-	-	-	-	-	-	1	-	-
Female	13	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	8	-	-	-	-	-	1	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	1	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Florida												
All races, both sexes	356	1	-	-	-	-	2	6	1	7	5	17
Male	153	-	-	-	-	-	-	-	1	4	3	6
Female	203	1	-	-	-	-	2	6	-	3	2	11
White, both sexes	291	-	-	-	-	-	2	4	1	4	3	9
Male	131	-	-	-	-	-	-	-	1	2	2	3
Female	160	-	-	-	-	-	2	4	-	2	1	6
Black, both sexes	65	1	-	-	-	-	-	2	-	3	2	8
Male	22	-	-	-	-	-	-	-	-	2	1	3
Female	43	1	-	-	-	-	-	2	-	1	1	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	229	-	-	-	-	1	4	5	5	5	11	12
Male	106	-	-	-	-	-	-	3	5	1	6	6
Female	123	-	-	-	-	1	4	2	-	4	5	6
White, both sexes	160	-	-	-	-	-	3	3	3	3	5	6
Male	74	-	-	-	-	-	-	2	3	1	3	3
Female	86	-	-	-	-	-	3	1	-	2	2	3
Black, both sexes	68	-	-	-	-	1	1	2	2	2	5	6
Male	32	-	-	-	-	-	-	1	2	-	3	3
Female	36	-	-	-	-	1	1	1	-	2	2	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Hawaii												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	1	2
Male	8	-	-	-	-	-	-	-	-	-	1	1
Female	8	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Idaho												
All races, both sexes	41	-	-	-	-	-	-	1	-	2	2	6
Male	15	-	-	-	-	-	-	1	-	-	1	-
Female	26	-	-	-	-	-	-	-	-	2	1	6
White, both sexes	40	-	-	-	-	-	-	1	-	2	2	6
Male	14	-	-	-	-	-	-	1	-	-	1	-
Female	26	-	-	-	-	-	-	-	-	2	1	6
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Illinois												
All races, both sexes	366	1	-	-	-	2	2	7	1	7	13	15
Male	149	-	-	-	-	1	2	3	-	5	3	8
Female	217	1	-	-	-	1	-	4	1	2	10	7
White, both sexes	289	1	-	-	-	1	1	6	1	6	9	10
Male	116	-	-	-	-	1	1	3	-	5	2	5
Female	173	1	-	-	-	-	-	3	1	1	7	5
Black, both sexes	73	-	-	-	-	1	-	1	-	1	4	5
Male	31	-	-	-	-	-	-	-	-	-	1	3
Female	42	-	-	-	-	1	-	1	-	1	3	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	1	-	-	-	-	-
Male	2	-	-	-	-	-	1	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	271	-	-	-	-	1	1	2	5	9	11	16
Male	123	-	-	-	-	-	1	-	3	6	9	9
Female	148	-	-	-	-	1	-	2	2	3	2	7
White, both sexes	248	-	-	-	-	1	1	2	5	6	9	16
Male	111	-	-	-	-	-	1	-	3	3	8	9
Female	137	-	-	-	-	1	-	2	2	3	1	7
Black, both sexes	23	-	-	-	-	-	-	-	-	3	2	-
Male	12	-	-	-	-	-	-	-	-	3	1	-
Female	11	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Iowa												
All races, both sexes	86	1	-	-	-	-	2	-	-	2	2	3
Male	28	1	-	-	-	-	-	-	-	-	2	1
Female	58	-	-	-	-	-	2	-	-	2	-	2
White, both sexes	82	1	-	-	-	-	2	-	-	2	2	3
Male	27	1	-	-	-	-	-	-	-	-	2	1
Female	55	-	-	-	-	-	2	-	-	2	-	2
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	100	-	-	-	-	2	1	1	1	3	4	6
Male	40	-	-	-	-	1	-	-	-	1	2	3
Female	60	-	-	-	-	1	1	1	1	2	2	3
White, both sexes	96	-	-	-	-	1	1	-	1	2	4	6
Male	40	-	-	-	-	1	-	-	-	1	2	3
Female	56	-	-	-	-	-	1	-	1	1	2	3
Black, both sexes	3	-	-	-	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	1	-	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Kentucky												
All races, both sexes	144	-	-	-	-	1	-	1	-	1	8	6
Male	65	-	-	-	-	-	-	-	-	-	2	2
Female	79	-	-	-	-	1	-	1	-	1	6	4
White, both sexes	134	-	-	-	-	1	-	1	-	1	6	5
Male	59	-	-	-	-	-	-	-	-	-	1	1
Female	75	-	-	-	-	1	-	1	-	1	5	4
Black, both sexes	10	-	-	-	-	-	-	-	-	-	2	1
Male	6	-	-	-	-	-	-	-	-	-	1	1
Female	4	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	162	1	-	-	1	-	-	1	4	8	3	11
Male	67	1	-	-	-	-	-	-	1	3	3	5
Female	95	-	-	-	1	-	-	1	3	5	-	6
White, both sexes	101	1	-	-	-	-	-	1	2	2	2	3
Male	41	1	-	-	-	-	-	-	-	-	2	1
Female	60	-	-	-	-	-	-	1	2	2	-	2
Black, both sexes	60	-	-	-	1	-	-	-	2	6	1	8
Male	25	-	-	-	-	-	-	-	1	3	1	4
Female	35	-	-	-	1	-	-	-	1	3	-	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Maine												
All races, both sexes	34	-	-	-	-	-	-	-	1	-	1	3
Male	12	-	-	-	-	-	-	-	-	-	1	2
Female	22	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	34	-	-	-	-	-	-	-	1	-	1	3
Male	12	-	-	-	-	-	-	-	-	-	1	2
Female	22	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	139	-	-	-	-	-	-	4	1	4	6	12
Male	62	-	-	-	-	-	-	3	1	4	3	9
Female	77	-	-	-	-	-	-	1	-	-	3	3
White, both sexes	90	-	-	-	-	-	-	2	1	2	2	6
Male	41	-	-	-	-	-	-	1	1	2	1	5
Female	49	-	-	-	-	-	-	1	-	-	1	1
Black, both sexes	45	-	-	-	-	-	-	1	-	2	4	6
Male	18	-	-	-	-	-	-	1	-	2	2	4
Female	27	-	-	-	-	-	-	-	-	-	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	1	-	-	-	-
Male	3	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Massachusetts												
All races, both sexes	121	-	-	-	-	-	-	1	-	3	1	1
Male	54	-	-	-	-	-	-	-	-	2	-	-
Female	67	-	-	-	-	-	-	1	-	1	1	1
White, both sexes	110	-	-	-	-	-	-	1	-	2	1	-
Male	48	-	-	-	-	-	-	-	-	1	-	-
Female	62	-	-	-	-	-	-	1	-	1	1	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	230	-	-	-	-	-	1	4	1	5	4	8
Male	101	-	-	-	-	-	-	3	1	3	1	4
Female	129	-	-	-	-	-	1	1	-	2	3	4
White, both sexes	184	-	-	-	-	-	1	3	-	4	2	6
Male	78	-	-	-	-	-	-	2	-	3	-	3
Female	106	-	-	-	-	-	1	1	-	1	2	3
Black, both sexes	45	-	-	-	-	-	-	1	1	-	2	2
Male	23	-	-	-	-	-	-	1	1	-	1	1
Female	22	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Minnesota												
All races, both sexes	104	-	-	-	-	1	3	-	4	4	7	5
Male	42	-	-	-	-	-	-	-	2	1	2	-
Female	62	-	-	-	-	1	3	-	2	3	5	5
White, both sexes	101	-	-	-	-	1	3	-	3	4	7	5
Male	40	-	-	-	-	-	-	-	2	1	2	-
Female	61	-	-	-	-	1	3	-	1	3	5	5
Black, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	111	-	-	-	-	1	3	1	1	3	6	6
Male	47	-	-	-	-	-	1	-	-	2	2	4
Female	64	-	-	-	-	1	2	1	1	1	4	2
White, both sexes	65	-	-	-	-	1	1	-	-	1	3	2
Male	27	-	-	-	-	-	1	-	-	-	1	2
Female	38	-	-	-	-	1	-	-	-	1	2	-
Black, both sexes	46	-	-	-	-	-	2	1	1	2	3	4
Male	20	-	-	-	-	-	-	-	-	2	1	2
Female	26	-	-	-	-	-	2	1	1	-	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Missouri												
All races, both sexes	193	-	-	1	-	-	-	1	3	7	9	16
Male	78	-	-	1	-	-	-	-	1	2	1	12
Female	115	-	-	-	-	-	-	1	2	5	8	4
White, both sexes	163	-	-	1	-	-	-	-	2	6	7	13
Male	63	-	-	1	-	-	-	-	1	2	-	9
Female	100	-	-	-	-	-	-	-	1	4	7	4
Black, both sexes	28	-	-	-	-	-	-	1	1	1	2	3
Male	15	-	-	-	-	-	-	-	-	-	1	3
Female	13	-	-	-	-	-	-	1	1	1	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	21	-	-	-	-	-	-	1	-	1	1	4
Male	9	-	-	-	-	-	-	-	-	1	-	2
Female	12	-	-	-	-	-	-	1	-	-	1	2
White, both sexes	21	-	-	-	-	-	-	1	-	1	1	4
Male	9	-	-	-	-	-	-	-	-	1	-	2
Female	12	-	-	-	-	-	-	1	-	-	1	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Nebraska												
All races, both sexes	53	-	-	-	-	-	-	1	1	3	3	1
Male	21	-	-	-	-	-	-	1	-	3	2	1
Female	32	-	-	-	-	-	-	-	1	-	1	-
White, both sexes	50	-	-	-	-	-	-	-	1	3	3	1
Male	19	-	-	-	-	-	-	-	-	3	2	1
Female	31	-	-	-	-	-	-	-	1	-	1	-
Black, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	59	-	-	-	-	-	-	2	1	3	3	2
Male	30	-	-	-	-	-	-	1	-	3	2	-
Female	29	-	-	-	-	-	-	1	1	-	1	2
White, both sexes	42	-	-	-	-	-	-	1	1	2	2	1
Male	19	-	-	-	-	-	-	-	-	2	1	-
Female	23	-	-	-	-	-	-	1	1	-	1	1
Black, both sexes	15	-	-	-	-	-	-	1	-	1	-	1
Male	9	-	-	-	-	-	-	1	-	1	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
New Hampshire												
All races, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	27	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	205	-	-	-	-	1	1	4	3	7	14	12
Male	75	-	-	-	-	-	-	1	2	4	7	3
Female	130	-	-	-	-	1	1	3	1	3	7	9
White, both sexes	160	-	-	-	-	-	-	1	2	5	9	7
Male	56	-	-	-	-	-	-	-	2	3	5	2
Female	104	-	-	-	-	-	-	1	-	2	4	5
Black, both sexes	43	-	-	-	-	1	1	3	1	2	5	5
Male	17	-	-	-	-	-	-	1	-	1	2	1
Female	26	-	-	-	-	1	1	2	1	1	3	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
New Mexico												
All races, both sexes	41	-	-	-	-	-	-	1	-	2	1	-
Male	18	-	-	-	-	-	-	1	-	1	-	-
Female	23	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	40	-	-	-	-	-	-	1	-	2	1	-
Male	18	-	-	-	-	-	-	1	-	1	-	-
Female	22	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	253	-	-	-	-	1	1	2	2	10	8	9
Male	121	-	-	-	-	1	1	2	2	8	3	4
Female	132	-	-	-	-	-	-	-	-	2	5	5
White, both sexes	222	-	-	-	-	1	1	1	1	8	8	5
Male	108	-	-	-	-	1	1	1	1	6	3	2
Female	114	-	-	-	-	-	-	-	-	2	5	3
Black, both sexes	30	-	-	-	-	-	-	1	1	2	-	4
Male	13	-	-	-	-	-	-	1	1	2	-	2
Female	17	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
North Carolina												
All races, both sexes	263	-	-	-	-	-	4	4	8	10	7	14
Male	103	-	-	-	-	-	1	1	2	4	2	7
Female	160	-	-	-	-	-	3	3	6	6	5	7
White, both sexes	187	-	-	-	-	-	3	2	3	2	3	6
Male	75	-	-	-	-	-	-	1	2	1	-	4
Female	112	-	-	-	-	-	3	1	1	1	3	2
Black, both sexes	72	-	-	-	-	-	1	2	5	7	4	8
Male	28	-	-	-	-	-	1	-	-	3	2	3
Female	44	-	-	-	-	-	-	2	5	4	2	5
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	18	-	-	-	-	-	-	1	-	-	1	1
Male	8	-	-	-	-	-	-	1	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	18	-	-	-	-	-	-	1	-	-	1	1
Male	8	-	-	-	-	-	-	1	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Ohio												
All races, both sexes	234	-	-	-	-	2	-	3	6	7	12	11
Male	87	-	-	-	-	-	-	-	2	6	4	4
Female	147	-	-	-	-	2	-	3	4	1	8	7
White, both sexes	203	-	-	-	-	2	-	2	5	4	10	9
Male	76	-	-	-	-	-	-	-	2	3	3	4
Female	127	-	-	-	-	2	-	2	3	1	7	5
Black, both sexes	29	-	-	-	-	-	-	1	1	2	2	2
Male	9	-	-	-	-	-	-	-	-	2	1	-
Female	20	-	-	-	-	-	-	1	1	-	1	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	102	-	-	-	-	1	-	5	-	3	5	4
Male	43	-	-	-	-	-	-	2	-	1	2	3
Female	59	-	-	-	-	1	-	3	-	2	3	1
White, both sexes	89	-	-	-	-	1	-	4	-	2	4	3
Male	38	-	-	-	-	-	-	1	-	1	2	3
Female	51	-	-	-	-	1	-	3	-	1	2	-
Black, both sexes	7	-	-	-	-	-	-	-	-	1	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	6	-	-	-	-	-	-	1	-	-	1	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Oregon												
All races, both sexes	68	-	-	-	-	1	2	-	-	2	-	4
Male	27	-	-	-	-	-	1	-	-	-	-	1
Female	41	-	-	-	-	1	1	-	-	2	-	3
White, both sexes	67	-	-	-	-	1	2	-	-	2	-	4
Male	26	-	-	-	-	-	1	-	-	-	-	1
Female	41	-	-	-	-	1	1	-	-	2	-	3
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	374	-	1	-	-	-	3	6	8	8	22	23
Male	170	-	-	-	-	-	2	1	4	4	11	10
Female	204	-	1	-	-	-	1	5	4	4	11	13
White, both sexes	318	-	1	-	-	-	2	3	6	7	16	18
Male	141	-	-	-	-	-	2	1	2	4	7	9
Female	177	-	1	-	-	-	-	2	4	3	9	9
Black, both sexes	56	-	-	-	-	-	1	3	2	1	6	5
Male	29	-	-	-	-	-	-	-	2	-	4	1
Female	27	-	-	-	-	-	1	3	-	1	2	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Rhode Island												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	151	-	-	-	-	-	1	1	6	4	6	9
Male	62	-	-	-	-	-	-	-	2	2	2	2
Female	89	-	-	-	-	-	1	1	4	2	4	7
White, both sexes	94	-	-	-	-	-	-	-	2	-	3	5
Male	33	-	-	-	-	-	-	-	-	-	2	-
Female	61	-	-	-	-	-	-	-	2	-	1	5
Black, both sexes	56	-	-	-	-	-	1	1	4	4	3	4
Male	28	-	-	-	-	-	-	-	2	2	-	2
Female	28	-	-	-	-	-	1	1	2	2	3	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
South Dakota												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	2
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	24	-	-	-	-	-	-	-	-	-	-	2
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	222	-	-	-	-	1	4	2	7	5	7	16
Male	91	-	-	-	-	1	3	-	3	2	2	5
Female	131	-	-	-	-	-	1	2	4	3	5	11
White, both sexes	182	-	-	-	-	1	1	2	5	3	6	9
Male	75	-	-	-	-	1	1	-	2	2	2	2
Female	107	-	-	-	-	-	-	2	3	1	4	7
Black, both sexes	39	-	-	-	-	-	3	-	2	2	1	7
Male	15	-	-	-	-	-	2	-	1	-	-	3
Female	24	-	-	-	-	-	1	-	1	2	1	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Texas												
All races, both sexes	610	-	-	-	-	1	5	7	5	13	19	17
Male	252	-	-	-	-	-	2	1	1	7	9	10
Female	358	-	-	-	-	1	3	6	4	6	10	7
White, both sexes	483	-	-	-	-	-	3	4	3	11	16	15
Male	200	-	-	-	-	-	1	-	-	6	7	10
Female	283	-	-	-	-	-	2	4	3	5	9	5
Black, both sexes	124	-	-	-	-	1	2	3	2	2	3	2
Male	51	-	-	-	-	-	1	1	1	1	2	-
Female	73	-	-	-	-	1	1	2	1	1	1	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	72	-	-	-	-	-	2	2	1	1	3	4
Male	27	-	-	-	-	-	1	1	-	-	-	-
Female	45	-	-	-	-	-	1	1	1	1	3	4
White, both sexes	69	-	-	-	-	-	2	2	1	1	2	3
Male	26	-	-	-	-	-	1	1	-	-	-	-
Female	43	-	-	-	-	-	1	1	1	1	2	3
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Vermont												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	205	-	-	-	1	2	1	3	2	2	7	10
Male	94	-	-	-	-	-	-	2	1	1	1	5
Female	111	-	-	-	1	2	1	1	1	1	6	5
White, both sexes	157	-	-	-	-	2	1	2	2	2	5	5
Male	71	-	-	-	-	-	-	2	1	1	-	4
Female	86	-	-	-	-	2	1	-	1	1	5	1
Black, both sexes	47	-	-	-	1	-	-	1	-	-	2	5
Male	22	-	-	-	-	-	-	-	-	-	1	1
Female	25	-	-	-	1	-	-	1	-	-	1	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Washington												
All races, both sexes	123	-	-	-	-	-	-	2	1	3	3	6
Male	44	-	-	-	-	-	-	1	-	-	1	2
Female	79	-	-	-	-	-	-	1	1	3	2	4
White, both sexes	111	-	-	-	-	-	-	2	1	3	1	4
Male	42	-	-	-	-	-	-	1	-	-	1	1
Female	69	-	-	-	-	-	-	1	1	3	-	3
Black, both sexes	8	-	-	-	-	-	-	-	-	-	2	2
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	2	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	68	-	-	-	-	-	-	1	1	1	1	5
Male	34	-	-	-	-	-	-	-	1	1	1	2
Female	34	-	-	-	-	-	-	1	-	-	-	3
White, both sexes	64	-	-	-	-	-	-	1	1	1	-	5
Male	31	-	-	-	-	-	-	-	1	1	-	2
Female	33	-	-	-	-	-	-	1	-	-	-	3
Black, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Wisconsin												
All races, both sexes	189	-	-	-	-	-	-	4	2	4	5	12
Male	87	-	-	-	-	-	-	2	2	1	3	8
Female	102	-	-	-	-	-	-	2	-	3	2	4
White, both sexes	172	-	-	-	-	-	-	4	1	4	4	9
Male	81	-	-	-	-	-	-	2	1	1	3	6
Female	91	-	-	-	-	-	-	2	-	3	1	3
Black, both sexes	13	-	-	-	-	-	-	-	1	-	1	3
Male	6	-	-	-	-	-	-	-	1	-	-	2
Female	7	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	22	-	-	-	-	-	-	1	1	-	2	1
Male	11	-	-	-	-	-	-	1	1	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	20	-	-	-	-	-	-	-	1	-	2	1
Male	9	-	-	-	-	-	-	-	1	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
United States												
All races, both sexes	431	470	548	607	679	810	994	843	476	145	24	-
Male	227	257	277	279	305	361	372	283	141	30	5	-
Female	204	213	271	328	374	449	622	560	335	115	19	-
White, both sexes	332	373	459	493	560	688	875	740	425	131	17	-
Male	169	204	231	220	254	315	328	245	123	30	3	-
Female	163	169	228	273	306	373	547	495	302	101	14	-
Black, both sexes	92	92	85	105	115	111	110	96	48	13	7	-
Male	52	50	43	55	50	45	42	34	17	-	2	-
Female	40	42	42	50	65	66	68	62	31	13	5	-
American Indian 1/, both sexes	3	2	1	2	-	5	1	1	1	-	-	-
Male	3	1	1	-	-	1	-	-	-	-	-	-
Female	-	1	-	2	-	4	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	4	3	3	7	4	6	8	6	2	1	-	-
Male	3	2	2	4	1	-	2	4	1	-	-	-
Female	1	1	1	3	3	6	6	2	1	1	-	-
Alabama												
All races, both sexes	9	10	12	13	19	22	29	19	12	1	-	-
Male	4	7	7	9	10	8	9	9	3	-	-	-
Female	5	3	5	4	9	14	20	10	9	1	-	-
White, both sexes	6	9	11	9	11	16	23	14	7	1	-	-
Male	2	6	7	6	6	5	6	6	1	-	-	-
Female	4	3	4	3	5	11	17	8	6	1	-	-
Black, both sexes	3	1	1	4	7	6	6	5	5	-	-	-
Male	2	1	-	3	4	3	3	3	2	-	-	-
Female	1	-	1	1	3	3	3	2	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Alaska												
All races, both sexes	1	-	-	-	2	1	-	-	-	-	-	-
Male	-	-	-	-	2	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	2	1	-	-	-	-	-	-
Male	-	-	-	-	2	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2	6	7	10	13	17	10	9	4	4	-	-
Male	1	4	3	6	5	8	4	3	1	2	-	-
Female	1	2	4	4	8	9	6	6	3	2	-	-
White, both sexes	-	4	7	10	12	14	9	9	4	4	-	-
Male	-	2	3	6	4	7	4	3	1	2	-	-
Female	-	2	4	4	8	7	5	6	3	2	-	-
Black, both sexes	2	1	-	-	1	2	1	-	-	-	-	-
Male	1	1	-	-	1	1	-	-	-	-	-	-
Female	1	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Arkansas												
All races, both sexes	9	4	7	14	15	17	14	17	10	2	1	-
Male	2	3	3	6	7	8	5	5	4	1	-	-
Female	7	1	4	8	8	9	9	12	6	1	1	-
White, both sexes	7	3	6	10	15	14	10	17	9	2	1	-
Male	1	2	3	3	7	8	3	5	3	1	-	-
Female	6	1	3	7	8	6	7	12	6	1	1	-
Black, both sexes	2	1	1	4	-	3	4	-	1	-	-	-
Male	1	1	-	3	-	-	2	-	1	-	-	-
Female	1	-	1	1	-	3	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	22	27	23	20	22	36	41	37	30	10	-	-
Male	8	8	10	6	9	17	18	12	10	3	-	-
Female	14	19	13	14	13	19	23	25	20	7	-	-
White, both sexes	19	19	17	18	19	34	35	35	29	8	-	-
Male	7	5	7	5	8	16	16	11	10	3	-	-
Female	12	14	10	13	11	18	19	24	19	5	-	-
Black, both sexes	2	8	6	-	3	2	4	1	-	1	-	-
Male	1	3	3	-	1	1	1	-	-	-	-	-
Female	1	5	3	-	2	1	3	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	2	-	-	2	1	1	1	-	-
Male	-	-	-	1	-	-	1	1	-	-	-	-
Female	1	-	-	1	-	-	1	-	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Colorado												
All races, both sexes	11	9	8	10	14	17	12	17	8	1	1	-
Male	8	4	5	6	10	10	5	8	4	-	-	-
Female	3	5	3	4	4	7	7	9	4	1	1	-
White, both sexes	10	8	8	10	12	17	12	17	7	1	1	-
Male	7	4	5	6	9	10	5	8	3	-	-	-
Female	3	4	3	4	3	7	7	9	4	1	1	-
Black, both sexes	1	1	-	-	2	-	-	-	1	-	-	-
Male	1	-	-	-	1	-	-	-	1	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	3	5	7	7	6	12	9	11	8	5	1	-
Male	2	4	3	2	3	4	2	2	1	1	1	-
Female	1	1	4	5	3	8	7	9	7	4	-	-
White, both sexes	2	3	5	5	6	9	9	11	7	5	1	-
Male	1	3	2	1	3	4	2	2	1	1	1	-
Female	1	-	3	4	3	5	7	9	6	4	-	-
Black, both sexes	1	2	2	1	-	2	-	-	1	-	-	-
Male	1	1	1	-	-	-	-	-	-	-	-	-
Female	-	1	1	1	-	2	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Delaware												
All races, both sexes	4	2	1	-	5	2	3	4	5	2	-	-
Male	2	2	-	-	4	1	2	2	2	-	-	-
Female	2	-	1	-	1	1	1	2	3	2	-	-
White, both sexes	2	2	-	-	4	1	2	4	5	2	-	-
Male	1	2	-	-	3	-	1	2	2	-	-	-
Female	1	-	-	-	1	1	1	2	3	2	-	-
Black, both sexes	2	-	1	-	1	1	1	-	-	-	-	-
Male	1	-	-	-	1	1	1	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	3	1	-	-	1	2	-	-	-	-
Male	-	-	2	-	-	-	-	1	-	-	-	-
Female	-	-	1	1	-	-	1	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	3	1	-	-	1	2	-	-	-	-
Male	-	-	2	-	-	-	-	1	-	-	-	-
Female	-	-	1	1	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Florida												
All races, both sexes	20	17	22	30	38	55	66	53	10	5	1	-
Male	7	9	12	19	13	28	28	17	6	-	-	-
Female	13	8	10	11	25	27	38	36	4	5	1	-
White, both sexes	13	14	20	22	35	46	60	46	7	4	1	-
Male	4	8	11	13	13	27	26	16	5	-	-	-
Female	9	6	9	9	22	19	34	30	2	4	1	-
Black, both sexes	7	3	2	8	3	9	6	7	3	1	-	-
Male	3	1	1	6	-	1	2	1	1	-	-	-
Female	4	2	1	2	3	8	4	6	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	17	24	17	19	21	25	21	29	9	3	1	-
Male	10	12	9	11	8	15	11	6	2	1	-	-
Female	7	12	8	8	13	10	10	23	7	2	1	-
White, both sexes	10	19	12	16	14	18	16	23	7	2	-	-
Male	5	11	7	9	6	9	9	3	2	1	-	-
Female	5	8	5	7	8	9	7	20	5	1	-	-
Black, both sexes	7	5	5	3	7	7	5	6	2	1	1	-
Male	5	1	2	2	2	6	2	3	-	-	-	-
Female	2	4	3	1	5	1	3	3	2	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Hawaii												
All races, both sexes	1	-	3	2	1	2	1	2	-	1	-	-
Male	1	-	3	1	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	2	1	2	-	1	-	-
White, both sexes	1	-	3	2	-	-	1	1	-	1	-	-
Male	1	-	3	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	2	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	1	-	-	-	-
Idaho												
All races, both sexes	1	2	4	3	3	5	3	5	3	-	1	-
Male	-	2	1	2	1	3	1	1	2	-	-	-
Female	1	-	3	1	2	2	2	4	1	-	1	-
White, both sexes	1	2	4	3	3	5	2	5	3	-	1	-
Male	-	2	1	2	1	3	-	1	2	-	-	-
Female	1	-	3	1	2	2	2	4	1	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Illinois												
All races, both sexes	24	18	38	33	30	39	55	39	28	12	2	-
Male	11	12	18	16	14	13	19	16	6	2	-	-
Female	13	6	20	17	16	26	36	23	22	10	2	-
White, both sexes	20	11	31	25	22	30	44	31	26	12	2	-
Male	9	6	15	13	12	10	14	12	6	2	-	-
Female	11	5	16	12	10	20	30	19	20	10	2	-
Black, both sexes	4	7	6	8	8	9	9	8	2	-	-	-
Male	2	6	2	3	2	3	5	4	-	-	-	-
Female	2	1	4	5	6	6	4	4	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	2	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Indiana												
All races, both sexes	18	29	32	20	24	26	36	20	14	6	1	-
Male	11	13	14	8	12	10	13	7	4	3	-	-
Female	7	16	18	12	12	16	23	13	10	3	1	-
White, both sexes	16	24	31	19	22	25	35	18	13	5	-	-
Male	9	11	14	7	10	10	13	7	3	3	-	-
Female	7	13	17	12	12	15	22	11	10	2	-	-
Black, both sexes	2	5	1	1	2	1	1	2	1	1	1	-
Male	2	2	-	1	2	-	-	-	1	-	-	-
Female	-	3	1	-	-	1	1	2	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Iowa												
All races, both sexes	5	10	4	5	7	11	18	5	10	1	-	-
Male	1	5	1	3	1	1	6	4	2	-	-	-
Female	4	5	3	2	6	10	12	1	8	1	-	-
White, both sexes	4	8	3	5	7	11	18	5	10	1	-	-
Male	1	4	1	3	1	1	6	4	2	-	-	-
Female	3	4	2	2	6	10	12	1	8	1	-	-
Black, both sexes	1	2	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	10	8	10	3	8	12	9	9	10	3	-	-
Male	7	3	6	1	4	3	1	5	3	-	-	-
Female	3	5	4	2	4	9	8	4	7	3	-	-
White, both sexes	10	8	10	3	8	11	9	9	10	3	-	-
Male	7	3	6	1	4	3	1	5	3	-	-	-
Female	3	5	4	2	4	8	8	4	7	3	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Kentucky												
All races, both sexes	10	11	13	15	16	20	15	19	6	2	-	-
Male	7	5	5	10	10	10	6	5	2	1	-	-
Female	3	6	8	5	6	10	9	14	4	1	-	-
White, both sexes	9	11	12	15	13	20	13	19	6	2	-	-
Male	6	5	5	10	8	10	5	5	2	1	-	-
Female	3	6	7	5	5	10	8	14	4	1	-	-
Black, both sexes	1	-	1	-	3	-	2	-	-	-	-	-
Male	1	-	-	-	2	-	1	-	-	-	-	-
Female	-	-	1	-	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	16	10	20	11	15	11	26	14	8	2	-	-
Male	9	6	9	5	5	4	7	5	4	-	-	-
Female	7	4	11	6	10	7	19	9	4	2	-	-
White, both sexes	10	7	13	6	9	5	20	11	8	1	-	-
Male	6	4	7	2	4	2	4	4	4	-	-	-
Female	4	3	6	4	5	3	16	7	4	1	-	-
Black, both sexes	6	3	7	5	6	6	6	2	-	1	-	-
Male	3	2	2	3	1	2	3	-	-	-	-	-
Female	3	1	5	2	5	4	3	2	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Maine												
All races, both sexes	-	3	1	2	1	10	4	5	2	1	-	-
Male	-	1	-	1	-	5	-	2	-	-	-	-
Female	-	2	1	1	1	5	4	3	2	1	-	-
White, both sexes	-	3	1	2	1	10	4	5	2	1	-	-
Male	-	1	-	1	-	5	-	2	-	-	-	-
Female	-	2	1	1	1	5	4	3	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	12	13	7	11	12	10	19	17	8	3	-	-
Male	7	9	2	7	2	3	3	6	3	-	-	-
Female	5	4	5	4	10	7	16	11	5	3	-	-
White, both sexes	6	9	5	7	8	7	12	14	6	3	-	-
Male	4	6	1	4	2	3	2	6	3	-	-	-
Female	2	3	4	3	6	4	10	8	3	3	-	-
Black, both sexes	5	4	2	3	4	2	7	3	2	-	-	-
Male	2	3	1	2	-	-	1	-	-	-	-	-
Female	3	1	1	1	4	2	6	3	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	1	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Massachusetts												
All races, both sexes	3	7	8	8	9	14	28	23	13	1	1	-
Male	2	5	4	6	5	7	9	9	5	-	-	-
Female	1	2	4	2	4	7	19	14	8	1	1	-
White, both sexes	3	7	7	7	7	12	27	22	12	1	1	-
Male	2	5	4	5	3	7	9	8	4	-	-	-
Female	1	2	3	2	4	5	18	14	8	1	1	-
Black, both sexes	-	-	1	1	2	1	1	1	1	-	-	-
Male	-	-	-	1	2	-	-	1	1	-	-	-
Female	-	-	1	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Michigan												
All races, both sexes	18	14	14	17	25	30	29	37	15	7	1	-
Male	9	10	10	7	12	13	7	13	5	3	-	-
Female	9	4	4	10	13	17	22	24	10	4	1	-
White, both sexes	13	11	10	13	21	25	26	29	13	6	1	-
Male	7	8	6	7	9	9	7	10	4	3	-	-
Female	6	3	4	6	12	16	19	19	9	3	1	-
Black, both sexes	5	3	4	4	4	5	3	8	2	1	-	-
Male	2	2	4	-	3	4	-	3	1	-	-	-
Female	3	1	-	4	1	1	3	5	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Minnesota												
All races, both sexes	8	3	7	5	14	9	13	9	9	3	-	-
Male	4	-	5	3	7	2	8	4	4	-	-	-
Female	4	3	2	2	7	7	5	5	5	3	-	-
White, both sexes	6	3	7	5	14	9	13	9	9	3	-	-
Male	2	-	5	3	7	2	8	4	4	-	-	-
Female	4	3	2	2	7	7	5	5	5	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	5	3	11	11	11	17	15	10	6	1	-	-
Male	4	2	6	3	4	8	5	4	2	-	-	-
Female	1	1	5	8	7	9	10	6	4	1	-	-
White, both sexes	2	3	6	6	6	10	13	6	5	-	-	-
Male	1	2	3	2	2	6	4	2	1	-	-	-
Female	1	1	3	4	4	4	9	4	4	-	-	-
Black, both sexes	3	-	5	5	5	7	2	4	1	1	-	-
Male	3	-	3	1	2	2	1	2	1	-	-	-
Female	-	-	2	4	3	5	1	2	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Missouri												
All races, both sexes	15	10	21	16	20	17	26	16	10	4	1	-
Male	6	8	11	8	9	5	8	4	1	1	-	-
Female	9	2	10	8	11	12	18	12	9	3	1	-
White, both sexes	11	8	16	14	17	15	24	15	9	4	1	-
Male	4	7	8	7	7	4	7	4	1	1	-	-
Female	7	1	8	7	10	11	17	11	8	3	1	-
Black, both sexes	4	1	5	2	3	2	2	1	-	-	-	-
Male	2	1	3	1	2	1	1	-	-	-	-	-
Female	2	-	2	1	1	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	-	2	4	1	3	1	2	-	-	-	-
Male	1	-	-	1	1	2	-	1	-	-	-	-
Female	-	-	2	3	-	1	1	1	-	-	-	-
White, both sexes	1	-	2	4	1	3	1	2	-	-	-	-
Male	1	-	-	1	1	2	-	1	-	-	-	-
Female	-	-	2	3	-	1	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Nebraska												
All races, both sexes	1	3	3	1	4	7	12	6	6	1	-	-
Male	1	1	1	1	1	3	1	4	1	-	-	-
Female	-	2	2	-	3	4	11	2	5	1	-	-
White, both sexes	1	3	3	1	4	6	12	5	6	1	-	-
Male	1	1	1	1	1	3	1	3	1	-	-	-
Female	-	2	2	-	3	3	11	2	5	1	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	4	5	3	6	2	10	9	2	7	-	-	-
Male	4	2	2	2	1	7	4	1	1	-	-	-
Female	-	3	1	4	1	3	5	1	6	-	-	-
White, both sexes	2	4	2	5	1	5	7	2	7	-	-	-
Male	2	2	1	2	-	3	4	1	1	-	-	-
Female	-	2	1	3	1	2	3	1	6	-	-	-
Black, both sexes	1	1	1	1	1	5	2	-	-	-	-	-
Male	1	-	1	-	1	4	-	-	-	-	-	-
Female	-	1	-	1	-	1	2	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
New Hampshire												
All races, both sexes	-	-	5	1	4	4	7	3	3	-	1	-
Male	-	-	2	-	3	1	2	1	2	-	-	-
Female	-	-	3	1	1	3	5	2	1	-	1	-
White, both sexes	-	-	4	1	4	4	7	3	3	-	1	-
Male	-	-	1	-	3	1	2	1	2	-	-	-
Female	-	-	3	1	1	3	5	2	1	-	1	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	9	9	13	19	20	20	30	25	13	4	1	-
Male	4	5	7	6	9	7	10	7	3	-	-	-
Female	5	4	6	13	11	13	20	18	10	4	1	-
White, both sexes	9	6	11	13	16	19	27	21	11	3	-	-
Male	4	2	5	2	8	6	9	6	2	-	-	-
Female	5	4	6	11	8	13	18	15	9	3	-	-
Black, both sexes	-	2	1	6	4	1	3	4	2	1	1	-
Male	-	2	1	4	1	1	1	1	1	-	-	-
Female	-	-	-	2	3	-	2	3	1	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
New Mexico												
All races, both sexes	1	5	6	2	4	5	4	6	2	2	-	-
Male	1	2	1	-	4	2	1	2	1	2	-	-
Female	-	3	5	2	-	3	3	4	1	-	-	-
White, both sexes	1	5	6	2	4	5	3	6	2	2	-	-
Male	1	2	1	-	4	2	1	2	1	2	-	-
Female	-	3	5	2	-	3	2	4	1	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	18	22	13	26	18	30	43	24	18	6	2	-
Male	12	9	7	11	11	12	23	10	5	-	-	-
Female	6	13	6	15	7	18	20	14	13	6	2	-
White, both sexes	15	19	13	20	16	28	40	23	18	4	1	-
Male	11	8	7	7	10	12	23	10	5	-	-	-
Female	4	11	6	13	6	16	17	13	13	4	1	-
Black, both sexes	3	3	-	6	2	2	2	1	-	2	1	-
Male	1	1	-	4	1	-	-	-	-	-	-	-
Female	2	2	-	2	1	2	2	1	-	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
North Carolina												
All races, both sexes	11	15	20	26	25	30	34	31	18	6	-	-
Male	4	10	12	10	12	14	12	9	2	1	-	-
Female	7	5	8	16	13	16	22	22	16	5	-	-
White, both sexes	9	9	15	22	15	23	32	22	16	5	-	-
Male	4	7	8	9	7	13	11	6	1	1	-	-
Female	5	2	7	13	8	10	21	16	15	4	-	-
Black, both sexes	2	6	5	3	9	6	2	9	2	1	-	-
Male	-	3	4	1	5	1	1	3	1	-	-	-
Female	2	3	1	2	4	5	1	6	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	1	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	1	-	2	-	4	2	5	1	-	-	-
Male	-	-	-	1	-	1	2	2	-	-	-	-
Female	-	1	-	1	-	3	-	3	1	-	-	-
White, both sexes	-	1	-	2	-	4	2	5	1	-	-	-
Male	-	-	-	1	-	1	2	2	-	-	-	-
Female	-	1	-	1	-	3	-	3	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Ohio												
All races, both sexes	18	15	17	15	27	17	34	29	18	3	-	-
Male	7	9	8	6	6	6	16	8	4	1	-	-
Female	11	6	9	9	21	11	18	21	14	2	-	-
White, both sexes	16	10	15	14	26	15	32	25	15	3	-	-
Male	5	7	8	6	6	6	14	7	4	1	-	-
Female	11	3	7	8	20	9	18	18	11	2	-	-
Black, both sexes	1	5	2	1	1	2	2	4	3	-	-	-
Male	1	2	-	-	-	-	2	1	-	-	-	-
Female	-	3	2	1	1	2	-	3	3	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	5	8	9	14	10	13	12	8	3	2	-	-
Male	3	3	5	8	5	4	4	3	-	-	-	-
Female	2	5	4	6	5	9	8	5	3	2	-	-
White, both sexes	4	6	8	14	10	10	11	7	3	2	-	-
Male	2	3	4	8	5	3	4	2	-	-	-	-
Female	2	3	4	6	5	7	7	5	3	2	-	-
Black, both sexes	-	1	1	-	-	1	1	1	-	-	-	-
Male	-	-	1	-	-	1	-	1	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	2	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	2	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Oregon												
All races, both sexes	9	2	5	6	7	6	11	7	5	1	-	-
Male	4	1	2	2	2	3	5	3	2	1	-	-
Female	5	1	3	4	5	3	6	4	3	-	-	-
White, both sexes	9	2	5	6	6	6	11	7	5	1	-	-
Male	4	1	2	2	1	3	5	3	2	1	-	-
Female	5	1	3	4	5	3	6	4	3	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	22	26	21	37	31	36	57	46	23	4	-	-
Male	14	14	10	18	18	17	22	15	9	1	-	-
Female	8	12	11	19	13	19	35	31	14	3	-	-
White, both sexes	18	20	20	30	26	35	48	41	23	4	-	-
Male	12	10	9	12	15	17	17	14	9	1	-	-
Female	6	10	11	18	11	18	31	27	14	3	-	-
Black, both sexes	4	6	1	7	5	1	9	5	-	-	-	-
Male	2	4	1	6	3	-	5	1	-	-	-	-
Female	2	2	-	1	2	1	4	4	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Rhode Island												
All races, both sexes	1	1	1	1	3	4	2	1	2	2	-	-
Male	1	1	1	-	2	-	-	-	-	-	-	-
Female	-	-	-	1	1	4	2	1	2	2	-	-
White, both sexes	1	1	1	1	3	4	2	1	2	2	-	-
Male	1	1	1	-	2	-	-	-	-	-	-	-
Female	-	-	-	1	1	4	2	1	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	8	14	12	21	6	17	16	18	9	3	-	-
Male	4	11	3	9	2	7	7	7	3	1	-	-
Female	4	3	9	12	4	10	9	11	6	2	-	-
White, both sexes	3	11	7	14	3	13	12	13	5	3	-	-
Male	1	8	2	5	-	4	6	3	1	1	-	-
Female	2	3	5	9	3	9	6	10	4	2	-	-
Black, both sexes	5	3	5	7	3	4	4	5	3	-	-	-
Male	3	3	1	4	2	3	1	4	1	-	-	-
Female	2	-	4	3	1	1	3	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
South Dakota												
All races, both sexes	-	1	2	1	4	4	4	4	1	-	1	-
Male	-	-	2	-	3	4	-	-	-	-	-	-
Female	-	1	-	1	1	-	4	4	1	-	1	-
White, both sexes	-	1	2	1	4	4	4	4	1	-	1	-
Male	-	-	2	-	3	4	-	-	-	-	-	-
Female	-	1	-	1	1	-	4	4	1	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	7	14	17	18	22	27	33	29	6	5	2	-
Male	7	8	8	4	12	14	12	5	2	2	1	-
Female	-	6	9	14	10	13	21	24	4	3	1	-
White, both sexes	5	13	17	13	20	25	28	25	4	5	-	-
Male	5	8	8	3	10	13	10	5	1	2	-	-
Female	-	5	9	10	10	12	18	20	3	3	-	-
Black, both sexes	2	1	-	5	2	1	5	4	2	-	2	-
Male	2	-	-	1	2	-	2	-	1	-	1	-
Female	-	1	-	4	-	1	3	4	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Texas												
All races, both sexes	36	39	49	49	73	66	76	81	54	16	4	-
Male	19	19	30	21	23	32	30	29	15	1	3	-
Female	17	20	19	28	50	34	46	52	39	15	1	-
White, both sexes	27	30	38	39	53	49	64	72	42	14	3	-
Male	15	16	22	18	16	22	26	26	12	1	2	-
Female	12	14	16	21	37	27	38	46	30	13	1	-
Black, both sexes	9	9	11	9	19	17	12	8	12	2	1	-
Male	4	3	8	3	7	10	4	2	3	-	1	-
Female	5	6	3	6	12	7	8	6	9	2	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Utah												
All races, both sexes	6	6	7	6	5	8	5	9	6	1	-	-
Male	3	3	3	2	2	6	1	4	1	-	-	-
Female	3	3	4	4	3	2	4	5	5	1	-	-
White, both sexes	6	5	7	6	5	8	5	9	6	1	-	-
Male	3	2	3	2	2	6	1	4	1	-	-	-
Female	3	3	4	4	3	2	4	5	5	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Vermont												
All races, both sexes	-	-	-	3	1	1	2	3	-	-	-	-
Male	-	-	-	2	1	-	1	1	-	-	-	-
Female	-	-	-	1	-	1	1	2	-	-	-	-
White, both sexes	-	-	-	3	1	1	2	3	-	-	-	-
Male	-	-	-	2	1	-	1	1	-	-	-	-
Female	-	-	-	1	-	1	1	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	12	16	9	28	26	19	29	23	10	4	1	-
Male	8	10	6	14	10	11	11	9	4	1	-	-
Female	4	6	3	14	16	8	18	14	6	3	1	-
White, both sexes	8	10	6	18	21	17	25	19	9	4	1	-
Male	5	5	5	7	10	11	10	6	3	1	-	-
Female	3	5	1	11	11	6	15	13	6	3	1	-
Black, both sexes	4	6	3	9	5	2	4	4	1	-	-	-
Male	3	5	1	6	-	-	1	3	1	-	-	-
Female	1	1	2	3	5	2	3	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Washington												
All races, both sexes	8	7	16	7	12	12	19	14	11	2	-	-
Male	2	4	8	3	4	3	8	4	4	-	-	-
Female	6	3	8	4	8	9	11	10	7	2	-	-
White, both sexes	7	7	16	6	11	11	18	11	11	2	-	-
Male	2	4	8	3	3	3	8	4	4	-	-	-
Female	5	3	8	3	8	8	10	7	7	2	-	-
Black, both sexes	1	-	-	-	1	1	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	1	-	-	-	-
West Virginia												
All races, both sexes	3	3	6	9	6	6	14	8	4	-	-	-
Male	1	2	5	4	2	5	8	2	-	-	-	-
Female	2	1	1	5	4	1	6	6	4	-	-	-
White, both sexes	3	2	6	9	6	6	13	8	3	-	-	-
Male	1	1	5	4	2	5	7	2	-	-	-	-
Female	2	1	1	5	4	1	6	6	3	-	-	-
Black, both sexes	-	1	-	-	-	-	1	-	1	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Wisconsin												
All races, both sexes	6	12	10	18	14	23	33	26	17	3	-	-
Male	4	8	4	7	11	12	14	5	5	1	-	-
Female	2	4	6	11	3	11	19	21	12	2	-	-
White, both sexes	4	11	9	16	13	21	30	26	17	3	-	-
Male	2	8	4	7	11	12	13	5	5	1	-	-
Female	2	3	5	9	2	9	17	21	12	2	-	-
Black, both sexes	2	1	1	1	1	1	1	-	-	-	-	-
Male	2	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	1	1	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Wyoming												
All races, both sexes	1	1	2	1	3	1	2	5	1	-	-	-
Male	-	1	1	1	2	1	1	1	1	-	-	-
Female	1	-	1	-	1	-	1	4	-	-	-	-
White, both sexes	1	1	1	1	3	1	2	5	1	-	-	-
Male	-	1	-	1	2	1	1	1	1	-	-	-
Female	1	-	1	-	1	-	1	4	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
United States												
All races, both sexes	5,455	96	24	3	11	18	21	43	50	81	94	190
Male	1,862	59	8	2	3	8	4	16	13	28	46	72
Female	3,593	37	16	1	8	10	17	27	37	53	48	118
White, both sexes	4,450	58	15	3	8	16	18	25	25	49	61	126
Male	1,543	41	2	2	3	7	4	8	5	20	28	50
Female	2,907	17	13	1	5	9	14	17	20	29	33	76
Black, both sexes	875	29	5	-	2	2	3	15	19	28	31	60
Male	272	13	4	-	-	1	-	8	7	8	17	20
Female	603	16	1	-	2	1	3	7	12	20	14	40
American Indian 1/, both sexes	30	4	1	-	-	-	-	2	3	-	1	2
Male	13	3	1	-	-	-	-	-	-	-	-	1
Female	17	1	-	-	-	-	-	2	3	-	1	1
Asian or Pacific Islander 2/, both sexes	100	5	3	-	1	-	-	1	3	4	1	2
Male	34	2	1	-	-	-	-	-	1	-	1	1
Female	66	3	2	-	1	-	-	1	2	4	-	1
Alabama												
All races, both sexes	67	2	1	-	-	-	-	1	-	1	1	2
Male	29	-	1	-	-	-	-	-	-	1	1	2
Female	38	2	-	-	-	-	-	1	-	-	-	-
White, both sexes	38	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	28	2	1	-	-	-	-	1	-	1	1	2
Male	15	-	1	-	-	-	-	-	-	1	1	2
Female	13	2	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Alaska												
All races, both sexes	18	-	1	-	-	1	-	1	2	-	-	-
Male	5	-	1	-	-	1	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	1	2	-	-	-
White, both sexes	12	-	-	-	-	1	-	-	-	-	-	-
Male	3	-	-	-	-	1	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	1	-	-	-	-	1	2	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	84	3	-	-	-	-	-	3	1	1	7	-
Male	31	1	-	-	-	-	-	1	-	1	4	-
Female	53	2	-	-	-	-	-	2	1	-	3	-
White, both sexes	71	1	-	-	-	-	-	3	-	1	5	-
Male	26	-	-	-	-	-	-	1	-	1	3	-
Female	45	1	-	-	-	-	-	2	-	-	2	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	2	-	-	-	-	-	-	1	-	1	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	4	1	-	-	-	-	-	-	1	-	1	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Arkansas												
All races, both sexes	73	2	-	-	-	-	-	2	-	-	1	1
Male	27	2	-	-	-	-	-	1	-	-	1	-
Female	46	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	63	-	-	-	-	-	-	1	-	-	-	1
Male	22	-	-	-	-	-	-	1	-	-	-	-
Female	41	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	10	2	-	-	-	-	-	1	-	-	1	-
Male	5	2	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	478	5	7	1	2	4	1	2	13	8	10	26
Male	179	4	2	1	1	2	-	-	3	2	7	6
Female	299	1	5	-	1	2	1	2	10	6	3	20
White, both sexes	387	4	6	1	2	4	-	2	9	5	9	20
Male	142	3	1	1	1	2	-	-	1	2	6	4
Female	245	1	5	-	1	2	-	2	8	3	3	16
Black, both sexes	55	-	1	-	-	-	1	-	2	-	1	6
Male	23	-	1	-	-	-	-	-	1	-	1	2
Female	32	-	-	-	-	-	1	-	1	-	-	4
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	33	1	-	-	-	-	-	-	2	3	-	-
Male	11	1	-	-	-	-	-	-	1	-	-	-
Female	22	-	-	-	-	-	-	-	1	3	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Colorado												
All races, both sexes	89	1	1	-	-	-	-	-	1	-	2	1
Male	30	1	-	-	-	-	-	-	-	-	-	1
Female	59	-	1	-	-	-	-	-	1	-	2	-
White, both sexes	81	1	1	-	-	-	-	-	1	-	2	1
Male	27	1	-	-	-	-	-	-	-	-	-	1
Female	54	-	1	-	-	-	-	-	1	-	2	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	65	1	-	-	-	-	-	-	2	1	-	2
Male	24	1	-	-	-	-	-	-	1	-	-	1
Female	41	-	-	-	-	-	-	-	1	1	-	1
White, both sexes	59	1	-	-	-	-	-	-	1	1	-	1
Male	22	1	-	-	-	-	-	-	1	-	-	-
Female	37	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	6	-	-	-	-	-	-	-	1	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Delaware												
All races, both sexes	19	-	1	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	1	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	16	-	-	-	-	-	1	-	-	-	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	1	-	-	-	-	2
White, both sexes	4	-	-	-	-	-	1	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	1	-	-	-	-	1
Black, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Florida												
All races, both sexes	281	7	-	1	-	1	2	1	3	-	8	10
Male	97	2	-	-	-	1	-	-	1	-	2	4
Female	184	5	-	1	-	-	2	1	2	-	6	6
White, both sexes	228	4	-	1	-	1	1	1	-	-	6	6
Male	87	2	-	-	-	1	-	-	-	-	1	3
Female	141	2	-	1	-	-	1	1	-	-	5	3
Black, both sexes	50	2	-	-	-	-	1	-	3	-	2	4
Male	8	-	-	-	-	-	-	-	1	-	1	1
Female	42	2	-	-	-	-	1	-	2	-	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	1	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	201	3	1	-	-	1	-	4	1	5	4	13
Male	65	1	-	-	-	-	-	3	-	1	1	3
Female	136	2	1	-	-	1	-	1	1	4	3	10
White, both sexes	125	-	-	-	-	1	-	1	-	3	-	3
Male	42	-	-	-	-	-	-	1	-	1	-	-
Female	83	-	-	-	-	1	-	-	-	2	-	3
Black, both sexes	75	3	1	-	-	-	-	3	1	2	4	10
Male	23	1	-	-	-	-	-	2	-	-	1	3
Female	52	2	1	-	-	-	-	1	1	2	3	7
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Hawaii												
All races, both sexes	21	1	1	-	-	-	-	-	1	1	-	1
Male	9	1	1	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	1	1	-	1
White, both sexes	5	-	1	-	-	-	-	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	1	-	-	-	-	-	-	1	1	-	1
Male	7	1	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	1	1	-	1
Idaho												
All races, both sexes	40	-	-	-	-	-	-	-	-	1	-	2
Male	14	-	-	-	-	-	-	-	-	1	-	1
Female	26	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	39	-	-	-	-	-	-	-	-	1	-	2
Male	14	-	-	-	-	-	-	-	-	1	-	1
Female	25	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Illinois												
All races, both sexes	228	2	1	-	-	-	1	2	1	6	6	9
Male	70	-	-	-	-	-	-	1	-	2	2	7
Female	158	2	1	-	-	-	1	1	1	4	4	2
White, both sexes	181	1	-	-	-	-	1	-	-	2	3	7
Male	54	-	-	-	-	-	-	-	-	-	1	5
Female	127	1	-	-	-	-	1	-	-	2	2	2
Black, both sexes	42	-	-	-	-	-	-	1	1	4	3	2
Male	16	-	-	-	-	-	-	1	-	2	1	2
Female	26	-	-	-	-	-	-	-	1	2	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	1	1	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	1	1	-	-	-	-	1	-	-	-	-
Indiana												
All races, both sexes	134	2	2	-	2	-	-	2	-	2	4	4
Male	49	2	-	-	1	-	-	1	-	-	3	2
Female	85	-	2	-	1	-	-	1	-	2	1	2
White, both sexes	116	2	2	-	2	-	-	-	-	1	3	3
Male	42	2	-	-	1	-	-	-	-	-	2	2
Female	74	-	2	-	1	-	-	-	-	1	1	1
Black, both sexes	18	-	-	-	-	-	-	2	-	1	1	1
Male	7	-	-	-	-	-	-	1	-	-	1	-
Female	11	-	-	-	-	-	-	1	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Iowa												
All races, both sexes	66	-	-	-	-	-	-	-	-	1	-	1
Male	23	-	-	-	-	-	-	-	-	1	-	1
Female	43	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	66	-	-	-	-	-	-	-	-	1	-	1
Male	23	-	-	-	-	-	-	-	-	1	-	1
Female	43	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	70	2	-	-	-	1	1	-	-	1	2	2
Male	26	1	-	-	-	-	1	-	-	-	1	2
Female	44	1	-	-	-	1	-	-	-	1	1	-
White, both sexes	66	1	-	-	-	1	1	-	-	1	2	2
Male	25	-	-	-	-	-	1	-	-	-	1	2
Female	41	1	-	-	-	1	-	-	-	1	1	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Kentucky												
All races, both sexes	78	-	-	-	-	-	-	-	1	1	1	4
Male	29	-	-	-	-	-	-	-	-	1	-	2
Female	49	-	-	-	-	-	-	-	1	-	1	2
White, both sexes	72	-	-	-	-	-	-	-	1	1	-	4
Male	26	-	-	-	-	-	-	-	-	1	-	2
Female	46	-	-	-	-	-	-	-	1	-	-	2
Black, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	82	2	-	-	1	1	-	1	-	2	1	3
Male	25	-	-	-	1	-	-	-	-	-	1	1
Female	57	2	-	-	-	1	-	1	-	2	-	2
White, both sexes	50	1	-	-	1	1	-	1	-	1	-	2
Male	15	-	-	-	1	-	-	-	-	-	-	1
Female	35	1	-	-	-	1	-	1	-	1	-	1
Black, both sexes	32	1	-	-	-	-	-	-	-	1	1	1
Male	10	-	-	-	-	-	-	-	-	-	1	-
Female	22	1	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Maine												
All races, both sexes	41	-	-	-	-	-	1	-	-	1	-	3
Male	19	-	-	-	-	-	-	-	-	1	-	2
Female	22	-	-	-	-	-	1	-	-	-	-	1
White, both sexes	41	-	-	-	-	-	1	-	-	1	-	3
Male	19	-	-	-	-	-	-	-	-	1	-	2
Female	22	-	-	-	-	-	1	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	103	2	-	-	-	-	-	1	1	2	1	2
Male	26	1	-	-	-	-	-	-	-	-	-	2
Female	77	1	-	-	-	-	-	1	1	2	1	-
White, both sexes	65	1	-	-	-	-	-	-	-	1	-	1
Male	19	1	-	-	-	-	-	-	-	-	-	1
Female	46	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	35	1	-	-	-	-	-	1	1	1	1	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	29	1	-	-	-	-	-	1	1	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Massachusetts												
All races, both sexes	111	1	-	-	1	-	-	1	-	-	1	4
Male	30	-	-	-	-	-	-	1	-	-	-	1
Female	81	1	-	-	1	-	-	-	-	-	1	3
White, both sexes	101	-	-	-	1	-	-	1	-	-	1	4
Male	25	-	-	-	-	-	-	1	-	-	-	1
Female	76	-	-	-	1	-	-	-	-	-	1	3
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	168	3	1	-	-	-	2	2	4	2	3	5
Male	47	1	1	-	-	-	-	1	2	1	1	-
Female	121	2	-	-	-	-	2	1	2	1	2	5
White, both sexes	123	1	-	-	-	-	1	2	-	-	1	4
Male	39	-	-	-	-	-	-	1	-	-	1	-
Female	84	1	-	-	-	-	1	1	-	-	-	4
Black, both sexes	41	2	-	-	-	-	1	-	4	2	2	1
Male	7	1	-	-	-	-	-	-	2	1	-	-
Female	34	1	-	-	-	-	1	-	2	1	2	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Minnesota												
All races, both sexes	142	1	-	-	-	-	-	1	-	1	1	2
Male	47	1	-	-	-	-	-	-	-	-	1	1
Female	95	-	-	-	-	-	-	1	-	1	-	1
White, both sexes	135	1	-	-	-	-	-	1	-	1	1	1
Male	46	1	-	-	-	-	-	-	-	-	1	1
Female	89	-	-	-	-	-	-	1	-	1	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	62	3	-	-	-	2	1	-	-	-	-	1
Male	17	2	-	-	-	-	-	-	-	-	-	1
Female	45	1	-	-	-	2	1	-	-	-	-	-
White, both sexes	35	1	-	-	-	1	1	-	-	-	-	1
Male	10	1	-	-	-	-	-	-	-	-	-	1
Female	25	-	-	-	-	1	1	-	-	-	-	-
Black, both sexes	27	2	-	-	-	1	-	-	-	-	-	-
Male	7	1	-	-	-	-	-	-	-	-	-	-
Female	20	1	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Missouri												
All races, both sexes	110	-	-	-	-	1	1	2	2	3	2	2
Male	44	-	-	-	-	1	1	1	1	-	-	1
Female	66	-	-	-	-	-	-	1	1	3	2	1
White, both sexes	92	-	-	-	-	-	1	2	1	1	2	2
Male	39	-	-	-	-	-	1	1	-	-	-	1
Female	53	-	-	-	-	-	-	1	1	1	2	1
Black, both sexes	18	-	-	-	-	1	-	-	1	2	-	-
Male	5	-	-	-	-	1	-	-	1	-	-	-
Female	13	-	-	-	-	-	-	-	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	20	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Nebraska												
All races, both sexes	60	-	-	-	-	-	-	-	1	-	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	40	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	57	-	-	-	-	-	-	-	1	-	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	38	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	26	1	-	-	1	-	-	2	-	2	1	1
Male	7	1	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	1	-	-	2	-	2	1	1
White, both sexes	19	-	-	-	-	-	-	1	-	1	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	1	-	1	-	1
Black, both sexes	5	1	-	-	-	-	-	1	-	1	1	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	1	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
New Hampshire												
All races, both sexes	27	-	-	-	-	-	-	-	-	-	1	-
Male	12	-	-	-	-	-	-	-	-	-	1	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	26	-	-	-	-	-	-	-	-	-	1	-
Male	12	-	-	-	-	-	-	-	-	-	1	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	154	4	-	-	-	-	2	-	-	2	2	3
Male	61	4	-	-	-	-	-	-	-	1	1	2
Female	93	-	-	-	-	-	2	-	-	1	1	1
White, both sexes	125	2	-	-	-	-	2	-	-	1	1	-
Male	45	2	-	-	-	-	-	-	-	1	-	-
Female	80	-	-	-	-	-	2	-	-	-	1	-
Black, both sexes	23	2	-	-	-	-	-	-	-	1	1	3
Male	13	2	-	-	-	-	-	-	-	-	1	2
Female	10	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
New Mexico												
All races, both sexes	43	1	-	-	-	-	-	1	-	2	-	3
Male	16	1	-	-	-	-	-	-	-	1	-	3
Female	27	-	-	-	-	-	-	1	-	1	-	-
White, both sexes	39	1	-	-	-	-	-	-	-	2	-	2
Male	14	1	-	-	-	-	-	-	-	1	-	2
Female	25	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	1	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	291	13	1	-	-	-	1	1	-	3	4	7
Male	97	8	-	-	-	-	-	1	-	2	2	3
Female	194	5	1	-	-	-	1	-	-	1	2	4
White, both sexes	234	8	-	-	-	-	1	-	-	2	2	5
Male	83	6	-	-	-	-	-	-	-	2	1	2
Female	151	2	-	-	-	-	1	-	-	-	1	3
Black, both sexes	49	5	-	-	-	-	-	1	-	1	2	2
Male	12	2	-	-	-	-	-	1	-	-	1	1
Female	37	3	-	-	-	-	-	-	-	1	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	1	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
North Carolina												
All races, both sexes	214	5	3	-	1	-	-	1	-	5	1	9
Male	77	4	2	-	-	-	-	1	-	2	1	-
Female	137	1	1	-	1	-	-	-	-	3	-	9
White, both sexes	143	3	1	-	-	-	-	-	-	3	-	2
Male	52	3	-	-	-	-	-	-	-	1	-	-
Female	91	-	1	-	-	-	-	-	-	2	-	2
Black, both sexes	71	2	2	-	1	-	-	1	-	2	1	7
Male	25	1	2	-	-	-	-	1	-	1	1	-
Female	46	1	-	-	1	-	-	-	-	1	-	7
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	11	1	-	-	-	-	-	-	-	-	-	-
Male	4	1	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	1	-	-	-	-	-	-	-	-	-	-
Male	4	1	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Ohio												
All races, both sexes	221	1	-	-	-	-	1	2	4	2	2	9
Male	63	-	-	-	-	-	-	1	-	-	2	2
Female	158	1	-	-	-	-	1	1	4	2	-	7
White, both sexes	187	-	-	-	-	-	1	1	3	1	1	7
Male	56	-	-	-	-	-	-	-	-	-	1	2
Female	131	-	-	-	-	-	1	1	3	1	-	5
Black, both sexes	34	1	-	-	-	-	-	1	1	1	1	2
Male	7	-	-	-	-	-	-	1	-	-	1	-
Female	27	1	-	-	-	-	-	-	1	1	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	59	1	-	-	-	-	-	1	1	1	-	4
Male	14	1	-	-	-	-	-	-	-	-	-	-
Female	45	-	-	-	-	-	-	1	1	1	-	4
White, both sexes	49	-	-	-	-	-	-	1	1	1	-	2
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	38	-	-	-	-	-	-	1	1	1	-	2
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	6	1	-	-	-	-	-	-	-	-	-	1
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Oregon												
All races, both sexes	99	2	-	-	-	-	1	1	-	-	2	4
Male	39	1	-	-	-	-	1	1	-	-	1	2
Female	60	1	-	-	-	-	-	-	-	-	1	2
White, both sexes	96	2	-	-	-	-	1	-	-	-	2	3
Male	37	1	-	-	-	-	1	-	-	-	1	1
Female	59	1	-	-	-	-	-	-	-	-	1	2
Black, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	304	2	1	-	-	-	1	2	3	3	1	8
Male	100	1	-	-	-	-	-	1	2	2	-	5
Female	204	1	1	-	-	-	1	1	1	1	1	3
White, both sexes	269	2	1	-	-	-	1	2	2	1	-	5
Male	87	1	-	-	-	-	-	1	1	1	-	3
Female	182	1	1	-	-	-	1	1	1	-	-	2
Black, both sexes	34	-	-	-	-	-	-	-	1	2	1	3
Male	13	-	-	-	-	-	-	-	1	1	-	2
Female	21	-	-	-	-	-	-	-	-	1	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Rhode Island												
All races, both sexes	31	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	30	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	103	3	1	-	-	-	-	3	1	4	1	4
Male	32	3	-	-	-	-	-	-	1	2	1	3
Female	71	-	1	-	-	-	-	3	-	2	-	1
White, both sexes	73	1	1	-	-	-	-	3	-	1	-	2
Male	23	1	-	-	-	-	-	-	-	1	-	2
Female	50	-	1	-	-	-	-	3	-	-	-	-
Black, both sexes	28	2	-	-	-	-	-	-	1	3	1	2
Male	8	2	-	-	-	-	-	-	1	1	1	1
Female	20	-	-	-	-	-	-	-	-	2	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
South Dakota												
All races, both sexes	8	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	93	2	-	-	-	-	2	-	-	1	3	1
Male	33	1	-	-	-	-	1	-	-	1	1	-
Female	60	1	-	-	-	-	1	-	-	-	2	1
White, both sexes	70	2	-	-	-	-	2	-	-	1	2	1
Male	26	1	-	-	-	-	1	-	-	1	1	-
Female	44	1	-	-	-	-	1	-	-	-	1	1
Black, both sexes	23	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Texas												
All races, both sexes	342	7	1	-	2	1	1	1	4	11	8	20
Male	122	5	-	-	-	-	-	-	2	3	5	7
Female	220	2	1	-	2	1	1	1	2	8	3	13
White, both sexes	281	7	1	-	2	1	1	1	3	9	5	15
Male	104	5	-	-	-	-	-	-	2	3	2	5
Female	177	2	1	-	2	1	1	1	1	6	3	10
Black, both sexes	53	-	-	-	-	-	-	-	1	2	2	5
Male	16	-	-	-	-	-	-	-	-	-	2	2
Female	37	-	-	-	-	-	-	-	1	2	-	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	49	-	-	-	-	1	-	-	1	1	1	1
Male	16	-	-	-	-	1	-	-	-	-	1	1
Female	33	-	-	-	-	-	-	-	1	1	-	-
White, both sexes	47	-	-	-	-	1	-	-	1	1	1	1
Male	16	-	-	-	-	1	-	-	-	-	1	1
Female	31	-	-	-	-	-	-	-	1	1	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Vermont												
All races, both sexes	17	1	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	10	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	1	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	10	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	138	5	-	-	1	-	-	-	1	2	1	5
Male	41	4	-	-	-	-	-	-	-	1	-	-
Female	97	1	-	-	1	-	-	-	1	1	1	5
White, both sexes	96	4	-	-	-	-	-	-	-	1	1	1
Male	30	4	-	-	-	-	-	-	-	-	-	-
Female	66	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	42	1	-	-	1	-	-	-	1	1	-	4
Male	11	-	-	-	-	-	-	-	-	1	-	-
Female	31	1	-	-	1	-	-	-	1	-	-	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Washington												
All races, both sexes	127	1	-	1	-	2	-	2	1	-	7	2
Male	45	1	-	1	-	1	-	1	-	-	3	1
Female	82	-	-	-	-	1	-	1	1	-	4	1
White, both sexes	122	1	-	1	-	2	-	1	1	-	6	2
Male	43	1	-	1	-	1	-	1	-	-	2	1
Female	79	-	-	-	-	1	-	-	1	-	4	1
Black, both sexes	3	-	-	-	-	-	-	1	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	41	1	-	-	-	2	1	-	-	1	1	4
Male	16	-	-	-	-	1	-	-	-	1	1	2
Female	25	1	-	-	-	1	1	-	-	-	-	2
White, both sexes	39	1	-	-	-	2	1	-	-	1	1	4
Male	15	-	-	-	-	1	-	-	-	1	1	2
Female	24	1	-	-	-	1	1	-	-	-	-	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Wisconsin												
All races, both sexes	116	1	-	-	-	-	-	-	-	-	3	2
Male	47	1	-	-	-	-	-	-	-	-	1	-
Female	69	-	-	-	-	-	-	-	-	-	2	2
White, both sexes	108	1	-	-	-	-	-	-	-	-	3	2
Male	43	1	-	-	-	-	-	-	-	-	1	-
Female	65	-	-	-	-	-	-	-	-	-	2	2
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	14	1	-	-	-	-	-	-	-	-	-	-
Male	4	1	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	1	-	-	-	-	-	-	-	-	-	-
Male	4	1	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
United States												
All races, both sexes	243	325	371	430	536	699	845	747	464	146	18	-
Male	95	130	145	169	212	254	273	191	98	32	4	-
Female	148	195	226	261	324	445	572	556	366	114	14	-
White, both sexes	167	234	294	328	440	585	736	687	427	136	12	-
Male	69	103	121	136	183	220	246	174	91	29	1	-
Female	98	131	173	192	257	365	490	513	336	107	11	-
Black, both sexes	70	78	70	92	80	101	96	47	32	9	6	-
Male	24	21	22	27	22	28	24	14	6	3	3	-
Female	46	57	48	65	58	73	72	33	26	6	3	-
American Indian 1/, both sexes	3	3	2	2	3	2	1	1	-	-	-	-
Male	1	1	-	2	1	1	1	1	-	-	-	-
Female	2	2	2	-	2	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	10	5	8	13	11	12	12	5	1	-	-
Male	1	5	2	4	6	5	2	2	1	-	-	-
Female	2	5	3	4	7	6	10	10	4	1	-	-
Alabama												
All races, both sexes	4	5	3	8	10	13	6	9	-	1	-	-
Male	3	3	2	3	3	5	2	3	-	-	-	-
Female	1	2	1	5	7	8	4	6	-	1	-	-
White, both sexes	2	2	2	5	7	8	4	7	-	1	-	-
Male	1	-	1	3	2	4	1	2	-	-	-	-
Female	1	2	1	2	5	4	3	5	-	1	-	-
Black, both sexes	2	3	1	2	3	5	2	2	-	-	-	-
Male	2	3	1	-	1	1	1	1	-	-	-	-
Female	-	-	-	2	2	4	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Alaska												
All races, both sexes	1	2	-	-	2	3	2	-	3	-	-	-
Male	-	-	-	-	2	1	-	-	-	-	-	-
Female	1	2	-	-	-	2	2	-	3	-	-	-
White, both sexes	1	2	-	-	2	2	1	-	3	-	-	-
Male	-	-	-	-	2	-	-	-	-	-	-	-
Female	1	2	-	-	-	2	1	-	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Arizona												
All races, both sexes	4	5	7	7	11	10	9	6	9	1	-	-
Male	2	4	3	2	2	4	1	2	3	1	-	-
Female	2	1	4	5	9	6	8	4	6	-	-	-
White, both sexes	2	3	4	7	10	10	9	6	9	1	-	-
Male	1	3	2	2	2	4	1	2	3	1	-	-
Female	1	-	2	5	8	6	8	4	6	-	-	-
Black, both sexes	1	2	1	-	-	-	-	-	-	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Arkansas												
All races, both sexes	3	8	7	9	7	7	15	3	6	1	1	-
Male	2	5	2	3	2	2	5	-	2	-	-	-
Female	1	3	5	6	5	5	10	3	4	1	1	-
White, both sexes	3	8	6	6	7	5	15	3	6	1	1	-
Male	2	5	2	2	2	1	5	-	2	-	-	-
Female	1	3	4	4	5	4	10	3	4	1	1	-
Black, both sexes	-	-	1	3	-	2	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	1	2	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	33	28	28	31	47	55	71	58	33	14	1	-
Male	14	15	10	17	20	20	26	21	6	1	1	-
Female	19	13	18	14	27	35	45	37	27	13	-	-
White, both sexes	26	23	22	17	37	46	63	50	27	14	-	-
Male	11	13	7	9	16	17	24	19	4	1	-	-
Female	15	10	15	8	21	29	39	31	23	13	-	-
Black, both sexes	6	3	3	8	4	7	3	4	5	-	1	-
Male	3	2	1	4	1	2	1	1	2	-	1	-
Female	3	1	2	4	3	5	2	3	3	-	-	-
American Indian 1/, both sexes	-	-	-	2	1	-	-	-	-	-	-	-
Male	-	-	-	2	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	3	4	5	2	5	4	1	-	-	-
Male	-	-	2	2	2	1	1	1	-	-	-	-
Female	1	2	1	2	3	1	4	3	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Colorado												
All races, both sexes	4	5	6	7	7	17	17	9	9	2	-	-
Male	2	-	3	3	2	7	6	2	2	1	-	-
Female	2	5	3	4	5	10	11	7	7	1	-	-
White, both sexes	4	4	6	6	5	14	16	9	9	2	-	-
Male	2	-	3	3	-	6	6	2	2	1	-	-
Female	2	4	3	3	5	8	10	7	7	1	-	-
Black, both sexes	-	1	-	1	2	2	1	-	-	-	-	-
Male	-	-	-	-	2	-	-	-	-	-	-	-
Female	-	1	-	1	-	2	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	3	2	5	5	12	13	12	2	5	-	-
Male	-	1	-	4	2	1	6	5	1	1	-	-
Female	-	2	2	1	3	11	7	7	1	4	-	-
White, both sexes	-	2	2	4	4	12	13	12	2	4	-	-
Male	-	1	-	3	2	1	6	5	1	1	-	-
Female	-	1	2	1	2	11	7	7	1	3	-	-
Black, both sexes	-	1	-	1	1	-	-	-	-	1	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Delaware												
All races, both sexes	1	1	3	1	2	1	2	4	1	1	1	-
Male	1	-	-	1	1	-	1	-	-	1	1	-
Female	-	1	3	-	1	1	1	4	1	-	-	-
White, both sexes	1	1	1	1	1	1	2	4	1	1	-	-
Male	1	-	-	1	1	-	1	-	-	1	-	-
Female	-	1	1	-	-	1	1	4	1	-	-	-
Black, both sexes	-	-	2	-	1	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	2	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	-	2	2	1	4	3	-	-	-	-
Male	1	-	-	1	-	-	1	1	-	-	-	-
Female	-	-	-	1	2	1	3	2	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	2	2	1	4	2	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	1	2	1	3	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Florida												
All races, both sexes	8	19	9	19	32	40	43	47	21	9	1	-
Male	-	8	4	7	13	10	20	15	8	1	1	-
Female	8	11	5	12	19	30	23	32	13	8	-	-
White, both sexes	5	10	5	18	27	34	36	44	19	9	1	-
Male	-	4	3	6	13	10	20	14	8	1	1	-
Female	5	6	2	12	14	24	16	30	11	8	-	-
Black, both sexes	3	8	4	-	5	6	7	3	2	-	-	-
Male	-	3	1	-	-	-	-	1	-	-	-	-
Female	3	5	3	-	5	6	7	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	13	9	16	16	16	28	37	15	17	1	1	-
Male	5	3	8	5	7	11	10	5	2	-	-	-
Female	8	6	8	11	9	17	27	10	15	1	1	-
White, both sexes	6	5	10	10	13	17	29	12	14	1	-	-
Male	2	3	5	4	6	5	8	5	2	-	-	-
Female	4	2	5	6	7	12	21	7	12	1	-	-
Black, both sexes	7	3	6	6	3	11	8	3	3	-	1	-
Male	3	-	3	1	1	6	2	-	-	-	-	-
Female	4	3	3	5	2	5	6	3	3	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Hawaii												
All races, both sexes	-	1	1	-	3	2	4	2	3	-	-	-
Male	-	1	-	-	2	1	2	1	-	-	-	-
Female	-	-	1	-	1	1	2	1	3	-	-	-
White, both sexes	-	-	1	-	-	-	1	-	2	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	3	2	3	2	1	-	-	-
Male	-	1	-	-	2	1	1	1	-	-	-	-
Female	-	-	-	-	1	1	2	1	1	-	-	-
Idaho												
All races, both sexes	1	2	4	5	4	2	5	6	5	3	-	-
Male	-	2	-	1	1	1	3	1	2	1	-	-
Female	1	-	4	4	3	1	2	5	3	2	-	-
White, both sexes	1	2	4	5	3	2	5	6	5	3	-	-
Male	-	2	-	1	1	1	3	1	2	1	-	-
Female	1	-	4	4	2	1	2	5	3	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Illinois												
All races, both sexes	6	11	14	17	25	32	37	28	25	4	1	-
Male	2	2	5	5	12	11	9	8	2	1	1	-
Female	4	9	9	12	13	21	28	20	23	3	-	-
White, both sexes	4	6	13	12	20	27	31	25	25	4	-	-
Male	2	2	5	5	8	11	5	7	2	1	-	-
Female	2	4	8	7	12	16	26	18	23	3	-	-
Black, both sexes	2	5	1	5	5	4	6	2	-	-	1	-
Male	-	-	-	-	4	-	4	1	-	-	1	-
Female	2	5	1	5	1	4	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Indiana												
All races, both sexes	5	6	14	19	12	20	21	14	5	-	-	-
Male	2	2	5	11	6	9	2	2	1	-	-	-
Female	3	4	9	8	6	11	19	12	4	-	-	-
White, both sexes	4	4	13	17	9	18	19	14	5	-	-	-
Male	1	2	5	10	4	8	2	2	1	-	-	-
Female	3	2	8	7	5	10	17	12	4	-	-	-
Black, both sexes	1	2	1	2	3	2	2	-	-	-	-	-
Male	1	-	-	1	2	1	-	-	-	-	-	-
Female	-	2	1	1	1	1	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Iowa												
All races, both sexes	3	5	2	7	9	6	11	11	4	6	-	-
Male	1	1	-	3	2	3	3	5	1	2	-	-
Female	2	4	2	4	7	3	8	6	3	4	-	-
White, both sexes	3	5	2	7	9	6	11	11	4	6	-	-
Male	1	1	-	3	2	3	3	5	1	2	-	-
Female	2	4	2	4	7	3	8	6	3	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	5	3	4	5	4	15	8	6	8	2	1	-
Male	3	2	2	2	1	5	3	1	2	-	-	-
Female	2	1	2	3	3	10	5	5	6	2	1	-
White, both sexes	5	3	4	4	4	13	8	6	8	2	1	-
Male	3	2	2	2	1	5	3	1	2	-	-	-
Female	2	1	2	2	3	8	5	5	6	2	1	-
Black, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Kentucky												
All races, both sexes	3	8	7	3	11	8	10	13	6	1	1	-
Male	2	4	2	2	6	1	5	1	3	-	-	-
Female	1	4	5	1	5	7	5	12	3	1	1	-
White, both sexes	2	6	7	2	11	7	10	13	6	1	1	-
Male	1	3	2	1	6	1	5	1	3	-	-	-
Female	1	3	5	1	5	6	5	12	3	1	1	-
Black, both sexes	1	2	-	1	-	1	-	-	-	-	-	-
Male	1	1	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	8	7	8	8	7	7	17	5	4	-	-	-
Male	5	3	-	3	3	1	6	1	-	-	-	-
Female	3	4	8	5	4	6	11	4	4	-	-	-
White, both sexes	4	5	3	3	5	4	12	4	3	-	-	-
Male	3	2	-	1	2	1	3	1	-	-	-	-
Female	1	3	3	2	3	3	9	3	3	-	-	-
Black, both sexes	4	2	5	5	2	3	5	1	1	-	-	-
Male	2	1	-	2	1	-	3	-	-	-	-	-
Female	2	1	5	3	1	3	2	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Maine												
All races, both sexes	1	1	1	6	5	10	2	7	2	1	-	-
Male	-	-	1	3	2	7	-	2	1	-	-	-
Female	1	1	-	3	3	3	2	5	1	1	-	-
White, both sexes	1	1	1	6	5	10	2	7	2	1	-	-
Male	-	-	1	3	2	7	-	2	1	-	-	-
Female	1	1	-	3	3	3	2	5	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	9	4	9	11	6	11	19	14	9	2	-	-
Male	4	1	6	1	1	5	2	2	-	1	-	-
Female	5	3	3	10	5	6	17	12	9	1	-	-
White, both sexes	6	1	8	6	4	6	11	12	8	-	-	-
Male	3	-	6	-	1	3	2	2	-	-	-	-
Female	3	1	2	6	3	3	9	10	8	-	-	-
Black, both sexes	3	2	1	5	2	5	7	1	1	2	-	-
Male	1	-	-	1	-	2	-	-	-	1	-	-
Female	2	2	1	4	2	3	7	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	1	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Massachusetts												
All races, both sexes	5	9	5	8	6	15	26	17	9	2	1	-
Male	-	2	2	4	3	5	9	2	1	-	-	-
Female	5	7	3	4	3	10	17	15	8	2	1	-
White, both sexes	5	7	4	5	5	15	24	17	9	2	1	-
Male	-	1	1	2	3	5	8	2	1	-	-	-
Female	5	6	3	3	2	10	16	15	8	2	1	-
Black, both sexes	-	1	1	3	-	-	2	-	-	-	-	-
Male	-	-	1	2	-	-	1	-	-	-	-	-
Female	-	1	-	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Michigan												
All races, both sexes	5	4	15	20	16	27	22	19	12	5	1	-
Male	4	2	5	6	2	9	6	3	2	1	-	-
Female	1	2	10	14	14	18	16	16	10	4	1	-
White, both sexes	4	3	10	12	11	24	16	18	12	3	1	-
Male	4	2	4	5	2	9	5	3	2	1	-	-
Female	-	1	6	7	9	15	11	15	10	2	1	-
Black, both sexes	1	1	3	8	5	3	6	-	-	2	-	-
Male	-	-	1	1	-	-	1	-	-	-	-	-
Female	1	1	2	7	5	3	5	-	-	2	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Minnesota												
All races, both sexes	3	1	13	10	8	19	25	37	18	2	-	-
Male	2	-	5	3	4	8	7	10	5	-	-	-
Female	1	1	8	7	4	11	18	27	13	2	-	-
White, both sexes	2	1	12	9	8	18	24	36	18	2	-	-
Male	2	-	5	2	4	8	7	10	5	-	-	-
Female	-	1	7	7	4	10	17	26	13	2	-	-
Black, both sexes	1	-	1	1	-	1	1	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	9	1	7	2	11	9	4	5	7	-	-	-
Male	4	-	2	-	1	3	1	1	2	-	-	-
Female	5	1	5	2	10	6	3	4	5	-	-	-
White, both sexes	4	-	3	1	8	4	3	4	4	-	-	-
Male	1	-	1	-	1	1	1	1	2	-	-	-
Female	3	-	2	1	7	3	2	3	2	-	-	-
Black, both sexes	5	1	4	1	3	5	1	1	3	-	-	-
Male	3	-	1	-	-	2	-	-	-	-	-	-
Female	2	1	3	1	3	3	1	1	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Missouri												
All races, both sexes	6	7	11	12	10	15	10	10	9	7	-	-
Male	3	4	7	7	4	6	3	2	1	2	-	-
Female	3	3	4	5	6	9	7	8	8	5	-	-
White, both sexes	5	3	11	11	10	13	8	9	6	7	-	-
Male	2	3	7	7	4	6	3	2	-	2	-	-
Female	3	-	4	4	6	7	5	7	6	5	-	-
Black, both sexes	1	4	-	1	-	2	2	1	3	-	-	-
Male	1	1	-	-	-	-	-	-	1	-	-	-
Female	-	3	-	1	-	2	2	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	2	-	-	3	2	2	5	5	-	-	-
Male	-	-	-	-	2	2	2	1	1	-	-	-
Female	-	2	-	-	1	-	-	4	4	-	-	-
White, both sexes	-	2	-	-	3	2	2	5	5	-	-	-
Male	-	-	-	-	2	2	2	1	1	-	-	-
Female	-	2	-	-	1	-	-	4	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Nebraska												
All races, both sexes	2	1	2	3	6	5	10	11	16	2	1	-
Male	1	1	1	2	1	2	3	2	6	1	-	-
Female	1	-	1	1	5	3	7	9	10	1	1	-
White, both sexes	2	1	2	3	5	5	9	11	15	2	1	-
Male	1	1	1	2	1	2	3	2	5	1	-	-
Female	1	-	1	1	4	3	6	9	10	1	1	-
Black, both sexes	-	-	-	-	1	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	3	1	2	3	2	5	1	1	-	-	-
Male	-	1	1	-	1	1	1	1	-	-	-	-
Female	-	2	-	2	2	1	4	-	1	-	-	-
White, both sexes	-	2	1	2	2	2	5	1	1	-	-	-
Male	-	-	1	-	1	1	1	1	-	-	-	-
Female	-	2	-	2	1	1	4	-	1	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
New Hampshire												
All races, both sexes	4	1	2	1	4	2	5	2	3	2	-	-
Male	1	-	1	1	4	-	1	-	2	1	-	-
Female	3	1	1	-	-	2	4	2	1	1	-	-
White, both sexes	3	1	2	1	4	2	5	2	3	2	-	-
Male	1	-	1	1	4	-	1	-	2	1	-	-
Female	2	1	1	-	-	2	4	2	1	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	7	11	14	17	10	20	21	22	10	9	-	-
Male	4	6	7	4	5	9	9	5	2	2	-	-
Female	3	5	7	13	5	11	12	17	8	7	-	-
White, both sexes	4	9	12	12	7	17	19	21	9	9	-	-
Male	2	6	5	2	4	6	9	5	1	2	-	-
Female	2	3	7	10	3	11	10	16	8	7	-	-
Black, both sexes	3	1	2	3	3	2	1	1	-	-	-	-
Male	2	-	2	1	1	2	-	-	-	-	-	-
Female	1	1	-	2	2	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	-	1	1	-	1	-	-	-
Male	-	-	-	1	-	1	-	-	1	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
New Mexico												
All races, both sexes	1	2	4	2	6	9	4	6	1	1	-	-
Male	1	2	1	-	-	4	1	1	1	-	-	-
Female	-	-	3	2	6	5	3	5	-	1	-	-
White, both sexes	1	2	4	2	5	9	4	5	1	1	-	-
Male	1	2	1	-	-	4	1	-	1	-	-	-
Female	-	-	3	2	5	5	3	5	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	13	15	9	21	34	39	43	53	23	11	-	-
Male	4	5	4	8	13	13	12	14	4	4	-	-
Female	9	10	5	13	21	26	31	39	19	7	-	-
White, both sexes	7	12	5	15	27	33	39	47	20	11	-	-
Male	4	5	3	7	12	11	12	10	4	4	-	-
Female	3	7	2	8	15	22	27	37	16	7	-	-
Black, both sexes	5	3	4	6	5	4	3	5	3	-	-	-
Male	-	-	1	1	-	1	-	4	-	-	-	-
Female	5	3	3	5	5	3	3	1	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	2	2	1	1	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	1	-	-	-	1	1	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
North Carolina												
All races, both sexes	11	15	17	14	21	20	43	28	16	4	-	-
Male	4	4	7	7	9	8	16	7	3	2	-	-
Female	7	11	10	7	12	12	27	21	13	2	-	-
White, both sexes	6	5	9	10	15	14	33	25	14	3	-	-
Male	1	2	4	6	8	5	12	6	3	1	-	-
Female	5	3	5	4	7	9	21	19	11	2	-	-
Black, both sexes	5	10	8	4	6	6	10	3	2	1	-	-
Male	3	2	3	1	1	3	4	1	-	1	-	-
Female	2	8	5	3	5	3	6	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	2	-	2	1	1	1	1	2	-	-
Male	-	-	-	-	1	1	-	1	-	-	-	-
Female	-	-	2	-	1	-	1	-	1	2	-	-
White, both sexes	-	-	2	-	2	1	1	1	1	2	-	-
Male	-	-	-	-	1	1	-	1	-	-	-	-
Female	-	-	2	-	1	-	1	-	1	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Ohio												
All races, both sexes	10	13	14	18	28	27	30	36	19	5	-	-
Male	3	6	3	7	9	9	8	11	2	-	-	-
Female	7	7	11	11	19	18	22	25	17	5	-	-
White, both sexes	6	12	11	14	23	23	30	31	19	4	-	-
Male	3	6	3	6	8	8	8	9	2	-	-	-
Female	3	6	8	8	15	15	22	22	17	4	-	-
Black, both sexes	4	1	3	4	5	4	-	5	-	1	-	-
Male	-	-	-	1	1	1	-	2	-	-	-	-
Female	4	1	3	3	4	3	-	3	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2	4	12	3	4	10	7	4	4	-	1	-
Male	-	1	3	1	1	3	1	1	2	-	-	-
Female	2	3	9	2	3	7	6	3	2	-	1	-
White, both sexes	1	3	12	3	3	7	6	4	4	-	1	-
Male	-	1	3	1	1	2	-	1	2	-	-	-
Female	1	2	9	2	2	5	6	3	2	-	1	-
Black, both sexes	-	-	-	-	1	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	1	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Oregon												
All races, both sexes	-	12	3	9	9	8	17	15	9	6	1	-
Male	-	6	1	5	5	1	7	5	2	1	-	-
Female	-	6	2	4	4	7	10	10	7	5	1	-
White, both sexes	-	12	3	9	9	8	17	14	9	6	1	-
Male	-	6	1	5	5	1	7	5	2	1	-	-
Female	-	6	2	4	4	7	10	9	7	5	1	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	6	18	22	17	29	42	56	52	34	5	2	-
Male	3	8	9	5	16	14	21	8	4	1	-	-
Female	3	10	13	12	13	28	35	44	30	4	2	-
White, both sexes	6	16	19	15	25	35	50	51	32	4	2	-
Male	3	7	8	5	14	11	20	8	4	-	-	-
Female	3	9	11	10	11	24	30	43	28	4	2	-
Black, both sexes	-	2	3	2	4	7	6	1	1	1	-	-
Male	-	1	1	-	2	3	1	-	-	1	-	-
Female	-	1	2	2	2	4	5	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Rhode Island												
All races, both sexes	1	5	-	3	3	4	5	7	2	1	-	-
Male	-	2	-	-	1	1	2	-	-	-	-	-
Female	1	3	-	3	2	3	3	7	2	1	-	-
White, both sexes	-	5	-	3	3	4	5	7	2	1	-	-
Male	-	2	-	-	1	1	2	-	-	-	-	-
Female	-	3	-	3	2	3	3	7	2	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	6	5	11	6	7	15	11	14	6	2	3	-
Male	2	2	3	2	1	6	1	5	-	-	-	-
Female	4	3	8	4	6	9	10	9	6	2	3	-
White, both sexes	3	1	7	3	6	14	10	13	5	2	1	-
Male	1	1	3	1	1	6	1	5	-	-	-	-
Female	2	-	4	2	5	8	9	8	5	2	1	-
Black, both sexes	3	3	4	3	1	1	1	-	1	-	2	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	2	3	4	2	1	1	1	-	1	-	2	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
South Dakota												
All races, both sexes	-	1	2	-	-	1	2	-	1	-	-	-
Male	-	-	1	-	-	1	1	-	-	-	-	-
Female	-	1	1	-	-	-	1	-	1	-	-	-
White, both sexes	-	1	2	-	-	1	2	-	1	-	-	-
Male	-	-	1	-	-	1	1	-	-	-	-	-
Female	-	1	1	-	-	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	3	10	6	11	7	12	11	16	7	1	-	-
Male	-	5	3	3	3	4	3	5	2	1	-	-
Female	3	5	3	8	4	8	8	11	5	-	-	-
White, both sexes	1	5	4	9	4	9	9	13	7	1	-	-
Male	-	2	2	3	3	3	3	3	2	1	-	-
Female	1	3	2	6	1	6	6	10	5	-	-	-
Black, both sexes	2	5	2	2	3	3	2	3	-	-	-	-
Male	-	3	1	-	-	1	-	2	-	-	-	-
Female	2	2	1	2	3	2	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Texas												
All races, both sexes	19	19	26	32	39	33	48	41	22	7	-	-
Male	3	5	9	14	18	15	17	12	5	2	-	-
Female	16	14	17	18	21	18	31	29	17	5	-	-
White, both sexes	11	13	20	24	36	28	42	39	18	5	-	-
Male	3	5	6	10	15	15	16	12	3	2	-	-
Female	8	8	14	14	21	13	26	27	15	3	-	-
Black, both sexes	7	4	6	8	2	4	6	1	4	1	-	-
Male	-	-	3	4	2	-	1	-	2	-	-	-
Female	7	4	3	4	-	4	5	1	2	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	-	1	1	-	1	-	1	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	2	-	-	-	1	-	1	-	1	-	-
Utah												
All races, both sexes	1	4	3	-	3	7	4	14	7	1	-	-
Male	-	2	2	-	-	2	1	3	2	1	-	-
Female	1	2	1	-	3	5	3	11	5	-	-	-
White, both sexes	1	3	3	-	3	7	3	14	7	1	-	-
Male	-	2	2	-	-	2	1	3	2	1	-	-
Female	1	1	1	-	3	5	2	11	5	-	-	-
Black, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Vermont												
All races, both sexes	-	1	2	1	1	3	3	1	3	1	-	-
Male	-	-	2	-	-	2	-	-	2	1	-	-
Female	-	1	-	1	1	1	3	1	1	-	-	-
White, both sexes	-	1	2	1	1	3	3	1	3	1	-	-
Male	-	-	2	-	-	2	-	-	2	1	-	-
Female	-	1	-	1	1	1	3	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	5	13	9	7	13	19	24	15	12	6	-	-
Male	2	4	5	2	6	4	7	2	3	1	-	-
Female	3	9	4	5	7	15	17	13	9	5	-	-
White, both sexes	4	10	6	5	7	14	18	9	10	6	-	-
Male	1	3	4	2	3	3	5	1	3	1	-	-
Female	3	7	2	3	4	11	13	8	7	5	-	-
Black, both sexes	1	3	3	2	6	5	6	6	2	-	-	-
Male	1	1	1	-	3	1	2	1	-	-	-	-
Female	-	2	2	2	3	4	4	5	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Washington												
All races, both sexes	6	5	7	13	8	17	10	21	16	8	-	-
Male	2	1	4	5	4	9	4	5	3	-	-	-
Female	4	4	3	8	4	8	6	16	13	8	-	-
White, both sexes	5	5	7	13	8	16	10	21	15	8	-	-
Male	1	1	4	5	4	9	4	5	3	-	-	-
Female	4	4	3	8	4	7	6	16	12	8	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
West Virginia												
All races, both sexes	1	2	2	4	6	4	5	5	2	-	-	-
Male	1	-	1	3	2	3	-	1	-	-	-	-
Female	-	2	1	1	4	1	5	4	2	-	-	-
White, both sexes	1	2	2	3	5	4	5	5	2	-	-	-
Male	1	-	1	2	2	3	-	1	-	-	-	-
Female	-	2	1	1	3	1	5	4	2	-	-	-
Black, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Wisconsin												
All races, both sexes	3	7	5	8	11	9	35	15	15	2	-	-
Male	2	4	3	3	6	3	14	5	5	-	-	-
Female	1	3	2	5	5	6	21	10	10	2	-	-
White, both sexes	3	3	5	7	11	8	33	15	15	2	-	-
Male	2	2	3	3	6	3	12	5	5	-	-	-
Female	1	1	2	4	5	5	21	10	10	2	-	-
Black, both sexes	-	4	-	1	-	1	2	-	-	-	-	-
Male	-	2	-	-	-	-	2	-	-	-	-	-
Female	-	2	-	1	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	1	-	-	1	3	3	2	2	-	-	-
Male	-	-	-	-	-	1	2	-	-	-	-	-
Female	1	1	-	-	1	2	1	2	2	-	-	-
White, both sexes	1	1	-	-	1	3	3	2	2	-	-	-
Male	-	-	-	-	-	1	2	-	-	-	-	-
Female	1	1	-	-	1	2	1	2	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
United States												
All races, both sexes	160,196	297	137	94	108	256	508	654	828	1,349	2,131	3,336
Male	72,310	164	69	54	61	183	332	453	535	886	1,415	2,183
Female	87,886	133	68	40	47	73	176	201	293	463	716	1,153
White, both sexes	139,016	172	89	64	72	162	338	416	540	883	1,448	2,352
Male	61,880	89	42	34	41	112	216	294	363	584	962	1,585
Female	77,136	83	47	30	31	50	122	122	177	299	486	767
Black, both sexes	18,295	106	36	22	34	88	152	210	249	422	608	901
Male	8,945	65	23	15	18	66	105	137	145	268	399	539
Female	9,350	41	13	7	16	22	47	73	104	154	209	362
American Indian 1/, both sexes	654	4	4	1	-	2	7	7	18	12	27	44
Male	358	2	3	1	-	2	1	7	14	9	20	33
Female	296	2	1	-	-	-	6	-	4	3	7	11
Asian or Pacific Islander 2/, both sexes	2,231	15	8	7	2	4	11	21	21	32	48	39
Male	1,127	8	1	4	2	3	10	15	13	25	34	26
Female	1,104	7	7	3	-	1	1	6	8	7	14	13
Alabama												
All races, both sexes	5,688	8	3	4	2	4	20	19	29	47	115	179
Male	2,643	4	3	3	-	2	14	11	17	21	70	120
Female	3,045	4	-	1	2	2	6	8	12	26	45	59
White, both sexes	4,284	5	-	2	1	3	13	11	15	25	80	104
Male	1,975	2	-	2	-	2	11	6	11	14	48	65
Female	2,309	3	-	-	1	1	2	5	4	11	32	39
Black, both sexes	1,389	3	3	2	1	1	7	8	14	22	35	74
Male	660	2	3	1	-	-	3	5	6	7	22	54
Female	729	1	-	1	1	1	4	3	8	15	13	20
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Alaska												
All races, both sexes	176	-	-	1	-	1	-	-	3	-	7	6
Male	88	-	-	-	-	-	-	-	3	-	5	4
Female	88	-	-	1	-	1	-	-	-	-	2	2
White, both sexes	133	-	-	1	-	1	-	-	1	-	4	5
Male	62	-	-	-	-	-	-	-	1	-	3	3
Female	71	-	-	1	-	1	-	-	-	-	1	2
Black, both sexes	3	-	-	-	-	-	-	-	-	-	2	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	36	-	-	-	-	-	-	-	2	-	1	-
Male	22	-	-	-	-	-	-	-	2	-	1	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2,098	6	2	2	1	6	11	14	12	15	33	56
Male	1,027	2	1	1	1	4	7	10	9	11	21	38
Female	1,071	4	1	1	-	2	4	4	3	4	12	18
White, both sexes	1,986	5	2	1	1	5	8	14	10	15	27	51
Male	974	2	1	1	1	4	7	10	7	11	17	35
Female	1,012	3	1	-	-	1	1	4	3	4	10	16
Black, both sexes	48	1	-	1	-	1	1	-	1	-	2	3
Male	22	-	-	-	-	-	-	-	1	-	1	1
Female	26	1	-	1	-	1	1	-	-	-	1	2
American Indian 1/, both sexes	55	-	-	-	-	-	2	-	1	-	3	1
Male	28	-	-	-	-	-	-	-	1	-	2	1
Female	27	-	-	-	-	-	2	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	1	1
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Arkansas												
All races, both sexes	1,828	4	3	-	3	5	7	8	6	13	22	42
Male	767	2	2	-	3	4	5	6	4	10	14	26
Female	1,061	2	1	-	-	1	2	2	2	3	8	16
White, both sexes	1,575	3	1	-	2	3	5	2	3	10	13	33
Male	656	1	1	-	2	2	3	2	2	9	7	21
Female	919	2	-	-	-	1	2	-	1	1	6	12
Black, both sexes	240	1	2	-	1	2	1	6	3	3	8	9
Male	107	1	1	-	1	2	1	4	2	1	7	5
Female	133	-	1	-	-	-	-	2	1	2	1	4
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	1	-	-	-	-	-
Male	3	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	12,866	32	13	8	9	21	50	78	86	133	216	331
Male	6,014	19	6	5	6	18	36	56	64	96	160	227
Female	6,852	13	7	3	3	3	14	22	22	37	56	104
White, both sexes	11,205	23	10	6	8	12	39	50	63	103	151	247
Male	5,143	12	6	3	6	11	26	38	47	71	112	174
Female	6,062	11	4	3	2	1	13	12	16	32	39	73
Black, both sexes	905	5	2	-	1	7	8	16	17	21	43	62
Male	464	4	-	-	-	5	7	8	12	17	32	40
Female	441	1	2	-	1	2	1	8	5	4	11	22
American Indian 1/, both sexes	69	-	-	-	-	-	-	2	1	-	6	7
Male	42	-	-	-	-	-	-	2	1	-	5	4
Female	27	-	-	-	-	-	-	-	-	-	1	3
Asian or Pacific Islander 2/, both sexes	687	4	1	2	-	2	3	10	5	9	16	15
Male	365	3	-	2	-	2	3	8	4	8	11	9
Female	322	1	1	-	-	-	-	2	1	1	5	6

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Colorado												
All races, both sexes	1,635	5	3	3	-	4	6	6	7	18	28	37
Male	717	2	3	2	-	2	4	5	3	14	18	28
Female	918	3	-	1	-	2	2	1	4	4	10	9
White, both sexes	1,570	4	3	3	-	3	4	6	6	17	24	33
Male	685	2	3	2	-	1	2	5	2	14	17	25
Female	885	2	-	1	-	2	2	1	4	3	7	8
Black, both sexes	43	1	-	-	-	-	2	-	1	1	3	3
Male	22	-	-	-	-	-	2	-	1	-	1	2
Female	21	1	-	-	-	-	-	-	-	1	2	1
American Indian 1/, both sexes	8	-	-	-	-	1	-	-	-	-	-	1
Male	6	-	-	-	-	1	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1	-
Connecticut												
All races, both sexes	2,507	1	3	2	-	6	6	3	13	17	21	51
Male	1,150	1	1	1	-	5	3	-	7	14	18	32
Female	1,357	-	2	1	-	1	3	3	6	3	3	19
White, both sexes	2,330	1	1	2	-	2	4	3	6	10	17	44
Male	1,064	1	-	1	-	2	2	-	4	8	14	29
Female	1,266	-	1	1	-	-	2	3	2	2	3	15
Black, both sexes	167	-	1	-	-	3	2	-	7	6	4	7
Male	81	-	1	-	-	3	1	-	3	5	4	3
Female	86	-	-	-	-	-	1	-	4	1	-	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	1	-	-	1	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	1	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Delaware												
All races, both sexes	505	-	1	-	-	1	4	1	2	10	11	12
Male	221	-	-	-	-	1	2	-	-	5	8	6
Female	284	-	1	-	-	-	2	1	2	5	3	6
White, both sexes	429	-	-	-	-	1	2	1	1	9	7	7
Male	184	-	-	-	-	1	-	-	-	5	5	5
Female	245	-	-	-	-	-	2	1	1	4	2	2
Black, both sexes	73	-	1	-	-	-	2	-	1	1	4	5
Male	35	-	-	-	-	-	2	-	-	-	3	1
Female	38	-	1	-	-	-	-	-	1	1	1	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	227	-	-	1	-	-	1	1	-	4	2	7
Male	96	-	-	1	-	-	1	-	-	3	2	5
Female	131	-	-	-	-	-	-	1	-	1	-	2
White, both sexes	59	-	-	-	-	-	-	-	-	-	1	1
Male	24	-	-	-	-	-	-	-	-	-	1	1
Female	35	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	168	-	-	1	-	-	1	1	-	4	1	6
Male	72	-	-	1	-	-	1	-	-	3	1	4
Female	96	-	-	-	-	-	-	1	-	1	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Florida												
All races, both sexes	8,755	24	4	5	10	16	25	18	44	82	96	155
Male	4,407	13	3	3	6	12	19	10	29	63	63	98
Female	4,348	11	1	2	4	4	6	8	15	19	33	57
White, both sexes	7,816	6	3	4	3	9	17	8	24	47	71	105
Male	3,906	3	2	3	1	7	11	5	16	41	45	70
Female	3,910	3	1	1	2	2	6	3	8	6	26	35
Black, both sexes	884	17	1	1	6	7	8	10	18	33	22	47
Male	466	9	1	-	4	5	8	5	12	21	17	25
Female	418	8	-	1	2	2	-	5	6	12	5	22
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	1	-	-	2
Male	7	-	-	-	-	-	-	-	1	-	-	2
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	47	1	-	-	1	-	-	-	1	2	3	1
Male	28	1	-	-	1	-	-	-	-	1	1	1
Female	19	-	-	-	-	-	-	-	1	1	2	-
Georgia												
All races, both sexes	6,501	23	9	6	5	12	24	24	34	77	116	149
Male	3,020	13	2	5	3	9	18	14	17	54	68	99
Female	3,481	10	7	1	2	3	6	10	17	23	48	50
White, both sexes	4,781	8	4	3	2	8	9	12	15	46	59	80
Male	2,197	6	-	2	1	5	4	8	7	32	29	56
Female	2,584	2	4	1	1	3	5	4	8	14	30	24
Black, both sexes	1,679	15	2	3	3	4	14	12	19	30	56	69
Male	809	7	2	3	2	4	13	6	10	21	38	43
Female	870	8	-	-	1	-	1	6	9	9	18	26
American Indian 1/, both sexes	6	-	1	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	35	-	2	-	-	-	1	-	-	1	-	-
Male	12	-	-	-	-	-	1	-	-	1	-	-
Female	23	-	2	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Hawaii												
All races, both sexes	916	2	-	1	-	1	2	1	8	10	14	21
Male	502	-	-	1	-	1	1	1	3	6	10	14
Female	414	2	-	-	-	-	1	-	5	4	4	7
White, both sexes	230	1	-	-	-	-	1	-	-	1	5	3
Male	131	-	-	-	-	-	-	-	-	1	4	2
Female	99	1	-	-	-	-	1	-	-	-	1	1
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	677	1	-	1	-	1	1	1	8	9	9	17
Male	366	-	-	1	-	1	1	1	3	5	6	11
Female	311	1	-	-	-	-	-	-	5	4	3	6
Idaho												
All races, both sexes	694	1	1	-	-	-	2	1	2	2	8	12
Male	324	1	-	-	-	-	1	1	1	1	3	11
Female	370	-	1	-	-	-	1	-	1	1	5	1
White, both sexes	680	1	1	-	-	-	2	1	2	2	7	12
Male	314	1	-	-	-	-	1	1	1	1	2	11
Female	366	-	1	-	-	-	1	-	1	1	5	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Illinois												
All races, both sexes	7,085	16	8	4	4	10	18	38	40	72	88	151
Male	3,101	11	4	2	1	9	12	29	24	41	60	102
Female	3,984	5	4	2	3	1	6	9	16	31	28	49
White, both sexes	6,085	9	5	3	4	7	16	24	26	39	47	91
Male	2,579	7	3	1	1	6	11	17	17	27	34	65
Female	3,506	2	2	2	3	1	5	7	9	12	13	26
Black, both sexes	947	7	3	1	-	3	2	13	14	32	39	59
Male	495	4	1	1	-	3	1	12	7	14	24	36
Female	452	3	2	-	-	-	1	1	7	18	15	23
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	1	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	47	-	-	-	-	-	-	1	-	-	1	1
Male	24	-	-	-	-	-	-	-	-	-	1	1
Female	23	-	-	-	-	-	-	1	-	-	-	-
Indiana												
All races, both sexes	4,271	9	2	-	3	7	15	18	18	29	65	92
Male	1,892	6	-	-	1	5	11	8	10	14	39	64
Female	2,379	3	2	-	2	2	4	10	8	15	26	28
White, both sexes	3,965	5	2	-	3	6	10	12	13	22	52	83
Male	1,738	3	-	-	1	5	9	5	8	12	30	57
Female	2,227	2	2	-	2	1	1	7	5	10	22	26
Black, both sexes	299	4	-	-	-	1	5	6	5	6	12	9
Male	150	3	-	-	-	-	2	3	2	1	8	7
Female	149	1	-	-	-	1	3	3	3	5	4	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	1	1	-
Male	4	-	-	-	-	-	-	-	-	1	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Iowa												
All races, both sexes	1,201	2	1	-	-	2	2	7	9	15	9	20
Male	525	-	1	-	-	2	2	5	5	10	7	14
Female	676	2	-	-	-	-	-	2	4	5	2	6
White, both sexes	1,184	1	1	-	-	2	2	4	9	14	9	18
Male	516	-	1	-	-	2	2	3	5	9	7	12
Female	668	1	-	-	-	-	-	1	4	5	2	6
Black, both sexes	16	1	-	-	-	-	-	3	-	-	-	2
Male	8	-	-	-	-	-	-	2	-	-	-	2
Female	8	1	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	2,058	3	1	-	-	-	6	3	9	8	18	39
Male	843	2	-	-	-	-	3	3	6	6	9	26
Female	1,215	1	1	-	-	-	3	-	3	2	9	13
White, both sexes	1,924	2	1	-	-	-	5	3	7	7	17	30
Male	779	1	-	-	-	-	3	3	4	5	9	21
Female	1,145	1	1	-	-	-	2	-	3	2	8	9
Black, both sexes	106	1	-	-	-	-	1	-	2	1	1	5
Male	54	1	-	-	-	-	-	-	2	1	-	2
Female	52	-	-	-	-	-	1	-	-	-	1	3
American Indian 1/, both sexes	20	-	-	-	-	-	-	-	-	-	-	4
Male	9	-	-	-	-	-	-	-	-	-	-	3
Female	11	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Kentucky												
All races, both sexes	3,055	4	2	1	3	1	9	8	12	27	43	61
Male	1,311	1	2	-	1	-	4	5	6	10	27	33
Female	1,744	3	-	1	2	1	5	3	6	17	16	28
White, both sexes	2,848	4	2	1	3	1	6	8	9	23	36	48
Male	1,218	1	2	-	1	-	2	5	5	9	24	27
Female	1,630	3	-	1	2	1	4	3	4	14	12	21
Black, both sexes	202	-	-	-	-	-	3	-	3	4	7	13
Male	89	-	-	-	-	-	2	-	1	1	3	6
Female	113	-	-	-	-	-	1	-	2	3	4	7
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	3,092	5	4	1	4	7	13	7	16	31	39	59
Male	1,350	3	4	1	2	6	6	3	9	22	23	33
Female	1,742	2	-	-	2	1	7	4	7	9	16	26
White, both sexes	2,277	1	-	-	2	4	5	2	5	21	19	33
Male	971	1	-	-	1	4	3	1	2	15	11	16
Female	1,306	-	-	-	1	-	2	1	3	6	8	17
Black, both sexes	801	4	3	1	2	3	8	4	11	10	20	26
Male	374	2	3	1	1	2	3	2	7	7	12	17
Female	427	2	-	-	1	1	5	2	4	3	8	9
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	1	-	-	-	-	1	-	-	-	-
Male	4	-	1	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Maine												
All races, both sexes	920	2	-	1	2	2	3	-	3	4	4	9
Male	432	1	-	-	1	-	2	-	2	3	2	6
Female	488	1	-	1	1	2	1	-	1	1	2	3
White, both sexes	913	2	-	1	2	2	3	-	3	4	3	7
Male	426	1	-	-	1	-	2	-	2	3	2	4
Female	487	1	-	1	1	2	1	-	1	1	1	3
Black, both sexes	3	-	-	-	-	-	-	-	-	-	1	2
Male	2	-	-	-	-	-	-	-	-	-	-	2
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	2,284	3	2	2	4	8	11	15	19	37	52	47
Male	1,061	1	1	2	-	7	9	8	11	24	40	22
Female	1,223	2	1	-	4	1	2	7	8	13	12	25
White, both sexes	1,683	1	1	1	3	1	6	9	7	17	29	24
Male	758	-	-	1	-	1	5	4	5	11	23	12
Female	925	1	1	-	3	-	1	5	2	6	6	12
Black, both sexes	563	1	1	1	1	7	5	6	12	20	21	23
Male	286	1	1	1	-	6	4	4	6	13	15	10
Female	277	-	-	-	1	1	1	2	6	7	6	13
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	36	1	-	-	-	-	-	-	-	-	2	-
Male	17	-	-	-	-	-	-	-	-	-	2	-
Female	19	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Massachusetts												
All races, both sexes	3,785	4	1	1	3	10	11	16	12	15	40	40
Male	1,583	2	-	-	2	6	7	10	11	8	29	30
Female	2,202	2	1	1	1	4	4	6	1	7	11	10
White, both sexes	3,601	4	1	1	2	8	10	13	11	13	32	35
Male	1,483	2	-	-	1	5	6	8	10	7	21	26
Female	2,118	2	1	1	1	3	4	5	1	6	11	9
Black, both sexes	150	-	-	-	1	2	1	2	1	1	7	5
Male	81	-	-	-	1	1	1	1	1	-	7	4
Female	69	-	-	-	-	1	-	1	-	1	-	1
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	30	-	-	-	-	-	-	1	-	1	1	-
Male	17	-	-	-	-	-	-	1	-	1	1	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	5,527	10	1	1	3	11	15	26	36	58	71	134
Male	2,427	6	1	-	2	7	9	20	19	38	49	87
Female	3,100	4	-	1	1	4	6	6	17	20	22	47
White, both sexes	4,791	9	1	1	2	7	7	13	20	29	42	90
Male	2,052	5	1	-	2	3	4	12	10	21	31	62
Female	2,739	4	-	1	-	4	3	1	10	8	11	28
Black, both sexes	698	1	-	-	1	4	8	13	16	29	26	44
Male	358	1	-	-	-	4	5	8	9	17	15	25
Female	340	-	-	-	1	-	3	5	7	12	11	19
American Indian 1/, both sexes	14	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	24	-	-	-	-	-	-	-	-	-	2	-
Male	13	-	-	-	-	-	-	-	-	-	2	-
Female	11	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Minnesota												
All races, both sexes	2,645	6	5	2	-	2	3	11	9	8	23	34
Male	1,174	6	2	2	-	-	2	8	5	3	19	20
Female	1,471	-	3	-	-	2	1	3	4	5	4	14
White, both sexes	2,556	4	5	1	-	2	3	4	6	4	17	28
Male	1,125	4	2	1	-	-	2	4	3	1	14	16
Female	1,431	-	3	-	-	2	1	-	3	3	3	12
Black, both sexes	54	-	-	-	-	-	-	6	1	3	5	3
Male	35	-	-	-	-	-	-	3	1	2	4	3
Female	19	-	-	-	-	-	-	3	-	1	1	-
American Indian 1/, both sexes	24	1	-	-	-	-	-	-	2	1	1	3
Male	10	1	-	-	-	-	-	-	1	-	1	1
Female	14	-	-	-	-	-	-	-	1	1	-	2
Asian or Pacific Islander 2/, both sexes	11	1	-	1	-	-	-	1	-	-	-	-
Male	4	1	-	1	-	-	-	1	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	2,931	5	1	1	3	3	9	12	10	26	41	76
Male	1,279	3	1	1	2	1	7	8	5	15	24	47
Female	1,652	2	-	-	1	2	2	4	5	11	17	29
White, both sexes	1,959	2	-	1	1	-	4	2	5	9	15	34
Male	861	1	-	1	1	-	3	2	3	5	9	21
Female	1,098	1	-	-	-	-	1	-	2	4	6	13
Black, both sexes	961	3	1	-	2	3	5	10	5	17	25	42
Male	412	2	1	-	1	1	4	6	2	10	14	26
Female	549	1	-	-	1	2	1	4	3	7	11	16
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Missouri												
All races, both sexes	3,747	5	3	4	2	6	11	12	22	28	42	75
Male	1,654	2	2	2	1	6	7	9	14	23	28	56
Female	2,093	3	1	2	1	-	4	3	8	5	14	19
White, both sexes	3,456	2	3	3	1	5	10	10	16	19	35	57
Male	1,509	1	2	1	1	5	6	8	11	14	23	46
Female	1,947	1	1	2	-	-	4	2	5	5	12	11
Black, both sexes	280	2	-	1	1	1	1	2	5	9	7	18
Male	140	-	-	1	-	1	1	1	3	9	5	10
Female	140	2	-	-	1	-	-	1	2	-	2	8
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	7	1	-	-	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	698	-	1	-	1	2	2	3	1	-	5	7
Male	320	-	1	-	1	2	-	1	-	-	4	3
Female	378	-	-	-	-	-	2	2	1	-	1	4
White, both sexes	676	-	-	-	1	2	1	3	-	-	5	5
Male	307	-	-	-	1	2	-	1	-	-	4	1
Female	369	-	-	-	-	-	1	2	-	-	1	4
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	21	-	1	-	-	-	1	-	1	-	-	2
Male	13	-	1	-	-	-	-	-	-	-	-	2
Female	8	-	-	-	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Nebraska												
All races, both sexes	1,269	1	1	-	1	-	1	-	1	9	15	19
Male	537	-	-	-	-	-	-	-	1	7	9	11
Female	732	1	1	-	1	-	1	-	-	2	6	8
White, both sexes	1,224	1	1	-	1	-	1	-	1	8	12	17
Male	521	-	-	-	-	-	-	-	1	6	7	10
Female	703	1	1	-	1	-	1	-	-	2	5	7
Black, both sexes	37	-	-	-	-	-	-	-	-	1	3	1
Male	13	-	-	-	-	-	-	-	-	1	2	1
Female	24	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1,738	2	2	2	2	3	10	8	12	22	36	52
Male	968	1	-	2	2	1	7	6	9	18	28	37
Female	770	1	2	-	-	2	3	2	3	4	8	15
White, both sexes	1,501	1	-	2	2	3	6	4	9	14	30	41
Male	842	-	-	2	2	1	5	3	7	12	22	30
Female	659	1	-	-	-	2	1	1	2	2	8	11
Black, both sexes	166	1	1	-	-	-	4	3	1	8	4	8
Male	96	1	-	-	-	-	2	2	-	6	4	5
Female	70	-	1	-	-	-	2	1	1	2	-	3
American Indian 1/, both sexes	16	-	-	-	-	-	-	1	1	-	2	1
Male	9	-	-	-	-	-	-	1	1	-	2	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	55	-	1	-	-	-	-	-	1	-	-	2
Male	21	-	-	-	-	-	-	-	1	-	-	2
Female	34	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
New Hampshire												
All races, both sexes	632	-	-	-	1	1	1	4	-	-	2	8
Male	275	-	-	-	-	1	-	2	-	-	-	4
Female	357	-	-	-	1	-	1	2	-	-	2	4
White, both sexes	625	-	-	-	1	1	1	4	-	-	2	7
Male	274	-	-	-	-	1	-	2	-	-	-	4
Female	351	-	-	-	1	-	1	2	-	-	2	3
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	4,617	7	1	2	2	5	9	12	17	33	48	83
Male	2,058	4	-	1	1	4	4	10	16	19	24	53
Female	2,559	3	1	1	1	1	5	2	1	14	24	30
White, both sexes	4,061	3	1	-	1	1	6	6	11	21	29	55
Male	1,796	2	-	-	1	1	3	5	10	11	19	38
Female	2,265	1	1	-	-	-	3	1	1	10	10	17
Black, both sexes	494	4	-	1	1	4	3	4	5	11	18	27
Male	233	2	-	1	-	3	1	4	5	7	5	15
Female	261	2	-	-	1	1	2	-	-	4	13	12
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	60	-	-	1	-	-	-	2	1	1	1	1
Male	28	-	-	-	-	-	-	1	1	1	-	-
Female	32	-	-	1	-	-	-	1	-	-	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
New Mexico												
All races, both sexes	826	1	-	-	-	4	1	3	4	7	6	14
Male	377	1	-	-	-	4	-	3	1	4	6	12
Female	449	-	-	-	-	-	1	-	3	3	-	2
White, both sexes	779	1	-	-	-	3	1	2	3	7	5	12
Male	345	1	-	-	-	3	-	2	-	4	5	10
Female	434	-	-	-	-	-	1	-	3	3	-	2
Black, both sexes	13	-	-	-	-	1	-	-	-	-	1	-
Male	9	-	-	-	-	1	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	30	-	-	-	-	-	-	1	1	-	-	2
Male	21	-	-	-	-	-	-	1	1	-	-	2
Female	9	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	7,874	12	8	6	9	13	30	28	37	55	97	143
Male	3,455	7	4	2	3	8	21	20	31	41	66	98
Female	4,419	5	4	4	6	5	9	8	6	14	31	45
White, both sexes	7,106	7	3	4	4	7	18	16	29	33	67	103
Male	3,075	3	-	1	1	4	13	11	26	23	46	76
Female	4,031	4	3	3	3	3	5	5	3	10	21	27
Black, both sexes	683	4	5	2	5	6	10	12	7	19	29	37
Male	337	3	4	1	2	4	6	9	4	16	19	19
Female	346	1	1	1	3	2	4	3	3	3	10	18
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	1	-	3
Male	7	-	-	-	-	-	-	-	-	1	-	3
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	74	1	-	-	-	-	2	-	1	2	1	-
Male	36	1	-	-	-	-	2	-	1	1	1	-
Female	38	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
North Carolina												
All races, both sexes	4,856	16	9	7	4	6	12	19	25	52	62	112
Male	2,181	10	4	4	3	4	4	15	19	37	37	70
Female	2,675	6	5	3	1	2	8	4	6	15	25	42
White, both sexes	3,801	9	6	6	2	1	5	8	15	26	37	58
Male	1,690	4	2	3	1	-	2	6	11	19	20	37
Female	2,111	5	4	3	1	1	3	2	4	7	17	21
Black, both sexes	1,001	6	2	1	2	5	5	10	9	25	24	50
Male	466	6	1	1	2	4	1	8	7	17	16	30
Female	535	-	1	-	-	1	4	2	2	8	8	20
American Indian 1/, both sexes	39	1	1	-	-	-	2	-	-	1	1	4
Male	19	-	1	-	-	-	1	-	-	1	1	3
Female	20	1	-	-	-	-	1	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	1	1	-	-	-
Male	6	-	-	-	-	-	-	1	1	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	270	2	1	-	-	-	1	3	3	2	2	6
Male	118	1	-	-	-	-	-	2	1	2	2	3
Female	152	1	1	-	-	-	1	1	2	-	-	3
White, both sexes	254	2	1	-	-	-	-	2	3	1	1	4
Male	107	1	-	-	-	-	-	1	1	1	1	2
Female	147	1	1	-	-	-	-	1	2	-	-	2
Black, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	15	-	-	-	-	-	1	-	-	1	1	2
Male	10	-	-	-	-	-	-	-	-	1	1	1
Female	5	-	-	-	-	-	1	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Ohio												
All races, both sexes	5,682	7	10	4	3	5	22	22	34	47	72	124
Male	2,476	5	6	2	2	4	15	17	22	31	43	82
Female	3,206	2	4	2	1	1	7	5	12	16	29	42
White, both sexes	5,135	3	8	4	1	3	16	15	27	37	54	97
Male	2,204	2	4	2	-	2	10	12	18	23	33	66
Female	2,931	1	4	2	1	1	6	3	9	14	21	31
Black, both sexes	535	4	2	-	1	2	6	6	7	10	18	27
Male	267	3	2	-	1	2	5	4	4	8	10	16
Female	268	1	-	-	-	-	1	2	3	2	8	11
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	1	-	-	1	-	-	-	-
Male	4	-	-	-	1	-	-	1	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2,534	4	3	2	2	2	2	11	15	19	37	55
Male	1,123	1	1	2	2	1	2	10	14	14	21	39
Female	1,411	3	2	-	-	1	-	1	1	5	16	16
White, both sexes	2,233	3	1	1	2	2	1	8	9	14	29	34
Male	982	-	-	1	2	1	1	7	8	10	16	24
Female	1,251	3	1	-	-	1	-	1	1	4	13	10
Black, both sexes	182	-	1	-	-	-	1	-	3	1	5	15
Male	78	-	-	-	-	-	1	-	3	1	3	10
Female	104	-	1	-	-	-	-	-	-	-	2	5
American Indian 1/, both sexes	111	1	1	1	-	-	-	3	3	4	2	6
Male	59	1	1	1	-	-	-	3	3	3	1	5
Female	52	-	-	-	-	-	-	-	-	1	1	1
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Oregon												
All races, both sexes	2,192	3	1	1	1	4	3	8	12	12	12	25
Male	941	2	-	1	1	3	1	7	9	6	8	19
Female	1,251	1	1	-	-	1	2	1	3	6	4	6
White, both sexes	2,143	2	1	1	1	3	3	6	10	9	12	25
Male	916	2	-	1	1	2	1	5	7	4	8	19
Female	1,227	-	1	-	-	1	2	1	3	5	4	6
Black, both sexes	11	-	-	-	-	1	-	1	1	3	-	-
Male	8	-	-	-	-	1	-	1	1	2	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	25	1	-	-	-	-	-	1	1	-	-	-
Male	8	-	-	-	-	-	-	1	1	-	-	-
Female	17	1	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	9,801	5	2	5	2	12	29	31	43	54	81	164
Male	4,241	1	-	1	2	8	20	20	28	29	62	99
Female	5,560	4	2	4	-	4	9	11	15	25	19	65
White, both sexes	9,014	5	1	2	2	7	20	21	26	33	59	125
Male	3,856	1	-	-	2	5	14	12	18	15	46	79
Female	5,158	4	1	2	-	2	6	9	8	18	13	46
Black, both sexes	748	-	-	3	-	5	8	10	17	20	21	39
Male	366	-	-	1	-	3	5	8	10	13	15	20
Female	382	-	-	2	-	2	3	2	7	7	6	19
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	38	-	1	-	-	-	1	-	-	1	1	-
Male	19	-	-	-	-	-	1	-	-	1	1	-
Female	19	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Rhode Island												
All races, both sexes	402	2	1	-	-	1	-	1	2	-	4	4
Male	171	1	1	-	-	-	-	1	2	-	3	4
Female	231	1	-	-	-	1	-	-	-	-	1	-
White, both sexes	389	1	1	-	-	-	-	1	2	-	3	4
Male	166	-	1	-	-	-	-	1	2	-	2	4
Female	223	1	-	-	-	-	-	-	-	-	1	-
Black, both sexes	9	-	-	-	-	1	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	3,016	7	-	-	5	5	11	18	23	35	69	72
Male	1,449	4	-	-	5	4	5	13	7	26	46	40
Female	1,567	3	-	-	-	1	6	5	16	9	23	32
White, both sexes	2,108	4	-	-	3	2	7	8	10	17	40	36
Male	996	3	-	-	3	1	2	5	3	12	29	22
Female	1,112	1	-	-	-	1	5	3	7	5	11	14
Black, both sexes	900	3	-	-	2	3	4	10	13	18	29	36
Male	450	1	-	-	2	3	3	8	4	14	17	18
Female	450	2	-	-	-	-	1	2	9	4	12	18
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
South Dakota												
All races, both sexes	293	-	2	-	-	2	-	2	4	1	2	4
Male	125	-	2	-	-	1	-	2	2	1	1	2
Female	168	-	-	-	-	1	-	-	2	-	1	2
White, both sexes	276	-	2	-	-	1	-	1	3	1	1	3
Male	116	-	2	-	-	-	-	1	1	1	1	1
Female	160	-	-	-	-	1	-	-	2	-	-	2
Black, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	15	-	-	-	-	1	-	-	1	-	1	1
Male	8	-	-	-	-	1	-	-	1	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	2,590	6	1	2	1	5	12	15	18	28	61	75
Male	1,136	5	-	-	-	4	9	12	11	19	32	55
Female	1,454	1	1	2	1	1	3	3	7	9	29	20
White, both sexes	2,160	2	1	1	1	4	9	13	13	17	40	55
Male	942	1	-	-	-	4	8	10	9	11	19	38
Female	1,218	1	1	1	1	-	1	3	4	6	21	17
Black, both sexes	422	4	-	1	-	1	3	2	5	11	21	20
Male	192	4	-	-	-	-	1	2	2	8	13	17
Female	230	-	-	1	-	1	2	-	3	3	8	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Texas												
All races, both sexes	10,317	25	12	6	4	25	46	64	53	107	151	232
Male	4,752	10	6	3	2	15	31	44	37	72	103	149
Female	5,565	15	6	3	2	10	15	20	16	35	48	83
White, both sexes	8,974	18	10	3	2	17	32	44	39	77	108	179
Male	4,083	8	6	1	2	9	20	33	27	53	73	121
Female	4,891	10	4	2	-	8	12	11	12	24	35	58
Black, both sexes	1,234	6	2	2	2	8	13	19	14	29	41	51
Male	616	2	-	2	-	6	11	11	10	18	28	26
Female	618	4	2	-	2	2	2	8	4	11	13	25
American Indian 1/, both sexes	17	-	-	-	-	-	1	-	-	-	-	2
Male	11	-	-	-	-	-	-	-	-	-	-	2
Female	6	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	92	1	-	1	-	-	-	1	-	1	2	-
Male	42	-	-	-	-	-	-	-	-	1	2	-
Female	50	1	-	1	-	-	-	1	-	-	-	-
Utah												
All races, both sexes	1,268	3	2	-	1	-	1	5	7	9	12	18
Male	575	2	-	-	1	-	1	3	5	6	7	11
Female	693	1	2	-	-	-	-	2	2	3	5	7
White, both sexes	1,237	3	2	-	1	-	1	5	6	8	10	17
Male	557	2	-	-	1	-	1	3	4	5	5	10
Female	680	1	2	-	-	-	-	2	2	3	5	7
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	1
Male	5	-	-	-	-	-	-	-	-	-	1	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	1	1	1	-
Male	6	-	-	-	-	-	-	-	1	1	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Vermont												
All races, both sexes	235	-	-	-	-	-	1	2	2	-	-	1
Male	106	-	-	-	-	-	-	1	2	-	-	1
Female	129	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	235	-	-	-	-	-	1	2	2	-	-	1
Male	106	-	-	-	-	-	-	1	2	-	-	1
Female	129	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	4,686	5	2	3	3	5	20	23	23	34	69	105
Male	2,124	4	2	2	2	4	12	15	16	17	47	66
Female	2,562	1	-	1	1	1	8	8	7	17	22	39
White, both sexes	3,681	1	1	2	2	3	10	13	12	24	31	71
Male	1,627	1	1	2	1	2	5	9	9	12	22	46
Female	2,054	-	-	-	1	1	5	4	3	12	9	25
Black, both sexes	943	3	1	-	1	2	8	10	10	9	34	34
Male	466	3	1	-	1	2	6	6	7	4	23	20
Female	477	-	-	-	-	-	2	4	3	5	11	14
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	56	1	-	1	-	-	2	-	1	1	4	-
Male	28	-	-	-	-	-	1	-	-	1	2	-
Female	28	1	-	1	-	-	1	-	1	-	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Washington												
All races, both sexes	2,276	3	2	1	2	2	3	12	15	15	22	49
Male	998	-	1	1	-	1	3	10	9	9	16	31
Female	1,278	3	1	-	2	1	-	2	6	6	6	18
White, both sexes	2,155	1	1	1	2	2	3	11	12	12	18	44
Male	941	-	1	1	-	1	3	9	8	6	13	28
Female	1,214	1	-	-	2	1	-	2	4	6	5	16
Black, both sexes	46	-	-	-	-	-	-	1	1	1	2	5
Male	26	-	-	-	-	-	-	1	-	1	2	3
Female	20	-	-	-	-	-	-	-	1	-	-	2
American Indian 1/, both sexes	22	1	-	-	-	-	-	-	2	1	1	-
Male	11	-	-	-	-	-	-	-	1	1	-	-
Female	11	1	-	-	-	-	-	-	1	-	1	-
Asian or Pacific Islander 2/, both sexes	53	1	1	-	-	-	-	-	-	1	1	-
Male	20	-	-	-	-	-	-	-	-	1	1	-
Female	33	1	1	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	1,435	1	2	1	1	2	2	5	4	7	9	18
Male	622	-	2	1	1	2	1	3	1	3	6	13
Female	813	1	-	-	-	-	1	2	3	4	3	5
White, both sexes	1,390	-	1	1	1	2	2	5	4	5	9	16
Male	602	-	1	1	1	2	1	3	1	2	6	12
Female	788	-	-	-	-	-	1	2	3	3	3	4
Black, both sexes	43	1	1	-	-	-	-	-	-	1	-	2
Male	18	-	1	-	-	-	-	-	-	-	-	1
Female	25	1	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Wisconsin												
All races, both sexes	3,400	5	1	1	2	6	5	8	11	13	28	49
Male	1,446	4	-	-	1	5	4	6	8	9	24	31
Female	1,954	1	1	1	1	1	1	2	3	4	4	18
White, both sexes	3,252	2	-	1	2	6	3	6	10	11	23	38
Male	1,371	1	-	-	1	5	2	5	7	7	19	22
Female	1,881	1	-	1	1	1	1	1	3	4	4	16
Black, both sexes	125	3	1	-	-	-	2	2	-	2	5	10
Male	63	3	-	-	-	-	2	1	-	2	5	8
Female	62	-	1	-	-	-	-	1	-	-	-	2
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	1	-	-	-
Male	7	-	-	-	-	-	-	-	1	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	292	-	-	-	-	-	-	-	1	2	5	2
Male	128	-	-	-	-	-	-	-	-	1	4	2
Female	164	-	-	-	-	-	-	-	1	1	1	-
White, both sexes	287	-	-	-	-	-	-	-	1	2	4	2
Male	128	-	-	-	-	-	-	-	-	1	4	2
Female	159	-	-	-	-	-	-	-	1	1	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
United States												
All races, both sexes	4,541	5,708	6,824	8,260	11,133	17,255	26,331	31,669	25,470	10,929	2,374	4
Male	2,978	3,717	4,236	4,900	6,255	8,782	11,919	12,239	8,003	2,556	386	4
Female	1,563	1,991	2,588	3,360	4,878	8,473	14,412	19,430	17,467	8,373	1,988	-
White, both sexes	3,202	4,173	5,286	6,543	9,218	15,079	23,689	29,302	23,771	10,097	2,116	4
Male	2,152	2,749	3,324	3,943	5,220	7,749	10,786	11,428	7,511	2,357	335	4
Female	1,050	1,424	1,962	2,600	3,998	7,330	12,903	17,874	16,260	7,740	1,781	-
Black, both sexes	1,246	1,413	1,370	1,515	1,697	1,860	2,170	1,889	1,381	696	230	-
Male	761	886	812	849	905	861	915	615	377	144	40	-
Female	485	527	558	666	792	999	1,255	1,274	1,004	552	190	-
American Indian 1/, both sexes	29	49	51	66	49	72	79	67	42	16	8	-
Male	19	28	31	36	30	42	35	23	14	4	4	-
Female	10	21	20	30	19	30	44	44	28	12	4	-
Asian or Pacific Islander 2/, both sexes	64	73	117	136	169	244	393	411	276	120	20	-
Male	46	54	69	72	100	130	183	173	101	51	7	-
Female	18	19	48	64	69	114	210	238	175	69	13	-
Alabama												
All races, both sexes	221	313	309	362	494	666	890	960	693	272	78	-
Male	142	195	196	212	287	331	395	348	205	57	10	-
Female	79	118	113	150	207	335	495	612	488	215	68	-
White, both sexes	128	210	202	258	356	510	708	812	573	219	49	-
Male	81	125	134	158	218	260	317	301	169	45	6	-
Female	47	85	68	100	138	250	391	511	404	174	43	-
Black, both sexes	91	102	106	103	137	152	181	147	119	53	28	-
Male	60	70	61	54	68	67	78	47	36	12	4	-
Female	31	32	45	49	69	85	103	100	83	41	24	-
American Indian 1/, both sexes	-	1	1	1	-	2	-	1	-	-	1	-
Male	-	-	1	-	-	2	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	1	-	-	1	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	1	2	1	-	1	-	-	-
Male	1	-	-	-	1	2	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Alaska												
All races, both sexes	6	12	4	15	17	22	28	25	20	8	1	-
Male	4	9	3	7	6	15	13	12	6	1	-	-
Female	2	3	1	8	11	7	15	13	14	7	1	-
White, both sexes	2	8	4	8	11	17	22	23	19	6	1	-
Male	2	5	3	4	4	11	9	10	6	1	-	-
Female	-	3	1	4	7	6	13	13	13	5	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	3	-	7	5	5	6	2	1	2	-	-
Male	1	3	-	3	2	4	4	2	-	-	-	-
Female	1	-	-	4	3	1	2	-	1	2	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	1	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
Arizona												
All races, both sexes	66	79	102	99	154	235	323	420	310	127	25	-
Male	48	57	61	60	94	131	154	180	105	28	4	-
Female	18	22	41	39	60	104	169	240	205	99	21	-
White, both sexes	58	69	92	93	146	225	307	409	303	121	24	-
Male	41	51	56	56	91	126	149	174	104	26	4	-
Female	17	18	36	37	55	99	158	235	199	95	20	-
Black, both sexes	5	6	1	4	5	4	4	3	3	2	1	-
Male	4	3	-	2	2	2	2	2	1	1	-	-
Female	1	3	1	2	3	2	2	1	2	1	1	-
American Indian 1/, both sexes	3	4	7	2	3	5	9	7	4	4	-	-
Male	3	3	4	2	1	3	3	4	-	1	-	-
Female	-	1	3	-	2	2	6	3	4	3	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	-	1	3	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	3	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Arkansas												
All races, both sexes	43	71	97	90	138	206	304	325	265	148	28	-
Male	25	42	49	52	84	106	119	105	76	27	6	-
Female	18	29	48	38	54	100	185	220	189	121	22	-
White, both sexes	28	42	76	73	125	182	280	294	241	138	21	-
Male	17	28	40	43	77	92	111	98	69	26	5	-
Female	11	14	36	30	48	90	169	196	172	112	16	-
Black, both sexes	15	28	18	17	13	23	22	28	24	9	7	-
Male	8	14	8	9	7	13	7	7	7	1	1	-
Female	7	14	10	8	6	10	15	21	17	8	6	-
American Indian 1/, both sexes	-	1	2	-	-	1	1	3	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	2	-	-	-	1	3	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	1	-	-	1	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
California												
All races, both sexes	362	429	447	540	797	1,267	2,078	2,619	2,218	930	202	-
Male	238	283	288	327	441	645	955	1,078	770	264	32	-
Female	124	146	159	213	356	622	1,123	1,541	1,448	666	170	-
White, both sexes	272	330	348	432	665	1,108	1,864	2,378	2,045	860	191	-
Male	188	220	223	265	368	566	855	970	707	243	32	-
Female	84	110	125	167	297	542	1,009	1,408	1,338	617	159	-
Black, both sexes	67	79	61	67	77	80	89	93	75	30	5	-
Male	35	47	37	42	40	33	37	41	23	4	-	-
Female	32	32	24	25	37	47	52	52	52	26	5	-
American Indian 1/, both sexes	3	4	5	6	4	7	10	6	6	1	1	-
Male	2	3	4	3	3	5	3	3	3	1	-	-
Female	1	1	1	3	1	2	7	3	3	-	1	-
Asian or Pacific Islander 2/, both sexes	20	16	33	35	51	72	115	142	92	39	5	-
Male	13	13	24	17	30	41	60	64	37	16	-	-
Female	7	3	9	18	21	31	55	78	55	23	5	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Colorado												
All races, both sexes	42	47	63	61	101	143	257	359	299	122	24	-
Male	28	30	41	37	59	67	113	139	91	26	5	-
Female	14	17	22	24	42	76	144	220	208	96	19	-
White, both sexes	38	44	57	58	96	139	248	352	292	119	24	-
Male	25	28	36	35	56	65	111	138	87	26	5	-
Female	13	16	21	23	40	74	137	214	205	93	19	-
Black, both sexes	3	3	4	2	3	2	6	4	3	2	-	-
Male	2	2	4	1	2	1	1	1	2	-	-	-
Female	1	1	-	1	1	1	5	3	1	2	-	-
American Indian 1/, both sexes	-	-	2	1	-	1	-	-	2	-	-	-
Male	-	-	1	1	-	1	-	-	1	-	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	2	1	3	3	2	1	-	-
Male	1	-	-	-	1	-	1	-	1	-	-	-
Female	-	-	-	-	1	1	2	3	1	1	-	-
Connecticut												
All races, both sexes	69	71	94	111	158	246	390	558	447	197	43	-
Male	50	51	59	69	99	136	191	208	145	52	8	-
Female	19	20	35	42	59	110	199	350	302	145	35	-
White, both sexes	57	61	81	91	137	230	378	538	433	192	42	-
Male	44	45	52	58	88	125	187	202	143	51	8	-
Female	13	16	29	33	49	105	191	336	290	141	34	-
Black, both sexes	11	10	13	19	21	16	11	18	13	4	1	-
Male	6	6	7	10	11	11	3	5	1	1	-	-
Female	5	4	6	9	10	5	8	13	12	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	-	1	2	1	-	-	-
Male	-	-	-	1	-	-	1	1	1	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Delaware												
All races, both sexes	17	20	16	37	33	67	86	89	70	24	3	1
Male	11	14	12	23	19	25	29	36	23	6	-	1
Female	6	6	4	14	14	42	57	53	47	18	3	-
White, both sexes	13	11	11	29	29	61	74	83	63	23	3	1
Male	9	8	7	16	17	24	25	34	21	6	-	1
Female	4	3	4	13	12	37	49	49	42	17	3	-
Black, both sexes	4	8	5	8	4	6	11	5	7	1	-	-
Male	2	5	5	7	2	1	4	1	2	-	-	-
Female	2	3	-	1	2	5	7	4	5	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	9	12	13	14	18	11	36	34	40	20	4	-
Male	6	6	8	7	10	7	12	12	10	4	2	-
Female	3	6	5	7	8	4	24	22	30	16	2	-
White, both sexes	-	1	3	1	2	1	10	12	20	5	2	-
Male	-	-	1	1	1	1	3	5	8	1	1	-
Female	-	1	2	-	1	-	7	7	12	4	1	-
Black, both sexes	9	11	10	13	16	10	26	22	20	15	2	-
Male	6	6	7	6	9	6	9	7	2	3	1	-
Female	3	5	3	7	7	4	17	15	18	12	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Florida												
All races, both sexes	254	313	358	479	598	1,009	1,492	1,750	1,357	569	97	-
Male	163	208	222	311	350	575	769	778	515	165	32	-
Female	91	105	136	168	248	434	723	972	842	404	65	-
White, both sexes	166	244	286	397	519	916	1,384	1,668	1,307	547	85	-
Male	111	171	185	258	300	530	716	753	494	157	27	-
Female	55	73	101	139	219	386	668	915	813	390	58	-
Black, both sexes	87	67	69	77	76	86	98	76	48	19	11	-
Male	52	35	35	49	47	41	48	23	20	5	4	-
Female	35	32	34	28	29	45	50	53	28	14	7	-
American Indian 1/, both sexes	-	1	1	1	-	-	1	-	-	-	1	-
Male	-	1	1	1	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	2	4	3	7	9	6	2	3	-	-
Male	-	1	1	3	3	4	5	2	1	3	-	-
Female	1	-	1	1	-	3	4	4	1	-	-	-
Georgia												
All races, both sexes	268	342	412	471	541	691	999	1,060	812	349	77	-
Male	169	238	263	255	293	348	447	388	235	65	17	-
Female	99	104	149	216	248	343	552	672	577	284	60	-
White, both sexes	163	214	264	309	363	535	801	877	686	270	53	-
Male	108	145	168	179	216	271	372	327	201	47	13	-
Female	55	69	96	130	147	264	429	550	485	223	40	-
Black, both sexes	104	128	145	161	174	147	191	179	122	77	24	-
Male	60	93	94	76	75	73	73	60	34	18	4	-
Female	44	35	51	85	99	74	118	119	88	59	20	-
American Indian 1/, both sexes	-	-	-	-	1	1	1	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	3	1	3	8	6	3	4	2	-	-
Male	1	-	1	-	1	4	2	1	-	-	-	-
Female	-	-	2	1	2	4	4	2	4	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Hawaii												
All races, both sexes	35	39	39	62	46	95	144	195	131	55	15	-
Male	30	28	22	37	30	54	76	100	58	23	7	-
Female	5	11	17	25	16	41	68	95	73	32	8	-
White, both sexes	14	18	9	14	8	20	29	46	36	20	5	-
Male	12	12	6	10	4	8	17	27	21	5	2	-
Female	2	6	3	4	4	12	12	19	15	15	3	-
Black, both sexes	-	1	-	-	2	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	2	-	-	-	-
Female	-	1	-	-	2	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	1	-	1	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	21	20	30	47	36	75	114	147	94	35	10	-
Male	18	16	16	26	26	46	58	71	37	18	5	-
Female	3	4	14	21	10	29	56	76	57	17	5	-
Idaho												
All races, both sexes	10	27	22	37	42	64	119	160	114	60	10	-
Male	6	18	17	24	23	35	59	62	44	16	1	-
Female	4	9	5	13	19	29	60	98	70	44	9	-
White, both sexes	9	25	22	36	42	61	116	158	113	60	10	-
Male	6	17	17	24	23	33	56	60	43	16	1	-
Female	3	8	5	12	19	28	60	98	70	44	9	-
Black, both sexes	1	1	-	-	-	1	-	1	-	-	-	-
Male	-	1	-	-	-	1	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	1	-	2	1	-	1	-	-	-
Male	-	-	-	-	-	1	1	-	1	-	-	-
Female	-	1	-	1	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	1	-	-	-	-
Male	-	-	-	-	-	-	2	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Illinois												
All races, both sexes	179	243	258	355	458	729	1,131	1,394	1,185	575	129	-
Male	110	160	157	219	246	353	501	548	375	122	15	-
Female	69	83	101	136	212	376	630	846	810	453	114	-
White, both sexes	123	177	187	263	357	620	1,014	1,307	1,114	538	114	-
Male	76	116	105	159	194	294	447	515	353	118	13	-
Female	47	61	82	104	163	326	567	792	761	420	101	-
Black, both sexes	56	64	67	90	92	103	105	83	65	35	14	-
Male	34	43	48	60	47	56	47	32	20	4	1	-
Female	22	21	19	30	45	47	58	51	45	31	13	-
American Indian 1/, both sexes	-	1	-	1	-	-	1	-	-	-	1	-
Male	-	-	-	-	-	-	1	-	-	-	1	-
Female	-	1	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	4	1	9	6	11	4	6	2	-	-
Male	-	1	4	-	5	3	6	1	2	-	-	-
Female	-	-	-	1	4	3	5	3	4	2	-	-
Indiana												
All races, both sexes	117	165	214	229	311	465	653	812	673	305	69	-
Male	76	110	135	133	170	224	307	324	184	58	13	-
Female	41	55	79	96	141	241	346	488	489	247	56	-
White, both sexes	93	141	188	199	284	438	607	790	650	298	69	-
Male	61	92	119	116	155	214	288	315	180	55	13	-
Female	32	49	69	83	129	224	319	475	470	243	56	-
Black, both sexes	24	23	26	30	26	26	46	22	22	6	-	-
Male	15	17	16	17	14	10	19	9	4	3	-	-
Female	9	6	10	13	12	16	27	13	18	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	1	-	-	1	1	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Iowa												
All races, both sexes	16	38	43	39	65	118	177	239	253	123	23	-
Male	11	25	25	26	32	69	82	94	84	30	1	-
Female	5	13	18	13	33	49	95	145	169	93	22	-
White, both sexes	15	38	41	38	63	115	176	239	253	123	23	-
Male	11	25	25	26	31	67	81	94	84	30	1	-
Female	4	13	16	12	32	48	95	145	169	93	22	-
Black, both sexes	1	-	2	1	2	3	1	-	-	-	-	-
Male	-	-	-	-	1	2	1	-	-	-	-	-
Female	1	-	2	1	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	50	46	65	68	126	193	329	434	409	195	56	-
Male	33	32	39	36	78	82	139	166	133	45	5	-
Female	17	14	26	32	48	111	190	268	276	150	51	-
White, both sexes	40	36	55	56	115	180	313	419	397	189	52	-
Male	26	25	34	29	72	77	133	161	131	41	4	-
Female	14	11	21	27	43	103	180	258	266	148	48	-
Black, both sexes	8	8	9	9	8	10	15	11	9	4	4	-
Male	6	6	5	6	4	3	6	5	2	4	1	-
Female	2	2	4	3	4	7	9	6	7	-	3	-
American Indian 1/, both sexes	1	1	1	3	2	3	-	3	-	2	-	-
Male	1	-	-	1	2	2	-	-	-	-	-	-
Female	-	1	1	2	-	1	-	3	-	2	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	1	-	1	1	3	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	1	1	3	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Kentucky												
All races, both sexes	93	139	136	210	233	384	535	530	421	169	34	-
Male	67	87	78	121	125	190	223	178	116	32	5	-
Female	26	52	58	89	108	194	312	352	305	137	29	-
White, both sexes	84	122	129	189	217	358	505	511	399	163	30	-
Male	64	75	74	111	113	181	212	170	106	31	5	-
Female	20	47	55	78	104	177	293	341	293	132	25	-
Black, both sexes	9	15	7	19	16	26	30	19	21	6	4	-
Male	3	10	4	9	12	9	11	8	9	1	-	-
Female	6	5	3	10	4	17	19	11	12	5	4	-
American Indian 1/, both sexes	-	2	-	1	-	-	-	-	1	-	-	-
Male	-	2	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	99	122	143	196	245	351	557	570	421	164	38	-
Male	63	73	80	130	131	178	224	208	109	35	7	-
Female	36	49	63	66	114	173	333	362	312	129	31	-
White, both sexes	44	70	86	121	167	262	436	488	340	138	33	-
Male	28	42	51	78	89	143	175	180	96	29	6	-
Female	16	28	35	43	78	119	261	308	244	109	27	-
Black, both sexes	55	51	56	74	76	89	118	81	79	25	5	-
Male	35	31	29	51	42	35	47	27	13	6	1	-
Female	20	20	27	23	34	54	71	54	66	19	4	-
American Indian 1/, both sexes	-	1	1	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	2	-	3	-	2	1	-	-
Male	-	-	-	1	-	-	2	-	-	-	-	-
Female	-	-	-	-	2	-	1	-	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Maine												
All races, both sexes	22	27	50	43	70	106	151	196	143	66	16	-
Male	17	17	34	24	45	56	74	76	48	22	2	-
Female	5	10	16	19	25	50	77	120	95	44	14	-
White, both sexes	22	27	49	43	69	106	150	196	143	65	16	-
Male	17	17	33	24	44	56	73	76	48	21	2	-
Female	5	10	16	19	25	50	77	120	95	44	14	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	1	-	1	-	-	-	-	-
Male	-	-	1	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	62	76	104	131	157	238	369	453	318	149	27	-
Male	44	41	71	75	87	119	164	178	114	36	7	-
Female	18	35	33	56	70	119	205	275	204	113	20	-
White, both sexes	22	46	56	74	103	171	301	382	287	119	23	-
Male	18	25	38	41	54	93	134	154	105	28	6	-
Female	4	21	18	33	49	78	167	228	182	91	17	-
Black, both sexes	40	30	45	54	51	59	63	63	28	28	4	-
Male	26	16	31	32	32	21	29	21	8	8	1	-
Female	14	14	14	22	19	38	34	42	20	20	3	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	3	3	8	5	8	2	2	-	-
Male	-	-	2	2	1	5	1	3	1	-	-	-
Female	-	-	-	1	2	3	4	5	1	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Massachusetts												
All races, both sexes	66	74	106	146	207	373	663	853	740	317	87	-
Male	40	55	60	92	130	204	284	306	222	71	14	-
Female	26	19	46	54	77	169	379	547	518	246	73	-
White, both sexes	58	64	87	131	190	357	637	829	730	305	83	-
Male	35	46	48	82	121	194	272	299	219	68	13	-
Female	23	18	39	49	69	163	365	530	511	237	70	-
Black, both sexes	7	9	16	11	15	13	19	20	7	9	4	-
Male	4	8	11	7	8	7	11	4	3	1	1	-
Female	3	1	5	4	7	6	8	16	4	8	3	-
American Indian 1/, both sexes	-	-	-	2	-	1	1	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	3	2	2	2	6	4	3	3	-	-
Male	1	1	1	2	1	2	1	3	-	2	-	-
Female	-	-	2	-	1	-	5	1	3	1	-	-
Michigan												
All races, both sexes	180	173	200	241	351	560	899	1,124	948	397	88	-
Male	127	109	123	149	175	290	408	419	288	91	10	-
Female	53	64	77	92	176	270	491	705	660	306	78	-
White, both sexes	114	111	155	198	293	491	798	1,065	895	374	76	-
Male	79	75	99	119	148	254	367	399	269	84	8	-
Female	35	36	56	79	145	237	431	666	626	290	68	-
Black, both sexes	65	61	43	40	55	67	92	54	49	20	10	-
Male	47	33	22	29	26	35	39	19	16	7	1	-
Female	18	28	21	11	29	32	53	35	33	13	9	-
American Indian 1/, both sexes	1	-	-	-	2	1	4	3	1	-	1	-
Male	1	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	4	3	1	-	1	-
Asian or Pacific Islander 2/, both sexes	-	1	2	3	1	1	5	2	3	3	1	-
Male	-	1	2	1	-	-	2	1	3	-	1	-
Female	-	-	-	2	1	1	3	1	-	3	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Minnesota												
All races, both sexes	47	47	81	86	128	264	393	611	565	266	54	-
Male	35	32	49	55	79	141	208	257	178	67	6	-
Female	12	15	32	31	49	123	185	354	387	199	48	-
White, both sexes	41	41	77	78	124	254	383	606	560	265	53	-
Male	31	29	47	50	77	136	202	257	176	67	6	-
Female	10	12	30	28	47	118	181	349	384	198	47	-
Black, both sexes	4	4	2	4	2	9	4	2	4	-	1	-
Male	3	3	1	3	2	5	3	-	2	-	-	-
Female	1	1	1	1	-	4	1	2	2	-	1	-
American Indian 1/, both sexes	2	2	2	4	1	-	3	1	1	-	-	-
Male	1	-	1	2	-	-	2	-	-	-	-	-
Female	1	2	1	2	1	-	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	3	2	-	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	1	2	2	-	1	-	-
Mississippi												
All races, both sexes	100	123	152	191	250	328	485	498	367	200	49	1
Male	61	83	86	105	137	148	212	183	91	44	14	1
Female	39	40	66	86	113	180	273	315	276	156	35	-
White, both sexes	47	60	84	113	163	225	370	380	280	139	24	1
Male	29	40	49	64	96	101	176	148	75	29	7	1
Female	18	20	35	49	67	124	194	232	205	110	17	-
Black, both sexes	53	63	67	76	85	101	114	117	87	60	25	-
Male	32	43	37	40	39	45	36	35	16	15	7	-
Female	21	20	30	36	46	56	78	82	71	45	18	-
American Indian 1/, both sexes	-	-	1	2	1	1	-	1	-	1	-	-
Male	-	-	-	1	1	1	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	1	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Missouri												
All races, both sexes	90	104	161	193	269	425	598	709	652	274	62	-
Male	60	69	84	112	147	223	277	251	205	68	8	-
Female	30	35	77	81	122	202	321	458	447	206	54	-
White, both sexes	73	83	141	177	239	389	569	673	630	262	59	-
Male	48	57	73	101	130	209	266	241	195	64	7	-
Female	25	26	68	76	109	180	303	432	435	198	52	-
Black, both sexes	16	21	20	15	29	34	28	34	21	12	3	-
Male	11	12	11	10	16	13	11	10	10	4	1	-
Female	5	9	9	5	13	21	17	24	11	8	2	-
American Indian 1/, both sexes	1	-	-	-	1	-	-	1	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	2	1	1	1	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	1	-	-	-
Montana												
All races, both sexes	10	15	19	34	43	68	113	155	150	60	9	-
Male	9	10	12	22	25	33	55	67	55	17	3	-
Female	1	5	7	12	18	35	58	88	95	43	6	-
White, both sexes	10	11	18	30	41	64	111	155	150	60	9	-
Male	9	8	11	19	23	32	54	67	55	17	3	-
Female	1	3	7	11	18	32	57	88	95	43	6	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	4	1	4	2	4	1	-	-	-	-	-
Male	-	2	1	3	2	1	1	-	-	-	-	-
Female	-	2	-	1	-	3	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Nebraska												
All races, both sexes	33	37	41	72	75	108	190	248	261	129	27	-
Male	22	26	24	55	46	62	86	87	77	21	3	-
Female	11	11	17	17	29	46	104	161	184	108	24	-
White, both sexes	29	33	38	67	73	104	185	245	254	127	27	-
Male	19	25	23	53	44	59	86	87	77	21	3	-
Female	10	8	15	14	29	45	99	158	177	106	24	-
Black, both sexes	3	4	2	3	1	4	5	1	7	2	-	-
Male	2	1	1	1	1	3	-	-	-	-	-	-
Female	1	3	1	2	-	1	5	1	7	2	-	-
American Indian 1/, both sexes	1	-	1	2	-	-	-	1	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Nevada												
All races, both sexes	67	100	114	158	166	224	301	259	146	40	11	1
Male	48	66	74	104	99	130	166	110	47	11	1	1
Female	19	34	40	54	67	94	135	149	99	29	10	-
White, both sexes	47	73	98	130	139	197	278	242	139	36	9	1
Male	35	49	65	88	84	121	154	106	44	10	1	1
Female	12	24	33	42	55	76	124	136	95	26	8	-
Black, both sexes	16	20	12	19	20	20	15	6	4	3	1	-
Male	11	13	7	12	11	9	10	1	2	-	-	-
Female	5	7	5	7	9	11	5	5	2	3	1	-
American Indian 1/, both sexes	1	2	1	1	1	1	1	1	2	-	-	-
Male	-	2	1	1	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	1	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes	3	5	3	8	6	6	7	10	1	1	1	-
Male	2	2	1	3	3	-	2	3	1	1	-	-
Female	1	3	2	5	3	6	5	7	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
New Hampshire												
All races, both sexes	13	15	32	25	41	65	104	148	108	46	18	-
Male	13	12	22	16	22	32	42	69	29	8	3	-
Female	-	3	10	9	19	33	62	79	79	38	15	-
White, both sexes	13	15	32	24	41	61	104	148	107	46	18	-
Male	13	12	22	15	22	32	42	69	29	8	3	-
Female	-	3	10	9	19	29	62	79	78	38	15	-
Black, both sexes	-	-	-	1	-	1	-	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	3	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	3	-	-	-	-	-	-
New Jersey												
All races, both sexes	93	124	153	184	310	461	825	1,063	787	330	68	-
Male	57	73	98	111	184	232	389	422	263	78	15	-
Female	36	51	55	73	126	229	436	641	524	252	53	-
White, both sexes	69	90	120	132	248	387	757	999	753	309	63	-
Male	47	58	81	81	149	200	352	398	254	71	15	-
Female	22	32	39	51	99	187	405	601	499	238	48	-
Black, both sexes	23	33	29	46	57	68	58	56	28	14	4	-
Male	9	14	16	25	31	30	32	21	8	4	-	-
Female	14	19	13	21	26	38	26	35	20	10	4	-
American Indian 1/, both sexes	-	-	-	2	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	4	4	5	6	10	8	6	7	1	-
Male	1	1	1	4	4	2	5	3	1	3	-	-
Female	-	-	3	-	1	4	5	5	5	4	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
New Mexico												
All races, both sexes	19	32	34	36	46	94	157	171	134	50	13	-
Male	11	18	17	23	29	48	73	62	45	15	5	-
Female	8	14	17	13	17	46	84	109	89	35	8	-
White, both sexes	18	27	32	34	42	86	148	168	131	48	11	-
Male	10	16	15	22	27	41	67	60	43	15	4	-
Female	8	11	17	12	15	45	81	108	88	33	7	-
Black, both sexes	-	1	-	-	2	2	4	-	-	2	-	-
Male	-	-	-	-	2	2	3	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	2	-	-
American Indian 1/, both sexes	1	3	2	2	2	5	4	2	3	-	2	-
Male	1	2	2	1	-	5	2	1	2	-	1	-
Female	-	1	-	1	2	-	2	1	1	-	1	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
New York												
All races, both sexes	196	226	289	315	502	783	1,315	1,661	1,406	597	146	-
Male	136	144	188	180	291	395	597	619	452	135	17	-
Female	60	82	101	135	211	388	718	1,042	954	462	129	-
White, both sexes	150	174	234	259	434	697	1,233	1,573	1,347	578	136	-
Male	106	114	151	148	246	356	560	602	439	132	17	-
Female	44	60	83	111	188	341	673	971	908	446	119	-
Black, both sexes	41	46	51	51	57	76	67	77	56	15	10	-
Male	26	27	36	29	37	36	31	13	13	2	-	-
Female	15	19	15	22	20	40	36	64	43	13	10	-
American Indian 1/, both sexes	1	1	-	1	1	1	1	1	-	-	-	-
Male	-	1	-	-	1	-	-	1	-	-	-	-
Female	1	-	-	1	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	5	4	4	10	9	14	10	3	4	-	-
Male	4	2	1	3	7	3	6	3	-	1	-	-
Female	-	3	3	1	3	6	8	7	3	3	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
North Carolina												
All races, both sexes	148	180	245	276	397	562	833	884	669	271	67	-
Male	98	111	159	149	228	277	358	338	198	52	6	-
Female	50	69	86	127	169	285	475	546	471	219	61	-
White, both sexes	88	114	173	180	292	451	693	766	599	218	54	-
Male	65	74	113	104	169	223	308	307	177	42	3	-
Female	23	40	60	76	123	228	385	459	422	176	51	-
Black, both sexes	60	64	67	92	101	106	132	109	66	53	12	-
Male	33	36	43	43	57	51	48	29	20	10	3	-
Female	27	28	24	49	44	55	84	80	46	43	9	-
American Indian 1/, both sexes	-	1	2	4	4	3	5	7	3	-	-	-
Male	-	-	2	2	2	1	2	2	1	-	-	-
Female	-	1	-	2	2	2	3	5	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	3	-	-	2	3	2	1	-	1	-
Male	-	1	1	-	-	2	-	-	-	-	-	-
Female	-	-	2	-	-	-	3	2	1	-	1	-
North Dakota												
All races, both sexes	9	9	9	8	10	17	27	68	57	28	8	-
Male	7	7	6	2	6	15	14	27	18	5	-	-
Female	2	2	3	6	4	2	13	41	39	23	8	-
White, both sexes	8	6	6	8	10	16	27	66	57	28	8	-
Male	6	4	5	2	6	14	14	26	18	5	-	-
Female	2	2	1	6	4	2	13	40	39	23	8	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	3	3	-	-	1	-	2	-	-	-	-
Male	1	3	1	-	-	1	-	1	-	-	-	-
Female	-	-	2	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Ohio												
All races, both sexes	177	211	249	269	402	621	986	1,118	902	336	61	-
Male	102	144	158	145	195	303	428	407	278	79	8	-
Female	75	67	91	124	207	318	558	711	624	257	53	-
White, both sexes	135	173	201	224	349	575	917	1,065	853	320	58	-
Male	82	123	131	123	169	278	391	389	263	76	7	-
Female	53	50	70	101	180	297	526	676	590	244	51	-
Black, both sexes	41	38	48	44	52	46	66	51	47	16	3	-
Male	19	21	27	22	26	25	35	18	15	3	1	-
Female	22	17	21	22	26	21	31	33	32	13	2	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	2	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	3	2	-	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	1	1	-	1	2	-	-	-	-
Oklahoma												
All races, both sexes	78	85	123	150	201	296	427	480	358	151	33	-
Male	41	51	79	78	94	145	191	190	113	27	7	-
Female	37	34	44	72	107	151	236	290	245	124	26	-
White, both sexes	56	63	99	125	174	266	392	446	335	142	31	-
Male	33	41	63	67	83	132	176	182	106	23	6	-
Female	23	22	36	58	91	134	216	264	229	119	25	-
Black, both sexes	18	12	12	14	18	17	18	24	16	6	1	-
Male	7	7	7	6	7	8	7	5	4	2	-	-
Female	11	5	5	8	11	9	11	19	12	4	1	-
American Indian 1/, both sexes	4	9	10	9	9	13	16	10	7	2	1	-
Male	1	3	8	5	4	5	7	3	3	1	1	-
Female	3	6	2	4	5	8	9	7	4	1	-	-
Asian or Pacific Islander 2/, both sexes	-	1	2	2	-	-	1	-	-	1	-	-
Male	-	-	1	-	-	-	1	-	-	1	-	-
Female	-	1	1	2	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Oregon												
All races, both sexes	46	51	54	87	117	207	375	497	449	183	44	-
Male	36	27	28	53	63	103	167	187	156	56	8	-
Female	10	24	26	34	54	104	208	310	293	127	36	-
White, both sexes	44	51	52	84	113	206	362	488	445	181	44	-
Male	34	27	26	51	61	102	161	186	155	55	8	-
Female	10	24	26	33	52	104	201	302	290	126	36	-
Black, both sexes	1	-	-	-	1	-	1	1	1	-	-	-
Male	1	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
American Indian 1/, both sexes	1	-	2	3	2	1	2	2	-	-	-	-
Male	1	-	2	2	1	1	2	-	-	-	-	-
Female	-	-	-	1	1	-	-	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	10	6	3	2	-	-
Male	-	-	-	-	-	-	3	1	1	1	-	-
Female	-	-	-	-	1	-	7	5	2	1	-	-
Pennsylvania												
All races, both sexes	230	313	385	454	597	1,067	1,684	2,226	1,647	636	134	-
Male	149	204	253	260	335	536	728	846	494	149	17	-
Female	81	109	132	194	262	531	956	1,380	1,153	487	117	-
White, both sexes	170	248	337	400	543	989	1,582	2,135	1,573	609	127	-
Male	112	162	222	235	300	505	681	812	474	145	16	-
Female	58	86	115	165	243	484	901	1,323	1,099	464	111	-
Black, both sexes	59	63	46	52	48	75	98	85	66	26	7	-
Male	36	40	30	24	32	30	47	31	17	3	1	-
Female	23	23	16	28	16	45	51	54	49	23	6	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	2	2	6	3	4	5	8	1	-	-
Male	1	2	1	1	3	1	-	3	3	1	-	-
Female	-	-	1	1	3	2	4	2	5	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Rhode Island												
All races, both sexes	5	4	19	18	23	42	73	92	67	39	5	-
Male	4	3	13	14	16	15	29	32	20	13	-	-
Female	1	1	6	4	7	27	44	60	47	26	5	-
White, both sexes	5	3	19	16	22	42	70	90	67	38	5	-
Male	4	3	13	13	15	15	29	32	20	12	-	-
Female	1	-	6	3	7	27	41	58	47	26	5	-
Black, both sexes	-	1	-	2	1	-	2	1	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
South Carolina												
All races, both sexes	135	180	209	219	260	359	448	469	332	134	25	1
Male	92	127	129	138	146	175	202	167	90	28	4	1
Female	43	53	80	81	114	184	246	302	242	106	21	-
White, both sexes	67	87	122	133	178	262	353	387	266	107	18	1
Male	43	62	77	86	98	134	162	151	73	25	4	1
Female	24	25	45	47	80	128	191	236	193	82	14	-
Black, both sexes	68	91	86	86	82	96	93	81	65	27	7	-
Male	49	64	52	52	48	40	39	16	17	3	-	-
Female	19	27	34	34	34	56	54	65	48	24	7	-
American Indian 1/, both sexes	-	1	1	-	-	-	1	-	1	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
South Dakota												
All races, both sexes	8	5	5	14	25	27	38	56	60	32	6	-
Male	3	3	2	10	18	16	14	16	20	11	1	-
Female	5	2	3	4	7	11	24	40	40	21	5	-
White, both sexes	7	5	3	13	23	26	37	54	58	32	6	-
Male	3	3	2	9	17	15	14	15	19	11	1	-
Female	4	2	1	4	6	11	23	39	39	21	5	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	1	2	1	1	2	2	-	-	-
Male	-	-	-	1	1	1	-	1	1	-	-	-
Female	1	-	1	-	1	-	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	87	104	135	168	196	312	426	457	325	132	24	-
Male	53	74	78	96	114	154	180	137	82	19	2	-
Female	34	30	57	72	82	158	246	320	243	113	22	-
White, both sexes	56	69	106	134	162	278	359	408	298	113	21	-
Male	36	51	62	78	103	139	157	121	77	17	1	-
Female	20	18	44	56	59	139	202	287	221	96	20	-
Black, both sexes	31	35	29	33	34	33	63	49	27	17	3	-
Male	17	23	16	18	11	15	22	16	5	1	1	-
Female	14	12	13	15	23	18	41	33	22	16	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	4	-	-	2	-	-
Male	-	-	-	-	-	-	1	-	-	1	-	-
Female	-	-	-	1	-	1	3	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Texas												
All races, both sexes	335	422	483	605	796	1,265	1,627	1,878	1,413	636	132	-
Male	215	266	303	344	465	634	732	729	429	136	27	-
Female	120	156	180	261	331	631	895	1,149	984	500	105	-
White, both sexes	252	308	382	504	668	1,136	1,470	1,723	1,315	573	114	-
Male	161	189	248	289	392	572	661	669	404	123	22	-
Female	91	119	134	215	276	564	809	1,054	911	450	92	-
Black, both sexes	79	107	97	96	119	114	137	140	80	60	18	-
Male	52	70	54	52	68	54	59	56	20	12	5	-
Female	27	37	43	44	51	60	78	84	60	48	13	-
American Indian 1/, both sexes	1	2	-	1	1	3	3	2	1	-	-	-
Male	1	2	-	1	1	2	1	1	-	-	-	-
Female	-	-	-	-	-	1	2	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	3	5	4	4	8	12	17	13	17	3	-	-
Male	1	5	1	2	4	6	11	3	5	1	-	-
Female	2	-	3	2	4	6	6	10	12	2	-	-
Utah												
All races, both sexes	23	32	33	76	87	145	263	243	205	96	7	-
Male	17	16	22	50	37	75	118	90	72	39	3	-
Female	6	16	11	26	50	70	145	153	133	57	4	-
White, both sexes	21	30	32	72	85	143	258	240	200	96	7	-
Male	15	16	21	48	36	74	115	89	70	39	3	-
Female	6	14	11	24	49	69	143	151	130	57	4	-
Black, both sexes	1	-	-	1	-	-	-	-	1	-	-	-
Male	1	-	-	1	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	1	-	-	1	-	-	-	-	-
Male	1	-	-	1	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	2	2	2	4	3	4	-	-	-
Male	-	-	1	-	1	1	2	1	1	-	-	-
Female	-	1	-	2	1	1	2	2	3	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Vermont												
All races, both sexes	4	7	9	11	13	26	39	51	43	22	4	-
Male	2	6	5	7	9	9	19	25	13	5	2	-
Female	2	1	4	4	4	17	20	26	30	17	2	-
White, both sexes	4	7	9	11	13	26	39	51	43	22	4	-
Male	2	6	5	7	9	9	19	25	13	5	2	-
Female	2	1	4	4	4	17	20	26	30	17	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	134	194	229	253	355	539	764	914	669	282	61	-
Male	88	119	138	146	199	278	347	348	203	64	7	-
Female	46	75	91	107	156	261	417	566	466	218	54	-
White, both sexes	82	108	143	181	254	416	639	803	586	247	52	-
Male	51	65	92	110	142	216	290	311	178	57	5	-
Female	31	43	51	71	112	200	349	492	408	190	47	-
Black, both sexes	51	85	81	70	93	117	116	104	75	31	8	-
Male	36	54	41	36	52	59	53	35	20	6	1	-
Female	15	31	40	34	41	58	63	69	55	25	7	-
American Indian 1/, both sexes	-	-	-	-	1	3	-	-	-	2	-	-
Male	-	-	-	-	1	1	-	-	-	1	-	-
Female	-	-	-	-	-	2	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	1	5	2	7	3	9	7	8	2	1	-
Male	1	-	5	-	4	2	4	2	5	-	1	-
Female	-	1	-	2	3	1	5	5	3	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Washington												
All races, both sexes	60	66	96	92	142	187	354	479	444	195	35	-
Male	39	44	49	61	85	92	149	203	140	48	7	-
Female	21	22	47	31	57	95	205	276	304	147	28	-
White, both sexes	52	60	84	85	132	174	338	462	433	194	34	-
Male	34	39	44	55	80	87	144	195	138	48	7	-
Female	18	21	40	30	52	87	194	267	295	146	27	-
Black, both sexes	5	2	6	2	7	4	3	3	2	1	1	-
Male	3	2	3	2	3	3	1	2	-	-	-	-
Female	2	-	3	-	4	1	2	1	2	1	1	-
American Indian 1/, both sexes	2	-	2	-	1	4	3	4	1	-	-	-
Male	1	-	1	-	1	1	2	2	1	-	-	-
Female	1	-	1	-	-	3	1	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	4	4	5	2	5	10	10	8	-	-	-
Male	1	3	1	4	1	1	2	4	1	-	-	-
Female	-	1	3	1	1	4	8	6	7	-	-	-
West Virginia												
All races, both sexes	32	48	73	90	105	185	252	259	227	90	22	-
Male	23	31	51	53	54	89	111	101	62	11	3	-
Female	9	17	22	37	51	96	141	158	165	79	19	-
White, both sexes	31	44	69	88	104	179	246	251	224	87	21	-
Male	22	28	48	52	53	86	108	100	61	11	3	-
Female	9	16	21	36	51	93	138	151	163	76	18	-
Black, both sexes	1	4	3	2	1	6	6	8	3	3	1	-
Male	1	3	2	1	1	3	3	1	1	-	-	-
Female	-	1	1	1	-	3	3	7	2	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Wisconsin												
All races, both sexes	67	87	83	126	193	312	576	759	686	308	74	-
Male	44	56	57	77	106	168	264	309	201	66	6	-
Female	23	31	26	49	87	144	312	450	485	242	68	-
White, both sexes	55	72	73	116	172	300	563	746	676	304	73	-
Male	38	48	52	73	93	162	262	304	198	66	6	-
Female	17	24	21	43	79	138	301	442	478	238	67	-
Black, both sexes	12	14	9	7	17	8	12	9	9	2	1	-
Male	6	7	4	3	10	3	2	4	3	-	-	-
Female	6	7	5	4	7	5	10	5	6	2	1	-
American Indian 1/, both sexes	-	-	-	2	2	2	-	-	1	1	-	-
Male	-	-	-	1	2	1	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	1	1	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	1	2	2	1	4	-	1	-	-
Male	-	1	1	-	1	2	-	1	-	-	-	-
Female	-	-	-	1	1	-	1	3	-	1	-	-
Wyoming												
All races, both sexes	9	9	14	14	24	27	48	57	54	25	1	-
Male	5	7	9	8	12	14	25	17	16	8	-	-
Female	4	2	5	6	12	13	23	40	38	17	1	-
White, both sexes	9	9	14	14	23	27	48	56	53	24	1	-
Male	5	7	9	8	12	14	25	17	16	8	-	-
Female	4	2	5	6	11	13	23	39	37	16	1	-
Black, both sexes	-	-	-	-	1	-	-	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
United States												
All races, both sexes	1,225	3	1	1	-	1	8	16	16	36	59	82
Male	718	2	1	1	-	-	4	14	14	21	44	46
Female	507	1	-	-	-	1	4	2	2	15	15	36
White, both sexes	967	2	-	1	-	1	8	12	11	27	43	53
Male	554	1	-	1	-	-	4	11	9	18	32	27
Female	413	1	-	-	-	1	4	1	2	9	11	26
Black, both sexes	223	1	1	-	-	-	-	1	4	8	15	26
Male	144	1	1	-	-	-	-	1	4	3	11	18
Female	79	-	-	-	-	-	-	-	-	5	4	8
American Indian 1/, both sexes	14	-	-	-	-	-	-	2	1	-	-	3
Male	9	-	-	-	-	-	-	2	1	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	21	-	-	-	-	-	-	1	-	1	1	-
Male	11	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	1	-	1	-	-
Alabama												
All races, both sexes	18	-	-	-	-	-	1	-	-	-	1	-
Male	10	-	-	-	-	-	1	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	14	-	-	-	-	-	1	-	-	-	1	-
Male	8	-	-	-	-	-	1	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Alaska												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	35	-	-	-	-	-	-	-	1	3	1	-
Male	21	-	-	-	-	-	-	-	-	2	1	-
Female	14	-	-	-	-	-	-	-	1	1	-	-
White, both sexes	33	-	-	-	-	-	-	-	1	3	1	-
Male	20	-	-	-	-	-	-	-	-	2	1	-
Female	13	-	-	-	-	-	-	-	1	1	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Arkansas												
All races, both sexes	17	-	-	-	-	-	1	-	1	-	2	1
Male	9	-	-	-	-	-	1	-	1	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	14	-	-	-	-	-	1	-	-	-	2	1
Male	7	-	-	-	-	-	1	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	3	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	141	-	-	-	-	-	1	1	3	1	8	14
Male	86	-	-	-	-	-	-	1	3	1	6	4
Female	55	-	-	-	-	-	1	-	-	-	2	10
White, both sexes	114	-	-	-	-	-	1	1	2	1	6	12
Male	73	-	-	-	-	-	-	1	2	1	4	4
Female	41	-	-	-	-	-	1	-	-	-	2	8
Black, both sexes	15	-	-	-	-	-	-	-	1	-	1	1
Male	7	-	-	-	-	-	-	-	1	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Colorado												
All races, both sexes	11	-	-	-	-	1	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	1	1
Female	7	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	1	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Delaware												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Florida												
All races, both sexes	74	-	-	-	-	-	-	-	1	5	3	6
Male	40	-	-	-	-	-	-	-	1	2	1	4
Female	34	-	-	-	-	-	-	-	-	3	2	2
White, both sexes	57	-	-	-	-	-	-	-	-	1	3	2
Male	28	-	-	-	-	-	-	-	-	1	1	-
Female	29	-	-	-	-	-	-	-	-	-	2	2
Black, both sexes	16	-	-	-	-	-	-	-	1	3	-	4
Male	12	-	-	-	-	-	-	-	1	1	-	4
Female	4	-	-	-	-	-	-	-	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Georgia												
All races, both sexes	28	1	1	-	-	-	-	1	-	-	2	1
Male	18	1	1	-	-	-	-	1	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	18	1	-	-	-	-	-	1	-	-	2	1
Male	11	1	-	-	-	-	-	1	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	10	-	1	-	-	-	-	-	-	-	-	-
Male	7	-	1	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Hawaii												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	7	-	-	-	-	-	-	-	1	-	-	1
Male	4	-	-	-	-	-	-	-	1	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	1	-	-	1
Male	4	-	-	-	-	-	-	-	1	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Illinois												
All races, both sexes	47	-	-	-	-	-	-	-	1	1	1	1
Male	30	-	-	-	-	-	-	-	1	-	1	1
Female	17	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	30	-	-	-	-	-	-	-	-	-	1	1
Male	19	-	-	-	-	-	-	-	-	-	1	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	17	-	-	-	-	-	-	-	1	1	-	-
Male	11	-	-	-	-	-	-	-	1	-	-	-
Female	6	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	25	-	-	-	-	-	-	-	-	-	3	-
Male	12	-	-	-	-	-	-	-	-	-	1	-
Female	13	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	21	-	-	-	-	-	-	-	-	-	2	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Iowa												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	1	4
Male	7	-	-	-	-	-	-	-	-	-	1	2
Female	9	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	12	-	-	-	-	-	-	-	-	-	1	1
Male	6	-	-	-	-	-	-	-	-	-	1	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	2
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Kentucky												
All races, both sexes	12	1	-	-	-	-	-	1	-	-	-	-
Male	8	-	-	-	-	-	-	1	-	-	-	-
Female	4	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	1	-	-	-	-	-	1	-	-	-	-
Male	7	-	-	-	-	-	-	1	-	-	-	-
Female	4	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	22	-	-	-	-	-	-	1	-	1	3	1
Male	12	-	-	-	-	-	-	-	-	-	1	1
Female	10	-	-	-	-	-	-	1	-	1	2	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	10	-	-	-	-	-	-	-	-	1	3	1
Male	6	-	-	-	-	-	-	-	-	-	1	1
Female	4	-	-	-	-	-	-	-	-	1	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Maine												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	23	-	-	-	-	-	-	-	-	1	2	2
Male	11	-	-	-	-	-	-	-	-	-	1	2
Female	12	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	11	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	12	-	-	-	-	-	-	-	-	-	2	2
Male	8	-	-	-	-	-	-	-	-	-	1	2
Female	4	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Massachusetts												
All races, both sexes	23	-	-	-	-	-	-	2	-	2	1	1
Male	14	-	-	-	-	-	-	1	-	1	1	1
Female	9	-	-	-	-	-	-	1	-	1	-	-
White, both sexes	21	-	-	-	-	-	-	2	-	2	-	1
Male	12	-	-	-	-	-	-	1	-	1	-	1
Female	9	-	-	-	-	-	-	1	-	1	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	44	-	-	-	-	-	-	1	-	1	1	3
Male	24	-	-	-	-	-	-	1	-	-	1	1
Female	20	-	-	-	-	-	-	-	-	1	-	2
White, both sexes	32	-	-	-	-	-	-	1	-	1	-	1
Male	17	-	-	-	-	-	-	1	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	12	-	-	-	-	-	-	-	-	-	1	2
Male	7	-	-	-	-	-	-	-	-	-	1	1
Female	5	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Minnesota												
All races, both sexes	21	-	-	-	-	-	-	-	-	-	-	3
Male	12	-	-	-	-	-	-	-	-	-	-	2
Female	9	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	3
Male	11	-	-	-	-	-	-	-	-	-	-	2
Female	8	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	13	-	-	1	-	-	-	-	-	1	2	-
Male	8	-	-	1	-	-	-	-	-	-	2	-
Female	5	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	9	-	-	1	-	-	-	-	-	1	1	-
Male	5	-	-	1	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Missouri												
All races, both sexes	22	-	-	-	-	-	-	-	-	1	-	2
Male	14	-	-	-	-	-	-	-	-	1	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	18	-	-	-	-	-	-	-	-	1	-	1
Male	10	-	-	-	-	-	-	-	-	1	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Nebraska												
All races, both sexes	9	-	-	-	-	-	-	-	1	-	2	-
Male	6	-	-	-	-	-	-	-	1	-	2	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	1	-	2	-
Male	6	-	-	-	-	-	-	-	1	-	2	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	11	-	-	-	-	-	1	1	1	-	1	1
Male	9	-	-	-	-	-	-	1	1	-	1	1
Female	2	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	1	-	1	-	1	1
Male	7	-	-	-	-	-	-	-	1	-	1	1
Female	2	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
New Hampshire												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	38	-	-	-	-	-	-	-	-	3	-	1
Male	21	-	-	-	-	-	-	-	-	3	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	31	-	-	-	-	-	-	-	-	3	-	1
Male	18	-	-	-	-	-	-	-	-	3	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
New Mexico												
All races, both sexes	10	-	-	-	-	-	-	2	1	-	1	-
Male	8	-	-	-	-	-	-	2	1	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	1	-	-	-	-
Male	4	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	1	1	-	-	-
Male	3	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	87	-	-	-	-	-	-	-	-	2	2	3
Male	53	-	-	-	-	-	-	-	-	1	2	2
Female	34	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	70	-	-	-	-	-	-	-	-	2	1	3
Male	40	-	-	-	-	-	-	-	-	1	1	2
Female	30	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	16	-	-	-	-	-	-	-	-	-	1	-
Male	12	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
North Carolina												
All races, both sexes	29	1	-	-	-	-	-	-	-	-	-	2
Male	16	1	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	12	1	-	-	-	-	-	-	-	-	-	2
Male	8	1	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Ohio												
All races, both sexes	58	-	-	-	-	-	1	-	1	2	1	4
Male	27	-	-	-	-	-	1	-	-	1	1	1
Female	31	-	-	-	-	-	-	-	1	1	-	3
White, both sexes	46	-	-	-	-	-	1	-	1	1	1	4
Male	21	-	-	-	-	-	1	-	-	-	1	1
Female	25	-	-	-	-	-	-	-	1	1	-	3
Black, both sexes	12	-	-	-	-	-	-	-	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	1	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	14	-	-	-	-	-	-	2	1	-	-	1
Male	9	-	-	-	-	-	-	2	1	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	10	-	-	-	-	-	-	2	1	-	-	-
Male	7	-	-	-	-	-	-	2	1	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Oregon												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	1	1
Male	10	-	-	-	-	-	-	-	-	-	1	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	1	1
Male	10	-	-	-	-	-	-	-	-	-	1	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	61	-	-	-	-	-	-	-	1	1	4	4
Male	37	-	-	-	-	-	-	-	1	-	4	3
Female	24	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	57	-	-	-	-	-	-	-	1	1	3	4
Male	33	-	-	-	-	-	-	-	1	-	3	3
Female	24	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Rhode Island												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	1	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	1	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	20	-	-	-	-	-	-	1	-	1	1	1
Male	10	-	-	-	-	-	-	1	-	1	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	12	-	-	-	-	-	-	-	-	1	1	-
Male	6	-	-	-	-	-	-	-	-	1	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	1	-	-	-	1
Male	4	-	-	-	-	-	-	1	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
South Dakota												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	33	-	-	-	-	-	-	1	-	1	2	5
Male	17	-	-	-	-	-	-	1	-	1	1	1
Female	16	-	-	-	-	-	-	-	-	-	1	4
White, both sexes	28	-	-	-	-	-	-	1	-	1	1	3
Male	16	-	-	-	-	-	-	1	-	1	1	-
Female	12	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	2
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Texas												
All races, both sexes	72	-	-	-	-	-	3	1	1	5	7	7
Male	46	-	-	-	-	-	1	1	1	5	6	6
Female	26	-	-	-	-	-	2	-	-	-	1	1
White, both sexes	61	-	-	-	-	-	3	1	1	4	7	4
Male	38	-	-	-	-	-	1	1	1	4	6	4
Female	23	-	-	-	-	-	2	-	-	-	1	-
Black, both sexes	11	-	-	-	-	-	-	-	-	1	-	3
Male	8	-	-	-	-	-	-	-	-	1	-	2
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	6	-	-	-	-	-	-	-	-	1	-	1
Male	4	-	-	-	-	-	-	-	-	1	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	1	-	1
Male	4	-	-	-	-	-	-	-	-	1	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Vermont												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	32	-	-	-	-	-	-	-	1	1	2	3
Male	22	-	-	-	-	-	-	-	1	-	2	3
Female	10	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	21	-	-	-	-	-	-	-	1	1	1	2
Male	14	-	-	-	-	-	-	-	1	-	1	2
Female	7	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	10	-	-	-	-	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	1	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Washington												
All races, both sexes	26	-	-	-	-	-	-	-	-	1	-	3
Male	15	-	-	-	-	-	-	-	-	1	-	2
Female	11	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	24	-	-	-	-	-	-	-	-	1	-	2
Male	13	-	-	-	-	-	-	-	-	1	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	7	-	-	-	-	-	-	1	-	1	-	-
Male	6	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	5	-	-	-	-	-	-	1	-	-	-	-
Male	5	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Wisconsin												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	2
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
United States												
All races, both sexes	116	105	102	117	119	151	157	90	31	12	1	1
Male	78	67	65	63	64	89	83	40	18	3	-	1
Female	38	38	37	54	55	62	74	50	13	9	1	-
White, both sexes	79	71	80	89	92	129	148	80	28	11	1	1
Male	51	42	52	45	51	77	77	35	17	3	-	1
Female	28	29	28	44	41	52	71	45	11	8	1	-
Black, both sexes	32	29	21	27	23	18	6	8	2	1	-	-
Male	23	22	13	18	11	9	4	4	1	-	-	-
Female	9	7	8	9	12	9	2	4	1	1	-	-
American Indian 1/, both sexes	2	3	-	-	1	1	1	-	-	-	-	-
Male	2	2	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	2	1	1	3	3	2	2	1	-	-	-
Male	2	1	-	-	2	2	2	1	-	-	-	-
Female	1	1	1	1	1	1	-	1	1	-	-	-
Alabama												
All races, both sexes	1	4	-	4	5	-	1	1	-	-	-	-
Male	1	3	-	1	2	-	1	1	-	-	-	-
Female	-	1	-	3	3	-	-	-	-	-	-	-
White, both sexes	1	4	-	3	3	-	1	-	-	-	-	-
Male	1	3	-	1	1	-	1	-	-	-	-	-
Female	-	1	-	2	2	-	-	-	-	-	-	-
Black, both sexes	-	-	-	1	2	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Alaska												
All races, both sexes	-	1	1	-	1	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	4	6	3	6	5	3	2	1	-	-	-	-
Male	2	4	2	4	3	2	1	-	-	-	-	-
Female	2	2	1	2	2	1	1	1	-	-	-	-
White, both sexes	4	5	3	6	4	3	2	1	-	-	-	-
Male	2	3	2	4	3	2	1	-	-	-	-	-
Female	2	2	1	2	1	1	1	1	-	-	-	-
Black, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Arkansas												
All races, both sexes	1	2	1	3	-	4	1	-	-	-	-	-
Male	-	1	1	1	-	2	1	-	-	-	-	-
Female	1	1	-	2	-	2	-	-	-	-	-	-
White, both sexes	-	2	-	3	-	4	1	-	-	-	-	-
Male	-	1	-	1	-	2	1	-	-	-	-	-
Female	-	1	-	2	-	2	-	-	-	-	-	-
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	12	12	13	11	13	17	22	10	2	-	1	-
Male	9	8	10	6	9	12	11	5	1	-	-	-
Female	3	4	3	5	4	5	11	5	1	-	1	-
White, both sexes	9	8	12	8	10	14	20	7	2	-	1	-
Male	7	5	10	5	8	10	11	4	1	-	-	-
Female	2	3	2	3	2	4	9	3	1	-	1	-
Black, both sexes	2	3	1	2	1	1	1	1	-	-	-	-
Male	2	2	-	1	-	-	-	-	-	-	-	-
Female	-	1	1	1	1	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	2	2	-	2	-	-	-	-
Male	-	-	-	-	1	2	-	1	-	-	-	-
Female	1	-	-	1	1	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Colorado												
All races, both sexes	2	-	1	-	-	3	-	2	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	1	-	-	2	-	2	-	-	-	-
White, both sexes	1	-	1	-	-	2	-	2	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	1	-	-	1	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	-	2	3	2	2	3	1	1	-	-	-
Male	1	-	-	1	1	2	3	-	-	-	-	-
Female	-	-	2	2	1	-	-	1	1	-	-	-
White, both sexes	1	-	1	2	2	1	3	-	1	-	-	-
Male	1	-	-	-	1	1	3	-	-	-	-	-
Female	-	-	1	2	1	-	-	-	1	-	-	-
Black, both sexes	-	-	1	1	-	1	-	1	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Delaware												
All races, both sexes	-	1	-	1	-	1	1	-	-	-	-	1
Male	-	1	-	1	-	-	1	-	-	-	-	1
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	1	-	-	-	-	1
Male	-	1	-	-	-	-	1	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Florida												
All races, both sexes	10	2	5	2	8	10	13	5	4	-	-	-
Male	7	1	3	-	4	7	6	2	2	-	-	-
Female	3	1	2	2	4	3	7	3	2	-	-	-
White, both sexes	7	1	5	2	7	8	13	4	4	-	-	-
Male	4	-	3	-	4	5	6	2	2	-	-	-
Female	3	1	2	2	3	3	7	2	2	-	-	-
Black, both sexes	3	1	-	-	1	2	-	1	-	-	-	-
Male	3	1	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	2	3	2	2	3	4	3	3	-	-	-	-
Male	2	3	1	1	1	3	2	1	-	-	-	-
Female	-	-	1	1	2	1	1	2	-	-	-	-
White, both sexes	1	2	1	-	-	4	2	3	-	-	-	-
Male	1	2	-	-	-	3	1	1	-	-	-	-
Female	-	-	1	-	-	1	1	2	-	-	-	-
Black, both sexes	1	1	1	2	3	-	1	-	-	-	-	-
Male	1	1	1	1	1	-	1	-	-	-	-	-
Female	-	-	-	1	2	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Hawaii												
All races, both sexes	1	1	-	1	1	-	1	1	1	-	-	-
Male	1	1	-	-	1	-	1	1	-	-	-	-
Female	-	-	-	1	-	-	-	-	1	-	-	-
White, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	-	1	-	1	-	-	-
Male	-	1	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Idaho												
All races, both sexes	-	1	-	1	-	-	2	1	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	-	-	1	-	-	2	1	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Illinois												
All races, both sexes	3	4	4	10	7	8	2	3	1	1	-	-
Male	3	3	2	7	3	5	2	2	-	-	-	-
Female	-	1	2	3	4	3	-	1	1	1	-	-
White, both sexes	2	1	3	6	3	7	2	2	1	1	-	-
Male	2	1	2	3	1	5	2	1	-	-	-	-
Female	-	-	1	3	2	2	-	1	1	1	-	-
Black, both sexes	1	3	1	4	4	1	-	1	-	-	-	-
Male	1	2	-	4	2	-	-	1	-	-	-	-
Female	-	1	1	-	2	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1	6	1	-	3	1	4	4	2	-	-	-
Male	-	4	1	-	2	1	1	1	1	-	-	-
Female	1	2	-	-	1	-	3	3	1	-	-	-
White, both sexes	1	4	-	-	3	1	4	4	2	-	-	-
Male	-	2	-	-	2	1	1	1	1	-	-	-
Female	1	2	-	-	1	-	3	3	1	-	-	-
Black, both sexes	-	2	1	-	-	-	-	-	-	-	-	-
Male	-	2	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Iowa												
All races, both sexes	3	1	2	3	1	1	2	1	1	-	-	-
Male	2	1	1	1	-	-	-	-	1	-	-	-
Female	1	-	1	2	1	1	2	1	-	-	-	-
White, both sexes	3	1	2	3	-	1	2	1	1	-	-	-
Male	2	1	1	1	-	-	-	-	1	-	-	-
Female	1	-	1	2	-	1	2	1	-	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	1	2	1	1	3	2	1	-	-	-	-
Male	-	-	2	1	-	-	1	-	-	-	-	-
Female	-	1	-	-	1	3	1	1	-	-	-	-
White, both sexes	-	1	2	1	1	2	2	1	-	-	-	-
Male	-	-	2	1	-	-	1	-	-	-	-	-
Female	-	1	-	-	1	2	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Kentucky												
All races, both sexes	1	1	1	3	3	1	-	-	-	-	-	-
Male	1	1	1	2	1	1	-	-	-	-	-	-
Female	-	-	-	1	2	-	-	-	-	-	-	-
White, both sexes	1	1	1	3	2	1	-	-	-	-	-	-
Male	1	1	1	2	-	1	-	-	-	-	-	-
Female	-	-	-	1	2	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	2	2	3	4	3	1	-	-	1	-	-
Male	-	2	1	3	2	2	-	-	-	-	-	-
Female	-	-	1	-	2	1	1	-	-	1	-	-
White, both sexes	-	1	2	2	2	2	1	-	-	1	-	-
Male	-	1	1	2	-	2	-	-	-	-	-	-
Female	-	-	1	-	2	-	1	-	-	1	-	-
Black, both sexes	-	1	-	1	2	1	-	-	-	-	-	-
Male	-	1	-	1	2	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Maine												
All races, both sexes	-	-	1	-	2	1	1	1	-	-	-	-
Male	-	-	1	-	2	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	1	-	2	1	1	1	-	-	-	-
Male	-	-	1	-	2	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	3	2	2	2	1	4	2	2	-	-	-	-
Male	2	1	-	1	1	3	-	-	-	-	-	-
Female	1	1	2	1	-	1	2	2	-	-	-	-
White, both sexes	2	-	-	1	-	3	2	2	-	-	-	-
Male	1	-	-	-	-	2	-	-	-	-	-	-
Female	1	-	-	1	-	1	2	2	-	-	-	-
Black, both sexes	1	2	2	1	1	1	-	-	-	-	-	-
Male	1	1	-	1	1	1	-	-	-	-	-	-
Female	-	1	2	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Massachusetts												
All races, both sexes	1	3	1	1	-	3	3	3	1	1	-	-
Male	1	3	-	1	-	1	1	2	-	1	-	-
Female	-	-	1	-	-	2	2	1	1	-	-	-
White, both sexes	1	3	1	-	-	3	3	3	1	1	-	-
Male	1	3	-	-	-	1	1	2	-	1	-	-
Female	-	-	1	-	-	2	2	1	1	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	10	3	2	4	6	5	7	1	-	-	-	-
Male	5	1	1	2	3	4	4	1	-	-	-	-
Female	5	2	1	2	3	1	3	-	-	-	-	-
White, both sexes	5	1	2	4	5	5	6	1	-	-	-	-
Male	3	-	1	2	2	4	3	1	-	-	-	-
Female	2	1	1	2	3	1	3	-	-	-	-	-
Black, both sexes	5	2	-	-	1	-	1	-	-	-	-	-
Male	2	1	-	-	1	-	1	-	-	-	-	-
Female	3	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Minnesota												
All races, both sexes	2	-	1	-	3	3	6	-	3	-	-	-
Male	2	-	1	-	-	-	4	-	3	-	-	-
Female	-	-	-	-	3	3	2	-	-	-	-	-
White, both sexes	1	-	1	-	2	3	6	-	3	-	-	-
Male	1	-	1	-	-	-	4	-	3	-	-	-
Female	-	-	-	-	2	3	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	2	-	1	2	1	3	-	-	-	-	-	-
Male	1	-	-	2	-	2	-	-	-	-	-	-
Female	1	-	1	-	1	1	-	-	-	-	-	-
White, both sexes	1	-	1	1	1	2	-	-	-	-	-	-
Male	1	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	1	-	1	1	-	-	-	-	-	-
Black, both sexes	1	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Missouri												
All races, both sexes	2	3	2	2	2	2	4	-	2	-	-	-
Male	2	2	1	1	2	1	2	-	1	-	-	-
Female	-	1	1	1	-	1	2	-	1	-	-	-
White, both sexes	2	1	2	2	1	2	4	-	2	-	-	-
Male	2	-	1	1	1	1	2	-	1	-	-	-
Female	-	1	1	1	-	1	2	-	1	-	-	-
Black, both sexes	-	2	-	-	1	-	-	-	-	-	-	-
Male	-	2	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Nebraska												
All races, both sexes	1	1	-	-	-	1	2	1	-	-	-	-
Male	1	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
White, both sexes	1	1	-	-	-	1	2	1	-	-	-	-
Male	1	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	1	1	3	-	-	1	-	-	-	-	-
Male	-	1	1	3	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	1	1	2	-	-	1	-	-	-	-	-
Male	-	1	1	2	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
New Hampshire												
All races, both sexes	2	-	-	1	1	1	1	1	-	-	-	-
Male	2	-	-	1	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	2	-	-	1	1	1	1	1	-	-	-	-
Male	2	-	-	1	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	2	2	8	4	8	8	1	1	-	-	-
Male	-	1	-	4	2	3	6	1	1	-	-	-
Female	-	1	2	4	2	5	2	-	-	-	-	-
White, both sexes	-	2	1	5	4	6	7	1	1	-	-	-
Male	-	1	-	3	2	2	5	1	1	-	-	-
Female	-	1	1	2	2	4	2	-	-	-	-	-
Black, both sexes	-	-	-	3	-	2	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	2	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
New Mexico												
All races, both sexes	2	-	-	-	1	2	-	-	-	1	-	-
Male	1	-	-	-	1	1	-	-	-	1	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	1	-	-	-	1	1	-	-	-	1	-	-
Male	-	-	-	-	1	1	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
New York												
All races, both sexes	10	6	8	4	4	15	14	12	5	2	-	-
Male	9	4	6	2	1	11	6	6	3	-	-	-
Female	1	2	2	2	3	4	8	6	2	2	-	-
White, both sexes	4	5	4	2	4	14	13	12	4	2	-	-
Male	3	3	3	1	1	11	5	6	3	-	-	-
Female	1	2	1	1	3	3	8	6	1	2	-	-
Black, both sexes	5	1	4	2	-	1	1	-	1	-	-	-
Male	5	1	3	1	-	-	1	-	-	-	-	-
Female	-	-	1	1	-	1	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
North Carolina												
All races, both sexes	1	4	4	5	3	3	4	-	-	2	-	-
Male	1	2	4	1	1	2	3	-	-	-	-	-
Female	-	2	-	4	2	1	1	-	-	2	-	-
White, both sexes	1	2	1	3	2	2	4	-	-	2	-	-
Male	1	-	1	1	1	1	3	-	-	-	-	-
Female	-	2	-	2	1	1	1	-	-	2	-	-
Black, both sexes	-	2	3	2	1	1	-	-	-	-	-	-
Male	-	2	3	-	-	1	-	-	-	-	-	-
Female	-	-	-	2	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	1	-	2	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	2	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Ohio												
All races, both sexes	5	4	8	7	6	8	9	2	-	-	-	-
Male	2	-	5	3	5	2	5	1	-	-	-	-
Female	3	4	3	4	1	6	4	1	-	-	-	-
White, both sexes	3	3	4	7	4	6	9	2	-	-	-	-
Male	1	-	3	3	4	1	5	1	-	-	-	-
Female	2	3	1	4	-	5	4	1	-	-	-	-
Black, both sexes	2	1	4	-	2	2	-	-	-	-	-	-
Male	1	-	2	-	1	1	-	-	-	-	-	-
Female	1	1	2	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1	1	1	-	2	3	2	-	-	-	-	-
Male	-	1	1	-	1	2	1	-	-	-	-	-
Female	1	-	-	-	1	1	1	-	-	-	-	-
White, both sexes	-	1	-	-	2	2	2	-	-	-	-	-
Male	-	1	-	-	1	1	1	-	-	-	-	-
Female	-	-	-	-	1	1	1	-	-	-	-	-
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Oregon												
All races, both sexes	1	2	3	1	3	1	1	1	-	-	-	-
Male	1	1	2	-	2	1	1	-	-	-	-	-
Female	-	1	1	1	1	-	-	1	-	-	-	-
White, both sexes	1	2	3	1	3	1	1	1	-	-	-	-
Male	1	1	2	-	2	1	1	-	-	-	-	-
Female	-	1	1	1	1	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	5	1	8	5	6	8	12	6	-	-	-	-
Male	4	1	5	2	5	3	6	3	-	-	-	-
Female	1	-	3	3	1	5	6	3	-	-	-	-
White, both sexes	3	1	8	4	6	8	12	6	-	-	-	-
Male	2	1	5	1	5	3	6	3	-	-	-	-
Female	1	-	3	3	1	5	6	3	-	-	-	-
Black, both sexes	2	-	-	1	-	-	-	-	-	-	-	-
Male	2	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	2	2	4	2	1	-	2	3	-	-	-	-
Male	-	2	2	1	-	-	-	2	-	-	-	-
Female	2	-	2	1	1	-	2	1	-	-	-	-
White, both sexes	-	1	4	1	-	-	2	2	-	-	-	-
Male	-	1	2	-	-	-	-	1	-	-	-	-
Female	-	-	2	1	-	-	2	1	-	-	-	-
Black, both sexes	2	1	-	1	1	-	-	1	-	-	-	-
Male	-	1	-	1	-	-	-	1	-	-	-	-
Female	2	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
South Dakota												
All races, both sexes	-	-	-	-	1	-	-	2	1	-	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	2	1	-	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	4	1	2	1	3	5	2	4	1	1	-	-
Male	2	-	2	1	2	4	1	1	-	-	-	-
Female	2	1	-	-	1	1	1	3	1	1	-	-
White, both sexes	4	1	2	1	3	5	1	4	1	-	-	-
Male	2	-	2	1	2	4	1	1	-	-	-	-
Female	2	1	-	-	1	1	-	3	1	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Texas												
All races, both sexes	10	6	4	3	6	6	6	6	1	-	-	-
Male	5	3	3	1	4	4	4	1	1	-	-	-
Female	5	3	1	2	2	2	2	5	-	-	-	-
White, both sexes	8	5	3	3	6	6	5	5	-	-	-	-
Male	3	3	2	1	4	4	3	1	-	-	-	-
Female	5	2	1	2	2	2	2	4	-	-	-	-
Black, both sexes	2	1	1	-	-	-	1	1	1	-	-	-
Male	2	-	1	-	-	-	1	-	1	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	2	-	-	-	2	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	2	-	-	-	2	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Vermont												
All races, both sexes	1	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	1	-	-	-	-	-	-
White, both sexes	1	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	5	5	1	3	-	1	4	5	-	1	-	-
Male	4	2	-	2	-	1	2	4	-	1	-	-
Female	1	3	1	1	-	-	2	1	-	-	-	-
White, both sexes	3	1	1	1	-	-	4	5	-	1	-	-
Male	2	-	-	1	-	-	2	4	-	1	-	-
Female	1	1	1	-	-	-	2	1	-	-	-	-
Black, both sexes	2	3	-	2	-	1	-	-	-	-	-	-
Male	2	2	-	1	-	1	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Washington												
All races, both sexes	1	5	1	2	2	3	4	2	2	-	-	-
Male	1	3	1	1	1	1	2	1	1	-	-	-
Female	-	2	-	1	1	2	2	1	1	-	-	-
White, both sexes	-	5	1	2	2	3	4	2	2	-	-	-
Male	-	3	1	1	1	1	2	1	1	-	-	-
Female	-	2	-	1	1	2	2	1	1	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	2	-	-	1	1	-	-	1	-	-	-
Male	-	2	-	-	1	1	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	1	1	-	-	1	-	-	-
Male	-	1	-	-	1	1	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Wisconsin												
All races, both sexes	2	2	3	3	1	2	2	1	-	2	-	-
Male	2	2	2	2	-	2	2	1	-	-	-	-
Female	-	-	1	1	1	-	-	-	-	2	-	-
White, both sexes	2	1	3	3	1	2	2	1	-	2	-	-
Male	2	1	2	2	-	2	2	1	-	-	-	-
Female	-	-	1	1	1	-	-	-	-	2	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	-	2	-	-	-	-	-	-	-	-	-
Male	-	-	2	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	2	-	-	-	-	-	-	-	-	-
Male	-	-	2	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
United States												
All races, both sexes	867	25	19	15	9	17	31	25	28	47	40	58
Male	454	16	13	10	7	10	16	14	18	31	26	31
Female	413	9	6	5	2	7	15	11	10	16	14	27
White, both sexes	698	13	14	10	8	14	23	18	23	32	32	45
Male	366	8	10	6	6	9	10	13	15	21	22	23
Female	332	5	4	4	2	5	13	5	8	11	10	22
Black, both sexes	145	11	3	3	1	3	7	7	4	13	6	12
Male	74	7	2	3	1	1	6	1	2	9	3	7
Female	71	4	1	-	-	2	1	6	2	4	3	5
American Indian 1/, both sexes	4	-	1	-	-	-	1	-	-	-	1	-
Male	3	-	1	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	20	1	1	2	-	-	-	-	1	2	1	1
Male	11	1	-	1	-	-	-	-	1	1	-	1
Female	9	-	1	1	-	-	-	-	-	1	1	-
Alabama												
All races, both sexes	17	-	-	-	-	1	-	1	1	-	-	1
Male	9	-	-	-	-	1	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	1	1	-	-	-
White, both sexes	13	-	-	-	-	1	-	1	-	-	-	1
Male	7	-	-	-	-	1	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Alaska												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	15	1	1	1	-	-	-	-	1	2	-	-
Male	8	-	1	1	-	-	-	-	-	2	-	-
Female	7	1	-	-	-	-	-	-	1	-	-	-
White, both sexes	14	1	1	1	-	-	-	-	1	2	-	-
Male	7	-	1	1	-	-	-	-	-	2	-	-
Female	7	1	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Arkansas												
All races, both sexes	9	1	-	-	-	-	1	1	-	-	-	-
Male	5	1	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	6	-	-	-	-	-	1	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	3	1	-	-	-	-	-	1	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	73	3	1	3	1	-	3	2	4	3	7	8
Male	39	1	1	2	1	-	1	2	2	1	6	4
Female	34	2	-	1	-	-	2	-	2	2	1	4
White, both sexes	62	3	1	3	1	-	3	2	4	2	5	7
Male	33	1	1	2	1	-	1	2	2	-	4	3
Female	29	2	-	1	-	-	2	-	2	2	1	4
Black, both sexes	5	-	-	-	-	-	-	-	-	1	1	-
Male	2	-	-	-	-	-	-	-	-	1	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Colorado												
All races, both sexes	13	-	1	-	-	-	1	-	-	1	2	1
Male	7	-	1	-	-	-	1	-	-	1	2	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	1	-	-	-	-	-	-	1	2	1
Male	6	-	1	-	-	-	-	-	-	1	2	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	11	-	-	-	-	-	-	-	1	1	-	2
Male	8	-	-	-	-	-	-	-	-	1	-	2
Female	3	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	1	1	-	1
Male	2	-	-	-	-	-	-	-	-	1	-	1
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Delaware												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Florida												
All races, both sexes	60	3	1	-	1	-	2	1	2	4	1	5
Male	36	2	1	-	1	-	2	-	1	3	-	3
Female	24	1	-	-	-	-	-	1	1	1	1	2
White, both sexes	47	1	1	-	-	-	2	-	2	3	1	3
Male	28	1	1	-	-	-	2	-	1	2	-	1
Female	19	-	-	-	-	-	-	-	1	1	1	2
Black, both sexes	13	2	-	-	1	-	-	1	-	1	-	2
Male	8	1	-	-	1	-	-	-	-	1	-	2
Female	5	1	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	31	2	1	-	-	1	1	1	1	3	1	2
Male	19	1	-	-	-	1	1	1	1	3	-	2
Female	12	1	1	-	-	-	-	-	-	-	1	-
White, both sexes	19	1	-	-	-	1	-	-	-	2	1	1
Male	11	1	-	-	-	1	-	-	-	2	-	1
Female	8	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	10	1	-	-	-	-	1	1	1	1	-	1
Male	8	-	-	-	-	-	1	1	1	1	-	1
Female	2	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Hawaii												
All races, both sexes	5	-	-	1	-	-	1	-	-	1	-	-
Male	2	-	-	1	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	1	-	-	1	-	-
White, both sexes	3	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	1	-	-	-	-	-	1	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Idaho												
All races, both sexes	5	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Illinois												
All races, both sexes	51	1	-	1	-	-	1	1	1	2	1	2
Male	26	-	-	-	-	-	1	-	1	2	-	-
Female	25	1	-	1	-	-	-	1	-	-	1	2
White, both sexes	39	-	-	1	-	-	-	1	1	2	-	1
Male	21	-	-	-	-	-	-	-	1	2	-	-
Female	18	-	-	1	-	-	-	1	-	-	-	1
Black, both sexes	11	1	-	-	-	-	1	-	-	-	1	1
Male	4	-	-	-	-	-	1	-	-	-	-	-
Female	7	1	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	24	1	-	-	1	1	1	-	1	-	-	-
Male	12	1	-	-	-	-	1	-	1	-	-	-
Female	12	-	-	-	1	1	-	-	-	-	-	-
White, both sexes	22	1	-	-	1	1	1	-	1	-	-	-
Male	11	1	-	-	-	-	1	-	1	-	-	-
Female	11	-	-	-	1	1	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Iowa												
All races, both sexes	15	-	-	-	-	-	-	2	2	1	1	1
Male	6	-	-	-	-	-	-	-	2	1	1	-
Female	9	-	-	-	-	-	-	2	-	-	-	1
White, both sexes	14	-	-	-	-	-	-	1	2	1	1	1
Male	6	-	-	-	-	-	-	-	2	1	1	-
Female	8	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	10	1	1	-	-	-	1	-	-	-	-	1
Male	5	1	-	-	-	-	-	-	-	-	-	-
Female	5	-	1	-	-	-	1	-	-	-	-	1
White, both sexes	9	-	1	-	-	-	1	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	1	-	-	-	1	-	-	-	-	1
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Kentucky												
All races, both sexes	10	-	-	-	-	-	1	-	1	2	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	1	-	1	2	1	-
White, both sexes	7	-	-	-	-	-	-	-	1	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	1	-	1	-
Black, both sexes	3	-	-	-	-	-	1	-	-	2	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	1	-	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Maine												
All races, both sexes	6	-	-	-	-	1	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	1	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	18	-	-	-	-	-	-	-	1	2	-	1
Male	8	-	-	-	-	-	-	-	1	2	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	1	1	-	1
Male	7	-	-	-	-	-	-	-	1	1	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Massachusetts												
All races, both sexes	24	1	-	-	-	3	2	1	-	-	5	1
Male	14	-	-	-	-	2	1	1	-	-	2	-
Female	10	1	-	-	-	1	1	-	-	-	3	1
White, both sexes	21	1	-	-	-	2	2	1	-	-	4	1
Male	11	-	-	-	-	1	1	1	-	-	1	-
Female	10	1	-	-	-	1	1	-	-	-	3	1
Black, both sexes	3	-	-	-	-	1	-	-	-	-	1	-
Male	3	-	-	-	-	1	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	16	-	-	-	1	2	-	-	-	-	-	1
Male	10	-	-	-	1	1	-	-	-	-	-	-
Female	6	-	-	-	-	1	-	-	-	-	-	1
White, both sexes	16	-	-	-	1	2	-	-	-	-	-	1
Male	10	-	-	-	1	1	-	-	-	-	-	-
Female	6	-	-	-	-	1	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Minnesota												
All races, both sexes	13	1	-	-	-	-	-	-	-	1	1	3
Male	7	1	-	-	-	-	-	-	-	-	1	3
Female	6	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	1	1	3
Male	6	-	-	-	-	-	-	-	-	-	1	3
Female	6	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	8	1	-	-	-	-	-	1	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	5	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Missouri												
All races, both sexes	25	2	2	-	-	1	-	-	2	1	1	-
Male	12	1	1	-	-	1	-	-	1	1	1	-
Female	13	1	1	-	-	-	-	-	1	-	-	-
White, both sexes	19	1	2	-	-	1	-	-	2	-	1	-
Male	9	1	1	-	-	1	-	-	1	-	1	-
Female	10	-	1	-	-	-	-	-	1	-	-	-
Black, both sexes	6	1	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	3	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	3	-	1	-	-	-	-	-	-	-	-	1
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Nebraska												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	8	-	-	-	1	-	-	1	-	1	-	-
Male	3	-	-	-	1	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	1	-	1	-	-
White, both sexes	3	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	1	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
New Hampshire												
All races, both sexes	3	-	-	-	-	1	-	-	-	-	-	1
Male	2	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	3	-	-	-	-	1	-	-	-	-	-	1
Male	2	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	19	-	-	1	-	-	-	-	1	-	2	-
Male	6	-	-	-	-	-	-	-	1	-	1	-
Female	13	-	-	1	-	-	-	-	-	-	1	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	1	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
New Mexico												
All races, both sexes	6	-	-	-	-	1	-	-	-	-	-	-
Male	4	-	-	-	-	1	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	1	-	-	-	-	-	-
Male	4	-	-	-	-	1	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	71	-	-	1	1	-	2	2	3	6	5	5
Male	39	-	-	-	-	-	2	1	3	6	2	3
Female	32	-	-	1	1	-	-	1	-	-	3	2
White, both sexes	55	-	-	1	1	-	2	1	3	4	4	3
Male	30	-	-	-	-	-	2	1	3	4	2	2
Female	25	-	-	1	1	-	-	-	-	-	2	1
Black, both sexes	15	-	-	-	-	-	-	1	-	1	1	2
Male	8	-	-	-	-	-	-	-	-	1	-	1
Female	7	-	-	-	-	-	-	1	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
North Carolina												
All races, both sexes	29	-	1	1	-	2	2	1	-	1	-	3
Male	15	-	1	1	-	-	1	-	-	1	-	2
Female	14	-	-	-	-	2	1	1	-	-	-	1
White, both sexes	22	-	1	-	-	1	1	1	-	1	-	3
Male	11	-	1	-	-	-	-	-	-	1	-	2
Female	11	-	-	-	-	1	1	1	-	-	-	1
Black, both sexes	7	-	-	1	-	1	1	-	-	-	-	-
Male	4	-	-	1	-	-	1	-	-	-	-	-
Female	3	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	4	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Ohio												
All races, both sexes	47	2	3	1	-	1	1	2	-	-	1	7
Male	22	2	3	1	-	1	-	2	-	-	1	2
Female	25	-	-	-	-	-	1	-	-	-	-	5
White, both sexes	40	-	2	1	-	1	1	2	-	-	1	5
Male	18	-	2	1	-	1	-	2	-	-	1	2
Female	22	-	-	-	-	-	1	-	-	-	-	3
Black, both sexes	7	2	1	-	-	-	-	-	-	-	-	2
Male	4	2	1	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Oregon												
All races, both sexes	10	1	-	-	1	1	-	1	-	-	-	-
Male	7	1	-	-	1	1	-	1	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	1	-	-	1	1	-	1	-	-	-	-
Male	7	1	-	-	1	1	-	1	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	40	-	-	-	-	-	2	-	1	4	2	1
Male	16	-	-	-	-	-	1	-	-	1	1	-
Female	24	-	-	-	-	-	1	-	1	3	1	1
White, both sexes	33	-	-	-	-	-	2	-	1	3	2	1
Male	15	-	-	-	-	-	1	-	-	1	1	-
Female	18	-	-	-	-	-	1	-	1	2	1	1
Black, both sexes	7	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Rhode Island												
All races, both sexes	5	-	1	-	-	1	-	1	-	-	1	-
Male	3	-	1	-	-	-	-	1	-	-	1	-
Female	2	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	3	-	1	-	-	-	-	1	-	-	1	-
Male	3	-	1	-	-	-	-	1	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	19	-	-	-	-	-	1	-	-	4	1	1
Male	10	-	-	-	-	-	1	-	-	3	1	1
Female	9	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	2	1	1
Male	7	-	-	-	-	-	-	-	-	1	1	1
Female	6	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	6	-	-	-	-	-	1	-	-	2	-	-
Male	3	-	-	-	-	-	1	-	-	2	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
South Dakota												
All races, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	12	1	-	1	-	-	1	-	2	-	-	2
Male	7	1	-	-	-	-	1	-	2	-	-	1
Female	5	-	-	1	-	-	-	-	-	-	-	1
White, both sexes	9	-	-	1	-	-	1	-	1	-	-	1
Male	4	-	-	-	-	-	1	-	1	-	-	-
Female	5	-	-	1	-	-	-	-	-	-	-	1
Black, both sexes	3	1	-	-	-	-	-	-	1	-	-	1
Male	3	1	-	-	-	-	-	-	1	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Texas												
All races, both sexes	41	2	3	1	-	-	4	2	1	2	2	2
Male	20	1	1	1	-	-	1	1	1	-	2	1
Female	21	1	2	-	-	-	3	1	-	2	-	1
White, both sexes	36	2	2	-	-	-	4	2	1	2	2	1
Male	17	1	1	-	-	-	1	1	1	-	2	1
Female	19	1	1	-	-	-	3	1	-	2	-	-
Black, both sexes	5	-	1	1	-	-	-	-	-	-	-	1
Male	3	-	-	1	-	-	-	-	-	-	-	-
Female	2	-	1	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	6	-	-	-	1	-	-	1	1	3	-	-
Male	5	-	-	-	1	-	-	1	1	2	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	6	-	-	-	1	-	-	1	1	3	-	-
Male	5	-	-	-	1	-	-	1	1	2	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Vermont												
All races, both sexes	4	-	-	-	-	-	1	1	-	-	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	2	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	1	1	-	-	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	2	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	17	-	1	-	-	-	1	-	-	-	3	1
Male	12	-	1	-	-	-	1	-	-	-	2	1
Female	5	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	1	-	-	-	1	-	-	-	1	1
Male	4	-	1	-	-	-	1	-	-	-	1	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Washington												
All races, both sexes	14	-	-	1	-	-	-	-	-	-	-	2
Male	10	-	-	1	-	-	-	-	-	-	-	2
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	1	-	-	-	-	-	-	-	2
Male	8	-	-	1	-	-	-	-	-	-	-	2
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	6	-	-	1	1	-	-	1	-	1	1	-
Male	5	-	-	1	1	-	-	1	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	6	-	-	1	1	-	-	1	-	1	1	-
Male	5	-	-	1	1	-	-	1	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Wisconsin												
All races, both sexes	19	1	-	-	-	-	-	1	-	1	-	1
Male	11	1	-	-	-	-	-	1	-	1	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	17	-	-	-	-	-	-	1	-	1	-	1
Male	10	-	-	-	-	-	-	1	-	1	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
United States												
All races, both sexes	58	65	58	63	56	73	64	69	30	14	3	-
Male	26	36	35	33	33	40	19	23	16	1	-	-
Female	32	29	23	30	23	33	45	46	14	13	3	-
White, both sexes	43	53	42	49	50	65	62	60	27	12	3	-
Male	20	31	26	25	31	36	18	21	14	1	-	-
Female	23	22	16	24	19	29	44	39	13	11	3	-
Black, both sexes	14	12	14	12	5	7	1	6	2	2	-	-
Male	5	5	8	6	2	3	1	1	1	-	-	-
Female	9	7	6	6	3	4	-	5	1	2	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	2	1	1	1	3	1	-	-	-
Male	-	-	1	2	-	1	-	1	1	-	-	-
Female	-	-	1	-	1	-	1	2	-	-	-	-
Alabama												
All races, both sexes	2	2	1	2	2	1	1	1	1	-	-	-
Male	1	1	1	1	-	1	-	1	1	-	-	-
Female	1	1	-	1	2	-	1	-	-	-	-	-
White, both sexes	1	2	-	1	2	1	1	1	1	-	-	-
Male	-	1	-	1	-	1	-	1	1	-	-	-
Female	1	1	-	-	2	-	1	-	-	-	-	-
Black, both sexes	1	-	1	1	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Alaska												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2	-	-	3	1	1	-	2	-	-	-	-
Male	1	-	-	2	-	1	-	-	-	-	-	-
Female	1	-	-	1	1	-	-	2	-	-	-	-
White, both sexes	1	-	-	3	1	1	-	2	-	-	-	-
Male	-	-	-	2	-	1	-	-	-	-	-	-
Female	1	-	-	1	1	-	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Arkansas												
All races, both sexes	1	1	-	-	1	1	1	1	-	-	-	-
Male	1	1	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
White, both sexes	-	1	-	-	1	1	1	1	-	-	-	-
Male	-	1	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	4	4	5	5	4	3	5	7	-	1	-	-
Male	1	2	3	2	3	3	2	2	-	-	-	-
Female	3	2	2	3	1	-	3	5	-	1	-	-
White, both sexes	3	4	3	4	3	2	5	6	-	1	-	-
Male	1	2	3	1	3	2	2	2	-	-	-	-
Female	2	2	-	3	-	-	3	4	-	1	-	-
Black, both sexes	1	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	1	1	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Colorado												
All races, both sexes	-	2	1	-	1	-	1	1	1	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	1	-	1	1	1	-	-	-
White, both sexes	-	2	1	-	1	-	1	1	1	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	1	-	1	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	1	-	2	-	1	2	1	-	-
Male	-	-	-	1	-	2	-	-	2	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-
White, both sexes	-	-	-	1	-	2	-	1	2	1	-	-
Male	-	-	-	1	-	2	-	-	2	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Florida												
All races, both sexes	2	8	3	5	4	12	3	3	-	-	-	-
Male	1	8	2	3	2	6	1	-	-	-	-	-
Female	1	-	1	2	2	6	2	3	-	-	-	-
White, both sexes	2	7	3	4	3	10	3	2	-	-	-	-
Male	1	7	2	2	2	5	1	-	-	-	-	-
Female	1	-	1	2	1	5	2	2	-	-	-	-
Black, both sexes	-	1	-	1	1	2	-	1	-	-	-	-
Male	-	1	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	3	1	4	3	1	3	1	1	-	1	-	-
Male	2	-	2	2	1	2	-	-	-	-	-	-
Female	1	1	2	1	-	1	1	1	-	1	-	-
White, both sexes	2	1	2	3	1	2	1	-	-	1	-	-
Male	1	-	1	2	1	1	-	-	-	-	-	-
Female	1	1	1	1	-	1	1	-	-	1	-	-
Black, both sexes	1	-	2	-	-	1	-	-	-	-	-	-
Male	1	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Hawaii												
All races, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	1	2	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	1	-	-	-
White, both sexes	-	-	-	1	2	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Illinois												
All races, both sexes	5	1	2	8	1	5	6	8	5	-	-	-
Male	2	1	-	3	-	4	3	6	3	-	-	-
Female	3	-	2	5	1	1	3	2	2	-	-	-
White, both sexes	4	1	1	7	1	4	6	6	3	-	-	-
Male	2	1	-	3	-	3	3	5	1	-	-	-
Female	2	-	1	4	1	1	3	1	2	-	-	-
Black, both sexes	1	-	1	1	-	1	-	2	1	-	-	-
Male	-	-	-	-	-	1	-	1	1	-	-	-
Female	1	-	1	1	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1	3	4	4	1	2	2	-	1	1	-	-
Male	-	3	3	3	-	-	-	-	-	-	-	-
Female	1	-	1	1	1	2	2	-	1	1	-	-
White, both sexes	1	2	4	4	1	2	2	-	-	1	-	-
Male	-	2	3	3	-	-	-	-	-	-	-	-
Female	1	-	1	1	1	2	2	-	-	1	-	-
Black, both sexes	-	1	-	-	-	-	-	-	1	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Iowa												
All races, both sexes	2	-	1	1	-	-	2	-	1	1	-	-
Male	-	-	1	-	-	-	-	-	1	-	-	-
Female	2	-	-	1	-	-	2	-	-	1	-	-
White, both sexes	2	-	1	1	-	-	2	-	1	1	-	-
Male	-	-	1	-	-	-	-	-	1	-	-	-
Female	2	-	-	1	-	-	2	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	1	1	2	1	-	1	-	-
Male	-	-	-	-	1	1	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-
White, both sexes	-	-	-	-	1	1	2	1	-	1	-	-
Male	-	-	-	-	1	1	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Kentucky												
All races, both sexes	2	-	1	-	1	-	-	1	-	-	-	-
Male	2	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-
White, both sexes	2	-	1	-	1	-	-	1	-	-	-	-
Male	2	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	2	1	-	1	1	-	-	-	-	-	-
Male	-	1	1	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	2	-	-	1	1	-	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Maine												
All races, both sexes	-	-	-	-	3	1	-	-	-	1	-	-
Male	-	-	-	-	1	1	-	-	-	1	-	-
Female	-	-	-	-	2	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	3	1	-	-	-	1	-	-
Male	-	-	-	-	1	1	-	-	-	1	-	-
Female	-	-	-	-	2	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	3	2	-	2	2	-	3	1	1	-	-
Male	-	1	1	-	-	1	-	-	1	-	-	-
Female	-	2	1	-	2	1	-	3	-	1	-	-
White, both sexes	-	3	1	-	1	1	-	1	1	1	-	-
Male	-	1	1	-	-	1	-	-	1	-	-	-
Female	-	2	-	-	1	-	-	1	-	1	-	-
Black, both sexes	-	-	1	-	1	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Massachusetts												
All races, both sexes	-	-	1	1	2	1	3	3	-	-	-	-
Male	-	-	1	-	2	1	2	2	-	-	-	-
Female	-	-	-	1	-	-	1	1	-	-	-	-
White, both sexes	-	-	-	1	2	1	3	3	-	-	-	-
Male	-	-	-	-	2	1	2	2	-	-	-	-
Female	-	-	-	1	-	-	1	1	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	1	4	-	3	1	-	2	1	-	-	-
Male	-	-	2	-	3	1	-	1	1	-	-	-
Female	-	1	2	-	-	-	-	1	-	-	-	-
White, both sexes	-	1	4	-	3	1	-	2	1	-	-	-
Male	-	-	2	-	3	1	-	1	1	-	-	-
Female	-	1	2	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Minnesota												
All races, both sexes	-	-	-	1	1	2	1	-	1	1	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	1	1	1	1	-	-	1	-	-
White, both sexes	-	-	-	1	1	2	1	-	1	1	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	1	1	1	1	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1	1	-	1	1	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	1	1	1	1	-	-	-	-	-
White, both sexes	-	1	-	1	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	1	1	-	-	-	-	-
Black, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Missouri												
All races, both sexes	1	4	-	3	1	1	1	5	-	-	-	-
Male	1	1	-	2	-	1	-	1	-	-	-	-
Female	-	3	-	1	1	-	1	4	-	-	-	-
White, both sexes	1	2	-	1	1	1	1	5	-	-	-	-
Male	1	1	-	-	-	1	-	1	-	-	-	-
Female	-	1	-	1	1	-	1	4	-	-	-	-
Black, both sexes	-	2	-	2	-	-	-	-	-	-	-	-
Male	-	-	-	2	-	-	-	-	-	-	-	-
Female	-	2	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Nebraska												
All races, both sexes	1	-	1	1	-	-	1	2	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	2	-	-	-	-
White, both sexes	1	-	1	1	-	-	1	2	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	2	-	-	1	-	-	2	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	2	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Black, both sexes	2	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
New Hampshire												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1	4	1	-	-	2	3	1	2	1	-	-
Male	-	1	1	-	-	-	1	1	-	-	-	-
Female	1	3	-	-	-	2	2	-	2	1	-	-
White, both sexes	1	2	1	-	-	1	2	1	2	1	-	-
Male	-	1	1	-	-	-	-	1	-	-	-	-
Female	1	1	-	-	-	1	2	-	2	1	-	-
Black, both sexes	-	2	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	2	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
New Mexico												
All races, both sexes	-	1	1	-	1	1	1	-	-	-	-	-
Male	-	1	-	-	1	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
White, both sexes	-	1	1	-	1	1	1	-	-	-	-	-
Male	-	1	-	-	1	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	6	5	10	4	6	2	4	5	3	-	1	-
Male	1	4	5	2	5	1	1	2	1	-	-	-
Female	5	1	5	2	1	1	3	3	2	-	1	-
White, both sexes	5	3	8	1	4	2	4	5	3	-	1	-
Male	1	2	4	1	3	1	1	2	1	-	-	-
Female	4	1	4	-	1	1	3	3	2	-	1	-
Black, both sexes	1	2	2	3	2	-	-	-	-	-	-	-
Male	-	2	1	1	2	-	-	-	-	-	-	-
Female	1	-	1	2	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
North Carolina												
All races, both sexes	3	2	2	1	-	5	2	1	2	-	-	-
Male	2	-	2	-	-	3	1	-	1	-	-	-
Female	1	2	-	1	-	2	1	1	1	-	-	-
White, both sexes	1	2	1	-	-	5	2	1	2	-	-	-
Male	1	-	1	-	-	3	1	-	1	-	-	-
Female	-	2	-	-	-	2	1	1	1	-	-	-
Black, both sexes	2	-	1	1	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	1	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	2	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Ohio												
All races, both sexes	4	6	1	1	3	4	3	4	1	1	1	-
Male	2	3	-	1	2	-	1	-	1	-	-	-
Female	2	3	1	-	1	4	2	4	-	1	1	-
White, both sexes	4	4	1	1	3	4	3	4	1	1	1	-
Male	2	2	-	1	2	-	1	-	1	-	-	-
Female	2	2	1	-	1	4	2	4	-	1	1	-
Black, both sexes	-	2	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	1	-	-	3	1	1	-	-	-	-
Male	-	-	1	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
White, both sexes	-	-	1	-	-	3	1	1	-	-	-	-
Male	-	-	1	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Oregon												
All races, both sexes	1	-	-	2	-	1	2	-	-	-	-	-
Male	1	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	1	1	-	-	-	-	-
White, both sexes	1	-	-	2	-	1	2	-	-	-	-	-
Male	1	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	3	4	4	3	3	1	4	1	4	2	1	-
Male	-	1	3	3	2	1	2	-	1	-	-	-
Female	3	3	1	-	1	-	2	1	3	2	1	-
White, both sexes	1	3	4	2	3	1	4	1	4	-	1	-
Male	-	1	3	2	2	1	2	-	1	-	-	-
Female	1	2	1	-	1	-	2	1	3	-	1	-
Black, both sexes	2	1	-	1	-	-	-	-	-	2	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Rhode Island												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	-	2	2	1	3	1	1	1	-	-	-
Male	-	-	1	1	-	1	-	1	-	-	-	-
Female	1	-	1	1	1	2	1	-	1	-	-	-
White, both sexes	1	-	1	1	1	2	1	1	1	-	-	-
Male	-	-	1	1	-	1	-	1	-	-	-	-
Female	1	-	-	-	1	1	1	-	1	-	-	-
Black, both sexes	-	-	1	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
South Dakota												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	1	1	1	2	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
White, both sexes	-	-	-	-	1	1	1	2	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Texas												
All races, both sexes	4	3	3	1	1	3	5	2	-	-	-	-
Male	2	2	3	1	1	1	-	1	-	-	-	-
Female	2	1	-	-	-	2	5	1	-	-	-	-
White, both sexes	4	3	1	1	1	3	5	2	-	-	-	-
Male	2	2	1	1	1	1	-	1	-	-	-	-
Female	2	1	-	-	-	2	5	1	-	-	-	-
Black, both sexes	-	-	2	-	-	-	-	-	-	-	-	-
Male	-	-	2	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Vermont												
All races, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	2	1	2	1	5	-	-	-	-
Male	-	-	-	1	1	2	-	3	-	-	-	-
Female	-	-	-	1	-	-	1	2	-	-	-	-
White, both sexes	-	-	-	2	1	2	1	3	-	-	-	-
Male	-	-	-	1	1	2	-	2	-	-	-	-
Female	-	-	-	1	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Washington												
All races, both sexes	2	2	1	2	2	1	1	-	-	-	-	-
Male	1	2	1	2	1	-	-	-	-	-	-	-
Female	1	-	-	-	1	1	1	-	-	-	-	-
White, both sexes	2	2	-	1	2	1	-	-	-	-	-	-
Male	1	2	-	1	1	-	-	-	-	-	-	-
Female	1	-	-	-	1	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	-	1	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Wisconsin												
All races, both sexes	2	2	1	2	1	3	1	1	1	1	-	-
Male	1	2	1	-	1	1	1	-	1	-	-	-
Female	1	-	-	2	-	2	-	1	-	1	-	-
White, both sexes	1	2	1	2	1	3	1	1	1	1	-	-
Male	1	2	1	-	1	1	1	-	1	-	-	-
Female	-	-	-	2	-	2	-	1	-	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
United States												
All races, both sexes	23,313	3	4	2	2	8	28	56	69	117	165	308
Male	9,490	1	3	2	1	6	13	28	41	68	102	188
Female	13,823	2	1	-	1	2	15	28	28	49	63	120
White, both sexes	21,538	2	2	2	2	6	24	47	52	83	118	236
Male	8,709	1	2	2	1	4	11	24	31	44	76	151
Female	12,829	1	-	-	1	2	13	23	21	39	42	85
Black, both sexes	1,388	-	2	-	-	2	4	6	15	31	43	62
Male	614	-	1	-	-	2	2	3	10	22	26	30
Female	774	-	1	-	-	-	2	3	5	9	17	32
American Indian 1/, both sexes	68	-	-	-	-	-	-	1	2	1	2	2
Male	28	-	-	-	-	-	-	1	-	-	-	2
Female	40	-	-	-	-	-	-	-	2	1	2	-
Asian or Pacific Islander 2/, both sexes	319	1	-	-	-	-	-	2	-	2	2	8
Male	139	-	-	-	-	-	-	-	-	2	-	5
Female	180	1	-	-	-	-	-	2	-	-	2	3
Alabama												
All races, both sexes	282	-	-	-	-	-	2	1	2	1	3	3
Male	121	-	-	-	-	-	2	-	2	-	-	2
Female	161	-	-	-	-	-	-	1	-	1	3	1
White, both sexes	228	-	-	-	-	-	2	-	1	1	1	1
Male	94	-	-	-	-	-	2	-	1	-	-	-
Female	134	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	54	-	-	-	-	-	-	1	1	-	2	2
Male	27	-	-	-	-	-	-	-	1	-	-	2
Female	27	-	-	-	-	-	-	1	-	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Alaska												
All races, both sexes	42	-	-	-	-	-	-	-	-	-	1	1
Male	21	-	-	-	-	-	-	-	-	-	-	1
Female	21	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	33	-	-	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	388	1	-	-	-	-	-	1	2	-	3	7
Male	154	-	-	-	-	-	-	1	2	-	1	4
Female	234	1	-	-	-	-	-	-	-	-	2	3
White, both sexes	374	1	-	-	-	-	-	1	1	-	3	5
Male	146	-	-	-	-	-	-	1	1	-	1	3
Female	228	1	-	-	-	-	-	-	-	-	2	2
Black, both sexes	7	-	-	-	-	-	-	-	1	-	-	1
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Arkansas												
All races, both sexes	173	-	-	-	-	-	-	1	-	-	2	1
Male	62	-	-	-	-	-	-	1	-	-	-	1
Female	111	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	159	-	-	-	-	-	-	1	-	-	1	-
Male	59	-	-	-	-	-	-	1	-	-	-	-
Female	100	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	2,547	-	1	-	-	1	2	8	5	15	22	24
Male	1,051	-	-	-	-	1	-	4	2	8	14	17
Female	1,496	-	1	-	-	-	2	4	3	7	8	7
White, both sexes	2,306	-	-	-	-	1	1	6	4	13	19	18
Male	946	-	-	-	-	1	-	4	1	6	14	15
Female	1,360	-	-	-	-	-	1	2	3	7	5	3
Black, both sexes	108	-	1	-	-	-	1	1	1	2	1	1
Male	45	-	-	-	-	-	-	-	1	2	-	-
Female	63	-	1	-	-	-	1	1	-	-	1	1
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	124	-	-	-	-	-	-	1	-	-	2	4
Male	55	-	-	-	-	-	-	-	-	-	-	1
Female	69	-	-	-	-	-	-	1	-	-	2	3

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Colorado												
All races, both sexes	284	-	-	-	-	-	-	-	1	1	1	1
Male	126	-	-	-	-	-	-	-	-	-	-	1
Female	158	-	-	-	-	-	-	-	1	1	1	-
White, both sexes	276	-	-	-	-	-	-	-	1	1	-	1
Male	125	-	-	-	-	-	-	-	-	-	-	1
Female	151	-	-	-	-	-	-	-	1	1	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	310	-	-	-	-	-	1	2	2	1	1	7
Male	124	-	-	-	-	-	-	-	1	1	1	2
Female	186	-	-	-	-	-	1	2	1	-	-	5
White, both sexes	294	-	-	-	-	-	1	2	2	-	1	5
Male	118	-	-	-	-	-	-	-	1	-	1	2
Female	176	-	-	-	-	-	1	2	1	-	-	3
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Delaware												
All races, both sexes	82	-	-	-	-	-	1	-	-	1	1	2
Male	34	-	-	-	-	-	-	-	-	-	1	1
Female	48	-	-	-	-	-	1	-	-	1	-	1
White, both sexes	72	-	-	-	-	-	1	-	-	1	1	-
Male	26	-	-	-	-	-	-	-	-	-	1	-
Female	46	-	-	-	-	-	1	-	-	1	-	-
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	2
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	33	-	-	-	-	-	-	-	-	1	-	-
Male	19	-	-	-	-	-	-	-	-	1	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	21	-	-	-	-	-	-	-	-	1	-	-
Male	13	-	-	-	-	-	-	-	-	1	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Florida												
All races, both sexes	1,666	-	-	-	-	-	2	2	4	11	5	26
Male	781	-	-	-	-	-	2	1	3	9	4	16
Female	885	-	-	-	-	-	-	1	1	2	1	10
White, both sexes	1,548	-	-	-	-	-	2	2	2	8	3	16
Male	728	-	-	-	-	-	2	1	1	7	3	12
Female	820	-	-	-	-	-	-	1	1	1	-	4
Black, both sexes	110	-	-	-	-	-	-	-	2	3	2	9
Male	48	-	-	-	-	-	-	-	2	2	1	3
Female	62	-	-	-	-	-	-	-	-	1	1	6
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	427	-	-	-	1	-	1	2	3	2	6	9
Male	183	-	-	-	-	-	1	-	-	2	3	5
Female	244	-	-	-	1	-	-	2	3	-	3	4
White, both sexes	344	-	-	-	1	-	-	1	3	2	5	4
Male	145	-	-	-	-	-	-	-	-	2	2	1
Female	199	-	-	-	1	-	-	1	3	-	3	3
Black, both sexes	81	-	-	-	-	-	1	1	-	-	1	5
Male	38	-	-	-	-	-	1	-	-	-	1	4
Female	43	-	-	-	-	-	-	1	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Hawaii												
All races, both sexes	98	-	-	-	-	-	-	-	-	-	-	1
Male	48	-	-	-	-	-	-	-	-	-	-	1
Female	50	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	27	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	71	-	-	-	-	-	-	-	-	-	-	1
Male	35	-	-	-	-	-	-	-	-	-	-	1
Female	36	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	134	-	-	-	-	-	-	-	-	-	-	1
Male	61	-	-	-	-	-	-	-	-	-	-	1
Female	73	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	134	-	-	-	-	-	-	-	-	-	-	1
Male	61	-	-	-	-	-	-	-	-	-	-	1
Female	73	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Illinois												
All races, both sexes	811	-	2	-	-	-	2	1	1	4	4	19
Male	334	-	2	-	-	-	1	1	1	4	1	10
Female	477	-	-	-	-	-	1	-	-	-	3	9
White, both sexes	739	-	1	-	-	-	2	1	1	3	1	15
Male	298	-	1	-	-	-	1	1	1	3	-	10
Female	441	-	-	-	-	-	1	-	-	-	1	5
Black, both sexes	67	-	1	-	-	-	-	-	-	1	3	4
Male	32	-	1	-	-	-	-	-	-	1	1	-
Female	35	-	-	-	-	-	-	-	-	-	2	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	513	-	-	-	-	-	-	3	1	4	4	9
Male	232	-	-	-	-	-	-	1	1	2	4	7
Female	281	-	-	-	-	-	-	2	-	2	-	2
White, both sexes	494	-	-	-	-	-	-	3	1	3	3	7
Male	221	-	-	-	-	-	-	1	1	1	3	6
Female	273	-	-	-	-	-	-	2	-	2	-	1
Black, both sexes	18	-	-	-	-	-	-	-	-	-	1	2
Male	10	-	-	-	-	-	-	-	-	-	1	1
Female	8	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Iowa												
All races, both sexes	319	-	-	-	-	-	-	-	1	-	2	2
Male	129	-	-	-	-	-	-	-	1	-	1	2
Female	190	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	317	-	-	-	-	-	-	-	1	-	2	2
Male	129	-	-	-	-	-	-	-	1	-	1	2
Female	188	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	240	-	-	-	-	-	-	-	1	1	2	3
Male	83	-	-	-	-	-	-	-	-	1	2	1
Female	157	-	-	-	-	-	-	-	1	-	-	2
White, both sexes	231	-	-	-	-	-	-	-	1	1	2	3
Male	82	-	-	-	-	-	-	-	-	1	2	1
Female	149	-	-	-	-	-	-	-	1	-	-	2
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Kentucky												
All races, both sexes	248	-	-	-	-	-	-	-	1	1	1	5
Male	97	-	-	-	-	-	-	-	-	-	1	3
Female	151	-	-	-	-	-	-	-	1	1	-	2
White, both sexes	240	-	-	-	-	-	-	-	1	1	1	5
Male	94	-	-	-	-	-	-	-	-	-	1	3
Female	146	-	-	-	-	-	-	-	1	1	-	2
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	281	-	-	-	-	1	1	-	-	2	1	7
Male	104	-	-	-	-	1	1	-	-	1	-	5
Female	177	-	-	-	-	-	-	-	-	1	1	2
White, both sexes	230	-	-	-	-	1	1	-	-	2	1	3
Male	84	-	-	-	-	1	1	-	-	1	-	2
Female	146	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	48	-	-	-	-	-	-	-	-	-	-	4
Male	19	-	-	-	-	-	-	-	-	-	-	3
Female	29	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Maine												
All races, both sexes	181	-	-	-	-	-	-	-	1	-	1	-
Male	74	-	-	-	-	-	-	-	1	-	1	-
Female	107	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	181	-	-	-	-	-	-	-	1	-	1	-
Male	74	-	-	-	-	-	-	-	1	-	1	-
Female	107	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	381	-	-	-	-	-	-	1	1	4	4	5
Male	180	-	-	-	-	-	-	-	1	3	3	2
Female	201	-	-	-	-	-	-	1	-	1	1	3
White, both sexes	317	-	-	-	-	-	-	1	-	1	1	3
Male	153	-	-	-	-	-	-	-	-	1	1	2
Female	164	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	59	-	-	-	-	-	-	-	1	3	3	2
Male	26	-	-	-	-	-	-	-	1	2	2	-
Female	33	-	-	-	-	-	-	-	-	1	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Massachusetts												
All races, both sexes	566	-	-	-	-	2	1	3	3	4	7	4
Male	207	-	-	-	-	1	-	1	3	2	7	3
Female	359	-	-	-	-	1	1	2	-	2	-	1
White, both sexes	547	-	-	-	-	2	1	3	2	3	6	4
Male	200	-	-	-	-	1	-	1	2	2	6	3
Female	347	-	-	-	-	1	1	2	-	1	-	1
Black, both sexes	15	-	-	-	-	-	-	-	1	1	1	-
Male	6	-	-	-	-	-	-	-	1	-	1	-
Female	9	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	727	-	-	-	-	1	1	1	3	11	1	17
Male	322	-	-	-	-	-	1	1	1	6	1	12
Female	405	-	-	-	-	1	-	-	2	5	-	5
White, both sexes	663	-	-	-	-	1	1	1	2	8	1	15
Male	290	-	-	-	-	-	1	1	1	4	1	11
Female	373	-	-	-	-	1	-	-	1	4	-	4
Black, both sexes	58	-	-	-	-	-	-	-	1	3	-	2
Male	29	-	-	-	-	-	-	-	-	2	-	1
Female	29	-	-	-	-	-	-	-	1	1	-	1
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Minnesota												
All races, both sexes	473	-	-	-	-	-	-	-	1	1	2	5
Male	209	-	-	-	-	-	-	-	-	-	2	3
Female	264	-	-	-	-	-	-	-	1	1	-	2
White, both sexes	464	-	-	-	-	-	-	-	1	-	2	5
Male	209	-	-	-	-	-	-	-	-	-	2	3
Female	255	-	-	-	-	-	-	-	1	-	-	2
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	129	-	-	-	1	1	-	-	-	2	2	3
Male	52	-	-	-	1	1	-	-	-	2	-	-
Female	77	-	-	-	-	-	-	-	-	-	2	3
White, both sexes	97	-	-	-	1	-	-	-	-	1	1	2
Male	39	-	-	-	1	-	-	-	-	1	-	-
Female	58	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	30	-	-	-	-	1	-	-	-	1	1	1
Male	12	-	-	-	-	1	-	-	-	1	-	-
Female	18	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Missouri												
All races, both sexes	512	-	1	-	-	-	2	-	3	3	7	7
Male	199	-	1	-	-	-	1	-	2	2	5	4
Female	313	-	-	-	-	-	1	-	1	1	2	3
White, both sexes	483	-	1	-	-	-	2	-	2	1	7	6
Male	189	-	1	-	-	-	1	-	2	-	5	4
Female	294	-	-	-	-	-	1	-	-	1	2	2
Black, both sexes	27	-	-	-	-	-	-	-	-	2	-	1
Male	10	-	-	-	-	-	-	-	-	2	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	114	-	-	-	-	-	-	1	1	-	-	2
Male	52	-	-	-	-	-	-	-	-	-	-	1
Female	62	-	-	-	-	-	-	1	1	-	-	1
White, both sexes	112	-	-	-	-	-	-	1	-	-	-	1
Male	51	-	-	-	-	-	-	-	-	-	-	-
Female	61	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	1	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Nebraska												
All races, both sexes	195	-	-	-	-	-	1	-	-	-	1	4
Male	72	-	-	-	-	-	-	-	-	-	-	3
Female	123	-	-	-	-	-	1	-	-	-	1	1
White, both sexes	193	-	-	-	-	-	1	-	-	-	1	4
Male	71	-	-	-	-	-	-	-	-	-	-	3
Female	122	-	-	-	-	-	1	-	-	-	1	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	131	-	-	-	-	-	1	1	-	1	5	2
Male	67	-	-	-	-	-	1	1	-	1	4	2
Female	64	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	117	-	-	-	-	-	1	1	-	1	3	1
Male	60	-	-	-	-	-	1	1	-	1	2	1
Female	57	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	2	1
Male	4	-	-	-	-	-	-	-	-	-	2	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
New Hampshire												
All races, both sexes	133	-	-	-	-	-	-	-	-	-	-	-
Male	50	-	-	-	-	-	-	-	-	-	-	-
Female	83	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	133	-	-	-	-	-	-	-	-	-	-	-
Male	50	-	-	-	-	-	-	-	-	-	-	-
Female	83	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	861	-	-	-	-	-	1	3	1	6	6	14
Male	344	-	-	-	-	-	-	2	1	1	3	10
Female	517	-	-	-	-	-	1	1	-	5	3	4
White, both sexes	802	-	-	-	-	-	1	1	-	4	2	9
Male	314	-	-	-	-	-	-	1	-	1	2	5
Female	488	-	-	-	-	-	1	-	-	3	-	4
Black, both sexes	51	-	-	-	-	-	-	1	1	2	4	5
Male	27	-	-	-	-	-	-	1	1	-	1	5
Female	24	-	-	-	-	-	-	-	-	2	3	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
New Mexico												
All races, both sexes	121	-	-	-	-	-	-	1	-	-	-	1
Male	56	-	-	-	-	-	-	1	-	-	-	1
Female	65	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	113	-	-	-	-	-	-	1	-	-	-	1
Male	52	-	-	-	-	-	-	1	-	-	-	1
Female	61	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1,325	-	-	-	-	-	1	2	3	2	11	8
Male	485	-	-	-	-	-	1	1	2	2	4	6
Female	840	-	-	-	-	-	-	1	1	-	7	2
White, both sexes	1,250	-	-	-	-	-	1	2	3	-	7	7
Male	458	-	-	-	-	-	1	1	2	-	2	5
Female	792	-	-	-	-	-	-	1	1	-	5	2
Black, both sexes	55	-	-	-	-	-	-	-	-	2	4	1
Male	21	-	-	-	-	-	-	-	-	2	2	1
Female	34	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
North Carolina												
All races, both sexes	632	-	-	-	-	-	-	-	2	5	3	10
Male	257	-	-	-	-	-	-	-	2	3	1	6
Female	375	-	-	-	-	-	-	-	-	2	2	4
White, both sexes	534	-	-	-	-	-	-	-	1	3	2	5
Male	214	-	-	-	-	-	-	-	1	1	-	4
Female	320	-	-	-	-	-	-	-	-	2	2	1
Black, both sexes	92	-	-	-	-	-	-	-	1	2	1	5
Male	42	-	-	-	-	-	-	-	1	2	1	2
Female	50	-	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	67	-	-	-	-	-	-	1	1	-	-	-
Male	27	-	-	-	-	-	-	-	1	-	-	-
Female	40	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	65	-	-	-	-	-	-	1	1	-	-	-
Male	25	-	-	-	-	-	-	-	1	-	-	-
Female	40	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Ohio												
All races, both sexes	1,106	-	-	-	-	-	1	-	2	1	7	11
Male	404	-	-	-	-	-	-	-	-	-	6	7
Female	702	-	-	-	-	-	1	-	2	1	1	4
White, both sexes	1,041	-	-	-	-	-	1	-	2	1	5	7
Male	376	-	-	-	-	-	-	-	-	-	4	4
Female	665	-	-	-	-	-	1	-	2	1	1	3
Black, both sexes	63	-	-	-	-	-	-	-	-	-	2	4
Male	28	-	-	-	-	-	-	-	-	-	2	3
Female	35	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	246	-	-	-	-	-	-	2	1	1	6	4
Male	112	-	-	-	-	-	-	1	1	1	2	2
Female	134	-	-	-	-	-	-	1	-	-	4	2
White, both sexes	217	-	-	-	-	-	-	1	1	1	4	4
Male	101	-	-	-	-	-	-	-	1	1	1	2
Female	116	-	-	-	-	-	-	1	-	-	3	2
Black, both sexes	19	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	10	-	-	-	-	-	-	1	-	-	1	-
Male	6	-	-	-	-	-	-	1	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Oregon												
All races, both sexes	576	-	-	1	-	-	1	-	-	2	-	3
Male	247	-	-	1	-	-	-	-	-	1	-	3
Female	329	-	-	-	-	-	1	-	-	1	-	-
White, both sexes	570	-	-	1	-	-	1	-	-	2	-	3
Male	244	-	-	1	-	-	-	-	-	1	-	3
Female	326	-	-	-	-	-	1	-	-	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1,563	-	-	-	-	-	2	3	2	7	9	19
Male	564	-	-	-	-	-	1	1	1	1	4	9
Female	999	-	-	-	-	-	1	2	1	6	5	10
White, both sexes	1,508	-	-	-	-	-	1	3	1	5	8	17
Male	539	-	-	-	-	-	1	1	-	-	3	9
Female	969	-	-	-	-	-	-	2	1	5	5	8
Black, both sexes	53	-	-	-	-	-	1	-	1	2	1	2
Male	23	-	-	-	-	-	-	-	1	1	1	-
Female	30	-	-	-	-	-	1	-	-	1	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Rhode Island												
All races, both sexes	91	-	-	-	-	-	-	-	-	-	-	2
Male	32	-	-	-	-	-	-	-	-	-	-	2
Female	59	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	90	-	-	-	-	-	-	-	-	-	-	2
Male	32	-	-	-	-	-	-	-	-	-	-	2
Female	58	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	330	-	-	-	-	1	-	-	2	4	2	2
Male	122	-	-	-	-	1	-	-	-	2	2	1
Female	208	-	-	-	-	-	-	-	2	2	-	1
White, both sexes	263	-	-	-	-	-	-	-	-	3	1	-
Male	101	-	-	-	-	-	-	-	-	1	1	-
Female	162	-	-	-	-	-	-	-	-	2	-	-
Black, both sexes	67	-	-	-	-	1	-	-	2	1	1	2
Male	21	-	-	-	-	1	-	-	-	1	1	1
Female	46	-	-	-	-	-	-	-	2	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
South Dakota												
All races, both sexes	76	-	-	-	-	-	-	-	1	-	-	2
Male	28	-	-	-	-	-	-	-	1	-	-	1
Female	48	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	75	-	-	-	-	-	-	-	1	-	-	2
Male	28	-	-	-	-	-	-	-	1	-	-	1
Female	47	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	432	-	-	-	-	-	1	3	1	3	6	9
Male	185	-	-	-	-	-	-	2	1	2	5	7
Female	247	-	-	-	-	-	1	1	-	1	1	2
White, both sexes	390	-	-	-	-	-	1	2	1	2	4	8
Male	164	-	-	-	-	-	-	1	1	2	3	6
Female	226	-	-	-	-	-	1	1	-	-	1	2
Black, both sexes	42	-	-	-	-	-	-	1	-	1	2	1
Male	21	-	-	-	-	-	-	1	-	-	2	1
Female	21	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Texas												
All races, both sexes	1,162	-	-	-	-	-	1	8	5	7	11	24
Male	519	-	-	-	-	-	-	7	4	6	7	11
Female	643	-	-	-	-	-	1	1	1	1	4	13
White, both sexes	1,054	-	-	-	-	-	1	7	4	4	7	23
Male	461	-	-	-	-	-	-	6	4	3	3	11
Female	593	-	-	-	-	-	1	1	-	1	4	12
Black, both sexes	96	-	-	-	-	-	-	1	1	3	4	1
Male	51	-	-	-	-	-	-	1	-	3	4	-
Female	45	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	152	-	-	-	-	-	-	1	1	1	-	2
Male	59	-	-	-	-	-	-	1	1	1	-	2
Female	93	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	148	-	-	-	-	-	-	1	1	1	-	2
Male	58	-	-	-	-	-	-	1	1	1	-	2
Female	90	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Vermont												
All races, both sexes	69	-	-	-	-	-	-	1	-	-	-	-
Male	32	-	-	-	-	-	-	-	-	-	-	-
Female	37	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	69	-	-	-	-	-	-	1	-	-	-	-
Male	32	-	-	-	-	-	-	-	-	-	-	-
Female	37	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	516	1	-	1	-	-	2	1	2	2	5	7
Male	190	-	-	1	-	-	1	-	1	-	3	2
Female	326	1	-	-	-	-	1	1	1	2	2	5
White, both sexes	453	-	-	1	-	-	1	1	1	1	1	6
Male	162	-	-	1	-	-	-	-	1	-	1	2
Female	291	-	-	-	-	-	1	1	-	1	-	4
Black, both sexes	56	-	-	-	-	-	1	-	1	1	4	1
Male	25	-	-	-	-	-	1	-	-	-	2	-
Female	31	-	-	-	-	-	-	-	1	1	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	1	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Washington												
All races, both sexes	788	-	-	-	-	1	-	-	4	3	2	7
Male	304	-	-	-	-	1	-	-	2	3	2	4
Female	484	-	-	-	-	-	-	-	2	-	-	3
White, both sexes	764	-	-	-	-	1	-	-	4	3	2	7
Male	300	-	-	-	-	1	-	-	2	3	2	4
Female	464	-	-	-	-	-	-	-	2	-	-	3
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	144	-	-	-	-	-	-	1	1	-	3	3
Male	56	-	-	-	-	-	-	-	-	-	2	2
Female	88	-	-	-	-	-	-	1	1	-	1	1
White, both sexes	141	-	-	-	-	-	-	1	1	-	3	3
Male	56	-	-	-	-	-	-	-	-	-	2	2
Female	85	-	-	-	-	-	-	1	1	-	1	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Wisconsin												
All races, both sexes	594	1	-	-	-	-	-	1	3	1	5	3
Male	201	1	-	-	-	-	-	-	2	-	5	2
Female	393	-	-	-	-	-	-	1	1	1	-	1
White, both sexes	584	1	-	-	-	-	-	1	3	1	5	3
Male	196	1	-	-	-	-	-	-	2	-	5	2
Female	388	-	-	-	-	-	-	1	1	1	-	1
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	42	-	-	-	-	-	-	-	-	1	-	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	42	-	-	-	-	-	-	-	-	1	-	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
United States												
All races, both sexes	431	524	644	913	1,374	2,367	4,036	5,640	4,596	1,742	284	-
Male	282	328	363	486	685	1,091	1,754	2,135	1,454	411	48	-
Female	149	196	281	427	689	1,276	2,282	3,505	3,142	1,331	236	-
White, both sexes	317	402	537	779	1,227	2,176	3,787	5,392	4,406	1,672	269	-
Male	210	253	305	421	622	1,008	1,651	2,054	1,406	390	42	-
Female	107	149	232	358	605	1,168	2,136	3,338	3,000	1,282	227	-
Black, both sexes	106	108	93	111	116	148	183	170	126	50	12	-
Male	66	67	51	55	47	63	78	50	27	10	4	-
Female	40	41	42	56	69	85	105	120	99	40	8	-
American Indian 1/, both sexes	4	5	3	6	6	7	10	7	8	2	2	-
Male	4	3	1	2	3	5	1	2	2	1	1	-
Female	-	2	2	4	3	2	9	5	6	1	1	-
Asian or Pacific Islander 2/, both sexes	4	9	11	17	25	36	56	71	56	18	1	-
Male	2	5	6	8	13	15	24	29	19	10	1	-
Female	2	4	5	9	12	21	32	42	37	8	-	-
Alabama												
All races, both sexes	8	10	9	14	28	42	49	53	37	17	3	-
Male	5	5	4	10	13	19	20	18	16	5	-	-
Female	3	5	5	4	15	23	29	35	21	12	3	-
White, both sexes	3	6	6	7	24	35	41	48	33	16	3	-
Male	-	2	1	6	13	16	18	15	15	5	-	-
Female	3	4	5	1	11	19	23	33	18	11	3	-
Black, both sexes	5	4	3	7	4	7	8	5	4	1	-	-
Male	5	3	3	4	-	3	2	3	1	-	-	-
Female	-	1	-	3	4	4	6	2	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Alaska												
All races, both sexes	4	2	1	2	7	6	5	5	5	3	-	-
Male	3	2	1	1	3	3	2	3	1	1	-	-
Female	1	-	-	1	4	3	3	2	4	2	-	-
White, both sexes	1	2	1	2	5	5	4	5	5	3	-	-
Male	1	2	1	1	3	2	2	3	1	1	-	-
Female	-	-	-	1	2	3	2	2	4	2	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	1	1	1	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
Arizona												
All races, both sexes	9	14	13	19	23	46	66	87	71	23	3	-
Male	8	9	7	11	9	22	30	28	18	4	-	-
Female	1	5	6	8	14	24	36	59	53	19	3	-
White, both sexes	8	13	12	17	22	45	64	86	70	23	3	-
Male	7	8	6	10	8	22	29	28	18	4	-	-
Female	1	5	6	7	14	23	35	58	52	19	3	-
Black, both sexes	-	1	-	1	1	1	-	-	1	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	1	-	-	-
American Indian 1/, both sexes	1	-	-	1	-	-	2	1	-	-	-	-
Male	1	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Arkansas												
All races, both sexes	3	7	8	9	15	19	32	35	22	16	3	-
Male	-	5	3	5	6	8	11	11	7	3	1	-
Female	3	2	5	4	9	11	21	24	15	13	2	-
White, both sexes	2	7	6	8	15	17	31	33	20	16	2	-
Male	-	5	2	5	6	8	10	11	7	3	1	-
Female	2	2	4	3	9	9	21	22	13	13	1	-
Black, both sexes	1	-	-	1	-	2	1	2	2	-	1	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	1	-	2	-	2	2	-	1	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	37	52	55	78	122	266	425	620	556	209	49	-
Male	29	33	38	42	56	117	192	246	190	53	9	-
Female	8	19	17	36	66	149	233	374	366	156	40	-
White, both sexes	27	41	48	65	103	237	385	580	516	195	47	-
Male	23	28	32	34	48	104	173	229	176	49	9	-
Female	4	13	16	31	55	133	212	351	340	146	38	-
Black, both sexes	10	8	5	7	7	12	16	14	14	6	1	-
Male	6	3	4	5	3	5	5	6	5	-	-	-
Female	4	5	1	2	4	7	11	8	9	6	1	-
American Indian 1/, both sexes	-	-	-	-	-	1	1	2	2	1	1	-
Male	-	-	-	-	-	1	-	1	1	1	-	-
Female	-	-	-	-	-	-	1	1	1	-	1	-
Asian or Pacific Islander 2/, both sexes	-	3	2	6	12	16	23	24	24	7	-	-
Male	-	2	2	3	5	7	14	10	8	3	-	-
Female	-	1	-	3	7	9	9	14	16	4	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Colorado												
All races, both sexes	5	1	10	12	18	36	53	69	51	22	3	-
Male	5	1	5	9	10	18	26	29	16	6	-	-
Female	-	-	5	3	8	18	27	40	35	16	3	-
White, both sexes	5	1	9	12	17	35	51	67	51	22	3	-
Male	5	1	4	9	10	18	26	29	16	6	-	-
Female	-	-	5	3	7	17	25	38	35	16	3	-
Black, both sexes	-	-	1	-	-	-	1	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	1	1	-	-	-	-
Connecticut												
All races, both sexes	6	7	10	7	11	20	55	77	77	22	4	-
Male	5	5	6	5	6	12	24	24	26	6	-	-
Female	1	2	4	2	5	8	31	53	51	16	4	-
White, both sexes	6	5	7	6	10	19	54	74	77	21	4	-
Male	5	4	5	4	5	11	24	24	26	6	-	-
Female	1	1	2	2	5	8	30	50	51	15	4	-
Black, both sexes	-	2	3	1	1	1	1	3	-	-	-	-
Male	-	1	1	1	1	1	-	-	-	-	-	-
Female	-	1	2	-	-	-	1	3	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Delaware												
All races, both sexes	-	5	3	6	4	8	17	15	14	4	1	-
Male	-	4	2	3	3	5	7	3	4	1	-	-
Female	-	1	1	3	1	3	10	12	10	3	1	-
White, both sexes	-	3	2	5	4	7	16	15	12	4	1	-
Male	-	2	1	2	3	4	6	3	3	1	-	-
Female	-	1	1	3	1	3	10	12	9	3	1	-
Black, both sexes	-	2	1	1	-	1	1	-	2	-	-	-
Male	-	2	1	1	-	1	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	3	5	1	1	2	5	4	1	5	5	-	-
Male	2	4	-	1	2	3	1	1	4	-	-	-
Female	1	1	1	-	-	2	3	-	1	5	-	-
White, both sexes	-	1	1	-	-	1	1	-	5	3	-	-
Male	-	-	-	-	-	1	1	-	4	-	-	-
Female	-	1	1	-	-	-	-	-	1	3	-	-
Black, both sexes	3	4	-	1	2	4	3	1	-	2	-	-
Male	2	4	-	1	2	2	-	1	-	-	-	-
Female	1	-	-	-	-	2	3	-	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Florida												
All races, both sexes	39	41	49	82	111	184	300	375	305	116	14	-
Male	22	25	19	50	60	95	152	168	114	37	4	-
Female	17	16	30	32	51	89	148	207	191	79	10	-
White, both sexes	28	34	36	71	101	173	284	363	299	113	13	-
Male	17	21	16	43	53	93	142	166	112	36	3	-
Female	11	13	20	28	48	80	142	197	187	77	10	-
Black, both sexes	11	7	12	10	10	9	15	11	6	2	1	-
Male	5	4	2	7	7	1	9	2	2	-	1	-
Female	6	3	10	3	3	8	6	9	4	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	2	1	1	-	1	-	-
Male	-	-	1	-	-	1	1	-	-	1	-	-
Female	-	-	-	1	-	1	-	1	-	-	-	-
Georgia												
All races, both sexes	11	16	16	23	38	52	57	100	60	27	3	-
Male	7	12	9	11	19	30	21	36	15	11	1	-
Female	4	4	7	12	19	22	36	64	45	16	2	-
White, both sexes	7	9	10	19	27	43	47	91	53	20	2	-
Male	4	7	6	10	14	27	16	35	14	7	-	-
Female	3	2	4	9	13	16	31	56	39	13	2	-
Black, both sexes	4	7	6	4	11	8	10	9	6	7	1	-
Male	3	5	3	1	5	3	5	1	1	4	1	-
Female	1	2	3	3	6	5	5	8	5	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Hawaii												
All races, both sexes	1	4	2	4	1	6	16	37	19	6	1	-
Male	-	3	1	2	1	4	5	17	10	3	1	-
Female	1	1	1	2	-	2	11	20	9	3	-	-
White, both sexes	-	2	1	1	-	1	6	8	7	1	-	-
Male	-	2	1	1	-	-	2	3	4	-	-	-
Female	-	-	-	-	-	1	4	5	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	1	3	1	5	10	29	12	5	1	-
Male	-	1	-	1	1	4	3	14	6	3	1	-
Female	1	1	1	2	-	1	7	15	6	2	-	-
Idaho												
All races, both sexes	-	3	2	6	8	9	34	39	21	10	1	-
Male	-	1	1	4	7	4	16	16	11	-	-	-
Female	-	2	1	2	1	5	18	23	10	10	1	-
White, both sexes	-	3	2	6	8	9	34	39	21	10	1	-
Male	-	1	1	4	7	4	16	16	11	-	-	-
Female	-	2	1	2	1	5	18	23	10	10	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Illinois												
All races, both sexes	11	18	16	34	49	80	134	192	167	71	6	-
Male	7	15	11	18	22	38	56	86	45	16	-	-
Female	4	3	5	16	27	42	78	106	122	55	6	-
White, both sexes	8	15	11	28	43	72	121	183	161	68	5	-
Male	6	12	6	14	21	34	48	82	43	15	-	-
Female	2	3	5	14	22	38	73	101	118	53	5	-
Black, both sexes	3	3	4	6	6	7	11	9	5	3	1	-
Male	1	3	4	4	1	3	6	4	2	1	-	-
Female	2	-	-	2	5	4	5	5	3	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	2	-	1	-	-	-
Male	-	-	1	-	-	1	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Indiana												
All races, both sexes	9	8	19	25	37	54	97	113	92	33	5	-
Male	5	6	12	13	15	23	49	52	32	10	-	-
Female	4	2	7	12	22	31	48	61	60	23	5	-
White, both sexes	8	7	18	24	34	54	94	111	89	33	5	-
Male	4	5	11	13	14	23	48	50	31	10	-	-
Female	4	2	7	11	20	31	46	61	58	23	5	-
Black, both sexes	1	1	1	1	3	-	3	2	3	-	-	-
Male	1	1	1	-	1	-	1	2	1	-	-	-
Female	-	-	-	1	2	-	2	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Iowa												
All races, both sexes	3	5	5	5	14	23	50	84	89	33	3	-
Male	3	3	1	1	5	16	23	35	28	10	-	-
Female	-	2	4	4	9	7	27	49	61	23	3	-
White, both sexes	3	5	4	5	14	22	50	84	89	33	3	-
Male	3	3	1	1	5	16	23	35	28	10	-	-
Female	-	2	3	4	9	6	27	49	61	23	3	-
Black, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	3	3	2	3	4	21	45	59	62	26	5	-
Male	1	2	1	1	2	9	13	28	15	6	1	-
Female	2	1	1	2	2	12	32	31	47	20	4	-
White, both sexes	3	2	2	3	3	21	43	58	59	25	5	-
Male	1	2	1	1	2	9	13	28	15	5	1	-
Female	2	-	1	2	1	12	30	30	44	20	4	-
Black, both sexes	-	1	-	-	1	-	2	1	2	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	1	-	-	1	-	2	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Kentucky												
All races, both sexes	5	7	8	13	18	34	49	50	35	18	3	-
Male	4	3	4	6	7	19	25	14	8	3	-	-
Female	1	4	4	7	11	15	24	36	27	15	3	-
White, both sexes	5	6	8	11	18	32	47	50	34	18	3	-
Male	4	2	4	5	7	19	24	14	8	3	-	-
Female	1	4	4	6	11	13	23	36	26	15	3	-
Black, both sexes	-	-	-	2	-	2	2	-	1	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	2	1	-	1	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	9	7	8	20	12	29	64	60	42	15	3	-
Male	5	3	3	11	5	15	24	16	11	2	1	-
Female	4	4	5	9	7	14	40	44	31	13	2	-
White, both sexes	5	5	4	12	12	24	53	55	34	15	3	-
Male	3	2	1	6	5	12	20	16	11	2	1	-
Female	2	3	3	6	7	12	33	39	23	13	2	-
Black, both sexes	4	2	4	7	-	5	10	5	7	-	-	-
Male	2	1	2	4	-	3	4	-	-	-	-	-
Female	2	1	2	3	-	2	6	5	7	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	1	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Maine												
All races, both sexes	3	2	8	9	9	18	27	48	38	14	3	-
Male	2	1	4	2	5	7	14	19	12	5	1	-
Female	1	1	4	7	4	11	13	29	26	9	2	-
White, both sexes	3	2	8	9	9	18	27	48	38	14	3	-
Male	2	1	4	2	5	7	14	19	12	5	1	-
Female	1	1	4	7	4	11	13	29	26	9	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	8	9	12	14	18	29	74	101	71	25	5	-
Male	5	5	9	9	9	15	30	48	33	8	-	-
Female	3	4	3	5	9	14	44	53	38	17	5	-
White, both sexes	2	7	10	8	10	25	67	88	68	21	5	-
Male	2	4	7	7	7	14	26	44	32	6	-	-
Female	-	3	3	1	3	11	41	44	36	15	5	-
Black, both sexes	6	2	2	6	7	4	5	12	3	3	-	-
Male	3	1	2	2	2	1	3	4	1	2	-	-
Female	3	1	-	4	5	3	2	8	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	2	1	-	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	1	1	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Massachusetts												
All races, both sexes	11	5	9	16	23	43	96	149	132	47	11	-
Male	6	3	4	6	15	24	45	49	32	5	1	-
Female	5	2	5	10	8	19	51	100	100	42	10	-
White, both sexes	11	4	9	16	23	41	93	144	130	44	11	-
Male	6	2	4	6	15	23	43	48	32	5	1	-
Female	5	2	5	10	8	18	50	96	98	39	10	-
Black, both sexes	-	1	-	-	-	2	2	4	1	2	-	-
Male	-	1	-	-	-	1	2	-	-	-	-	-
Female	-	-	-	-	-	1	-	4	1	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	1	1	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	1	-	-
Michigan												
All races, both sexes	21	25	21	28	37	74	113	164	143	59	7	-
Male	19	17	15	15	18	36	49	67	48	15	1	-
Female	2	8	6	13	19	38	64	97	95	44	6	-
White, both sexes	9	18	17	27	34	66	103	158	136	59	7	-
Male	9	14	14	15	15	33	46	64	45	15	1	-
Female	-	4	3	12	19	33	57	94	91	44	6	-
Black, both sexes	12	7	4	1	2	8	7	5	6	-	-	-
Male	10	3	1	-	2	3	2	3	2	-	-	-
Female	2	4	3	1	-	5	5	2	4	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Minnesota												
All races, both sexes	4	7	7	8	23	42	79	131	113	45	5	-
Male	2	5	3	4	13	19	48	58	41	11	-	-
Female	2	2	4	4	10	23	31	73	72	34	5	-
White, both sexes	4	6	7	8	23	40	79	128	112	45	4	-
Male	2	5	3	4	13	19	48	58	41	11	-	-
Female	2	1	4	4	10	21	31	70	71	34	4	-
Black, both sexes	-	-	-	-	-	2	-	2	1	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	2	1	-	1	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	3	4	8	7	12	12	30	25	13	5	1	-
Male	2	3	4	4	5	5	11	8	3	2	1	-
Female	1	1	4	3	7	7	19	17	10	3	-	-
White, both sexes	1	2	6	2	11	9	27	19	10	4	1	-
Male	1	1	3	1	4	3	11	7	3	2	1	-
Female	-	1	3	1	7	6	16	12	7	2	-	-
Black, both sexes	2	2	2	5	1	2	2	6	3	1	-	-
Male	1	2	1	3	1	1	-	1	-	-	-	-
Female	1	-	1	2	-	1	2	5	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Missouri												
All races, both sexes	7	6	19	24	33	57	81	117	109	32	4	-
Male	5	2	10	10	16	27	32	43	29	8	2	-
Female	2	4	9	14	17	30	49	74	80	24	2	-
White, both sexes	6	5	17	24	29	53	80	110	106	31	3	-
Male	4	2	8	10	15	25	32	41	29	8	2	-
Female	2	3	9	14	14	28	48	69	77	23	1	-
Black, both sexes	1	1	2	-	4	4	1	7	2	1	1	-
Male	1	-	2	-	1	2	-	2	-	-	-	-
Female	-	1	-	-	3	2	1	5	2	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Montana												
All races, both sexes	2	3	4	5	7	7	22	25	24	10	1	-
Male	1	2	2	5	4	4	6	12	12	3	-	-
Female	1	1	2	-	3	3	16	13	12	7	1	-
White, both sexes	2	3	4	5	7	7	22	25	24	10	1	-
Male	1	2	2	5	4	4	6	12	12	3	-	-
Female	1	1	2	-	3	3	16	13	12	7	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Nebraska												
All races, both sexes	5	2	-	4	4	11	30	58	52	21	2	-
Male	3	2	-	3	3	3	14	15	20	6	-	-
Female	2	-	-	1	1	8	16	43	32	15	2	-
White, both sexes	5	2	-	4	3	11	30	58	51	21	2	-
Male	3	2	-	3	2	3	14	15	20	6	-	-
Female	2	-	-	1	1	8	16	43	31	15	2	-
Black, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	2	4	8	6	13	14	25	22	19	7	1	-
Male	2	3	6	3	7	6	13	9	7	2	-	-
Female	-	1	2	3	6	8	12	13	12	5	1	-
White, both sexes	2	4	8	5	11	11	23	22	18	5	1	-
Male	2	3	6	3	6	5	13	9	6	1	-	-
Female	-	1	2	2	5	6	10	13	12	4	1	-
Black, both sexes	-	-	-	-	1	1	2	-	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	2	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	2	-	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	1	1	-	-
Female	-	-	-	1	-	2	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
New Hampshire												
All races, both sexes	-	1	4	3	5	7	23	46	34	8	2	-
Male	-	1	1	2	1	2	9	22	8	3	1	-
Female	-	-	3	1	4	5	14	24	26	5	1	-
White, both sexes	-	1	4	3	5	7	23	46	34	8	2	-
Male	-	1	1	2	1	2	9	22	8	3	1	-
Female	-	-	3	1	4	5	14	24	26	5	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	15	16	15	19	42	87	147	242	174	61	12	-
Male	10	9	10	13	23	39	73	78	57	12	3	-
Female	5	7	5	6	19	48	74	164	117	49	9	-
White, both sexes	14	11	12	13	39	79	141	234	170	60	12	-
Male	10	7	8	9	22	33	69	76	57	11	3	-
Female	4	4	4	4	17	46	72	158	113	49	9	-
Black, both sexes	1	5	2	5	3	8	6	5	3	-	-	-
Male	-	2	2	3	1	6	4	1	-	-	-	-
Female	1	3	-	2	2	2	2	4	3	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	3	1	1	-	-
Male	-	-	-	-	-	-	-	1	-	1	-	-
Female	-	-	1	-	-	-	-	2	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
New Mexico												
All races, both sexes	5	4	5	7	6	16	24	29	14	9	-	-
Male	3	3	4	2	3	8	13	12	3	3	-	-
Female	2	1	1	5	3	8	11	17	11	6	-	-
White, both sexes	5	1	5	6	6	16	21	29	13	9	-	-
Male	3	1	4	2	3	8	11	12	3	3	-	-
Female	2	-	1	4	3	8	10	17	10	6	-	-
Black, both sexes	-	1	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	2	-	1	-	-	1	-	1	-	-	-
Male	-	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	17	31	32	42	82	143	214	327	280	109	21	-
Male	13	17	21	18	43	58	89	108	79	21	2	-
Female	4	14	11	24	39	85	125	219	201	88	19	-
White, both sexes	14	24	28	37	72	135	200	317	274	108	21	-
Male	11	15	19	16	37	57	85	107	77	21	2	-
Female	3	9	9	21	35	78	115	210	197	87	19	-
Black, both sexes	1	6	3	3	7	5	10	7	5	1	-	-
Male	-	2	2	1	4	1	4	-	2	-	-	-
Female	1	4	1	2	3	4	6	7	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	1	2	3	3	3	3	1	-	-	-
Male	2	-	-	1	2	-	-	1	-	-	-	-
Female	-	1	1	1	1	3	3	2	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
North Carolina												
All races, both sexes	22	9	25	25	46	60	127	153	101	34	10	-
Male	10	4	14	12	26	27	52	61	27	8	4	-
Female	12	5	11	13	20	33	75	92	74	26	6	-
White, both sexes	10	7	18	15	38	49	111	143	96	30	6	-
Male	6	4	10	7	22	21	48	55	25	8	2	-
Female	4	3	8	8	16	28	63	88	71	22	4	-
Black, both sexes	12	2	7	9	7	11	13	10	4	4	4	-
Male	4	-	4	5	4	6	4	6	1	-	2	-
Female	8	2	3	4	3	5	9	4	3	4	2	-
American Indian 1/, both sexes	-	-	-	1	1	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	1	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
North Dakota												
All races, both sexes	2	2	2	1	3	4	10	20	19	2	-	-
Male	2	2	2	-	1	3	5	7	4	-	-	-
Female	-	-	-	1	2	1	5	13	15	2	-	-
White, both sexes	1	2	2	1	3	4	10	19	19	2	-	-
Male	1	2	2	-	1	3	5	6	4	-	-	-
Female	-	-	-	1	2	1	5	13	15	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Ohio												
All races, both sexes	17	22	30	37	64	102	226	276	201	92	17	-
Male	8	12	14	21	28	43	88	97	56	21	3	-
Female	9	10	16	16	36	59	138	179	145	71	14	-
White, both sexes	13	15	26	31	58	99	214	271	192	89	17	-
Male	7	8	12	20	27	42	81	95	53	20	3	-
Female	6	7	14	11	31	57	133	176	139	69	14	-
Black, both sexes	4	7	4	6	6	3	12	5	7	3	-	-
Male	1	4	2	1	1	1	7	2	3	1	-	-
Female	3	3	2	5	5	2	5	3	4	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	8	7	9	13	21	35	41	54	31	11	2	-
Male	7	3	6	7	10	19	17	19	12	3	2	-
Female	1	4	3	6	11	16	24	35	19	8	-	-
White, both sexes	6	5	8	11	15	31	38	52	28	11	1	-
Male	5	3	5	7	9	17	16	19	11	3	1	-
Female	1	2	3	4	6	14	22	33	17	8	-	-
Black, both sexes	2	2	-	2	4	1	3	2	2	-	-	-
Male	2	-	-	-	-	-	1	-	1	-	-	-
Female	-	2	-	2	4	1	2	2	1	-	-	-
American Indian 1/, both sexes	-	-	1	-	2	3	-	-	1	-	1	-
Male	-	-	1	-	1	2	-	-	-	-	1	-
Female	-	-	-	-	1	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Oregon												
All races, both sexes	5	6	8	19	31	60	102	153	136	39	10	-
Male	4	2	3	10	19	27	50	61	52	13	1	-
Female	1	4	5	9	12	33	52	92	84	26	9	-
White, both sexes	3	6	8	18	31	60	101	152	136	38	10	-
Male	2	2	3	10	19	27	50	61	52	12	1	-
Female	1	4	5	8	12	33	51	91	84	26	9	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Pennsylvania												
All races, both sexes	18	30	43	61	92	136	263	410	328	124	16	-
Male	9	22	24	32	43	58	102	129	98	28	2	-
Female	9	8	19	29	49	78	161	281	230	96	14	-
White, both sexes	15	25	39	58	88	133	256	400	323	121	15	-
Male	6	18	22	32	41	56	98	125	97	28	2	-
Female	9	7	17	26	47	77	158	275	226	93	13	-
Black, both sexes	3	5	4	3	2	3	7	10	5	3	1	-
Male	3	4	2	-	-	2	4	4	1	-	-	-
Female	-	1	2	3	2	1	3	6	4	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	-	-	-	-	-	-	-
Male	-	-	-	-	2	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Rhode Island												
All races, both sexes	1	-	6	1	3	12	13	23	21	8	1	-
Male	1	-	5	-	2	4	4	6	7	1	-	-
Female	-	-	1	1	1	8	9	17	14	7	1	-
White, both sexes	1	-	6	1	3	12	12	23	21	8	1	-
Male	1	-	5	-	2	4	4	6	7	1	-	-
Female	-	-	1	1	1	8	8	17	14	7	1	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	10	6	10	22	23	43	52	75	62	15	1	-
Male	6	5	4	12	7	21	20	25	14	2	-	-
Female	4	1	6	10	16	22	32	50	48	13	1	-
White, both sexes	5	5	8	16	14	35	47	67	48	13	1	-
Male	3	4	4	8	5	17	19	24	13	2	-	-
Female	2	1	4	8	9	18	28	43	35	11	1	-
Black, both sexes	5	1	2	6	9	8	5	8	14	2	-	-
Male	3	1	-	4	2	4	1	1	1	-	-	-
Female	2	-	2	2	7	4	4	7	13	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
South Dakota												
All races, both sexes	1	1	2	-	6	7	12	15	17	10	2	-
Male	1	1	1	-	4	4	4	4	5	1	1	-
Female	-	-	1	-	2	3	8	11	12	9	1	-
White, both sexes	1	1	1	-	6	7	12	15	17	10	2	-
Male	1	1	1	-	4	4	4	4	5	1	1	-
Female	-	-	-	-	2	3	8	11	12	9	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	9	7	22	29	29	52	83	79	78	19	2	-
Male	7	5	12	17	14	21	41	23	25	3	-	-
Female	2	2	10	12	15	31	42	56	53	16	2	-
White, both sexes	6	4	17	27	25	48	76	76	72	19	2	-
Male	4	2	9	17	13	19	37	23	24	3	-	-
Female	2	2	8	10	12	29	39	53	48	16	2	-
Black, both sexes	3	3	5	2	4	4	7	3	6	-	-	-
Male	3	3	3	-	1	2	4	-	1	-	-	-
Female	-	-	2	2	3	2	3	3	5	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Texas												
All races, both sexes	31	54	44	66	94	154	181	247	166	63	6	-
Male	18	34	30	30	46	72	77	111	51	14	1	-
Female	13	20	14	36	48	82	104	136	115	49	5	-
White, both sexes	25	41	34	56	85	139	174	232	159	57	6	-
Male	13	24	22	24	41	65	74	106	50	14	1	-
Female	12	17	12	32	44	74	100	126	109	43	5	-
Black, both sexes	6	12	10	9	8	14	5	12	4	6	-	-
Male	5	9	8	5	4	6	2	4	-	-	-	-
Female	1	3	2	4	4	8	3	8	4	6	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	1	1	1	3	3	-	-	-
Male	-	1	-	1	1	1	1	1	1	-	-	-
Female	-	-	-	-	-	-	-	2	2	-	-	-
Utah												
All races, both sexes	3	10	1	9	15	16	33	37	17	6	-	-
Male	3	5	1	5	6	4	11	12	5	2	-	-
Female	-	5	-	4	9	12	22	25	12	4	-	-
White, both sexes	3	8	1	9	15	16	33	36	16	6	-	-
Male	3	5	1	5	6	4	11	12	4	2	-	-
Female	-	3	-	4	9	12	22	24	12	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Vermont												
All races, both sexes	1	2	3	2	-	11	12	17	12	7	1	-
Male	1	1	3	2	-	2	5	9	6	2	1	-
Female	-	1	-	-	-	9	7	8	6	5	-	-
White, both sexes	1	2	3	2	-	11	12	17	12	7	1	-
Male	1	1	3	2	-	2	5	9	6	2	1	-
Female	-	1	-	-	-	9	7	8	6	5	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	9	15	12	24	28	50	88	123	110	29	7	-
Male	6	9	8	10	18	22	32	40	29	8	-	-
Female	3	6	4	14	10	28	56	83	81	21	7	-
White, both sexes	6	6	10	19	22	41	84	115	103	28	7	-
Male	4	2	8	8	14	18	30	38	28	7	-	-
Female	2	4	2	11	8	23	54	77	75	21	7	-
Black, both sexes	3	9	2	5	4	8	4	7	5	1	-	-
Male	2	7	-	2	2	4	2	2	-	1	-	-
Female	1	2	2	3	2	4	2	5	5	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	1	-	1	2	-	-	-
Male	-	-	-	-	2	-	-	-	1	-	-	-
Female	-	-	-	-	-	1	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Washington												
All races, both sexes	12	10	23	25	38	59	125	185	188	91	15	-
Male	7	5	7	14	23	26	49	80	61	17	3	-
Female	5	5	16	11	15	33	76	105	127	74	12	-
White, both sexes	12	10	20	23	37	55	118	182	184	91	15	-
Male	7	5	7	13	23	25	48	79	61	17	3	-
Female	5	5	13	10	14	30	70	103	123	74	12	-
Black, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	2	1	3	5	2	4	-	-	-
Male	-	-	-	1	-	1	1	1	-	-	-	-
Female	-	-	1	1	1	2	4	1	4	-	-	-
West Virginia												
All races, both sexes	3	2	4	9	13	17	26	36	20	4	2	-
Male	2	-	1	6	9	5	11	11	7	-	-	-
Female	1	2	3	3	4	12	15	25	13	4	2	-
White, both sexes	3	2	3	9	13	17	25	35	20	4	2	-
Male	2	-	1	6	9	5	11	11	7	-	-	-
Female	1	2	2	3	4	12	14	24	13	4	2	-
Black, both sexes	-	-	1	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Wisconsin												
All races, both sexes	9	6	10	13	29	43	97	175	134	57	7	-
Male	6	4	6	7	11	18	39	59	35	6	-	-
Female	3	2	4	6	18	25	58	116	99	51	7	-
White, both sexes	8	5	9	12	29	43	94	174	132	57	7	-
Male	5	3	6	7	11	18	38	58	34	6	-	-
Female	3	2	3	5	18	25	56	116	98	51	7	-
Black, both sexes	1	-	1	-	-	-	3	1	1	-	-	-
Male	1	-	-	-	-	-	1	1	1	-	-	-
Female	-	-	1	-	-	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	1	1	2	3	6	8	7	9	3	-	-
Male	-	-	1	1	2	2	4	2	5	1	-	-
Female	1	1	-	1	1	4	4	5	4	2	-	-
White, both sexes	1	1	1	2	3	6	8	7	9	3	-	-
Male	-	-	1	1	2	2	4	2	5	1	-	-
Female	1	1	-	1	1	4	4	5	4	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
United States												
All races, both sexes	2,644	-	2	-	-	6	11	14	18	34	32	66
Male	949	-	2	-	-	4	3	3	6	19	15	30
Female	1,695	-	-	-	-	2	8	11	12	15	17	36
White, both sexes	2,383	-	1	-	-	5	10	12	15	24	26	53
Male	859	-	1	-	-	3	3	3	4	12	12	26
Female	1,524	-	-	-	-	2	7	9	11	12	14	27
Black, both sexes	184	-	1	-	-	1	1	2	2	9	5	11
Male	67	-	1	-	-	1	-	-	2	6	3	3
Female	117	-	-	-	-	-	1	2	-	3	2	8
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	67	-	-	-	-	-	-	-	-	1	1	2
Male	21	-	-	-	-	-	-	-	-	1	-	1
Female	46	-	-	-	-	-	-	-	-	-	1	1
Alabama												
All races, both sexes	33	-	-	-	-	-	-	1	2	-	-	1
Male	14	-	-	-	-	-	-	-	2	-	-	1
Female	19	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	28	-	-	-	-	-	-	-	1	-	-	-
Male	12	-	-	-	-	-	-	-	1	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	1	1	-	-	1
Male	2	-	-	-	-	-	-	-	1	-	-	1
Female	3	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Alaska												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	41	-	-	-	-	-	-	-	-	-	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	27	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	38	-	-	-	-	-	-	-	-	-	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	24	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Arkansas												
All races, both sexes	21	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	285	-	-	-	-	1	2	2	3	3	5	5
Male	97	-	-	-	-	1	-	1	1	1	1	2
Female	188	-	-	-	-	-	2	1	2	2	4	3
White, both sexes	239	-	-	-	-	1	1	2	3	2	4	2
Male	85	-	-	-	-	1	-	1	1	-	1	1
Female	154	-	-	-	-	-	1	1	2	2	3	1
Black, both sexes	17	-	-	-	-	-	1	-	-	1	-	1
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	12	-	-	-	-	-	1	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	28	-	-	-	-	-	-	-	-	-	1	2
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	21	-	-	-	-	-	-	-	-	-	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Colorado												
All races, both sexes	34	-	-	-	-	-	-	-	1	-	1	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	1	-	1	-
White, both sexes	31	-	-	-	-	-	-	-	1	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	46	-	-	-	-	-	1	1	1	-	-	3
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	31	-	-	-	-	-	1	1	1	-	-	3
White, both sexes	43	-	-	-	-	-	1	1	1	-	-	2
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	-	-	1	1	1	-	-	2
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Delaware												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	6	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Florida												
All races, both sexes	201	-	-	-	-	-	1	1	1	2	-	7
Male	96	-	-	-	-	-	1	-	-	1	-	3
Female	105	-	-	-	-	-	-	1	1	1	-	4
White, both sexes	186	-	-	-	-	-	1	1	1	1	-	4
Male	91	-	-	-	-	-	1	-	-	1	-	3
Female	95	-	-	-	-	-	-	1	1	-	-	1
Black, both sexes	13	-	-	-	-	-	-	-	-	1	-	3
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	1	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	52	-	-	-	-	-	-	1	1	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	41	-	-	-	-	-	-	1	1	-	-	1
White, both sexes	40	-	-	-	-	-	-	-	1	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	30	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	11	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Hawaii												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Illinois												
All races, both sexes	82	-	2	-	-	-	1	-	1	3	1	2
Male	30	-	2	-	-	-	1	-	1	3	-	-
Female	52	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	70	-	1	-	-	-	1	-	1	2	1	1
Male	23	-	1	-	-	-	1	-	1	2	-	-
Female	47	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	10	-	1	-	-	-	-	-	-	1	-	1
Male	5	-	1	-	-	-	-	-	-	1	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	63	-	-	-	-	-	-	1	-	1	-	3
Male	19	-	-	-	-	-	-	-	-	1	-	3
Female	44	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	59	-	-	-	-	-	-	1	-	-	-	2
Male	16	-	-	-	-	-	-	-	-	-	-	2
Female	43	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Iowa												
All races, both sexes	46	-	-	-	-	-	-	-	-	-	1	-
Male	19	-	-	-	-	-	-	-	-	-	1	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	46	-	-	-	-	-	-	-	-	-	1	-
Male	19	-	-	-	-	-	-	-	-	-	1	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Kentucky												
All races, both sexes	32	-	-	-	-	-	-	-	1	-	-	2
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	19	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	29	-	-	-	-	-	-	-	1	-	-	2
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	17	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	31	-	-	-	-	-	-	-	-	-	-	2
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	27	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Maine												
All races, both sexes	29	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	29	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	35	-	-	-	-	-	-	-	-	-	-	2
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	20	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	28	-	-	-	-	-	-	-	-	-	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Massachusetts												
All races, both sexes	69	-	-	-	-	2	1	1	-	2	4	-
Male	22	-	-	-	-	1	-	-	-	1	4	-
Female	47	-	-	-	-	1	1	1	-	1	-	-
White, both sexes	67	-	-	-	-	2	1	1	-	2	3	-
Male	21	-	-	-	-	1	-	-	-	1	3	-
Female	46	-	-	-	-	1	1	1	-	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	76	-	-	-	-	1	-	1	-	3	-	4
Male	21	-	-	-	-	-	-	1	-	2	-	2
Female	55	-	-	-	-	1	-	-	-	1	-	2
White, both sexes	69	-	-	-	-	1	-	1	-	2	-	4
Male	19	-	-	-	-	-	-	1	-	2	-	2
Female	50	-	-	-	-	1	-	-	-	-	-	2
Black, both sexes	7	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Minnesota												
All races, both sexes	65	-	-	-	-	-	-	-	1	-	-	3
Male	27	-	-	-	-	-	-	-	-	-	-	2
Female	38	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	64	-	-	-	-	-	-	-	1	-	-	3
Male	27	-	-	-	-	-	-	-	-	-	-	2
Female	37	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	15	-	-	-	-	1	-	-	-	-	-	2
Male	8	-	-	-	-	1	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	6	-	-	-	-	1	-	-	-	-	-	-
Male	3	-	-	-	-	1	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Missouri												
All races, both sexes	73	-	-	-	-	-	-	-	1	1	2	1
Male	27	-	-	-	-	-	-	-	-	-	1	1
Female	46	-	-	-	-	-	-	-	1	1	1	-
White, both sexes	67	-	-	-	-	-	-	-	-	1	2	1
Male	26	-	-	-	-	-	-	-	-	-	1	1
Female	41	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	11	-	-	-	-	-	-	1	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	1	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Nebraska												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	8	-	-	-	-	-	-	1	-	-	1	-
Male	3	-	-	-	-	-	-	1	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	7	-	-	-	-	-	-	1	-	-	1	-
Male	3	-	-	-	-	-	-	1	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
New Hampshire												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	78	-	-	-	-	-	1	-	-	2	1	3
Male	24	-	-	-	-	-	-	-	-	-	-	1
Female	54	-	-	-	-	-	1	-	-	2	1	2
White, both sexes	72	-	-	-	-	-	1	-	-	2	-	3
Male	21	-	-	-	-	-	-	-	-	-	-	1
Female	51	-	-	-	-	-	1	-	-	2	-	2
Black, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
New Mexico												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	132	-	-	-	-	-	1	1	-	-	3	2
Male	49	-	-	-	-	-	1	-	-	-	2	1
Female	83	-	-	-	-	-	-	1	-	-	1	1
White, both sexes	122	-	-	-	-	-	1	1	-	-	1	2
Male	45	-	-	-	-	-	1	-	-	-	-	1
Female	77	-	-	-	-	-	-	1	-	-	1	1
Black, both sexes	6	-	-	-	-	-	-	-	-	-	2	-
Male	3	-	-	-	-	-	-	-	-	-	2	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
North Carolina												
All races, both sexes	77	-	-	-	-	-	-	-	-	2	-	-
Male	28	-	-	-	-	-	-	-	-	2	-	-
Female	49	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	65	-	-	-	-	-	-	-	-	1	-	-
Male	24	-	-	-	-	-	-	-	-	1	-	-
Female	41	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	11	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Ohio												
All races, both sexes	150	-	-	-	-	-	1	-	1	-	3	5
Male	49	-	-	-	-	-	-	-	-	-	2	3
Female	101	-	-	-	-	-	1	-	1	-	1	2
White, both sexes	137	-	-	-	-	-	1	-	1	-	3	4
Male	43	-	-	-	-	-	-	-	-	-	2	2
Female	94	-	-	-	-	-	1	-	1	-	1	2
Black, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	22	-	-	-	-	-	-	-	-	1	2	-
Male	10	-	-	-	-	-	-	-	-	1	1	-
Female	12	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	16	-	-	-	-	-	-	-	-	1	2	-
Male	8	-	-	-	-	-	-	-	-	1	1	-
Female	8	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Oregon												
All races, both sexes	75	-	-	-	-	-	-	-	-	1	-	1
Male	33	-	-	-	-	-	-	-	-	-	-	1
Female	42	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	74	-	-	-	-	-	-	-	-	1	-	1
Male	33	-	-	-	-	-	-	-	-	-	-	1
Female	41	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	190	-	-	-	-	-	-	1	1	3	4	5
Male	64	-	-	-	-	-	-	-	1	-	1	2
Female	126	-	-	-	-	-	-	1	-	3	3	3
White, both sexes	182	-	-	-	-	-	-	1	-	2	4	5
Male	60	-	-	-	-	-	-	-	-	-	1	2
Female	122	-	-	-	-	-	-	1	-	2	3	3
Black, both sexes	8	-	-	-	-	-	-	-	1	1	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	4	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Rhode Island												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	38	-	-	-	-	-	-	-	-	1	1	-
Male	13	-	-	-	-	-	-	-	-	1	1	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	35	-	-	-	-	-	-	-	-	1	1	-
Male	13	-	-	-	-	-	-	-	-	1	1	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
South Dakota												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	42	-	-	-	-	-	-	-	-	1	1	-
Male	15	-	-	-	-	-	-	-	-	1	1	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	35	-	-	-	-	-	-	-	-	1	1	-
Male	11	-	-	-	-	-	-	-	-	1	1	-
Female	24	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Texas												
All races, both sexes	121	-	-	-	-	-	1	-	1	3	-	5
Male	53	-	-	-	-	-	-	-	1	3	-	2
Female	68	-	-	-	-	-	1	-	-	-	-	3
White, both sexes	109	-	-	-	-	-	1	-	1	1	-	5
Male	45	-	-	-	-	-	-	-	1	1	-	2
Female	64	-	-	-	-	-	1	-	-	-	-	3
Black, both sexes	10	-	-	-	-	-	-	-	-	2	-	-
Male	7	-	-	-	-	-	-	-	-	2	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Vermont												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	52	-	-	-	-	-	1	1	-	1	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	35	-	-	-	-	-	1	1	-	1	-	-
White, both sexes	43	-	-	-	-	-	1	1	-	1	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	1	1	-	1	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Washington												
All races, both sexes	83	-	-	-	-	1	-	-	1	1	-	2
Male	22	-	-	-	-	1	-	-	-	1	-	-
Female	61	-	-	-	-	-	-	-	1	-	-	2
White, both sexes	78	-	-	-	-	1	-	-	1	1	-	2
Male	21	-	-	-	-	1	-	-	-	1	-	-
Female	57	-	-	-	-	-	-	-	1	-	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	11	-	-	-	-	-	-	-	1	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	10	-	-	-	-	-	-	-	1	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Wisconsin												
All races, both sexes	61	-	-	-	-	-	-	-	-	1	-	2
Male	22	-	-	-	-	-	-	-	-	-	-	2
Female	39	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	61	-	-	-	-	-	-	-	-	1	-	2
Male	22	-	-	-	-	-	-	-	-	-	-	2
Female	39	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
United States												
All races, both sexes	61	78	99	133	194	299	485	553	400	140	19	-
Male	32	41	39	58	73	112	193	175	118	24	2	-
Female	29	37	60	75	121	187	292	378	282	116	17	-
White, both sexes	43	57	80	113	170	272	449	519	381	136	17	-
Male	21	29	33	51	67	105	179	169	115	24	2	-
Female	22	28	47	62	103	167	270	350	266	112	15	-
Black, both sexes	18	18	13	14	19	17	19	15	14	3	2	-
Male	11	11	3	5	4	4	9	2	2	-	-	-
Female	7	7	10	9	15	13	10	13	12	3	2	-
American Indian 1/, both sexes	-	1	1	1	-	-	3	3	-	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	1	-	-	3	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	5	5	5	10	14	16	5	1	-	-
Male	-	-	3	2	2	3	5	3	1	-	-	-
Female	-	2	2	3	3	7	9	13	4	1	-	-
Alabama												
All races, both sexes	1	1	1	1	5	4	4	9	2	1	-	-
Male	-	-	1	-	2	3	2	3	-	-	-	-
Female	1	1	-	1	3	1	2	6	2	1	-	-
White, both sexes	1	1	1	1	4	4	4	8	2	1	-	-
Male	-	-	1	-	2	3	2	3	-	-	-	-
Female	1	1	-	1	2	1	2	5	2	1	-	-
Black, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Alaska												
All races, both sexes	-	-	1	-	2	1	-	1	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	1	-	1	1	-	-	-
White, both sexes	-	-	1	-	1	1	-	1	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1	2	1	5	2	3	8	10	6	2	-	-
Male	1	1	1	2	-	2	1	4	1	-	-	-
Female	-	1	-	3	2	1	7	6	5	2	-	-
White, both sexes	1	2	1	4	2	3	7	10	5	2	-	-
Male	1	1	1	2	-	2	1	4	1	-	-	-
Female	-	1	-	2	2	1	6	6	4	2	-	-
Black, both sexes	-	-	-	1	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Arkansas												
All races, both sexes	-	-	1	-	3	4	6	1	3	2	-	-
Male	-	-	-	-	-	1	2	-	-	-	-	-
Female	-	-	1	-	3	3	4	1	3	2	-	-
White, both sexes	-	-	-	-	3	3	5	1	2	2	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	3	2	4	1	2	2	-	-
Black, both sexes	-	-	-	-	-	1	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	5	10	9	17	13	26	50	64	49	16	5	-
Male	4	5	5	8	7	8	14	19	17	2	1	-
Female	1	5	4	9	6	18	36	45	32	14	4	-
White, both sexes	3	8	8	13	10	21	42	52	47	15	5	-
Male	2	5	4	7	5	8	11	18	17	2	1	-
Female	1	3	4	6	5	13	31	34	30	13	4	-
Black, both sexes	2	1	-	2	1	3	2	2	-	1	-	-
Male	2	-	-	1	1	-	-	-	-	-	-	-
Female	-	1	-	1	-	3	2	2	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	2	2	2	6	9	2	-	-	-
Male	-	-	1	-	1	-	3	1	-	-	-	-
Female	-	1	-	2	1	2	3	8	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Colorado												
All races, both sexes	-	-	2	2	5	4	7	5	4	3	-	-
Male	-	-	1	2	3	2	2	3	-	1	-	-
Female	-	-	1	-	2	2	5	2	4	2	-	-
White, both sexes	-	-	2	2	4	4	6	5	4	3	-	-
Male	-	-	1	2	3	2	2	3	-	1	-	-
Female	-	-	1	-	1	2	4	2	4	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Connecticut												
All races, both sexes	-	2	1	-	3	2	11	9	11	-	1	-
Male	-	1	1	-	1	-	7	3	2	-	-	-
Female	-	1	-	-	2	2	4	6	9	-	1	-
White, both sexes	-	1	1	-	3	2	10	9	11	-	1	-
Male	-	1	1	-	1	-	7	3	2	-	-	-
Female	-	-	-	-	2	2	3	6	9	-	1	-
Black, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Delaware												
All races, both sexes	-	-	-	-	1	1	1	3	2	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	3	2	-	-	-
White, both sexes	-	-	-	-	1	1	1	3	2	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	3	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	2	-	-	-	1	-	-	1	1	-	-
Male	-	2	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	2	-	-	-	1	-	-	-	1	-	-
Male	-	2	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Florida												
All races, both sexes	6	5	10	10	18	26	37	40	28	9	-	-
Male	3	4	2	6	8	14	21	19	12	2	-	-
Female	3	1	8	4	10	12	16	21	16	7	-	-
White, both sexes	4	3	6	9	18	24	37	40	28	9	-	-
Male	2	2	1	5	8	14	21	19	12	2	-	-
Female	2	1	5	4	10	10	16	21	16	7	-	-
Black, both sexes	2	2	3	1	-	1	-	-	-	-	-	-
Male	1	2	-	1	-	-	-	-	-	-	-	-
Female	1	-	3	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Georgia												
All races, both sexes	4	1	3	5	7	6	6	13	4	-	-	-
Male	2	1	1	2	-	2	1	2	-	-	-	-
Female	2	-	2	3	7	4	5	11	4	-	-	-
White, both sexes	2	1	2	3	3	5	6	12	4	-	-	-
Male	1	1	1	2	-	2	1	2	-	-	-	-
Female	1	-	1	1	3	3	5	10	4	-	-	-
Black, both sexes	2	-	1	2	4	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	2	4	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Hawaii												
All races, both sexes	-	-	1	2	-	3	3	4	2	-	-	-
Male	-	-	-	1	-	2	2	2	1	-	-	-
Female	-	-	1	1	-	1	1	2	1	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	2	-	3	2	4	1	-	-	-
Male	-	-	-	1	-	2	1	2	-	-	-	-
Female	-	-	1	1	-	1	1	2	1	-	-	-
Idaho												
All races, both sexes	-	-	1	1	-	1	4	2	3	1	-	-
Male	-	-	-	1	-	-	1	-	2	-	-	-
Female	-	-	1	-	-	1	3	2	1	1	-	-
White, both sexes	-	-	1	1	-	1	4	2	3	1	-	-
Male	-	-	-	1	-	-	1	-	2	-	-	-
Female	-	-	1	-	-	1	3	2	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Illinois												
All races, both sexes	2	3	2	5	4	8	16	12	11	8	1	-
Male	1	3	1	2	1	2	7	4	1	1	-	-
Female	1	-	1	3	3	6	9	8	10	7	1	-
White, both sexes	2	2	1	5	1	8	15	10	11	8	-	-
Male	1	2	-	2	-	2	6	3	1	1	-	-
Female	1	-	1	3	1	6	9	7	10	7	-	-
Black, both sexes	-	1	-	-	3	-	-	2	-	-	1	-
Male	-	1	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	2	-	-	1	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	2	2	3	2	8	6	12	10	8	3	2	-
Male	-	1	1	1	-	2	2	3	4	1	-	-
Female	2	1	2	1	8	4	10	7	4	2	2	-
White, both sexes	2	1	3	2	8	6	11	10	8	3	2	-
Male	-	-	1	1	-	2	2	3	4	1	-	-
Female	2	1	2	1	8	4	9	7	4	2	2	-
Black, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Iowa												
All races, both sexes	-	-	1	3	1	2	11	11	15	1	-	-
Male	-	-	-	-	-	2	7	3	6	-	-	-
Female	-	-	1	3	1	-	4	8	9	1	-	-
White, both sexes	-	-	1	3	1	2	11	11	15	1	-	-
Male	-	-	-	-	-	2	7	3	6	-	-	-
Female	-	-	1	3	1	-	4	8	9	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	1	1	1	-	2	5	1	4	1	-	-
Male	-	-	1	-	-	2	2	1	-	-	-	-
Female	-	1	-	1	-	-	3	-	4	1	-	-
White, both sexes	-	-	1	1	-	2	5	1	4	1	-	-
Male	-	-	1	-	-	2	2	1	-	-	-	-
Female	-	-	-	1	-	-	3	-	4	1	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Kentucky												
All races, both sexes	-	1	2	2	5	4	8	6	-	1	-	-
Male	-	-	2	-	2	2	4	1	-	1	-	-
Female	-	1	-	2	3	2	4	5	-	-	-	-
White, both sexes	-	1	2	1	5	3	7	6	-	1	-	-
Male	-	-	2	-	2	2	3	1	-	1	-	-
Female	-	1	-	1	3	1	4	5	-	-	-	-
Black, both sexes	-	-	-	1	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	3	-	1	3	1	3	5	6	5	2	-	-
Male	2	-	-	1	-	2	2	1	1	-	-	-
Female	1	-	1	2	1	1	3	5	4	2	-	-
White, both sexes	3	-	1	1	1	3	5	6	4	2	-	-
Male	2	-	-	-	-	2	2	1	1	-	-	-
Female	1	-	1	1	1	1	3	5	3	2	-	-
Black, both sexes	-	-	-	1	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Maine												
All races, both sexes	2	-	3	1	1	5	6	4	5	2	-	-
Male	1	-	1	-	-	3	2	-	2	1	-	-
Female	1	-	2	1	1	2	4	4	3	1	-	-
White, both sexes	2	-	3	1	1	5	6	4	5	2	-	-
Male	1	-	1	-	-	3	2	-	2	1	-	-
Female	1	-	2	1	1	2	4	4	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	2	1	-	4	2	6	8	6	4	-	-
Male	-	1	-	-	2	-	1	6	2	2	-	-
Female	-	1	1	-	2	2	5	2	4	2	-	-
White, both sexes	-	1	1	-	2	2	5	7	5	4	-	-
Male	-	1	-	-	1	-	1	6	2	2	-	-
Female	-	-	1	-	1	2	4	1	3	2	-	-
Black, both sexes	-	1	-	-	2	-	-	1	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Massachusetts												
All races, both sexes	3	1	2	3	3	7	7	12	13	6	2	-
Male	1	-	-	1	2	2	6	2	1	1	-	-
Female	2	1	2	2	1	5	1	10	12	5	2	-
White, both sexes	3	1	2	3	3	7	7	12	13	5	2	-
Male	1	-	-	1	2	2	6	2	1	1	-	-
Female	2	1	2	2	1	5	1	10	12	4	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Michigan												
All races, both sexes	3	2	2	2	7	7	15	12	11	6	-	-
Male	3	-	1	-	1	-	3	7	1	-	-	-
Female	-	2	1	2	6	7	12	5	10	6	-	-
White, both sexes	1	1	1	2	7	7	14	12	10	6	-	-
Male	1	-	1	-	1	-	3	7	1	-	-	-
Female	-	1	-	2	6	7	11	5	9	6	-	-
Black, both sexes	2	1	1	-	-	-	1	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Minnesota												
All races, both sexes	3	-	2	2	6	8	8	13	11	6	2	-
Male	2	-	-	-	2	6	6	5	2	2	-	-
Female	1	-	2	2	4	2	2	8	9	4	2	-
White, both sexes	3	-	2	2	6	8	8	12	11	6	2	-
Male	2	-	-	-	2	6	6	5	2	2	-	-
Female	1	-	2	2	4	2	2	7	9	4	2	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	1	3	-	-	2	1	4	1	-	-	-
Male	-	1	2	-	-	2	-	2	-	-	-	-
Female	-	-	1	-	-	-	1	2	1	-	-	-
White, both sexes	-	-	1	-	-	2	1	3	-	-	-	-
Male	-	-	1	-	-	2	-	2	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	1	2	-	-	-	-	1	1	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Missouri												
All races, both sexes	1	3	3	4	4	10	14	16	10	3	-	-
Male	1	2	1	2	2	5	5	4	2	1	-	-
Female	-	1	2	2	2	5	9	12	8	2	-	-
White, both sexes	-	3	3	4	4	9	14	14	9	3	-	-
Male	-	2	1	2	2	5	5	4	2	1	-	-
Female	-	1	2	2	2	4	9	10	7	2	-	-
Black, both sexes	1	-	-	-	-	1	-	2	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	-	-	-	-	1	2	3	2	1	-	-
Male	-	-	-	-	-	1	-	1	1	-	-	-
Female	1	-	-	-	-	-	2	2	1	1	-	-
White, both sexes	1	-	-	-	-	1	2	3	2	1	-	-
Male	-	-	-	-	-	1	-	1	1	-	-	-
Female	1	-	-	-	-	-	2	2	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Nebraska												
All races, both sexes	1	-	-	1	-	1	1	3	1	3	-	-
Male	1	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	1	2	1	3	-	-
White, both sexes	1	-	-	1	-	1	1	3	1	3	-	-
Male	1	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	1	2	1	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	1	-	3	-	-	1	-	1	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	1	-	2	-	-	-	-	1	-	-
White, both sexes	-	-	1	-	3	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	1	-	2	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
New Hampshire												
All races, both sexes	-	-	1	1	-	1	3	5	2	1	-	-
Male	-	-	-	1	-	-	-	3	-	1	-	-
Female	-	-	1	-	-	1	3	2	2	-	-	-
White, both sexes	-	-	1	1	-	1	3	5	2	1	-	-
Male	-	-	-	1	-	-	-	3	-	1	-	-
Female	-	-	1	-	-	1	3	2	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1	2	2	2	6	9	14	22	12	1	-	-
Male	1	1	1	-	2	2	8	3	5	-	-	-
Female	-	1	1	2	4	7	6	19	7	1	-	-
White, both sexes	1	1	1	2	6	8	13	21	12	1	-	-
Male	1	-	1	-	2	1	7	3	5	-	-	-
Female	-	1	-	2	4	7	6	18	7	1	-	-
Black, both sexes	-	1	-	-	-	1	1	-	-	-	-	-
Male	-	1	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
New Mexico												
All races, both sexes	2	1	1	1	2	4	1	2	-	2	-	-
Male	1	1	-	-	-	3	1	1	-	1	-	-
Female	1	-	1	1	2	1	-	1	-	1	-	-
White, both sexes	2	-	1	-	2	4	1	2	-	2	-	-
Male	1	-	-	-	-	3	1	1	-	1	-	-
Female	1	-	1	-	2	1	-	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	3	3	6	8	21	27	30	20	7	-	-
Male	-	1	1	5	6	6	10	5	11	-	-	-
Female	-	2	2	1	2	15	17	25	9	7	-	-
White, both sexes	-	3	3	5	7	19	24	29	20	7	-	-
Male	-	1	1	4	5	6	10	5	11	-	-	-
Female	-	2	2	1	2	13	14	24	9	7	-	-
Black, both sexes	-	-	-	1	-	1	2	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
North Carolina												
All races, both sexes	4	2	4	4	7	7	17	15	10	3	2	-
Male	1	1	3	3	1	2	5	7	2	-	1	-
Female	3	1	1	1	6	5	12	8	8	3	1	-
White, both sexes	1	2	4	2	6	7	13	15	10	3	1	-
Male	1	1	3	1	1	2	4	7	2	-	1	-
Female	-	1	1	1	5	5	9	8	8	3	-	-
Black, both sexes	3	-	-	2	1	-	3	-	-	-	1	-
Male	-	-	-	2	-	-	1	-	-	-	-	-
Female	3	-	-	-	1	-	2	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	1	1	2	-	4	1	-	-	-
Male	-	-	-	-	-	2	-	1	-	-	-	-
Female	-	-	-	1	1	-	-	3	1	-	-	-
White, both sexes	-	-	-	1	1	2	-	3	1	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	3	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Ohio												
All races, both sexes	2	6	7	5	12	16	27	29	22	14	-	-
Male	1	2	2	2	5	5	13	6	6	2	-	-
Female	1	4	5	3	7	11	14	23	16	12	-	-
White, both sexes	1	5	5	3	12	15	25	28	20	14	-	-
Male	1	2	1	2	5	5	11	5	5	2	-	-
Female	-	3	4	1	7	10	14	23	15	12	-	-
Black, both sexes	1	1	2	2	-	1	2	1	2	-	-	-
Male	-	-	1	-	-	-	2	1	1	-	-	-
Female	1	1	1	2	-	1	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1	1	1	-	2	2	3	3	6	-	-	-
Male	1	-	1	-	-	-	3	1	2	-	-	-
Female	-	1	-	-	2	2	-	2	4	-	-	-
White, both sexes	-	-	1	-	-	1	3	3	5	-	-	-
Male	-	-	1	-	-	-	3	1	1	-	-	-
Female	-	-	-	-	-	1	-	2	4	-	-	-
Black, both sexes	1	1	-	-	2	1	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	1	-	-	2	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Oregon												
All races, both sexes	1	-	-	3	2	6	21	18	16	5	1	-
Male	-	-	-	2	1	1	10	9	9	-	-	-
Female	1	-	-	1	1	5	11	9	7	5	1	-
White, both sexes	1	-	-	3	2	6	20	18	16	5	1	-
Male	-	-	-	2	1	1	10	9	9	-	-	-
Female	1	-	-	1	1	5	10	9	7	5	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Pennsylvania												
All races, both sexes	3	5	7	16	16	28	29	47	19	5	1	-
Male	1	5	3	9	8	9	8	11	5	1	-	-
Female	2	-	4	7	8	19	21	36	14	4	1	-
White, both sexes	2	4	7	15	15	26	29	47	19	5	1	-
Male	-	4	3	9	8	8	8	11	5	1	-	-
Female	2	-	4	6	7	18	21	36	14	4	1	-
Black, both sexes	1	1	-	1	1	2	-	-	-	-	-	-
Male	1	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Rhode Island												
All races, both sexes	1	-	1	1	-	3	1	3	4	1	-	-
Male	1	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	1	1	-	1	1	3	4	1	-	-
White, both sexes	1	-	1	1	-	3	1	3	4	1	-	-
Male	1	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	1	1	-	1	1	3	4	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	-	1	3	4	5	7	10	4	-	1	-
Male	-	-	-	-	2	3	3	3	-	-	-	-
Female	1	-	1	3	2	2	4	7	4	-	1	-
White, both sexes	-	-	-	3	4	5	7	10	3	-	1	-
Male	-	-	-	-	2	3	3	3	-	-	-	-
Female	-	-	-	3	2	2	4	7	3	-	1	-
Black, both sexes	1	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
South Dakota												
All races, both sexes	-	-	1	-	1	2	1	-	1	3	-	-
Male	-	-	1	-	1	-	-	-	-	1	-	-
Female	-	-	-	-	-	2	1	-	1	2	-	-
White, both sexes	-	-	1	-	1	2	1	-	1	3	-	-
Male	-	-	1	-	1	-	-	-	-	1	-	-
Female	-	-	-	-	-	2	1	-	1	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	1	2	2	1	2	6	10	7	9	-	-	-
Male	1	1	2	-	-	1	4	1	3	-	-	-
Female	-	1	-	1	2	5	6	6	6	-	-	-
White, both sexes	-	2	1	1	-	5	9	7	8	-	-	-
Male	-	1	1	-	-	-	3	1	3	-	-	-
Female	-	1	-	1	-	5	6	6	5	-	-	-
Black, both sexes	1	-	1	-	2	1	1	-	1	-	-	-
Male	1	-	1	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	2	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Texas												
All races, both sexes	2	8	2	8	11	13	28	22	13	3	1	-
Male	1	5	1	2	5	5	13	10	4	1	-	-
Female	1	3	1	6	6	8	15	12	9	2	1	-
White, both sexes	1	6	2	8	9	12	26	22	11	3	1	-
Male	-	3	1	2	4	5	12	10	3	1	-	-
Female	1	3	1	6	5	7	14	12	8	2	1	-
Black, both sexes	1	2	-	-	2	1	1	-	1	-	-	-
Male	1	2	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	4	-	-	3	2	3	-	1	1	-	-
Male	-	1	-	-	2	-	1	-	-	-	-	-
Female	-	3	-	-	1	2	2	-	1	1	-	-
White, both sexes	-	3	-	-	3	2	3	-	1	1	-	-
Male	-	1	-	-	2	-	1	-	-	-	-	-
Female	-	2	-	-	1	2	2	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Vermont												
All races, both sexes	-	-	-	-	-	-	1	5	1	-	-	-
Male	-	-	-	-	-	-	1	2	1	-	-	-
Female	-	-	-	-	-	-	-	3	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	5	1	-	-	-
Male	-	-	-	-	-	-	1	2	1	-	-	-
Female	-	-	-	-	-	-	-	3	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	2	1	2	4	6	12	11	9	2	-	-
Male	-	1	-	1	2	1	3	3	5	1	-	-
Female	-	1	1	1	2	5	9	8	4	1	-	-
White, both sexes	-	2	-	2	4	4	10	8	8	2	-	-
Male	-	1	-	1	2	1	2	3	5	1	-	-
Female	-	1	-	1	2	3	8	5	3	1	-	-
Black, both sexes	-	-	1	-	-	1	2	3	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	1	1	3	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Washington												
All races, both sexes	3	2	4	3	-	8	14	20	18	6	-	-
Male	1	-	-	-	-	2	5	6	5	1	-	-
Female	2	2	4	3	-	6	9	14	13	5	-	-
White, both sexes	3	2	4	3	-	7	13	18	17	6	-	-
Male	1	-	-	-	-	1	5	6	5	1	-	-
Female	2	2	4	3	-	6	8	12	12	5	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	1	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	1	-	-	-
West Virginia												
All races, both sexes	-	1	2	-	1	1	1	3	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	1	2	-	-	1	1	3	-	-	-	-
White, both sexes	-	1	1	-	1	1	1	3	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	1	1	-	-	1	1	3	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Wisconsin												
All races, both sexes	1	-	1	4	6	6	10	14	13	3	-	-
Male	-	-	1	3	2	2	4	6	2	-	-	-
Female	1	-	-	1	4	4	6	8	11	3	-	-
White, both sexes	1	-	1	4	6	6	10	14	13	3	-	-
Male	-	-	1	3	2	2	4	6	2	-	-	-
Female	1	-	-	1	4	4	6	8	11	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
United States												
All races, both sexes	15,183	2	1	2	2	2	6	20	22	26	57	93
Male	6,283	-	1	2	1	2	4	14	18	20	39	66
Female	8,900	2	-	-	1	-	2	6	4	6	18	27
White, both sexes	14,295	1	1	2	2	1	5	16	18	18	44	78
Male	5,903	-	1	2	1	1	3	10	14	13	30	56
Female	8,392	1	-	-	1	-	2	6	4	5	14	22
Black, both sexes	682	-	-	-	-	1	1	3	4	8	13	12
Male	287	-	-	-	-	1	1	3	4	7	9	8
Female	395	-	-	-	-	-	-	-	-	1	4	4
American Indian 1/, both sexes	34	-	-	-	-	-	-	1	-	-	-	2
Male	15	-	-	-	-	-	-	1	-	-	-	2
Female	19	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	172	1	-	-	-	-	-	-	-	-	-	1
Male	78	-	-	-	-	-	-	-	-	-	-	-
Female	94	1	-	-	-	-	-	-	-	-	-	1
Alabama												
All races, both sexes	162	-	-	-	-	-	1	-	-	-	-	-
Male	70	-	-	-	-	-	1	-	-	-	-	-
Female	92	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	137	-	-	-	-	-	1	-	-	-	-	-
Male	61	-	-	-	-	-	1	-	-	-	-	-
Female	76	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Alaska												
All races, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	242	1	-	-	-	-	-	-	-	-	2	1
Male	90	-	-	-	-	-	-	-	-	-	1	1
Female	152	1	-	-	-	-	-	-	-	-	1	-
White, both sexes	237	1	-	-	-	-	-	-	-	-	2	1
Male	86	-	-	-	-	-	-	-	-	-	1	1
Female	151	1	-	-	-	-	-	-	-	-	1	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Arkansas												
All races, both sexes	101	-	-	-	-	-	-	-	-	-	-	-
Male	42	-	-	-	-	-	-	-	-	-	-	-
Female	59	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	97	-	-	-	-	-	-	-	-	-	-	-
Male	42	-	-	-	-	-	-	-	-	-	-	-
Female	55	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1,839	-	-	-	-	-	-	3	1	4	7	9
Male	776	-	-	-	-	-	-	2	1	2	6	8
Female	1,063	-	-	-	-	-	-	1	-	2	1	1
White, both sexes	1,710	-	-	-	-	-	-	3	-	4	7	7
Male	717	-	-	-	-	-	-	2	-	2	6	7
Female	993	-	-	-	-	-	-	1	-	2	1	-
Black, both sexes	56	-	-	-	-	-	-	-	1	-	-	-
Male	21	-	-	-	-	-	-	-	1	-	-	-
Female	35	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	68	-	-	-	-	-	-	-	-	-	-	1
Male	35	-	-	-	-	-	-	-	-	-	-	-
Female	33	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Colorado												
All races, both sexes	177	-	-	-	-	-	-	-	-	-	-	-
Male	82	-	-	-	-	-	-	-	-	-	-	-
Female	95	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	174	-	-	-	-	-	-	-	-	-	-	-
Male	82	-	-	-	-	-	-	-	-	-	-	-
Female	92	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	198	-	-	-	-	-	-	-	1	-	-	-
Male	82	-	-	-	-	-	-	-	1	-	-	-
Female	116	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	191	-	-	-	-	-	-	-	1	-	-	-
Male	79	-	-	-	-	-	-	-	1	-	-	-
Female	112	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Delaware												
All races, both sexes	56	-	-	-	-	-	-	-	-	-	1	-
Male	23	-	-	-	-	-	-	-	-	-	1	-
Female	33	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	51	-	-	-	-	-	-	-	-	-	1	-
Male	19	-	-	-	-	-	-	-	-	-	1	-
Female	32	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Florida												
All races, both sexes	990	-	-	-	-	-	-	1	2	2	2	3
Male	471	-	-	-	-	-	-	1	2	2	1	2
Female	519	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	937	-	-	-	-	-	-	1	-	2	1	2
Male	446	-	-	-	-	-	-	1	-	2	1	1
Female	491	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	49	-	-	-	-	-	-	-	2	-	1	1
Male	22	-	-	-	-	-	-	-	2	-	-	1
Female	27	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	262	-	-	-	1	-	1	1	-	-	3	2
Male	114	-	-	-	-	-	1	-	-	-	1	1
Female	148	-	-	-	1	-	-	1	-	-	2	1
White, both sexes	217	-	-	-	1	-	-	1	-	-	3	1
Male	95	-	-	-	-	-	-	-	-	-	1	1
Female	122	-	-	-	1	-	-	1	-	-	2	-
Black, both sexes	44	-	-	-	-	-	1	-	-	-	-	1
Male	19	-	-	-	-	-	1	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Hawaii												
All races, both sexes	63	-	-	-	-	-	-	-	-	-	-	-
Male	26	-	-	-	-	-	-	-	-	-	-	-
Female	37	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	44	-	-	-	-	-	-	-	-	-	-	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	26	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	85	-	-	-	-	-	-	-	-	-	-	-
Male	44	-	-	-	-	-	-	-	-	-	-	-
Female	41	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	85	-	-	-	-	-	-	-	-	-	-	-
Male	44	-	-	-	-	-	-	-	-	-	-	-
Female	41	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Illinois												
All races, both sexes	543	-	-	-	-	-	-	-	-	-	1	3
Male	218	-	-	-	-	-	-	-	-	-	-	2
Female	325	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	509	-	-	-	-	-	-	-	-	-	-	3
Male	202	-	-	-	-	-	-	-	-	-	-	2
Female	307	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	32	-	-	-	-	-	-	-	-	-	1	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	325	-	-	-	-	-	-	-	1	1	1	2
Male	152	-	-	-	-	-	-	-	1	1	1	2
Female	173	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	315	-	-	-	-	-	-	-	1	1	-	2
Male	147	-	-	-	-	-	-	-	1	1	-	2
Female	168	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	10	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Iowa												
All races, both sexes	242	-	-	-	-	-	-	-	1	-	1	-
Male	95	-	-	-	-	-	-	-	1	-	-	-
Female	147	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	240	-	-	-	-	-	-	-	1	-	1	-
Male	95	-	-	-	-	-	-	-	1	-	-	-
Female	145	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	174	-	-	-	-	-	-	-	1	1	1	2
Male	65	-	-	-	-	-	-	-	-	1	1	1
Female	109	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	167	-	-	-	-	-	-	-	1	1	1	2
Male	64	-	-	-	-	-	-	-	-	1	1	1
Female	103	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Kentucky												
All races, both sexes	146	-	-	-	-	-	-	-	-	-	1	1
Male	64	-	-	-	-	-	-	-	-	-	1	1
Female	82	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	143	-	-	-	-	-	-	-	-	-	1	1
Male	62	-	-	-	-	-	-	-	-	-	1	1
Female	81	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	162	-	-	-	-	1	-	-	-	-	-	3
Male	61	-	-	-	-	1	-	-	-	-	-	3
Female	101	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	134	-	-	-	-	1	-	-	-	-	-	1
Male	51	-	-	-	-	1	-	-	-	-	-	1
Female	83	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	27	-	-	-	-	-	-	-	-	-	-	2
Male	10	-	-	-	-	-	-	-	-	-	-	2
Female	17	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Maine												
All races, both sexes	111	-	-	-	-	-	-	-	1	-	-	-
Male	53	-	-	-	-	-	-	-	1	-	-	-
Female	58	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	111	-	-	-	-	-	-	-	1	-	-	-
Male	53	-	-	-	-	-	-	-	1	-	-	-
Female	58	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	266	-	-	-	-	-	-	-	-	2	1	1
Male	125	-	-	-	-	-	-	-	-	2	-	1
Female	141	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	235	-	-	-	-	-	-	-	-	1	-	1
Male	112	-	-	-	-	-	-	-	-	1	-	1
Female	123	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	29	-	-	-	-	-	-	-	-	1	1	-
Male	12	-	-	-	-	-	-	-	-	1	-	-
Female	17	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Massachusetts												
All races, both sexes	400	-	-	-	-	-	-	-	1	1	-	3
Male	145	-	-	-	-	-	-	-	1	1	-	3
Female	255	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	391	-	-	-	-	-	-	-	1	1	-	3
Male	142	-	-	-	-	-	-	-	1	1	-	3
Female	249	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	460	-	-	-	-	-	1	-	2	4	1	5
Male	211	-	-	-	-	-	1	-	1	4	1	4
Female	249	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	432	-	-	-	-	-	1	-	2	2	1	5
Male	196	-	-	-	-	-	1	-	1	2	1	4
Female	236	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	24	-	-	-	-	-	-	-	-	2	-	-
Male	13	-	-	-	-	-	-	-	-	2	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Minnesota												
All races, both sexes	344	-	-	-	-	-	-	-	-	-	1	1
Male	151	-	-	-	-	-	-	-	-	-	1	-
Female	193	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	339	-	-	-	-	-	-	-	-	-	1	1
Male	151	-	-	-	-	-	-	-	-	-	1	-
Female	188	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	80	-	-	-	1	-	-	-	-	2	2	-
Male	29	-	-	-	1	-	-	-	-	2	-	-
Female	51	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	65	-	-	-	1	-	-	-	-	1	1	-
Male	24	-	-	-	1	-	-	-	-	1	-	-
Female	41	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	13	-	-	-	-	-	-	-	-	1	1	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	9	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Missouri												
All races, both sexes	314	-	1	-	-	-	-	-	1	-	4	4
Male	124	-	1	-	-	-	-	-	1	-	3	2
Female	190	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	302	-	1	-	-	-	-	-	1	-	4	3
Male	121	-	1	-	-	-	-	-	1	-	3	2
Female	181	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	71	-	-	-	-	-	-	-	-	-	-	1
Male	34	-	-	-	-	-	-	-	-	-	-	1
Female	37	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	70	-	-	-	-	-	-	-	-	-	-	-
Male	33	-	-	-	-	-	-	-	-	-	-	-
Female	37	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Nebraska												
All races, both sexes	138	-	-	-	-	-	1	-	-	-	-	4
Male	51	-	-	-	-	-	-	-	-	-	-	3
Female	87	-	-	-	-	-	1	-	-	-	-	1
White, both sexes	136	-	-	-	-	-	1	-	-	-	-	4
Male	50	-	-	-	-	-	-	-	-	-	-	3
Female	86	-	-	-	-	-	1	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	86	-	-	-	-	-	-	-	-	-	2	1
Male	41	-	-	-	-	-	-	-	-	-	2	1
Female	45	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	77	-	-	-	-	-	-	-	-	-	1	-
Male	35	-	-	-	-	-	-	-	-	-	1	-
Female	42	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	1	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
New Hampshire												
All races, both sexes	85	-	-	-	-	-	-	-	-	-	-	-
Male	32	-	-	-	-	-	-	-	-	-	-	-
Female	53	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	85	-	-	-	-	-	-	-	-	-	-	-
Male	32	-	-	-	-	-	-	-	-	-	-	-
Female	53	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	551	-	-	-	-	-	-	2	1	1	2	7
Male	232	-	-	-	-	-	-	2	1	-	2	5
Female	319	-	-	-	-	-	-	-	-	1	-	2
White, both sexes	519	-	-	-	-	-	-	1	-	1	1	4
Male	214	-	-	-	-	-	-	1	-	-	1	2
Female	305	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	28	-	-	-	-	-	-	1	1	-	1	3
Male	15	-	-	-	-	-	-	1	1	-	1	3
Female	13	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
New Mexico												
All races, both sexes	72	-	-	-	-	-	-	1	-	-	-	-
Male	35	-	-	-	-	-	-	1	-	-	-	-
Female	37	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	68	-	-	-	-	-	-	1	-	-	-	-
Male	33	-	-	-	-	-	-	1	-	-	-	-
Female	35	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	842	-	-	-	-	-	-	-	-	-	2	4
Male	297	-	-	-	-	-	-	-	-	-	-	4
Female	545	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	809	-	-	-	-	-	-	-	-	-	2	4
Male	284	-	-	-	-	-	-	-	-	-	-	4
Female	525	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
North Carolina												
All races, both sexes	394	-	-	-	-	-	-	-	1	1	1	3
Male	172	-	-	-	-	-	-	-	1	-	1	3
Female	222	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	343	-	-	-	-	-	-	-	1	1	-	3
Male	147	-	-	-	-	-	-	-	1	-	-	3
Female	196	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	48	-	-	-	-	-	-	-	-	-	1	-
Male	25	-	-	-	-	-	-	-	-	-	1	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	41	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	41	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Ohio												
All races, both sexes	690	-	-	-	-	-	-	-	1	-	1	-
Male	265	-	-	-	-	-	-	-	-	-	1	-
Female	425	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	663	-	-	-	-	-	-	-	1	-	-	-
Male	255	-	-	-	-	-	-	-	-	-	-	-
Female	408	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	25	-	-	-	-	-	-	-	-	-	1	-
Male	10	-	-	-	-	-	-	-	-	-	1	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	147	-	-	-	-	-	-	2	-	-	-	2
Male	65	-	-	-	-	-	-	1	-	-	-	1
Female	82	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	130	-	-	-	-	-	-	1	-	-	-	2
Male	60	-	-	-	-	-	-	-	-	-	-	1
Female	70	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	1	-	-	-	-
Male	4	-	-	-	-	-	-	1	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Oregon												
All races, both sexes	415	-	-	1	-	-	1	-	-	1	-	2
Male	170	-	-	1	-	-	-	-	-	1	-	2
Female	245	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	411	-	-	1	-	-	1	-	-	1	-	2
Male	168	-	-	1	-	-	-	-	-	1	-	2
Female	243	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	996	-	-	-	-	-	1	-	1	1	2	4
Male	357	-	-	-	-	-	1	-	-	1	1	1
Female	639	-	-	-	-	-	-	-	1	-	1	3
White, both sexes	969	-	-	-	-	-	1	-	1	-	2	3
Male	346	-	-	-	-	-	1	-	-	-	1	1
Female	623	-	-	-	-	-	-	-	1	-	1	2
Black, both sexes	26	-	-	-	-	-	-	-	-	1	-	1
Male	10	-	-	-	-	-	-	-	-	1	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Rhode Island												
All races, both sexes	58	-	-	-	-	-	-	-	-	-	-	-
Male	23	-	-	-	-	-	-	-	-	-	-	-
Female	35	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	58	-	-	-	-	-	-	-	-	-	-	-
Male	23	-	-	-	-	-	-	-	-	-	-	-
Female	35	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	217	-	-	-	-	1	-	-	-	1	1	1
Male	88	-	-	-	-	1	-	-	-	1	1	-
Female	129	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	174	-	-	-	-	-	-	-	-	-	-	-
Male	75	-	-	-	-	-	-	-	-	-	-	-
Female	99	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	43	-	-	-	-	1	-	-	-	1	1	1
Male	13	-	-	-	-	1	-	-	-	1	1	-
Female	30	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
South Dakota												
All races, both sexes	45	-	-	-	-	-	-	-	1	-	-	2
Male	18	-	-	-	-	-	-	-	1	-	-	1
Female	27	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	45	-	-	-	-	-	-	-	1	-	-	2
Male	18	-	-	-	-	-	-	-	1	-	-	1
Female	27	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	286	-	-	-	-	-	-	3	-	2	2	6
Male	120	-	-	-	-	-	-	2	-	1	1	5
Female	166	-	-	-	-	-	-	1	-	1	1	1
White, both sexes	260	-	-	-	-	-	-	2	-	1	2	5
Male	108	-	-	-	-	-	-	1	-	1	1	4
Female	152	-	-	-	-	-	-	1	-	-	1	1
Black, both sexes	26	-	-	-	-	-	-	1	-	1	-	1
Male	12	-	-	-	-	-	-	1	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Texas												
All races, both sexes	739	-	-	-	-	-	-	5	2	2	7	8
Male	341	-	-	-	-	-	-	4	2	1	4	5
Female	398	-	-	-	-	-	-	1	-	1	3	3
White, both sexes	686	-	-	-	-	-	-	4	2	1	4	8
Male	313	-	-	-	-	-	-	3	2	-	1	5
Female	373	-	-	-	-	-	-	1	-	1	3	3
Black, both sexes	47	-	-	-	-	-	-	1	-	1	3	-
Male	24	-	-	-	-	-	-	1	-	1	3	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	89	-	-	-	-	-	-	1	1	-	-	-
Male	38	-	-	-	-	-	-	1	1	-	-	-
Female	51	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	88	-	-	-	-	-	-	1	1	-	-	-
Male	38	-	-	-	-	-	-	1	1	-	-	-
Female	50	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Vermont												
All races, both sexes	46	-	-	-	-	-	-	-	-	-	-	-
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	46	-	-	-	-	-	-	-	-	-	-	-
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	341	1	-	1	-	-	-	-	1	-	1	3
Male	129	-	-	1	-	-	-	-	1	-	1	-
Female	212	1	-	-	-	-	-	-	-	-	-	3
White, both sexes	316	-	-	1	-	-	-	-	1	-	1	3
Male	116	-	-	1	-	-	-	-	1	-	1	-
Female	200	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	1	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Washington												
All races, both sexes	526	-	-	-	-	-	-	-	-	-	2	3
Male	219	-	-	-	-	-	-	-	-	-	2	2
Female	307	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	513	-	-	-	-	-	-	-	-	-	2	3
Male	217	-	-	-	-	-	-	-	-	-	2	2
Female	296	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	103	-	-	-	-	-	-	1	-	-	2	1
Male	44	-	-	-	-	-	-	-	-	-	2	1
Female	59	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	102	-	-	-	-	-	-	1	-	-	2	1
Male	44	-	-	-	-	-	-	-	-	-	2	1
Female	58	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Wisconsin												
All races, both sexes	400	-	-	-	-	-	-	-	1	-	3	1
Male	126	-	-	-	-	-	-	-	1	-	3	-
Female	274	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	396	-	-	-	-	-	-	-	1	-	3	1
Male	125	-	-	-	-	-	-	-	1	-	3	-
Female	271	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
United States												
All races, both sexes	157	216	269	458	780	1,436	2,679	4,018	3,401	1,303	233	-
Male	117	162	170	258	427	717	1,205	1,586	1,119	314	41	-
Female	40	54	99	200	353	719	1,474	2,432	2,282	989	192	-
White, both sexes	133	190	237	400	700	1,341	2,528	3,854	3,260	1,246	220	-
Male	100	145	150	228	389	673	1,142	1,529	1,085	296	35	-
Female	33	45	87	172	311	668	1,386	2,325	2,175	950	185	-
Black, both sexes	23	22	28	46	67	77	114	122	91	40	10	-
Male	16	13	18	25	31	33	48	40	18	8	4	-
Female	7	9	10	21	36	44	66	82	73	32	6	-
American Indian 1/, both sexes	1	1	1	3	4	4	4	2	7	2	2	-
Male	1	1	1	1	1	3	1	-	1	1	1	-
Female	-	-	-	2	3	1	3	2	6	1	1	-
Asian or Pacific Islander 2/, both sexes	-	3	3	9	9	14	33	40	43	15	1	-
Male	-	3	1	4	6	8	14	17	15	9	1	-
Female	-	-	2	5	3	6	19	23	28	6	-	-
Alabama												
All races, both sexes	3	2	3	5	17	28	30	31	25	14	3	-
Male	3	1	1	5	8	11	11	11	13	5	-	-
Female	-	1	2	-	9	17	19	20	12	9	3	-
White, both sexes	-	2	2	3	14	22	25	29	23	13	3	-
Male	-	1	-	3	8	9	11	10	13	5	-	-
Female	-	1	2	-	6	13	14	19	10	8	3	-
Black, both sexes	3	-	1	2	3	6	5	2	2	1	-	-
Male	3	-	1	2	-	2	-	1	-	-	-	-
Female	-	-	-	-	3	4	5	1	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Alaska												
All races, both sexes	1	2	-	2	2	3	5	3	1	3	-	-
Male	1	2	-	1	1	2	2	2	1	1	-	-
Female	-	-	-	1	1	1	3	1	-	2	-	-
White, both sexes	1	2	-	2	1	2	4	3	1	3	-	-
Male	1	2	-	1	1	1	2	2	1	1	-	-
Female	-	-	-	1	-	1	2	1	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	4	7	6	8	15	31	43	58	52	12	2	-
Male	4	3	5	5	6	16	19	14	15	1	-	-
Female	-	4	1	3	9	15	24	44	37	11	2	-
White, both sexes	3	7	5	8	14	30	42	58	52	12	2	-
Male	3	3	4	5	5	16	18	14	15	1	-	-
Female	-	4	1	3	9	14	24	44	37	11	2	-
Black, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Arkansas												
All races, both sexes	1	3	2	3	10	9	20	28	12	10	3	-
Male	-	3	1	1	5	6	8	9	5	3	1	-
Female	1	-	1	2	5	3	12	19	7	7	2	-
White, both sexes	1	3	2	3	10	8	20	27	11	10	2	-
Male	-	3	1	1	5	6	8	9	5	3	1	-
Female	1	-	1	2	5	2	12	18	6	7	1	-
Black, both sexes	-	-	-	-	-	1	-	1	1	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	17	24	29	46	80	178	306	473	444	177	41	-
Male	14	20	21	25	37	87	147	196	154	48	8	-
Female	3	4	8	21	43	91	159	277	290	129	33	-
White, both sexes	14	20	27	39	73	166	281	454	411	165	39	-
Male	12	17	19	20	35	81	134	187	143	44	8	-
Female	2	3	8	19	38	85	147	267	268	121	31	-
Black, both sexes	3	2	2	3	3	3	12	10	11	5	1	-
Male	2	1	2	2	-	1	5	5	2	-	-	-
Female	1	1	-	1	3	2	7	5	9	5	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	2	1	1	-
Male	-	-	-	-	-	-	-	-	1	1	-	-
Female	-	-	-	-	-	-	-	-	1	-	1	-
Asian or Pacific Islander 2/, both sexes	-	2	-	4	4	9	13	9	20	6	-	-
Male	-	2	-	3	2	5	8	4	8	3	-	-
Female	-	-	-	1	2	4	5	5	12	3	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Colorado												
All races, both sexes	4	1	4	6	10	22	33	48	34	13	2	-
Male	4	1	1	4	6	12	17	20	13	4	-	-
Female	-	-	3	2	4	10	16	28	21	9	2	-
White, both sexes	4	1	4	6	10	21	33	46	34	13	2	-
Male	4	1	1	4	6	12	17	20	13	4	-	-
Female	-	-	3	2	4	9	16	26	21	9	2	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Connecticut												
All races, both sexes	4	-	3	3	7	14	34	51	59	19	3	-
Male	4	-	2	3	5	11	13	16	21	6	-	-
Female	-	-	1	-	2	3	21	35	38	13	3	-
White, both sexes	4	-	1	3	6	13	34	49	59	18	3	-
Male	4	-	1	3	4	10	13	16	21	6	-	-
Female	-	-	-	-	2	3	21	33	38	12	3	-
Black, both sexes	-	-	2	-	1	1	-	2	-	-	-	-
Male	-	-	1	-	1	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Delaware												
All races, both sexes	-	4	-	5	3	5	13	11	9	4	1	-
Male	-	3	-	2	2	4	5	3	2	1	-	-
Female	-	1	-	3	1	1	8	8	7	3	1	-
White, both sexes	-	3	-	5	3	4	12	11	7	4	1	-
Male	-	2	-	2	2	3	4	3	1	1	-	-
Female	-	1	-	3	1	1	8	8	6	3	1	-
Black, both sexes	-	1	-	-	-	1	1	-	2	-	-	-
Male	-	1	-	-	-	1	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	1	1	-	-	4	4	-	4	4	-	-
Male	-	1	-	-	-	2	1	-	4	-	-	-
Female	-	-	1	-	-	2	3	-	-	4	-	-
White, both sexes	-	-	1	-	-	1	1	-	4	3	-	-
Male	-	-	-	-	-	1	1	-	4	-	-	-
Female	-	-	1	-	-	-	-	-	-	3	-	-
Black, both sexes	-	1	-	-	-	3	3	-	-	1	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	2	3	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Florida												
All races, both sexes	17	16	20	37	51	106	188	239	213	82	11	-
Male	13	13	10	20	33	56	100	107	84	24	3	-
Female	4	3	10	17	18	50	88	132	129	58	8	-
White, both sexes	13	16	18	33	45	100	181	228	208	79	10	-
Male	11	13	10	18	27	55	95	105	82	23	2	-
Female	2	3	8	15	18	45	86	123	126	56	8	-
Black, both sexes	4	-	2	4	6	5	6	10	5	2	1	-
Male	2	-	-	2	6	-	4	2	2	-	1	-
Female	2	-	2	2	-	5	2	8	3	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	1	-	1	-	-
Male	-	-	-	-	-	1	1	-	-	1	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Georgia												
All races, both sexes	-	9	7	12	26	29	34	72	40	22	3	-
Male	-	5	4	6	15	19	13	28	12	8	1	-
Female	-	4	3	6	11	10	21	44	28	14	2	-
White, both sexes	-	5	6	10	19	23	28	66	35	17	2	-
Male	-	3	4	5	10	17	11	27	11	5	-	-
Female	-	2	2	5	9	6	17	39	24	12	2	-
Black, both sexes	-	4	1	2	7	6	6	6	4	5	1	-
Male	-	2	-	1	5	2	2	1	1	3	1	-
Female	-	2	1	1	2	4	4	5	3	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Hawaii												
All races, both sexes	-	1	1	2	-	2	10	27	15	4	1	-
Male	-	1	1	1	-	1	1	11	7	2	1	-
Female	-	-	-	1	-	1	9	16	8	2	-	-
White, both sexes	-	1	1	1	-	1	3	6	6	-	-	-
Male	-	1	1	1	-	-	-	2	3	-	-	-
Female	-	-	-	-	-	1	3	4	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	7	21	9	4	1	-
Male	-	-	-	-	-	1	1	9	4	2	1	-
Female	-	-	-	1	-	-	6	12	5	2	-	-
Idaho												
All races, both sexes	-	2	1	1	5	5	20	27	16	7	1	-
Male	-	1	1	1	5	2	13	12	9	-	-	-
Female	-	1	-	-	-	3	7	15	7	7	1	-
White, both sexes	-	2	1	1	5	5	20	27	16	7	1	-
Male	-	1	1	1	5	2	13	12	9	-	-	-
Female	-	1	-	-	-	3	7	15	7	7	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Illinois												
All races, both sexes	3	4	9	16	25	50	92	151	129	55	5	-
Male	1	3	6	8	11	26	40	70	38	13	-	-
Female	2	1	3	8	14	24	52	81	91	42	5	-
White, both sexes	2	4	7	13	23	47	83	146	124	52	5	-
Male	1	3	4	5	11	24	35	68	37	12	-	-
Female	1	1	3	8	12	23	48	78	87	40	5	-
Black, both sexes	1	-	2	3	2	3	8	5	4	3	-	-
Male	-	-	2	3	-	2	4	2	1	1	-	-
Female	1	-	-	-	2	1	4	3	3	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Indiana												
All races, both sexes	2	3	6	17	21	37	62	83	65	22	2	-
Male	1	2	3	12	10	18	36	39	21	5	-	-
Female	1	1	3	5	11	19	26	44	44	17	2	-
White, both sexes	2	3	6	17	18	37	61	81	62	22	2	-
Male	1	2	3	12	9	18	36	37	20	5	-	-
Female	1	1	3	5	9	19	25	44	42	17	2	-
Black, both sexes	-	-	-	-	3	-	1	2	3	-	-	-
Male	-	-	-	-	1	-	-	2	1	-	-	-
Female	-	-	-	-	2	-	1	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Iowa												
All races, both sexes	2	1	3	2	10	17	35	68	69	30	3	-
Male	2	1	-	1	4	11	15	30	21	9	-	-
Female	-	-	3	1	6	6	20	38	48	21	3	-
White, both sexes	2	1	2	2	10	16	35	68	69	30	3	-
Male	2	1	-	1	4	11	15	30	21	9	-	-
Female	-	-	2	1	6	5	20	38	48	21	3	-
Black, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	2	1	1	1	3	10	30	45	50	22	4	-
Male	1	1	-	1	2	5	7	24	14	6	1	-
Female	1	-	1	-	1	5	23	21	36	16	3	-
White, both sexes	2	1	1	1	2	10	29	44	47	21	4	-
Male	1	1	-	1	2	5	7	24	14	5	1	-
Female	1	-	1	-	-	5	22	20	33	16	3	-
Black, both sexes	-	-	-	-	1	-	1	1	2	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	1	-	1	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Kentucky												
All races, both sexes	2	3	3	7	9	21	23	35	25	14	2	-
Male	1	2	1	4	4	12	16	13	7	2	-	-
Female	1	1	2	3	5	9	7	22	18	12	2	-
White, both sexes	2	2	3	6	9	20	23	35	25	14	2	-
Male	1	1	1	3	4	12	16	13	7	2	-	-
Female	1	1	2	3	5	8	7	22	18	12	2	-
Black, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	3	5	4	6	7	19	37	37	28	9	3	-
Male	1	2	2	3	3	12	12	11	8	2	1	-
Female	2	3	2	3	4	7	25	26	20	7	2	-
White, both sexes	2	3	3	4	7	16	29	33	23	9	3	-
Male	1	1	1	2	3	10	9	11	8	2	1	-
Female	1	2	2	2	4	6	20	22	15	7	2	-
Black, both sexes	1	2	1	2	-	3	7	4	5	-	-	-
Male	-	1	1	1	-	2	3	-	-	-	-	-
Female	1	1	-	1	-	1	4	4	5	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Maine												
All races, both sexes	1	1	5	5	6	9	17	33	21	9	3	-
Male	1	-	3	2	3	3	11	17	8	3	1	-
Female	-	1	2	3	3	6	6	16	13	6	2	-
White, both sexes	1	1	5	5	6	9	17	33	21	9	3	-
Male	1	-	3	2	3	3	11	17	8	3	1	-
Female	-	1	2	3	3	6	6	16	13	6	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	2	4	5	9	10	22	53	78	57	19	3	-
Male	1	2	5	8	4	12	22	36	27	5	-	-
Female	1	2	-	1	6	10	31	42	30	14	3	-
White, both sexes	1	4	4	8	6	20	47	69	55	16	3	-
Male	1	2	4	7	4	12	18	32	26	4	-	-
Female	-	2	-	1	2	8	29	37	29	12	3	-
Black, both sexes	1	-	1	1	4	2	5	9	2	2	-	-
Male	-	-	1	1	-	-	3	4	1	1	-	-
Female	1	-	-	-	4	2	2	5	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Massachusetts												
All races, both sexes	6	2	3	10	16	28	74	109	103	36	8	-
Male	4	1	2	4	10	18	32	37	27	4	1	-
Female	2	1	1	6	6	10	42	72	76	32	7	-
White, both sexes	6	2	3	10	16	27	72	107	101	34	8	-
Male	4	1	2	4	10	17	30	37	27	4	1	-
Female	2	1	1	6	6	10	42	70	74	30	7	-
Black, both sexes	-	-	-	-	-	1	2	2	1	2	-	-
Male	-	-	-	-	-	1	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	1	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Michigan												
All races, both sexes	7	13	8	15	21	42	64	125	105	42	5	-
Male	7	10	6	10	14	23	32	50	38	9	1	-
Female	-	3	2	5	7	19	32	75	67	33	4	-
White, both sexes	4	13	7	15	19	40	56	121	99	42	5	-
Male	4	10	6	10	12	23	29	48	35	9	1	-
Female	-	3	1	5	7	17	27	73	64	33	4	-
Black, both sexes	3	-	1	-	2	2	6	3	5	-	-	-
Male	3	-	-	-	2	-	2	2	2	-	-	-
Female	-	-	1	-	-	2	4	1	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Minnesota												
All races, both sexes	1	3	3	3	12	29	57	102	93	36	3	-
Male	-	3	3	1	8	13	36	43	35	8	-	-
Female	1	-	-	2	4	16	21	59	58	28	3	-
White, both sexes	1	3	3	3	12	28	57	100	92	36	2	-
Male	-	3	3	1	8	13	36	43	35	8	-	-
Female	1	-	-	2	4	15	21	57	57	28	2	-
Black, both sexes	-	-	-	-	-	1	-	1	1	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	1	-	5	7	8	20	18	11	4	1	-
Male	-	1	-	2	2	2	8	5	3	2	1	-
Female	-	-	-	3	5	6	12	13	8	2	-	-
White, both sexes	-	1	-	1	7	7	18	14	9	4	1	-
Male	-	1	-	-	2	1	8	4	3	2	1	-
Female	-	-	-	1	5	6	10	10	6	2	-	-
Black, both sexes	-	-	-	4	-	-	1	4	2	-	-	-
Male	-	-	-	2	-	-	-	1	-	-	-	-
Female	-	-	-	2	-	-	1	3	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Missouri												
All races, both sexes	2	1	10	9	16	29	50	80	78	25	4	-
Male	2	-	5	3	10	15	22	32	19	7	2	-
Female	-	1	5	6	6	14	28	48	59	18	2	-
White, both sexes	2	1	10	9	15	27	50	75	77	24	3	-
Male	2	-	5	3	10	14	22	30	19	7	2	-
Female	-	1	5	6	5	13	28	45	58	17	1	-
Black, both sexes	-	-	-	-	1	2	-	5	-	1	1	-
Male	-	-	-	-	-	1	-	2	-	-	-	-
Female	-	-	-	-	1	1	-	3	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Montana												
All races, both sexes	1	1	1	5	3	4	11	18	17	8	1	-
Male	1	1	-	5	2	2	3	8	9	2	-	-
Female	-	-	1	-	1	2	8	10	8	6	1	-
White, both sexes	1	1	1	5	3	4	11	18	17	8	1	-
Male	1	1	-	5	2	2	3	8	9	2	-	-
Female	-	-	1	-	1	2	8	10	8	6	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Nebraska												
All races, both sexes	2	1	-	2	2	9	26	34	41	14	2	-
Male	1	1	-	1	2	2	14	7	17	3	-	-
Female	1	-	-	1	-	7	12	27	24	11	2	-
White, both sexes	2	1	-	2	1	9	26	34	40	14	2	-
Male	1	1	-	1	1	2	14	7	17	3	-	-
Female	1	-	-	1	-	7	12	27	23	11	2	-
Black, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	1	2	1	8	10	20	21	15	5	-	-
Male	-	1	2	-	4	5	11	8	5	2	-	-
Female	-	-	-	1	4	5	9	13	10	3	-	-
White, both sexes	-	1	2	-	6	9	19	21	14	4	-	-
Male	-	1	2	-	3	4	11	8	4	1	-	-
Female	-	-	-	-	3	5	8	13	10	3	-	-
Black, both sexes	-	-	-	-	1	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	1	1	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
New Hampshire												
All races, both sexes	-	-	3	2	3	3	15	29	24	4	2	-
Male	-	-	1	1	1	1	6	14	6	1	1	-
Female	-	-	2	1	2	2	9	15	18	3	1	-
White, both sexes	-	-	3	2	3	3	15	29	24	4	2	-
Male	-	-	1	1	1	1	6	14	6	1	1	-
Female	-	-	2	1	2	2	9	15	18	3	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	7	8	6	8	26	53	93	155	126	46	10	-
Male	5	7	5	7	14	29	48	51	43	11	2	-
Female	2	1	1	1	12	24	45	104	83	35	8	-
White, both sexes	6	7	5	5	23	49	90	150	122	45	10	-
Male	5	7	4	5	13	26	46	49	43	10	2	-
Female	1	-	1	-	10	23	44	101	79	35	8	-
Black, both sexes	1	1	1	2	3	4	3	4	3	-	-	-
Male	-	-	1	1	1	3	2	1	-	-	-	-
Female	1	1	-	1	2	1	1	3	3	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	1	1	-	-
Male	-	-	-	-	-	-	-	1	-	1	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
New Mexico												
All races, both sexes	-	1	2	2	1	9	18	23	9	6	-	-
Male	-	1	2	1	1	4	11	10	2	2	-	-
Female	-	-	-	1	-	5	7	13	7	4	-	-
White, both sexes	-	1	2	2	1	9	15	23	8	6	-	-
Male	-	1	2	1	1	4	9	10	2	2	-	-
Female	-	-	-	1	-	5	6	13	6	4	-	-
Black, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	4	13	13	16	42	68	145	234	207	74	20	-
Male	3	10	10	7	25	27	59	79	57	14	2	-
Female	1	3	3	9	17	41	86	155	150	60	18	-
White, both sexes	4	11	11	15	37	68	135	227	202	73	20	-
Male	3	9	9	7	21	27	55	78	55	14	2	-
Female	1	2	2	8	16	41	80	149	147	59	18	-
Black, both sexes	-	2	1	-	4	-	8	5	4	1	-	-
Male	-	1	1	-	3	-	4	-	2	-	-	-
Female	-	1	-	-	1	-	4	5	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	1	-	2	2	1	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	1	1	-	-	2	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
North Carolina												
All races, both sexes	7	2	8	13	24	36	82	112	70	26	8	-
Male	4	2	5	6	16	21	36	45	22	7	3	-
Female	3	-	3	7	8	15	46	67	48	19	5	-
White, both sexes	6	2	5	7	19	28	75	102	67	22	5	-
Male	3	2	3	4	13	16	34	39	21	7	1	-
Female	3	-	2	3	6	12	41	63	46	15	4	-
Black, both sexes	1	-	3	5	4	8	6	10	3	4	3	-
Male	1	-	2	2	3	5	2	6	1	-	2	-
Female	-	-	1	3	1	3	4	4	2	4	1	-
American Indian 1/, both sexes	-	-	-	1	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	1	-	-	2	2	7	11	16	2	-	-
Male	-	1	-	-	1	1	4	5	4	-	-	-
Female	-	-	-	-	1	1	3	6	12	2	-	-
White, both sexes	-	1	-	-	2	2	7	11	16	2	-	-
Male	-	1	-	-	1	1	4	5	4	-	-	-
Female	-	-	-	-	1	1	3	6	12	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Ohio												
All races, both sexes	6	4	9	20	35	58	148	196	136	63	13	-
Male	4	3	7	12	18	30	54	77	42	15	2	-
Female	2	1	2	8	17	28	94	119	94	48	11	-
White, both sexes	6	3	9	18	30	56	142	194	131	60	13	-
Male	4	3	7	11	17	29	51	76	41	14	2	-
Female	2	-	2	7	13	27	91	118	90	46	11	-
Black, both sexes	-	1	-	2	5	2	6	2	3	3	-	-
Male	-	-	-	1	1	1	3	1	1	1	-	-
Female	-	1	-	1	4	1	3	1	2	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	4	2	3	6	12	22	25	39	18	10	2	-
Male	3	1	3	3	7	14	7	14	6	3	2	-
Female	1	1	-	3	5	8	18	25	12	7	-	-
White, both sexes	4	1	2	5	9	20	22	37	16	10	1	-
Male	3	1	2	3	7	13	6	14	6	3	1	-
Female	1	-	-	2	2	7	16	23	10	7	-	-
Black, both sexes	-	1	-	1	2	-	3	2	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	1	2	-	2	2	1	-	-	-
American Indian 1/, both sexes	-	-	1	-	1	2	-	-	1	-	1	-
Male	-	-	1	-	-	1	-	-	-	-	1	-
Female	-	-	-	-	1	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Oregon												
All races, both sexes	3	5	4	9	25	43	66	112	105	29	9	-
Male	3	2	2	3	16	19	29	44	36	11	1	-
Female	-	3	2	6	9	24	37	68	69	18	8	-
White, both sexes	2	5	4	8	25	43	66	111	105	28	9	-
Male	2	2	2	3	16	19	29	44	36	10	1	-
Female	-	3	2	5	9	24	37	67	69	18	8	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Pennsylvania												
All races, both sexes	3	18	14	25	52	68	180	280	246	88	13	-
Male	2	14	8	10	23	29	74	99	72	20	2	-
Female	1	4	6	15	29	39	106	181	174	68	11	-
White, both sexes	3	16	14	24	50	67	177	273	241	85	12	-
Male	2	12	8	10	22	28	72	96	71	20	2	-
Female	1	4	6	14	28	39	105	177	170	65	10	-
Black, both sexes	-	2	-	1	1	1	3	7	5	3	1	-
Male	-	2	-	-	-	1	2	3	1	-	-	-
Female	-	-	-	1	1	-	1	4	4	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Rhode Island												
All races, both sexes	-	-	3	-	1	7	9	15	15	7	1	-
Male	-	-	3	-	1	2	4	6	6	1	-	-
Female	-	-	-	-	-	5	5	9	9	6	1	-
White, both sexes	-	-	3	-	1	7	9	15	15	7	1	-
Male	-	-	3	-	1	2	4	6	6	1	-	-
Female	-	-	-	-	-	5	5	9	9	6	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	4	3	6	13	14	28	34	52	46	13	-	-
Male	4	3	3	9	4	15	15	19	12	1	-	-
Female	-	-	3	4	10	13	19	33	34	12	-	-
White, both sexes	2	3	6	10	8	24	31	44	35	11	-	-
Male	2	3	3	7	2	13	15	18	11	1	-	-
Female	-	-	3	3	6	11	16	26	24	10	-	-
Black, both sexes	2	-	-	3	6	4	3	8	11	2	-	-
Male	2	-	-	2	2	2	-	1	1	-	-	-
Female	-	-	-	1	4	2	3	7	10	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
South Dakota												
All races, both sexes	-	-	-	-	2	4	8	11	11	4	2	-
Male	-	-	-	-	2	3	4	3	3	-	1	-
Female	-	-	-	-	-	1	4	8	8	4	1	-
White, both sexes	-	-	-	-	2	4	8	11	11	4	2	-
Male	-	-	-	-	2	3	4	3	3	-	1	-
Female	-	-	-	-	-	1	4	8	8	4	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	3	2	12	18	21	28	58	57	55	17	2	-
Male	2	2	6	11	11	11	29	17	20	2	-	-
Female	1	-	6	7	10	17	29	40	35	15	2	-
White, both sexes	2	1	9	17	19	26	52	55	50	17	2	-
Male	1	1	4	11	10	10	26	17	19	2	-	-
Female	1	-	5	6	9	16	26	38	31	15	2	-
Black, both sexes	1	1	3	1	2	2	6	2	5	-	-	-
Male	1	1	2	-	1	1	3	-	1	-	-	-
Female	-	-	1	1	1	1	3	2	4	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Texas												
All races, both sexes	13	25	23	38	47	103	110	178	123	50	5	-
Male	10	19	17	19	26	52	46	83	39	13	1	-
Female	3	6	6	19	21	51	64	95	84	37	4	-
White, both sexes	12	22	19	32	45	93	106	167	119	47	5	-
Male	9	16	13	16	25	47	45	78	39	13	1	-
Female	3	6	6	16	20	46	61	89	80	34	4	-
Black, both sexes	1	2	4	5	2	9	4	9	3	3	-	-
Male	1	2	4	2	1	4	1	4	-	-	-	-
Female	-	-	-	3	1	5	3	5	3	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	1	-	2	1	-	-	-
Male	-	1	-	1	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Utah												
All races, both sexes	1	2	-	5	3	6	23	29	13	5	-	-
Male	1	2	-	4	2	2	9	10	4	2	-	-
Female	-	-	-	1	1	4	14	19	9	3	-	-
White, both sexes	1	2	-	5	3	6	23	28	13	5	-	-
Male	1	2	-	4	2	2	9	10	4	2	-	-
Female	-	-	-	1	1	4	14	18	9	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Vermont												
All races, both sexes	1	1	-	2	-	9	9	10	9	5	-	-
Male	1	1	-	2	-	2	3	6	4	2	-	-
Female	-	-	-	-	-	7	6	4	5	3	-	-
White, both sexes	1	1	-	2	-	9	9	10	9	5	-	-
Male	1	1	-	2	-	2	3	6	4	2	-	-
Female	-	-	-	-	-	7	6	4	5	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	6	3	4	15	16	29	57	90	82	26	6	-
Male	4	2	2	8	11	15	25	33	19	7	-	-
Female	2	1	2	7	5	14	32	57	63	19	6	-
White, both sexes	6	1	3	11	12	26	56	86	78	25	6	-
Male	4	1	2	6	7	14	24	31	18	6	-	-
Female	2	-	1	5	5	12	32	55	60	19	6	-
Black, both sexes	-	2	1	4	2	3	1	4	2	1	-	-
Male	-	1	-	2	2	1	1	2	-	1	-	-
Female	-	1	1	2	-	2	-	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	-	-	-	2	-	-	-
Male	-	-	-	-	2	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Washington												
All races, both sexes	5	5	11	13	25	35	86	128	143	59	11	-
Male	3	5	5	9	18	14	36	62	50	11	2	-
Female	2	-	6	4	7	21	50	66	93	48	9	-
White, both sexes	5	5	9	12	24	34	81	127	141	59	11	-
Male	3	5	5	9	18	14	35	61	50	11	2	-
Female	2	-	4	3	6	20	46	66	91	48	9	-
Black, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	1	1	4	1	2	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	1	1	1	1	3	-	2	-	-	-
West Virginia												
All races, both sexes	1	1	1	6	10	12	19	27	18	2	2	-
Male	1	-	1	4	7	4	8	10	6	-	-	-
Female	-	1	-	2	3	8	11	17	12	2	2	-
White, both sexes	1	1	1	6	10	12	19	26	18	2	2	-
Male	1	-	1	4	7	4	8	10	6	-	-	-
Female	-	1	-	2	3	8	11	16	12	2	2	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Wisconsin												
All races, both sexes	1	3	6	3	14	30	72	124	91	44	7	-
Male	-	2	4	2	5	12	28	39	25	5	-	-
Female	1	1	2	1	9	18	44	85	66	39	7	-
White, both sexes	1	3	6	3	14	30	69	124	90	44	7	-
Male	-	2	4	2	5	12	27	39	25	5	-	-
Female	1	1	2	1	9	18	42	85	65	39	7	-
Black, both sexes	-	-	-	-	-	-	3	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	-	1	1	3	3	4	1	7	2	-	-
Male	-	-	1	1	2	2	3	1	4	1	-	-
Female	1	-	-	-	1	1	1	-	3	1	-	-
White, both sexes	1	-	1	1	3	3	4	1	7	2	-	-
Male	-	-	1	1	2	2	3	1	4	1	-	-
Female	1	-	-	-	1	1	1	-	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
United States												
All races, both sexes	5,486	1	1	-	-	-	11	22	29	57	76	149
Male	2,258	1	-	-	-	-	6	11	17	29	48	92
Female	3,228	-	1	-	-	-	5	11	12	28	28	57
White, both sexes	4,860	1	-	-	-	-	9	19	19	41	48	105
Male	1,947	1	-	-	-	-	5	11	13	19	34	69
Female	2,913	-	-	-	-	-	4	8	6	22	14	36
Black, both sexes	522	-	1	-	-	-	2	1	9	14	25	39
Male	260	-	-	-	-	-	1	-	4	9	14	19
Female	262	-	1	-	-	-	1	1	5	5	11	20
American Indian 1/, both sexes	24	-	-	-	-	-	-	-	1	1	2	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	1	1	2	-
Asian or Pacific Islander 2/, both sexes	80	-	-	-	-	-	-	2	-	1	1	5
Male	40	-	-	-	-	-	-	-	-	1	-	4
Female	40	-	-	-	-	-	-	2	-	-	1	1
Alabama												
All races, both sexes	87	-	-	-	-	-	1	-	-	1	3	2
Male	37	-	-	-	-	-	1	-	-	-	-	1
Female	50	-	-	-	-	-	-	-	-	1	3	1
White, both sexes	63	-	-	-	-	-	1	-	-	1	1	1
Male	21	-	-	-	-	-	1	-	-	-	-	-
Female	42	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	24	-	-	-	-	-	-	-	-	-	2	1
Male	16	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Alaska												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	105	-	-	-	-	-	-	1	2	-	1	5
Male	50	-	-	-	-	-	-	1	2	-	-	2
Female	55	-	-	-	-	-	-	-	-	-	1	3
White, both sexes	99	-	-	-	-	-	-	1	1	-	1	3
Male	46	-	-	-	-	-	-	1	1	-	-	1
Female	53	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	3	-	-	-	-	-	-	-	1	-	-	1
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Arkansas												
All races, both sexes	51	-	-	-	-	-	-	1	-	-	1	1
Male	17	-	-	-	-	-	-	1	-	-	-	1
Female	34	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	45	-	-	-	-	-	-	1	-	-	-	-
Male	15	-	-	-	-	-	-	1	-	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	423	-	1	-	-	-	-	3	1	8	10	10
Male	178	-	-	-	-	-	-	1	-	5	7	7
Female	245	-	1	-	-	-	-	2	1	3	3	3
White, both sexes	357	-	-	-	-	-	-	1	1	7	8	9
Male	144	-	-	-	-	-	-	1	-	4	7	7
Female	213	-	-	-	-	-	-	-	1	3	1	2
Black, both sexes	35	-	1	-	-	-	-	1	-	1	1	-
Male	19	-	-	-	-	-	-	-	-	1	-	-
Female	16	-	1	-	-	-	-	1	-	-	1	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	28	-	-	-	-	-	-	1	-	-	1	1
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	1	-	-	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Colorado												
All races, both sexes	73	-	-	-	-	-	-	-	-	1	-	1
Male	30	-	-	-	-	-	-	-	-	-	-	1
Female	43	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	71	-	-	-	-	-	-	-	-	1	-	1
Male	29	-	-	-	-	-	-	-	-	-	-	1
Female	42	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	66	-	-	-	-	-	-	1	-	1	1	4
Male	27	-	-	-	-	-	-	-	-	1	1	2
Female	39	-	-	-	-	-	-	1	-	-	-	2
White, both sexes	60	-	-	-	-	-	-	1	-	-	1	3
Male	24	-	-	-	-	-	-	-	-	-	1	2
Female	36	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Delaware												
All races, both sexes	18	-	-	-	-	-	1	-	-	1	-	2
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	1	-	-	1	-	1
White, both sexes	13	-	-	-	-	-	1	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	1	-	-	1	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Florida												
All races, both sexes	475	-	-	-	-	-	1	-	1	7	3	16
Male	214	-	-	-	-	-	1	-	1	6	3	11
Female	261	-	-	-	-	-	-	-	-	1	-	5
White, both sexes	425	-	-	-	-	-	1	-	1	5	2	10
Male	191	-	-	-	-	-	1	-	1	4	2	8
Female	234	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	48	-	-	-	-	-	-	-	-	2	1	5
Male	22	-	-	-	-	-	-	-	-	2	1	2
Female	26	-	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	113	-	-	-	-	-	-	-	2	2	3	6
Male	58	-	-	-	-	-	-	-	-	2	2	4
Female	55	-	-	-	-	-	-	-	2	-	1	2
White, both sexes	87	-	-	-	-	-	-	-	2	2	2	2
Male	40	-	-	-	-	-	-	-	-	2	1	-
Female	47	-	-	-	-	-	-	-	2	-	1	2
Black, both sexes	26	-	-	-	-	-	-	-	-	-	1	4
Male	18	-	-	-	-	-	-	-	-	-	1	4
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Hawaii												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	36	-	-	-	-	-	-	-	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	23	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	36	-	-	-	-	-	-	-	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	23	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Illinois												
All races, both sexes	186	-	-	-	-	-	1	1	-	1	2	14
Male	86	-	-	-	-	-	-	1	-	1	1	8
Female	100	-	-	-	-	-	1	-	-	-	1	6
White, both sexes	160	-	-	-	-	-	1	1	-	1	-	11
Male	73	-	-	-	-	-	-	1	-	1	-	8
Female	87	-	-	-	-	-	1	-	-	-	-	3
Black, both sexes	25	-	-	-	-	-	-	-	-	-	2	3
Male	12	-	-	-	-	-	-	-	-	-	1	-
Female	13	-	-	-	-	-	-	-	-	-	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	125	-	-	-	-	-	-	2	-	2	3	4
Male	61	-	-	-	-	-	-	1	-	-	3	2
Female	64	-	-	-	-	-	-	1	-	2	-	2
White, both sexes	120	-	-	-	-	-	-	2	-	2	3	3
Male	58	-	-	-	-	-	-	1	-	-	3	2
Female	62	-	-	-	-	-	-	1	-	2	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Iowa												
All races, both sexes	31	-	-	-	-	-	-	-	-	-	-	2
Male	15	-	-	-	-	-	-	-	-	-	-	2
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	31	-	-	-	-	-	-	-	-	-	-	2
Male	15	-	-	-	-	-	-	-	-	-	-	2
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	50	-	-	-	-	-	-	-	-	-	1	1
Male	12	-	-	-	-	-	-	-	-	-	1	-
Female	38	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	49	-	-	-	-	-	-	-	-	-	1	1
Male	12	-	-	-	-	-	-	-	-	-	1	-
Female	37	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Kentucky												
All races, both sexes	70	-	-	-	-	-	-	-	-	1	-	2
Male	20	-	-	-	-	-	-	-	-	-	-	1
Female	50	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	68	-	-	-	-	-	-	-	-	1	-	2
Male	20	-	-	-	-	-	-	-	-	-	-	1
Female	48	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	88	-	-	-	-	-	1	-	-	2	1	2
Male	34	-	-	-	-	-	1	-	-	1	-	2
Female	54	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	69	-	-	-	-	-	1	-	-	2	1	1
Male	25	-	-	-	-	-	1	-	-	1	-	1
Female	44	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	18	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Maine												
All races, both sexes	41	-	-	-	-	-	-	-	-	-	1	-
Male	11	-	-	-	-	-	-	-	-	-	1	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	41	-	-	-	-	-	-	-	-	-	1	-
Male	11	-	-	-	-	-	-	-	-	-	1	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	80	-	-	-	-	-	-	1	1	2	3	2
Male	40	-	-	-	-	-	-	-	1	1	3	-
Female	40	-	-	-	-	-	-	1	-	1	-	2
White, both sexes	54	-	-	-	-	-	-	1	-	-	1	1
Male	27	-	-	-	-	-	-	-	-	-	1	-
Female	27	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	24	-	-	-	-	-	-	-	1	2	2	1
Male	13	-	-	-	-	-	-	-	1	1	2	-
Female	11	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Massachusetts												
All races, both sexes	97	-	-	-	-	-	-	2	2	1	3	1
Male	40	-	-	-	-	-	-	1	2	-	3	-
Female	57	-	-	-	-	-	-	1	-	1	-	1
White, both sexes	89	-	-	-	-	-	-	2	1	-	3	1
Male	37	-	-	-	-	-	-	1	1	-	3	-
Female	52	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	6	-	-	-	-	-	-	-	1	1	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	4	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	191	-	-	-	-	-	-	-	1	4	-	8
Male	90	-	-	-	-	-	-	-	-	-	-	6
Female	101	-	-	-	-	-	-	-	1	4	-	2
White, both sexes	162	-	-	-	-	-	-	-	-	4	-	6
Male	75	-	-	-	-	-	-	-	-	-	-	5
Female	87	-	-	-	-	-	-	-	-	4	-	1
Black, both sexes	27	-	-	-	-	-	-	-	1	-	-	2
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Minnesota												
All races, both sexes	64	-	-	-	-	-	-	-	-	1	1	1
Male	31	-	-	-	-	-	-	-	-	-	1	1
Female	33	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	61	-	-	-	-	-	-	-	-	-	1	1
Male	31	-	-	-	-	-	-	-	-	-	1	1
Female	30	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	34	-	-	-	-	-	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Missouri												
All races, both sexes	125	-	-	-	-	-	2	-	1	2	1	2
Male	48	-	-	-	-	-	1	-	1	2	1	1
Female	77	-	-	-	-	-	1	-	-	-	-	1
White, both sexes	114	-	-	-	-	-	2	-	1	-	1	2
Male	42	-	-	-	-	-	1	-	1	-	1	1
Female	72	-	-	-	-	-	1	-	-	-	-	1
Black, both sexes	11	-	-	-	-	-	-	-	-	2	-	-
Male	6	-	-	-	-	-	-	-	-	2	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	32	-	-	-	-	-	-	-	1	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	31	-	-	-	-	-	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Nebraska												
All races, both sexes	45	-	-	-	-	-	-	-	-	-	-	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	45	-	-	-	-	-	-	-	-	-	-	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	37	-	-	-	-	-	1	-	-	1	2	1
Male	23	-	-	-	-	-	1	-	-	1	2	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	33	-	-	-	-	-	1	-	-	1	1	1
Male	22	-	-	-	-	-	1	-	-	1	1	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
New Hampshire												
All races, both sexes	34	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	34	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	232	-	-	-	-	-	-	1	-	3	3	4
Male	88	-	-	-	-	-	-	-	-	1	1	4
Female	144	-	-	-	-	-	-	1	-	2	2	-
White, both sexes	211	-	-	-	-	-	-	-	-	1	1	2
Male	79	-	-	-	-	-	-	-	-	1	1	2
Female	132	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	19	-	-	-	-	-	-	-	-	2	2	2
Male	9	-	-	-	-	-	-	-	-	-	-	2
Female	10	-	-	-	-	-	-	-	-	2	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
New Mexico												
All races, both sexes	33	-	-	-	-	-	-	-	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	20	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	31	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	351	-	-	-	-	-	-	1	3	2	6	2
Male	139	-	-	-	-	-	-	1	2	2	2	1
Female	212	-	-	-	-	-	-	-	1	-	4	1
White, both sexes	319	-	-	-	-	-	-	1	3	-	4	1
Male	129	-	-	-	-	-	-	1	2	-	2	-
Female	190	-	-	-	-	-	-	-	1	-	2	1
Black, both sexes	24	-	-	-	-	-	-	-	-	2	2	1
Male	7	-	-	-	-	-	-	-	-	2	-	1
Female	17	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
North Carolina												
All races, both sexes	161	-	-	-	-	-	-	-	1	2	2	7
Male	57	-	-	-	-	-	-	-	1	1	-	3
Female	104	-	-	-	-	-	-	-	-	1	2	4
White, both sexes	126	-	-	-	-	-	-	-	-	1	2	2
Male	43	-	-	-	-	-	-	-	-	-	-	1
Female	83	-	-	-	-	-	-	-	-	1	2	1
Black, both sexes	33	-	-	-	-	-	-	-	1	1	-	5
Male	13	-	-	-	-	-	-	-	1	1	-	2
Female	20	-	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	17	-	-	-	-	-	-	1	1	-	-	-
Male	8	-	-	-	-	-	-	-	1	-	-	-
Female	9	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	1	1	-	-	-
Male	7	-	-	-	-	-	-	-	1	-	-	-
Female	9	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Ohio												
All races, both sexes	266	-	-	-	-	-	-	-	-	1	3	6
Male	90	-	-	-	-	-	-	-	-	-	3	4
Female	176	-	-	-	-	-	-	-	-	1	-	2
White, both sexes	241	-	-	-	-	-	-	-	-	1	2	3
Male	78	-	-	-	-	-	-	-	-	-	2	2
Female	163	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	25	-	-	-	-	-	-	-	-	-	1	3
Male	12	-	-	-	-	-	-	-	-	-	1	2
Female	13	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	77	-	-	-	-	-	-	-	1	-	4	2
Male	37	-	-	-	-	-	-	-	1	-	1	1
Female	40	-	-	-	-	-	-	-	-	-	3	1
White, both sexes	71	-	-	-	-	-	-	-	1	-	2	2
Male	33	-	-	-	-	-	-	-	1	-	-	1
Female	38	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Oregon												
All races, both sexes	86	-	-	-	-	-	-	-	-	-	-	-
Male	44	-	-	-	-	-	-	-	-	-	-	-
Female	42	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	85	-	-	-	-	-	-	-	-	-	-	-
Male	43	-	-	-	-	-	-	-	-	-	-	-
Female	42	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	377	-	-	-	-	-	1	2	-	3	3	10
Male	143	-	-	-	-	-	-	1	-	-	2	6
Female	234	-	-	-	-	-	1	1	-	3	1	4
White, both sexes	357	-	-	-	-	-	-	2	-	3	2	9
Male	133	-	-	-	-	-	-	1	-	-	1	6
Female	224	-	-	-	-	-	-	1	-	3	1	3
Black, both sexes	19	-	-	-	-	-	1	-	-	-	1	1
Male	9	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	1	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Rhode Island												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	-	2
Male	6	-	-	-	-	-	-	-	-	-	-	2
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	2
Male	6	-	-	-	-	-	-	-	-	-	-	2
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	75	-	-	-	-	-	-	-	2	2	-	1
Male	21	-	-	-	-	-	-	-	-	-	-	1
Female	54	-	-	-	-	-	-	-	2	2	-	-
White, both sexes	54	-	-	-	-	-	-	-	-	2	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	41	-	-	-	-	-	-	-	-	2	-	-
Black, both sexes	21	-	-	-	-	-	-	-	2	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
South Dakota												
All races, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	104	-	-	-	-	-	1	-	1	-	3	3
Male	50	-	-	-	-	-	-	-	1	-	3	2
Female	54	-	-	-	-	-	1	-	-	-	-	1
White, both sexes	95	-	-	-	-	-	1	-	1	-	1	3
Male	45	-	-	-	-	-	-	-	1	-	1	2
Female	50	-	-	-	-	-	1	-	-	-	-	1
Black, both sexes	9	-	-	-	-	-	-	-	-	-	2	-
Male	5	-	-	-	-	-	-	-	-	-	2	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Texas												
All races, both sexes	302	-	-	-	-	-	-	3	2	2	4	11
Male	125	-	-	-	-	-	-	3	1	2	3	4
Female	177	-	-	-	-	-	-	-	1	-	1	7
White, both sexes	259	-	-	-	-	-	-	3	1	2	3	10
Male	103	-	-	-	-	-	-	3	1	2	2	4
Female	156	-	-	-	-	-	-	-	-	-	1	6
Black, both sexes	39	-	-	-	-	-	-	-	1	-	1	1
Male	20	-	-	-	-	-	-	-	-	-	1	-
Female	19	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	48	-	-	-	-	-	-	-	-	1	-	1
Male	16	-	-	-	-	-	-	-	-	1	-	1
Female	32	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	46	-	-	-	-	-	-	-	-	1	-	1
Male	15	-	-	-	-	-	-	-	-	1	-	1
Female	31	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Vermont												
All races, both sexes	16	-	-	-	-	-	-	1	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	1	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	123	-	-	-	-	-	1	-	1	1	4	4
Male	44	-	-	-	-	-	1	-	-	-	2	2
Female	79	-	-	-	-	-	-	-	1	1	2	2
White, both sexes	94	-	-	-	-	-	-	-	-	-	-	3
Male	30	-	-	-	-	-	-	-	-	-	-	2
Female	64	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	28	-	-	-	-	-	1	-	1	1	4	1
Male	14	-	-	-	-	-	1	-	-	-	2	-
Female	14	-	-	-	-	-	-	-	1	1	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Washington												
All races, both sexes	179	-	-	-	-	-	-	-	3	2	-	2
Male	63	-	-	-	-	-	-	-	2	2	-	2
Female	116	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	173	-	-	-	-	-	-	-	3	2	-	2
Male	62	-	-	-	-	-	-	-	2	2	-	2
Female	111	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	30	-	-	-	-	-	-	-	-	-	1	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	19	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	29	-	-	-	-	-	-	-	-	-	1	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	18	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Wisconsin												
All races, both sexes	133	1	-	-	-	-	-	1	2	-	2	-
Male	53	1	-	-	-	-	-	-	1	-	2	-
Female	80	-	-	-	-	-	-	1	1	-	-	-
White, both sexes	127	1	-	-	-	-	-	1	2	-	2	-
Male	49	1	-	-	-	-	-	-	1	-	2	-
Female	78	-	-	-	-	-	-	1	1	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
United States												
All races, both sexes	213	230	276	322	400	632	872	1,069	795	299	32	-
Male	133	125	154	170	185	262	356	374	217	73	5	-
Female	80	105	122	152	215	370	516	695	578	226	27	-
White, both sexes	141	155	220	266	357	563	810	1,019	765	290	32	-
Male	89	79	122	142	166	230	330	356	206	70	5	-
Female	52	76	98	124	191	333	480	663	559	220	27	-
Black, both sexes	65	68	52	51	30	54	50	33	21	7	-	-
Male	39	43	30	25	12	26	21	8	7	2	-	-
Female	26	25	22	26	18	28	29	25	14	5	-	-
American Indian 1/, both sexes	3	3	1	2	2	3	3	2	1	-	-	-
Male	3	1	-	1	2	2	-	1	1	-	-	-
Female	-	2	1	1	-	1	3	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	4	3	3	11	12	9	15	8	2	-	-
Male	2	2	2	2	5	4	5	9	3	1	-	-
Female	2	2	1	1	6	8	4	6	5	1	-	-
Alabama												
All races, both sexes	4	7	5	8	6	10	15	13	10	2	-	-
Male	2	4	2	5	3	5	7	4	3	-	-	-
Female	2	3	3	3	3	5	8	9	7	2	-	-
White, both sexes	2	3	3	3	6	9	12	11	8	2	-	-
Male	-	1	-	3	3	4	5	2	2	-	-	-
Female	2	2	3	-	3	5	7	9	6	2	-	-
Black, both sexes	2	4	2	5	-	1	3	2	2	-	-	-
Male	2	3	2	2	-	1	2	2	1	-	-	-
Female	-	1	-	3	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Alaska												
All races, both sexes	3	-	-	-	3	2	-	1	3	-	-	-
Male	2	-	-	-	2	1	-	1	-	-	-	-
Female	1	-	-	-	1	1	-	-	3	-	-	-
White, both sexes	-	-	-	-	3	2	-	1	3	-	-	-
Male	-	-	-	-	2	1	-	1	-	-	-	-
Female	-	-	-	-	1	1	-	-	3	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	4	5	6	6	6	12	15	19	13	9	1	-
Male	3	5	1	4	3	4	10	10	2	3	-	-
Female	1	-	5	2	3	8	5	9	11	6	1	-
White, both sexes	4	4	6	5	6	12	15	18	13	9	1	-
Male	3	4	1	3	3	4	10	10	2	3	-	-
Female	1	-	5	2	3	8	5	8	11	6	1	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Arkansas												
All races, both sexes	2	4	5	6	2	6	6	6	7	4	-	-
Male	-	2	2	4	1	1	1	2	2	-	-	-
Female	2	2	3	2	1	5	5	4	5	4	-	-
White, both sexes	1	4	4	5	2	6	6	5	7	4	-	-
Male	-	2	1	4	1	1	1	2	2	-	-	-
Female	1	2	3	1	1	5	5	3	5	4	-	-
Black, both sexes	1	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	15	18	17	15	29	62	69	83	63	16	3	-
Male	11	8	12	9	12	22	31	31	19	3	-	-
Female	4	10	5	6	17	40	38	52	44	13	3	-
White, both sexes	10	13	13	13	20	50	62	74	58	15	3	-
Male	9	6	9	7	8	15	28	24	16	3	-	-
Female	1	7	4	6	12	35	34	50	42	12	3	-
Black, both sexes	5	5	3	2	3	6	2	2	3	-	-	-
Male	2	2	2	2	2	4	-	1	3	-	-	-
Female	3	3	1	-	1	2	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	6	5	4	6	2	1	-	-
Male	-	-	1	-	2	2	3	5	-	-	-	-
Female	-	-	-	-	4	3	1	1	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Colorado												
All races, both sexes	1	-	4	4	3	10	13	16	13	6	1	-
Male	1	-	3	3	1	4	7	6	3	1	-	-
Female	-	-	1	1	2	6	6	10	10	5	1	-
White, both sexes	1	-	3	4	3	10	12	16	13	6	1	-
Male	1	-	2	3	1	4	7	6	3	1	-	-
Female	-	-	1	1	2	6	5	10	10	5	1	-
Black, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	2	5	6	4	1	4	10	17	7	3	-	-
Male	1	4	3	2	-	1	4	5	3	-	-	-
Female	1	1	3	2	1	3	6	12	4	3	-	-
White, both sexes	2	4	5	3	1	4	10	16	7	3	-	-
Male	1	3	3	1	-	1	4	5	3	-	-	-
Female	1	1	2	2	1	3	6	11	4	3	-	-
Black, both sexes	-	1	1	1	-	-	-	1	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Delaware												
All races, both sexes	-	1	3	1	-	2	3	1	3	-	-	-
Male	-	1	2	1	-	1	1	-	2	-	-	-
Female	-	-	1	-	-	1	2	1	1	-	-	-
White, both sexes	-	-	2	-	-	2	3	1	3	-	-	-
Male	-	-	1	-	-	1	1	-	2	-	-	-
Female	-	-	1	-	-	1	2	1	1	-	-	-
Black, both sexes	-	1	1	1	-	-	-	-	-	-	-	-
Male	-	1	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	3	2	-	1	2	-	-	1	-	-	-	-
Male	2	1	-	1	2	-	-	1	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	1	-	1	2	-	-	1	-	-	-	-
Male	2	1	-	1	2	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Florida												
All races, both sexes	16	20	19	35	42	52	75	96	64	25	3	-
Male	6	8	7	24	19	25	31	42	18	11	1	-
Female	10	12	12	11	23	27	44	54	46	14	2	-
White, both sexes	11	15	12	29	38	49	66	95	63	25	3	-
Male	4	6	5	20	18	24	26	42	18	11	1	-
Female	7	9	7	9	20	25	40	53	45	14	2	-
Black, both sexes	5	5	7	5	4	3	9	1	1	-	-	-
Male	2	2	2	4	1	1	5	-	-	-	-	-
Female	3	3	5	1	3	2	4	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	7	6	6	6	5	17	17	15	16	5	-	-
Male	5	6	4	3	4	9	7	6	3	3	-	-
Female	2	-	2	3	1	8	10	9	13	2	-	-
White, both sexes	5	3	2	6	5	15	13	13	14	3	-	-
Male	3	3	1	3	4	8	4	6	3	2	-	-
Female	2	-	1	3	1	7	9	7	11	1	-	-
Black, both sexes	2	3	4	-	-	2	4	2	2	2	-	-
Male	2	3	3	-	-	1	3	-	-	1	-	-
Female	-	-	1	-	-	1	1	2	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Hawaii												
All races, both sexes	1	3	-	-	1	1	3	6	2	2	-	-
Male	-	2	-	-	1	1	2	4	2	1	-	-
Female	1	1	-	-	-	-	1	2	-	1	-	-
White, both sexes	-	1	-	-	-	-	2	2	-	1	-	-
Male	-	1	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	-	-	1	1	1	4	2	1	-	-
Male	-	1	-	-	1	1	1	3	2	1	-	-
Female	1	1	-	-	-	-	-	1	-	-	-	-
Idaho												
All races, both sexes	-	1	-	4	3	3	10	10	2	2	-	-
Male	-	-	-	2	2	2	2	4	-	-	-	-
Female	-	1	-	2	1	1	8	6	2	2	-	-
White, both sexes	-	1	-	4	3	3	10	10	2	2	-	-
Male	-	-	-	2	2	2	2	4	-	-	-	-
Female	-	1	-	2	1	1	8	6	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Illinois												
All races, both sexes	6	11	5	13	20	22	26	29	27	8	-	-
Male	5	9	4	8	10	10	9	12	6	2	-	-
Female	1	2	1	5	10	12	17	17	21	6	-	-
White, both sexes	4	9	3	10	19	17	23	27	26	8	-	-
Male	4	7	2	7	10	8	7	11	5	2	-	-
Female	-	2	1	3	9	9	16	16	21	6	-	-
Black, both sexes	2	2	2	3	1	4	3	2	1	-	-	-
Male	1	2	2	1	-	1	2	1	1	-	-	-
Female	1	-	-	2	1	3	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	5	3	10	6	8	11	23	20	19	8	1	-
Male	4	3	8	-	5	3	11	10	7	4	-	-
Female	1	-	2	6	3	8	12	10	12	4	1	-
White, both sexes	4	3	9	5	8	11	22	20	19	8	1	-
Male	3	3	7	-	5	3	10	10	7	4	-	-
Female	1	-	2	5	3	8	12	10	12	4	1	-
Black, both sexes	1	-	1	1	-	-	1	-	-	-	-	-
Male	1	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Iowa												
All races, both sexes	1	4	1	-	3	4	4	5	5	2	-	-
Male	1	2	1	-	1	3	1	2	1	1	-	-
Female	-	2	-	-	2	1	3	3	4	1	-	-
White, both sexes	1	4	1	-	3	4	4	5	5	2	-	-
Male	1	2	1	-	1	3	1	2	1	1	-	-
Female	-	2	-	-	2	1	3	3	4	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	1	-	1	1	9	10	13	8	3	1	-
Male	-	1	-	-	-	2	4	3	1	-	-	-
Female	1	-	-	1	1	7	6	10	7	3	1	-
White, both sexes	1	1	-	1	1	9	9	13	8	3	1	-
Male	-	1	-	-	-	2	4	3	1	-	-	-
Female	1	-	-	1	1	7	5	10	7	3	1	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Kentucky												
All races, both sexes	3	3	3	4	4	9	18	9	10	3	1	-
Male	3	1	1	2	1	5	5	-	1	-	-	-
Female	-	2	2	2	3	4	13	9	9	3	1	-
White, both sexes	3	3	3	4	4	9	17	9	9	3	1	-
Male	3	1	1	2	1	5	5	-	1	-	-	-
Female	-	2	2	2	3	4	12	9	8	3	1	-
Black, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	3	2	3	11	4	7	22	17	9	4	-	-
Male	2	1	1	7	2	1	10	4	2	-	-	-
Female	1	1	2	4	2	6	12	13	7	4	-	-
White, both sexes	-	2	-	7	4	5	19	16	7	4	-	-
Male	-	1	-	4	2	-	9	4	2	-	-	-
Female	-	1	-	3	2	5	10	12	5	4	-	-
Black, both sexes	3	-	3	4	-	2	3	1	1	-	-	-
Male	2	-	1	3	-	1	1	-	-	-	-	-
Female	1	-	2	1	-	1	2	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Maine												
All races, both sexes	-	1	-	3	2	4	4	11	12	3	-	-
Male	-	1	-	-	2	1	1	2	2	1	-	-
Female	-	-	-	3	-	3	3	9	10	2	-	-
White, both sexes	-	1	-	3	2	4	4	11	12	3	-	-
Male	-	1	-	-	2	1	1	2	2	1	-	-
Female	-	-	-	3	-	3	3	9	10	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	6	3	6	5	4	5	15	15	8	2	2	-
Male	4	2	4	1	3	3	7	6	4	1	-	-
Female	2	1	2	4	1	2	8	9	4	1	2	-
White, both sexes	1	2	5	-	2	3	15	12	8	1	2	-
Male	1	1	3	-	2	2	7	6	4	-	-	-
Female	-	1	2	-	-	1	8	6	4	1	2	-
Black, both sexes	5	1	1	5	1	2	-	2	-	1	-	-
Male	3	1	1	1	1	1	-	-	-	1	-	-
Female	2	-	-	4	-	1	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Massachusetts												
All races, both sexes	2	2	4	3	4	8	15	28	16	5	1	-
Male	1	2	2	1	3	4	7	10	4	-	-	-
Female	1	-	2	2	1	4	8	18	12	5	1	-
White, both sexes	2	1	4	3	4	7	14	25	16	5	1	-
Male	1	1	2	1	3	4	7	9	4	-	-	-
Female	1	-	2	2	1	3	7	16	12	5	1	-
Black, both sexes	-	1	-	-	-	1	-	2	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Michigan												
All races, both sexes	11	10	11	11	9	25	34	27	27	11	2	-
Male	9	7	8	5	3	13	14	10	9	6	-	-
Female	2	3	3	6	6	12	20	17	18	5	2	-
White, both sexes	4	4	9	10	8	19	33	25	27	11	2	-
Male	4	4	7	5	2	10	14	9	9	6	-	-
Female	-	-	2	5	6	9	19	16	18	5	2	-
Black, both sexes	7	6	2	1	-	6	-	2	-	-	-	-
Male	5	3	1	-	-	3	-	1	-	-	-	-
Female	2	3	1	1	-	3	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Minnesota												
All races, both sexes	-	4	2	3	5	5	14	16	9	3	-	-
Male	-	2	-	3	3	-	6	10	4	1	-	-
Female	-	2	2	-	2	5	8	6	5	2	-	-
White, both sexes	-	3	2	3	5	4	14	16	9	3	-	-
Male	-	2	-	3	3	-	6	10	4	1	-	-
Female	-	1	2	-	2	4	8	6	5	2	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	3	2	5	2	5	2	9	3	1	1	-	-
Male	2	1	2	2	3	1	3	1	-	-	-	-
Female	1	1	3	-	2	1	6	2	1	1	-	-
White, both sexes	1	1	5	1	4	-	8	2	1	-	-	-
Male	1	-	2	1	2	-	3	1	-	-	-	-
Female	-	1	3	-	2	-	5	1	1	-	-	-
Black, both sexes	2	1	-	1	1	2	1	1	-	1	-	-
Male	1	1	-	1	1	1	-	-	-	-	-	-
Female	1	-	-	-	-	1	1	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Missouri												
All races, both sexes	4	2	6	11	13	18	17	21	21	4	-	-
Male	2	-	4	5	4	7	5	7	8	-	-	-
Female	2	2	2	6	9	11	12	14	13	4	-	-
White, both sexes	4	1	4	11	10	17	16	21	20	4	-	-
Male	2	-	2	5	3	6	5	7	8	-	-	-
Female	2	1	2	6	7	11	11	14	12	4	-	-
Black, both sexes	-	1	2	-	3	1	1	-	1	-	-	-
Male	-	-	2	-	1	1	-	-	-	-	-	-
Female	-	1	-	-	2	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	2	3	-	4	2	9	4	5	1	-	-
Male	-	1	2	-	2	1	3	3	2	1	-	-
Female	-	1	1	-	2	1	6	1	3	-	-	-
White, both sexes	-	2	3	-	4	2	9	4	5	1	-	-
Male	-	1	2	-	2	1	3	3	2	1	-	-
Female	-	1	1	-	2	1	6	1	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Nebraska												
All races, both sexes	2	1	-	1	2	1	3	21	10	4	-	-
Male	1	1	-	1	1	1	-	7	3	3	-	-
Female	1	-	-	-	1	-	3	14	7	1	-	-
White, both sexes	2	1	-	1	2	1	3	21	10	4	-	-
Male	1	1	-	1	1	1	-	7	3	3	-	-
Female	1	-	-	-	1	-	3	14	7	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	2	3	5	5	2	4	5	-	4	1	1	-
Male	2	2	4	3	2	1	2	-	2	-	-	-
Female	-	1	1	2	-	3	3	-	2	1	1	-
White, both sexes	2	3	5	5	2	2	4	-	4	1	1	-
Male	2	2	4	3	2	1	2	-	2	-	-	-
Female	-	1	1	2	-	1	2	-	2	1	1	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
New Hampshire												
All races, both sexes	-	1	-	-	2	3	5	12	8	3	-	-
Male	-	1	-	-	-	1	3	5	2	1	-	-
Female	-	-	-	-	2	2	2	7	6	2	-	-
White, both sexes	-	1	-	-	2	3	5	12	8	3	-	-
Male	-	1	-	-	-	1	3	5	2	1	-	-
Female	-	-	-	-	2	2	2	7	6	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	7	6	7	9	10	25	40	65	36	14	2	-
Male	4	1	4	6	7	8	17	24	9	1	1	-
Female	3	5	3	3	3	17	23	41	27	13	1	-
White, both sexes	7	3	6	6	10	22	38	63	36	14	2	-
Male	4	-	3	4	7	6	16	24	9	1	1	-
Female	3	3	3	2	3	16	22	39	27	13	1	-
Black, both sexes	-	3	1	3	-	3	2	1	-	-	-	-
Male	-	1	1	2	-	2	1	-	-	-	-	-
Female	-	2	-	1	-	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
New Mexico												
All races, both sexes	3	2	2	4	3	3	5	4	5	1	-	-
Male	2	1	2	1	2	1	1	1	1	-	-	-
Female	1	1	-	3	1	2	4	3	4	1	-	-
White, both sexes	3	-	2	4	3	3	5	4	5	1	-	-
Male	2	-	2	1	2	1	1	1	1	-	-	-
Female	1	-	-	3	1	2	4	3	4	1	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	13	15	16	20	32	54	42	63	53	28	1	-
Male	10	6	10	6	12	25	20	24	11	7	-	-
Female	3	9	6	14	20	29	22	39	42	21	1	-
White, both sexes	10	10	14	17	28	48	41	61	52	28	1	-
Male	8	5	9	5	11	24	20	24	11	7	-	-
Female	2	5	5	12	17	24	21	37	41	21	1	-
Black, both sexes	1	4	2	2	3	4	-	2	1	-	-	-
Male	-	1	1	-	1	1	-	-	-	-	-	-
Female	1	3	1	2	2	3	-	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	-	1	1	2	1	-	-	-	-	-
Male	2	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	2	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
North Carolina												
All races, both sexes	11	5	13	8	15	17	28	26	21	5	-	-
Male	5	1	6	3	9	4	11	9	3	1	-	-
Female	6	4	7	5	6	13	17	17	18	4	-	-
White, both sexes	3	3	9	6	13	14	23	26	19	5	-	-
Male	2	1	4	2	8	3	10	9	2	1	-	-
Female	1	2	5	4	5	11	13	17	17	4	-	-
Black, both sexes	8	2	4	2	2	3	4	-	1	-	-	-
Male	3	-	2	1	1	1	1	-	-	-	-	-
Female	5	2	2	1	1	2	3	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
North Dakota												
All races, both sexes	2	1	2	-	-	-	3	5	2	-	-	-
Male	2	1	2	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	2	4	2	-	-	-
White, both sexes	1	1	2	-	-	-	3	5	2	-	-	-
Male	1	1	2	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	2	4	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Ohio												
All races, both sexes	9	12	14	12	17	28	51	51	43	15	4	-
Male	3	7	5	7	5	8	21	14	8	4	1	-
Female	6	5	9	5	12	20	30	37	35	11	3	-
White, both sexes	6	7	12	10	16	28	47	49	41	15	4	-
Male	2	3	4	7	5	8	19	14	7	4	1	-
Female	4	4	8	3	11	20	28	35	34	11	3	-
Black, both sexes	3	5	2	2	1	-	4	2	2	-	-	-
Male	1	4	1	-	-	-	2	-	1	-	-	-
Female	2	1	1	2	1	-	2	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	3	4	5	7	7	11	13	12	7	1	-	-
Male	3	2	2	4	3	5	7	4	4	-	-	-
Female	-	2	3	3	4	6	6	8	3	1	-	-
White, both sexes	2	4	5	6	6	10	13	12	7	1	-	-
Male	2	2	2	4	2	4	7	4	4	-	-	-
Female	-	2	3	2	4	6	6	8	3	1	-	-
Black, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Oregon												
All races, both sexes	1	1	4	7	4	11	15	23	15	5	-	-
Male	1	-	1	5	2	7	11	8	7	2	-	-
Female	-	1	3	2	2	4	4	15	8	3	-	-
White, both sexes	-	1	4	7	4	11	15	23	15	5	-	-
Male	-	-	1	5	2	7	11	8	7	2	-	-
Female	-	1	3	2	2	4	4	15	8	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	12	7	22	20	24	40	54	83	63	31	2	-
Male	6	3	13	13	12	20	20	19	21	7	-	-
Female	6	4	9	7	12	20	34	64	42	24	2	-
White, both sexes	10	5	18	19	23	40	50	80	63	31	2	-
Male	4	2	11	13	11	20	18	18	21	7	-	-
Female	6	3	7	6	12	20	32	62	42	24	2	-
Black, both sexes	2	2	4	1	-	-	4	3	-	-	-	-
Male	2	1	2	-	-	-	2	1	-	-	-	-
Female	-	1	2	1	-	-	2	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Rhode Island												
All races, both sexes	-	-	2	-	2	2	3	5	2	-	-	-
Male	-	-	2	-	1	-	-	-	1	-	-	-
Female	-	-	-	-	1	2	3	5	1	-	-	-
White, both sexes	-	-	2	-	2	2	2	5	2	-	-	-
Male	-	-	2	-	1	-	-	-	1	-	-	-
Female	-	-	-	-	1	2	2	5	1	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	5	3	3	6	5	10	11	13	12	2	-	-
Male	2	2	1	3	1	3	2	3	2	1	-	-
Female	3	1	2	3	4	7	9	10	10	1	-	-
White, both sexes	3	2	2	3	2	6	9	13	10	2	-	-
Male	1	1	1	1	1	1	1	3	2	1	-	-
Female	2	1	1	2	1	5	8	10	8	1	-	-
Black, both sexes	2	1	1	3	3	4	2	-	2	-	-	-
Male	1	1	-	2	-	2	1	-	-	-	-	-
Female	1	-	1	1	3	2	1	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
South Dakota												
All races, both sexes	1	1	1	-	3	1	3	4	5	3	-	-
Male	1	1	-	-	1	1	-	1	2	-	-	-
Female	-	-	1	-	2	-	3	3	3	3	-	-
White, both sexes	1	1	-	-	3	1	3	4	5	3	-	-
Male	1	1	-	-	1	1	-	1	2	-	-	-
Female	-	-	-	-	2	-	3	3	3	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	5	3	8	10	6	18	15	15	14	2	-	-
Male	4	2	4	6	3	9	8	5	2	1	-	-
Female	1	1	4	4	3	9	7	10	12	1	-	-
White, both sexes	4	1	7	9	6	17	15	14	14	2	-	-
Male	3	-	4	6	3	9	8	5	2	1	-	-
Female	1	1	3	3	3	8	7	9	12	1	-	-
Black, both sexes	1	2	1	1	-	1	-	1	-	-	-	-
Male	1	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Texas												
All races, both sexes	16	21	19	20	36	38	43	47	30	10	-	-
Male	7	10	12	9	15	15	18	18	8	-	-	-
Female	9	11	7	11	21	23	25	29	22	10	-	-
White, both sexes	12	13	13	16	31	34	42	43	29	7	-	-
Male	4	5	8	6	12	13	17	18	8	-	-	-
Female	8	8	5	10	19	21	25	25	21	7	-	-
Black, both sexes	4	8	6	4	4	4	-	3	-	3	-	-
Male	3	5	4	3	2	2	-	-	-	-	-	-
Female	1	3	2	1	2	2	-	3	-	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	1	1	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Utah												
All races, both sexes	2	4	1	4	9	8	7	8	3	-	-	-
Male	2	2	1	1	2	2	1	2	1	-	-	-
Female	-	2	-	3	7	6	6	6	2	-	-	-
White, both sexes	2	3	1	4	9	8	7	8	2	-	-	-
Male	2	2	1	1	2	2	1	2	-	-	-	-
Female	-	1	-	3	7	6	6	6	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Vermont												
All races, both sexes	-	1	3	-	-	2	2	2	2	2	1	-
Male	-	-	3	-	-	-	1	1	1	-	1	-
Female	-	1	-	-	-	2	1	1	1	2	-	-
White, both sexes	-	1	3	-	-	2	2	2	2	2	1	-
Male	-	-	3	-	-	-	1	1	1	-	1	-
Female	-	1	-	-	-	2	1	1	1	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	3	10	7	7	8	15	19	22	19	1	1	-
Male	2	6	6	1	5	6	4	4	5	-	-	-
Female	1	4	1	6	3	9	15	18	14	1	1	-
White, both sexes	-	3	7	6	6	11	18	21	17	1	1	-
Male	-	-	6	1	5	3	4	4	5	-	-	-
Female	-	3	1	5	1	8	14	17	12	1	1	-
Black, both sexes	3	7	-	1	2	4	1	-	2	-	-	-
Male	2	6	-	-	-	3	-	-	-	-	-	-
Female	1	1	-	1	2	1	1	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Washington												
All races, both sexes	4	3	8	9	13	16	25	37	27	26	4	-
Male	3	-	2	5	5	10	8	12	6	5	1	-
Female	1	3	6	4	8	6	17	25	21	21	3	-
White, both sexes	4	3	7	8	13	14	24	37	26	26	4	-
Male	3	-	2	4	5	10	8	12	6	5	1	-
Female	1	3	5	4	8	4	16	25	20	21	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	-	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
West Virginia												
All races, both sexes	2	-	1	3	2	4	6	6	2	2	-	-
Male	1	-	-	2	1	1	3	1	1	-	-	-
Female	1	-	1	1	1	3	3	5	1	2	-	-
White, both sexes	2	-	1	3	2	4	5	6	2	2	-	-
Male	1	-	-	2	1	1	3	1	1	-	-	-
Female	1	-	1	1	1	3	2	5	1	2	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Wisconsin												
All races, both sexes	7	3	3	6	9	7	15	37	30	10	-	-
Male	6	2	1	2	4	4	7	14	8	1	-	-
Female	1	1	2	4	5	3	8	23	22	9	-	-
White, both sexes	6	2	2	5	9	7	15	36	29	10	-	-
Male	5	1	1	2	4	4	7	13	7	1	-	-
Female	1	1	1	3	5	3	8	23	22	9	-	-
Black, both sexes	1	-	1	-	-	-	-	1	1	-	-	-
Male	1	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	1	-	1	-	2	3	6	2	1	-	-
Male	-	-	-	-	-	-	1	1	1	-	-	-
Female	-	1	-	1	-	2	2	5	1	1	-	-
White, both sexes	-	1	-	1	-	2	3	6	2	1	-	-
Male	-	-	-	-	-	-	1	1	1	-	-	-
Female	-	1	-	1	-	2	2	5	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
United States												
All races, both sexes	24,267	120	38	24	33	87	186	264	332	555	775	1,141
Male	14,410	60	17	13	20	69	136	200	231	404	557	818
Female	9,857	60	21	11	13	18	50	64	101	151	218	323
White, both sexes	19,420	74	29	19	24	49	113	150	206	334	502	764
Male	11,492	35	13	10	15	37	84	120	152	246	361	581
Female	7,928	39	16	9	9	12	29	30	54	88	141	183
Black, both sexes	4,243	38	5	5	7	34	67	99	112	197	244	341
Male	2,525	21	2	3	3	28	47	68	68	139	174	212
Female	1,718	17	3	2	4	6	20	31	44	58	70	129
American Indian 1/, both sexes	138	1	1	-	-	2	1	2	7	7	11	20
Male	89	-	1	-	-	2	1	2	7	5	9	15
Female	49	1	-	-	-	-	-	-	-	2	2	5
Asian or Pacific Islander 2/, both sexes	466	7	3	-	2	2	5	13	7	17	18	16
Male	304	4	1	-	2	2	4	10	4	14	13	10
Female	162	3	2	-	-	-	1	3	3	3	5	6
Alabama												
All races, both sexes	393	3	-	-	-	2	4	3	5	11	11	14
Male	218	1	-	-	-	-	3	2	2	6	6	12
Female	175	2	-	-	-	2	1	1	3	5	5	2
White, both sexes	269	2	-	-	-	1	3	3	1	5	6	7
Male	147	-	-	-	-	-	2	2	1	3	3	6
Female	122	2	-	-	-	1	1	1	-	2	3	1
Black, both sexes	123	1	-	-	-	1	1	-	4	6	5	7
Male	71	1	-	-	-	-	1	-	1	3	3	6
Female	52	-	-	-	-	1	-	-	3	3	2	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Alaska												
All races, both sexes	19	-	-	1	-	-	-	-	2	-	4	-
Male	12	-	-	-	-	-	-	-	2	-	4	-
Female	7	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	1	-	-	-	-	1	-	3	-
Male	6	-	-	-	-	-	-	-	1	-	3	-
Female	6	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	1	-	-	-
Male	5	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	324	4	1	-	1	1	2	5	8	7	16	25
Male	213	2	-	-	1	1	1	3	7	5	13	20
Female	111	2	1	-	-	-	1	2	1	2	3	5
White, both sexes	303	3	1	-	1	1	2	5	7	7	13	25
Male	201	2	-	-	1	1	1	3	6	5	11	20
Female	102	1	1	-	-	-	1	2	1	2	2	5
Black, both sexes	11	1	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	1	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	1	-	1	-
Male	5	-	-	-	-	-	-	-	1	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Arkansas												
All races, both sexes	229	1	1	-	-	3	4	2	2	3	7	7
Male	118	-	1	-	-	2	3	2	1	3	6	4
Female	111	1	-	-	-	1	1	-	1	-	1	3
White, both sexes	186	1	1	-	-	2	2	-	-	3	3	5
Male	96	-	1	-	-	1	1	-	-	3	2	3
Female	90	1	-	-	-	1	1	-	-	-	1	2
Black, both sexes	42	-	-	-	-	1	1	2	2	-	4	2
Male	21	-	-	-	-	1	1	2	1	-	4	1
Female	21	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	2,770	15	8	2	4	10	32	47	50	74	104	168
Male	1,736	10	5	-	2	8	25	36	40	57	80	131
Female	1,034	5	3	2	2	2	7	11	10	17	24	37
White, both sexes	2,221	12	7	2	3	5	27	27	34	55	68	125
Male	1,387	7	5	-	2	5	20	23	29	40	53	103
Female	834	5	2	2	1	-	7	4	5	15	15	22
Black, both sexes	346	2	1	-	1	3	4	11	12	12	27	33
Male	207	2	-	-	-	1	4	5	7	11	20	22
Female	139	-	1	-	1	2	-	6	5	1	7	11
American Indian 1/, both sexes	20	-	-	-	-	-	-	1	1	-	3	3
Male	15	-	-	-	-	-	-	1	1	-	2	2
Female	5	-	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	183	1	-	-	-	2	1	8	3	7	6	7
Male	127	1	-	-	-	2	1	7	3	6	5	4
Female	56	-	-	-	-	-	-	1	-	1	1	3

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Colorado												
All races, both sexes	238	2	1	-	-	1	3	2	4	9	12	20
Male	155	-	1	-	-	1	2	2	2	7	10	17
Female	83	2	-	-	-	-	1	-	2	2	2	3
White, both sexes	223	1	1	-	-	-	3	2	3	8	11	18
Male	145	-	1	-	-	-	2	2	1	7	10	15
Female	78	1	-	-	-	-	1	-	2	1	1	3
Black, both sexes	11	1	-	-	-	-	-	-	1	1	-	1
Male	7	-	-	-	-	-	-	-	1	-	-	1
Female	4	1	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	3	-	-	-	-	1	-	-	-	-	-	1
Male	3	-	-	-	-	1	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Connecticut												
All races, both sexes	320	1	-	1	-	1	-	-	6	7	10	15
Male	211	1	-	1	-	1	-	-	5	7	10	9
Female	109	-	-	-	-	-	-	-	1	-	-	6
White, both sexes	286	1	-	1	-	1	-	-	2	6	8	12
Male	189	1	-	1	-	1	-	-	2	6	8	8
Female	97	-	-	-	-	-	-	-	-	-	-	4
Black, both sexes	32	-	-	-	-	-	-	-	4	1	2	3
Male	21	-	-	-	-	-	-	-	3	1	2	1
Female	11	-	-	-	-	-	-	-	1	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Delaware												
All races, both sexes	147	-	1	-	-	1	-	-	2	6	10	6
Male	76	-	-	-	-	1	-	-	-	4	7	4
Female	71	-	1	-	-	-	-	-	2	2	3	2
White, both sexes	123	-	-	-	-	1	-	-	1	6	6	4
Male	64	-	-	-	-	1	-	-	-	4	4	4
Female	59	-	-	-	-	-	-	-	1	2	2	-
Black, both sexes	24	-	1	-	-	-	-	-	1	-	4	2
Male	12	-	-	-	-	-	-	-	-	-	3	-
Female	12	-	1	-	-	-	-	-	1	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	50	-	-	-	-	-	1	-	-	3	2	2
Male	24	-	-	-	-	-	1	-	-	2	2	1
Female	26	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	11	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	39	-	-	-	-	-	1	-	-	3	1	2
Male	19	-	-	-	-	-	1	-	-	2	1	1
Female	20	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Florida												
All races, both sexes	2,096	13	-	4	4	2	11	8	22	39	51	62
Male	1,281	8	-	2	2	2	10	5	17	31	38	42
Female	815	5	-	2	2	-	1	3	5	8	13	20
White, both sexes	1,760	4	-	3	2	1	5	2	13	23	38	43
Male	1,080	1	-	2	-	1	4	1	10	20	27	31
Female	680	3	-	1	2	-	1	1	3	3	11	12
Black, both sexes	315	8	-	1	1	1	6	6	8	15	12	17
Male	186	6	-	-	1	1	6	4	6	10	11	9
Female	129	2	-	1	-	-	-	2	2	5	1	8
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	1	-	-	2
Male	5	-	-	-	-	-	-	-	1	-	-	2
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	1	-	-	1	-	-	-	-	1	1	-
Male	10	1	-	-	1	-	-	-	-	1	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
Georgia												
All races, both sexes	770	6	4	1	-	4	8	5	10	27	25	33
Male	462	3	-	1	-	3	6	4	5	21	16	24
Female	308	3	4	-	-	1	2	1	5	6	9	9
White, both sexes	496	1	3	-	-	1	1	1	3	13	10	15
Male	288	1	-	-	-	-	-	1	1	10	5	12
Female	208	-	3	-	-	1	1	-	2	3	5	3
Black, both sexes	268	5	-	1	-	3	6	4	7	13	15	18
Male	170	2	-	1	-	3	5	3	4	10	11	12
Female	98	3	-	-	-	-	1	1	3	3	4	6
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	1	-	-	-	1	-	-	1	-	-
Male	4	-	-	-	-	-	1	-	-	1	-	-
Female	2	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Hawaii												
All races, both sexes	198	1	-	-	-	-	1	-	3	6	6	9
Male	140	-	-	-	-	-	1	-	1	4	5	6
Female	58	1	-	-	-	-	-	-	2	2	1	3
White, both sexes	53	1	-	-	-	-	-	-	-	-	3	1
Male	37	-	-	-	-	-	-	-	-	-	3	1
Female	16	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	144	-	-	-	-	-	1	-	3	6	3	8
Male	103	-	-	-	-	-	1	-	1	4	2	5
Female	41	-	-	-	-	-	-	-	2	2	1	3
Idaho												
All races, both sexes	79	1	-	-	-	-	-	-	-	2	3	1
Male	47	1	-	-	-	-	-	-	-	1	2	1
Female	32	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	73	1	-	-	-	-	-	-	-	2	2	1
Male	43	1	-	-	-	-	-	-	-	1	1	1
Female	30	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Illinois												
All races, both sexes	1,052	4	1	-	-	5	7	18	16	32	34	53
Male	634	1	-	-	-	4	6	14	12	20	28	45
Female	418	3	1	-	-	1	1	4	4	12	6	8
White, both sexes	795	3	1	-	-	3	6	11	10	11	16	29
Male	472	1	-	-	-	2	6	8	8	10	14	25
Female	323	2	1	-	-	1	-	3	2	1	2	4
Black, both sexes	243	1	-	-	-	2	1	6	6	20	17	23
Male	154	-	-	-	-	2	-	6	4	10	13	19
Female	89	1	-	-	-	-	1	-	2	10	4	4
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	1	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	1	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	1	-	-	-	-
Indiana												
All races, both sexes	579	1	-	-	1	3	9	4	5	10	21	31
Male	353	1	-	-	1	2	6	2	1	7	12	28
Female	226	-	-	-	-	1	3	2	4	3	9	3
White, both sexes	520	1	-	-	1	2	5	2	4	10	17	27
Male	317	1	-	-	1	2	4	-	1	7	9	25
Female	203	-	-	-	-	-	1	2	3	3	8	2
Black, both sexes	57	-	-	-	-	1	4	2	1	-	3	4
Male	35	-	-	-	-	-	2	2	-	-	2	3
Female	22	-	-	-	-	1	2	-	1	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Iowa												
All races, both sexes	203	-	1	-	-	1	1	3	2	6	2	8
Male	121	-	1	-	-	1	1	3	2	4	2	6
Female	82	-	-	-	-	-	-	-	-	2	-	2
White, both sexes	195	-	1	-	-	1	1	2	2	5	2	7
Male	115	-	1	-	-	1	1	2	2	3	2	5
Female	80	-	-	-	-	-	-	-	-	2	-	2
Black, both sexes	7	-	-	-	-	-	-	1	-	-	-	1
Male	5	-	-	-	-	-	-	1	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	202	-	-	-	-	-	2	2	2	1	2	12
Male	126	-	-	-	-	-	1	2	2	1	1	9
Female	76	-	-	-	-	-	1	-	-	-	1	3
White, both sexes	177	-	-	-	-	-	1	2	2	-	1	10
Male	110	-	-	-	-	-	1	2	2	-	1	9
Female	67	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	23	-	-	-	-	-	1	-	-	1	1	2
Male	16	-	-	-	-	-	-	-	-	1	-	-
Female	7	-	-	-	-	-	1	-	-	-	1	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Kentucky												
All races, both sexes	306	1	-	-	-	-	1	3	5	2	5	12
Male	176	-	-	-	-	-	-	3	2	1	4	6
Female	130	1	-	-	-	-	1	-	3	1	1	6
White, both sexes	270	1	-	-	-	-	1	3	2	-	2	6
Male	151	-	-	-	-	-	-	3	1	-	1	2
Female	119	1	-	-	-	-	1	-	1	-	1	4
Black, both sexes	35	-	-	-	-	-	-	-	3	2	3	6
Male	24	-	-	-	-	-	-	-	1	1	3	4
Female	11	-	-	-	-	-	-	-	2	1	-	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	476	2	3	1	2	2	5	4	5	9	14	14
Male	259	-	3	1	2	1	3	2	4	7	11	5
Female	217	2	-	-	-	1	2	2	1	2	3	9
White, both sexes	323	-	-	-	1	-	1	1	1	5	7	8
Male	171	-	-	-	1	-	1	1	1	4	5	2
Female	152	-	-	-	-	-	-	-	-	1	2	6
Black, both sexes	150	2	2	1	1	2	4	3	4	4	7	6
Male	86	-	2	1	1	1	2	1	3	3	6	3
Female	64	2	-	-	-	1	2	2	1	1	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	1	-	-	-	-	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Maine												
All races, both sexes	114	1	-	1	1	-	2	-	1	2	-	3
Male	64	-	-	-	1	-	2	-	-	1	-	1
Female	50	1	-	1	-	-	-	-	1	1	-	2
White, both sexes	113	1	-	1	1	-	2	-	1	2	-	3
Male	63	-	-	-	1	-	2	-	-	1	-	1
Female	50	1	-	1	-	-	-	-	1	1	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	516	2	-	-	-	1	4	3	4	8	15	16
Male	292	1	-	-	-	1	3	1	2	6	13	7
Female	224	1	-	-	-	-	1	2	2	2	2	9
White, both sexes	341	-	-	-	-	-	-	1	1	3	6	8
Male	189	-	-	-	-	-	-	-	1	2	5	4
Female	152	-	-	-	-	-	-	1	-	1	1	4
Black, both sexes	167	1	-	-	-	1	4	2	3	5	9	8
Male	100	1	-	-	-	1	3	1	1	4	8	3
Female	67	-	-	-	-	-	1	1	2	1	1	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	1	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Massachusetts												
All races, both sexes	418	2	-	-	1	2	3	3	5	4	15	13
Male	262	1	-	-	1	1	3	3	5	3	11	12
Female	156	1	-	-	-	1	-	-	-	1	4	1
White, both sexes	391	2	-	-	1	2	3	2	5	4	14	10
Male	243	1	-	-	1	1	3	2	5	3	10	10
Female	148	1	-	-	-	1	-	-	-	1	4	-
Black, both sexes	24	-	-	-	-	-	-	-	-	-	1	3
Male	18	-	-	-	-	-	-	-	-	-	1	2
Female	6	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1,061	7	-	-	-	6	11	20	23	33	48	80
Male	630	3	-	-	-	5	6	14	15	24	33	54
Female	431	4	-	-	-	1	5	6	8	9	15	26
White, both sexes	777	6	-	-	-	2	4	7	14	14	26	49
Male	463	2	-	-	-	1	2	6	9	13	20	37
Female	314	4	-	-	-	1	2	1	5	1	6	12
Black, both sexes	276	1	-	-	-	4	7	13	9	19	20	31
Male	161	1	-	-	-	4	4	8	6	11	11	17
Female	115	-	-	-	-	-	3	5	3	8	9	14
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Minnesota												
All races, both sexes	306	-	-	-	-	-	-	4	2	1	10	8
Male	179	-	-	-	-	-	-	3	1	1	8	4
Female	127	-	-	-	-	-	-	1	1	-	2	4
White, both sexes	283	-	-	-	-	-	-	2	2	-	8	5
Male	162	-	-	-	-	-	-	2	1	-	6	2
Female	121	-	-	-	-	-	-	-	1	-	2	3
Black, both sexes	19	-	-	-	-	-	-	2	-	1	2	2
Male	16	-	-	-	-	-	-	1	-	1	2	2
Female	3	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	282	2	-	-	1	-	3	4	1	8	5	11
Male	166	-	-	-	-	-	2	3	-	6	3	9
Female	116	2	-	-	1	-	1	1	1	2	2	2
White, both sexes	159	1	-	-	-	-	1	2	-	-	1	3
Male	99	-	-	-	-	-	1	2	-	-	-	3
Female	60	1	-	-	-	-	-	-	-	-	1	-
Black, both sexes	121	1	-	-	1	-	2	2	1	8	3	8
Male	66	-	-	-	-	-	1	1	-	6	2	6
Female	55	1	-	-	1	-	1	1	1	2	1	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Missouri												
All races, both sexes	511	2	-	1	1	2	4	4	5	14	13	32
Male	284	1	-	1	-	2	3	2	1	12	10	24
Female	227	1	-	-	1	-	1	2	4	2	3	8
White, both sexes	425	1	-	1	-	2	4	2	4	8	10	20
Male	234	-	-	1	-	2	3	1	1	6	8	17
Female	191	1	-	-	-	-	1	1	3	2	2	3
Black, both sexes	83	-	-	-	1	-	-	2	1	6	3	12
Male	48	-	-	-	-	-	-	1	-	6	2	7
Female	35	-	-	-	1	-	-	1	1	-	1	5
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	39	-	-	-	1	-	-	-	-	-	1	1
Male	22	-	-	-	1	-	-	-	-	-	1	1
Female	17	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	38	-	-	-	1	-	-	-	-	-	1	-
Male	21	-	-	-	1	-	-	-	-	-	1	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Nebraska												
All races, both sexes	103	-	-	-	-	-	-	-	-	4	4	4
Male	53	-	-	-	-	-	-	-	-	4	2	1
Female	50	-	-	-	-	-	-	-	-	-	2	3
White, both sexes	91	-	-	-	-	-	-	-	-	3	3	4
Male	52	-	-	-	-	-	-	-	-	3	2	1
Female	39	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	10	-	-	-	-	-	-	-	-	1	1	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	9	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	197	1	-	-	-	1	3	2	5	7	13	11
Male	141	1	-	-	-	1	2	1	4	7	12	6
Female	56	-	-	-	-	-	1	1	1	-	1	5
White, both sexes	159	-	-	-	-	1	1	1	4	5	10	9
Male	116	-	-	-	-	1	1	-	3	5	9	5
Female	43	-	-	-	-	-	-	1	1	-	1	4
Black, both sexes	31	1	-	-	-	-	2	1	-	2	2	1
Male	22	1	-	-	-	-	1	1	-	2	2	1
Female	9	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	1	-	1	1
Male	3	-	-	-	-	-	-	-	1	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
New Hampshire												
All races, both sexes	83	-	-	-	-	-	-	3	-	-	-	2
Male	58	-	-	-	-	-	-	2	-	-	-	1
Female	25	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	81	-	-	-	-	-	-	3	-	-	-	2
Male	57	-	-	-	-	-	-	2	-	-	-	1
Female	24	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	768	3	-	-	2	3	3	8	8	17	16	33
Male	446	2	-	-	1	3	1	7	8	9	8	25
Female	322	1	-	-	1	-	2	1	-	8	8	8
White, both sexes	618	1	-	-	1	1	2	4	5	12	11	23
Male	366	1	-	-	1	1	1	3	5	6	7	19
Female	252	-	-	-	-	-	1	1	-	6	4	4
Black, both sexes	137	2	-	-	1	2	1	3	3	5	5	10
Male	75	1	-	-	-	2	-	3	3	3	1	6
Female	62	1	-	-	1	-	1	-	-	2	4	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	1	-	-	-	-
Male	5	-	-	-	-	-	-	1	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
New Mexico												
All races, both sexes	109	1	-	-	-	1	-	-	2	5	4	8
Male	65	1	-	-	-	1	-	-	-	2	4	6
Female	44	-	-	-	-	-	-	-	2	3	-	2
White, both sexes	103	1	-	-	-	1	-	-	2	5	4	6
Male	60	1	-	-	-	1	-	-	-	2	4	4
Female	43	-	-	-	-	-	-	-	2	3	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	2
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1,191	2	4	3	2	2	12	11	14	22	38	62
Male	705	-	-	2	-	2	9	8	11	17	27	43
Female	486	2	4	1	2	-	3	3	3	5	11	19
White, both sexes	1,009	1	3	2	2	1	7	6	10	11	24	41
Male	593	-	-	1	-	1	6	4	9	8	15	30
Female	416	1	3	1	2	-	1	2	1	3	9	11
Black, both sexes	167	1	1	1	-	1	5	5	4	11	14	19
Male	103	-	-	1	-	1	3	4	2	9	12	11
Female	64	1	1	-	-	-	2	1	2	2	2	8
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	2
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
North Carolina												
All races, both sexes	886	4	3	2	3	2	3	11	11	31	34	52
Male	511	3	1	2	2	2	1	10	8	23	18	36
Female	375	1	2	-	1	-	2	1	3	8	16	16
White, both sexes	611	3	3	2	2	-	1	4	6	14	22	21
Male	349	2	1	2	1	-	-	4	5	11	11	14
Female	262	1	2	-	1	-	1	-	1	3	11	7
Black, both sexes	259	1	-	-	1	2	1	7	5	16	12	28
Male	153	1	-	-	1	2	-	6	3	11	7	20
Female	106	-	-	-	-	-	1	1	2	5	5	8
American Indian 1/, both sexes	14	-	-	-	-	-	1	-	-	1	-	3
Male	8	-	-	-	-	-	1	-	-	1	-	2
Female	6	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	34	-	-	-	-	-	-	1	1	2	-	3
Male	22	-	-	-	-	-	-	1	-	2	-	1
Female	12	-	-	-	-	-	-	-	1	-	-	2
White, both sexes	29	-	-	-	-	-	-	1	1	1	-	2
Male	20	-	-	-	-	-	-	1	-	1	-	1
Female	9	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	1	-	1
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Ohio												
All races, both sexes	1,038	4	5	2	2	1	11	12	12	23	23	40
Male	575	3	2	1	1	1	8	8	10	17	13	30
Female	463	1	3	1	1	-	3	4	2	6	10	10
White, both sexes	863	3	5	2	1	-	6	7	10	17	12	30
Male	464	2	2	1	-	-	4	4	8	11	7	23
Female	399	1	3	1	1	-	2	3	2	6	5	7
Black, both sexes	173	1	-	-	-	1	5	4	2	6	11	10
Male	109	1	-	-	-	1	4	3	2	6	6	7
Female	64	-	-	-	-	-	1	1	-	-	5	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	1	-	-	1	-	-	-	-
Male	2	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	257	3	1	1	-	1	1	4	6	11	11	20
Male	152	-	1	1	-	1	1	4	6	8	6	17
Female	105	3	-	-	-	-	-	-	-	3	5	3
White, both sexes	202	3	-	1	-	1	-	3	3	9	10	10
Male	117	-	-	1	-	1	-	3	3	7	5	8
Female	85	3	-	-	-	-	-	-	-	2	5	2
Black, both sexes	37	-	-	-	-	-	1	-	2	-	1	8
Male	24	-	-	-	-	-	1	-	2	-	1	7
Female	13	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	17	-	1	-	-	-	-	1	1	2	-	2
Male	11	-	1	-	-	-	-	1	1	1	-	2
Female	6	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Oregon												
All races, both sexes	193	1	-	-	-	1	-	4	8	4	5	7
Male	117	1	-	-	-	1	-	3	7	2	2	6
Female	76	-	-	-	-	-	-	1	1	2	3	1
White, both sexes	182	1	-	-	-	-	-	3	7	3	5	7
Male	109	1	-	-	-	-	-	2	6	1	2	6
Female	73	-	-	-	-	-	-	1	1	2	3	1
Black, both sexes	7	-	-	-	-	1	-	1	1	1	-	-
Male	5	-	-	-	-	1	-	1	1	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1,473	3	1	3	1	7	8	15	19	18	31	46
Male	838	1	-	-	1	6	5	13	13	11	24	34
Female	635	2	1	3	-	1	3	2	6	7	7	12
White, both sexes	1,283	3	-	2	1	4	5	8	10	7	23	31
Male	724	1	-	-	1	4	3	8	7	3	19	26
Female	559	2	-	2	-	-	2	-	3	4	4	5
Black, both sexes	186	-	-	1	-	3	3	7	9	11	8	15
Male	112	-	-	-	-	2	2	5	6	8	5	8
Female	74	-	-	1	-	1	1	2	3	3	3	7
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	1	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Rhode Island												
All races, both sexes	38	1	-	-	-	-	-	-	1	-	-	-
Male	29	1	-	-	-	-	-	-	1	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	36	-	-	-	-	-	-	-	1	-	-	-
Male	27	-	-	-	-	-	-	-	1	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	447	3	-	-	2	3	-	5	2	9	18	17
Male	258	2	-	-	2	2	-	4	1	6	11	7
Female	189	1	-	-	-	1	-	1	1	3	7	10
White, both sexes	265	2	-	-	2	2	-	2	-	3	9	8
Male	156	2	-	-	2	1	-	2	-	2	6	4
Female	109	-	-	-	-	1	-	-	-	1	3	4
Black, both sexes	182	1	-	-	-	1	-	3	2	6	9	9
Male	102	-	-	-	-	1	-	2	1	4	5	3
Female	80	1	-	-	-	-	-	1	1	2	4	6
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
South Dakota												
All races, both sexes	33	-	-	-	-	1	-	-	-	-	-	1
Male	14	-	-	-	-	1	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	31	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	1	-	-	-	-	-	-
Male	2	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	387	2	-	-	-	1	1	1	5	4	16	18
Male	213	2	-	-	-	1	1	1	1	3	11	15
Female	174	-	-	-	-	-	-	-	4	1	5	3
White, both sexes	271	1	-	-	-	1	1	1	3	2	8	11
Male	149	1	-	-	-	1	1	1	1	1	5	9
Female	122	-	-	-	-	-	-	-	2	1	3	2
Black, both sexes	114	1	-	-	-	-	-	-	2	2	8	7
Male	63	1	-	-	-	-	-	-	-	2	6	6
Female	51	-	-	-	-	-	-	-	2	-	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Texas												
All races, both sexes	1,397	13	2	-	-	9	14	22	27	42	64	77
Male	813	6	2	-	-	5	11	16	18	29	47	52
Female	584	7	-	-	-	4	3	6	9	13	17	25
White, both sexes	1,173	10	2	-	-	6	11	17	17	30	45	58
Male	689	6	2	-	-	2	9	14	11	22	35	41
Female	484	4	-	-	-	4	2	3	6	8	10	17
Black, both sexes	210	2	-	-	-	3	3	4	10	12	19	19
Male	119	-	-	-	-	3	2	2	7	7	12	11
Female	91	2	-	-	-	-	1	2	3	5	7	8
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	1	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	1	-	-	-	-	-	1	-	-	-	-
Utah												
All races, both sexes	99	-	1	-	-	-	1	1	3	1	5	6
Male	58	-	-	-	-	-	1	1	3	-	2	3
Female	41	-	1	-	-	-	-	-	-	1	3	3
White, both sexes	95	-	1	-	-	-	1	1	2	1	4	6
Male	54	-	-	-	-	-	1	1	2	-	1	3
Female	41	-	1	-	-	-	-	-	-	1	3	3
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	1	-	1	-
Male	2	-	-	-	-	-	-	-	1	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Vermont												
All races, both sexes	38	-	-	-	-	-	-	-	-	-	-	1
Male	24	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	38	-	-	-	-	-	-	-	-	-	-	1
Male	24	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	709	1	-	1	1	2	7	11	10	14	21	29
Male	422	1	-	1	1	2	4	8	6	10	14	20
Female	287	-	-	-	-	-	3	3	4	4	7	9
White, both sexes	478	-	-	1	1	1	4	6	4	8	11	13
Male	290	-	-	1	1	1	2	5	3	6	7	11
Female	188	-	-	-	-	-	2	1	1	2	4	2
Black, both sexes	221	1	-	-	-	1	2	5	5	5	7	16
Male	127	1	-	-	-	1	2	3	3	3	5	9
Female	94	-	-	-	-	-	-	2	2	2	2	7
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	1	-	1	1	3	-
Male	4	-	-	-	-	-	-	-	-	1	2	-
Female	4	-	-	-	-	-	1	-	1	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Washington												
All races, both sexes	423	3	-	-	1	-	1	5	6	8	13	26
Male	256	-	-	-	-	-	1	4	4	5	9	19
Female	167	3	-	-	1	-	-	1	2	3	4	7
White, both sexes	373	1	-	-	1	-	1	4	6	5	9	23
Male	224	-	-	-	-	-	1	3	4	2	6	18
Female	149	1	-	-	1	-	-	1	2	3	3	5
Black, both sexes	30	-	-	-	-	-	-	1	-	1	2	3
Male	20	-	-	-	-	-	-	1	-	1	2	1
Female	10	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	5	1	-	-	-	-	-	-	-	1	1	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	3	1	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	15	1	-	-	-	-	-	-	-	1	1	-
Male	10	-	-	-	-	-	-	-	-	1	1	-
Female	5	1	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	157	1	-	-	-	-	-	-	-	1	2	3
Male	91	-	-	-	-	-	-	-	-	1	2	2
Female	66	1	-	-	-	-	-	-	-	-	-	1
White, both sexes	150	-	-	-	-	-	-	-	-	1	2	2
Male	88	-	-	-	-	-	-	-	-	1	2	2
Female	62	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	1	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	1	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Wisconsin												
All races, both sexes	489	3	-	-	2	5	3	4	2	8	11	18
Male	277	2	-	-	1	4	3	3	1	7	9	10
Female	212	1	-	-	1	1	-	1	1	1	2	8
White, both sexes	436	1	-	-	2	5	1	2	2	7	7	13
Male	243	-	-	-	1	4	1	2	1	6	5	7
Female	193	1	-	-	1	1	-	-	1	1	2	6
Black, both sexes	48	2	-	-	-	-	2	2	-	1	4	4
Male	32	2	-	-	-	-	2	1	-	1	4	2
Female	16	-	-	-	-	-	-	1	-	-	-	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	20	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
United States												
All races, both sexes	1,367	1,484	1,602	1,704	1,984	2,695	3,461	3,386	2,152	737	139	1
Male	979	1,018	1,086	1,130	1,292	1,665	1,963	1,657	859	205	30	1
Female	388	466	516	574	692	1,030	1,498	1,729	1,293	532	109	-
White, both sexes	906	1,046	1,181	1,297	1,588	2,291	3,015	3,064	1,975	668	124	1
Male	686	738	820	859	1,024	1,445	1,722	1,525	802	189	27	1
Female	220	308	361	438	564	846	1,293	1,539	1,173	479	97	-
Black, both sexes	427	413	367	361	355	337	364	250	149	57	14	-
Male	268	260	233	245	238	179	193	87	43	12	2	-
Female	159	153	134	116	117	158	171	163	106	45	12	-
American Indian 1/, both sexes	9	6	14	12	5	18	15	4	1	2	-	-
Male	5	4	10	6	3	8	8	3	-	-	-	-
Female	4	2	4	6	2	10	7	1	1	2	-	-
Asian or Pacific Islander 2/, both sexes	25	19	40	34	36	49	67	68	27	10	1	-
Male	20	16	23	20	27	33	40	42	14	4	1	-
Female	5	3	17	14	9	16	27	26	13	6	-	-
Alabama												
All races, both sexes	17	25	28	24	31	58	67	56	25	6	3	-
Male	13	15	17	15	25	32	36	26	5	-	2	-
Female	4	10	11	9	6	26	31	30	20	6	1	-
White, both sexes	6	16	17	16	23	42	49	47	19	3	3	-
Male	6	8	8	8	21	24	27	23	3	-	2	-
Female	-	8	9	8	2	18	22	24	16	3	1	-
Black, both sexes	11	8	11	8	8	16	18	9	6	3	-	-
Male	7	7	9	7	4	8	9	3	2	-	-	-
Female	4	1	2	1	4	8	9	6	4	3	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Alaska												
All races, both sexes	-	-	-	2	5	2	2	1	-	-	-	-
Male	-	-	-	1	2	2	1	-	-	-	-	-
Female	-	-	-	1	3	-	1	1	-	-	-	-
White, both sexes	-	-	-	1	4	-	1	1	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	3	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	1	2	1	-	-	-	-	-
Male	-	-	-	-	1	2	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	24	26	23	19	25	31	34	38	25	9	-	-
Male	19	20	13	13	17	22	22	20	12	2	-	-
Female	5	6	10	6	8	9	12	18	13	7	-	-
White, both sexes	20	22	22	19	24	27	33	38	24	9	-	-
Male	16	18	13	13	16	20	22	20	11	2	-	-
Female	4	4	9	6	8	7	11	18	13	7	-	-
Black, both sexes	4	3	-	-	1	-	-	-	1	-	-	-
Male	3	1	-	-	1	-	-	-	1	-	-	-
Female	1	2	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	1	-	-	3	1	-	-	-	-	-
Male	-	1	-	-	-	2	-	-	-	-	-	-
Female	-	-	1	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Arkansas												
All races, both sexes	4	7	19	12	22	25	42	35	20	11	2	-
Male	3	2	10	5	13	18	21	15	8	1	-	-
Female	1	5	9	7	9	7	21	20	12	10	2	-
White, both sexes	4	1	11	10	20	22	39	31	19	10	2	-
Male	3	-	7	3	12	16	21	14	8	1	-	-
Female	1	1	4	7	8	6	18	17	11	9	2	-
Black, both sexes	-	6	8	2	2	3	3	4	1	1	-	-
Male	-	2	3	2	1	2	-	1	-	-	-	-
Female	-	4	5	-	1	1	3	3	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	171	199	185	195	227	268	349	364	207	77	14	-
Male	123	142	126	140	147	170	181	196	88	28	1	-
Female	48	57	59	55	80	98	168	168	119	49	13	-
White, both sexes	122	154	135	150	184	222	309	309	187	72	12	-
Male	91	111	94	111	119	145	159	163	79	27	1	-
Female	31	43	41	39	65	77	150	146	108	45	11	-
Black, both sexes	35	38	29	32	25	24	18	24	11	2	2	-
Male	21	25	17	21	14	10	9	14	4	-	-	-
Female	14	13	12	11	11	14	9	10	7	2	2	-
American Indian 1/, both sexes	2	-	4	2	1	2	-	1	-	-	-	-
Male	2	-	4	1	-	1	-	1	-	-	-	-
Female	-	-	-	1	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	7	17	11	17	20	22	30	9	3	-	-
Male	9	6	11	7	14	14	13	18	5	1	-	-
Female	3	1	6	4	3	6	9	12	4	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Colorado												
All races, both sexes	21	21	21	8	11	20	35	23	18	5	1	-
Male	15	15	16	7	8	13	18	10	11	-	-	-
Female	6	6	5	1	3	7	17	13	7	5	1	-
White, both sexes	19	20	19	8	11	19	34	23	17	5	1	-
Male	14	14	14	7	8	12	18	10	10	-	-	-
Female	5	6	5	1	3	7	16	13	7	5	1	-
Black, both sexes	2	1	1	-	-	1	1	-	1	-	-	-
Male	1	1	1	-	-	1	-	-	1	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	22	13	16	17	26	34	53	48	34	11	5	-
Male	17	10	14	13	21	23	36	28	12	2	1	-
Female	5	3	2	4	5	11	17	20	22	9	4	-
White, both sexes	17	12	15	13	20	32	51	45	34	11	5	-
Male	15	10	13	11	16	21	35	26	12	2	1	-
Female	2	2	2	2	4	11	16	19	22	9	4	-
Black, both sexes	4	1	1	4	6	2	2	2	-	-	-	-
Male	2	-	1	2	5	2	1	1	-	-	-	-
Female	2	1	-	2	1	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Delaware												
All races, both sexes	9	7	5	10	11	25	16	19	13	6	-	-
Male	6	4	5	8	6	10	7	8	5	1	-	-
Female	3	3	-	2	5	15	9	11	8	5	-	-
White, both sexes	7	5	3	7	10	23	14	17	13	6	-	-
Male	4	4	3	6	5	10	5	8	5	1	-	-
Female	3	1	-	1	5	13	9	9	8	5	-	-
Black, both sexes	2	2	2	3	1	2	2	2	-	-	-	-
Male	2	-	2	2	1	-	2	-	-	-	-	-
Female	-	2	-	1	-	2	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	3	3	8	4	5	1	5	9	4	-	-	-
Male	2	1	5	1	3	-	3	3	-	-	-	-
Female	1	2	3	3	2	1	2	6	4	-	-	-
White, both sexes	-	-	2	-	1	-	2	3	2	-	-	-
Male	-	-	1	-	1	-	1	1	-	-	-	-
Female	-	-	1	-	-	-	1	2	2	-	-	-
Black, both sexes	3	3	6	4	4	1	3	6	2	-	-	-
Male	2	1	4	1	2	-	2	2	-	-	-	-
Female	1	2	2	3	2	1	1	4	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Florida												
All races, both sexes	101	128	130	156	140	258	308	340	234	76	9	-
Male	75	87	90	112	93	172	191	171	109	20	4	-
Female	26	41	40	44	47	86	117	169	125	56	5	-
White, both sexes	60	94	99	127	118	227	274	320	226	73	8	-
Male	52	70	72	90	80	154	173	165	105	18	4	-
Female	8	24	27	37	38	73	101	155	121	55	4	-
Black, both sexes	41	32	30	28	21	27	32	18	7	3	1	-
Male	23	15	17	21	12	15	18	5	4	2	-	-
Female	18	17	13	7	9	12	14	13	3	1	1	-
American Indian 1/, both sexes	-	1	1	-	-	-	1	-	-	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	1	4	1	2	1	-	-	-
Male	-	1	-	1	1	3	-	1	-	-	-	-
Female	-	-	-	-	-	1	1	1	1	-	-	-
Georgia												
All races, both sexes	36	51	68	66	70	76	105	87	62	21	5	-
Male	27	37	47	40	46	53	62	36	23	6	2	-
Female	9	14	21	26	24	23	43	51	39	15	3	-
White, both sexes	19	25	39	45	43	60	75	70	53	15	4	-
Male	15	17	27	25	27	39	49	32	21	5	1	-
Female	4	8	12	20	16	21	26	38	32	10	3	-
Black, both sexes	17	26	29	21	26	15	29	17	9	6	1	-
Male	12	20	20	15	18	13	13	4	2	1	1	-
Female	5	6	9	6	8	2	16	13	7	5	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	1	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Hawaii												
All races, both sexes	9	7	15	18	15	18	35	31	16	7	1	-
Male	9	5	8	13	10	13	27	25	11	1	1	-
Female	-	2	7	5	5	5	8	6	5	6	-	-
White, both sexes	1	5	3	4	4	6	9	9	4	3	-	-
Male	1	4	2	4	1	3	6	8	4	-	-	-
Female	-	1	1	-	3	3	3	1	-	3	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	1	12	14	11	12	26	22	12	4	1	-
Male	8	1	6	9	9	10	21	17	7	1	1	-
Female	-	-	6	5	2	2	5	5	5	3	-	-
Idaho												
All races, both sexes	3	6	4	6	6	11	13	17	5	-	1	-
Male	3	4	3	4	5	8	5	6	4	-	-	-
Female	-	2	1	2	1	3	8	11	1	-	1	-
White, both sexes	3	5	4	5	6	9	12	17	5	-	1	-
Male	3	3	3	4	5	7	4	6	4	-	-	-
Female	-	2	1	1	1	2	8	11	1	-	1	-
Black, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Illinois												
All races, both sexes	64	68	58	72	84	119	140	126	105	35	11	-
Male	42	47	41	53	58	71	72	68	41	8	3	-
Female	22	21	17	19	26	48	68	58	64	27	8	-
White, both sexes	40	38	40	49	68	92	124	116	93	35	10	-
Male	29	27	27	36	46	54	64	65	39	8	3	-
Female	11	11	13	13	22	38	60	51	54	27	7	-
Black, both sexes	24	28	16	22	14	26	14	10	12	-	1	-
Male	13	19	12	17	10	16	8	3	2	-	-	-
Female	11	9	4	5	4	10	6	7	10	-	1	-
American Indian 1/, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	2	-	2	1	2	-	-	-	-	-
Male	-	1	2	-	2	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Indiana												
All races, both sexes	35	39	40	36	49	72	78	81	46	14	4	-
Male	28	24	25	22	34	41	50	42	23	3	1	-
Female	7	15	15	14	15	31	28	39	23	11	3	-
White, both sexes	27	34	35	30	43	68	73	79	44	14	4	-
Male	22	22	23	17	29	39	47	41	23	3	1	-
Female	5	12	12	13	14	29	26	38	21	11	3	-
Black, both sexes	8	5	5	6	6	3	5	2	2	-	-	-
Male	6	2	2	5	5	2	3	1	-	-	-	-
Female	2	3	3	1	1	1	2	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Iowa												
All races, both sexes	4	9	13	9	12	30	37	24	27	12	2	-
Male	4	7	6	8	7	18	23	12	12	4	-	-
Female	-	2	7	1	5	12	14	12	15	8	2	-
White, both sexes	4	9	12	8	11	29	36	24	27	12	2	-
Male	4	7	6	8	6	17	22	12	12	4	-	-
Female	-	2	6	-	5	12	14	12	15	8	2	-
Black, both sexes	-	-	1	1	1	1	1	-	-	-	-	-
Male	-	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	13	7	12	16	17	20	38	32	17	6	3	-
Male	11	6	7	10	12	10	27	19	7	1	-	-
Female	2	1	5	6	5	10	11	13	10	5	3	-
White, both sexes	11	4	10	11	15	18	35	31	17	6	3	-
Male	9	3	5	6	11	10	25	18	7	1	-	-
Female	2	1	5	5	4	8	10	13	10	5	3	-
Black, both sexes	2	3	2	5	1	1	3	1	-	-	-	-
Male	2	3	2	4	1	-	2	1	-	-	-	-
Female	-	-	-	1	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Kentucky												
All races, both sexes	14	16	20	29	30	52	46	35	25	7	3	-
Male	9	13	13	21	18	29	31	16	9	-	1	-
Female	5	3	7	8	12	23	15	19	16	7	2	-
White, both sexes	11	11	20	26	27	49	44	34	23	7	3	-
Male	8	9	13	19	15	27	29	16	7	-	1	-
Female	3	2	7	7	12	22	15	18	16	7	2	-
Black, both sexes	3	4	-	3	3	3	2	1	2	-	-	-
Male	1	3	-	2	3	2	2	-	2	-	-	-
Female	2	1	-	1	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	22	31	32	25	41	63	68	70	48	11	4	-
Male	17	19	23	22	19	36	35	31	13	3	2	-
Female	5	12	9	3	22	27	33	39	35	8	2	-
White, both sexes	10	16	22	14	26	52	50	58	39	8	4	-
Male	7	11	15	14	13	31	22	27	12	2	2	-
Female	3	5	7	-	13	21	28	31	27	6	2	-
Black, both sexes	12	15	10	11	14	11	17	12	9	3	-	-
Male	10	8	8	8	6	5	12	4	1	1	-	-
Female	2	7	2	3	8	6	5	8	8	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Maine												
All races, both sexes	3	6	5	10	9	13	20	21	10	5	1	-
Male	2	5	4	5	7	7	14	8	5	2	-	-
Female	1	1	1	5	2	6	6	13	5	3	1	-
White, both sexes	3	6	5	10	9	13	20	21	10	4	1	-
Male	2	5	4	5	7	7	14	8	5	1	-	-
Female	1	1	1	5	2	6	6	13	5	3	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	27	25	33	39	56	55	75	90	36	22	5	-
Male	19	17	24	22	32	34	41	44	15	8	2	-
Female	8	8	9	17	24	21	34	46	21	14	3	-
White, both sexes	8	12	14	21	43	41	57	74	33	15	4	-
Male	7	9	9	13	24	27	31	37	14	4	2	-
Female	1	3	5	8	19	14	26	37	19	11	2	-
Black, both sexes	19	13	18	17	13	12	17	14	3	7	1	-
Male	12	8	14	9	8	5	10	7	1	4	-	-
Female	7	5	4	8	5	7	7	7	2	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	2	1	2	-	-	-	-
Male	-	-	1	-	-	2	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Massachusetts												
All races, both sexes	16	26	25	34	28	46	71	74	30	16	4	-
Male	13	19	19	21	18	32	44	36	16	4	-	-
Female	3	7	6	13	10	14	27	38	14	12	4	-
White, both sexes	14	24	18	31	26	45	69	71	30	16	4	-
Male	12	17	14	19	16	31	42	36	16	4	-	-
Female	2	7	4	12	10	14	27	35	14	12	4	-
Black, both sexes	2	2	6	3	2	1	2	2	-	-	-	-
Male	1	2	5	2	2	1	2	-	-	-	-	-
Female	1	-	1	1	-	-	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-
Michigan												
All races, both sexes	84	70	68	70	75	94	112	131	92	31	6	-
Male	62	45	42	49	44	65	66	56	37	10	-	-
Female	22	25	26	21	31	29	46	75	55	21	6	-
White, both sexes	52	36	53	54	56	74	90	119	84	31	6	-
Male	40	26	34	35	34	53	53	53	35	10	-	-
Female	12	10	19	19	22	21	37	66	49	21	6	-
Black, both sexes	31	33	15	16	19	20	21	10	7	-	-	-
Male	21	18	8	14	10	12	13	2	1	-	-	-
Female	10	15	7	2	9	8	8	8	6	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	1	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	1	1	-	-	-
Male	-	1	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Minnesota												
All races, both sexes	18	15	19	24	22	45	43	53	33	8	1	-
Male	15	12	12	19	17	29	18	22	15	3	-	-
Female	3	3	7	5	5	16	25	31	18	5	1	-
White, both sexes	15	12	17	21	21	44	42	52	33	8	1	-
Male	12	10	10	17	16	28	18	22	15	3	-	-
Female	3	2	7	4	5	16	24	30	18	5	1	-
Black, both sexes	3	3	1	3	1	1	-	-	-	-	-	-
Male	3	2	1	2	1	1	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Mississippi												
All races, both sexes	18	15	18	15	36	33	48	29	23	10	2	-
Male	14	12	15	12	25	16	28	10	8	3	-	-
Female	4	3	3	3	11	17	20	19	15	7	2	-
White, both sexes	7	7	10	9	19	21	35	24	14	5	-	-
Male	7	5	8	8	15	11	23	8	6	2	-	-
Female	-	2	2	1	4	10	12	16	8	3	-	-
Black, both sexes	11	8	8	6	17	12	13	5	9	4	2	-
Male	7	7	7	4	10	5	5	2	2	1	-	-
Female	4	1	1	2	7	7	8	3	7	3	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Missouri												
All races, both sexes	27	28	29	44	38	63	73	59	46	23	3	-
Male	15	21	17	28	23	42	36	26	14	5	1	-
Female	12	7	12	16	15	21	37	33	32	18	2	-
White, both sexes	16	19	22	41	29	54	69	54	45	21	3	-
Male	8	14	14	26	16	38	36	23	14	5	1	-
Female	8	5	8	15	13	16	33	31	31	16	2	-
Black, both sexes	11	9	7	3	8	8	4	5	1	2	-	-
Male	7	7	3	2	6	4	-	3	-	-	-	-
Female	4	2	4	1	2	4	4	2	1	2	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Montana												
All races, both sexes	4	-	2	4	1	4	6	9	5	1	-	-
Male	4	-	-	1	-	2	4	6	2	-	-	-
Female	-	-	2	3	1	2	2	3	3	1	-	-
White, both sexes	4	-	2	4	1	4	6	9	5	1	-	-
Male	4	-	-	1	-	2	4	6	2	-	-	-
Female	-	-	2	3	1	2	2	3	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Nebraska												
All races, both sexes	2	4	3	8	8	12	15	16	15	8	-	-
Male	-	2	1	5	4	7	11	10	5	1	-	-
Female	2	2	2	3	4	5	4	6	10	7	-	-
White, both sexes	2	3	2	6	8	11	14	15	14	6	-	-
Male	-	2	1	5	4	7	11	10	5	1	-	-
Female	2	1	1	1	4	4	3	5	9	5	-	-
Black, both sexes	-	1	-	2	-	1	1	-	1	2	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	2	-	1	1	-	1	2	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Nevada												
All races, both sexes	19	15	9	14	20	20	33	17	4	3	-	-
Male	15	7	7	14	15	11	25	9	3	1	-	-
Female	4	8	2	-	5	9	8	8	1	2	-	-
White, both sexes	16	12	7	12	14	18	28	15	4	2	-	-
Male	13	7	5	12	10	11	21	9	3	1	-	-
Female	3	5	2	-	4	7	7	6	1	1	-	-
Black, both sexes	3	3	1	2	6	-	4	2	-	1	-	-
Male	2	-	1	2	5	-	4	-	-	-	-	-
Female	1	3	-	-	1	-	-	2	-	1	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
New Hampshire												
All races, both sexes	5	5	13	4	14	13	11	10	3	-	-	-
Male	5	5	11	3	9	9	8	4	1	-	-	-
Female	-	-	2	1	5	4	3	6	2	-	-	-
White, both sexes	5	5	13	3	14	12	11	10	3	-	-	-
Male	5	5	11	2	9	9	8	4	1	-	-	-
Female	-	-	2	1	5	3	3	6	2	-	-	-
Black, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	29	41	50	37	65	78	134	133	81	25	2	-
Male	20	25	31	26	46	50	80	67	29	7	1	-
Female	9	16	19	11	19	28	54	66	52	18	1	-
White, both sexes	20	33	40	18	49	59	116	127	71	23	2	-
Male	16	22	26	13	36	41	69	64	27	7	1	-
Female	4	11	14	5	13	18	47	63	44	16	1	-
Black, both sexes	8	7	9	17	16	18	15	6	8	1	-	-
Male	3	2	4	12	10	9	11	3	2	-	-	-
Female	5	5	5	5	6	9	4	3	6	1	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	1	-	1	3	-	2	1	-	-
Male	1	1	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	3	-	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
New Mexico												
All races, both sexes	4	10	11	4	7	14	11	14	11	2	-	-
Male	3	6	6	1	3	11	7	7	5	2	-	-
Female	1	4	5	3	4	3	4	7	6	-	-	-
White, both sexes	4	10	10	4	7	13	9	14	11	2	-	-
Male	3	6	5	1	3	10	6	7	5	2	-	-
Female	1	4	5	3	4	3	3	7	6	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	1	2	-	-	-	-	-
Male	-	-	1	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	70	58	75	73	100	120	167	189	122	39	6	-
Male	47	39	53	48	69	71	105	93	51	10	-	-
Female	23	19	22	25	31	49	62	96	71	29	6	-
White, both sexes	48	48	59	56	90	106	159	176	116	39	4	-
Male	34	33	40	38	62	63	101	88	50	10	-	-
Female	14	15	19	18	28	43	58	88	66	29	4	-
Black, both sexes	20	8	15	15	9	13	7	10	6	-	2	-
Male	12	5	12	9	7	8	3	3	1	-	-	-
Female	8	3	3	6	2	5	4	7	5	-	2	-
American Indian 1/, both sexes	1	-	-	1	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	1	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	1	1	1	-	1	2	-	-	-	-
Male	1	1	1	1	-	-	1	1	-	-	-	-
Female	-	1	-	-	1	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
North Carolina												
All races, both sexes	52	52	51	71	65	97	125	111	74	27	5	-
Male	37	39	34	43	40	54	63	63	26	6	-	-
Female	15	13	17	28	25	43	62	48	48	21	5	-
White, both sexes	29	30	38	48	46	69	96	96	60	16	5	-
Male	20	24	26	28	27	41	51	56	20	5	-	-
Female	9	6	12	20	19	28	45	40	40	11	5	-
Black, both sexes	23	22	12	22	18	24	27	14	13	11	-	-
Male	17	15	8	14	13	11	11	6	6	1	-	-
Female	6	7	4	8	5	13	16	8	7	10	-	-
American Indian 1/, both sexes	-	-	-	1	1	3	2	1	1	-	-	-
Male	-	-	-	1	-	1	1	1	-	-	-	-
Female	-	-	-	-	1	2	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	2	3	3	-	1	5	6	5	1	1	-	-
Male	1	2	1	-	-	5	4	4	1	-	-	-
Female	1	1	2	-	1	-	2	1	-	1	-	-
White, both sexes	2	2	1	-	1	5	6	5	1	1	-	-
Male	1	1	1	-	-	5	4	4	1	-	-	-
Female	1	1	-	-	1	-	2	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	2	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	2	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Ohio												
All races, both sexes	71	85	75	75	77	112	133	143	102	27	3	-
Male	46	65	53	38	51	62	63	62	33	8	-	-
Female	25	20	22	37	26	50	70	81	69	19	3	-
White, both sexes	48	68	56	60	60	103	116	135	95	26	3	-
Male	34	54	42	27	37	57	55	58	30	8	-	-
Female	14	14	14	33	23	46	61	77	65	18	3	-
Black, both sexes	23	17	19	15	17	9	17	8	7	1	-	-
Male	12	11	11	11	14	5	8	4	3	-	-	-
Female	11	6	8	4	3	4	9	4	4	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	21	14	17	20	14	21	31	31	20	7	2	-
Male	11	11	13	13	9	10	14	18	8	-	-	-
Female	10	3	4	7	5	11	17	13	12	7	2	-
White, both sexes	13	11	13	14	10	16	27	29	20	7	2	-
Male	8	9	10	9	6	10	12	17	8	-	-	-
Female	5	2	3	5	4	6	15	12	12	7	2	-
Black, both sexes	5	3	3	4	4	3	1	2	-	-	-	-
Male	2	2	2	3	3	-	-	1	-	-	-	-
Female	3	1	1	1	1	3	1	1	-	-	-	-
American Indian 1/, both sexes	3	-	1	1	-	2	3	-	-	-	-	-
Male	1	-	1	1	-	-	2	-	-	-	-	-
Female	2	-	-	-	-	2	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Oregon												
All races, both sexes	12	14	11	8	13	19	30	30	18	6	2	-
Male	9	8	7	6	5	12	18	18	7	5	-	-
Female	3	6	4	2	8	7	12	12	11	1	2	-
White, both sexes	12	14	11	7	13	18	27	29	17	6	2	-
Male	9	8	7	5	5	11	16	18	7	5	-	-
Female	3	6	4	2	8	7	11	11	10	1	2	-
Black, both sexes	-	-	-	-	-	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	1	1	-	-	-	-	-
Male	-	-	-	1	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Pennsylvania												
All races, both sexes	62	73	79	90	118	187	239	262	153	50	8	-
Male	44	42	58	46	79	118	133	135	53	18	4	-
Female	18	31	21	44	39	69	106	127	100	32	4	-
White, both sexes	43	59	65	78	107	168	221	247	145	48	8	-
Male	31	31	47	42	69	110	123	129	48	18	4	-
Female	12	28	18	36	38	58	98	118	97	30	4	-
Black, both sexes	19	14	14	12	11	17	18	15	7	2	-	-
Male	13	11	11	4	10	7	10	6	4	-	-	-
Female	6	3	3	8	1	10	8	9	3	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Rhode Island												
All races, both sexes	1	-	5	4	3	4	8	7	3	1	-	-
Male	1	-	5	4	3	2	5	6	-	1	-	-
Female	-	-	-	-	-	2	3	1	3	-	-	-
White, both sexes	1	-	5	4	3	4	8	7	3	-	-	-
Male	1	-	5	4	3	2	5	6	-	-	-	-
Female	-	-	-	-	-	2	3	1	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	34	29	45	44	39	53	60	43	27	13	-	1
Male	25	20	27	27	30	36	29	18	5	5	-	1
Female	9	9	18	17	9	17	31	25	22	8	-	-
White, both sexes	15	10	18	25	22	39	43	30	22	12	-	1
Male	11	8	10	16	16	29	20	17	4	5	-	1
Female	4	2	8	9	6	10	23	13	18	7	-	-
Black, both sexes	19	19	27	19	17	14	17	13	5	1	-	-
Male	14	12	17	11	14	7	9	1	1	-	-	-
Female	5	7	10	8	3	7	8	12	4	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
South Dakota												
All races, both sexes	1	1	-	6	2	2	4	5	9	1	-	-
Male	1	1	-	4	2	-	2	1	1	1	-	-
Female	-	-	-	2	-	2	2	4	8	-	-	-
White, both sexes	1	1	-	5	2	2	4	5	9	1	-	-
Male	1	1	-	3	2	-	2	1	1	1	-	-
Female	-	-	-	2	-	2	2	4	8	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	24	24	27	33	42	43	53	45	35	9	4	-
Male	16	19	18	20	25	30	23	14	10	2	1	-
Female	8	5	9	13	17	13	30	31	25	7	3	-
White, both sexes	8	7	20	24	32	38	37	36	30	8	3	-
Male	7	5	15	14	20	27	17	12	9	2	1	-
Female	1	2	5	10	12	11	20	24	21	6	2	-
Black, both sexes	16	17	7	8	10	5	15	9	5	1	1	-
Male	9	14	3	6	5	3	5	2	1	-	-	-
Female	7	3	4	2	5	2	10	7	4	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Texas												
All races, both sexes	96	98	104	116	120	149	162	153	92	29	8	-
Male	64	64	71	72	73	82	87	61	44	7	2	-
Female	32	34	33	44	47	67	75	92	48	22	6	-
White, both sexes	71	69	89	94	104	136	147	146	88	28	5	-
Male	48	45	66	60	64	74	80	58	44	7	1	-
Female	23	24	23	34	40	62	67	88	44	21	4	-
Black, both sexes	23	28	13	20	16	13	12	6	3	1	3	-
Male	15	18	5	11	9	8	5	3	-	-	1	-
Female	8	10	8	9	7	5	7	3	3	1	2	-
American Indian 1/, both sexes	1	-	-	1	-	-	2	-	-	-	-	-
Male	1	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	2	1	-	-	1	1	1	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	1	-	2	1	-	-	-	1	1	-	-	-
Utah												
All races, both sexes	5	4	7	13	6	8	18	12	5	3	-	-
Male	4	2	6	11	3	3	11	4	2	2	-	-
Female	1	2	1	2	3	5	7	8	3	1	-	-
White, both sexes	4	4	7	12	6	8	18	12	5	3	-	-
Male	3	2	6	10	3	3	11	4	2	2	-	-
Female	1	2	1	2	3	5	7	8	3	1	-	-
Black, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Vermont												
All races, both sexes	1	2	2	4	4	4	11	4	4	-	1	-
Male	1	2	2	2	3	1	7	3	1	-	1	-
Female	-	-	-	2	1	3	4	1	3	-	-	-
White, both sexes	1	2	2	4	4	4	11	4	4	-	1	-
Male	1	2	2	2	3	1	7	3	1	-	1	-
Female	-	-	-	2	1	3	4	1	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	30	40	56	47	78	81	102	98	54	23	3	-
Male	23	25	36	37	49	52	64	38	26	5	-	-
Female	7	15	20	10	29	29	38	60	28	18	3	-
White, both sexes	18	19	23	27	50	59	83	83	46	18	3	-
Male	15	13	18	21	29	41	52	36	25	3	-	-
Female	3	6	5	6	21	18	31	47	21	15	3	-
Black, both sexes	12	21	32	20	27	21	18	15	8	5	-	-
Male	8	12	17	16	19	11	12	2	1	2	-	-
Female	4	9	15	4	8	10	6	13	7	3	-	-
American Indian 1/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Washington												
All races, both sexes	28	29	33	29	43	33	63	55	35	10	2	-
Male	20	22	20	21	31	22	35	24	14	5	-	-
Female	8	7	13	8	12	11	28	31	21	5	2	-
White, both sexes	22	23	28	26	39	30	58	50	35	10	2	-
Male	17	17	18	18	29	20	32	20	14	5	-	-
Female	5	6	10	8	10	10	26	30	21	5	2	-
Black, both sexes	4	2	4	2	4	3	2	2	-	-	-	-
Male	2	2	2	2	2	2	1	2	-	-	-	-
Female	2	-	2	-	2	1	1	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	4	1	1	-	-	2	3	-	-	-	-
Male	1	3	-	1	-	-	1	2	-	-	-	-
Female	-	1	1	-	-	-	1	1	-	-	-	-
West Virginia												
All races, both sexes	10	8	11	14	15	27	27	17	19	2	-	-
Male	10	6	9	9	8	15	14	7	8	-	-	-
Female	-	2	2	5	7	12	13	10	11	2	-	-
White, both sexes	9	7	10	14	15	26	27	16	19	2	-	-
Male	9	5	8	9	8	15	14	7	8	-	-	-
Female	-	2	2	5	7	11	13	9	11	2	-	-
Black, both sexes	1	1	1	-	-	1	-	1	-	-	-	-
Male	1	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Wisconsin												
All races, both sexes	19	26	18	25	34	56	97	78	56	21	3	-
Male	9	16	14	15	22	33	55	49	20	4	-	-
Female	10	10	4	10	12	23	42	29	36	17	3	-
White, both sexes	14	21	14	21	26	53	95	75	54	20	3	-
Male	8	13	11	13	16	31	54	47	19	4	-	-
Female	6	8	3	8	10	22	41	28	35	16	3	-
Black, both sexes	5	5	4	3	7	3	2	2	2	-	-	-
Male	1	3	3	2	6	2	1	1	1	-	-	-
Female	4	2	1	1	1	1	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	1	1	1	4	1	2	6	3	-	-	-
Male	-	1	1	-	3	1	1	2	1	-	-	-
Female	-	-	-	1	1	-	1	4	2	-	-	-
White, both sexes	-	1	1	1	4	1	2	6	3	-	-	-
Male	-	1	1	-	3	1	1	2	1	-	-	-
Female	-	-	-	1	1	-	1	4	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
United States												
All races, both sexes	41,727	53	24	22	31	76	115	129	175	258	533	870
Male	18,241	29	10	15	20	52	69	81	100	135	311	544
Female	23,486	24	14	7	11	24	46	48	75	123	222	326
White, both sexes	35,669	30	17	15	19	53	80	85	112	176	387	639
Male	15,375	17	7	11	11	36	49	56	65	100	223	414
Female	20,294	13	10	4	8	17	31	29	47	76	164	225
Black, both sexes	5,187	22	6	4	12	22	30	40	52	77	129	217
Male	2,424	12	3	3	9	16	17	21	26	32	75	118
Female	2,763	10	3	1	3	6	13	19	26	45	54	99
American Indian 1/, both sexes	169	-	1	-	-	-	2	1	3	1	6	8
Male	101	-	-	-	-	-	-	1	2	1	5	7
Female	68	-	1	-	-	-	2	-	1	-	1	1
Asian or Pacific Islander 2/, both sexes	702	1	-	3	-	1	3	3	8	4	11	6
Male	341	-	-	1	-	-	3	3	7	2	8	5
Female	361	1	-	2	-	1	-	-	1	2	3	1
Alabama												
All races, both sexes	2,553	5	1	3	2	1	8	11	13	27	76	119
Male	1,241	3	1	2	-	1	6	6	8	11	48	76
Female	1,312	2	-	1	2	-	2	5	5	16	28	43
White, both sexes	1,856	3	-	1	1	1	4	7	10	17	61	73
Male	900	2	-	1	-	1	4	4	7	9	38	46
Female	956	1	-	-	1	-	-	3	3	8	23	27
Black, both sexes	687	2	1	2	1	-	4	4	3	10	15	45
Male	336	1	1	1	-	-	2	2	1	2	10	29
Female	351	1	-	1	1	-	2	2	2	8	5	16
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Alaska												
All races, both sexes	40	-	-	-	-	-	-	-	-	-	1	2
Male	21	-	-	-	-	-	-	-	-	-	-	2
Female	19	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	30	-	-	-	-	-	-	-	-	-	1	2
Male	15	-	-	-	-	-	-	-	-	-	-	2
Female	15	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	411	-	-	-	-	3	5	2	-	-	5	4
Male	187	-	-	-	-	3	3	2	-	-	2	2
Female	224	-	-	-	-	-	2	-	-	-	3	2
White, both sexes	387	-	-	-	-	3	3	2	-	-	4	3
Male	177	-	-	-	-	3	3	2	-	-	1	2
Female	210	-	-	-	-	-	-	-	-	-	3	1
Black, both sexes	8	-	-	-	-	-	1	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	1	-	-	-	-	1
American Indian 1/, both sexes	14	-	-	-	-	-	1	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Arkansas												
All races, both sexes	492	-	1	-	2	1	1	2	1	7	6	19
Male	223	-	-	-	2	1	1	1	1	4	3	11
Female	269	-	1	-	-	-	-	1	-	3	3	8
White, both sexes	404	-	-	-	1	1	1	1	1	4	3	15
Male	187	-	-	-	1	1	1	1	1	3	1	9
Female	217	-	-	-	-	-	-	-	-	1	2	6
Black, both sexes	86	-	1	-	1	-	-	1	-	3	3	4
Male	36	-	-	-	1	-	-	-	-	1	2	2
Female	50	-	1	-	-	-	-	1	-	2	1	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1,945	4	-	1	2	4	2	4	3	6	10	17
Male	760	2	-	1	2	4	1	3	3	4	3	8
Female	1,185	2	-	-	-	-	1	1	-	2	7	9
White, both sexes	1,756	1	-	-	2	2	2	3	2	6	9	12
Male	672	1	-	-	2	2	1	2	2	4	2	5
Female	1,084	-	-	-	-	-	1	1	-	2	7	7
Black, both sexes	67	2	-	-	-	2	-	-	-	-	-	4
Male	27	1	-	-	-	2	-	-	-	-	-	2
Female	40	1	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	114	1	-	1	-	-	-	1	1	-	1	1
Male	57	-	-	1	-	-	-	1	1	-	1	1
Female	57	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Colorado												
All races, both sexes	345	3	-	2	-	2	2	1	1	3	4	5
Male	126	2	-	2	-	1	1	1	-	3	2	2
Female	219	1	-	-	-	1	1	-	1	-	2	3
White, both sexes	338	3	-	2	-	2	1	1	1	3	4	5
Male	120	2	-	2	-	1	-	1	-	3	2	2
Female	218	1	-	-	-	1	1	-	1	-	2	3
Black, both sexes	3	-	-	-	-	-	1	-	-	-	-	-
Male	2	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1,025	-	1	-	-	2	2	1	4	6	6	20
Male	437	-	1	-	-	1	-	-	1	3	4	16
Female	588	-	-	-	-	1	2	1	3	3	2	4
White, both sexes	943	-	-	-	-	1	1	1	2	2	4	19
Male	398	-	-	-	-	1	-	-	1	-	2	15
Female	545	-	-	-	-	-	1	1	1	2	2	4
Black, both sexes	78	-	1	-	-	-	1	-	2	4	2	1
Male	37	-	1	-	-	-	-	-	-	3	2	1
Female	41	-	-	-	-	-	1	-	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	1	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Delaware												
All races, both sexes	99	-	-	-	-	-	2	-	-	2	-	4
Male	34	-	-	-	-	-	1	-	-	-	-	1
Female	65	-	-	-	-	-	1	-	-	2	-	3
White, both sexes	85	-	-	-	-	-	1	-	-	1	-	3
Male	29	-	-	-	-	-	-	-	-	-	-	1
Female	56	-	-	-	-	-	1	-	-	1	-	2
Black, both sexes	12	-	-	-	-	-	1	-	-	1	-	1
Male	4	-	-	-	-	-	1	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	61	-	-	-	-	-	-	1	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	44	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	50	-	-	-	-	-	-	1	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	34	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Florida												
All races, both sexes	1,867	2	-	1	3	9	3	1	4	5	17	29
Male	883	1	-	1	3	6	1	1	2	3	8	18
Female	984	1	-	-	-	3	2	-	2	2	9	11
White, both sexes	1,707	1	-	1	1	5	3	1	1	3	15	25
Male	806	1	-	1	1	4	1	1	-	3	7	17
Female	901	-	-	-	-	1	2	-	1	-	8	8
Black, both sexes	153	1	-	-	2	4	-	-	2	2	1	4
Male	73	-	-	-	2	2	-	-	2	-	1	1
Female	80	1	-	-	-	2	-	-	-	2	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	1	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	1	-	1	-
Georgia												
All races, both sexes	2,264	8	2	1	2	2	8	9	11	18	42	41
Male	994	3	-	1	2	2	6	4	9	12	23	21
Female	1,270	5	2	-	-	-	2	5	2	6	19	20
White, both sexes	1,670	3	1	1	-	1	4	6	5	12	26	26
Male	737	1	-	1	-	1	2	3	4	8	15	14
Female	933	2	1	-	-	-	2	3	1	4	11	12
Black, both sexes	581	5	-	-	2	1	4	3	6	6	15	15
Male	250	2	-	-	2	1	4	1	5	4	7	7
Female	331	3	-	-	-	-	-	2	1	2	8	8
American Indian 1/, both sexes	5	-	1	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Hawaii												
All races, both sexes	393	-	-	-	-	-	-	-	2	1	2	4
Male	198	-	-	-	-	-	-	-	2	-	-	3
Female	195	-	-	-	-	-	-	-	-	1	2	1
White, both sexes	73	-	-	-	-	-	-	-	-	-	-	1
Male	39	-	-	-	-	-	-	-	-	-	-	-
Female	34	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	316	-	-	-	-	-	-	-	2	1	2	2
Male	156	-	-	-	-	-	-	-	2	-	-	2
Female	160	-	-	-	-	-	-	-	-	1	2	-
Idaho												
All races, both sexes	175	-	-	-	-	-	-	-	-	-	3	5
Male	91	-	-	-	-	-	-	-	-	-	1	5
Female	84	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	173	-	-	-	-	-	-	-	-	-	3	5
Male	90	-	-	-	-	-	-	-	-	-	1	5
Female	83	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Illinois												
All races, both sexes	1,736	2	3	3	3	4	4	11	15	16	27	31
Male	724	2	1	2	1	4	2	8	6	7	17	19
Female	1,012	-	2	1	2	-	2	3	9	9	10	12
White, both sexes	1,513	2	2	2	3	3	4	6	10	13	19	21
Male	620	2	1	1	1	3	2	4	5	7	12	14
Female	893	-	1	1	2	-	2	2	5	6	7	7
Black, both sexes	207	-	1	1	-	1	-	5	5	3	7	10
Male	97	-	-	1	-	1	-	4	1	-	4	5
Female	110	-	1	-	-	-	-	1	4	3	3	5
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1,125	5	1	-	-	-	2	4	7	11	20	40
Male	497	3	-	-	-	-	1	2	4	3	12	23
Female	628	2	1	-	-	-	1	2	3	8	8	17
White, both sexes	1,039	1	1	-	-	-	1	3	5	6	17	38
Male	456	-	-	-	-	-	1	2	3	3	10	21
Female	583	1	1	-	-	-	-	1	2	3	7	17
Black, both sexes	84	4	-	-	-	-	1	1	2	5	3	2
Male	40	3	-	-	-	-	-	-	1	-	2	2
Female	44	1	-	-	-	-	1	1	1	5	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Iowa												
All races, both sexes	285	1	-	-	-	1	-	1	2	2	1	5
Male	123	-	-	-	-	1	-	1	-	1	1	5
Female	162	1	-	-	-	-	-	-	2	1	-	-
White, both sexes	283	1	-	-	-	1	-	1	2	2	1	4
Male	122	-	-	-	-	1	-	1	-	1	1	4
Female	161	1	-	-	-	-	-	-	2	1	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	448	1	-	-	-	-	2	1	-	2	7	4
Male	188	1	-	-	-	-	2	1	-	1	2	4
Female	260	-	-	-	-	-	-	-	-	1	5	-
White, both sexes	416	1	-	-	-	-	2	1	-	2	7	3
Male	171	1	-	-	-	-	2	1	-	1	2	3
Female	245	-	-	-	-	-	-	-	-	1	5	-
Black, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Kentucky												
All races, both sexes	942	2	2	-	2	1	3	1	3	9	17	24
Male	392	1	2	-	1	-	1	1	2	3	11	13
Female	550	1	-	-	1	1	2	-	1	6	6	11
White, both sexes	882	2	2	-	2	1	2	1	3	9	16	21
Male	367	1	2	-	1	-	-	1	2	3	11	12
Female	515	1	-	-	1	1	2	-	1	6	5	9
Black, both sexes	59	-	-	-	-	-	1	-	-	-	1	3
Male	25	-	-	-	-	-	1	-	-	-	-	1
Female	34	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	702	1	-	-	1	2	3	1	3	12	12	19
Male	326	1	-	-	-	2	-	-	1	8	5	13
Female	376	-	-	-	1	-	3	1	2	4	7	6
White, both sexes	479	-	-	-	1	1	2	1	1	8	4	10
Male	211	-	-	-	-	1	-	-	-	5	1	6
Female	268	-	-	-	1	-	2	1	1	3	3	4
Black, both sexes	221	1	-	-	-	1	1	-	2	4	8	9
Male	115	1	-	-	-	1	-	-	1	3	4	7
Female	106	-	-	-	-	-	1	-	1	1	4	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Maine												
All races, both sexes	259	-	-	-	-	-	-	-	1	1	1	3
Male	125	-	-	-	-	-	-	-	1	1	1	3
Female	134	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	256	-	-	-	-	-	-	-	1	1	1	2
Male	122	-	-	-	-	-	-	-	1	1	1	2
Female	134	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	609	-	2	1	-	4	5	5	4	5	13	11
Male	250	-	1	1	-	3	4	4	1	2	8	4
Female	359	-	1	-	-	1	1	1	3	3	5	7
White, both sexes	442	-	1	1	-	1	5	4	1	3	7	6
Male	170	-	-	1	-	1	4	4	-	2	3	3
Female	272	-	1	-	-	-	1	-	1	1	4	3
Black, both sexes	151	-	1	-	-	3	-	1	3	2	4	5
Male	71	-	1	-	-	2	-	-	1	-	3	1
Female	80	-	-	-	-	1	-	1	2	2	1	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	-	-	2	-
Male	9	-	-	-	-	-	-	-	-	-	2	-
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Massachusetts												
All races, both sexes	979	1	1	1	-	2	3	4	4	1	9	13
Male	421	1	-	-	-	1	2	2	3	-	6	9
Female	558	-	1	1	-	1	1	2	1	1	3	4
White, both sexes	929	1	1	1	-	1	2	2	4	1	6	11
Male	391	1	-	-	-	1	1	1	3	-	3	7
Female	538	-	1	1	-	-	1	1	1	1	3	4
Black, both sexes	38	-	-	-	-	1	1	2	-	-	2	2
Male	21	-	-	-	-	-	1	1	-	-	2	2
Female	17	-	-	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	975	-	1	-	1	1	1	2	3	3	7	8
Male	387	-	1	-	1	-	1	2	-	1	4	7
Female	588	-	-	-	-	1	-	-	3	2	3	1
White, both sexes	878	-	1	-	1	1	1	2	3	1	7	6
Male	338	-	1	-	1	-	1	2	-	1	4	5
Female	540	-	-	-	-	1	-	-	3	-	3	1
Black, both sexes	94	-	-	-	-	-	-	-	-	2	-	2
Male	49	-	-	-	-	-	-	-	-	-	-	2
Female	45	-	-	-	-	-	-	-	-	2	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Minnesota												
All races, both sexes	619	2	1	1	-	-	-	1	1	2	4	5
Male	271	2	-	1	-	-	-	1	1	1	4	3
Female	348	-	1	-	-	-	-	-	-	1	-	2
White, both sexes	599	2	1	1	-	-	-	-	-	2	3	3
Male	262	2	-	1	-	-	-	-	-	1	3	2
Female	337	-	1	-	-	-	-	-	-	1	-	1
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	1	-	1	2
Male	3	-	-	-	-	-	-	-	1	-	1	1
Female	4	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	863	2	-	-	-	-	2	4	6	8	15	33
Male	404	2	-	-	-	-	2	3	4	2	10	20
Female	459	-	-	-	-	-	-	1	2	6	5	13
White, both sexes	525	-	-	-	-	-	2	-	4	4	5	16
Male	248	-	-	-	-	-	2	-	3	2	4	11
Female	277	-	-	-	-	-	-	-	1	2	1	5
Black, both sexes	333	2	-	-	-	-	-	4	2	4	10	17
Male	153	2	-	-	-	-	-	3	1	-	6	9
Female	180	-	-	-	-	-	-	1	1	4	4	8
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Missouri												
All races, both sexes	955	-	-	1	1	1	3	3	7	3	11	20
Male	432	-	-	-	1	1	1	3	5	2	7	16
Female	523	-	-	1	-	-	2	-	2	1	4	4
White, both sexes	899	-	-	1	1	1	3	3	4	3	9	18
Male	404	-	-	-	1	1	1	3	3	2	6	15
Female	495	-	-	1	-	-	2	-	1	1	3	3
Black, both sexes	51	-	-	-	-	-	-	-	3	-	2	2
Male	26	-	-	-	-	-	-	-	2	-	1	1
Female	25	-	-	-	-	-	-	-	1	-	1	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	176	-	-	-	-	1	1	2	-	-	3	1
Male	87	-	-	-	-	1	-	1	-	-	2	1
Female	89	-	-	-	-	-	1	1	-	-	1	-
White, both sexes	168	-	-	-	-	1	1	2	-	-	3	1
Male	81	-	-	-	-	1	-	1	-	-	2	1
Female	87	-	-	-	-	-	1	1	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Nebraska												
All races, both sexes	406	-	-	-	1	-	-	-	-	2	3	6
Male	182	-	-	-	-	-	-	-	-	1	1	4
Female	224	-	-	-	1	-	-	-	-	1	2	2
White, both sexes	394	-	-	-	1	-	-	-	-	2	3	6
Male	176	-	-	-	-	-	-	-	-	1	1	4
Female	218	-	-	-	1	-	-	-	-	1	2	2
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	800	-	1	2	1	1	3	3	4	8	13	26
Male	441	-	-	2	1	-	2	3	2	7	9	19
Female	359	-	1	-	-	1	1	-	2	1	4	7
White, both sexes	671	-	-	2	1	1	1	2	2	4	12	20
Male	373	-	-	2	1	-	1	2	1	4	8	15
Female	298	-	-	-	-	1	-	-	1	-	4	5
Black, both sexes	94	-	1	-	-	-	2	1	1	4	-	4
Male	54	-	-	-	-	-	1	1	-	3	-	2
Female	40	-	1	-	-	-	1	-	1	1	-	2
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	31	-	-	-	-	-	-	-	1	-	-	2
Male	12	-	-	-	-	-	-	-	1	-	-	2
Female	19	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
New Hampshire												
All races, both sexes	127	-	-	-	1	-	-	1	-	-	1	4
Male	53	-	-	-	-	-	-	-	-	-	-	2
Female	74	-	-	-	1	-	-	1	-	-	1	2
White, both sexes	126	-	-	-	1	-	-	1	-	-	1	3
Male	53	-	-	-	-	-	-	-	-	-	-	2
Female	73	-	-	-	1	-	-	1	-	-	1	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1,114	3	1	-	-	1	2	-	3	5	9	20
Male	480	2	-	-	-	1	1	-	2	4	5	13
Female	634	1	1	-	-	-	1	-	1	1	4	7
White, both sexes	985	1	1	-	-	-	2	-	2	2	5	15
Male	420	1	-	-	-	-	1	-	1	1	3	11
Female	565	-	1	-	-	-	1	-	1	1	2	4
Black, both sexes	111	2	-	-	-	1	-	-	1	2	4	4
Male	51	1	-	-	-	1	-	-	1	2	2	2
Female	60	1	-	-	-	-	-	-	-	-	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	18	-	-	-	-	-	-	-	-	1	-	1
Male	9	-	-	-	-	-	-	-	-	1	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
New Mexico												
All races, both sexes	157	-	-	-	-	1	-	-	-	-	1	1
Male	65	-	-	-	-	1	-	-	-	-	1	1
Female	92	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	146	-	-	-	-	-	-	-	-	-	1	1
Male	56	-	-	-	-	-	-	-	-	-	1	1
Female	90	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	1	-	-	-	-	-	-
Male	4	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1,999	4	-	-	3	1	9	6	7	14	16	30
Male	817	1	-	-	1	-	4	4	6	7	12	21
Female	1,182	3	-	-	2	1	5	2	1	7	4	9
White, both sexes	1,823	4	-	-	-	1	4	4	6	11	14	24
Male	738	1	-	-	-	-	1	3	5	6	11	19
Female	1,085	3	-	-	-	1	3	1	1	5	3	5
Black, both sexes	155	-	-	-	3	-	3	2	-	2	1	5
Male	67	-	-	-	1	-	1	1	-	1	-	1
Female	88	-	-	-	2	-	2	1	-	1	1	4
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	19	-	-	-	-	-	2	-	1	1	1	-
Male	10	-	-	-	-	-	2	-	1	-	1	-
Female	9	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
North Carolina												
All races, both sexes	1,331	1	-	-	-	1	2	1	5	6	11	25
Male	556	-	-	-	-	1	1	1	3	3	7	13
Female	775	1	-	-	-	-	1	-	2	3	4	12
White, both sexes	1,041	1	-	-	-	-	1	1	4	4	7	14
Male	438	-	-	-	-	-	1	1	2	2	6	8
Female	603	1	-	-	-	-	-	-	2	2	1	6
Black, both sexes	275	-	-	-	-	1	-	-	1	2	4	10
Male	109	-	-	-	-	1	-	-	1	1	1	4
Female	166	-	-	-	-	-	-	-	-	1	3	6
American Indian 1/, both sexes	13	-	-	-	-	-	1	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	59	-	1	-	-	-	-	1	-	-	1	1
Male	27	-	-	-	-	-	-	1	-	-	1	-
Female	32	-	1	-	-	-	-	-	-	-	-	1
White, both sexes	57	-	1	-	-	-	-	-	-	-	-	1
Male	25	-	-	-	-	-	-	-	-	-	-	-
Female	32	-	1	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Ohio												
All races, both sexes	1,580	1	-	1	-	2	2	2	8	11	22	40
Male	645	-	-	-	-	1	-	2	4	6	10	25
Female	935	1	-	1	-	1	2	-	4	5	12	15
White, both sexes	1,454	-	-	1	-	2	2	2	5	10	20	32
Male	592	-	-	-	-	1	-	2	3	5	10	20
Female	862	-	-	1	-	1	2	-	2	5	10	12
Black, both sexes	120	1	-	-	-	-	-	-	3	1	2	8
Male	51	-	-	-	-	-	-	-	1	1	-	5
Female	69	1	-	-	-	-	-	-	2	-	2	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	512	-	-	-	1	1	1	2	1	2	10	11
Male	213	-	-	-	1	-	1	2	-	2	5	9
Female	299	-	-	-	-	1	-	-	1	-	5	2
White, both sexes	439	-	-	-	1	1	1	1	1	1	8	7
Male	183	-	-	-	1	-	1	1	-	1	4	6
Female	256	-	-	-	-	1	-	-	1	-	4	1
Black, both sexes	37	-	-	-	-	-	-	-	-	-	1	2
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	26	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	34	-	-	-	-	-	-	1	-	1	1	2
Male	19	-	-	-	-	-	-	1	-	1	1	2
Female	15	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Oregon												
All races, both sexes	591	-	-	-	-	2	1	1	3	3	1	3
Male	222	-	-	-	-	1	1	1	2	2	1	1
Female	369	-	-	-	-	1	-	-	1	1	-	2
White, both sexes	572	-	-	-	-	2	1	-	2	1	1	3
Male	212	-	-	-	-	1	1	-	1	1	1	1
Female	360	-	-	-	-	1	-	-	1	-	-	2
Black, both sexes	3	-	-	-	-	-	-	-	-	2	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	1	1	-	-	-
Male	5	-	-	-	-	-	-	1	1	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	2,857	-	-	-	-	5	8	7	13	15	15	53
Male	1,252	-	-	-	-	2	7	4	8	10	12	30
Female	1,605	-	-	-	-	3	1	3	5	5	3	23
White, both sexes	2,596	-	-	-	-	3	6	5	9	9	8	42
Male	1,130	-	-	-	-	1	5	2	7	6	7	25
Female	1,466	-	-	-	-	2	1	3	2	3	1	17
Black, both sexes	245	-	-	-	-	2	2	2	4	5	6	11
Male	116	-	-	-	-	1	2	2	1	3	4	5
Female	129	-	-	-	-	1	-	-	3	2	2	6
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	-	-	1	1	-
Male	6	-	-	-	-	-	-	-	-	1	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Rhode Island												
All races, both sexes	112	-	-	-	-	-	-	-	-	-	2	-
Male	36	-	-	-	-	-	-	-	-	-	1	-
Female	76	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	106	-	-	-	-	-	-	-	-	-	1	-
Male	33	-	-	-	-	-	-	-	-	-	-	-
Female	73	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	946	1	-	-	3	-	5	4	6	4	19	19
Male	460	-	-	-	3	-	2	1	2	3	12	10
Female	486	1	-	-	-	-	3	3	4	1	7	9
White, both sexes	647	1	-	-	1	-	4	3	2	3	12	9
Male	299	-	-	-	1	-	1	1	1	2	8	5
Female	348	1	-	-	-	-	3	2	1	1	4	4
Black, both sexes	295	-	-	-	2	-	1	1	4	1	7	10
Male	159	-	-	-	2	-	1	-	1	1	4	5
Female	136	-	-	-	-	-	-	1	3	-	3	5
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
South Dakota												
All races, both sexes	73	-	1	-	-	1	-	1	1	-	-	-
Male	33	-	1	-	-	-	-	1	-	-	-	-
Female	40	-	-	-	-	1	-	-	1	-	-	-
White, both sexes	71	-	1	-	-	1	-	1	1	-	-	-
Male	31	-	1	-	-	-	-	1	-	-	-	-
Female	40	-	-	-	-	1	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	831	1	1	-	-	2	5	2	6	11	22	28
Male	367	-	-	-	-	2	4	-	4	7	7	20
Female	464	1	1	-	-	-	1	2	2	4	15	8
White, both sexes	700	1	1	-	-	2	4	2	4	6	17	24
Male	319	-	-	-	-	2	4	-	3	4	6	17
Female	381	1	1	-	-	-	-	2	1	2	11	7
Black, both sexes	128	-	-	-	-	-	1	-	2	5	5	4
Male	48	-	-	-	-	-	-	-	1	3	1	3
Female	80	-	-	-	-	-	1	-	1	2	4	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Texas												
All races, both sexes	2,318	2	2	3	-	11	7	12	6	13	28	48
Male	1,009	1	1	2	-	6	5	5	5	6	16	25
Female	1,309	1	1	1	-	5	2	7	1	7	12	23
White, both sexes	1,941	-	2	1	-	8	5	5	4	8	17	35
Male	823	-	1	1	-	4	4	2	3	4	8	21
Female	1,118	-	1	-	-	4	1	3	1	4	9	14
Black, both sexes	335	2	-	1	-	3	2	7	2	5	10	13
Male	164	1	-	1	-	2	1	3	2	2	7	4
Female	171	1	-	-	-	1	1	4	-	3	3	9
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	38	-	-	1	-	-	-	-	-	-	1	-
Male	18	-	-	-	-	-	-	-	-	-	1	-
Female	20	-	-	1	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	219	-	-	-	-	-	-	1	2	-	2	3
Male	97	-	-	-	-	-	-	-	-	-	2	3
Female	122	-	-	-	-	-	-	1	2	-	-	-
White, both sexes	213	-	-	-	-	-	-	1	2	-	1	2
Male	92	-	-	-	-	-	-	-	-	-	1	2
Female	121	-	-	-	-	-	-	1	2	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	1	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Vermont												
All races, both sexes	65	-	-	-	-	-	-	-	-	-	-	-
Male	27	-	-	-	-	-	-	-	-	-	-	-
Female	38	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	65	-	-	-	-	-	-	-	-	-	-	-
Male	27	-	-	-	-	-	-	-	-	-	-	-
Female	38	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1,537	1	-	1	2	2	6	7	5	10	24	43
Male	701	1	-	-	1	2	3	4	3	4	15	27
Female	836	-	-	1	1	-	3	3	2	6	9	16
White, both sexes	1,203	1	-	-	1	1	2	3	2	8	11	34
Male	536	1	-	-	-	1	1	2	-	3	6	22
Female	667	-	-	-	1	-	1	1	2	5	5	12
Black, both sexes	313	-	-	-	1	1	3	4	3	2	13	9
Male	154	-	-	-	1	1	1	2	3	1	9	5
Female	159	-	-	-	-	-	2	2	-	1	4	4
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	19	-	-	1	-	-	1	-	-	-	-	-
Male	11	-	-	-	-	-	1	-	-	-	-	-
Female	8	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Washington												
All races, both sexes	460	-	1	-	-	1	1	2	5	1	4	5
Male	171	-	1	-	-	-	1	2	3	-	2	3
Female	289	-	-	-	-	1	-	-	2	1	2	2
White, both sexes	441	-	1	-	-	1	1	2	2	1	4	4
Male	162	-	1	-	-	-	1	2	2	-	2	2
Female	279	-	-	-	-	1	-	-	-	1	2	2
Black, both sexes	6	-	-	-	-	-	-	-	1	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	2	-	-	-
Male	5	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	491	-	-	-	-	2	1	2	2	1	2	7
Male	209	-	-	-	-	2	1	1	-	1	1	6
Female	282	-	-	-	-	-	-	1	2	-	1	1
White, both sexes	475	-	-	-	-	2	1	2	2	1	2	6
Male	201	-	-	-	-	2	1	1	-	1	1	5
Female	274	-	-	-	-	-	-	1	2	-	1	1
Black, both sexes	15	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Wisconsin												
All races, both sexes	719	-	-	-	-	1	-	2	2	2	5	11
Male	307	-	-	-	-	1	-	2	2	-	5	7
Female	412	-	-	-	-	-	-	-	-	2	-	4
White, both sexes	684	-	-	-	-	1	-	2	1	2	5	8
Male	290	-	-	-	-	1	-	2	1	-	5	4
Female	394	-	-	-	-	-	-	-	-	2	-	4
Black, both sexes	27	-	-	-	-	-	-	-	-	-	-	3
Male	10	-	-	-	-	-	-	-	-	-	-	3
Female	17	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	1	-	-	-
Male	5	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	80	-	-	-	-	-	-	-	1	-	3	-
Male	32	-	-	-	-	-	-	-	-	-	2	-
Female	48	-	-	-	-	-	-	-	1	-	1	-
White, both sexes	78	-	-	-	-	-	-	-	1	-	2	-
Male	32	-	-	-	-	-	-	-	-	-	2	-
Female	46	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
United States												
All races, both sexes	1,324	1,746	2,124	2,465	3,254	4,736	6,905	7,807	5,972	2,524	583	1
Male	863	1,115	1,315	1,438	1,793	2,326	2,922	2,746	1,710	558	88	1
Female	461	631	809	1,027	1,461	2,410	3,983	5,061	4,262	1,966	495	-
White, both sexes	987	1,306	1,646	1,936	2,628	4,072	6,097	7,112	5,486	2,289	496	1
Male	650	839	1,033	1,160	1,455	2,016	2,583	2,513	1,559	503	74	1
Female	337	467	613	776	1,173	2,056	3,514	4,599	3,927	1,786	422	-
Black, both sexes	315	393	421	469	546	566	663	544	382	198	79	-
Male	199	243	242	245	289	260	271	179	115	40	9	-
Female	116	150	179	224	257	306	392	365	267	158	70	-
American Indian 1/, both sexes	6	18	16	21	18	16	18	21	9	3	1	-
Male	4	10	12	14	15	7	12	6	3	1	1	-
Female	2	8	4	7	3	9	6	15	6	2	-	-
Asian or Pacific Islander 2/, both sexes	16	29	41	39	62	82	127	130	95	34	7	-
Male	10	23	28	19	34	43	56	48	33	14	4	-
Female	6	6	13	20	28	39	71	82	62	20	3	-
Alabama												
All races, both sexes	133	168	167	176	224	286	365	373	252	106	37	-
Male	91	105	118	98	128	138	162	136	77	20	6	-
Female	42	63	49	78	96	148	203	237	175	86	31	-
White, both sexes	89	122	109	123	155	212	271	295	202	81	19	-
Male	58	74	82	72	89	104	121	113	58	15	2	-
Female	31	48	27	51	66	108	150	182	144	66	17	-
Black, both sexes	43	46	57	52	69	71	93	77	49	25	18	-
Male	33	31	35	26	39	31	41	23	19	5	4	-
Female	10	15	22	26	30	40	52	54	30	20	14	-
American Indian 1/, both sexes	-	-	1	1	-	1	-	1	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	2	1	-	1	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Alaska												
All races, both sexes	-	4	1	4	1	5	7	8	6	1	-	-
Male	-	3	1	1	1	4	4	4	1	-	-	-
Female	-	1	-	3	-	1	3	4	5	1	-	-
White, both sexes	-	1	1	2	-	4	5	8	5	1	-	-
Male	-	-	1	1	-	4	2	4	1	-	-	-
Female	-	1	-	1	-	-	3	4	4	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	2	-	2	1	1	2	-	1	-	-	-
Male	-	2	-	-	1	-	2	-	-	-	-	-
Female	-	-	-	2	-	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	15	12	22	14	39	49	81	83	46	25	6	-
Male	8	8	12	9	22	27	36	31	15	6	1	-
Female	7	4	10	5	17	22	45	52	31	19	5	-
White, both sexes	13	11	18	14	37	48	75	81	45	25	5	-
Male	6	7	10	9	22	26	33	31	15	6	1	-
Female	7	4	8	5	15	22	42	50	30	19	4	-
Black, both sexes	1	-	-	-	1	-	2	-	-	-	1	-
Male	1	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	1	3	-	1	1	3	2	1	-	-	-
Male	1	1	2	-	-	1	1	-	-	-	-	-
Female	-	-	1	-	1	-	2	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Arkansas												
All races, both sexes	19	24	29	26	43	53	74	87	63	25	9	-
Male	12	14	17	15	31	22	31	27	24	4	2	-
Female	7	10	12	11	12	31	43	60	39	21	7	-
White, both sexes	12	14	23	20	36	44	68	76	57	22	5	-
Male	8	10	14	13	27	17	28	25	21	4	2	-
Female	4	4	9	7	9	27	40	51	36	18	3	-
Black, both sexes	7	10	6	6	7	9	6	10	6	2	4	-
Male	4	4	3	2	4	5	3	2	3	-	-	-
Female	3	6	3	4	3	4	3	8	3	2	4	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
California												
All races, both sexes	39	38	41	58	102	198	345	475	406	161	29	-
Male	25	20	22	33	55	81	145	181	120	41	6	-
Female	14	18	19	25	47	117	200	294	286	120	23	-
White, both sexes	30	30	35	54	90	177	309	442	374	148	28	-
Male	20	16	17	33	48	75	124	163	111	38	6	-
Female	10	14	18	21	42	102	185	279	263	110	22	-
Black, both sexes	6	5	1	1	3	11	10	7	9	5	1	-
Male	3	1	-	-	2	3	7	4	2	-	-	-
Female	3	4	1	1	1	8	3	3	7	5	1	-
American Indian 1/, both sexes	-	-	-	-	1	1	2	2	2	-	-	-
Male	-	-	-	-	1	-	1	1	1	-	-	-
Female	-	-	-	-	-	1	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	3	3	5	3	8	9	24	24	21	8	-	-
Male	2	3	5	-	4	3	13	13	6	3	-	-
Female	1	-	-	3	4	6	11	11	15	5	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Colorado												
All races, both sexes	1	7	11	17	23	35	43	89	73	20	3	-
Male	-	4	8	7	14	13	16	28	19	3	-	-
Female	1	3	3	10	9	22	27	61	54	17	3	-
White, both sexes	1	7	11	16	21	34	42	89	72	20	3	-
Male	-	4	8	6	12	12	16	28	18	3	-	-
Female	1	3	3	10	9	22	26	61	54	17	3	-
Black, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	-	1	-	-	-
Male	-	-	-	1	-	1	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	29	36	48	61	71	110	149	236	158	73	12	-
Male	20	27	27	37	42	56	70	72	42	18	-	-
Female	9	9	21	24	29	54	79	164	116	55	12	-
White, both sexes	24	33	43	49	61	102	143	227	149	71	11	-
Male	16	24	24	31	38	51	68	69	40	18	-	-
Female	8	9	19	18	23	51	75	158	109	53	11	-
Black, both sexes	5	3	5	11	10	8	6	8	8	2	1	-
Male	4	3	3	5	4	5	2	3	1	-	-	-
Female	1	-	2	6	6	3	4	5	7	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	1	1	-	-	-
Male	-	-	-	1	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Delaware												
All races, both sexes	1	4	3	9	6	8	22	20	13	3	2	-
Male	1	2	2	6	1	2	4	9	5	-	-	-
Female	-	2	1	3	5	6	18	11	8	3	2	-
White, both sexes	1	2	3	7	4	8	19	19	12	3	2	-
Male	1	1	2	4	1	2	4	8	5	-	-	-
Female	-	1	1	3	3	6	15	11	7	3	2	-
Black, both sexes	-	2	-	2	2	-	2	-	1	-	-	-
Male	-	1	-	2	-	-	-	-	-	-	-	-
Female	-	1	-	-	2	-	2	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	1	1	6	7	4	16	11	7	4	2	-
Male	-	-	-	4	5	3	2	2	-	1	-	-
Female	1	1	1	2	2	1	14	9	7	3	2	-
White, both sexes	-	-	-	1	1	-	3	3	2	-	1	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	3	3	2	-	1	-
Black, both sexes	1	1	1	5	6	4	13	8	5	4	1	-
Male	-	-	-	3	5	3	2	2	-	1	-	-
Female	1	1	1	2	1	1	11	6	5	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Florida												
All races, both sexes	56	62	82	117	140	236	338	372	259	112	19	-
Male	30	40	52	70	77	133	164	145	92	29	7	-
Female	26	22	30	47	63	103	174	227	167	83	12	-
White, both sexes	39	51	68	98	124	220	322	354	251	107	17	-
Male	20	36	43	61	67	127	156	138	88	28	6	-
Female	19	15	25	37	57	93	166	216	163	79	11	-
Black, both sexes	16	11	14	18	15	16	14	18	8	5	2	-
Male	10	4	9	8	9	6	6	7	4	1	1	-
Female	6	7	5	10	6	10	8	11	4	4	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	1	-	2	-	-	-	-	-
Male	-	-	-	1	1	-	2	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	100	131	145	173	200	247	351	356	275	121	21	-
Male	56	84	88	95	97	118	162	117	75	15	4	-
Female	44	47	57	78	103	129	189	239	200	106	17	-
White, both sexes	62	88	95	115	131	189	289	283	223	95	15	-
Male	36	55	58	65	72	92	141	93	60	12	4	-
Female	26	33	37	50	59	97	148	190	163	83	11	-
Black, both sexes	38	43	47	58	67	55	60	72	52	26	6	-
Male	20	29	29	30	24	24	20	23	15	3	-	-
Female	18	14	18	28	43	31	40	49	37	23	6	-
American Indian 1/, both sexes	-	-	-	-	1	1	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	3	-	1	2	1	1	-	-	-	-
Male	-	-	1	-	-	2	1	1	-	-	-	-
Female	-	-	2	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Hawaii												
All races, both sexes	7	18	17	25	20	52	61	85	64	27	8	-
Male	6	13	11	15	13	28	27	36	26	15	3	-
Female	1	5	6	10	7	24	34	49	38	12	5	-
White, both sexes	1	5	3	5	2	9	8	15	13	8	3	-
Male	1	2	2	3	2	5	5	8	7	4	-	-
Female	-	3	1	2	-	4	3	7	6	4	3	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	13	14	19	17	43	52	70	51	19	5	-
Male	5	11	9	11	11	23	21	28	19	11	3	-
Female	1	2	5	8	6	20	31	42	32	8	2	-
Idaho												
All races, both sexes	5	9	7	13	8	12	30	42	24	15	2	-
Male	2	9	5	9	4	7	16	18	6	8	1	-
Female	3	-	2	4	4	5	14	24	18	7	1	-
White, both sexes	4	9	7	13	8	12	30	41	24	15	2	-
Male	2	9	5	9	4	7	16	17	6	8	1	-
Female	2	-	2	4	4	5	14	24	18	7	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Illinois												
All races, both sexes	48	69	71	95	120	179	285	339	241	132	38	-
Male	26	42	41	52	66	84	122	115	74	28	5	-
Female	22	27	30	43	54	95	163	224	167	104	33	-
White, both sexes	40	60	55	74	91	150	255	319	229	121	34	-
Male	21	38	30	41	52	67	110	108	69	28	4	-
Female	19	22	25	33	39	83	145	211	160	93	30	-
Black, both sexes	8	9	15	20	25	28	28	18	10	10	3	-
Male	5	4	10	11	13	17	10	7	4	-	-	-
Female	3	5	5	9	12	11	18	11	6	10	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	4	1	2	2	2	1	-	-
Male	-	-	1	-	1	-	2	-	1	-	-	-
Female	-	-	-	1	3	1	-	2	1	1	-	-
Indiana												
All races, both sexes	38	63	85	84	91	111	153	192	143	62	13	-
Male	27	43	54	52	52	50	61	63	32	12	3	-
Female	11	20	31	32	39	61	92	129	111	50	10	-
White, both sexes	34	55	74	78	83	100	144	188	138	60	13	-
Male	25	36	47	49	48	44	59	63	31	11	3	-
Female	9	19	27	29	35	56	85	125	107	49	10	-
Black, both sexes	4	8	11	6	7	11	9	4	5	1	-	-
Male	2	7	7	3	3	6	2	-	1	1	-	-
Female	2	1	4	3	4	5	7	4	4	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	1	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Iowa												
All races, both sexes	1	17	12	15	17	32	44	41	54	31	8	-
Male	-	11	10	11	10	17	21	15	12	7	-	-
Female	1	6	2	4	7	15	23	26	42	24	8	-
White, both sexes	-	17	12	15	17	32	44	41	54	31	8	-
Male	-	11	10	11	10	17	21	15	12	7	-	-
Female	-	6	2	4	7	15	23	26	42	24	8	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	17	16	16	18	43	42	72	77	86	41	3	-
Male	12	13	8	9	24	16	28	27	30	10	-	-
Female	5	3	8	9	19	26	44	50	56	31	3	-
White, both sexes	14	11	14	15	41	42	68	71	84	37	3	-
Male	10	9	7	7	24	16	26	25	30	7	-	-
Female	4	2	7	8	17	26	42	46	54	30	3	-
Black, both sexes	3	3	2	2	2	-	4	4	2	3	-	-
Male	2	3	1	2	-	-	2	2	-	3	-	-
Female	1	-	1	-	2	-	2	2	2	-	-	-
American Indian 1/, both sexes	-	1	-	1	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Kentucky												
All races, both sexes	41	48	46	63	78	104	152	168	111	59	8	-
Male	29	22	28	32	45	54	59	49	31	7	1	-
Female	12	26	18	31	33	50	93	119	80	52	7	-
White, both sexes	38	41	42	56	74	96	143	162	106	58	7	-
Male	29	17	26	29	41	52	56	47	29	7	1	-
Female	9	24	16	27	33	44	87	115	77	51	6	-
Black, both sexes	3	7	4	6	4	8	9	6	5	1	1	-
Male	-	5	2	3	4	2	3	2	2	-	-	-
Female	3	2	2	3	-	6	6	4	3	1	1	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	29	33	39	49	75	80	133	103	67	34	6	-
Male	20	17	19	31	46	38	55	40	19	11	-	-
Female	9	16	20	18	29	42	78	63	48	23	6	-
White, both sexes	13	19	22	30	48	55	95	85	52	27	5	-
Male	9	9	12	17	29	25	41	32	16	8	-	-
Female	4	10	10	13	19	30	54	53	36	19	5	-
Black, both sexes	16	13	17	19	26	25	38	18	15	7	1	-
Male	11	8	7	14	17	13	14	8	3	3	-	-
Female	5	5	10	5	9	12	24	10	12	4	1	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Maine												
All races, both sexes	8	9	26	13	18	37	39	49	35	15	4	-
Male	6	5	19	8	13	22	15	17	8	6	-	-
Female	2	4	7	5	5	15	24	32	27	9	4	-
White, both sexes	8	9	25	13	17	37	39	49	35	15	4	-
Male	6	5	18	8	12	22	15	17	8	6	-	-
Female	2	4	7	5	5	15	24	32	27	9	4	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	13	22	33	41	38	71	109	101	90	34	7	-
Male	10	10	22	26	23	28	41	30	21	9	2	-
Female	3	12	11	15	15	43	68	71	69	25	5	-
White, both sexes	7	15	17	24	20	44	86	82	81	30	7	-
Male	7	7	11	13	10	18	33	25	19	7	2	-
Female	-	8	6	11	10	26	53	57	62	23	5	-
Black, both sexes	6	7	14	16	16	23	21	18	7	4	-	-
Male	3	3	10	12	12	8	8	4	1	2	-	-
Female	3	4	4	4	4	15	13	14	6	2	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	2	4	2	1	2	-	-	-
Male	-	-	1	1	1	2	-	1	1	-	-	-
Female	-	-	-	-	1	2	2	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Massachusetts												
All races, both sexes	20	24	39	45	55	114	166	195	182	79	21	-
Male	11	19	28	25	35	62	69	67	57	23	1	-
Female	9	5	11	20	20	52	97	128	125	56	20	-
White, both sexes	19	19	35	39	49	107	160	193	181	78	19	-
Male	11	15	25	21	32	58	66	65	57	23	1	-
Female	8	4	10	18	17	49	94	128	124	55	18	-
Black, both sexes	1	5	3	4	4	5	4	1	-	1	2	-
Male	-	4	2	2	2	2	2	1	-	-	-	-
Female	1	1	1	2	2	3	2	-	-	1	2	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	2	1	2	1	1	-	-	-
Male	-	-	1	1	1	1	1	1	-	-	-	-
Female	-	-	-	-	1	-	1	-	1	-	-	-
Michigan												
All races, both sexes	29	22	45	35	78	99	169	214	171	67	19	-
Male	19	13	27	24	29	54	64	73	52	13	2	-
Female	10	9	18	11	49	45	105	141	119	54	17	-
White, both sexes	22	14	33	32	69	89	150	201	165	63	17	-
Male	14	8	19	22	26	48	56	68	50	10	2	-
Female	8	6	14	10	43	41	94	133	115	53	15	-
Black, both sexes	7	8	12	3	8	9	18	13	6	4	2	-
Male	5	5	8	2	3	6	8	5	2	3	-	-
Female	2	3	4	1	5	3	10	8	4	1	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Minnesota												
All races, both sexes	14	10	29	27	38	75	92	133	122	51	11	-
Male	9	7	16	16	24	35	48	55	31	15	2	-
Female	5	3	13	11	14	40	44	78	91	36	9	-
White, both sexes	13	9	27	25	36	73	89	132	121	51	11	-
Male	9	7	16	15	23	34	46	55	31	15	2	-
Female	4	2	11	10	13	39	43	77	90	36	9	-
Black, both sexes	1	-	1	1	1	2	1	-	1	-	-	-
Male	-	-	-	1	1	1	1	-	-	-	-	-
Female	1	-	1	-	-	1	-	-	1	-	-	-
American Indian 1/, both sexes	-	1	1	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	2	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	1	1	-	-	-	-
Mississippi												
All races, both sexes	43	49	70	67	67	90	132	122	94	45	14	-
Male	30	34	41	35	34	42	64	44	22	12	3	-
Female	13	15	29	32	33	48	68	78	72	33	11	-
White, both sexes	22	22	34	33	42	60	98	85	67	27	4	-
Male	12	15	21	18	25	29	50	32	16	7	1	-
Female	10	7	13	15	17	31	48	53	51	20	3	-
Black, both sexes	21	27	36	32	23	30	34	36	27	18	10	-
Male	18	19	20	16	7	13	14	12	6	5	2	-
Female	3	8	16	16	16	17	20	24	21	13	8	-
American Indian 1/, both sexes	-	-	-	2	1	-	-	1	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Missouri												
All races, both sexes	33	33	49	48	87	115	160	158	161	51	10	-
Male	28	22	27	31	48	61	73	44	51	11	-	-
Female	5	11	22	17	39	54	87	114	110	40	10	-
White, both sexes	30	30	46	44	81	108	148	153	157	50	9	-
Male	25	21	25	28	44	58	68	44	49	10	-	-
Female	5	9	21	16	37	50	80	109	108	40	9	-
Black, both sexes	2	3	3	4	6	6	11	3	4	1	1	-
Male	2	1	2	3	4	2	5	-	2	1	-	-
Female	-	2	1	1	2	4	6	3	2	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Montana												
All races, both sexes	-	7	8	14	15	16	34	39	29	4	2	-
Male	-	6	7	10	10	6	15	17	9	1	1	-
Female	-	1	1	4	5	10	19	22	20	3	1	-
White, both sexes	-	4	7	11	15	15	34	39	29	4	2	-
Male	-	4	6	7	10	6	15	17	9	1	1	-
Female	-	-	1	4	5	9	19	22	20	3	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	3	1	3	-	1	-	-	-	-	-	-
Male	-	2	1	3	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Nebraska												
All races, both sexes	15	17	23	35	31	37	57	65	68	39	7	-
Male	11	12	13	29	19	22	25	27	12	5	1	-
Female	4	5	10	6	12	15	32	38	56	34	6	-
White, both sexes	12	16	23	33	31	36	55	64	66	39	7	-
Male	8	12	13	27	19	21	25	27	12	5	1	-
Female	4	4	10	6	12	15	30	37	54	34	6	-
Black, both sexes	2	1	-	1	-	1	2	1	2	-	-	-
Male	2	-	-	1	-	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	2	1	2	-	-	-
American Indian 1/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	35	68	70	94	92	101	115	97	53	10	2	1
Male	23	45	43	55	52	60	56	45	14	2	-	1
Female	12	23	27	39	40	41	59	52	39	8	2	-
White, both sexes	21	47	61	76	78	89	103	91	49	9	1	1
Male	14	31	38	47	45	55	50	44	12	2	-	1
Female	7	16	23	29	33	34	53	47	37	7	1	-
Black, both sexes	11	15	8	13	9	11	7	2	3	1	1	-
Male	8	11	4	6	5	5	5	1	2	-	-	-
Female	3	4	4	7	4	6	2	1	1	1	1	-
American Indian 1/, both sexes	1	1	-	-	-	-	-	-	1	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	5	1	5	5	1	5	4	-	-	-	-
Male	1	2	1	2	2	-	1	-	-	-	-	-
Female	1	3	-	3	3	1	4	4	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
New Hampshire												
All races, both sexes	3	5	9	7	9	11	17	29	19	9	2	-
Male	3	5	6	4	4	6	5	12	5	1	-	-
Female	-	-	3	3	5	5	12	17	14	8	2	-
White, both sexes	3	5	9	7	9	11	17	29	19	9	2	-
Male	3	5	6	4	4	6	5	12	5	1	-	-
Female	-	-	3	3	5	5	12	17	14	8	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	22	34	45	58	86	121	208	224	181	73	18	-
Male	12	19	31	37	46	59	94	81	57	14	2	-
Female	10	15	14	21	40	62	114	143	124	59	16	-
White, both sexes	15	26	35	47	72	108	188	212	171	67	16	-
Male	10	15	25	29	37	55	83	80	52	14	2	-
Female	5	11	10	18	35	53	105	132	119	53	14	-
Black, both sexes	7	8	9	9	12	11	15	11	9	5	1	-
Male	2	4	6	6	7	4	7	1	5	-	-	-
Female	5	4	3	3	5	7	8	10	4	5	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	2	2	2	5	1	1	1	1	-
Male	-	-	-	2	2	-	4	-	-	-	-	-
Female	-	-	1	-	-	2	1	1	1	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
New Mexico												
All races, both sexes	2	4	6	8	9	23	30	34	27	9	2	-
Male	-	2	2	7	6	12	10	12	7	3	1	-
Female	2	2	4	1	3	11	20	22	20	6	1	-
White, both sexes	2	4	5	7	8	20	28	32	27	9	2	-
Male	-	2	1	6	5	9	9	11	7	3	1	-
Female	2	2	4	1	3	11	19	21	20	6	1	-
Black, both sexes	-	-	-	-	1	2	1	-	-	-	-	-
Male	-	-	-	-	1	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	1	-	1	1	1	-	-	-	-
Male	-	-	1	1	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	41	53	92	93	138	190	346	414	352	156	34	-
Male	30	33	58	50	78	100	138	145	95	29	5	-
Female	11	20	34	43	60	90	208	269	257	127	29	-
White, both sexes	36	47	78	75	115	169	325	390	340	149	31	-
Male	28	29	51	39	63	89	129	140	91	28	5	-
Female	8	18	27	36	52	80	196	250	249	121	26	-
Black, both sexes	5	5	12	18	20	19	17	23	12	5	3	-
Male	2	3	7	11	12	9	8	5	4	1	-	-
Female	3	2	5	7	8	10	9	18	8	4	3	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	2	-	2	2	4	1	-	2	-	-
Male	-	1	-	-	2	2	1	-	-	-	-	-
Female	-	-	2	-	-	-	3	1	-	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
North Carolina												
All races, both sexes	35	45	70	78	129	182	245	230	166	77	22	-
Male	25	29	45	37	73	85	97	76	44	15	1	-
Female	10	16	25	41	56	97	148	154	122	62	21	-
White, both sexes	22	29	47	42	92	153	200	195	151	59	19	-
Male	19	20	31	24	51	75	80	68	40	9	1	-
Female	3	9	16	18	41	78	120	127	111	50	18	-
Black, both sexes	13	16	19	35	35	29	44	31	14	18	3	-
Male	6	9	11	12	20	10	16	7	4	6	-	-
Female	7	7	8	23	15	19	28	24	10	12	3	-
American Indian 1/, both sexes	-	-	2	1	2	-	1	4	1	-	-	-
Male	-	-	2	1	2	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	3	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	5	-	1	3	1	3	7	13	14	4	4	-
Male	4	-	1	-	1	2	4	4	7	2	-	-
Female	1	-	-	3	-	1	3	9	7	2	4	-
White, both sexes	5	-	1	3	1	3	7	13	14	4	4	-
Male	4	-	1	-	1	2	4	4	7	2	-	-
Female	1	-	-	3	-	1	3	9	7	2	4	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Ohio												
All races, both sexes	45	56	73	81	127	187	278	295	247	86	16	-
Male	27	36	47	44	54	89	121	94	63	21	1	-
Female	18	20	26	37	73	98	157	201	184	65	15	-
White, both sexes	39	50	64	70	112	174	256	282	238	81	14	-
Male	25	33	44	41	48	80	108	90	60	21	1	-
Female	14	17	20	29	64	94	148	192	178	60	13	-
Black, both sexes	6	6	9	10	14	13	19	12	9	5	2	-
Male	2	3	3	3	6	9	11	4	3	-	-	-
Female	4	3	6	7	8	4	8	8	6	5	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	3	1	-	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	1	1	-	1	1	-	-	-	-
Oklahoma												
All races, both sexes	16	23	29	27	43	59	92	90	69	28	7	-
Male	8	12	20	11	17	26	39	32	19	6	3	-
Female	8	11	9	16	26	33	53	58	50	22	4	-
White, both sexes	14	16	22	21	35	49	85	81	63	25	7	-
Male	8	10	16	9	15	22	35	29	18	4	3	-
Female	6	6	6	12	20	27	50	52	45	21	4	-
Black, both sexes	1	1	5	3	4	5	3	6	4	2	-	-
Male	-	1	2	-	-	3	1	2	-	1	-	-
Female	1	-	3	3	4	2	2	4	4	1	-	-
American Indian 1/, both sexes	1	5	2	2	4	5	4	3	2	1	-	-
Male	-	1	2	2	2	1	3	1	1	1	-	-
Female	1	4	-	-	2	4	1	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Oregon												
All races, both sexes	17	16	15	18	33	54	107	127	127	50	13	-
Male	14	11	7	11	17	22	39	41	40	9	2	-
Female	3	5	8	7	16	32	68	86	87	41	11	-
White, both sexes	17	16	14	17	31	54	101	124	125	50	13	-
Male	14	11	6	10	16	22	36	41	39	9	2	-
Female	3	5	8	7	15	32	65	83	86	41	11	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	1	-	-	1	1	-	-	-	-
Male	-	-	1	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	5	2	2	-	-	-
Male	-	-	-	-	-	-	2	-	1	-	-	-
Female	-	-	-	-	1	-	3	2	1	-	-	-
Pennsylvania												
All races, both sexes	94	116	137	155	199	352	492	611	398	153	34	-
Male	66	79	88	103	105	161	195	225	120	33	4	-
Female	28	37	49	52	94	191	297	386	278	120	30	-
White, both sexes	78	88	120	137	177	322	457	589	367	147	32	-
Male	54	64	80	94	93	151	177	216	112	32	4	-
Female	24	24	40	43	84	171	280	373	255	115	28	-
Black, both sexes	16	27	15	17	21	29	34	20	26	6	2	-
Male	12	14	7	9	12	10	18	8	7	1	-	-
Female	4	13	8	8	9	19	16	12	19	5	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	2	1	1	1	1	2	5	-	-	-
Male	-	1	1	-	-	-	-	1	1	-	-	-
Female	-	-	1	1	1	1	1	1	4	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Rhode Island												
All races, both sexes	-	1	5	9	8	12	25	22	16	11	1	-
Male	-	1	1	7	5	2	10	3	2	4	-	-
Female	-	-	4	2	3	10	15	19	14	7	1	-
White, both sexes	-	1	5	7	7	12	25	20	16	11	1	-
Male	-	1	1	6	4	2	10	3	2	4	-	-
Female	-	-	4	1	3	10	15	17	14	7	1	-
Black, both sexes	-	-	-	2	1	-	-	1	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
South Carolina												
All races, both sexes	42	62	73	67	89	124	141	148	100	30	9	-
Male	29	43	48	44	56	58	64	51	26	6	2	-
Female	13	19	25	23	33	66	77	97	74	24	7	-
White, both sexes	23	26	40	40	59	86	109	120	78	24	7	-
Male	16	14	28	25	35	42	52	43	17	6	2	-
Female	7	12	12	15	24	44	57	77	61	18	5	-
Black, both sexes	19	35	32	27	30	38	30	28	22	6	2	-
Male	13	28	20	19	21	16	11	8	9	-	-	-
Female	6	7	12	8	9	22	19	20	13	6	2	-
American Indian 1/, both sexes	-	1	1	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
South Dakota												
All races, both sexes	-	2	1	2	5	10	8	18	14	8	1	-
Male	-	1	-	2	4	6	3	5	7	3	-	-
Female	-	1	1	-	1	4	5	13	7	5	1	-
White, both sexes	-	2	1	2	4	10	8	17	14	8	1	-
Male	-	1	-	2	3	6	3	4	7	3	-	-
Female	-	1	1	-	1	4	5	13	7	5	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	31	46	48	59	63	104	143	137	80	33	9	-
Male	19	30	23	36	40	50	49	47	20	8	1	-
Female	12	16	25	23	23	54	94	90	60	25	8	-
White, both sexes	24	36	38	48	50	94	122	121	72	27	7	-
Male	17	27	16	30	37	46	44	40	19	7	-	-
Female	7	9	22	18	13	48	78	81	53	20	7	-
Black, both sexes	7	10	10	11	13	10	18	16	8	6	2	-
Male	2	3	7	6	3	4	5	7	1	1	1	-
Female	5	7	3	5	10	6	13	9	7	5	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	3	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	3	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Texas												
All races, both sexes	74	111	107	152	194	293	398	390	302	136	29	-
Male	45	70	64	87	109	150	164	137	77	28	6	-
Female	29	41	43	65	85	143	234	253	225	108	23	-
White, both sexes	56	82	79	128	146	258	341	348	273	119	26	-
Male	33	49	49	73	82	131	136	120	70	26	6	-
Female	23	33	30	55	64	127	205	228	203	93	20	-
Black, both sexes	16	24	28	24	42	28	48	36	25	16	3	-
Male	11	16	15	14	23	15	23	17	5	2	-	-
Female	5	8	13	10	19	13	25	19	20	14	3	-
American Indian 1/, both sexes	-	2	-	-	1	1	-	-	-	-	-	-
Male	-	2	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	3	-	-	5	6	9	6	4	1	-	-
Male	1	3	-	-	3	3	5	-	2	-	-	-
Female	1	-	-	-	2	3	4	6	2	1	-	-
Utah												
All races, both sexes	7	6	5	15	16	27	41	45	31	14	4	-
Male	5	1	3	12	6	15	16	16	10	7	1	-
Female	2	5	2	3	10	12	25	29	21	7	3	-
White, both sexes	7	6	4	14	16	26	41	45	30	14	4	-
Male	5	1	2	11	6	15	16	16	9	7	1	-
Female	2	5	2	3	10	11	25	29	21	7	3	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Vermont												
All races, both sexes	1	1	1	4	4	8	7	15	16	8	-	-
Male	-	1	-	2	3	5	3	8	4	1	-	-
Female	1	-	1	2	1	3	4	7	12	7	-	-
White, both sexes	1	1	1	4	4	8	7	15	16	8	-	-
Male	-	1	-	2	3	5	3	8	4	1	-	-
Female	1	-	1	2	1	3	4	7	12	7	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	59	80	95	109	138	185	237	278	168	63	24	-
Male	36	51	58	53	77	92	98	96	59	17	4	-
Female	23	29	37	56	61	93	139	182	109	46	20	-
White, both sexes	37	53	69	79	103	141	199	241	144	57	17	-
Male	19	35	43	42	56	69	84	83	50	17	2	-
Female	18	18	26	37	47	72	115	158	94	40	15	-
Black, both sexes	21	27	22	28	33	41	38	34	22	5	6	-
Male	16	16	11	11	20	21	14	13	8	-	1	-
Female	5	11	11	17	13	20	24	21	14	5	5	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	-	4	2	2	2	-	3	2	-	1	-
Male	1	-	4	-	1	2	-	-	1	-	1	-
Female	-	-	-	2	1	-	-	3	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Washington												
All races, both sexes	11	12	14	8	33	48	65	108	99	35	7	-
Male	6	6	7	5	18	24	22	41	22	6	2	-
Female	5	6	7	3	15	24	43	67	77	29	5	-
White, both sexes	10	12	13	8	30	46	65	104	96	34	7	-
Male	5	6	6	5	16	23	22	39	22	6	2	-
Female	5	6	7	3	14	23	43	65	74	28	5	-
Black, both sexes	-	-	-	-	1	1	-	-	1	1	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	1	-	-
American Indian 1/, both sexes	1	-	1	-	1	-	-	1	-	-	-	-
Male	1	-	1	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	3	2	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	2	2	-	-	-
West Virginia												
All races, both sexes	9	23	31	30	36	69	88	83	67	31	7	-
Male	6	16	21	16	15	35	35	36	14	2	1	-
Female	3	7	10	14	21	34	53	47	53	29	6	-
White, both sexes	9	21	29	29	35	66	87	82	65	29	7	-
Male	6	15	19	16	14	33	35	35	14	2	1	-
Female	3	6	10	13	21	33	52	47	51	27	6	-
Black, both sexes	-	2	1	1	1	3	1	1	2	2	-	-
Male	-	1	1	-	1	2	-	1	-	-	-	-
Female	-	1	-	1	-	1	1	-	2	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Wisconsin												
All races, both sexes	17	26	27	35	60	67	126	149	116	57	16	-
Male	14	16	17	22	36	39	56	46	32	11	1	-
Female	3	10	10	13	24	28	70	103	84	46	15	-
White, both sexes	17	21	24	31	53	61	124	147	114	57	16	-
Male	14	14	16	21	31	36	56	46	31	11	1	-
Female	3	7	8	10	22	25	68	101	83	46	15	-
Black, both sexes	-	5	2	4	4	4	2	1	2	-	-	-
Male	-	2	-	1	2	1	-	-	1	-	-	-
Female	-	3	2	3	2	3	2	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	2	-	-	-	-	-	-	-
Male	-	-	-	-	2	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	2	-	1	-	-	-	-
Male	-	-	1	-	1	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Wyoming												
All races, both sexes	3	3	5	5	7	9	10	17	10	6	1	-
Male	2	3	2	3	3	5	5	4	3	-	-	-
Female	1	-	3	2	4	4	5	13	7	6	1	-
White, both sexes	3	3	5	5	7	9	10	17	10	5	1	-
Male	2	3	2	3	3	5	5	4	3	-	-	-
Female	1	-	3	2	4	4	5	13	7	5	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
United States												
All races, both sexes	56,565	21	11	6	6	12	31	27	60	105	212	383
Male	22,914	11	3	3	3	10	15	15	39	63	144	248
Female	33,651	10	8	3	3	2	16	12	21	42	68	135
White, both sexes	50,367	10	8	3	6	6	18	17	30	59	117	237
Male	20,262	6	1	1	3	4	7	9	20	36	77	155
Female	30,105	4	7	2	3	2	11	8	10	23	40	82
Black, both sexes	5,497	8	3	2	-	6	12	9	29	43	91	142
Male	2,341	4	2	1	-	6	8	5	18	24	63	90
Female	3,156	4	1	1	-	-	4	4	11	19	28	52
American Indian 1/, both sexes	188	1	-	-	-	-	1	-	1	-	2	3
Male	85	-	-	-	-	-	-	-	1	-	2	2
Female	103	1	-	-	-	-	1	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	513	2	-	1	-	-	-	1	-	3	2	1
Male	226	1	-	1	-	-	-	1	-	3	2	1
Female	287	1	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	1,980	-	-	-	-	-	3	-	6	8	18	29
Male	812	-	-	-	-	-	2	-	4	4	10	19
Female	1,168	-	-	-	-	-	1	-	2	4	8	10
White, both sexes	1,544	-	-	-	-	-	2	-	2	2	8	14
Male	626	-	-	-	-	-	2	-	2	2	4	7
Female	918	-	-	-	-	-	-	-	-	-	4	7
Black, both sexes	434	-	-	-	-	-	1	-	4	6	10	15
Male	185	-	-	-	-	-	-	-	2	2	6	12
Female	249	-	-	-	-	-	1	-	2	4	4	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Alaska												
All races, both sexes	50	-	-	-	-	-	-	-	-	-	1	-
Male	24	-	-	-	-	-	-	-	-	-	1	-
Female	26	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	38	-	-	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	677	-	-	1	-	1	-	-	-	-	3	6
Male	307	-	-	-	-	-	-	-	-	-	1	3
Female	370	-	-	1	-	1	-	-	-	-	2	3
White, both sexes	646	-	-	-	-	1	-	-	-	-	2	6
Male	293	-	-	-	-	-	-	-	-	-	-	3
Female	353	-	-	-	-	1	-	-	-	-	2	3
Black, both sexes	10	-	-	1	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	18	-	-	-	-	-	-	-	-	-	1	-
Male	9	-	-	-	-	-	-	-	-	-	1	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Arkansas												
All races, both sexes	711	-	1	-	-	-	-	1	-	1	3	5
Male	269	-	1	-	-	-	-	1	-	1	3	3
Female	442	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	627	-	-	-	-	-	-	-	-	1	2	3
Male	231	-	-	-	-	-	-	-	-	1	2	2
Female	396	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	77	-	1	-	-	-	-	1	-	-	1	2
Male	36	-	1	-	-	-	-	1	-	-	1	1
Female	41	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	4,198	1	1	1	1	-	1	3	4	5	14	25
Male	1,771	1	-	1	1	-	1	1	3	2	12	18
Female	2,427	-	1	-	-	-	-	2	1	3	2	7
White, both sexes	3,743	-	1	-	1	-	1	2	4	4	9	17
Male	1,563	-	-	-	1	-	1	-	3	1	7	12
Female	2,180	-	1	-	-	-	-	2	1	3	2	5
Black, both sexes	255	-	-	-	-	-	-	1	-	1	3	6
Male	119	-	-	-	-	-	-	1	-	1	3	5
Female	136	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	19	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	181	1	-	1	-	-	-	-	-	-	2	1
Male	81	1	-	1	-	-	-	-	-	-	2	1
Female	100	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Colorado												
All races, both sexes	618	-	-	-	-	-	-	1	-	2	2	-
Male	229	-	-	-	-	-	-	1	-	1	1	-
Female	389	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	591	-	-	-	-	-	-	1	-	2	-	-
Male	220	-	-	-	-	-	-	1	-	1	-	-
Female	371	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	18	-	-	-	-	-	-	-	-	-	2	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	706	-	1	-	-	1	-	-	-	-	1	2
Male	304	-	-	-	-	1	-	-	-	-	1	-
Female	402	-	1	-	-	-	-	-	-	-	-	2
White, both sexes	675	-	1	-	-	-	-	-	-	-	1	2
Male	293	-	-	-	-	-	-	-	-	-	1	-
Female	382	-	1	-	-	-	-	-	-	-	-	2
Black, both sexes	30	-	-	-	-	1	-	-	-	-	-	-
Male	10	-	-	-	-	1	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Delaware												
All races, both sexes	145	-	-	-	-	-	-	-	-	-	-	-
Male	58	-	-	-	-	-	-	-	-	-	-	-
Female	87	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	121	-	-	-	-	-	-	-	-	-	-	-
Male	49	-	-	-	-	-	-	-	-	-	-	-
Female	72	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	65	-	-	-	-	-	-	-	-	-	-	2
Male	25	-	-	-	-	-	-	-	-	-	-	1
Female	40	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	44	-	-	-	-	-	-	-	-	-	-	2
Male	17	-	-	-	-	-	-	-	-	-	-	1
Female	27	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Florida												
All races, both sexes	2,129	2	-	-	-	1	3	1	4	3	5	13
Male	948	-	-	-	-	1	-	-	2	2	4	7
Female	1,181	2	-	-	-	-	3	1	2	1	1	6
White, both sexes	1,936	-	-	-	-	-	3	1	3	-	1	4
Male	852	-	-	-	-	-	-	-	2	-	1	2
Female	1,084	-	-	-	-	-	3	1	1	-	-	2
Black, both sexes	183	2	-	-	-	1	-	-	1	3	4	9
Male	90	-	-	-	-	1	-	-	-	2	3	5
Female	93	2	-	-	-	-	-	-	1	1	1	4
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	2,045	-	-	2	-	2	1	-	4	11	18	28
Male	838	-	-	1	-	1	1	-	-	7	12	22
Female	1,207	-	-	1	-	1	-	-	4	4	6	6
White, both sexes	1,542	-	-	2	-	2	1	-	-	7	5	9
Male	625	-	-	1	-	1	1	-	-	6	2	8
Female	917	-	-	1	-	1	-	-	-	1	3	1
Black, both sexes	490	-	-	-	-	-	-	-	4	4	13	19
Male	212	-	-	-	-	-	-	-	-	1	10	14
Female	278	-	-	-	-	-	-	-	4	3	3	5
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Hawaii												
All races, both sexes	139	-	-	-	-	-	-	1	-	-	-	-
Male	67	-	-	-	-	-	-	1	-	-	-	-
Female	72	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	46	-	-	-	-	-	-	-	-	-	-	-
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	91	-	-	-	-	-	-	1	-	-	-	-
Male	45	-	-	-	-	-	-	1	-	-	-	-
Female	46	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	249	-	-	-	-	-	1	1	-	-	1	1
Male	93	-	-	-	-	-	-	1	-	-	-	1
Female	156	-	-	-	-	-	1	-	-	-	1	-
White, both sexes	245	-	-	-	-	-	1	1	-	-	1	1
Male	89	-	-	-	-	-	-	1	-	-	-	1
Female	156	-	-	-	-	-	1	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Illinois												
All races, both sexes	2,910	3	-	-	-	-	-	2	2	8	12	22
Male	1,125	3	-	-	-	-	-	2	-	4	8	14
Female	1,785	-	-	-	-	-	-	-	2	4	4	8
White, both sexes	2,562	2	-	-	-	-	-	1	1	1	4	9
Male	961	2	-	-	-	-	-	1	-	1	3	6
Female	1,601	-	-	-	-	-	-	-	1	-	1	3
Black, both sexes	335	1	-	-	-	-	-	1	1	7	8	13
Male	160	1	-	-	-	-	-	1	-	3	5	8
Female	175	-	-	-	-	-	-	-	1	4	3	5
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1,798	1	1	-	1	1	-	1	2	1	11	7
Male	683	1	-	-	-	1	-	1	2	1	5	3
Female	1,115	-	1	-	1	-	-	-	-	-	6	4
White, both sexes	1,677	1	1	-	1	1	-	1	1	-	8	7
Male	632	1	-	-	-	1	-	1	1	-	4	3
Female	1,045	-	1	-	1	-	-	-	-	-	4	4
Black, both sexes	119	-	-	-	-	-	-	-	1	1	3	-
Male	50	-	-	-	-	-	-	-	1	1	1	-
Female	69	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Iowa												
All races, both sexes	297	-	-	-	-	-	-	-	1	1	1	2
Male	109	-	-	-	-	-	-	-	-	1	1	1
Female	188	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	296	-	-	-	-	-	-	-	1	1	1	2
Male	108	-	-	-	-	-	-	-	-	1	1	1
Female	188	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1,043	-	-	-	-	-	1	-	1	-	2	3
Male	390	-	-	-	-	-	-	-	1	-	1	3
Female	653	-	-	-	-	-	1	-	-	-	1	-
White, both sexes	990	-	-	-	-	-	1	-	-	-	2	3
Male	370	-	-	-	-	-	-	-	-	-	1	3
Female	620	-	-	-	-	-	1	-	-	-	1	-
Black, both sexes	42	-	-	-	-	-	-	-	1	-	-	-
Male	17	-	-	-	-	-	-	-	1	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Kentucky												
All races, both sexes	1,272	-	-	-	1	-	1	1	-	5	4	13
Male	504	-	-	-	-	-	1	-	-	2	4	7
Female	768	-	-	-	1	-	-	1	-	3	-	6
White, both sexes	1,189	-	-	-	1	-	-	1	-	5	4	9
Male	471	-	-	-	-	-	-	-	-	2	4	6
Female	718	-	-	-	1	-	-	1	-	3	-	3
Black, both sexes	81	-	-	-	-	-	1	-	-	-	-	4
Male	31	-	-	-	-	-	1	-	-	-	-	1
Female	50	-	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1,294	1	-	-	-	-	2	-	5	4	6	11
Male	512	1	-	-	-	-	1	-	4	3	4	5
Female	782	-	-	-	-	-	1	-	1	1	2	6
White, both sexes	984	-	-	-	-	-	-	-	1	3	5	6
Male	390	-	-	-	-	-	-	-	1	2	4	2
Female	594	-	-	-	-	-	-	-	-	1	1	4
Black, both sexes	307	1	-	-	-	-	2	-	4	1	1	5
Male	122	1	-	-	-	-	1	-	3	1	-	3
Female	185	-	-	-	-	-	1	-	1	-	1	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Maine												
All races, both sexes	283	1	-	-	1	-	1	-	-	-	-	-
Male	114	1	-	-	-	-	-	-	-	-	-	-
Female	169	-	-	-	1	-	1	-	-	-	-	-
White, both sexes	283	1	-	-	1	-	1	-	-	-	-	-
Male	114	1	-	-	-	-	-	-	-	-	-	-
Female	169	-	-	-	1	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	603	-	-	-	-	1	1	-	-	1	3	3
Male	245	-	-	-	-	1	1	-	-	1	2	2
Female	358	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	476	-	-	-	-	-	-	-	-	-	2	1
Male	191	-	-	-	-	-	-	-	-	-	2	-
Female	285	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	120	-	-	-	-	1	1	-	-	1	1	2
Male	51	-	-	-	-	1	1	-	-	1	-	2
Female	69	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Massachusetts												
All races, both sexes	1,582	-	-	-	-	1	-	-	-	2	2	2
Male	580	-	-	-	-	1	-	-	-	2	1	2
Female	1,002	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	1,509	-	-	-	-	1	-	-	-	1	1	2
Male	549	-	-	-	-	1	-	-	-	1	-	2
Female	960	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	59	-	-	-	-	-	-	-	-	-	1	-
Male	24	-	-	-	-	-	-	-	-	-	1	-
Female	35	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	1	-	-
Male	7	-	-	-	-	-	-	-	-	1	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	2,422	-	-	-	-	-	-	-	3	5	7	15
Male	930	-	-	-	-	-	-	-	1	3	5	10
Female	1,492	-	-	-	-	-	-	-	2	2	2	5
White, both sexes	2,180	-	-	-	-	-	-	-	1	1	3	12
Male	830	-	-	-	-	-	-	-	-	-	2	8
Female	1,350	-	-	-	-	-	-	-	1	1	1	4
Black, both sexes	222	-	-	-	-	-	-	-	2	4	4	3
Male	93	-	-	-	-	-	-	-	1	3	3	2
Female	129	-	-	-	-	-	-	-	1	1	1	1
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Minnesota												
All races, both sexes	1,061	-	1	-	-	-	-	1	1	-	-	1
Male	423	-	-	-	-	-	-	-	1	-	-	1
Female	638	-	1	-	-	-	-	1	-	-	-	-
White, both sexes	1,041	-	1	-	-	-	-	-	-	-	-	1
Male	410	-	-	-	-	-	-	-	-	-	-	1
Female	631	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	12	-	-	-	-	-	-	1	1	-	-	-
Male	9	-	-	-	-	-	-	-	1	-	-	-
Female	3	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1,355	-	-	-	-	-	3	2	1	5	12	20
Male	518	-	-	-	-	-	2	1	1	3	6	10
Female	837	-	-	-	-	-	1	1	-	2	6	10
White, both sexes	982	-	-	-	-	-	1	-	-	1	5	9
Male	380	-	-	-	-	-	-	-	-	-	2	4
Female	602	-	-	-	-	-	1	-	-	1	3	5
Black, both sexes	371	-	-	-	-	-	2	2	1	4	7	11
Male	137	-	-	-	-	-	2	1	1	3	4	6
Female	234	-	-	-	-	-	-	1	-	1	3	5
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Missouri												
All races, both sexes	1,451	-	-	-	-	1	-	-	1	3	5	9
Male	576	-	-	-	-	1	-	-	1	3	3	7
Female	875	-	-	-	-	-	-	-	-	-	2	2
White, both sexes	1,360	-	-	-	-	-	-	-	-	3	3	7
Male	534	-	-	-	-	-	-	-	-	3	1	6
Female	826	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	90	-	-	-	-	1	-	-	1	-	2	2
Male	41	-	-	-	-	1	-	-	1	-	2	1
Female	49	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	327	-	-	-	-	-	1	-	-	-	-	1
Male	140	-	-	-	-	-	-	-	-	-	-	-
Female	187	-	-	-	-	-	1	-	-	-	-	1
White, both sexes	318	-	-	-	-	-	-	-	-	-	-	1
Male	136	-	-	-	-	-	-	-	-	-	-	-
Female	182	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	-	-	-	-	-	1	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Nebraska												
All races, both sexes	465	-	-	-	-	-	-	-	-	1	1	1
Male	181	-	-	-	-	-	-	-	-	1	1	1
Female	284	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	452	-	-	-	-	-	-	-	-	1	-	1
Male	176	-	-	-	-	-	-	-	-	1	-	1
Female	276	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	12	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	447	-	-	-	-	-	-	-	-	1	-	2
Male	236	-	-	-	-	-	-	-	-	-	-	2
Female	211	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	415	-	-	-	-	-	-	-	-	1	-	2
Male	219	-	-	-	-	-	-	-	-	-	-	2
Female	196	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
New Hampshire												
All races, both sexes	231	-	-	-	-	-	-	-	-	-	-	-
Male	82	-	-	-	-	-	-	-	-	-	-	-
Female	149	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	228	-	-	-	-	-	-	-	-	-	-	-
Male	82	-	-	-	-	-	-	-	-	-	-	-
Female	146	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1,522	-	-	1	-	-	1	-	1	-	2	6
Male	620	-	-	1	-	-	1	-	1	-	-	3
Female	902	-	-	-	-	-	-	-	-	-	2	3
White, both sexes	1,357	-	-	-	-	-	-	-	1	-	1	2
Male	551	-	-	-	-	-	-	-	1	-	-	1
Female	806	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	152	-	-	1	-	-	1	-	-	-	1	4
Male	64	-	-	1	-	-	1	-	-	-	-	2
Female	88	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
New Mexico												
All races, both sexes	375	-	-	-	-	-	-	-	-	1	-	3
Male	156	-	-	-	-	-	-	-	-	1	-	3
Female	219	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	360	-	-	-	-	-	-	-	-	1	-	3
Male	148	-	-	-	-	-	-	-	-	1	-	3
Female	212	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	2,675	1	-	-	1	1	-	2	2	-	7	12
Male	1,085	1	-	-	1	1	-	1	2	-	5	9
Female	1,590	-	-	-	-	-	-	1	-	-	2	3
White, both sexes	2,461	1	-	-	1	1	-	2	1	-	5	8
Male	999	1	-	-	1	1	-	1	1	-	4	8
Female	1,462	-	-	-	-	-	-	1	-	-	1	-
Black, both sexes	195	-	-	-	-	-	-	-	1	-	2	4
Male	75	-	-	-	-	-	-	-	1	-	1	1
Female	120	-	-	-	-	-	-	-	-	-	1	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
North Carolina												
All races, both sexes	1,540	6	1	1	-	-	-	1	1	3	5	12
Male	613	2	-	-	-	-	-	-	1	1	5	8
Female	927	4	1	1	-	-	-	1	-	2	-	4
White, both sexes	1,277	3	1	1	-	-	-	-	-	-	-	8
Male	514	-	-	-	-	-	-	-	-	-	-	6
Female	763	3	1	1	-	-	-	-	-	-	-	2
Black, both sexes	254	2	-	-	-	-	-	1	1	3	5	4
Male	97	2	-	-	-	-	-	-	1	1	5	2
Female	157	-	-	-	-	-	-	1	-	2	-	2
American Indian 1/, both sexes	4	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	81	-	-	-	-	-	-	-	1	-	-	-
Male	29	-	-	-	-	-	-	-	-	-	-	-
Female	52	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	77	-	-	-	-	-	-	-	1	-	-	-
Male	26	-	-	-	-	-	-	-	-	-	-	-
Female	51	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Ohio												
All races, both sexes	1,514	-	1	-	-	-	1	-	2	1	6	7
Male	637	-	1	-	-	-	1	-	2	-	5	5
Female	877	-	-	-	-	-	-	-	-	1	1	2
White, both sexes	1,386	-	-	-	-	-	1	-	1	-	5	5
Male	582	-	-	-	-	-	1	-	1	-	4	5
Female	804	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	127	-	1	-	-	-	-	-	1	1	1	2
Male	55	-	1	-	-	-	-	-	1	-	1	-
Female	72	-	-	-	-	-	-	-	-	1	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1,392	-	1	-	1	-	-	1	3	2	7	15
Male	585	-	-	-	1	-	-	1	3	1	5	9
Female	807	-	1	-	-	-	-	-	-	1	2	6
White, both sexes	1,272	-	1	-	1	-	-	1	2	1	5	11
Male	533	-	-	-	1	-	-	1	2	-	4	6
Female	739	-	1	-	-	-	-	-	-	1	1	5
Black, both sexes	78	-	-	-	-	-	-	-	1	1	2	3
Male	34	-	-	-	-	-	-	-	1	1	1	2
Female	44	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	38	-	-	-	-	-	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	23	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Oregon												
All races, both sexes	705	1	-	-	-	-	1	-	1	2	1	2
Male	284	-	-	-	-	-	-	-	-	-	-	2
Female	421	1	-	-	-	-	1	-	1	2	1	-
White, both sexes	693	-	-	-	-	-	1	-	1	2	1	2
Male	280	-	-	-	-	-	-	-	-	-	-	2
Female	413	-	-	-	-	-	1	-	1	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	1	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	1	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	3,287	1	-	-	-	-	3	1	3	3	8	21
Male	1,275	-	-	-	-	-	-	-	2	2	7	11
Female	2,012	1	-	-	-	-	3	1	1	1	1	10
White, both sexes	3,068	1	-	-	-	-	2	1	1	2	6	12
Male	1,185	-	-	-	-	-	-	-	1	1	5	6
Female	1,883	1	-	-	-	-	2	1	-	1	1	6
Black, both sexes	204	-	-	-	-	-	1	-	2	1	2	9
Male	82	-	-	-	-	-	-	-	1	1	2	5
Female	122	-	-	-	-	-	1	-	1	-	-	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Rhode Island												
All races, both sexes	136	-	-	-	-	-	-	-	-	-	-	-
Male	57	-	-	-	-	-	-	-	-	-	-	-
Female	79	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	134	-	-	-	-	-	-	-	-	-	-	-
Male	57	-	-	-	-	-	-	-	-	-	-	-
Female	77	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	910	2	-	-	-	1	2	-	-	1	6	10
Male	384	1	-	-	-	1	1	-	-	-	5	6
Female	526	1	-	-	-	-	1	-	-	1	1	4
White, both sexes	674	1	-	-	-	-	1	-	-	-	2	7
Male	291	1	-	-	-	-	1	-	-	-	2	4
Female	383	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes	233	1	-	-	-	1	1	-	-	1	4	3
Male	92	-	-	-	-	1	-	-	-	-	3	2
Female	141	1	-	-	-	-	1	-	-	1	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
South Dakota												
All races, both sexes	63	-	-	-	-	-	-	-	1	-	-	-
Male	27	-	-	-	-	-	-	-	1	-	-	-
Female	36	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	58	-	-	-	-	-	-	-	-	-	-	-
Male	25	-	-	-	-	-	-	-	-	-	-	-
Female	33	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	722	-	-	-	-	-	1	1	-	3	8	6
Male	260	-	-	-	-	-	1	1	-	2	5	5
Female	462	-	-	-	-	-	-	-	-	1	3	1
White, both sexes	617	-	-	-	-	-	-	1	-	2	5	3
Male	216	-	-	-	-	-	-	1	-	1	2	2
Female	401	-	-	-	-	-	-	-	-	1	3	1
Black, both sexes	102	-	-	-	-	-	1	-	-	1	3	3
Male	43	-	-	-	-	-	1	-	-	1	3	3
Female	59	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Texas												
All races, both sexes	4,275	1	-	-	-	1	2	5	6	14	18	40
Male	1,850	-	-	-	-	1	1	2	3	11	13	28
Female	2,425	1	-	-	-	-	1	3	3	3	5	12
White, both sexes	3,802	-	-	-	-	-	2	4	5	11	14	29
Male	1,630	-	-	-	-	-	1	2	2	8	10	20
Female	2,172	-	-	-	-	-	1	2	3	3	4	9
Black, both sexes	442	1	-	-	-	1	-	1	1	2	4	10
Male	204	-	-	-	-	1	-	-	1	2	3	7
Female	238	1	-	-	-	-	-	1	-	-	1	3
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	27	-	-	-	-	-	-	-	-	1	-	-
Male	13	-	-	-	-	-	-	-	-	1	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	612	-	1	-	-	-	-	-	-	2	1	1
Male	262	-	-	-	-	-	-	-	-	1	1	-
Female	350	-	1	-	-	-	-	-	-	1	-	1
White, both sexes	601	-	1	-	-	-	-	-	-	2	1	1
Male	256	-	-	-	-	-	-	-	-	1	1	-
Female	345	-	1	-	-	-	-	-	-	1	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Vermont												
All races, both sexes	46	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	46	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1,657	-	1	-	-	-	1	1	2	4	9	14
Male	683	-	1	-	-	-	1	1	2	2	6	9
Female	974	-	-	-	-	-	-	-	-	2	3	5
White, both sexes	1,339	-	1	-	-	-	-	-	1	3	3	9
Male	540	-	1	-	-	-	-	-	1	2	3	6
Female	799	-	-	-	-	-	-	-	-	1	-	3
Black, both sexes	300	-	-	-	-	-	1	1	1	1	6	5
Male	133	-	-	-	-	-	1	1	1	-	3	3
Female	167	-	-	-	-	-	-	-	-	1	3	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Washington												
All races, both sexes	413	-	-	-	-	-	-	-	-	-	-	1
Male	163	-	-	-	-	-	-	-	-	-	-	-
Female	250	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	397	-	-	-	-	-	-	-	-	-	-	1
Male	157	-	-	-	-	-	-	-	-	-	-	-
Female	240	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	552	-	-	-	-	-	-	-	1	2	1	2
Male	221	-	-	-	-	-	-	-	1	1	-	2
Female	331	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	537	-	-	-	-	-	-	-	1	1	1	2
Male	214	-	-	-	-	-	-	-	1	-	-	2
Female	323	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Wisconsin												
All races, both sexes	1,420	-	1	-	-	-	-	-	1	-	1	8
Male	573	-	-	-	-	-	-	-	1	-	1	6
Female	847	-	1	-	-	-	-	-	-	-	-	2
White, both sexes	1,381	-	-	-	-	-	-	-	1	-	1	6
Male	561	-	-	-	-	-	-	-	1	-	1	4
Female	820	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	33	-	1	-	-	-	-	-	-	-	-	2
Male	11	-	-	-	-	-	-	-	-	-	-	2
Female	22	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	115	-	-	-	-	-	-	-	-	-	-	-
Male	41	-	-	-	-	-	-	-	-	-	-	-
Female	74	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	113	-	-	-	-	-	-	-	-	-	-	-
Male	41	-	-	-	-	-	-	-	-	-	-	-
Female	72	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
United States												
All races, both sexes	690	1,139	1,619	2,291	3,458	5,985	9,950	12,855	11,224	5,230	1,249	1
Male	407	716	927	1,312	1,889	2,926	4,360	4,916	3,489	1,219	198	1
Female	283	423	692	979	1,569	3,059	5,590	7,939	7,735	4,011	1,051	-
White, both sexes	443	786	1,244	1,820	2,912	5,243	9,002	11,925	10,505	4,844	1,131	1
Male	272	498	717	1,070	1,612	2,585	3,989	4,615	3,281	1,129	174	1
Female	171	288	527	750	1,300	2,658	5,013	7,310	7,224	3,715	957	-
Black, both sexes	236	333	346	415	505	665	808	792	620	329	103	-
Male	130	206	196	210	257	299	312	249	171	69	21	-
Female	106	127	150	205	248	366	496	543	449	260	82	-
American Indian 1/, both sexes	4	13	14	22	15	24	26	28	20	9	5	-
Male	1	7	6	12	5	17	10	10	8	2	2	-
Female	3	6	8	10	10	7	16	18	12	7	3	-
Asian or Pacific Islander 2/, both sexes	7	7	15	34	26	53	114	110	79	48	10	-
Male	4	5	8	20	15	25	49	42	29	19	1	-
Female	3	2	7	14	11	28	65	68	50	29	9	-
Alabama												
All races, both sexes	45	65	71	103	162	228	336	418	326	132	30	-
Male	21	38	34	65	92	115	137	145	94	31	1	-
Female	24	27	37	38	70	113	199	273	232	101	29	-
White, both sexes	17	35	48	74	114	176	282	367	273	110	20	-
Male	9	17	25	49	70	90	115	128	81	24	1	-
Female	8	18	23	25	44	86	167	239	192	86	19	-
Black, both sexes	28	30	23	29	48	51	54	51	53	22	9	-
Male	12	21	9	16	22	24	22	17	13	7	-	-
Female	16	9	14	13	26	27	32	34	40	15	9	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	1	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Alaska												
All races, both sexes	-	1	1	4	3	6	12	9	8	4	1	-
Male	-	1	1	3	-	5	5	4	4	-	-	-
Female	-	-	-	1	3	1	7	5	4	4	1	-
White, both sexes	-	1	1	2	1	5	10	7	8	2	1	-
Male	-	1	1	1	-	4	4	2	4	-	-	-
Female	-	-	-	1	1	1	6	5	4	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	2	2	1	2	2	-	2	-	-
Male	-	-	-	2	-	1	1	2	-	-	-	-
Female	-	-	-	-	2	-	1	-	-	2	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2	8	16	22	44	71	111	175	141	61	15	-
Male	1	6	13	11	31	37	53	83	49	16	3	-
Female	1	2	3	11	13	34	58	92	92	45	12	-
White, both sexes	2	6	15	21	42	68	105	169	138	56	15	-
Male	1	5	12	10	30	35	53	78	49	14	3	-
Female	1	1	3	11	12	33	52	91	89	42	12	-
Black, both sexes	-	1	-	-	-	2	2	2	1	1	-	-
Male	-	-	-	-	-	2	-	2	-	1	-	-
Female	-	1	-	-	-	-	2	-	1	-	-	-
American Indian 1/, both sexes	-	1	1	1	2	1	2	3	2	4	-	-
Male	-	1	1	1	1	-	-	3	-	1	-	-
Female	-	-	-	-	1	1	2	-	2	3	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Arkansas												
All races, both sexes	8	26	28	30	44	83	120	131	136	81	13	-
Male	5	16	14	19	24	42	43	40	36	18	3	-
Female	3	10	14	11	20	41	77	91	100	63	10	-
White, both sexes	5	15	24	24	41	73	108	122	122	76	11	-
Male	2	10	13	16	22	35	39	38	32	17	2	-
Female	3	5	11	8	19	38	69	84	90	59	9	-
Black, both sexes	3	10	3	6	3	9	10	7	14	5	2	-
Male	3	6	1	3	2	6	3	2	4	1	1	-
Female	-	4	2	3	1	3	7	5	10	4	1	-
American Indian 1/, both sexes	-	1	1	-	-	1	1	2	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	1	-	-	-	1	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	44	64	109	153	264	421	767	948	883	397	92	-
Male	24	42	62	79	141	205	352	374	322	119	11	-
Female	20	22	47	74	123	216	415	574	561	278	81	-
White, both sexes	37	45	86	118	227	375	689	858	815	367	87	-
Male	22	30	48	60	120	182	321	342	294	108	11	-
Female	15	15	38	58	107	193	368	516	521	259	76	-
Black, both sexes	4	18	17	20	31	28	36	40	35	14	1	-
Male	1	11	10	11	18	12	15	15	12	4	-	-
Female	3	7	7	9	13	16	21	25	23	10	1	-
American Indian 1/, both sexes	1	1	1	4	-	3	5	1	2	-	-	-
Male	-	1	-	2	-	3	1	-	1	-	-	-
Female	1	-	1	2	-	-	4	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	5	11	6	15	37	49	31	16	4	-
Male	1	-	4	6	3	8	15	17	15	7	-	-
Female	1	-	1	5	3	7	22	32	16	9	4	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Colorado												
All races, both sexes	3	10	7	15	40	43	104	156	149	70	16	-
Male	2	5	4	7	23	18	41	60	44	17	5	-
Female	1	5	3	8	17	25	63	96	105	53	11	-
White, both sexes	3	8	5	13	38	43	100	151	144	67	16	-
Male	2	4	3	6	22	18	40	59	42	17	5	-
Female	1	4	2	7	16	25	60	92	102	50	11	-
Black, both sexes	-	2	1	2	2	-	2	3	2	2	-	-
Male	-	1	1	1	1	-	-	1	1	-	-	-
Female	-	1	-	1	1	-	2	2	1	2	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	2	2	1	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	1	2	1	1	-	-
Connecticut												
All races, both sexes	3	9	16	18	38	70	113	176	159	78	21	-
Male	1	4	10	9	21	38	53	78	57	25	6	-
Female	2	5	6	9	17	32	60	98	102	53	15	-
White, both sexes	2	6	13	16	35	66	110	172	154	76	21	-
Male	1	2	8	8	21	36	52	77	57	24	6	-
Female	1	4	5	8	14	30	58	95	97	52	15	-
Black, both sexes	1	3	3	2	3	4	2	4	5	2	-	-
Male	-	2	2	1	-	2	-	1	-	1	-	-
Female	1	1	1	1	3	2	2	3	5	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Delaware												
All races, both sexes	5	3	5	8	10	19	25	31	28	11	-	-
Male	3	3	3	4	7	5	8	14	7	4	-	-
Female	2	-	2	4	3	14	17	17	21	7	-	-
White, both sexes	3	-	3	8	9	17	19	28	24	10	-	-
Male	3	-	1	4	6	5	7	13	6	4	-	-
Female	-	-	2	4	3	12	12	15	18	6	-	-
Black, both sexes	2	2	2	-	1	2	6	3	4	1	-	-
Male	-	2	2	-	1	-	1	1	1	-	-	-
Female	2	-	-	-	-	2	5	2	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	3	2	2	3	1	8	11	23	7	2	-
Male	1	1	2	1	-	1	4	4	6	2	2	-
Female	-	2	-	1	3	-	4	7	17	5	-	-
White, both sexes	-	-	-	-	-	-	4	4	11	1	1	-
Male	-	-	-	-	-	-	1	2	4	-	1	-
Female	-	-	-	-	-	-	3	2	7	1	-	-
Black, both sexes	1	3	2	2	3	1	4	7	12	6	1	-
Male	1	1	2	1	-	1	3	2	2	2	1	-
Female	-	2	-	1	3	-	1	5	10	4	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Florida												
All races, both sexes	33	40	61	83	144	222	378	492	417	184	43	-
Male	20	26	36	55	78	112	176	221	142	55	11	-
Female	13	14	25	28	66	110	202	271	275	129	32	-
White, both sexes	20	36	51	66	124	198	350	468	397	177	37	-
Male	11	22	31	44	63	100	167	213	133	53	10	-
Female	9	14	20	22	61	98	183	255	264	124	27	-
Black, both sexes	13	4	10	15	19	23	26	22	20	6	5	-
Male	9	4	5	9	14	12	9	7	9	1	-	-
Female	4	-	5	6	5	11	17	15	11	5	5	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	1	-
Male	-	-	-	1	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	1	2	2	-	1	-	-
Male	-	-	-	1	1	-	-	1	-	1	-	-
Female	-	-	-	-	-	1	2	1	-	-	-	-
Georgia												
All races, both sexes	44	69	88	123	145	210	371	407	332	150	40	-
Male	27	55	52	55	81	100	146	151	90	27	10	-
Female	17	14	36	68	64	110	225	256	242	123	30	-
White, both sexes	26	33	52	67	101	158	295	344	288	122	30	-
Male	17	27	30	38	64	74	119	129	79	21	8	-
Female	9	6	22	29	37	84	176	215	209	101	22	-
Black, both sexes	18	36	36	55	44	49	73	61	42	26	10	-
Male	10	28	22	17	17	26	26	22	11	6	2	-
Female	8	8	14	38	27	23	47	39	31	20	8	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	3	3	1	2	2	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	3	2	1	2	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Hawaii												
All races, both sexes	1	3	3	8	7	15	24	36	25	11	5	-
Male	-	2	1	5	4	7	14	18	10	3	2	-
Female	1	1	2	3	3	8	10	18	15	8	3	-
White, both sexes	-	1	2	1	2	4	4	13	10	7	2	-
Male	-	1	1	1	1	-	3	7	5	-	2	-
Female	-	-	1	-	1	4	1	6	5	7	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	1	7	5	11	20	22	14	4	3	-
Male	-	1	-	4	3	7	11	10	5	3	-	-
Female	1	1	1	3	2	4	9	12	9	1	3	-
Idaho												
All races, both sexes	-	5	7	9	15	20	36	54	60	33	6	-
Male	-	2	6	5	5	8	19	17	22	7	-	-
Female	-	3	1	4	10	12	17	37	38	26	6	-
White, both sexes	-	5	7	9	15	20	34	53	59	33	6	-
Male	-	2	6	5	5	8	17	16	21	7	-	-
Female	-	3	1	4	10	12	17	37	38	26	6	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Illinois												
All races, both sexes	26	61	80	103	163	284	509	643	616	308	68	-
Male	16	35	44	64	79	128	222	236	199	64	7	-
Female	10	26	36	39	84	156	287	407	417	244	61	-
White, both sexes	12	42	55	76	123	245	457	602	583	288	61	-
Male	7	22	26	47	59	109	199	222	189	62	6	-
Female	5	20	29	29	64	136	258	380	394	226	55	-
Black, both sexes	14	19	25	27	38	38	46	40	31	19	7	-
Male	9	13	18	17	19	19	20	14	10	2	1	-
Female	5	6	7	10	19	19	26	26	21	17	6	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	1	5	1	2	1	-	-
Male	-	-	-	-	1	-	2	-	-	-	-	-
Female	-	-	-	-	1	1	3	1	2	1	-	-
Indiana												
All races, both sexes	23	35	52	68	119	208	286	391	361	183	46	-
Male	11	22	33	36	63	100	127	150	90	29	8	-
Female	12	13	19	32	56	108	159	241	271	154	38	-
White, both sexes	16	29	44	54	109	197	257	377	349	178	46	-
Male	8	18	28	30	58	98	114	144	88	27	8	-
Female	8	11	16	24	51	99	143	233	261	151	38	-
Black, both sexes	7	5	8	14	10	11	29	14	11	5	-	-
Male	3	3	5	6	5	2	13	6	2	2	-	-
Female	4	2	3	8	5	9	16	8	9	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	1	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Iowa												
All races, both sexes	1	2	7	4	16	30	36	78	68	41	9	-
Male	1	2	4	3	7	16	14	27	24	7	1	-
Female	-	-	3	1	9	14	22	51	44	34	8	-
White, both sexes	1	2	7	4	16	29	36	78	68	41	9	-
Male	1	2	4	3	7	15	14	27	24	7	1	-
Female	-	-	3	1	9	14	22	51	44	34	8	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	13	15	24	26	51	99	159	256	236	115	42	-
Male	6	9	15	13	34	42	64	92	79	27	4	-
Female	7	6	9	13	17	57	95	164	157	88	38	-
White, both sexes	9	14	20	22	46	90	152	250	229	114	38	-
Male	4	9	14	12	30	38	62	90	77	27	3	-
Female	5	5	6	10	16	52	90	160	152	87	35	-
Black, both sexes	3	1	3	2	4	8	6	5	5	-	4	-
Male	2	-	1	-	3	3	2	2	2	-	1	-
Female	1	1	2	2	1	5	4	3	3	-	3	-
American Indian 1/, both sexes	-	-	1	2	1	1	-	1	-	1	-	-
Male	-	-	-	1	1	1	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Kentucky												
All races, both sexes	13	36	47	78	83	153	254	257	231	76	19	-
Male	10	23	23	44	42	70	95	95	64	21	3	-
Female	3	13	24	34	41	83	159	162	167	55	16	-
White, both sexes	10	35	44	70	76	143	239	246	219	71	16	-
Male	8	23	21	41	39	67	90	89	58	20	3	-
Female	2	12	23	29	37	76	149	157	161	51	13	-
Black, both sexes	3	1	3	7	7	10	15	11	11	5	3	-
Male	2	-	2	2	3	3	5	6	5	1	-	-
Female	1	1	1	5	4	7	10	5	6	4	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	28	36	50	81	89	149	226	278	219	86	23	-
Male	14	22	25	52	46	75	85	99	58	14	4	-
Female	14	14	25	29	43	74	141	179	161	72	19	-
White, both sexes	13	20	27	51	62	108	182	237	178	72	19	-
Male	6	11	15	31	34	65	69	86	49	12	3	-
Female	7	9	12	20	28	43	113	151	129	60	16	-
Black, both sexes	15	16	22	30	27	41	44	41	40	13	4	-
Male	8	11	10	21	12	10	16	13	9	2	1	-
Female	7	5	12	9	15	31	28	28	31	11	3	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Maine												
All races, both sexes	2	5	5	6	22	27	58	73	47	29	6	-
Male	2	2	2	4	12	13	26	28	16	7	1	-
Female	-	3	3	2	10	14	32	45	31	22	5	-
White, both sexes	2	5	5	6	22	27	58	73	47	29	6	-
Male	2	2	2	4	12	13	26	28	16	7	1	-
Female	-	3	3	2	10	14	32	45	31	22	5	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	7	12	14	28	37	73	97	145	111	61	9	-
Male	5	6	9	13	19	36	46	52	41	9	2	-
Female	2	6	5	15	18	37	51	93	70	52	7	-
White, both sexes	2	6	9	16	28	53	80	126	98	48	7	-
Male	-	3	6	5	13	29	40	45	37	9	2	-
Female	2	3	3	11	15	24	40	81	61	39	5	-
Black, both sexes	5	6	5	11	9	18	17	17	12	12	2	-
Male	5	3	3	7	6	6	6	6	4	-	-	-
Female	-	3	2	4	3	12	11	11	8	12	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	2	-	2	-	1	-	-
Male	-	-	-	1	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Massachusetts												
All races, both sexes	9	11	25	37	79	153	295	394	365	158	49	-
Male	5	7	6	28	48	78	113	139	105	35	10	-
Female	4	4	19	9	31	75	182	255	260	123	39	-
White, both sexes	6	10	20	33	70	147	281	381	359	150	47	-
Male	3	6	4	26	44	74	109	135	103	32	9	-
Female	3	4	16	7	26	73	172	246	256	118	38	-
Black, both sexes	2	1	4	2	9	5	10	12	5	6	2	-
Male	1	1	2	1	4	3	4	3	2	1	1	-
Female	1	-	2	1	5	2	6	9	3	5	1	-
American Indian 1/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	1	-	1	3	1	1	2	-	-
Male	1	-	-	1	-	1	-	1	-	2	-	-
Female	-	-	1	-	-	-	3	-	1	-	-	-
Michigan												
All races, both sexes	18	37	45	85	129	253	461	571	509	232	52	-
Male	12	22	25	49	70	113	212	206	144	51	7	-
Female	6	15	20	36	59	140	249	365	365	181	45	-
White, both sexes	12	27	33	65	106	227	416	544	477	214	42	-
Male	6	16	19	38	60	102	198	197	132	47	5	-
Female	6	11	14	27	46	125	218	347	345	167	37	-
Black, both sexes	6	10	10	17	22	25	41	25	30	15	8	-
Male	6	6	4	10	10	10	13	9	11	4	1	-
Female	-	4	6	7	12	15	28	16	19	11	7	-
American Indian 1/, both sexes	-	-	-	-	1	1	1	1	1	-	1	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	1	1	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	2	3	-	-	3	1	1	3	1	-
Male	-	-	2	1	-	-	1	-	1	-	1	-
Female	-	-	-	2	-	-	2	1	-	3	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Minnesota												
All races, both sexes	4	10	18	22	36	86	149	271	275	153	33	-
Male	2	5	14	14	23	49	79	115	81	35	4	-
Female	2	5	4	8	13	37	70	156	194	118	29	-
White, both sexes	3	9	18	19	36	81	144	271	273	152	33	-
Male	2	4	14	12	23	46	75	115	79	35	4	-
Female	1	5	4	7	13	35	69	156	194	117	29	-
Black, both sexes	-	1	-	-	-	4	3	-	2	-	-	-
Male	-	1	-	-	-	3	2	-	2	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	3	-	-	2	-	-	-	-	-
Male	-	-	-	2	-	-	2	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	1	-	-
Mississippi												
All races, both sexes	28	39	42	80	99	160	225	281	203	124	30	1
Male	11	25	19	42	57	68	83	104	50	25	10	1
Female	17	14	23	38	42	92	142	177	153	99	20	-
White, both sexes	14	21	26	56	67	110	174	220	165	93	19	1
Male	6	15	13	30	39	45	70	87	45	18	5	1
Female	8	6	13	26	28	65	104	133	120	75	14	-
Black, both sexes	14	18	15	24	32	49	51	61	38	31	11	-
Male	5	10	6	12	18	22	13	17	5	7	5	-
Female	9	8	9	12	14	27	38	44	33	24	6	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Missouri												
All races, both sexes	11	22	44	60	92	149	231	332	301	147	43	-
Male	5	15	20	34	49	71	108	124	96	35	4	-
Female	6	7	24	26	43	78	123	208	205	112	39	-
White, both sexes	9	18	37	54	83	136	222	316	288	142	42	-
Male	4	13	17	30	45	66	105	120	88	33	3	-
Female	5	5	20	24	38	70	117	196	200	109	39	-
Black, both sexes	2	4	7	5	9	13	9	16	13	5	1	-
Male	1	2	3	3	4	5	3	4	8	2	1	-
Female	1	2	4	2	5	8	6	12	5	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	4	2	4	8	17	36	50	75	82	41	6	-
Male	4	1	2	5	9	19	30	26	29	13	2	-
Female	-	1	2	3	8	17	20	49	53	28	4	-
White, both sexes	4	1	4	7	15	34	48	75	82	41	6	-
Male	4	1	2	5	7	18	29	26	29	13	2	-
Female	-	-	2	2	8	16	19	49	53	28	4	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	1	2	2	1	-	-	-	-	-
Male	-	-	-	-	2	1	1	-	-	-	-	-
Female	-	1	-	1	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Nebraska												
All races, both sexes	4	13	11	16	25	41	73	93	114	56	16	-
Male	3	9	7	10	16	26	29	30	37	9	2	-
Female	1	4	4	6	9	15	44	63	77	47	14	-
White, both sexes	3	11	9	15	25	39	71	93	112	56	16	-
Male	3	8	6	10	16	24	29	30	37	9	2	-
Female	-	3	3	5	9	15	42	63	75	47	14	-
Black, both sexes	1	2	2	-	-	2	2	-	2	-	-	-
Male	-	1	1	-	-	2	-	-	-	-	-	-
Female	1	1	1	-	-	-	2	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	6	6	21	30	33	68	89	106	58	19	8	-
Male	6	4	14	21	21	43	56	43	19	6	1	-
Female	-	2	7	9	12	25	33	63	39	13	7	-
White, both sexes	5	4	17	26	31	62	86	99	56	19	7	-
Male	5	2	12	17	20	41	54	40	19	6	1	-
Female	-	2	5	9	11	21	32	59	37	13	6	-
Black, both sexes	-	2	2	2	2	4	2	-	1	-	-	-
Male	-	2	2	2	1	2	1	-	-	-	-	-
Female	-	-	-	-	1	2	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	1	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	2	1	-	1	1	6	-	-	1	-
Male	1	-	-	1	-	-	1	3	-	-	-	-
Female	-	-	2	-	-	1	-	3	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
New Hampshire												
All races, both sexes	-	3	5	7	9	28	44	50	47	25	13	-
Male	-	1	3	4	6	11	16	25	12	3	1	-
Female	-	2	2	3	3	17	28	25	35	22	12	-
White, both sexes	-	3	5	7	9	26	44	50	46	25	13	-
Male	-	1	3	4	6	11	16	25	12	3	1	-
Female	-	2	2	3	3	15	28	25	34	22	12	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-
New Jersey												
All races, both sexes	11	17	30	47	92	142	275	412	301	149	35	-
Male	8	10	18	22	56	67	111	171	100	42	9	-
Female	3	7	12	25	36	75	164	241	201	107	26	-
White, both sexes	6	8	22	37	68	117	255	378	292	138	32	-
Male	5	6	15	19	44	57	104	155	98	37	9	-
Female	1	2	7	18	24	60	151	223	194	101	23	-
Black, both sexes	5	9	8	10	22	23	19	31	7	8	3	-
Male	3	4	3	3	11	9	7	15	1	4	-	-
Female	2	5	5	7	11	14	12	16	6	4	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	2	1	3	2	3	-	-
Male	-	-	-	-	1	1	-	1	1	1	-	-
Female	-	-	-	-	1	1	1	2	1	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
New Mexico												
All races, both sexes	4	12	9	14	20	31	79	89	74	28	11	-
Male	2	6	4	10	14	12	36	31	28	5	4	-
Female	2	6	5	4	6	19	43	58	46	23	7	-
White, both sexes	4	10	9	14	18	29	77	88	72	26	9	-
Male	2	6	4	10	14	10	34	30	26	5	3	-
Female	2	4	5	4	4	19	43	58	46	21	6	-
Black, both sexes	-	-	-	-	-	-	1	-	-	2	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	2	-	-
American Indian 1/, both sexes	-	1	-	-	2	2	-	1	2	-	2	-
Male	-	-	-	-	-	2	-	1	2	-	1	-
Female	-	1	-	-	2	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	24	43	49	74	145	258	504	631	578	265	78	-
Male	16	30	31	47	81	124	219	241	203	65	8	-
Female	8	13	18	27	64	134	285	390	375	200	70	-
White, both sexes	21	26	37	65	127	223	471	597	548	254	73	-
Male	14	19	23	40	69	108	202	237	199	63	8	-
Female	7	7	14	25	58	115	269	360	349	191	65	-
Black, both sexes	3	16	12	8	14	32	28	32	29	9	5	-
Male	2	10	8	6	9	15	14	3	4	1	-	-
Female	1	6	4	2	5	17	14	29	25	8	5	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	4	3	5	2	1	2	-	-
Male	-	-	-	1	3	1	3	1	-	1	-	-
Female	-	-	-	-	1	2	2	1	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
North Carolina												
All races, both sexes	19	41	61	67	109	175	275	327	287	122	27	-
Male	12	20	36	43	63	89	119	110	84	20	-	-
Female	7	21	25	24	46	86	156	217	203	102	27	-
White, both sexes	12	27	45	55	84	138	240	280	257	102	24	-
Male	9	15	27	37	50	67	107	103	76	17	-	-
Female	3	12	18	18	34	71	133	177	181	85	24	-
Black, both sexes	7	12	16	12	25	36	35	43	30	20	2	-
Male	3	4	9	6	13	21	12	7	8	3	-	-
Female	4	8	7	6	12	15	23	36	22	17	2	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	-	2	-	-	1	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	1	-
North Dakota												
All races, both sexes	-	2	1	1	4	3	4	22	18	21	4	-
Male	-	2	1	1	3	3	1	10	5	3	-	-
Female	-	-	-	-	1	-	3	12	13	18	4	-
White, both sexes	-	-	-	1	4	3	4	21	18	21	4	-
Male	-	-	-	1	3	3	1	10	5	3	-	-
Female	-	-	-	-	1	-	3	11	13	18	4	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	2	1	-	-	-	-	1	-	-	-	-
Male	-	2	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Ohio												
All races, both sexes	19	22	46	56	99	176	292	343	307	113	23	-
Male	8	16	30	32	44	95	128	134	108	24	4	-
Female	11	6	16	24	55	81	164	209	199	89	19	-
White, both sexes	15	19	34	46	90	162	275	318	287	106	22	-
Male	7	15	21	26	40	89	120	126	102	22	3	-
Female	8	4	13	20	50	73	155	192	185	84	19	-
Black, both sexes	4	3	12	10	9	14	17	24	20	7	1	-
Male	1	1	9	6	4	6	8	8	6	2	1	-
Female	3	2	3	4	5	8	9	16	14	5	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Oklahoma												
All races, both sexes	27	35	57	83	111	164	250	289	227	97	22	-
Male	13	20	33	44	53	81	116	117	70	16	2	-
Female	14	15	24	39	58	83	134	172	157	81	20	-
White, both sexes	20	25	47	73	103	155	231	270	213	92	21	-
Male	11	14	27	39	49	75	108	114	65	15	2	-
Female	9	11	20	34	54	80	123	156	148	77	19	-
Black, both sexes	7	6	3	5	6	7	10	13	10	3	1	-
Male	2	4	2	3	4	5	5	1	3	-	-	-
Female	5	2	1	2	2	2	5	12	7	3	1	-
American Indian 1/, both sexes	-	4	5	5	2	2	8	6	4	1	-	-
Male	-	2	3	2	-	1	2	2	2	-	-	-
Female	-	2	2	3	2	1	6	4	2	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	-	-	1	-	-	1	-	-
Male	-	-	1	-	-	-	1	-	-	1	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Oregon												
All races, both sexes	5	10	14	28	33	61	125	168	155	80	18	-
Male	4	4	7	20	17	33	53	58	53	28	5	-
Female	1	6	7	8	16	28	72	110	102	52	13	-
White, both sexes	5	10	13	28	31	61	122	165	154	79	18	-
Male	4	4	6	20	16	33	52	57	53	28	5	-
Female	1	6	7	8	15	28	70	108	101	51	13	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	2	-	-	1	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	3	2	1	1	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	2	1	1	1	-	-
Pennsylvania												
All races, both sexes	28	58	84	115	148	331	588	839	699	286	71	-
Male	16	36	53	61	80	168	254	316	197	65	7	-
Female	12	22	31	54	68	163	334	523	502	221	64	-
White, both sexes	14	46	74	100	134	310	551	801	671	273	69	-
Male	10	28	46	53	71	160	240	305	191	62	6	-
Female	4	18	28	47	63	150	311	496	480	211	63	-
Black, both sexes	13	11	10	14	12	21	34	34	26	12	2	-
Male	5	7	7	7	8	8	14	9	5	2	1	-
Female	8	4	3	7	4	13	20	25	21	10	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	1	2	-	3	3	2	1	-	-
Male	1	1	-	1	1	-	-	2	1	1	-	-
Female	-	-	-	-	1	-	3	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Rhode Island												
All races, both sexes	-	1	2	4	8	12	24	38	25	19	3	-
Male	-	1	1	3	5	6	8	16	10	7	-	-
Female	-	-	1	1	3	6	16	22	15	12	3	-
White, both sexes	-	1	2	4	8	12	22	38	25	19	3	-
Male	-	1	1	3	5	6	8	16	10	7	-	-
Female	-	-	1	1	3	6	14	22	15	12	3	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
South Carolina												
All races, both sexes	21	39	44	60	76	107	164	172	128	64	13	-
Male	15	23	23	34	38	44	76	60	43	12	2	-
Female	6	16	21	26	38	63	88	112	85	52	11	-
White, both sexes	6	20	27	36	58	78	129	147	105	48	9	-
Male	4	15	12	24	30	34	60	56	37	9	2	-
Female	2	5	15	12	28	44	69	91	68	39	7	-
Black, both sexes	15	19	17	24	18	28	35	24	22	16	4	-
Male	11	8	11	10	8	9	16	4	6	3	-	-
Female	4	11	6	14	10	19	19	20	16	13	4	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
South Dakota												
All races, both sexes	3	-	-	4	6	5	5	12	15	9	3	-
Male	1	-	-	2	3	4	1	4	5	6	-	-
Female	2	-	-	2	3	1	4	8	10	3	3	-
White, both sexes	2	-	-	4	5	4	5	12	14	9	3	-
Male	1	-	-	2	3	3	1	4	5	6	-	-
Female	1	-	-	2	2	1	4	8	9	3	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	1	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	13	19	26	34	47	90	121	171	112	61	9	-
Male	7	16	18	14	26	40	51	44	25	5	-	-
Female	6	3	8	20	21	50	70	127	87	56	9	-
White, both sexes	8	14	21	25	41	77	102	152	105	52	9	-
Male	4	13	15	9	24	35	44	39	23	4	-	-
Female	4	1	6	16	17	42	58	113	82	48	9	-
Black, both sexes	5	5	5	9	6	12	19	19	7	7	-	-
Male	3	3	3	5	2	5	7	5	2	-	-	-
Female	2	2	2	4	4	7	12	14	5	7	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	2	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	1	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Texas												
All races, both sexes	83	107	163	212	299	528	738	929	708	343	78	-
Male	55	69	100	117	178	259	344	360	216	75	18	-
Female	28	38	63	95	121	269	394	569	492	268	60	-
White, both sexes	62	72	129	175	259	474	676	848	663	312	67	-
Male	43	47	81	99	152	233	317	331	203	67	14	-
Female	19	25	48	76	107	241	359	517	460	245	53	-
Black, both sexes	21	35	32	35	40	49	56	76	38	29	11	-
Male	12	22	18	17	26	23	23	26	12	7	4	-
Female	9	13	14	18	14	26	33	50	26	22	7	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	2	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	2	-	4	6	3	7	2	-	-
Male	-	-	1	1	-	2	4	2	1	1	-	-
Female	-	-	1	1	-	2	2	1	6	1	-	-
Utah												
All races, both sexes	3	9	11	22	36	65	125	128	138	67	3	-
Male	2	8	6	10	17	41	53	46	48	27	2	-
Female	1	1	5	12	19	24	72	82	90	40	1	-
White, both sexes	2	9	11	20	35	64	123	127	135	67	3	-
Male	1	8	6	10	16	40	51	45	48	27	2	-
Female	1	1	5	10	19	24	72	82	87	40	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	1	1	1	1	3	-	-	-
Male	-	-	-	-	1	1	1	1	-	-	-	-
Female	-	-	-	2	-	-	-	-	3	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Vermont												
All races, both sexes	-	1	2	1	1	2	6	14	10	7	2	-
Male	-	1	-	1	1	1	4	5	1	2	-	-
Female	-	-	2	-	-	1	2	9	9	5	2	-
White, both sexes	-	1	2	1	1	2	6	14	10	7	2	-
Male	-	1	-	1	1	1	4	5	1	2	-	-
Female	-	-	2	-	-	1	2	9	9	5	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	20	39	58	61	90	193	295	374	312	157	26	-
Male	15	23	31	38	42	96	140	158	83	32	3	-
Female	5	16	27	23	48	97	155	216	229	125	23	-
White, both sexes	10	19	33	46	64	150	234	331	273	137	25	-
Male	8	10	18	32	32	74	112	140	70	28	3	-
Female	2	9	15	14	32	76	122	191	203	109	22	-
Black, both sexes	10	20	25	15	23	42	55	42	35	17	1	-
Male	7	13	13	6	9	21	25	17	10	3	-	-
Female	3	7	12	9	14	21	30	25	25	14	1	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	3	-	6	1	4	2	-	-
Male	-	-	-	-	1	-	3	1	3	-	-	-
Female	-	-	-	-	2	-	3	-	1	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Washington												
All races, both sexes	3	4	13	16	17	34	79	94	99	43	10	-
Male	2	2	5	12	8	12	31	39	38	12	2	-
Female	1	2	8	4	9	22	48	55	61	31	8	-
White, both sexes	3	4	12	16	15	30	77	91	96	43	9	-
Male	2	2	4	12	7	11	31	38	36	12	2	-
Female	1	2	8	4	8	19	46	53	60	31	7	-
Black, both sexes	-	-	1	-	2	-	-	1	-	-	1	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	3	-	1	1	-	-	-
Male	-	-	-	-	-	1	-	1	1	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	2	1	2	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	1	2	1	1	-	-	-
West Virginia												
All races, both sexes	8	10	20	35	31	64	102	107	109	48	12	-
Male	4	4	14	21	16	29	47	42	29	9	2	-
Female	4	6	6	14	15	35	55	65	80	39	10	-
White, both sexes	8	10	20	34	31	62	98	103	108	47	11	-
Male	4	4	14	20	16	28	44	42	28	9	2	-
Female	4	6	6	14	15	34	54	61	80	38	9	-
Black, both sexes	-	-	-	1	-	2	4	4	1	1	1	-
Male	-	-	-	1	-	1	3	-	1	-	-	-
Female	-	-	-	-	-	1	1	4	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Wisconsin												
All races, both sexes	13	18	20	36	62	130	229	332	361	163	45	-
Male	9	13	13	23	34	68	102	146	109	43	5	-
Female	4	5	7	13	28	62	127	186	252	120	40	-
White, both sexes	9	16	18	36	57	127	224	325	357	160	44	-
Male	6	12	12	23	32	67	102	144	109	43	5	-
Female	3	4	6	13	25	60	122	181	248	117	39	-
Black, both sexes	4	2	2	-	5	1	4	5	4	2	1	-
Male	3	1	1	-	2	-	-	2	-	-	-	-
Female	1	1	1	-	3	1	4	3	4	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	2	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	2	-	1	-	-
Wyoming												
All races, both sexes	-	1	1	4	6	8	23	26	31	15	-	-
Male	-	1	-	3	2	3	10	8	7	7	-	-
Female	-	-	1	1	4	5	13	18	24	8	-	-
White, both sexes	-	1	1	4	6	8	23	25	30	15	-	-
Male	-	1	-	3	2	3	10	8	7	7	-	-
Female	-	-	1	1	4	5	13	17	23	8	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
United States												
All races, both sexes	51,967	8	6	3	2	10	17	20	42	82	172	311
Male	20,929	6	1	2	1	9	10	12	29	48	115	203
Female	31,038	2	5	1	1	1	7	8	13	34	57	108
White, both sexes	46,510	5	4	-	2	5	9	12	23	46	92	187
Male	18,607	4	-	-	1	4	4	6	16	27	59	122
Female	27,903	1	4	-	1	1	5	6	7	19	33	65
Black, both sexes	4,823	2	2	2	-	5	8	7	18	33	76	121
Male	2,051	2	1	1	-	5	6	5	12	18	52	79
Female	2,772	-	1	1	-	-	2	2	6	15	24	42
American Indian 1/, both sexes	174	-	-	-	-	-	-	-	1	-	2	2
Male	78	-	-	-	-	-	-	-	1	-	2	1
Female	96	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	460	1	-	1	-	-	-	1	-	3	2	1
Male	193	-	-	1	-	-	-	1	-	3	2	1
Female	267	1	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	1,740	-	-	-	-	-	1	-	2	3	14	24
Male	694	-	-	-	-	-	-	-	1	1	8	15
Female	1,046	-	-	-	-	-	1	-	1	2	6	9
White, both sexes	1,405	-	-	-	-	-	-	-	1	1	6	10
Male	553	-	-	-	-	-	-	-	1	1	3	4
Female	852	-	-	-	-	-	-	-	-	-	3	6
Black, both sexes	333	-	-	-	-	-	1	-	1	2	8	14
Male	140	-	-	-	-	-	-	-	-	-	5	11
Female	193	-	-	-	-	-	1	-	1	2	3	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Alaska												
All races, both sexes	45	-	-	-	-	-	-	-	-	-	1	-
Male	20	-	-	-	-	-	-	-	-	-	1	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	35	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	632	-	-	1	-	-	-	-	-	-	3	4
Male	291	-	-	-	-	-	-	-	-	-	1	2
Female	341	-	-	1	-	-	-	-	-	-	2	2
White, both sexes	603	-	-	-	-	-	-	-	-	-	2	4
Male	278	-	-	-	-	-	-	-	-	-	-	2
Female	325	-	-	-	-	-	-	-	-	-	2	2
Black, both sexes	10	-	-	1	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	16	-	-	-	-	-	-	-	-	-	1	-
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Arkansas												
All races, both sexes	673	-	1	-	-	-	-	1	-	1	3	5
Male	254	-	1	-	-	-	-	1	-	1	3	3
Female	419	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	595	-	-	-	-	-	-	-	-	1	2	3
Male	220	-	-	-	-	-	-	-	-	1	2	2
Female	375	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	71	-	1	-	-	-	-	1	-	-	1	2
Male	32	-	1	-	-	-	-	1	-	-	1	1
Female	39	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	4,057	-	1	1	-	-	-	2	4	4	13	24
Male	1,706	-	-	1	-	-	-	1	3	1	11	17
Female	2,351	-	1	-	-	-	-	1	1	3	2	7
White, both sexes	3,622	-	1	-	-	-	-	1	4	3	8	16
Male	1,514	-	-	-	-	-	-	-	3	-	6	11
Female	2,108	-	1	-	-	-	-	1	1	3	2	5
Black, both sexes	247	-	-	-	-	-	-	1	-	1	3	6
Male	112	-	-	-	-	-	-	1	-	1	3	5
Female	135	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	19	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	169	-	-	1	-	-	-	-	-	-	2	1
Male	72	-	-	1	-	-	-	-	-	-	2	1
Female	97	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Colorado												
All races, both sexes	575	-	-	-	-	-	-	1	-	1	1	-
Male	208	-	-	-	-	-	-	1	-	-	-	-
Female	367	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	552	-	-	-	-	-	-	1	-	1	-	-
Male	202	-	-	-	-	-	-	1	-	-	-	-
Female	350	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	15	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	601	-	1	-	-	1	-	-	-	-	-	2
Male	264	-	-	-	-	1	-	-	-	-	-	-
Female	337	-	1	-	-	-	-	-	-	-	-	2
White, both sexes	574	-	1	-	-	-	-	-	-	-	-	2
Male	255	-	-	-	-	-	-	-	-	-	-	-
Female	319	-	1	-	-	-	-	-	-	-	-	2
Black, both sexes	26	-	-	-	-	1	-	-	-	-	-	-
Male	8	-	-	-	-	1	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Delaware												
All races, both sexes	133	-	-	-	-	-	-	-	-	-	-	-
Male	51	-	-	-	-	-	-	-	-	-	-	-
Female	82	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	111	-	-	-	-	-	-	-	-	-	-	-
Male	44	-	-	-	-	-	-	-	-	-	-	-
Female	67	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	62	-	-	-	-	-	-	-	-	-	-	1
Male	24	-	-	-	-	-	-	-	-	-	-	1
Female	38	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	42	-	-	-	-	-	-	-	-	-	-	1
Male	16	-	-	-	-	-	-	-	-	-	-	1
Female	26	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Florida												
All races, both sexes	1,992	-	-	-	-	-	1	1	3	2	5	11
Male	886	-	-	-	-	-	-	-	2	1	4	6
Female	1,106	-	-	-	-	-	1	1	1	1	1	5
White, both sexes	1,816	-	-	-	-	-	1	1	3	-	1	3
Male	799	-	-	-	-	-	-	-	2	-	1	2
Female	1,017	-	-	-	-	-	1	1	1	-	-	1
Black, both sexes	166	-	-	-	-	-	-	-	-	2	4	8
Male	81	-	-	-	-	-	-	-	-	1	3	4
Female	85	-	-	-	-	-	-	-	-	1	1	4
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	1,824	-	-	-	-	2	1	-	2	9	14	24
Male	747	-	-	-	-	1	1	-	-	5	10	18
Female	1,077	-	-	-	-	1	-	-	2	4	4	6
White, both sexes	1,384	-	-	-	-	2	1	-	-	5	2	9
Male	562	-	-	-	-	1	1	-	-	4	1	8
Female	822	-	-	-	-	1	-	-	-	1	1	1
Black, both sexes	428	-	-	-	-	-	-	-	2	4	12	15
Male	184	-	-	-	-	-	-	-	-	1	9	10
Female	244	-	-	-	-	-	-	-	2	3	3	5
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Hawaii												
All races, both sexes	110	-	-	-	-	-	-	1	-	-	-	-
Male	53	-	-	-	-	-	-	1	-	-	-	-
Female	57	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	35	-	-	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	73	-	-	-	-	-	-	1	-	-	-	-
Male	35	-	-	-	-	-	-	1	-	-	-	-
Female	38	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	225	-	-	-	-	-	-	1	-	-	1	1
Male	83	-	-	-	-	-	-	1	-	-	-	1
Female	142	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	222	-	-	-	-	-	-	1	-	-	1	1
Male	80	-	-	-	-	-	-	1	-	-	-	1
Female	142	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Illinois												
All races, both sexes	2,747	2	-	-	-	-	-	2	1	7	11	18
Male	1,061	2	-	-	-	-	-	2	-	4	7	11
Female	1,686	-	-	-	-	-	-	-	1	3	4	7
White, both sexes	2,423	1	-	-	-	-	-	1	-	1	4	6
Male	905	1	-	-	-	-	-	1	-	1	3	3
Female	1,518	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	311	1	-	-	-	-	-	1	1	6	7	12
Male	152	1	-	-	-	-	-	1	-	3	4	8
Female	159	-	-	-	-	-	-	-	1	3	3	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1,697	1	1	-	-	1	-	1	2	1	10	6
Male	643	1	-	-	-	1	-	1	2	1	4	2
Female	1,054	-	1	-	-	-	-	-	-	-	6	4
White, both sexes	1,587	1	1	-	-	1	-	1	1	-	8	6
Male	596	1	-	-	-	1	-	1	1	-	4	2
Female	991	-	1	-	-	-	-	-	-	-	4	4
Black, both sexes	109	-	-	-	-	-	-	-	1	1	2	-
Male	46	-	-	-	-	-	-	-	1	1	-	-
Female	63	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Iowa												
All races, both sexes	264	-	-	-	-	-	-	-	-	1	1	2
Male	100	-	-	-	-	-	-	-	-	1	1	1
Female	164	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	263	-	-	-	-	-	-	-	-	1	1	2
Male	99	-	-	-	-	-	-	-	-	1	1	1
Female	164	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	995	-	-	-	-	-	1	-	-	-	1	2
Male	370	-	-	-	-	-	-	-	-	-	1	2
Female	625	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	946	-	-	-	-	-	1	-	-	-	1	2
Male	352	-	-	-	-	-	-	-	-	-	1	2
Female	594	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	38	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Kentucky												
All races, both sexes	1,184	-	-	-	-	-	1	1	-	5	2	9
Male	473	-	-	-	-	-	1	-	-	2	2	6
Female	711	-	-	-	-	-	-	1	-	3	-	3
White, both sexes	1,107	-	-	-	-	-	-	1	-	5	2	6
Male	441	-	-	-	-	-	-	-	-	2	2	5
Female	666	-	-	-	-	-	-	1	-	3	-	1
Black, both sexes	75	-	-	-	-	-	1	-	-	-	-	3
Male	30	-	-	-	-	-	1	-	-	-	-	1
Female	45	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1,248	1	-	-	-	-	2	-	5	4	6	9
Male	487	1	-	-	-	-	1	-	4	3	4	5
Female	761	-	-	-	-	-	1	-	1	1	2	4
White, both sexes	952	-	-	-	-	-	-	-	1	3	5	4
Male	372	-	-	-	-	-	-	-	1	2	4	2
Female	580	-	-	-	-	-	-	-	-	1	1	2
Black, both sexes	293	1	-	-	-	-	2	-	4	1	1	5
Male	115	1	-	-	-	-	1	-	3	1	-	3
Female	178	-	-	-	-	-	1	-	1	-	1	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Maine												
All races, both sexes	255	1	-	-	1	-	-	-	-	-	-	-
Male	104	1	-	-	-	-	-	-	-	-	-	-
Female	151	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	255	1	-	-	1	-	-	-	-	-	-	-
Male	104	1	-	-	-	-	-	-	-	-	-	-
Female	151	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	586	-	-	-	-	1	1	-	-	1	3	3
Male	237	-	-	-	-	1	1	-	-	1	2	2
Female	349	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	463	-	-	-	-	-	-	-	-	-	2	1
Male	185	-	-	-	-	-	-	-	-	-	2	-
Female	278	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	116	-	-	-	-	1	1	-	-	1	1	2
Male	49	-	-	-	-	1	1	-	-	1	-	2
Female	67	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Massachusetts												
All races, both sexes	1,440	-	-	-	-	1	-	-	-	2	1	2
Male	523	-	-	-	-	1	-	-	-	2	-	2
Female	917	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	1,380	-	-	-	-	1	-	-	-	1	1	2
Male	500	-	-	-	-	1	-	-	-	1	-	2
Female	880	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	48	-	-	-	-	-	-	-	-	-	-	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	2,275	-	-	-	-	-	-	-	2	3	5	12
Male	862	-	-	-	-	-	-	-	1	2	3	8
Female	1,413	-	-	-	-	-	-	-	1	1	2	4
White, both sexes	2,063	-	-	-	-	-	-	-	-	-	3	9
Male	776	-	-	-	-	-	-	-	-	-	2	6
Female	1,287	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	196	-	-	-	-	-	-	-	2	3	2	3
Male	81	-	-	-	-	-	-	-	1	2	1	2
Female	115	-	-	-	-	-	-	-	1	1	1	1
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Minnesota												
All races, both sexes	969	-	-	-	-	-	-	-	1	-	-	1
Male	381	-	-	-	-	-	-	-	1	-	-	1
Female	588	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	950	-	-	-	-	-	-	-	-	-	-	1
Male	368	-	-	-	-	-	-	-	-	-	-	1
Female	582	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	11	-	-	-	-	-	-	-	1	-	-	-
Male	9	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1,211	-	-	-	-	-	3	2	-	5	9	15
Male	464	-	-	-	-	-	2	1	-	3	4	9
Female	747	-	-	-	-	-	1	1	-	2	5	6
White, both sexes	886	-	-	-	-	-	1	-	-	1	3	8
Male	342	-	-	-	-	-	-	-	-	-	-	4
Female	544	-	-	-	-	-	1	-	-	1	3	4
Black, both sexes	323	-	-	-	-	-	2	2	-	4	6	7
Male	121	-	-	-	-	-	2	1	-	3	4	5
Female	202	-	-	-	-	-	-	1	-	1	2	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Missouri												
All races, both sexes	1,354	-	-	-	-	1	-	-	1	2	4	7
Male	533	-	-	-	-	1	-	-	1	2	3	6
Female	821	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	1,271	-	-	-	-	-	-	-	-	2	2	5
Male	493	-	-	-	-	-	-	-	-	2	1	5
Female	778	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	82	-	-	-	-	1	-	-	1	-	2	2
Male	39	-	-	-	-	1	-	-	1	-	2	1
Female	43	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	285	-	-	-	-	-	-	-	-	-	-	1
Male	126	-	-	-	-	-	-	-	-	-	-	-
Female	159	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	277	-	-	-	-	-	-	-	-	-	-	1
Male	122	-	-	-	-	-	-	-	-	-	-	-
Female	155	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Nebraska												
All races, both sexes	395	-	-	-	-	-	-	-	-	1	-	-
Male	145	-	-	-	-	-	-	-	-	1	-	-
Female	250	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	384	-	-	-	-	-	-	-	-	1	-	-
Male	142	-	-	-	-	-	-	-	-	1	-	-
Female	242	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	418	-	-	-	-	-	-	-	-	1	-	2
Male	219	-	-	-	-	-	-	-	-	-	-	2
Female	199	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	390	-	-	-	-	-	-	-	-	1	-	2
Male	203	-	-	-	-	-	-	-	-	-	-	2
Female	187	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
New Hampshire												
All races, both sexes	215	-	-	-	-	-	-	-	-	-	-	-
Male	76	-	-	-	-	-	-	-	-	-	-	-
Female	139	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	212	-	-	-	-	-	-	-	-	-	-	-
Male	76	-	-	-	-	-	-	-	-	-	-	-
Female	136	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1,386	-	-	1	-	-	1	-	-	-	1	4
Male	556	-	-	1	-	-	1	-	-	-	-	2
Female	830	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	1,242	-	-	-	-	-	-	-	-	-	-	2
Male	498	-	-	-	-	-	-	-	-	-	-	1
Female	744	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	132	-	-	1	-	-	1	-	-	-	1	2
Male	53	-	-	1	-	-	1	-	-	-	-	1
Female	79	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
New Mexico												
All races, both sexes	362	-	-	-	-	-	-	-	-	1	-	2
Male	153	-	-	-	-	-	-	-	-	1	-	2
Female	209	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	347	-	-	-	-	-	-	-	-	1	-	2
Male	145	-	-	-	-	-	-	-	-	1	-	2
Female	202	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	2,416	1	-	-	1	1	-	1	1	-	7	8
Male	976	1	-	-	1	1	-	-	1	-	5	5
Female	1,440	-	-	-	-	-	-	1	-	-	2	3
White, both sexes	2,224	1	-	-	1	1	-	1	1	-	5	5
Male	897	1	-	-	1	1	-	-	1	-	4	5
Female	1,327	-	-	-	-	-	-	1	-	-	1	-
Black, both sexes	174	-	-	-	-	-	-	-	-	-	2	3
Male	69	-	-	-	-	-	-	-	-	-	1	-
Female	105	-	-	-	-	-	-	-	-	-	1	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
North Carolina												
All races, both sexes	1,403	-	-	-	-	-	-	-	1	1	5	9
Male	556	-	-	-	-	-	-	-	1	-	5	7
Female	847	-	-	-	-	-	-	-	-	1	-	2
White, both sexes	1,171	-	-	-	-	-	-	-	-	-	-	6
Male	470	-	-	-	-	-	-	-	-	-	-	5
Female	701	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	224	-	-	-	-	-	-	-	1	1	5	3
Male	84	-	-	-	-	-	-	-	1	-	5	2
Female	140	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	75	-	-	-	-	-	-	-	1	-	-	-
Male	26	-	-	-	-	-	-	-	-	-	-	-
Female	49	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	71	-	-	-	-	-	-	-	1	-	-	-
Male	23	-	-	-	-	-	-	-	-	-	-	-
Female	48	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Ohio												
All races, both sexes	1,412	-	-	-	-	-	1	-	2	1	5	6
Male	588	-	-	-	-	-	1	-	2	-	4	4
Female	824	-	-	-	-	-	-	-	-	1	1	2
White, both sexes	1,293	-	-	-	-	-	1	-	1	-	4	4
Male	537	-	-	-	-	-	1	-	1	-	3	4
Female	756	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	118	-	-	-	-	-	-	-	1	1	1	2
Male	51	-	-	-	-	-	-	-	1	-	1	-
Female	67	-	-	-	-	-	-	-	-	1	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1,305	-	1	-	-	-	-	1	3	2	7	14
Male	557	-	-	-	-	-	-	1	3	1	5	8
Female	748	-	1	-	-	-	-	-	-	1	2	6
White, both sexes	1,191	-	1	-	-	-	-	1	2	1	5	10
Male	505	-	-	-	-	-	-	1	2	-	4	5
Female	686	-	1	-	-	-	-	-	-	1	1	5
Black, both sexes	74	-	-	-	-	-	-	-	1	1	2	3
Male	34	-	-	-	-	-	-	-	1	1	1	2
Female	40	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	36	-	-	-	-	-	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	21	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Oregon												
All races, both sexes	645	1	-	-	-	-	-	-	1	1	1	1
Male	254	-	-	-	-	-	-	-	-	-	-	1
Female	391	1	-	-	-	-	-	-	1	1	1	-
White, both sexes	635	-	-	-	-	-	-	-	1	1	1	1
Male	252	-	-	-	-	-	-	-	-	-	-	1
Female	383	-	-	-	-	-	-	-	1	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	1	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	2,999	1	-	-	-	-	1	1	2	2	4	20
Male	1,155	-	-	-	-	-	-	-	2	1	3	11
Female	1,844	1	-	-	-	-	1	1	-	1	1	9
White, both sexes	2,803	1	-	-	-	-	1	1	1	2	2	12
Male	1,072	-	-	-	-	-	-	-	1	1	1	6
Female	1,731	1	-	-	-	-	1	1	-	1	1	6
Black, both sexes	184	-	-	-	-	-	-	-	1	-	2	8
Male	77	-	-	-	-	-	-	-	1	-	2	5
Female	107	-	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Rhode Island												
All races, both sexes	133	-	-	-	-	-	-	-	-	-	-	-
Male	56	-	-	-	-	-	-	-	-	-	-	-
Female	77	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	131	-	-	-	-	-	-	-	-	-	-	-
Male	56	-	-	-	-	-	-	-	-	-	-	-
Female	75	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	785	-	-	-	-	1	1	-	-	1	6	8
Male	328	-	-	-	-	1	1	-	-	-	5	5
Female	457	-	-	-	-	-	-	-	-	1	1	3
White, both sexes	593	-	-	-	-	-	1	-	-	-	2	5
Male	254	-	-	-	-	-	1	-	-	-	2	3
Female	339	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	189	-	-	-	-	1	-	-	-	1	4	3
Male	73	-	-	-	-	1	-	-	-	-	3	2
Female	116	-	-	-	-	-	-	-	-	1	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
South Dakota												
All races, both sexes	56	-	-	-	-	-	-	-	1	-	-	-
Male	23	-	-	-	-	-	-	-	1	-	-	-
Female	33	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	51	-	-	-	-	-	-	-	-	-	-	-
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	640	-	-	-	-	-	-	-	-	3	7	5
Male	227	-	-	-	-	-	-	-	-	2	4	4
Female	413	-	-	-	-	-	-	-	-	1	3	1
White, both sexes	554	-	-	-	-	-	-	-	-	2	4	3
Male	188	-	-	-	-	-	-	-	-	1	1	2
Female	366	-	-	-	-	-	-	-	-	1	3	1
Black, both sexes	83	-	-	-	-	-	-	-	-	1	3	2
Male	38	-	-	-	-	-	-	-	-	1	3	2
Female	45	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Texas												
All races, both sexes	3,899	-	-	-	-	1	2	3	5	11	12	31
Male	1,667	-	-	-	-	1	1	1	2	9	8	22
Female	2,232	-	-	-	-	-	1	2	3	2	4	9
White, both sexes	3,478	-	-	-	-	-	2	2	4	8	9	22
Male	1,470	-	-	-	-	-	1	1	1	6	6	16
Female	2,008	-	-	-	-	-	1	1	3	2	3	6
Black, both sexes	395	-	-	-	-	1	-	1	1	2	3	9
Male	185	-	-	-	-	1	-	-	1	2	2	6
Female	210	-	-	-	-	-	-	1	-	-	1	3
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	23	-	-	-	-	-	-	-	-	1	-	-
Male	10	-	-	-	-	-	-	-	-	1	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	584	-	-	-	-	-	-	-	-	2	1	1
Male	249	-	-	-	-	-	-	-	-	1	1	-
Female	335	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	574	-	-	-	-	-	-	-	-	2	1	1
Male	244	-	-	-	-	-	-	-	-	1	1	-
Female	330	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Vermont												
All races, both sexes	43	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	43	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1,301	-	-	-	-	-	-	1	-	2	6	6
Male	530	-	-	-	-	-	-	1	-	1	5	4
Female	771	-	-	-	-	-	-	-	-	1	1	2
White, both sexes	1,080	-	-	-	-	-	-	-	-	1	3	2
Male	433	-	-	-	-	-	-	-	-	1	3	1
Female	647	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	205	-	-	-	-	-	-	1	-	1	3	4
Male	88	-	-	-	-	-	-	1	-	-	2	3
Female	117	-	-	-	-	-	-	-	-	1	1	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Washington												
All races, both sexes	382	-	-	-	-	-	-	-	-	-	-	1
Male	152	-	-	-	-	-	-	-	-	-	-	-
Female	230	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	366	-	-	-	-	-	-	-	-	-	-	1
Male	146	-	-	-	-	-	-	-	-	-	-	-
Female	220	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	447	-	-	-	-	-	-	-	1	2	1	2
Male	187	-	-	-	-	-	-	-	1	1	-	2
Female	260	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	433	-	-	-	-	-	-	-	1	1	1	2
Male	180	-	-	-	-	-	-	-	1	-	-	2
Female	253	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Wisconsin												
All races, both sexes	1,381	-	1	-	-	-	-	-	1	-	1	8
Male	553	-	-	-	-	-	-	-	1	-	1	6
Female	828	-	1	-	-	-	-	-	-	-	-	2
White, both sexes	1,343	-	-	-	-	-	-	-	1	-	1	6
Male	542	-	-	-	-	-	-	-	1	-	1	4
Female	801	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	32	-	1	-	-	-	-	-	-	-	-	2
Male	10	-	-	-	-	-	-	-	-	-	-	2
Female	22	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	106	-	-	-	-	-	-	-	-	-	-	-
Male	36	-	-	-	-	-	-	-	-	-	-	-
Female	70	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	104	-	-	-	-	-	-	-	-	-	-	-
Male	36	-	-	-	-	-	-	-	-	-	-	-
Female	68	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
United States												
All races, both sexes	591	962	1,405	2,035	3,111	5,464	9,208	12,003	10,507	4,860	1,147	1
Male	351	602	794	1,165	1,687	2,657	4,039	4,581	3,292	1,147	177	1
Female	240	360	611	870	1,424	2,807	5,169	7,422	7,215	3,713	970	-
White, both sexes	381	665	1,100	1,638	2,624	4,792	8,352	11,168	9,856	4,509	1,039	1
Male	235	424	625	958	1,437	2,346	3,706	4,311	3,103	1,064	154	1
Female	146	241	475	680	1,187	2,446	4,646	6,857	6,753	3,445	885	-
Black, both sexes	201	279	283	349	447	601	726	707	563	299	94	-
Male	113	168	160	180	230	273	281	222	158	65	20	-
Female	88	111	123	169	217	328	445	485	405	234	74	-
American Indian 1/, both sexes	3	13	11	21	15	23	25	26	19	9	4	-
Male	-	7	5	11	5	16	10	9	7	2	2	-
Female	3	6	6	10	10	7	15	17	12	7	2	-
Asian or Pacific Islander 2/, both sexes	6	5	11	27	25	48	105	102	69	43	10	-
Male	3	3	4	16	15	22	42	39	24	16	1	-
Female	3	2	7	11	10	26	63	63	45	27	9	-
Alabama												
All races, both sexes	36	50	57	81	137	197	304	386	300	122	26	-
Male	16	29	26	50	79	99	123	130	89	27	1	-
Female	20	21	31	31	58	98	181	256	211	95	25	-
White, both sexes	14	30	38	62	100	157	261	348	255	104	18	-
Male	7	14	19	40	61	79	106	118	78	21	1	-
Female	7	16	19	22	39	78	155	230	177	83	17	-
Black, both sexes	22	20	19	19	37	39	43	38	45	18	7	-
Male	9	15	7	10	18	19	17	12	11	6	-	-
Female	13	5	12	9	19	20	26	26	34	12	7	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	1	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Alaska												
All races, both sexes	-	1	1	3	3	4	12	8	7	4	1	-
Male	-	1	1	2	-	3	5	4	3	-	-	-
Female	-	-	-	1	3	1	7	4	4	4	1	-
White, both sexes	-	1	1	2	1	4	10	6	7	2	1	-
Male	-	1	1	1	-	3	4	2	3	-	-	-
Female	-	-	-	1	1	1	6	4	4	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	2	-	2	2	-	2	-	-
Male	-	-	-	1	-	-	1	2	-	-	-	-
Female	-	-	-	-	2	-	1	-	-	2	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2	7	15	20	41	67	103	166	131	59	13	-
Male	1	5	13	11	29	36	49	79	48	14	3	-
Female	1	2	2	9	12	31	54	87	83	45	10	-
White, both sexes	2	5	14	19	39	64	98	161	128	54	13	-
Male	1	4	12	10	28	34	49	75	48	12	3	-
Female	1	1	2	9	11	30	49	86	80	42	10	-
Black, both sexes	-	1	-	-	-	2	2	2	1	1	-	-
Male	-	-	-	-	-	2	-	2	-	1	-	-
Female	-	1	-	-	-	-	2	-	1	-	-	-
American Indian 1/, both sexes	-	1	1	1	2	1	1	2	2	4	-	-
Male	-	1	1	1	1	-	-	2	-	1	-	-
Female	-	-	-	-	1	1	1	-	2	3	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Arkansas												
All races, both sexes	8	23	28	28	42	76	114	129	124	77	13	-
Male	5	14	14	19	22	38	41	39	33	17	3	-
Female	3	9	14	9	20	38	73	90	91	60	10	-
White, both sexes	5	14	24	23	39	66	103	121	111	72	11	-
Male	2	10	13	16	20	31	38	37	30	16	2	-
Female	3	4	11	7	19	35	65	84	81	56	9	-
Black, both sexes	3	8	3	5	3	9	9	6	13	5	2	-
Male	3	4	1	3	2	6	2	2	3	1	1	-
Female	-	4	2	2	1	3	7	4	10	4	1	-
American Indian 1/, both sexes	-	1	1	-	-	1	1	2	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	1	-	-	-	1	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	44	61	105	148	251	405	739	929	857	381	88	-
Male	24	39	59	75	135	194	339	365	313	118	11	-
Female	20	22	46	73	116	211	400	564	544	263	77	-
White, both sexes	37	42	84	115	215	361	664	840	794	354	83	-
Male	22	27	47	58	115	172	311	334	289	108	11	-
Female	15	15	37	57	100	189	353	506	505	246	72	-
Black, both sexes	4	18	16	19	30	27	33	39	35	14	1	-
Male	1	11	9	10	17	12	12	14	12	4	-	-
Female	3	7	7	9	13	15	21	25	23	10	1	-
American Indian 1/, both sexes	1	1	1	4	-	3	5	1	2	-	-	-
Male	-	1	-	2	-	3	1	-	1	-	-	-
Female	1	-	1	2	-	-	4	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	4	10	6	14	37	49	26	13	4	-
Male	1	-	3	5	3	7	15	17	11	6	-	-
Female	1	-	1	5	3	7	22	32	15	7	4	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Colorado												
All races, both sexes	2	9	6	15	36	39	95	144	145	66	15	-
Male	2	4	4	7	19	15	39	56	43	14	4	-
Female	-	5	2	8	17	24	56	88	102	52	11	-
White, both sexes	2	8	5	13	34	39	91	140	140	63	15	-
Male	2	4	3	6	18	15	38	56	41	14	4	-
Female	-	4	2	7	16	24	53	84	99	49	11	-
Black, both sexes	-	1	1	2	2	-	2	2	2	2	-	-
Male	-	-	1	1	1	-	-	-	1	-	-	-
Female	-	1	-	1	1	-	2	2	1	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	2	2	1	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	1	2	1	1	-	-
Connecticut												
All races, both sexes	1	6	11	13	31	54	98	156	145	65	17	-
Male	-	2	7	6	17	33	46	73	55	20	4	-
Female	1	4	4	7	14	21	52	83	90	45	13	-
White, both sexes	1	4	8	12	28	50	95	152	141	63	17	-
Male	-	1	5	6	17	31	45	72	55	19	4	-
Female	1	3	3	6	11	19	50	80	86	44	13	-
Black, both sexes	-	2	3	1	3	4	2	4	4	2	-	-
Male	-	1	2	-	-	2	-	1	-	1	-	-
Female	-	1	1	1	3	2	2	3	4	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Delaware												
All races, both sexes	4	2	5	8	7	17	25	29	25	11	-	-
Male	2	2	3	4	4	3	8	14	7	4	-	-
Female	2	-	2	4	3	14	17	15	18	7	-	-
White, both sexes	2	-	3	8	7	15	19	26	21	10	-	-
Male	2	-	1	4	4	3	7	13	6	4	-	-
Female	-	-	2	4	3	12	12	13	15	6	-	-
Black, both sexes	2	2	2	-	-	2	6	3	4	1	-	-
Male	-	2	2	-	-	-	1	1	1	-	-	-
Female	2	-	-	-	-	2	5	2	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	3	2	2	3	1	7	11	22	7	2	-
Male	1	1	2	1	-	1	3	4	6	2	2	-
Female	-	2	-	1	3	-	4	7	16	5	-	-
White, both sexes	-	-	-	-	-	-	4	4	10	1	1	-
Male	-	-	-	-	-	-	1	2	4	-	1	-
Female	-	-	-	-	-	-	3	2	6	1	-	-
Black, both sexes	1	3	2	2	3	1	3	7	12	6	1	-
Male	1	1	2	1	-	1	2	2	2	2	1	-
Female	-	2	-	1	3	-	1	5	10	4	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Florida												
All races, both sexes	28	36	59	76	135	208	356	469	392	169	41	-
Male	17	24	34	52	72	103	166	209	132	53	11	-
Female	11	12	25	24	63	105	190	260	260	116	30	-
White, both sexes	17	32	50	59	117	185	329	447	373	163	35	-
Male	10	20	30	41	58	91	157	202	124	51	10	-
Female	7	12	20	18	59	94	172	245	249	112	25	-
Black, both sexes	11	4	9	15	17	22	25	20	19	5	5	-
Male	7	4	4	9	13	12	9	6	8	1	-	-
Female	4	-	5	6	4	10	16	14	11	4	5	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	1	-
Male	-	-	-	1	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	1	2	2	-	1	-	-
Male	-	-	-	1	1	-	-	1	-	1	-	-
Female	-	-	-	-	-	1	2	1	-	-	-	-
Georgia												
All races, both sexes	40	56	79	105	125	194	338	368	300	132	35	-
Male	25	46	47	45	68	96	132	138	81	26	8	-
Female	15	10	32	60	57	98	206	230	219	106	27	-
White, both sexes	23	27	48	59	88	145	267	312	264	106	26	-
Male	15	23	28	31	56	71	107	118	72	20	6	-
Female	8	4	20	28	32	74	160	194	192	86	20	-
Black, both sexes	17	29	31	46	37	46	68	54	34	24	9	-
Male	10	23	19	14	12	25	24	20	9	6	2	-
Female	7	6	12	32	25	21	44	34	25	18	7	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	3	3	1	2	2	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	3	2	1	2	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Hawaii												
All races, both sexes	1	2	3	5	6	9	20	25	23	10	5	-
Male	-	1	1	4	4	5	10	12	10	3	2	-
Female	1	1	2	1	2	4	10	13	13	7	3	-
White, both sexes	-	1	2	1	1	1	4	8	9	6	2	-
Male	-	1	1	1	1	-	3	3	5	-	2	-
Female	-	-	1	-	-	1	1	5	4	6	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	4	5	8	16	16	13	4	3	-
Male	-	-	-	3	3	5	7	8	5	3	-	-
Female	1	1	1	1	2	3	9	8	8	1	3	-
Idaho												
All races, both sexes	-	2	5	9	15	19	33	48	55	30	6	-
Male	-	-	4	5	5	8	18	15	19	7	-	-
Female	-	2	1	4	10	11	15	33	36	23	6	-
White, both sexes	-	2	5	9	15	19	31	47	55	30	6	-
Male	-	-	4	5	5	8	16	14	19	7	-	-
Female	-	2	1	4	10	11	15	33	36	23	6	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Illinois												
All races, both sexes	25	50	71	96	148	273	475	615	590	298	65	-
Male	16	28	40	59	72	124	211	226	193	61	5	-
Female	9	22	31	37	76	149	264	389	397	237	60	-
White, both sexes	12	32	50	72	111	236	425	577	559	278	58	-
Male	7	16	25	43	52	106	188	213	183	59	4	-
Female	5	16	25	29	59	130	237	364	376	219	54	-
Black, both sexes	13	18	21	24	35	36	44	37	29	19	7	-
Male	9	12	15	16	19	18	20	13	10	2	1	-
Female	4	6	6	8	16	18	24	24	19	17	6	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	1	5	1	2	1	-	-
Male	-	-	-	-	1	-	2	-	-	-	-	-
Female	-	-	-	-	1	1	3	1	2	1	-	-
Indiana												
All races, both sexes	19	31	49	62	111	194	267	378	347	172	44	-
Male	10	21	32	33	58	94	118	145	86	27	7	-
Female	9	10	17	29	53	100	149	233	261	145	37	-
White, both sexes	13	26	42	50	101	183	239	365	337	168	44	-
Male	7	17	27	28	53	92	106	139	84	26	7	-
Female	6	9	15	22	48	91	133	226	253	142	37	-
Black, both sexes	6	4	7	12	10	11	28	13	10	4	-	-
Male	3	3	5	5	5	2	12	6	2	1	-	-
Female	3	1	2	7	5	9	16	7	8	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Iowa												
All races, both sexes	1	2	7	4	15	27	30	70	61	37	6	-
Male	1	2	4	3	7	13	12	24	23	7	1	-
Female	-	-	3	1	8	14	18	46	38	30	5	-
White, both sexes	1	2	7	4	15	26	30	70	61	37	6	-
Male	1	2	4	3	7	12	12	24	23	7	1	-
Female	-	-	3	1	8	14	18	46	38	30	5	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	13	13	22	25	47	96	151	246	226	110	42	-
Male	6	8	13	12	30	41	61	89	76	27	4	-
Female	7	5	9	13	17	55	90	157	150	83	38	-
White, both sexes	9	12	18	21	43	88	144	240	220	109	38	-
Male	4	8	12	11	27	37	59	87	74	27	3	-
Female	5	4	6	10	16	51	85	153	146	82	35	-
Black, both sexes	3	1	3	2	3	7	6	5	4	-	4	-
Male	2	-	1	-	2	3	2	2	2	-	1	-
Female	1	1	2	2	1	4	4	3	2	-	3	-
American Indian 1/, both sexes	-	-	1	2	1	1	-	1	-	1	-	-
Male	-	-	-	1	1	1	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Kentucky												
All races, both sexes	12	33	40	73	77	147	233	238	226	69	18	-
Male	9	21	18	41	38	66	91	90	64	21	3	-
Female	3	12	22	32	39	81	142	148	162	48	15	-
White, both sexes	9	32	38	66	70	137	219	228	214	65	15	-
Male	7	21	17	38	35	63	86	84	58	20	3	-
Female	2	11	21	28	35	74	133	144	156	45	12	-
Black, both sexes	3	1	2	6	7	10	14	10	11	4	3	-
Male	2	-	1	2	3	3	5	6	5	1	-	-
Female	1	1	1	4	4	7	9	4	6	3	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	28	31	44	74	86	144	219	271	216	85	23	-
Male	14	19	21	48	44	70	83	95	57	14	4	-
Female	14	12	23	26	42	74	136	176	159	71	19	-
White, both sexes	13	18	24	47	61	104	177	230	175	71	19	-
Male	6	11	12	28	33	61	67	82	48	12	3	-
Female	7	7	12	19	28	43	110	148	127	59	16	-
Black, both sexes	15	13	19	27	25	40	42	41	40	13	4	-
Male	8	8	9	20	11	9	16	13	9	2	1	-
Female	7	5	10	7	14	31	26	28	31	11	3	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Maine												
All races, both sexes	2	5	5	4	18	23	53	66	45	27	5	-
Male	2	2	2	3	9	12	24	25	16	7	1	-
Female	-	3	3	1	9	11	29	41	29	20	4	-
White, both sexes	2	5	5	4	18	23	53	66	45	27	5	-
Male	2	2	2	3	9	12	24	25	16	7	1	-
Female	-	3	3	1	9	11	29	41	29	20	4	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	6	11	14	28	34	70	94	141	111	59	9	-
Male	5	5	9	13	18	34	45	49	41	9	2	-
Female	1	6	5	15	16	36	49	92	70	50	7	-
White, both sexes	1	6	9	16	25	51	77	123	98	47	7	-
Male	-	3	6	5	12	27	39	43	37	9	2	-
Female	1	3	3	11	13	24	38	80	61	38	5	-
Black, both sexes	5	5	5	11	9	17	17	16	12	11	2	-
Male	5	2	3	7	6	6	6	5	4	-	-	-
Female	-	3	2	4	3	11	11	11	8	11	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	2	-	2	-	1	-	-
Male	-	-	-	1	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Massachusetts												
All races, both sexes	5	6	18	29	70	139	275	358	344	145	45	-
Male	4	4	5	22	42	71	105	125	98	32	10	-
Female	1	2	13	7	28	68	170	233	246	113	35	-
White, both sexes	3	6	14	27	62	133	262	348	338	139	43	-
Male	2	4	4	22	38	67	101	122	96	31	9	-
Female	1	2	10	5	24	66	161	226	242	108	34	-
Black, both sexes	1	-	3	1	8	5	9	9	5	5	2	-
Male	1	-	1	-	4	3	4	2	2	-	1	-
Female	-	-	2	1	4	2	5	7	3	5	1	-
American Indian 1/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	1	3	1	1	1	-	-
Male	1	-	-	-	-	1	-	1	-	1	-	-
Female	-	-	1	-	-	-	3	-	1	-	-	-
Michigan												
All races, both sexes	14	33	39	79	117	231	432	549	490	223	46	-
Male	9	19	21	47	62	101	202	197	134	50	6	-
Female	5	14	18	32	55	130	230	352	356	173	40	-
White, both sexes	9	24	30	63	98	208	395	523	459	205	37	-
Male	4	14	17	37	54	92	189	189	122	46	4	-
Female	5	10	13	26	44	116	206	334	337	159	33	-
Black, both sexes	5	9	9	13	18	22	34	24	29	15	8	-
Male	5	5	4	9	8	8	12	8	11	4	1	-
Female	-	4	5	4	10	14	22	16	18	11	7	-
American Indian 1/, both sexes	-	-	-	-	1	1	1	1	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	3	-	-	2	1	1	3	1	-
Male	-	-	-	1	-	-	1	-	1	-	1	-
Female	-	-	-	2	-	-	1	1	-	3	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Minnesota												
All races, both sexes	4	9	16	19	29	75	137	249	252	146	31	-
Male	2	4	13	13	19	41	69	106	74	34	4	-
Female	2	5	3	6	10	34	68	143	178	112	27	-
White, both sexes	3	8	16	16	29	70	132	249	250	145	31	-
Male	2	3	13	11	19	38	65	106	72	34	4	-
Female	1	5	3	5	10	32	67	143	178	111	27	-
Black, both sexes	-	1	-	-	-	4	3	-	2	-	-	-
Male	-	1	-	-	-	3	2	-	2	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	3	-	-	2	-	-	-	-	-
Male	-	-	-	2	-	-	2	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	1	-	-
Mississippi												
All races, both sexes	26	33	38	67	87	142	213	249	184	112	25	1
Male	10	19	17	36	52	61	80	93	46	22	8	1
Female	16	14	21	31	35	81	133	156	138	90	17	-
White, both sexes	13	18	23	46	59	98	165	197	154	84	15	1
Male	5	12	12	26	35	40	69	78	42	15	3	1
Female	8	6	11	20	24	58	96	119	112	69	12	-
Black, both sexes	13	15	14	21	28	43	48	52	30	28	10	-
Male	5	7	5	10	17	20	11	15	4	7	5	-
Female	8	8	9	11	11	23	37	37	26	21	5	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Missouri												
All races, both sexes	7	20	37	50	83	141	219	316	288	137	41	-
Male	3	13	15	29	45	67	103	117	94	31	3	-
Female	4	7	22	21	38	74	116	199	194	106	38	-
White, both sexes	6	16	30	46	75	129	211	301	275	132	41	-
Male	3	11	12	25	41	62	100	113	86	29	3	-
Female	3	5	18	21	34	67	111	188	189	103	38	-
Black, both sexes	1	4	7	3	8	12	8	15	13	5	-	-
Male	-	2	3	3	4	5	3	4	8	2	-	-
Female	1	2	4	-	4	7	5	11	5	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	4	2	4	8	15	35	44	62	71	35	4	-
Male	4	1	2	5	8	18	27	22	26	12	1	-
Female	-	1	2	3	7	17	17	40	45	23	3	-
White, both sexes	4	1	4	7	13	33	42	62	71	35	4	-
Male	4	1	2	5	6	17	26	22	26	12	1	-
Female	-	-	2	2	7	16	16	40	45	23	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	1	2	2	1	-	-	-	-	-
Male	-	-	-	-	2	1	1	-	-	-	-	-
Female	-	1	-	1	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Nebraska												
All races, both sexes	4	9	6	14	20	33	62	86	98	48	14	-
Male	3	6	3	8	13	20	22	30	29	8	2	-
Female	1	3	3	6	7	13	40	56	69	40	12	-
White, both sexes	3	8	4	13	20	31	60	86	96	48	14	-
Male	3	6	2	8	13	18	22	30	29	8	2	-
Female	-	2	2	5	7	13	38	56	67	40	12	-
Black, both sexes	1	1	2	-	-	2	2	-	2	-	-	-
Male	-	-	1	-	-	2	-	-	-	-	-	-
Female	1	1	1	-	-	-	2	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	5	6	18	27	31	61	87	97	57	18	8	-
Male	5	4	12	19	20	37	54	41	18	6	1	-
Female	-	2	6	8	11	24	33	56	39	12	7	-
White, both sexes	5	4	14	23	30	56	84	91	55	18	7	-
Male	5	2	10	15	19	35	52	38	18	6	1	-
Female	-	2	4	8	11	21	32	53	37	12	6	-
Black, both sexes	-	2	2	2	1	4	2	-	1	-	-	-
Male	-	2	2	2	1	2	1	-	-	-	-	-
Female	-	-	-	-	-	2	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	1	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	1	-	-	1	5	-	-	1	-
Male	-	-	-	1	-	-	1	3	-	-	-	-
Female	-	-	2	-	-	-	-	2	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
New Hampshire												
All races, both sexes	-	3	4	5	9	27	44	44	45	22	12	-
Male	-	1	2	4	6	10	16	22	11	3	1	-
Female	-	2	2	1	3	17	28	22	34	19	11	-
White, both sexes	-	3	4	5	9	25	44	44	44	22	12	-
Male	-	1	2	4	6	10	16	22	11	3	1	-
Female	-	2	2	1	3	15	28	22	33	19	11	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-
New Jersey												
All races, both sexes	7	16	27	44	76	129	254	382	273	141	30	-
Male	4	9	16	21	45	60	102	154	93	40	8	-
Female	3	7	11	23	31	69	152	228	180	101	22	-
White, both sexes	4	7	19	35	57	105	236	352	266	130	29	-
Male	3	5	13	18	36	51	95	141	92	35	8	-
Female	1	2	6	17	21	54	141	211	174	95	21	-
Black, both sexes	3	9	8	9	18	22	17	27	5	8	1	-
Male	1	4	3	3	8	8	7	12	-	4	-	-
Female	2	5	5	6	10	14	10	15	5	4	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	2	1	3	2	3	-	-
Male	-	-	-	-	1	1	-	1	1	1	-	-
Female	-	-	-	-	-	1	1	2	1	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
New Mexico												
All races, both sexes	2	12	7	14	19	31	76	86	74	27	11	-
Male	2	6	3	10	13	12	36	31	28	5	4	-
Female	-	6	4	4	6	19	40	55	46	22	7	-
White, both sexes	2	10	7	14	17	29	74	85	72	25	9	-
Male	2	6	3	10	13	10	34	30	26	5	3	-
Female	-	4	4	4	4	19	40	55	46	20	6	-
Black, both sexes	-	-	-	-	-	-	1	-	-	2	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	2	-	-
American Indian 1/, both sexes	-	1	-	-	2	2	-	1	2	-	2	-
Male	-	-	-	-	-	2	-	1	2	-	1	-
Female	-	1	-	-	2	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	21	39	43	65	127	232	453	574	524	248	70	-
Male	13	27	28	42	73	110	197	219	182	63	8	-
Female	8	12	15	23	54	122	256	355	342	185	62	-
White, both sexes	18	22	33	57	110	198	425	542	499	240	65	-
Male	11	16	21	36	61	94	181	215	180	61	8	-
Female	7	6	12	21	49	104	244	327	319	179	57	-
Black, both sexes	3	16	10	7	13	31	24	30	24	6	5	-
Male	2	10	7	5	9	15	14	3	2	1	-	-
Female	1	6	3	2	4	16	10	27	22	5	5	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	4	3	4	2	1	2	-	-
Male	-	-	-	1	3	1	2	1	-	1	-	-
Female	-	-	-	-	1	2	2	1	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
North Carolina												
All races, both sexes	16	30	51	61	97	160	253	314	270	112	23	-
Male	11	14	30	41	56	80	107	108	79	17	-	-
Female	5	16	21	20	41	80	146	206	191	95	23	-
White, both sexes	11	19	40	50	75	128	221	267	241	93	20	-
Male	8	11	23	36	44	61	96	101	71	14	-	-
Female	3	8	17	14	31	67	125	166	170	79	20	-
Black, both sexes	5	9	11	11	22	31	32	43	29	19	2	-
Male	3	2	7	5	12	18	11	7	8	3	-	-
Female	2	7	4	6	10	13	21	36	21	16	2	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	-	2	-	-	1	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	1	-
North Dakota												
All races, both sexes	-	2	1	1	4	2	4	21	17	18	4	-
Male	-	2	1	1	3	2	1	9	5	2	-	-
Female	-	-	-	-	1	-	3	12	12	16	4	-
White, both sexes	-	-	-	1	4	2	4	20	17	18	4	-
Male	-	-	-	1	3	2	1	9	5	2	-	-
Female	-	-	-	-	1	-	3	11	12	16	4	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	2	1	-	-	-	-	1	-	-	-	-
Male	-	2	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Ohio												
All races, both sexes	18	20	43	48	90	156	273	329	293	105	22	-
Male	8	15	27	25	40	84	118	130	103	23	4	-
Female	10	5	16	23	50	72	155	199	190	82	18	-
White, both sexes	14	18	32	40	81	143	257	306	273	98	21	-
Male	7	14	19	21	36	78	110	122	97	21	3	-
Female	7	4	13	19	45	65	147	184	176	77	18	-
Black, both sexes	4	2	11	8	9	13	16	22	20	7	1	-
Male	1	1	8	4	4	6	8	8	6	2	1	-
Female	3	1	3	4	5	7	8	14	14	5	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Oklahoma												
All races, both sexes	25	35	52	78	104	155	240	266	210	90	22	-
Male	13	20	31	42	50	78	109	111	68	15	2	-
Female	12	15	21	36	54	77	131	155	142	75	20	-
White, both sexes	19	25	43	68	96	146	221	250	197	85	21	-
Male	11	14	25	37	46	72	101	108	63	14	2	-
Female	8	11	18	31	50	74	120	142	134	71	19	-
Black, both sexes	6	6	3	5	6	7	10	11	9	3	1	-
Male	2	4	2	3	4	5	5	1	3	-	-	-
Female	4	2	1	2	2	2	5	10	6	3	1	-
American Indian 1/, both sexes	-	4	4	5	2	2	8	5	4	1	-	-
Male	-	2	3	2	-	1	2	2	2	-	-	-
Female	-	2	1	3	2	1	6	3	2	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	-	-	1	-	-	1	-	-
Male	-	-	1	-	-	-	1	-	-	1	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Oregon												
All races, both sexes	4	8	13	25	28	56	115	156	145	72	18	-
Male	3	3	6	17	13	30	49	51	50	26	5	-
Female	1	5	7	8	15	26	66	105	95	46	13	-
White, both sexes	4	8	13	25	26	56	112	154	144	71	18	-
Male	3	3	6	17	12	30	48	51	50	26	5	-
Female	1	5	7	8	14	26	64	103	94	45	13	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	2	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	3	1	1	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	2	1	1	1	-	-
Pennsylvania												
All races, both sexes	24	46	71	101	130	297	540	780	651	262	66	-
Male	15	27	43	52	69	153	233	298	184	58	6	-
Female	9	19	28	49	61	144	307	482	467	204	60	-
White, both sexes	13	34	62	88	116	280	504	747	624	251	64	-
Male	9	19	37	45	60	147	220	287	178	56	5	-
Female	4	15	25	43	56	133	284	460	446	195	59	-
Black, both sexes	10	11	9	13	12	17	33	29	26	11	2	-
Male	5	7	6	7	8	6	13	9	5	2	1	-
Female	5	4	3	6	4	11	20	20	21	9	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	2	-	3	3	1	-	-	-
Male	1	1	-	-	1	-	-	2	1	-	-	-
Female	-	-	-	-	1	-	3	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Rhode Island												
All races, both sexes	-	1	2	3	8	11	24	38	25	18	3	-
Male	-	1	1	3	5	5	8	16	10	7	-	-
Female	-	-	1	-	3	6	16	22	15	11	3	-
White, both sexes	-	1	2	3	8	11	22	38	25	18	3	-
Male	-	1	1	3	5	5	8	16	10	7	-	-
Female	-	-	1	-	3	6	14	22	15	11	3	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
South Carolina												
All races, both sexes	18	32	32	53	67	92	145	148	113	55	13	-
Male	12	19	15	31	34	39	64	51	40	9	2	-
Female	6	13	17	22	33	53	81	97	73	46	11	-
White, both sexes	5	16	23	35	52	68	117	125	93	42	9	-
Male	3	13	9	23	28	30	52	47	34	7	2	-
Female	2	3	14	12	24	38	65	78	59	35	7	-
Black, both sexes	13	16	9	18	15	23	28	22	19	13	4	-
Male	9	6	6	8	6	8	12	4	6	2	-	-
Female	4	10	3	10	9	15	16	18	13	11	4	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
South Dakota												
All races, both sexes	2	-	-	3	5	5	5	11	14	8	2	-
Male	1	-	-	1	3	4	1	3	4	5	-	-
Female	1	-	-	2	2	1	4	8	10	3	2	-
White, both sexes	1	-	-	3	4	4	5	11	13	8	2	-
Male	1	-	-	1	3	3	1	3	4	5	-	-
Female	-	-	-	2	1	1	4	8	9	3	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	1	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	12	18	21	29	41	83	106	152	98	57	8	-
Male	7	15	14	12	24	35	46	38	22	4	-	-
Female	5	3	7	17	17	48	60	114	76	53	8	-
White, both sexes	7	13	17	22	36	71	91	138	93	49	8	-
Male	4	12	11	8	22	30	40	34	20	3	-	-
Female	3	1	6	14	14	41	51	104	73	46	8	-
Black, both sexes	5	5	4	7	5	11	15	14	5	6	-	-
Male	3	3	3	4	2	5	6	4	2	-	-	-
Female	2	2	1	3	3	6	9	10	3	6	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	2	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	1	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Texas												
All races, both sexes	67	93	140	190	271	472	684	866	658	326	67	-
Male	44	61	86	105	158	228	317	332	205	73	14	-
Female	23	32	54	85	113	244	367	534	453	253	53	-
White, both sexes	48	61	114	157	231	425	630	793	618	297	57	-
Male	33	40	71	89	132	207	295	304	193	65	10	-
Female	15	21	43	68	99	218	335	489	425	232	47	-
Black, both sexes	19	32	25	31	40	42	49	68	35	27	10	-
Male	11	21	15	15	26	18	19	25	12	7	4	-
Female	8	11	10	16	14	24	30	43	23	20	6	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	2	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	2	-	4	5	3	5	2	-	-
Male	-	-	-	1	-	2	3	2	-	1	-	-
Female	-	-	1	1	-	2	2	1	5	1	-	-
Utah												
All races, both sexes	2	8	11	20	36	59	121	122	134	64	3	-
Male	1	7	6	9	17	37	51	43	48	26	2	-
Female	1	1	5	11	19	22	70	79	86	38	1	-
White, both sexes	2	8	11	18	35	58	119	121	131	64	3	-
Male	1	7	6	9	16	36	49	42	48	26	2	-
Female	1	1	5	9	19	22	70	79	83	38	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	1	1	1	1	3	-	-	-
Male	-	-	-	-	1	1	1	1	-	-	-	-
Female	-	-	-	2	-	-	-	-	3	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Vermont												
All races, both sexes	-	-	2	1	1	2	6	14	10	5	2	-
Male	-	-	-	1	1	1	4	5	1	2	-	-
Female	-	-	2	-	-	1	2	9	9	3	2	-
White, both sexes	-	-	2	1	1	2	6	14	10	5	2	-
Male	-	-	-	1	1	1	4	5	1	2	-	-
Female	-	-	2	-	-	1	2	9	9	3	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	9	19	33	40	68	157	234	313	263	130	20	-
Male	7	13	16	25	29	82	116	125	74	30	2	-
Female	2	6	17	15	39	75	118	188	189	100	18	-
White, both sexes	5	10	23	32	49	121	188	280	232	115	19	-
Male	4	7	12	22	23	62	93	113	64	26	2	-
Female	1	3	11	10	26	59	95	167	168	89	17	-
Black, both sexes	4	9	10	8	16	35	42	32	27	12	1	-
Male	3	6	4	3	5	19	21	11	7	3	-	-
Female	1	3	6	5	11	16	21	21	20	9	1	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	3	-	4	1	4	2	-	-
Male	-	-	-	-	1	-	2	1	3	-	-	-
Female	-	-	-	-	2	-	2	-	1	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Washington												
All races, both sexes	3	4	12	16	17	32	74	88	86	39	10	-
Male	2	2	5	12	8	12	29	36	32	12	2	-
Female	1	2	7	4	9	20	45	52	54	27	8	-
White, both sexes	3	4	11	16	15	28	72	85	83	39	9	-
Male	2	2	4	12	7	11	29	35	30	12	2	-
Female	1	2	7	4	8	17	43	50	53	27	7	-
Black, both sexes	-	-	1	-	2	-	-	1	-	-	1	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	3	-	1	1	-	-	-
Male	-	-	-	-	-	1	-	1	1	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	2	1	2	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	1	2	1	1	-	-	-
West Virginia												
All races, both sexes	7	7	16	30	28	51	82	87	90	34	9	-
Male	4	4	10	16	14	21	44	36	25	8	1	-
Female	3	3	6	14	14	30	38	51	65	26	8	-
White, both sexes	7	7	16	29	28	49	78	83	89	33	9	-
Male	4	4	10	15	14	20	41	36	24	8	1	-
Female	3	3	6	14	14	29	37	47	65	25	8	-
Black, both sexes	-	-	-	1	-	2	4	4	1	1	-	-
Male	-	-	-	1	-	1	3	-	1	-	-	-
Female	-	-	-	-	-	1	1	4	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Wisconsin												
All races, both sexes	12	17	19	32	59	126	220	328	353	160	44	-
Male	8	12	12	20	33	67	96	144	107	41	5	-
Female	4	5	7	12	26	59	124	184	246	119	39	-
White, both sexes	9	15	17	32	54	123	215	321	349	157	43	-
Male	6	11	11	20	31	66	96	142	107	41	5	-
Female	3	4	6	12	23	57	119	179	242	116	38	-
Black, both sexes	3	2	2	-	5	1	4	5	4	2	1	-
Male	2	1	1	-	2	-	-	2	-	-	-	-
Female	1	1	1	-	3	1	4	3	4	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	2	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	2	-	1	-	-
Wyoming												
All races, both sexes	-	-	1	4	6	8	20	25	29	13	-	-
Male	-	-	-	3	2	3	9	7	7	5	-	-
Female	-	-	1	1	4	5	11	18	22	8	-	-
White, both sexes	-	-	1	4	6	8	20	24	28	13	-	-
Male	-	-	-	3	2	3	9	7	7	5	-	-
Female	-	-	1	1	4	5	11	17	21	8	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
United States												
All races, both sexes	4,598	13	5	3	4	2	14	7	18	23	40	72
Male	1,985	5	2	1	2	1	5	3	10	15	29	45
Female	2,613	8	3	2	2	1	9	4	8	8	11	27
White, both sexes	3,857	5	4	3	4	1	9	5	7	13	25	50
Male	1,655	2	1	1	2	-	3	3	4	9	18	33
Female	2,202	3	3	2	2	1	6	2	3	4	7	17
Black, both sexes	674	6	1	-	-	1	4	2	11	10	15	21
Male	290	2	1	-	-	1	2	-	6	6	11	11
Female	384	4	-	-	-	-	2	2	5	4	4	10
American Indian 1/, both sexes	14	1	-	-	-	-	1	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	7	1	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	53	1	-	-	-	-	-	-	-	-	-	-
Male	33	1	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	240	-	-	-	-	-	2	-	4	5	4	5
Male	118	-	-	-	-	-	2	-	3	3	2	4
Female	122	-	-	-	-	-	-	-	1	2	2	1
White, both sexes	139	-	-	-	-	-	2	-	1	1	2	4
Male	73	-	-	-	-	-	2	-	1	1	1	3
Female	66	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	101	-	-	-	-	-	-	-	3	4	2	1
Male	45	-	-	-	-	-	-	-	2	2	1	1
Female	56	-	-	-	-	-	-	-	1	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Alaska												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	45	-	-	-	-	1	-	-	-	-	-	2
Male	16	-	-	-	-	-	-	-	-	-	-	1
Female	29	-	-	-	-	1	-	-	-	-	-	1
White, both sexes	43	-	-	-	-	1	-	-	-	-	-	2
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	28	-	-	-	-	1	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Arkansas												
All races, both sexes	38	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	32	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	141	1	-	-	1	-	1	1	-	1	1	1
Male	65	1	-	-	1	-	1	-	-	1	1	1
Female	76	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	121	-	-	-	1	-	1	1	-	1	1	1
Male	49	-	-	-	1	-	1	-	-	1	1	1
Female	72	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	1	-	-	-	-	-	-	-	-	-	-
Male	9	1	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Colorado												
All races, both sexes	43	-	-	-	-	-	-	-	-	1	1	-
Male	21	-	-	-	-	-	-	-	-	1	1	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	39	-	-	-	-	-	-	-	-	1	-	-
Male	18	-	-	-	-	-	-	-	-	1	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	105	-	-	-	-	-	-	-	-	-	1	-
Male	40	-	-	-	-	-	-	-	-	-	1	-
Female	65	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	101	-	-	-	-	-	-	-	-	-	1	-
Male	38	-	-	-	-	-	-	-	-	-	1	-
Female	63	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Delaware												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Florida												
All races, both sexes	137	2	-	-	-	1	2	-	1	1	-	2
Male	62	-	-	-	-	1	-	-	-	1	-	1
Female	75	2	-	-	-	-	2	-	1	-	-	1
White, both sexes	120	-	-	-	-	-	2	-	-	-	-	1
Male	53	-	-	-	-	-	-	-	-	-	-	-
Female	67	-	-	-	-	-	2	-	-	-	-	1
Black, both sexes	17	2	-	-	-	1	-	-	1	1	-	1
Male	9	-	-	-	-	1	-	-	-	1	-	1
Female	8	2	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	221	-	-	2	-	-	-	-	2	2	4	4
Male	91	-	-	1	-	-	-	-	-	2	2	4
Female	130	-	-	1	-	-	-	-	2	-	2	-
White, both sexes	158	-	-	2	-	-	-	-	-	2	3	-
Male	63	-	-	1	-	-	-	-	-	2	1	-
Female	95	-	-	1	-	-	-	-	-	-	2	-
Black, both sexes	62	-	-	-	-	-	-	-	2	-	1	4
Male	28	-	-	-	-	-	-	-	-	-	1	4
Female	34	-	-	-	-	-	-	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Hawaii												
All races, both sexes	29	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	24	-	-	-	-	-	1	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	23	-	-	-	-	-	1	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Illinois												
All races, both sexes	163	1	-	-	-	-	-	-	1	1	1	4
Male	64	1	-	-	-	-	-	-	-	-	1	3
Female	99	-	-	-	-	-	-	-	1	1	-	1
White, both sexes	139	1	-	-	-	-	-	-	1	-	-	3
Male	56	1	-	-	-	-	-	-	-	-	-	3
Female	83	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	24	-	-	-	-	-	-	-	-	1	1	1
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	16	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	101	-	-	-	1	-	-	-	-	-	1	1
Male	40	-	-	-	-	-	-	-	-	-	1	1
Female	61	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	90	-	-	-	1	-	-	-	-	-	-	1
Male	36	-	-	-	-	-	-	-	-	-	-	1
Female	54	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	10	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Iowa												
All races, both sexes	33	-	-	-	-	-	-	-	1	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	33	-	-	-	-	-	-	-	1	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	48	-	-	-	-	-	-	-	1	-	1	1
Male	20	-	-	-	-	-	-	-	1	-	-	1
Female	28	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	44	-	-	-	-	-	-	-	-	-	1	1
Male	18	-	-	-	-	-	-	-	-	-	-	1
Female	26	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	4	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Kentucky												
All races, both sexes	88	-	-	-	1	-	-	-	-	-	2	4
Male	31	-	-	-	-	-	-	-	-	-	2	1
Female	57	-	-	-	1	-	-	-	-	-	-	3
White, both sexes	82	-	-	-	1	-	-	-	-	-	2	3
Male	30	-	-	-	-	-	-	-	-	-	2	1
Female	52	-	-	-	1	-	-	-	-	-	-	2
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	46	-	-	-	-	-	-	-	-	-	-	2
Male	25	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	32	-	-	-	-	-	-	-	-	-	-	2
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Maine												
All races, both sexes	28	-	-	-	-	-	1	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	28	-	-	-	-	-	1	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Massachusetts												
All races, both sexes	142	-	-	-	-	-	-	-	-	-	1	-
Male	57	-	-	-	-	-	-	-	-	-	1	-
Female	85	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	129	-	-	-	-	-	-	-	-	-	-	-
Male	49	-	-	-	-	-	-	-	-	-	-	-
Female	80	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	11	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	147	-	-	-	-	-	-	-	1	2	2	3
Male	68	-	-	-	-	-	-	-	-	1	2	2
Female	79	-	-	-	-	-	-	-	1	1	-	1
White, both sexes	117	-	-	-	-	-	-	-	1	1	-	3
Male	54	-	-	-	-	-	-	-	-	-	-	2
Female	63	-	-	-	-	-	-	-	1	1	-	1
Black, both sexes	26	-	-	-	-	-	-	-	-	1	2	-
Male	12	-	-	-	-	-	-	-	-	1	2	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Minnesota												
All races, both sexes	92	-	1	-	-	-	-	1	-	-	-	-
Male	42	-	-	-	-	-	-	-	-	-	-	-
Female	50	-	1	-	-	-	-	1	-	-	-	-
White, both sexes	91	-	1	-	-	-	-	-	-	-	-	-
Male	42	-	-	-	-	-	-	-	-	-	-	-
Female	49	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	144	-	-	-	-	-	-	-	1	-	3	5
Male	54	-	-	-	-	-	-	-	1	-	2	1
Female	90	-	-	-	-	-	-	-	-	-	1	4
White, both sexes	96	-	-	-	-	-	-	-	-	-	2	1
Male	38	-	-	-	-	-	-	-	-	-	2	-
Female	58	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	48	-	-	-	-	-	-	-	1	-	1	4
Male	16	-	-	-	-	-	-	-	1	-	-	1
Female	32	-	-	-	-	-	-	-	-	-	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Missouri												
All races, both sexes	97	-	-	-	-	-	-	-	-	1	1	2
Male	43	-	-	-	-	-	-	-	-	1	-	1
Female	54	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	89	-	-	-	-	-	-	-	-	1	1	2
Male	41	-	-	-	-	-	-	-	-	1	-	1
Female	48	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	42	-	-	-	-	-	1	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	41	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Nebraska												
All races, both sexes	70	-	-	-	-	-	-	-	-	-	1	1
Male	36	-	-	-	-	-	-	-	-	-	1	1
Female	34	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	68	-	-	-	-	-	-	-	-	-	-	1
Male	34	-	-	-	-	-	-	-	-	-	-	1
Female	34	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	29	-	-	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
New Hampshire												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	136	-	-	-	-	-	-	-	1	-	1	2
Male	64	-	-	-	-	-	-	-	1	-	-	1
Female	72	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	115	-	-	-	-	-	-	-	1	-	1	-
Male	53	-	-	-	-	-	-	-	1	-	-	-
Female	62	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	20	-	-	-	-	-	-	-	-	-	-	2
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
New Mexico												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	259	-	-	-	-	-	-	1	1	-	-	4
Male	109	-	-	-	-	-	-	1	1	-	-	4
Female	150	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	237	-	-	-	-	-	-	1	-	-	-	3
Male	102	-	-	-	-	-	-	1	-	-	-	3
Female	135	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	21	-	-	-	-	-	-	-	1	-	-	1
Male	6	-	-	-	-	-	-	-	1	-	-	1
Female	15	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
North Carolina												
All races, both sexes	137	6	1	1	-	-	-	1	-	2	-	3
Male	57	2	-	-	-	-	-	-	-	1	-	1
Female	80	4	1	1	-	-	-	1	-	1	-	2
White, both sexes	106	3	1	1	-	-	-	-	-	-	-	2
Male	44	-	-	-	-	-	-	-	-	-	-	1
Female	62	3	1	1	-	-	-	-	-	-	-	1
Black, both sexes	30	2	-	-	-	-	-	1	-	2	-	1
Male	13	2	-	-	-	-	-	-	-	1	-	-
Female	17	-	-	-	-	-	-	1	-	1	-	1
American Indian 1/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Ohio												
All races, both sexes	102	-	1	-	-	-	-	-	-	-	1	1
Male	49	-	1	-	-	-	-	-	-	-	1	1
Female	53	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	93	-	-	-	-	-	-	-	-	-	1	1
Male	45	-	-	-	-	-	-	-	-	-	1	1
Female	48	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	1	-	-	-	-	-	-	-	-	-
Male	4	-	1	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	87	-	-	-	1	-	-	-	-	-	-	1
Male	28	-	-	-	1	-	-	-	-	-	-	1
Female	59	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	81	-	-	-	1	-	-	-	-	-	-	1
Male	28	-	-	-	1	-	-	-	-	-	-	1
Female	53	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Oregon												
All races, both sexes	60	-	-	-	-	-	1	-	-	1	-	1
Male	30	-	-	-	-	-	-	-	-	-	-	1
Female	30	-	-	-	-	-	1	-	-	1	-	-
White, both sexes	58	-	-	-	-	-	1	-	-	1	-	1
Male	28	-	-	-	-	-	-	-	-	-	-	1
Female	30	-	-	-	-	-	1	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	288	-	-	-	-	-	2	-	1	1	4	1
Male	120	-	-	-	-	-	-	-	-	1	4	-
Female	168	-	-	-	-	-	2	-	1	-	-	1
White, both sexes	265	-	-	-	-	-	1	-	-	-	4	-
Male	113	-	-	-	-	-	-	-	-	-	4	-
Female	152	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	20	-	-	-	-	-	1	-	1	1	-	1
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	15	-	-	-	-	-	1	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Rhode Island												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	125	2	-	-	-	-	1	-	-	-	-	2
Male	56	1	-	-	-	-	-	-	-	-	-	1
Female	69	1	-	-	-	-	1	-	-	-	-	1
White, both sexes	81	1	-	-	-	-	-	-	-	-	-	2
Male	37	1	-	-	-	-	-	-	-	-	-	1
Female	44	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	44	1	-	-	-	-	1	-	-	-	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	25	1	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
South Dakota												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	82	-	-	-	-	-	1	1	-	-	1	1
Male	33	-	-	-	-	-	1	1	-	-	1	1
Female	49	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	63	-	-	-	-	-	-	1	-	-	1	-
Male	28	-	-	-	-	-	-	1	-	-	1	-
Female	35	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	19	-	-	-	-	-	1	-	-	-	-	1
Male	5	-	-	-	-	-	1	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Texas												
All races, both sexes	376	1	-	-	-	-	-	2	1	3	6	9
Male	183	-	-	-	-	-	-	1	1	2	5	6
Female	193	1	-	-	-	-	-	1	-	1	1	3
White, both sexes	324	-	-	-	-	-	-	2	1	3	5	7
Male	160	-	-	-	-	-	-	1	1	2	4	4
Female	164	-	-	-	-	-	-	1	-	1	1	3
Black, both sexes	47	1	-	-	-	-	-	-	-	-	1	1
Male	19	-	-	-	-	-	-	-	-	-	1	1
Female	28	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	28	-	1	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	27	-	1	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Vermont												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	356	-	1	-	-	-	1	-	2	2	3	8
Male	153	-	1	-	-	-	1	-	2	1	1	5
Female	203	-	-	-	-	-	-	-	-	1	2	3
White, both sexes	259	-	1	-	-	-	-	-	1	2	-	7
Male	107	-	1	-	-	-	-	-	1	1	-	5
Female	152	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	95	-	-	-	-	-	1	-	1	-	3	1
Male	45	-	-	-	-	-	1	-	1	-	1	-
Female	50	-	-	-	-	-	-	-	-	-	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Washington												
All races, both sexes	31	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	31	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	105	-	-	-	-	-	-	-	-	-	-	-
Male	34	-	-	-	-	-	-	-	-	-	-	-
Female	71	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	104	-	-	-	-	-	-	-	-	-	-	-
Male	34	-	-	-	-	-	-	-	-	-	-	-
Female	70	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Wisconsin												
All races, both sexes	39	-	-	-	-	-	-	-	-	-	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	38	-	-	-	-	-	-	-	-	-	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
United States												
All races, both sexes	99	177	214	256	347	521	742	852	717	370	102	-
Male	56	114	133	147	202	269	321	335	197	72	21	-
Female	43	63	81	109	145	252	421	517	520	298	81	-
White, both sexes	62	121	144	182	288	451	650	757	649	335	92	-
Male	37	74	92	112	175	239	283	304	178	65	20	-
Female	25	47	52	70	113	212	367	453	471	270	72	-
Black, both sexes	35	54	63	66	58	64	82	85	57	30	9	-
Male	17	38	36	30	27	26	31	27	13	4	1	-
Female	18	16	27	36	31	38	51	58	44	26	8	-
American Indian 1/, both sexes	1	-	3	1	-	1	1	2	1	-	1	-
Male	1	-	1	1	-	1	-	1	1	-	-	-
Female	-	-	2	-	-	-	1	1	-	-	1	-
Asian or Pacific Islander 2/, both sexes	1	2	4	7	1	5	9	8	10	5	-	-
Male	1	2	4	4	-	3	7	3	5	3	-	-
Female	-	-	-	3	1	2	2	5	5	2	-	-
Alabama												
All races, both sexes	9	15	14	22	25	31	32	32	26	10	4	-
Male	5	9	8	15	13	16	14	15	5	4	-	-
Female	4	6	6	7	12	15	18	17	21	6	4	-
White, both sexes	3	5	10	12	14	19	21	19	18	6	2	-
Male	2	3	6	9	9	11	9	10	3	3	-	-
Female	1	2	4	3	5	8	12	9	15	3	2	-
Black, both sexes	6	10	4	10	11	12	11	13	8	4	2	-
Male	3	6	2	6	4	5	5	5	2	1	-	-
Female	3	4	2	4	7	7	6	8	6	3	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Alaska												
All races, both sexes	-	-	-	1	-	2	-	1	1	-	-	-
Male	-	-	-	1	-	2	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	1	1	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	1	1	2	3	4	8	9	10	2	2	-
Male	-	1	-	-	2	1	4	4	1	2	-	-
Female	-	-	1	2	1	3	4	5	9	-	2	-
White, both sexes	-	1	1	2	3	4	7	8	10	2	2	-
Male	-	1	-	-	2	1	4	3	1	2	-	-
Female	-	-	1	2	1	3	3	5	9	-	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Arkansas												
All races, both sexes	-	3	-	2	2	7	6	2	12	4	-	-
Male	-	2	-	-	2	4	2	1	3	1	-	-
Female	-	1	-	2	-	3	4	1	9	3	-	-
White, both sexes	-	1	-	1	2	7	5	1	11	4	-	-
Male	-	-	-	-	2	4	1	1	2	1	-	-
Female	-	1	-	1	-	3	4	-	9	3	-	-
Black, both sexes	-	2	-	1	-	-	1	1	1	-	-	-
Male	-	2	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	3	4	5	13	16	28	19	26	16	4	-
Male	-	3	3	4	6	11	13	9	9	1	-	-
Female	-	-	1	1	7	5	15	10	17	15	4	-
White, both sexes	-	3	2	3	12	14	25	18	21	13	4	-
Male	-	3	1	2	5	10	10	8	5	-	-	-
Female	-	-	1	1	7	4	15	10	16	13	4	-
Black, both sexes	-	-	1	1	1	1	3	1	-	-	-	-
Male	-	-	1	1	1	-	3	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	1	-	-	5	3	-	-
Male	-	-	1	1	-	1	-	-	4	1	-	-
Female	-	-	-	-	-	-	-	-	1	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Colorado												
All races, both sexes	1	1	1	-	4	4	9	12	4	4	1	-
Male	-	1	-	-	4	3	2	4	1	3	1	-
Female	1	-	1	-	-	1	7	8	3	1	-	-
White, both sexes	1	-	-	-	4	4	9	11	4	4	1	-
Male	-	-	-	-	4	3	2	3	1	3	1	-
Female	1	-	-	-	-	1	7	8	3	1	-	-
Black, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	2	3	5	5	7	16	15	20	14	13	4	-
Male	1	2	3	3	4	5	7	5	2	5	2	-
Female	1	1	2	2	3	11	8	15	12	8	2	-
White, both sexes	1	2	5	4	7	16	15	20	13	13	4	-
Male	1	1	3	2	4	5	7	5	2	5	2	-
Female	-	1	2	2	3	11	8	15	11	8	2	-
Black, both sexes	1	1	-	1	-	-	-	-	1	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Delaware												
All races, both sexes	1	1	-	-	3	2	-	2	3	-	-	-
Male	1	1	-	-	3	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	3	-	-	-
White, both sexes	1	-	-	-	2	2	-	2	3	-	-	-
Male	1	-	-	-	2	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	3	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Florida												
All races, both sexes	5	4	2	7	9	14	22	23	25	15	2	-
Male	3	2	2	3	6	9	10	12	10	2	-	-
Female	2	2	-	4	3	5	12	11	15	13	2	-
White, both sexes	3	4	1	7	7	13	21	21	24	14	2	-
Male	1	2	1	3	5	9	10	11	9	2	-	-
Female	2	2	-	4	2	4	11	10	15	12	2	-
Black, both sexes	2	-	1	-	2	1	1	2	1	1	-	-
Male	2	-	1	-	1	-	-	1	1	-	-	-
Female	-	-	-	-	1	1	1	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	4	13	9	18	20	16	33	39	32	18	5	-
Male	2	9	5	10	13	4	14	13	9	1	2	-
Female	2	4	4	8	7	12	19	26	23	17	3	-
White, both sexes	3	6	4	8	13	13	28	32	24	16	4	-
Male	2	4	2	7	8	3	12	11	7	1	2	-
Female	1	2	2	1	5	10	16	21	17	15	2	-
Black, both sexes	1	7	5	9	7	3	5	7	8	2	1	-
Male	-	5	3	3	5	1	2	2	2	-	-	-
Female	1	2	2	6	2	2	3	5	6	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Hawaii												
All races, both sexes	-	1	-	3	1	6	4	11	2	1	-	-
Male	-	1	-	1	-	2	4	6	-	-	-	-
Female	-	-	-	2	1	4	-	5	2	1	-	-
White, both sexes	-	-	-	-	1	3	-	5	1	1	-	-
Male	-	-	-	-	-	-	-	4	-	-	-	-
Female	-	-	-	-	1	3	-	1	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	3	-	3	4	6	1	-	-	-
Male	-	1	-	1	-	2	4	2	-	-	-	-
Female	-	-	-	2	-	1	-	4	1	-	-	-
Idaho												
All races, both sexes	-	3	2	-	-	1	3	6	5	3	-	-
Male	-	2	2	-	-	-	1	2	3	-	-	-
Female	-	1	-	-	-	1	2	4	2	3	-	-
White, both sexes	-	3	2	-	-	1	3	6	4	3	-	-
Male	-	2	2	-	-	-	1	2	2	-	-	-
Female	-	1	-	-	-	1	2	4	2	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Illinois												
All races, both sexes	1	11	9	7	15	11	34	28	26	10	3	-
Male	-	7	4	5	7	4	11	10	6	3	2	-
Female	1	4	5	2	8	7	23	18	20	7	1	-
White, both sexes	-	10	5	4	12	9	32	25	24	10	3	-
Male	-	6	1	4	7	3	11	9	6	3	2	-
Female	-	4	4	-	5	6	21	16	18	7	1	-
Black, both sexes	1	1	4	3	3	2	2	3	2	-	-	-
Male	-	1	3	1	-	1	-	1	-	-	-	-
Female	1	-	1	2	3	1	2	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	4	4	3	6	8	14	19	13	14	11	2	-
Male	1	1	1	3	5	6	9	5	4	2	1	-
Female	3	3	2	3	3	8	10	8	10	9	1	-
White, both sexes	3	3	2	4	8	14	18	12	12	10	2	-
Male	1	1	1	2	5	6	8	5	4	1	1	-
Female	2	2	1	2	3	8	10	7	8	9	1	-
Black, both sexes	1	1	1	2	-	-	1	1	1	1	-	-
Male	-	-	-	1	-	-	1	-	-	1	-	-
Female	1	1	1	1	-	-	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Iowa												
All races, both sexes	-	-	-	-	1	3	6	8	7	4	3	-
Male	-	-	-	-	-	3	2	3	1	-	-	-
Female	-	-	-	-	1	-	4	5	6	4	3	-
White, both sexes	-	-	-	-	1	3	6	8	7	4	3	-
Male	-	-	-	-	-	3	2	3	1	-	-	-
Female	-	-	-	-	1	-	4	5	6	4	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	2	2	1	4	3	8	10	10	5	-	-
Male	-	1	2	1	4	1	3	3	3	-	-	-
Female	-	1	-	-	-	2	5	7	7	5	-	-
White, both sexes	-	2	2	1	3	2	8	10	9	5	-	-
Male	-	1	2	1	3	1	3	3	3	-	-	-
Female	-	1	-	-	-	1	5	7	6	5	-	-
Black, both sexes	-	-	-	-	1	1	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Kentucky												
All races, both sexes	1	3	7	5	6	6	21	19	5	7	1	-
Male	1	2	5	3	4	4	4	5	-	-	-	-
Female	-	1	2	2	2	2	17	14	5	7	1	-
White, both sexes	1	3	6	4	6	6	20	18	5	6	1	-
Male	1	2	4	3	4	4	4	5	-	-	-	-
Female	-	1	2	1	2	2	16	13	5	6	1	-
Black, both sexes	-	-	1	1	-	-	1	1	-	1	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	5	6	7	3	5	7	7	3	1	-	-
Male	-	3	4	4	2	5	2	4	1	-	-	-
Female	-	2	2	3	1	-	5	3	2	1	-	-
White, both sexes	-	2	3	4	1	4	5	7	3	1	-	-
Male	-	-	3	3	1	4	2	4	1	-	-	-
Female	-	2	-	1	-	-	3	3	2	1	-	-
Black, both sexes	-	3	3	3	2	1	2	-	-	-	-	-
Male	-	3	1	1	1	1	-	-	-	-	-	-
Female	-	-	2	2	1	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Maine												
All races, both sexes	-	-	-	2	4	4	5	7	2	2	1	-
Male	-	-	-	1	3	1	2	3	-	-	-	-
Female	-	-	-	1	1	3	3	4	2	2	1	-
White, both sexes	-	-	-	2	4	4	5	7	2	2	1	-
Male	-	-	-	1	3	1	2	3	-	-	-	-
Female	-	-	-	1	1	3	3	4	2	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	1	-	-	3	3	3	4	-	2	-	-
Male	-	1	-	-	1	2	1	3	-	-	-	-
Female	1	-	-	-	2	1	2	1	-	2	-	-
White, both sexes	1	-	-	-	3	2	3	3	-	1	-	-
Male	-	-	-	-	1	2	1	2	-	-	-	-
Female	1	-	-	-	2	-	2	1	-	1	-	-
Black, both sexes	-	1	-	-	-	1	-	1	-	1	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Massachusetts												
All races, both sexes	4	5	7	8	9	14	20	36	21	13	4	-
Male	1	3	1	6	6	7	8	14	7	3	-	-
Female	3	2	6	2	3	7	12	22	14	10	4	-
White, both sexes	3	4	6	6	8	14	19	33	21	11	4	-
Male	1	2	-	4	6	7	8	13	7	1	-	-
Female	2	2	6	2	2	7	11	20	14	10	4	-
Black, both sexes	1	1	1	1	1	-	1	3	-	1	-	-
Male	-	1	1	1	-	-	-	1	-	1	-	-
Female	1	-	-	-	1	-	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	1	-	-
Male	-	-	-	1	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	4	4	6	6	12	22	29	22	19	9	6	-
Male	3	3	4	2	8	12	10	9	10	1	1	-
Female	1	1	2	4	4	10	19	13	9	8	5	-
White, both sexes	3	3	3	2	8	19	21	21	18	9	5	-
Male	2	2	2	1	6	10	9	8	10	1	1	-
Female	1	1	1	1	2	9	12	13	8	8	4	-
Black, both sexes	1	1	1	4	4	3	7	1	1	-	-	-
Male	1	1	-	1	2	2	1	1	-	-	-	-
Female	-	-	1	3	2	1	6	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	-	-	1	-	-	-	-	-
Male	-	-	2	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Minnesota												
All races, both sexes	-	1	2	3	7	11	12	22	23	7	2	-
Male	-	1	1	1	4	8	10	9	7	1	-	-
Female	-	-	1	2	3	3	2	13	16	6	2	-
White, both sexes	-	1	2	3	7	11	12	22	23	7	2	-
Male	-	1	1	1	4	8	10	9	7	1	-	-
Female	-	-	1	2	3	3	2	13	16	6	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	2	6	4	13	12	18	12	32	19	12	5	-
Male	1	6	2	6	5	7	3	11	4	3	2	-
Female	1	-	2	7	7	11	9	21	15	9	3	-
White, both sexes	1	3	3	10	8	12	9	23	11	9	4	-
Male	1	3	1	4	4	5	1	9	3	3	2	-
Female	-	-	2	6	4	7	8	14	8	6	2	-
Black, both sexes	1	3	1	3	4	6	3	9	8	3	1	-
Male	-	3	1	2	1	2	2	2	1	-	-	-
Female	1	-	-	1	3	4	1	7	7	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Missouri												
All races, both sexes	4	2	7	10	9	8	12	16	13	10	2	-
Male	2	2	5	5	4	4	5	7	2	4	1	-
Female	2	-	2	5	5	4	7	9	11	6	1	-
White, both sexes	3	2	7	8	8	7	11	15	13	10	1	-
Male	1	2	5	5	4	4	5	7	2	4	-	-
Female	2	-	2	3	4	3	6	8	11	6	1	-
Black, both sexes	1	-	-	2	1	1	1	1	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	2	1	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	2	1	6	13	11	6	2	-
Male	-	-	-	-	1	1	3	4	3	1	1	-
Female	-	-	-	-	1	-	3	9	8	5	1	-
White, both sexes	-	-	-	-	2	1	6	13	11	6	2	-
Male	-	-	-	-	1	1	3	4	3	1	1	-
Female	-	-	-	-	1	-	3	9	8	5	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Nebraska												
All races, both sexes	-	4	5	2	5	8	11	7	16	8	2	-
Male	-	3	4	2	3	6	7	-	8	1	-	-
Female	-	1	1	-	2	2	4	7	8	7	2	-
White, both sexes	-	3	5	2	5	8	11	7	16	8	2	-
Male	-	2	4	2	3	6	7	-	8	1	-	-
Female	-	1	1	-	2	2	4	7	8	7	2	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1	-	3	3	2	7	2	9	1	1	-	-
Male	1	-	2	2	1	6	2	2	1	-	-	-
Female	-	-	1	1	1	1	-	7	-	1	-	-
White, both sexes	-	-	3	3	1	6	2	8	1	1	-	-
Male	-	-	2	2	1	6	2	2	1	-	-	-
Female	-	-	1	1	-	-	-	6	-	1	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	1	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
New Hampshire												
All races, both sexes	-	-	1	2	-	1	-	6	2	3	1	-
Male	-	-	1	-	-	1	-	3	1	-	-	-
Female	-	-	-	2	-	-	-	3	1	3	1	-
White, both sexes	-	-	1	2	-	1	-	6	2	3	1	-
Male	-	-	1	-	-	1	-	3	1	-	-	-
Female	-	-	-	2	-	-	-	3	1	3	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	4	1	3	3	16	13	21	30	28	8	5	-
Male	4	1	2	1	11	7	9	17	7	2	1	-
Female	-	-	1	2	5	6	12	13	21	6	4	-
White, both sexes	2	1	3	2	11	12	19	26	26	8	3	-
Male	2	1	2	1	8	6	9	14	6	2	1	-
Female	-	-	1	1	3	6	10	12	20	6	2	-
Black, both sexes	2	-	-	1	4	1	2	4	2	-	2	-
Male	2	-	-	-	3	1	-	3	1	-	-	-
Female	-	-	-	1	1	-	2	1	1	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
New Mexico												
All races, both sexes	2	-	2	-	1	-	3	3	-	1	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	2	-	1	-	-	-	3	3	-	1	-	-
White, both sexes	2	-	2	-	1	-	3	3	-	1	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	2	-	1	-	-	-	3	3	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	3	4	6	9	18	26	51	57	54	17	8	-
Male	3	3	3	5	8	14	22	22	21	2	-	-
Female	-	1	3	4	10	12	29	35	33	15	8	-
White, both sexes	3	4	4	8	17	25	46	55	49	14	8	-
Male	3	3	2	4	8	14	21	22	19	2	-	-
Female	-	1	2	4	9	11	25	33	30	12	8	-
Black, both sexes	-	-	2	1	1	1	4	2	5	3	-	-
Male	-	-	1	1	-	-	-	-	2	-	-	-
Female	-	-	1	-	1	1	4	2	3	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
North Carolina												
All races, both sexes	3	11	10	6	12	15	22	13	17	10	4	-
Male	1	6	6	2	7	9	12	2	5	3	-	-
Female	2	5	4	4	5	6	10	11	12	7	4	-
White, both sexes	1	8	5	5	9	10	19	13	16	9	4	-
Male	1	4	4	1	6	6	11	2	5	3	-	-
Female	-	4	1	4	3	4	8	11	11	6	4	-
Black, both sexes	2	3	5	1	3	5	3	-	1	1	-	-
Male	-	2	2	1	1	3	1	-	-	-	-	-
Female	2	1	3	-	2	2	2	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	1	-	1	1	3	-	-
Male	-	-	-	-	-	1	-	1	-	1	-	-
Female	-	-	-	-	-	-	-	-	1	2	-	-
White, both sexes	-	-	-	-	-	1	-	1	1	3	-	-
Male	-	-	-	-	-	1	-	1	-	1	-	-
Female	-	-	-	-	-	-	-	-	1	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Ohio												
All races, both sexes	1	2	3	8	9	20	19	14	14	8	1	-
Male	-	1	3	7	4	11	10	4	5	1	-	-
Female	1	1	-	1	5	9	9	10	9	7	1	-
White, both sexes	1	1	2	6	9	19	18	12	14	8	1	-
Male	-	1	2	5	4	11	10	4	5	1	-	-
Female	1	-	-	1	5	8	8	8	9	7	1	-
Black, both sexes	-	1	1	2	-	1	1	2	-	-	-	-
Male	-	-	1	2	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2	-	5	5	7	9	10	23	17	7	-	-
Male	-	-	2	2	3	3	7	6	2	1	-	-
Female	2	-	3	3	4	6	3	17	15	6	-	-
White, both sexes	1	-	4	5	7	9	10	20	16	7	-	-
Male	-	-	2	2	3	3	7	6	2	1	-	-
Female	1	-	2	3	4	6	3	14	14	6	-	-
Black, both sexes	1	-	-	-	-	-	-	2	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	2	1	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Oregon												
All races, both sexes	1	2	1	3	5	5	10	12	10	8	-	-
Male	1	1	1	3	4	3	4	7	3	2	-	-
Female	-	1	-	-	1	2	6	5	7	6	-	-
White, both sexes	1	2	-	3	5	5	10	11	10	8	-	-
Male	1	1	-	3	4	3	4	6	3	2	-	-
Female	-	1	-	-	1	2	6	5	7	6	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	4	12	13	14	18	34	48	59	48	24	5	-
Male	1	9	10	9	11	15	21	18	13	7	1	-
Female	3	3	3	5	7	19	27	41	35	17	4	-
White, both sexes	1	12	12	12	18	30	47	54	47	22	5	-
Male	1	9	9	8	11	13	20	18	13	6	1	-
Female	-	3	3	4	7	17	27	36	34	16	4	-
Black, both sexes	3	-	1	1	-	4	1	5	-	1	-	-
Male	-	-	1	-	-	2	1	-	-	-	-	-
Female	3	-	-	1	-	2	-	5	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	1	1	-	-
Male	-	-	-	1	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Rhode Island												
All races, both sexes	-	-	-	1	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	1	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	3	7	12	7	9	15	19	24	15	9	-	-
Male	3	4	8	3	4	5	12	9	3	3	-	-
Female	-	3	4	4	5	10	7	15	12	6	-	-
White, both sexes	1	4	4	1	6	10	12	22	12	6	-	-
Male	1	2	3	1	2	4	8	9	3	2	-	-
Female	-	2	1	-	4	6	4	13	9	4	-	-
Black, both sexes	2	3	8	6	3	5	7	2	3	3	-	-
Male	2	2	5	2	2	1	4	-	-	1	-	-
Female	-	1	3	4	1	4	3	2	3	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
South Dakota												
All races, both sexes	1	-	-	1	1	-	-	1	1	1	1	-
Male	-	-	-	1	-	-	-	1	1	1	-	-
Female	1	-	-	-	1	-	-	-	-	-	1	-
White, both sexes	1	-	-	1	1	-	-	1	1	1	1	-
Male	-	-	-	1	-	-	-	1	1	1	-	-
Female	1	-	-	-	1	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	1	1	5	5	6	7	15	19	14	4	1	-
Male	-	1	4	2	2	5	5	6	3	1	-	-
Female	1	-	1	3	4	2	10	13	11	3	1	-
White, both sexes	1	1	4	3	5	6	11	14	12	3	1	-
Male	-	1	4	1	2	5	4	5	3	1	-	-
Female	1	-	-	2	3	1	7	9	9	2	1	-
Black, both sexes	-	-	1	2	1	1	4	5	2	1	-	-
Male	-	-	-	1	-	-	1	1	-	-	-	-
Female	-	-	1	1	1	1	3	4	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Texas												
All races, both sexes	16	14	23	22	28	56	54	63	50	17	11	-
Male	11	8	14	12	20	31	27	28	11	2	4	-
Female	5	6	9	10	8	25	27	35	39	15	7	-
White, both sexes	14	11	15	18	28	49	46	55	45	15	10	-
Male	10	7	10	10	20	26	22	27	10	2	4	-
Female	4	4	5	8	8	23	24	28	35	13	6	-
Black, both sexes	2	3	7	4	-	7	7	8	3	2	1	-
Male	1	1	3	2	-	5	4	1	-	-	-	-
Female	1	2	4	2	-	2	3	7	3	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	1	-	2	-	-	-
Male	-	-	1	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Utah												
All races, both sexes	1	1	-	2	-	6	4	6	4	3	-	-
Male	1	1	-	1	-	4	2	3	-	1	-	-
Female	-	-	-	1	-	2	2	3	4	2	-	-
White, both sexes	-	1	-	2	-	6	4	6	4	3	-	-
Male	-	1	-	1	-	4	2	3	-	1	-	-
Female	-	-	-	1	-	2	2	3	4	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Vermont												
All races, both sexes	-	1	-	-	-	-	-	-	-	2	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	2	-	-
White, both sexes	-	1	-	-	-	-	-	-	-	2	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	11	20	25	21	22	36	61	61	49	27	6	-
Male	8	10	15	13	13	14	24	33	9	2	1	-
Female	3	10	10	8	9	22	37	28	40	25	5	-
White, both sexes	5	9	10	14	15	29	46	51	41	22	6	-
Male	4	3	6	10	9	12	19	27	6	2	1	-
Female	1	6	4	4	6	17	27	24	35	20	5	-
Black, both sexes	6	11	15	7	7	7	13	10	8	5	-	-
Male	4	7	9	3	4	2	4	6	3	-	-	-
Female	2	4	6	4	3	5	9	4	5	5	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Washington												
All races, both sexes	-	-	1	-	-	2	5	6	13	4	-	-
Male	-	-	-	-	-	-	2	3	6	-	-	-
Female	-	-	1	-	-	2	3	3	7	4	-	-
White, both sexes	-	-	1	-	-	2	5	6	13	4	-	-
Male	-	-	-	-	-	-	2	3	6	-	-	-
Female	-	-	1	-	-	2	3	3	7	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	1	3	4	5	3	13	20	20	19	14	3	-
Male	-	-	4	5	2	8	3	6	4	1	1	-
Female	1	3	-	-	1	5	17	14	15	13	2	-
White, both sexes	1	3	4	5	3	13	20	20	19	14	2	-
Male	-	-	4	5	2	8	3	6	4	1	1	-
Female	1	3	-	-	1	5	17	14	15	13	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Wisconsin												
All races, both sexes	1	1	1	4	3	4	9	4	8	3	1	-
Male	1	1	1	3	1	1	6	2	2	2	-	-
Female	-	-	-	1	2	3	3	2	6	1	1	-
White, both sexes	-	1	1	4	3	4	9	4	8	3	1	-
Male	-	1	1	3	1	1	6	2	2	2	-	-
Female	-	-	-	1	2	3	3	2	6	1	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	1	-	-	-	-	3	1	2	2	-	-
Male	-	1	-	-	-	-	1	1	-	2	-	-
Female	-	-	-	-	-	-	2	-	2	-	-	-
White, both sexes	-	1	-	-	-	-	3	1	2	2	-	-
Male	-	1	-	-	-	-	1	1	-	2	-	-
Female	-	-	-	-	-	-	2	-	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
United States												
All races, both sexes	690	40	34	21	19	27	31	33	25	41	60	73
Male	322	25	18	8	5	15	19	21	14	19	33	27
Female	368	15	16	13	14	12	12	12	11	22	27	46
White, both sexes	528	22	17	12	8	19	27	19	22	32	46	58
Male	234	11	8	2	3	10	16	12	12	17	26	20
Female	294	11	9	10	5	9	11	7	10	15	20	38
Black, both sexes	140	15	13	7	11	8	2	14	2	8	10	15
Male	73	11	9	4	2	5	2	9	1	1	5	7
Female	67	4	4	3	9	3	-	5	1	7	5	8
American Indian 1/, both sexes	8	2	1	1	-	-	1	-	-	-	2	-
Male	6	2	1	1	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	1	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	14	1	3	1	-	-	1	-	1	1	2	-
Male	9	1	-	1	-	-	1	-	1	1	1	-
Female	5	-	3	-	-	-	-	-	-	-	1	-
Alabama												
All races, both sexes	6	-	1	1	-	-	1	-	-	-	-	-
Male	5	-	1	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	5	-	-	1	-	-	1	-	-	-	-	-
Male	4	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Alaska												
All races, both sexes	2	-	-	-	-	1	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	1	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Arkansas												
All races, both sexes	11	2	-	-	1	1	-	-	1	-	-	1
Male	5	1	-	-	1	1	-	-	-	-	-	1
Female	6	1	-	-	-	-	-	-	1	-	-	-
White, both sexes	9	2	-	-	1	-	-	-	1	-	-	1
Male	4	1	-	-	1	-	-	-	-	-	-	1
Female	5	1	-	-	-	-	-	-	1	-	-	-
Black, both sexes	2	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	52	2	2	-	1	2	3	-	3	5	6	6
Male	21	1	-	-	-	1	3	-	-	4	4	1
Female	31	1	2	-	1	1	-	-	3	1	2	5
White, both sexes	41	2	1	-	1	2	2	-	3	3	4	6
Male	16	1	-	-	-	1	2	-	-	3	3	1
Female	25	1	1	-	1	1	-	-	3	-	1	5
Black, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	1	-	-	-	1	-	-	1	2	-
Male	4	-	-	-	-	-	1	-	-	1	1	-
Female	3	-	1	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Colorado												
All races, both sexes	11	-	1	1	-	-	-	1	-	-	1	-
Male	3	-	1	-	-	-	-	-	-	-	-	-
Female	8	-	-	1	-	-	-	1	-	-	1	-
White, both sexes	11	-	1	1	-	-	-	1	-	-	1	-
Male	3	-	1	-	-	-	-	-	-	-	-	-
Female	8	-	-	1	-	-	-	1	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	7	-	1	1	-	-	2	-	-	1	1	-
Male	3	-	-	-	-	-	2	-	-	1	-	-
Female	4	-	1	1	-	-	-	-	-	-	1	-
White, both sexes	6	-	-	1	-	-	2	-	-	1	1	-
Male	3	-	-	-	-	-	2	-	-	1	-	-
Female	3	-	-	1	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Delaware												
All races, both sexes	3	-	-	-	-	-	1	1	-	-	-	-
Male	2	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Florida												
All races, both sexes	54	3	3	-	2	2	1	4	2	4	4	2
Male	24	1	2	-	-	1	1	2	1	3	2	1
Female	30	2	1	-	2	1	-	2	1	1	2	1
White, both sexes	39	-	2	-	-	2	1	2	-	2	4	1
Male	17	-	1	-	-	1	1	2	-	2	2	-
Female	22	-	1	-	-	1	-	-	-	-	2	1
Black, both sexes	15	3	1	-	2	-	-	2	2	2	-	1
Male	7	1	1	-	-	-	-	-	1	1	-	1
Female	8	2	-	-	2	-	-	2	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	41	4	1	2	2	2	3	-	1	3	7	4
Male	21	3	1	2	1	1	1	-	-	-	4	1
Female	20	1	-	-	1	1	2	-	1	3	3	3
White, both sexes	29	2	-	-	1	2	3	-	1	3	4	3
Male	13	2	-	-	1	1	1	-	-	-	1	1
Female	16	-	-	-	-	1	2	-	1	3	3	2
Black, both sexes	12	2	1	2	1	-	-	-	-	-	3	1
Male	8	1	1	2	-	-	-	-	-	-	3	-
Female	4	1	-	-	1	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Hawaii												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Illinois												
All races, both sexes	24	4	2	-	1	-	1	-	-	2	1	5
Male	14	4	1	-	-	-	1	-	-	2	1	1
Female	10	-	1	-	1	-	-	-	-	-	-	4
White, both sexes	17	1	1	-	1	-	1	-	-	2	1	2
Male	10	1	1	-	-	-	1	-	-	2	1	-
Female	7	-	-	-	1	-	-	-	-	-	-	2
Black, both sexes	7	3	1	-	-	-	-	-	-	-	-	3
Male	4	3	-	-	-	-	-	-	-	-	-	1
Female	3	-	1	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	14	-	-	-	-	2	-	2	-	-	1	2
Male	10	-	-	-	-	2	-	1	-	-	1	2
Female	4	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	12	-	-	-	-	2	-	1	-	-	1	1
Male	8	-	-	-	-	2	-	-	-	-	1	1
Female	4	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	1	-	-	-	1
Male	2	-	-	-	-	-	-	1	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Iowa												
All races, both sexes	7	-	-	-	-	-	1	1	-	-	-	-
Male	5	-	-	-	-	-	1	1	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	1	-	-	-	-	-
Male	4	-	-	-	-	-	1	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	12	-	-	-	-	-	-	-	1	1	1	4
Male	5	-	-	-	-	-	-	-	-	1	1	1
Female	7	-	-	-	-	-	-	-	1	-	-	3
White, both sexes	11	-	-	-	-	-	-	-	1	1	1	4
Male	4	-	-	-	-	-	-	-	-	1	1	1
Female	7	-	-	-	-	-	-	-	1	-	-	3
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Kentucky												
All races, both sexes	8	-	-	1	-	-	-	-	-	1	2	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	1	-	-	-	-	-	1	2	-
White, both sexes	5	-	-	1	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	1	-	-	-	-	-	1	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	2	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	8	1	1	-	1	-	-	-	-	1	-	-
Male	5	1	1	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	5	1	-	-	-	-	-	-	-	1	-	-
Male	3	1	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	1	-	1	-	-	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Maine												
All races, both sexes	5	-	-	-	-	1	-	-	-	-	1	2
Male	3	-	-	-	-	-	-	-	-	-	-	2
Female	2	-	-	-	-	1	-	-	-	-	1	-
White, both sexes	3	-	-	-	-	1	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	17	-	-	-	3	-	-	-	2	4	3	2
Male	6	-	-	-	-	-	-	-	1	1	2	-
Female	11	-	-	-	3	-	-	-	1	3	1	2
White, both sexes	12	-	-	-	2	-	-	-	2	2	3	1
Male	6	-	-	-	-	-	-	-	1	1	2	-
Female	6	-	-	-	2	-	-	-	1	1	1	1
Black, both sexes	5	-	-	-	1	-	-	-	-	2	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	1	-	-	-	-	2	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Massachusetts												
All races, both sexes	12	-	-	-	1	-	-	3	-	2	-	3
Male	5	-	-	-	1	-	-	2	-	-	-	1
Female	7	-	-	-	-	-	-	1	-	2	-	2
White, both sexes	11	-	-	-	-	-	-	3	-	2	-	3
Male	4	-	-	-	-	-	-	2	-	-	-	1
Female	7	-	-	-	-	-	-	1	-	2	-	2
Black, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	23	2	-	1	1	1	1	1	-	1	-	3
Male	11	2	-	-	-	1	-	1	-	1	-	-
Female	12	-	-	1	1	-	1	-	-	-	-	3
White, both sexes	19	2	-	1	-	1	1	1	-	1	-	2
Male	10	2	-	-	-	1	-	1	-	1	-	-
Female	9	-	-	1	-	-	1	-	-	-	-	2
Black, both sexes	4	-	-	-	1	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	1	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Minnesota												
All races, both sexes	23	3	3	1	-	1	1	4	-	2	2	2
Male	12	3	2	1	-	-	1	3	-	-	-	-
Female	11	-	1	-	-	1	-	1	-	2	2	2
White, both sexes	17	2	3	-	-	1	1	2	-	1	1	2
Male	9	2	2	-	-	-	1	2	-	-	-	-
Female	8	-	1	-	-	1	-	-	-	1	1	2
Black, both sexes	4	-	-	-	-	-	-	2	-	1	1	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	3	-	-	-	-	-	-	1	-	1	1	-
American Indian 1/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	4	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Missouri												
All races, both sexes	11	1	-	1	-	1	-	-	1	-	-	-
Male	5	-	-	1	-	1	-	-	1	-	-	-
Female	6	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	1	-	-	1	-	-	-
Male	4	-	-	-	-	1	-	-	1	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	1	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	2	-	-	-	-	1	-	-	-	-	-	1
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	2	-	-	-	-	1	-	-	-	-	-	1
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Nebraska												
All races, both sexes	3	1	1	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	1	1	-	-	-	-	-	-	1	-	-
White, both sexes	3	1	1	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	1	1	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	2	1	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	1	-	-	-	-	-	-
White, both sexes	2	1	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
New Hampshire												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	22	1	-	-	-	1	-	-	3	-	3	5
Male	8	-	-	-	-	-	-	-	3	-	2	-
Female	14	1	-	-	-	1	-	-	-	-	1	5
White, both sexes	16	1	-	-	-	-	-	-	3	-	3	2
Male	7	-	-	-	-	-	-	-	3	-	2	-
Female	9	1	-	-	-	-	-	-	-	-	1	2
Black, both sexes	5	-	-	-	-	1	-	-	-	-	-	3
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	1	-	-	-	-	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
New Mexico												
All races, both sexes	4	-	-	-	-	1	1	-	1	-	-	1
Male	2	-	-	-	-	1	-	-	-	-	-	1
Female	2	-	-	-	-	-	1	-	1	-	-	-
White, both sexes	4	-	-	-	-	1	1	-	1	-	-	1
Male	2	-	-	-	-	1	-	-	-	-	-	1
Female	2	-	-	-	-	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	41	2	3	2	2	3	2	2	-	-	2	3
Male	20	2	3	-	1	2	1	2	-	-	2	1
Female	21	-	-	2	1	1	1	-	-	-	-	2
White, both sexes	24	-	-	1	-	1	1	-	-	-	2	2
Male	8	-	-	-	-	-	-	-	-	-	2	-
Female	16	-	-	1	-	1	1	-	-	-	-	2
Black, both sexes	16	1	3	1	2	2	1	2	-	-	-	1
Male	11	1	3	-	1	2	1	2	-	-	-	1
Female	5	-	-	1	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
North Carolina												
All races, both sexes	27	-	4	3	-	1	-	2	3	1	3	4
Male	15	-	2	1	-	1	-	2	3	1	3	-
Female	12	-	2	2	-	-	-	-	-	-	-	4
White, both sexes	20	-	1	3	-	-	-	1	2	1	2	4
Male	9	-	-	1	-	-	-	1	2	1	2	-
Female	11	-	1	2	-	-	-	-	-	-	-	4
Black, both sexes	4	-	2	-	-	1	-	1	-	-	-	-
Male	3	-	1	-	-	1	-	1	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	1	-	-	-	-	-	-	-	1	-
Male	2	-	1	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	2	2	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	2	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Ohio												
All races, both sexes	33	-	-	-	-	1	2	3	-	3	5	3
Male	13	-	-	-	-	1	2	2	-	1	1	2
Female	20	-	-	-	-	-	-	1	-	2	4	1
White, both sexes	26	-	-	-	-	-	2	1	-	2	4	3
Male	10	-	-	-	-	-	2	1	-	1	1	2
Female	16	-	-	-	-	-	-	-	-	1	3	1
Black, both sexes	7	-	-	-	-	1	-	2	-	1	1	-
Male	3	-	-	-	-	1	-	1	-	-	-	-
Female	4	-	-	-	-	-	-	1	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	7	1	1	1	-	-	-	-	-	1	-	-
Male	5	1	-	1	-	-	-	-	-	1	-	-
Female	2	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	1	-	1	-	-	-	-	-	-	-	-
Male	2	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Oregon												
All races, both sexes	10	-	1	-	-	-	-	-	-	-	2	2
Male	6	-	-	-	-	-	-	-	-	-	2	1
Female	4	-	1	-	-	-	-	-	-	-	-	1
White, both sexes	10	-	1	-	-	-	-	-	-	-	2	2
Male	6	-	-	-	-	-	-	-	-	-	2	1
Female	4	-	1	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	29	1	1	2	1	-	1	-	2	1	1	5
Male	11	-	-	1	1	-	1	-	1	-	1	2
Female	18	1	1	1	-	-	-	-	1	1	-	3
White, both sexes	24	1	1	-	1	-	1	-	2	1	1	4
Male	8	-	-	-	1	-	1	-	1	-	1	1
Female	16	1	1	-	-	-	-	-	1	1	-	3
Black, both sexes	5	-	-	2	-	-	-	-	-	-	-	1
Male	3	-	-	1	-	-	-	-	-	-	-	1
Female	2	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Rhode Island												
All races, both sexes	3	1	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	2	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	1	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	2	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	14	1	-	-	-	-	-	1	1	-	2	1
Male	8	1	-	-	-	-	-	1	-	-	1	1
Female	6	-	-	-	-	-	-	-	1	-	1	-
White, both sexes	8	-	-	-	-	-	-	-	1	-	2	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	1	-	1	-
Black, both sexes	6	1	-	-	-	-	-	1	-	-	-	1
Male	6	1	-	-	-	-	-	1	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
South Dakota												
All races, both sexes	2	-	1	-	-	-	-	-	-	-	1	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	18	2	-	1	-	-	1	3	2	-	1	-
Male	12	2	-	-	-	-	1	3	2	-	-	-
Female	6	-	-	1	-	-	-	-	-	-	1	-
White, both sexes	13	-	-	-	-	-	1	2	2	-	1	-
Male	8	-	-	-	-	-	1	2	2	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	5	2	-	1	-	-	-	1	-	-	-	-
Male	4	2	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Texas												
All races, both sexes	53	3	4	2	2	2	5	2	-	5	3	3
Male	21	1	1	-	-	1	2	-	-	2	3	3
Female	32	2	3	2	2	1	3	2	-	3	-	-
White, both sexes	40	2	3	2	-	1	4	1	-	4	2	3
Male	17	-	1	-	-	1	2	-	-	2	2	3
Female	23	2	2	2	-	-	2	1	-	2	-	-
Black, both sexes	12	1	1	-	2	1	-	1	-	1	1	-
Male	4	1	-	-	-	-	-	-	-	-	1	-
Female	8	-	1	-	2	1	-	1	-	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	10	1	-	-	-	-	-	1	-	-	2	2
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	5	1	-	-	-	-	-	1	-	-	2	1
White, both sexes	10	1	-	-	-	-	-	1	-	-	2	2
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	5	1	-	-	-	-	-	1	-	-	2	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Vermont												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	13	1	-	-	-	1	1	1	1	-	2	1
Male	6	1	-	-	-	-	-	1	1	-	2	-
Female	7	-	-	-	-	1	1	-	-	-	-	1
White, both sexes	8	-	-	-	-	1	1	1	1	-	1	1
Male	3	-	-	-	-	-	-	1	1	-	1	-
Female	5	-	-	-	-	1	1	-	-	-	-	1
Black, both sexes	5	1	-	-	-	-	-	-	-	-	1	-
Male	3	1	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Washington												
All races, both sexes	9	-	1	-	1	-	-	1	-	1	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	1	-	1	-	-	1	-	1	-	1
White, both sexes	8	-	-	-	1	-	-	1	-	1	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	1	-	-	1	-	1	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	6	-	2	-	-	-	1	-	-	1	-	1
Male	3	-	2	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	1	-	-	1	-	1
White, both sexes	5	-	1	-	-	-	1	-	-	1	-	1
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	1	-	-	1	-	1
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Wisconsin												
All races, both sexes	17	-	-	1	-	-	2	-	1	-	3	1
Male	9	-	-	-	-	-	1	-	1	-	1	1
Female	8	-	-	1	-	-	1	-	-	-	2	-
White, both sexes	17	-	-	1	-	-	2	-	1	-	3	1
Male	9	-	-	-	-	-	1	-	1	-	1	1
Female	8	-	-	1	-	-	1	-	-	-	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
United States												
All races, both sexes	50	36	32	11	13	19	36	29	32	22	6	-
Male	18	18	19	6	9	9	15	11	7	4	2	-
Female	32	18	13	5	4	10	21	18	25	18	4	-
White, both sexes	40	30	27	10	11	14	32	28	29	19	6	-
Male	12	15	15	5	7	6	13	11	7	4	2	-
Female	28	15	12	5	4	8	19	17	22	15	4	-
Black, both sexes	8	6	5	1	1	4	3	1	3	3	-	-
Male	4	3	4	1	1	3	1	-	-	-	-	-
Female	4	3	1	-	-	1	2	1	3	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	1	1	-	-	-	-	-	-
Male	2	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Alabama												
All races, both sexes	-	1	-	-	-	-	-	2	-	-	-	-
Male	-	1	-	-	-	-	-	2	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	2	-	-	-	-
Male	-	1	-	-	-	-	-	2	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	2	3	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	1	3	-	-
White, both sexes	-	-	-	-	-	-	-	-	2	2	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	1	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	7	5	1	1	2	1	1	1	1	1	1	-
Male	2	1	1	-	1	-	-	-	-	1	1	-
Female	5	4	-	1	1	1	1	1	1	-	-	-
White, both sexes	5	5	-	1	2	-	1	1	-	1	1	-
Male	1	1	-	-	1	-	-	-	-	1	1	-
Female	4	4	-	1	1	-	1	1	-	-	-	-
Black, both sexes	1	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Colorado												
All races, both sexes	4	-	2	-	-	-	-	-	1	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	3	-	1	-	-	-	-	-	1	-	-	-
White, both sexes	4	-	2	-	-	-	-	-	1	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	3	-	1	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Delaware												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Florida												
All races, both sexes	3	3	1	-	1	2	5	1	6	4	1	-
Male	1	3	1	-	1	1	1	1	-	1	-	-
Female	2	-	-	-	-	1	4	-	6	3	1	-
White, both sexes	2	3	-	-	1	2	5	1	6	4	1	-
Male	-	3	-	-	1	1	1	1	-	1	-	-
Female	2	-	-	-	-	1	4	-	6	3	1	-
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	2	3	2	1	1	-	1	1	1	-	-	-
Male	1	2	1	1	-	-	-	1	1	-	-	-
Female	1	1	1	-	1	-	1	-	-	-	-	-
White, both sexes	2	2	2	-	1	-	1	1	1	-	-	-
Male	1	2	1	-	-	-	-	1	1	-	-	-
Female	1	-	1	-	1	-	1	-	-	-	-	-
Black, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Hawaii												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Illinois												
All races, both sexes	1	-	1	1	-	2	1	1	1	-	-	-
Male	1	-	1	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	2	-	1	1	-	-	-
White, both sexes	1	-	1	1	-	2	1	1	1	-	-	-
Male	1	-	1	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	2	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1	-	2	1	-	-	1	2	-	-	-	-
Male	-	-	1	1	-	-	-	2	-	-	-	-
Female	1	-	1	-	-	-	1	-	-	-	-	-
White, both sexes	1	-	2	1	-	-	1	2	-	-	-	-
Male	-	-	1	1	-	-	-	2	-	-	-	-
Female	1	-	1	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Iowa												
All races, both sexes	-	1	1	-	2	-	-	1	-	-	-	-
Male	-	-	-	-	2	-	-	1	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	1	-	2	-	-	1	-	-	-	-
Male	-	-	-	-	2	-	-	1	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	1	1	-	-	1	-	1	1	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	1	1	-	-
White, both sexes	-	-	-	1	-	-	1	-	1	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	1	1	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Kentucky												
All races, both sexes	-	1	-	-	1	-	1	1	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	1	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	1	-	-	-	-	2	-	-	1	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	2	-	-	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	1	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Maine												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	2	1	-	-	-	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Massachusetts												
All races, both sexes	1	1	-	-	-	-	-	-	-	1	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	1	1	-	-	-	-	-	-	-	1	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1	3	1	-	-	2	-	4	1	-	-	-
Male	1	1	1	-	-	2	-	1	-	-	-	-
Female	-	2	-	-	-	-	-	3	1	-	-	-
White, both sexes	-	2	1	-	-	2	-	4	1	-	-	-
Male	-	1	1	-	-	2	-	1	-	-	-	-
Female	-	1	-	-	-	-	-	3	1	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Minnesota												
All races, both sexes	-	-	1	-	-	-	1	-	2	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	1	-	-	-	1	-	2	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	1	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	2	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Missouri												
All races, both sexes	1	-	-	-	-	2	1	1	2	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	1	-	-	-	-	2	-	-	2	-	-	-
White, both sexes	1	-	-	-	-	2	1	1	2	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	1	-	-	-	-	2	-	-	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
New Hampshire												
All races, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	6	-	-	-	2	-	-	1	-	-	-	-
Male	1	-	-	-	2	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	5	-	-	-	1	-	-	1	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	4	3	1	-	-	1	2	2	3	2	2	-
Male	2	1	1	-	-	-	-	-	1	-	1	-
Female	2	2	-	-	-	1	2	2	2	2	1	-
White, both sexes	3	3	1	-	-	-	1	2	3	2	2	-
Male	2	1	1	-	-	-	-	-	1	-	1	-
Female	1	2	-	-	-	-	1	2	2	2	1	-
Black, both sexes	1	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
North Carolina												
All races, both sexes	1	2	-	1	1	-	-	-	1	-	-	-
Male	-	1	-	-	-	-	-	-	1	-	-	-
Female	1	1	-	1	1	-	-	-	-	-	-	-
White, both sexes	1	2	-	1	1	-	-	-	1	-	-	-
Male	-	1	-	-	-	-	-	-	1	-	-	-
Female	1	1	-	1	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Ohio												
All races, both sexes	3	1	1	1	-	2	1	4	2	1	-	-
Male	1	-	-	1	-	-	-	-	1	1	-	-
Female	2	1	1	-	-	2	1	4	1	-	-	-
White, both sexes	2	1	1	1	-	2	1	3	2	1	-	-
Male	-	-	-	1	-	-	-	-	1	1	-	-
Female	2	1	1	-	-	2	1	3	1	-	-	-
Black, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	1	1	-	-	1	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	1	1	-	-	1	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Oregon												
All races, both sexes	-	-	-	1	1	-	1	-	1	-	1	-
Male	-	-	-	1	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	1	-
White, both sexes	-	-	-	1	1	-	1	-	1	-	1	-
Male	-	-	-	1	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	3	3	2	-	-	-	2	1	1	1	1	-
Male	1	1	1	-	-	-	-	1	-	-	-	-
Female	2	2	1	-	-	-	2	-	1	1	1	-
White, both sexes	2	3	1	-	-	-	2	1	1	1	1	-
Male	1	1	-	-	-	-	-	1	-	-	-	-
Female	1	2	1	-	-	-	2	-	1	1	1	-
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Rhode Island												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	1	1	-	-	2	1	-	1	1	-	-
Male	-	1	-	-	-	2	1	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	1	1	-	-
White, both sexes	1	-	1	-	-	-	1	-	1	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	1	1	-	-
Black, both sexes	-	1	-	-	-	2	-	-	-	-	-	-
Male	-	1	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	1	1	-	-	1	1	2	1	1	-	-
Male	-	-	1	-	-	1	1	1	-	-	-	-
Female	-	1	-	-	-	-	-	1	1	1	-	-
White, both sexes	-	1	1	-	-	1	-	2	1	1	-	-
Male	-	-	1	-	-	1	-	1	-	-	-	-
Female	-	1	-	-	-	-	-	1	1	1	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Texas												
All races, both sexes	6	2	4	-	2	2	1	2	2	1	-	-
Male	4	1	1	-	1	1	-	-	-	-	-	-
Female	2	1	3	-	1	1	1	2	2	1	-	-
White, both sexes	5	2	4	-	2	1	1	2	1	-	-	-
Male	3	1	1	-	1	-	-	-	-	-	-	-
Female	2	1	3	-	1	1	1	2	1	-	-	-
Black, both sexes	1	-	-	-	-	1	-	-	1	1	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	4	-	-	-	-	-	-	-	-	-
Male	-	-	4	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	4	-	-	-	-	-	-	-	-	-
Male	-	-	4	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Vermont												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	2	-	-	-	-	2	-	-	1	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	2	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Black, both sexes	-	2	-	-	-	-	-	-	-	1	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Washington												
All races, both sexes	-	1	-	-	-	1	1	-	1	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
White, both sexes	-	1	-	-	-	1	1	-	1	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Wisconsin												
All races, both sexes	1	-	1	-	-	-	4	1	-	2	-	-
Male	-	-	1	-	-	-	3	-	-	1	-	-
Female	1	-	-	-	-	-	1	1	-	1	-	-
White, both sexes	1	-	1	-	-	-	4	1	-	2	-	-
Male	-	-	1	-	-	-	3	-	-	1	-	-
Female	1	-	-	-	-	-	1	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
United States												
All races, both sexes	11,542	32	6	3	8	28	78	104	123	190	287	421
Male	5,761	20	4	2	5	21	60	80	78	145	198	281
Female	5,781	12	2	1	3	7	18	24	45	45	89	140
White, both sexes	9,829	19	2	2	5	14	45	68	84	140	203	320
Male	4,888	10	1	1	2	12	35	49	59	102	145	214
Female	4,941	9	1	1	3	2	10	19	25	38	58	106
Black, both sexes	1,472	11	3	1	3	13	30	34	31	45	70	86
Male	750	9	3	1	3	8	23	29	16	38	42	57
Female	722	2	-	-	-	5	7	5	15	7	28	29
American Indian 1/, both sexes	65	-	-	-	-	-	1	1	4	3	3	8
Male	37	-	-	-	-	-	-	1	3	3	2	6
Female	28	-	-	-	-	-	1	-	1	-	1	2
Asian or Pacific Islander 2/, both sexes	176	2	1	-	-	1	2	1	4	2	11	7
Male	86	1	-	-	-	1	2	1	-	2	9	4
Female	90	1	1	-	-	-	-	-	4	-	2	3
Alabama												
All races, both sexes	439	-	1	-	-	-	1	3	2	-	6	13
Male	227	-	1	-	-	-	-	3	1	-	6	10
Female	212	-	-	-	-	-	1	-	1	-	-	3
White, both sexes	355	-	-	-	-	-	-	-	1	-	3	8
Male	189	-	-	-	-	-	-	-	-	-	3	5
Female	166	-	-	-	-	-	-	-	1	-	-	3
Black, both sexes	82	-	1	-	-	-	1	3	1	-	3	5
Male	36	-	1	-	-	-	-	3	1	-	3	5
Female	46	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Alaska												
All races, both sexes	18	-	-	-	-	-	-	-	1	-	-	1
Male	8	-	-	-	-	-	-	-	1	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	245	-	-	-	-	1	4	6	-	3	5	13
Male	134	-	-	-	-	-	3	4	-	2	3	8
Female	111	-	-	-	-	1	1	2	-	1	2	5
White, both sexes	227	-	-	-	-	-	3	6	-	3	4	11
Male	128	-	-	-	-	-	3	4	-	2	3	6
Female	99	-	-	-	-	-	-	2	-	1	1	5
Black, both sexes	10	-	-	-	-	1	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	8	-	-	-	-	-	1	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	1	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Arkansas												
All races, both sexes	186	-	-	-	-	-	-	1	1	2	2	8
Male	76	-	-	-	-	-	-	1	1	2	1	6
Female	110	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	170	-	-	-	-	-	-	-	1	2	2	8
Male	69	-	-	-	-	-	-	-	1	2	1	6
Female	101	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	16	-	-	-	-	-	-	1	-	-	-	-
Male	7	-	-	-	-	-	-	1	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1,140	7	-	1	-	4	6	13	14	24	45	69
Male	550	4	-	1	-	4	5	9	11	19	35	44
Female	590	3	-	-	-	-	1	4	3	5	10	25
White, both sexes	962	5	-	1	-	2	2	9	10	19	31	50
Male	453	2	-	1	-	2	1	6	8	16	25	31
Female	509	3	-	-	-	-	1	3	2	3	6	19
Black, both sexes	105	1	-	-	-	2	3	3	3	4	10	17
Male	56	1	-	-	-	2	3	2	3	2	7	11
Female	49	-	-	-	-	-	-	1	-	2	3	6
American Indian 1/, both sexes	9	-	-	-	-	-	-	1	-	-	2	1
Male	8	-	-	-	-	-	-	1	-	-	2	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	64	1	-	-	-	-	1	-	1	1	2	1
Male	33	1	-	-	-	-	1	-	-	1	1	1
Female	31	-	-	-	-	-	-	-	1	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Colorado												
All races, both sexes	115	-	-	-	-	-	-	1	1	2	5	9
Male	67	-	-	-	-	-	-	1	1	2	2	6
Female	48	-	-	-	-	-	-	-	-	-	3	3
White, both sexes	111	-	-	-	-	-	-	1	1	2	5	8
Male	64	-	-	-	-	-	-	1	1	2	2	6
Female	47	-	-	-	-	-	-	-	-	-	3	2
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	113	-	-	-	-	2	1	-	-	1	2	5
Male	55	-	-	-	-	2	1	-	-	1	2	3
Female	58	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	107	-	-	-	-	-	-	-	-	1	2	5
Male	51	-	-	-	-	-	-	-	-	1	2	3
Female	56	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	6	-	-	-	-	2	1	-	-	-	-	-
Male	4	-	-	-	-	2	1	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Delaware												
All races, both sexes	23	-	-	-	-	-	-	-	-	1	-	-
Male	12	-	-	-	-	-	-	-	-	1	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	1	-	-
Male	11	-	-	-	-	-	-	-	-	1	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	2
Male	9	-	-	-	-	-	-	-	-	-	-	2
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Florida												
All races, both sexes	809	1	-	-	-	2	3	1	5	11	10	12
Male	414	1	-	-	-	2	3	1	2	10	6	7
Female	395	-	-	-	-	-	-	-	3	1	4	5
White, both sexes	722	-	-	-	-	1	1	-	3	7	6	11
Male	367	-	-	-	-	1	1	-	2	6	4	7
Female	355	-	-	-	-	-	-	-	1	1	2	4
Black, both sexes	79	1	-	-	-	1	2	1	2	4	3	1
Male	42	1	-	-	-	1	2	1	-	4	1	-
Female	37	-	-	-	-	-	-	-	2	-	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	895	2	-	-	-	1	2	6	4	13	15	31
Male	485	2	-	-	-	1	2	4	2	9	9	24
Female	410	-	-	-	-	-	-	2	2	4	6	7
White, both sexes	663	-	-	-	-	1	-	3	3	7	6	21
Male	367	-	-	-	-	1	-	3	2	4	3	19
Female	296	-	-	-	-	-	-	-	1	3	3	2
Black, both sexes	227	2	-	-	-	-	2	3	1	6	9	10
Male	116	2	-	-	-	-	2	1	-	5	6	5
Female	111	-	-	-	-	-	-	2	1	1	3	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Hawaii												
All races, both sexes	75	1	-	-	-	1	-	-	3	2	6	7
Male	41	-	-	-	-	1	-	-	-	2	5	4
Female	34	1	-	-	-	-	-	-	3	-	1	3
White, both sexes	26	-	-	-	-	-	-	-	-	1	2	1
Male	19	-	-	-	-	-	-	-	-	1	1	1
Female	7	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	48	1	-	-	-	1	-	-	3	1	4	6
Male	22	-	-	-	-	1	-	-	-	1	4	3
Female	26	1	-	-	-	-	-	-	3	-	-	3
Idaho												
All races, both sexes	45	-	-	-	-	-	1	-	1	-	1	3
Male	27	-	-	-	-	-	1	-	-	-	-	2
Female	18	-	-	-	-	-	-	-	1	-	1	1
White, both sexes	44	-	-	-	-	-	1	-	1	-	1	3
Male	26	-	-	-	-	-	1	-	-	-	-	2
Female	18	-	-	-	-	-	-	-	1	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Illinois												
All races, both sexes	454	2	-	-	-	1	3	5	4	7	8	18
Male	214	1	-	-	-	1	1	4	3	2	4	12
Female	240	1	-	-	-	-	2	1	1	5	4	6
White, both sexes	390	1	-	-	-	1	3	4	3	7	5	13
Male	178	1	-	-	-	1	1	3	2	2	3	9
Female	212	-	-	-	-	-	2	1	1	5	2	4
Black, both sexes	60	1	-	-	-	-	-	1	1	-	3	5
Male	33	-	-	-	-	-	-	1	1	-	1	3
Female	27	1	-	-	-	-	-	-	-	-	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	193	1	-	-	-	-	3	4	2	3	5	3
Male	93	-	-	-	-	-	3	1	1	1	4	1
Female	100	1	-	-	-	-	-	3	1	2	1	2
White, both sexes	180	1	-	-	-	-	3	2	1	3	4	3
Male	85	-	-	-	-	-	3	1	1	1	3	1
Female	95	1	-	-	-	-	-	1	-	2	1	2
Black, both sexes	13	-	-	-	-	-	-	2	1	-	1	-
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Iowa												
All races, both sexes	60	1	-	-	-	-	-	-	1	5	2	2
Male	26	-	-	-	-	-	-	-	-	3	1	-
Female	34	1	-	-	-	-	-	-	1	2	1	2
White, both sexes	59	-	-	-	-	-	-	-	1	5	2	2
Male	26	-	-	-	-	-	-	-	-	3	1	-
Female	33	-	-	-	-	-	-	-	1	2	1	2
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	87	1	-	-	-	-	-	-	4	3	3	8
Male	39	-	-	-	-	-	-	-	3	2	1	6
Female	48	1	-	-	-	-	-	-	1	1	2	2
White, both sexes	78	1	-	-	-	-	-	-	3	3	3	5
Male	32	-	-	-	-	-	-	-	2	2	1	3
Female	46	1	-	-	-	-	-	-	1	1	2	2
Black, both sexes	4	-	-	-	-	-	-	-	1	-	-	2
Male	3	-	-	-	-	-	-	-	1	-	-	2
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Kentucky												
All races, both sexes	257	-	-	-	-	-	3	2	2	7	13	7
Male	129	-	-	-	-	-	2	-	2	4	7	4
Female	128	-	-	-	-	-	1	2	-	3	6	3
White, both sexes	244	-	-	-	-	-	3	2	2	7	12	7
Male	124	-	-	-	-	-	2	-	2	4	7	4
Female	120	-	-	-	-	-	1	2	-	3	5	3
Black, both sexes	13	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	304	-	-	-	-	2	2	1	3	2	3	7
Male	129	-	-	-	-	2	1	1	-	2	2	4
Female	175	-	-	-	-	-	1	-	3	-	1	3
White, both sexes	241	-	-	-	-	2	1	-	2	2	2	6
Male	104	-	-	-	-	2	1	-	-	2	1	4
Female	137	-	-	-	-	-	-	-	2	-	1	2
Black, both sexes	61	-	-	-	-	-	1	1	1	-	1	1
Male	23	-	-	-	-	-	-	1	-	-	1	-
Female	38	-	-	-	-	-	1	-	1	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Maine												
All races, both sexes	66	-	-	-	-	-	-	-	-	1	1	1
Male	44	-	-	-	-	-	-	-	-	1	-	-
Female	22	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	65	-	-	-	-	-	-	-	-	1	1	1
Male	43	-	-	-	-	-	-	-	-	1	-	-
Female	22	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	117	1	-	1	1	2	1	6	7	12	12	7
Male	69	-	-	1	-	2	1	3	5	9	11	4
Female	48	1	-	-	1	-	-	3	2	3	1	3
White, both sexes	72	1	-	-	1	-	1	3	2	6	10	4
Male	39	-	-	-	-	-	1	-	2	4	10	2
Female	33	1	-	-	1	-	-	3	-	2	-	2
Black, both sexes	44	-	-	1	-	2	-	3	5	6	2	3
Male	29	-	-	1	-	2	-	3	3	5	1	2
Female	15	-	-	-	-	-	-	-	2	1	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Massachusetts												
All races, both sexes	181	-	-	-	1	-	2	-	-	-	1	3
Male	80	-	-	-	-	-	1	-	-	-	1	2
Female	101	-	-	-	1	-	1	-	-	-	-	1
White, both sexes	172	-	-	-	1	-	2	-	-	-	1	3
Male	73	-	-	-	-	-	1	-	-	-	1	2
Female	99	-	-	-	1	-	1	-	-	-	-	1
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	259	1	-	-	-	-	1	1	4	4	7	7
Male	113	1	-	-	-	-	1	1	2	3	5	3
Female	146	-	-	-	-	-	-	-	2	1	2	4
White, both sexes	226	1	-	-	-	-	-	1	-	3	5	4
Male	94	1	-	-	-	-	-	1	-	2	4	1
Female	132	-	-	-	-	-	-	-	-	1	1	3
Black, both sexes	32	-	-	-	-	-	1	-	4	1	1	3
Male	18	-	-	-	-	-	1	-	2	1	-	2
Female	14	-	-	-	-	-	-	-	2	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Minnesota												
All races, both sexes	129	-	-	-	-	1	2	1	4	1	4	7
Male	61	-	-	-	-	-	1	1	2	1	4	4
Female	68	-	-	-	-	1	1	-	2	-	-	3
White, both sexes	121	-	-	-	-	1	2	-	3	-	2	6
Male	56	-	-	-	-	-	1	-	2	-	2	3
Female	65	-	-	-	-	1	1	-	1	-	-	3
Black, both sexes	5	-	-	-	-	-	-	1	-	1	2	1
Male	5	-	-	-	-	-	-	1	-	1	2	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	277	-	1	-	1	1	1	1	2	2	5	9
Male	130	-	1	-	1	-	1	1	-	2	3	8
Female	147	-	-	-	-	1	-	-	2	-	2	1
White, both sexes	180	-	-	-	-	-	-	-	1	2	2	4
Male	89	-	-	-	-	-	-	-	-	2	2	3
Female	91	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	97	-	1	-	1	1	1	1	1	-	3	5
Male	41	-	1	-	1	-	1	1	-	-	1	5
Female	56	-	-	-	-	1	-	-	1	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Missouri												
All races, both sexes	260	-	-	1	-	-	2	5	3	3	5	5
Male	132	-	-	-	-	-	2	4	3	2	2	4
Female	128	-	-	1	-	-	-	1	-	1	3	1
White, both sexes	243	-	-	1	-	-	1	5	3	3	5	5
Male	125	-	-	-	-	-	1	4	3	2	2	4
Female	118	-	-	1	-	-	-	1	-	1	3	1
Black, both sexes	17	-	-	-	-	-	1	-	-	-	-	-
Male	7	-	-	-	-	-	1	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	35	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	34	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Nebraska												
All races, both sexes	81	-	-	-	-	-	-	-	-	1	4	3
Male	40	-	-	-	-	-	-	-	-	1	3	1
Female	41	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	75	-	-	-	-	-	-	-	-	1	3	1
Male	37	-	-	-	-	-	-	-	-	1	2	-
Female	38	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	1	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	142	-	1	-	-	-	2	-	2	4	4	10
Male	71	-	-	-	-	-	2	-	2	3	2	7
Female	71	-	1	-	-	-	-	-	-	1	2	3
White, both sexes	125	-	-	-	-	-	2	-	2	3	4	8
Male	66	-	-	-	-	-	2	-	2	2	2	6
Female	59	-	-	-	-	-	-	-	-	1	2	2
Black, both sexes	12	-	-	-	-	-	-	-	-	1	-	2
Male	3	-	-	-	-	-	-	-	-	1	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
New Hampshire												
All races, both sexes	45	-	-	-	-	-	1	-	-	-	-	1
Male	25	-	-	-	-	-	-	-	-	-	-	1
Female	20	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	44	-	-	-	-	-	1	-	-	-	-	1
Male	25	-	-	-	-	-	-	-	-	-	-	1
Female	19	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	273	-	-	-	-	-	2	1	-	2	10	4
Male	133	-	-	-	-	-	1	1	-	2	5	2
Female	140	-	-	-	-	-	1	-	-	-	5	2
White, both sexes	240	-	-	-	-	-	1	1	-	-	6	3
Male	116	-	-	-	-	-	1	1	-	-	4	2
Female	124	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	28	-	-	-	-	-	1	-	-	2	3	1
Male	13	-	-	-	-	-	-	-	-	2	1	-
Female	15	-	-	-	-	-	1	-	-	-	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
New Mexico												
All races, both sexes	44	-	-	-	-	-	-	-	-	1	-	-
Male	21	-	-	-	-	-	-	-	-	1	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	42	-	-	-	-	-	-	-	-	1	-	-
Male	19	-	-	-	-	-	-	-	-	1	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	485	3	1	-	-	6	4	3	8	9	16	20
Male	251	3	1	-	-	3	4	3	7	8	12	13
Female	234	-	-	-	-	3	-	-	1	1	4	7
White, both sexes	414	1	-	-	-	3	3	1	6	5	10	15
Male	209	1	-	-	-	2	3	1	6	4	9	10
Female	205	-	-	-	-	1	-	-	-	1	1	5
Black, both sexes	64	2	1	-	-	3	1	2	2	3	6	5
Male	40	2	1	-	-	1	1	2	1	3	3	3
Female	24	-	-	-	-	2	-	-	1	-	3	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
North Carolina												
All races, both sexes	382	4	-	-	1	-	5	3	3	5	6	4
Male	198	4	-	-	1	-	1	2	2	5	3	4
Female	184	-	-	-	-	-	4	1	1	-	3	-
White, both sexes	279	2	-	-	-	-	2	1	2	3	4	3
Male	147	2	-	-	-	-	1	-	1	3	1	3
Female	132	-	-	-	-	-	1	1	1	-	3	-
Black, both sexes	98	2	-	-	1	-	3	1	1	2	2	1
Male	50	2	-	-	1	-	-	1	1	2	2	1
Female	48	-	-	-	-	-	3	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	1	2
Male	11	-	-	-	-	-	-	-	-	-	1	2
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	18	-	-	-	-	-	-	-	-	-	1	1
Male	9	-	-	-	-	-	-	-	-	-	1	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Ohio												
All races, both sexes	306	-	1	-	1	-	3	3	9	6	7	12
Male	153	-	-	-	1	-	3	3	6	6	6	10
Female	153	-	1	-	-	-	-	-	3	-	1	2
White, both sexes	279	-	1	-	-	-	2	3	8	6	6	11
Male	141	-	-	-	-	-	2	3	6	6	5	9
Female	138	-	1	-	-	-	-	-	2	-	1	2
Black, both sexes	26	-	-	-	1	-	1	-	1	-	1	1
Male	11	-	-	-	1	-	1	-	-	-	1	1
Female	15	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	100	-	-	-	-	-	-	-	3	2	3	4
Male	44	-	-	-	-	-	-	-	3	1	3	2
Female	56	-	-	-	-	-	-	-	-	1	-	2
White, both sexes	83	-	-	-	-	-	-	-	1	1	2	2
Male	35	-	-	-	-	-	-	-	1	-	2	2
Female	48	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	2	1	-	1
Male	5	-	-	-	-	-	-	-	2	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Oregon												
All races, both sexes	92	-	-	-	-	-	-	2	-	1	2	7
Male	48	-	-	-	-	-	-	2	-	1	2	5
Female	44	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	91	-	-	-	-	-	-	2	-	1	2	7
Male	48	-	-	-	-	-	-	2	-	1	2	5
Female	43	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	491	-	-	-	-	-	5	5	2	5	11	15
Male	248	-	-	-	-	-	5	2	2	4	9	10
Female	243	-	-	-	-	-	-	3	-	1	2	5
White, both sexes	445	-	-	-	-	-	3	4	1	5	8	14
Male	222	-	-	-	-	-	3	1	1	4	7	9
Female	223	-	-	-	-	-	-	3	-	1	1	5
Black, both sexes	44	-	-	-	-	-	1	1	1	-	3	1
Male	25	-	-	-	-	-	1	1	1	-	2	1
Female	19	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Rhode Island												
All races, both sexes	14	-	-	-	-	-	-	-	1	-	-	-
Male	10	-	-	-	-	-	-	-	1	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	1	-	-	-
Male	10	-	-	-	-	-	-	-	1	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	330	-	-	-	-	-	3	7	12	12	20	21
Male	197	-	-	-	-	-	1	6	4	11	13	14
Female	133	-	-	-	-	-	2	1	8	1	7	7
White, both sexes	226	-	-	-	-	-	2	3	7	5	12	11
Male	134	-	-	-	-	-	-	2	2	5	9	8
Female	92	-	-	-	-	-	2	1	5	-	3	3
Black, both sexes	103	-	-	-	-	-	1	4	5	7	8	10
Male	63	-	-	-	-	-	1	4	2	6	4	6
Female	40	-	-	-	-	-	-	-	3	1	4	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
South Dakota												
All races, both sexes	40	-	-	-	-	-	-	1	-	1	1	1
Male	19	-	-	-	-	-	-	1	-	1	1	1
Female	21	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	34	-	-	-	-	-	-	-	-	1	1	-
Male	16	-	-	-	-	-	-	-	-	1	1	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	155	-	-	-	1	2	2	4	2	6	6	7
Male	75	-	-	-	-	1	1	4	1	4	3	6
Female	80	-	-	-	1	1	1	-	1	2	3	1
White, both sexes	132	-	-	-	1	1	1	4	2	4	4	5
Male	66	-	-	-	-	1	1	4	1	2	2	4
Female	66	-	-	-	1	-	-	-	1	2	2	1
Black, both sexes	23	-	-	-	-	1	1	-	-	2	2	2
Male	9	-	-	-	-	-	-	-	-	2	1	2
Female	14	-	-	-	-	1	1	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Texas												
All races, both sexes	999	4	1	-	2	2	10	12	7	19	18	31
Male	474	1	1	-	2	2	10	12	5	13	9	23
Female	525	3	-	-	-	-	-	-	2	6	9	8
White, both sexes	867	4	1	-	2	2	2	7	7	14	14	26
Male	408	1	1	-	2	2	2	7	5	10	7	20
Female	459	3	-	-	-	-	-	-	2	4	7	6
Black, both sexes	123	-	-	-	-	-	8	5	-	5	3	4
Male	63	-	-	-	-	-	8	5	-	3	1	2
Female	60	-	-	-	-	-	-	-	-	2	2	2
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	164	2	-	-	-	-	-	-	-	1	2	3
Male	85	2	-	-	-	-	-	-	-	1	2	1
Female	79	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	158	2	-	-	-	-	-	-	-	-	2	3
Male	83	2	-	-	-	-	-	-	-	-	2	1
Female	75	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Vermont												
All races, both sexes	9	-	-	-	-	-	-	-	2	-	-	-
Male	5	-	-	-	-	-	-	-	2	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	2	-	-	-
Male	5	-	-	-	-	-	-	-	2	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	205	1	-	-	-	-	2	2	2	3	3	7
Male	88	1	-	-	-	-	2	1	2	1	3	4
Female	117	-	-	-	-	-	-	1	-	2	-	3
White, both sexes	169	-	-	-	-	-	2	2	2	3	2	6
Male	75	-	-	-	-	-	2	1	2	1	2	3
Female	94	-	-	-	-	-	-	1	-	2	-	3
Black, both sexes	33	1	-	-	-	-	-	-	-	-	1	1
Male	12	1	-	-	-	-	-	-	-	-	1	1
Female	21	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Washington												
All races, both sexes	143	-	-	-	-	-	1	4	-	1	3	4
Male	77	-	-	-	-	-	1	4	-	-	3	1
Female	66	-	-	-	-	-	-	-	-	1	-	3
White, both sexes	137	-	-	-	-	-	1	4	-	1	3	4
Male	75	-	-	-	-	-	1	4	-	-	3	1
Female	62	-	-	-	-	-	-	-	-	1	-	3
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	72	-	-	-	-	-	-	-	-	-	-	2
Male	31	-	-	-	-	-	-	-	-	-	-	1
Female	41	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	71	-	-	-	-	-	-	-	-	-	-	2
Male	31	-	-	-	-	-	-	-	-	-	-	1
Female	40	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Wisconsin												
All races, both sexes	122	-	-	-	-	-	-	-	2	1	3	5
Male	54	-	-	-	-	-	-	-	1	1	3	4
Female	68	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	115	-	-	-	-	-	-	-	2	-	2	5
Male	50	-	-	-	-	-	-	-	1	-	2	4
Female	65	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	6	-	-	-	-	-	-	-	-	1	1	-
Male	3	-	-	-	-	-	-	-	-	1	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	31	-	-	-	-	-	-	-	-	1	1	1
Male	23	-	-	-	-	-	-	-	-	1	1	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	30	-	-	-	-	-	-	-	-	1	1	1
Male	23	-	-	-	-	-	-	-	-	1	1	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
United States												
All races, both sexes	505	609	643	696	875	1,229	1,722	1,793	1,433	648	109	-
Male	325	419	426	432	490	636	803	711	450	155	20	-
Female	180	190	217	264	385	593	919	1,082	983	493	89	-
White, both sexes	387	479	529	563	710	1,089	1,546	1,641	1,315	582	86	-
Male	251	333	356	358	418	576	733	654	425	138	16	-
Female	136	146	173	205	292	513	813	987	890	444	70	-
Black, both sexes	108	119	103	119	146	115	142	118	97	56	22	-
Male	66	80	65	69	60	45	55	45	19	13	4	-
Female	42	39	38	50	86	70	87	73	78	43	18	-
American Indian 1/, both sexes	3	4	4	5	4	6	8	7	4	-	-	-
Male	2	2	2	2	4	4	3	2	1	-	-	-
Female	1	2	2	3	-	2	5	5	3	-	-	-
Asian or Pacific Islander 2/, both sexes	7	7	7	9	15	19	26	27	17	10	1	-
Male	6	4	3	3	8	11	12	10	5	4	-	-
Female	1	3	4	6	7	8	14	17	12	6	1	-
Alabama												
All races, both sexes	15	38	33	39	42	51	71	56	52	11	5	-
Male	10	27	22	22	27	26	39	19	12	1	1	-
Female	5	11	11	17	15	25	32	37	40	10	4	-
White, both sexes	11	24	22	34	35	44	63	52	45	9	4	-
Male	7	19	18	21	24	25	35	19	11	1	1	-
Female	4	5	4	13	11	19	28	33	34	8	3	-
Black, both sexes	3	14	11	5	6	7	8	4	7	2	1	-
Male	2	8	4	1	2	1	4	-	1	-	-	-
Female	1	6	7	4	4	6	4	4	6	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Alaska												
All races, both sexes	2	3	-	3	-	3	2	2	1	-	-	-
Male	1	2	-	1	-	1	1	1	-	-	-	-
Female	1	1	-	2	-	2	1	1	1	-	-	-
White, both sexes	1	3	-	1	-	3	2	2	1	-	-	-
Male	1	2	-	-	-	1	1	1	-	-	-	-
Female	-	1	-	1	-	2	1	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	2	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	9	13	25	16	17	34	28	34	27	9	1	-
Male	8	10	14	10	12	20	11	18	11	-	-	-
Female	1	3	11	6	5	14	17	16	16	9	1	-
White, both sexes	9	12	22	13	16	33	28	32	26	8	1	-
Male	8	10	13	8	12	20	11	17	11	-	-	-
Female	1	2	9	5	4	13	17	15	15	8	1	-
Black, both sexes	-	-	1	3	1	1	-	1	-	1	-	-
Male	-	-	-	2	-	-	-	-	-	-	-	-
Female	-	-	1	1	1	1	-	1	-	1	-	-
American Indian 1/, both sexes	-	1	2	-	-	-	-	1	1	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	1	1	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Arkansas												
All races, both sexes	7	4	12	10	13	21	34	36	22	12	1	-
Male	4	3	4	7	9	14	11	12	-	1	-	-
Female	3	1	8	3	4	7	23	24	22	11	1	-
White, both sexes	5	2	12	8	12	21	32	31	21	12	1	-
Male	4	1	4	5	9	14	11	10	-	1	-	-
Female	1	1	8	3	3	7	21	21	21	11	1	-
Black, both sexes	2	2	-	2	1	-	2	5	1	-	-	-
Male	-	2	-	2	-	-	-	2	-	-	-	-
Female	2	-	-	-	1	-	2	3	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	48	55	38	39	63	93	164	194	163	84	16	-
Male	25	35	26	25	29	57	72	74	49	22	4	-
Female	23	20	12	14	34	36	92	120	114	62	12	-
White, both sexes	39	43	29	32	46	81	146	175	151	76	15	-
Male	23	27	19	21	21	48	65	67	46	20	4	-
Female	16	16	10	11	25	33	81	108	105	56	11	-
Black, both sexes	8	7	6	5	10	4	8	6	5	3	-	-
Male	2	5	5	4	3	3	1	2	-	-	-	-
Female	6	2	1	1	7	1	7	4	5	3	-	-
American Indian 1/, both sexes	-	2	-	-	2	-	1	-	-	-	-	-
Male	-	1	-	-	2	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	3	3	2	5	8	9	13	7	5	1	-
Male	-	2	2	-	3	6	5	5	3	2	-	-
Female	1	1	1	2	2	2	4	8	4	3	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Colorado												
All races, both sexes	6	6	10	9	8	6	21	19	6	5	1	-
Male	4	4	7	7	4	4	12	12	1	-	-	-
Female	2	2	3	2	4	2	9	7	5	5	1	-
White, both sexes	5	6	9	9	8	6	20	19	6	5	1	-
Male	3	4	6	7	4	4	11	12	1	-	-	-
Female	2	2	3	2	4	2	9	7	5	5	1	-
Black, both sexes	1	-	1	-	-	-	1	-	-	-	-	-
Male	1	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	7	6	2	4	10	8	17	19	16	12	1	-
Male	6	5	2	3	8	3	5	6	6	1	1	-
Female	1	1	-	1	2	5	12	13	10	11	-	-
White, both sexes	6	5	2	4	9	8	17	19	16	12	1	-
Male	6	5	2	3	7	3	5	6	6	1	1	-
Female	-	-	-	1	2	5	12	13	10	11	-	-
Black, both sexes	1	1	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Delaware												
All races, both sexes	2	-	-	3	2	6	4	4	1	-	-	-
Male	1	-	-	1	2	3	1	2	1	-	-	-
Female	1	-	-	2	-	3	3	2	-	-	-	-
White, both sexes	2	-	-	2	2	6	4	4	1	-	-	-
Male	1	-	-	-	2	3	1	2	1	-	-	-
Female	1	-	-	2	-	3	3	2	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	1	1	1	-	3	2	-	4	-	-
Male	1	-	1	-	-	-	2	2	-	1	-	-
Female	-	-	-	1	1	-	1	-	-	3	-	-
White, both sexes	-	-	-	-	-	-	-	2	-	1	-	-
Male	-	-	-	-	-	-	-	2	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	1	1	1	-	3	-	-	3	-	-
Male	1	-	1	-	-	-	2	-	-	-	-	-
Female	-	-	-	1	1	-	1	-	-	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Florida												
All races, both sexes	10	29	27	34	50	85	147	162	132	77	11	-
Male	7	18	19	21	35	49	78	70	56	23	6	-
Female	3	11	8	13	15	36	69	92	76	54	5	-
White, both sexes	8	18	24	29	41	78	133	156	124	73	9	-
Male	6	12	18	18	30	45	70	68	54	21	4	-
Female	2	6	6	11	11	33	63	88	70	52	5	-
Black, both sexes	2	11	2	5	9	7	11	5	7	3	2	-
Male	1	6	1	3	5	4	6	2	1	1	2	-
Female	1	5	1	2	4	3	5	3	6	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	3	1	1	1	-	-
Male	-	-	-	-	-	-	2	-	1	1	-	-
Female	-	-	1	-	-	-	1	1	-	-	-	-
Georgia												
All races, both sexes	70	68	87	80	83	99	110	105	82	29	8	-
Male	47	45	63	50	48	42	54	46	31	6	-	-
Female	23	23	24	30	35	57	56	59	51	23	8	-
White, both sexes	44	54	63	60	59	79	91	85	68	17	2	-
Male	33	35	45	39	38	35	46	36	26	2	-	-
Female	11	19	18	21	21	44	45	49	42	15	2	-
Black, both sexes	25	14	24	20	23	19	18	20	13	12	6	-
Male	13	10	18	11	10	6	8	10	5	4	-	-
Female	12	4	6	9	13	13	10	10	8	8	6	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	1	1	-	1	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Hawaii												
All races, both sexes	14	5	2	6	2	4	7	5	6	4	-	-
Male	12	4	1	2	1	2	2	3	1	1	-	-
Female	2	1	1	4	1	2	5	2	5	3	-	-
White, both sexes	10	4	-	2	-	-	2	1	2	1	-	-
Male	8	3	-	1	-	-	1	1	1	1	-	-
Female	2	1	-	1	-	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	1	2	4	1	4	5	4	4	3	-	-
Male	4	1	1	1	1	2	1	2	-	-	-	-
Female	-	-	1	3	-	2	4	2	4	3	-	-
Idaho												
All races, both sexes	2	3	2	1	3	12	4	7	3	2	-	-
Male	1	2	2	1	1	8	2	5	1	1	-	-
Female	1	1	-	-	2	4	2	2	2	1	-	-
White, both sexes	2	3	2	1	3	11	4	7	3	2	-	-
Male	1	2	2	1	1	7	2	5	1	1	-	-
Female	1	1	-	-	2	4	2	2	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Illinois												
All races, both sexes	21	22	26	32	34	52	54	82	49	28	6	-
Male	13	17	17	21	18	23	23	35	13	6	-	-
Female	8	5	9	11	16	29	31	47	36	22	6	-
White, both sexes	16	20	21	22	28	48	48	78	43	25	4	-
Male	8	15	13	14	15	22	20	32	12	5	-	-
Female	8	5	8	8	13	26	28	46	31	20	4	-
Black, both sexes	5	2	5	10	5	2	6	3	6	3	2	-
Male	5	2	4	7	2	-	3	2	1	1	-	-
Female	-	-	1	3	3	2	3	1	5	2	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	2	-	1	-	-	-	-
Male	-	-	-	-	1	1	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Indiana												
All races, both sexes	9	11	11	11	11	17	32	29	28	12	1	-
Male	5	8	6	6	4	9	19	14	6	4	1	-
Female	4	3	5	5	7	8	13	15	22	8	-	-
White, both sexes	5	10	11	8	11	16	32	29	28	12	1	-
Male	2	7	6	3	4	9	19	14	6	4	1	-
Female	3	3	5	5	7	7	13	15	22	8	-	-
Black, both sexes	4	1	-	3	-	1	-	-	-	-	-	-
Male	3	1	-	3	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Iowa												
All races, both sexes	2	3	2	2	3	2	6	10	13	5	1	-
Male	1	1	2	2	1	2	1	4	6	2	-	-
Female	1	2	-	-	2	-	5	6	7	3	1	-
White, both sexes	2	3	2	2	3	2	6	10	13	5	1	-
Male	1	1	2	2	1	2	1	4	6	2	-	-
Female	1	2	-	-	2	-	5	6	7	3	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	4	4	8	3	9	7	10	8	7	5	3	-
Male	3	2	5	2	5	4	3	-	2	1	-	-
Female	1	2	3	1	4	3	7	8	5	4	3	-
White, both sexes	3	4	7	3	8	6	10	7	7	5	3	-
Male	2	2	5	2	4	3	3	-	2	1	-	-
Female	1	2	2	1	4	3	7	7	5	4	3	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	1	1	-	1	-	-	-	-
Male	1	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Kentucky												
All races, both sexes	17	30	13	24	19	40	33	18	19	9	1	-
Male	12	25	9	16	10	17	12	4	4	1	-	-
Female	5	5	4	8	9	23	21	14	15	8	1	-
White, both sexes	17	27	13	23	19	37	31	17	17	9	1	-
Male	12	23	9	15	10	15	12	4	4	1	-	-
Female	5	4	4	8	9	22	19	13	13	8	1	-
Black, both sexes	-	3	-	1	-	3	2	1	2	-	-	-
Male	-	2	-	1	-	2	-	-	-	-	-	-
Female	-	1	-	-	-	1	2	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	11	10	11	18	23	26	63	59	45	16	2	-
Male	7	8	8	11	13	11	24	22	8	5	-	-
Female	4	2	3	7	10	15	39	37	37	11	2	-
White, both sexes	3	7	9	12	16	20	53	53	37	14	2	-
Male	3	7	7	8	8	7	22	19	8	5	-	-
Female	-	-	2	4	8	13	31	34	29	9	2	-
Black, both sexes	8	3	2	6	7	6	9	5	8	2	-	-
Male	4	1	1	3	5	4	1	2	-	-	-	-
Female	4	2	1	3	2	2	8	3	8	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Maine												
All races, both sexes	6	4	5	5	7	9	6	4	13	2	2	-
Male	5	3	4	5	5	5	5	3	7	1	-	-
Female	1	1	1	-	2	4	1	1	6	1	2	-
White, both sexes	6	4	5	5	7	9	5	4	13	2	2	-
Male	5	3	4	5	5	5	4	3	7	1	-	-
Female	1	1	1	-	2	4	1	1	6	1	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	4	3	6	6	5	4	12	11	9	6	1	-
Male	3	1	5	3	3	2	6	4	3	2	1	-
Female	1	2	1	3	2	2	6	7	6	4	-	-
White, both sexes	1	3	4	3	1	4	9	9	6	4	-	-
Male	1	1	3	2	-	2	4	3	2	2	-	-
Female	-	2	1	1	1	2	5	6	4	2	-	-
Black, both sexes	3	-	2	3	4	-	3	1	3	2	1	-
Male	2	-	2	1	3	-	2	-	1	-	1	-
Female	1	-	-	2	1	-	1	1	2	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Massachusetts												
All races, both sexes	8	4	6	12	20	13	29	35	30	15	2	-
Male	4	3	2	11	12	6	10	11	12	3	2	-
Female	4	1	4	1	8	7	19	24	18	12	-	-
White, both sexes	6	3	4	11	20	13	28	34	29	15	2	-
Male	2	2	1	10	12	6	9	11	11	3	2	-
Female	4	1	3	1	8	7	19	23	18	12	-	-
Black, both sexes	2	-	2	1	-	-	1	1	1	-	-	-
Male	2	-	1	1	-	-	1	-	1	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	17	12	14	19	23	32	37	37	31	8	4	-
Male	9	10	10	10	8	15	13	14	6	2	-	-
Female	8	2	4	9	15	17	24	23	25	6	4	-
White, both sexes	14	12	12	16	20	27	33	36	31	7	4	-
Male	7	10	9	7	8	11	11	14	6	2	-	-
Female	7	2	3	9	12	16	22	22	25	5	4	-
Black, both sexes	3	-	2	3	3	5	4	1	-	1	-	-
Male	2	-	1	3	-	4	2	-	-	-	-	-
Female	1	-	1	-	3	1	2	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Minnesota												
All races, both sexes	5	5	6	4	5	11	22	23	16	8	4	-
Male	5	3	3	2	2	8	10	7	5	3	-	-
Female	-	2	3	2	3	3	12	16	11	5	4	-
White, both sexes	5	5	6	4	5	11	21	23	15	8	4	-
Male	5	3	3	2	2	8	10	7	5	3	-	-
Female	-	2	3	2	3	3	11	16	10	5	4	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	5	15	13	18	34	29	47	41	34	16	2	-
Male	3	9	7	10	16	15	26	17	8	2	-	-
Female	2	6	6	8	18	14	21	24	26	14	2	-
White, both sexes	2	7	7	10	23	22	34	32	24	10	-	-
Male	2	4	4	6	13	12	22	14	5	-	-	-
Female	-	3	3	4	10	10	12	18	19	10	-	-
Black, both sexes	3	8	6	8	11	7	13	9	10	6	2	-
Male	1	5	3	4	3	3	4	3	3	2	-	-
Female	2	3	3	4	8	4	9	6	7	4	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Missouri												
All races, both sexes	8	8	18	12	16	36	47	37	31	21	2	-
Male	4	6	9	6	9	20	25	12	14	9	1	-
Female	4	2	9	6	7	16	22	25	17	12	1	-
White, both sexes	8	8	17	11	15	33	44	34	30	18	2	-
Male	4	6	8	6	9	20	22	11	14	8	1	-
Female	4	2	9	5	6	13	22	23	16	10	1	-
Black, both sexes	-	-	1	1	1	3	3	3	1	3	-	-
Male	-	-	1	-	-	-	3	1	-	1	-	-
Female	-	-	-	1	1	3	-	2	1	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	3	1	2	2	5	1	7	10	4	-	-
Male	-	1	1	1	2	2	-	6	3	-	-	-
Female	-	2	-	1	-	3	1	1	7	4	-	-
White, both sexes	-	3	1	2	2	4	1	7	10	4	-	-
Male	-	1	1	1	2	2	-	6	3	-	-	-
Female	-	2	-	1	-	2	1	1	7	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Nebraska												
All races, both sexes	5	-	3	8	7	6	12	13	12	5	2	-
Male	3	-	3	7	4	4	6	5	3	-	-	-
Female	2	-	-	1	3	2	6	8	9	5	2	-
White, both sexes	5	-	3	8	6	6	12	12	11	5	2	-
Male	3	-	3	7	3	4	6	5	3	-	-	-
Female	2	-	-	1	3	2	6	7	8	5	2	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	3	6	5	10	8	21	36	17	12	1	-	-
Male	1	6	3	7	4	10	16	4	4	-	-	-
Female	2	-	2	3	4	11	20	13	8	1	-	-
White, both sexes	3	5	4	9	5	17	35	15	12	1	-	-
Male	1	5	3	7	3	9	16	4	4	-	-	-
Female	2	-	1	2	2	8	19	11	8	1	-	-
Black, both sexes	-	-	1	-	2	4	-	2	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	2	3	-	2	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
New Hampshire												
All races, both sexes	2	1	1	2	3	5	8	11	5	4	1	-
Male	2	-	1	2	1	3	4	6	3	1	1	-
Female	-	1	-	-	2	2	4	5	2	3	-	-
White, both sexes	2	1	1	2	3	4	8	11	5	4	1	-
Male	2	-	1	2	1	3	4	6	3	1	1	-
Female	-	1	-	-	2	1	4	5	2	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
New Jersey												
All races, both sexes	9	10	10	15	19	23	50	49	47	21	1	-
Male	6	8	7	9	9	14	24	23	19	3	-	-
Female	3	2	3	6	10	9	26	26	28	18	1	-
White, both sexes	8	8	9	12	15	17	48	45	46	20	1	-
Male	5	6	6	8	7	12	22	21	19	2	-	-
Female	3	2	3	4	8	5	26	24	27	18	1	-
Black, both sexes	1	2	1	2	4	5	2	3	1	-	-	-
Male	1	2	1	-	2	1	2	1	-	-	-	-
Female	-	-	-	2	2	4	-	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	-	1	-	1	-	-
Male	-	-	-	1	-	1	-	1	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
New Mexico												
All races, both sexes	2	1	2	3	2	7	12	5	8	1	-	-
Male	2	-	1	3	1	3	7	-	2	1	-	-
Female	-	1	1	-	1	4	5	5	6	-	-	-
White, both sexes	2	1	2	3	1	6	12	5	8	1	-	-
Male	2	-	1	3	-	2	7	-	2	1	-	-
Female	-	1	1	-	1	4	5	5	6	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	24	27	22	25	27	54	64	81	63	24	4	-
Male	18	16	13	13	14	30	39	24	19	10	1	-
Female	6	11	9	12	13	24	25	57	44	14	3	-
White, both sexes	19	18	19	23	22	48	60	74	59	24	4	-
Male	13	12	10	13	11	27	37	22	17	10	1	-
Female	6	6	9	10	11	21	23	52	42	14	3	-
Black, both sexes	5	8	3	2	5	5	3	5	3	-	-	-
Male	5	4	3	-	3	3	1	2	2	-	-	-
Female	-	4	-	2	2	2	2	3	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	1	2	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	1	-	2	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
North Carolina												
All races, both sexes	15	25	32	28	44	40	55	62	38	9	3	-
Male	11	16	24	13	25	17	23	28	15	3	1	-
Female	4	9	8	15	19	23	32	34	23	6	2	-
White, both sexes	12	15	23	16	29	35	40	51	32	9	-	-
Male	9	10	17	7	18	15	18	25	14	3	-	-
Female	3	5	6	9	11	20	22	26	18	6	-	-
Black, both sexes	3	10	9	11	15	5	13	11	5	-	3	-
Male	2	6	7	6	7	2	5	3	1	-	1	-
Female	1	4	2	5	8	3	8	8	4	-	2	-
American Indian 1/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
North Dakota												
All races, both sexes	-	1	2	-	1	2	-	6	5	-	-	-
Male	-	1	1	-	1	2	-	2	1	-	-	-
Female	-	-	1	-	-	-	-	4	4	-	-	-
White, both sexes	-	1	2	-	1	1	-	6	5	-	-	-
Male	-	1	1	-	1	1	-	2	1	-	-	-
Female	-	-	1	-	-	-	-	4	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Ohio												
All races, both sexes	13	15	15	11	26	30	44	51	42	16	1	-
Male	8	12	9	5	11	12	22	19	16	4	-	-
Female	5	3	6	6	15	18	22	32	26	12	1	-
White, both sexes	11	13	15	8	22	25	43	50	38	16	1	-
Male	6	11	9	4	11	9	21	19	16	4	-	-
Female	5	2	6	4	11	16	22	31	22	12	1	-
Black, both sexes	1	2	-	3	4	5	1	1	4	-	-	-
Male	1	1	-	1	-	3	1	-	-	-	-	-
Female	-	1	-	2	4	2	-	1	4	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	5	5	9	7	10	10	9	15	11	7	-	-
Male	2	4	5	3	4	4	3	4	4	2	-	-
Female	3	1	4	4	6	6	6	11	7	5	-	-
White, both sexes	3	5	8	6	9	9	7	13	11	6	-	-
Male	1	4	4	3	3	4	3	3	4	1	-	-
Female	2	1	4	3	6	5	4	10	7	5	-	-
Black, both sexes	2	-	-	-	-	1	1	1	-	1	-	-
Male	1	-	-	-	-	-	-	1	-	1	-	-
Female	1	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	1	1	-	1	1	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Oregon												
All races, both sexes	5	3	3	10	3	11	7	18	12	8	-	-
Male	3	1	2	4	2	8	5	9	3	1	-	-
Female	2	2	1	6	1	3	2	9	9	7	-	-
White, both sexes	5	3	3	10	3	11	7	17	12	8	-	-
Male	3	1	2	4	2	8	5	9	3	1	-	-
Female	2	2	1	6	1	3	2	8	9	7	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Pennsylvania												
All races, both sexes	17	28	28	25	31	52	84	96	64	20	3	-
Male	9	22	21	13	21	27	36	37	25	5	-	-
Female	8	6	7	12	10	25	48	59	39	15	3	-
White, both sexes	14	23	26	21	28	47	79	90	62	19	1	-
Male	8	18	19	11	19	24	35	33	25	5	-	-
Female	6	5	7	10	9	23	44	57	37	14	1	-
Black, both sexes	3	5	2	4	2	5	5	6	2	1	2	-
Male	1	4	2	2	2	3	1	4	-	-	-	-
Female	2	1	-	2	-	2	4	2	2	1	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Rhode Island												
All races, both sexes	2	1	1	-	1	2	3	2	1	-	-	-
Male	2	1	1	-	1	1	2	1	-	-	-	-
Female	-	-	-	-	-	1	1	1	1	-	-	-
White, both sexes	2	1	1	-	1	2	3	2	1	-	-	-
Male	2	1	1	-	1	1	2	1	-	-	-	-
Female	-	-	-	-	-	1	1	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	24	41	30	22	31	27	27	27	13	11	2	-
Male	17	33	24	19	15	13	12	10	2	3	-	-
Female	7	8	6	3	16	14	15	17	11	8	2	-
White, both sexes	16	25	23	14	24	22	21	20	11	9	1	-
Male	9	20	20	12	12	11	10	9	2	3	-	-
Female	7	5	3	2	12	11	11	11	9	6	1	-
Black, both sexes	8	15	7	8	7	5	6	7	2	2	1	-
Male	8	13	4	7	3	2	2	1	-	-	-	-
Female	-	2	3	1	4	3	4	6	2	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
South Dakota												
All races, both sexes	3	1	2	2	4	3	9	4	4	4	-	-
Male	-	-	1	2	4	2	4	1	1	-	-	-
Female	3	1	1	-	-	1	5	3	3	4	-	-
White, both sexes	3	1	1	2	4	3	8	3	3	4	-	-
Male	-	-	1	2	4	2	4	1	-	-	-	-
Female	3	1	-	-	-	1	4	2	3	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	1	-	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	6	6	9	12	11	16	22	17	18	8	-	-
Male	2	4	4	8	6	8	14	6	2	1	-	-
Female	4	2	5	4	5	8	8	11	16	7	-	-
White, both sexes	6	6	7	9	10	14	20	15	17	6	-	-
Male	2	4	4	7	6	7	14	4	2	1	-	-
Female	4	2	3	2	4	7	6	11	15	5	-	-
Black, both sexes	-	-	2	3	1	2	2	2	1	2	-	-
Male	-	-	-	1	-	1	-	2	-	-	-	-
Female	-	-	2	2	1	1	2	-	1	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Texas												
All races, both sexes	31	41	54	55	80	130	136	149	142	64	11	-
Male	22	23	31	36	53	65	56	58	40	12	-	-
Female	9	18	23	19	27	65	80	91	102	52	11	-
White, both sexes	21	34	43	47	65	119	121	140	131	57	10	-
Male	16	18	26	31	47	64	51	52	37	9	-	-
Female	5	16	17	16	18	55	70	88	94	48	10	-
Black, both sexes	10	7	11	8	13	9	15	9	8	7	1	-
Male	6	5	5	5	6	1	5	6	2	3	-	-
Female	4	2	6	3	7	8	10	3	6	4	1	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	1	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	2	1	-	-	1	-	-	-
Utah												
All races, both sexes	5	3	5	15	14	29	46	19	14	6	-	-
Male	3	-	2	11	5	12	27	11	7	1	-	-
Female	2	3	3	4	9	17	19	8	7	5	-	-
White, both sexes	5	3	5	15	13	29	43	18	14	6	-	-
Male	3	-	2	11	5	12	26	11	7	1	-	-
Female	2	3	3	4	8	17	17	7	7	5	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	3	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	2	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Vermont												
All races, both sexes	-	1	-	-	2	-	2	1	1	-	-	-
Male	-	1	-	-	1	-	-	-	1	-	-	-
Female	-	-	-	-	1	-	2	1	-	-	-	-
White, both sexes	-	1	-	-	2	-	2	1	1	-	-	-
Male	-	1	-	-	1	-	-	-	1	-	-	-
Female	-	-	-	-	1	-	2	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	11	13	7	7	20	27	35	31	25	8	1	-
Male	4	8	5	5	12	13	11	9	6	1	-	-
Female	7	5	2	2	8	14	24	22	19	7	1	-
White, both sexes	8	10	7	7	14	23	32	25	20	6	-	-
Male	3	5	5	5	10	12	10	8	5	1	-	-
Female	5	5	2	2	4	11	22	17	15	5	-	-
Black, both sexes	3	3	-	-	6	4	1	5	5	2	1	-
Male	1	3	-	-	2	1	-	1	1	-	-	-
Female	2	-	-	-	4	3	1	4	4	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Washington												
All races, both sexes	3	3	11	10	7	8	16	35	20	16	1	-
Male	2	3	8	6	3	6	10	18	4	8	-	-
Female	1	-	3	4	4	2	6	17	16	8	1	-
White, both sexes	3	3	10	9	7	8	15	33	19	16	1	-
Male	2	3	8	5	3	6	9	18	4	8	-	-
Female	1	-	2	4	4	2	6	15	15	8	1	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-
West Virginia												
All races, both sexes	2	3	6	2	9	7	9	15	11	5	1	-
Male	1	3	5	1	5	4	4	4	3	-	-	-
Female	1	-	1	1	4	3	5	11	8	5	1	-
White, both sexes	2	3	6	2	9	7	9	14	11	5	1	-
Male	1	3	5	1	5	4	4	4	3	-	-	-
Female	1	-	1	1	4	3	5	10	8	5	1	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Wisconsin												
All races, both sexes	4	7	3	12	6	11	20	22	18	5	3	-
Male	3	3	3	8	2	7	6	8	4	1	-	-
Female	1	4	-	4	4	4	14	14	14	4	3	-
White, both sexes	3	6	3	11	5	11	19	22	18	5	3	-
Male	2	3	3	7	2	7	6	8	4	1	-	-
Female	1	3	-	4	3	4	13	14	14	4	3	-
Black, both sexes	1	1	-	-	1	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	4	3	4	2	4	3	5	1	1	1	-	-
Male	3	2	3	1	2	3	5	1	-	-	-	-
Female	1	1	1	1	2	-	-	-	1	1	-	-
White, both sexes	4	3	4	2	3	3	5	1	1	1	-	-
Male	3	2	3	1	2	3	5	1	-	-	-	-
Female	1	1	1	1	1	-	-	-	1	1	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
United States												
All races, both sexes	135,952	132	52	38	45	71	124	180	325	733	1,400	2,616
Male	54,111	77	20	17	20	40	73	90	162	401	742	1,410
Female	81,841	55	32	21	25	31	51	90	163	332	658	1,206
White, both sexes	114,695	94	34	28	35	52	86	129	203	470	886	1,682
Male	44,714	51	16	14	18	28	48	69	98	251	495	925
Female	69,981	43	18	14	17	24	38	60	105	219	391	757
Black, both sexes	17,085	34	16	6	7	16	30	43	105	216	439	795
Male	7,549	25	4	2	1	11	19	19	52	126	209	404
Female	9,536	9	12	4	6	5	11	24	53	90	230	391
American Indian 1/, both sexes	586	2	1	-	-	1	1	2	1	9	19	24
Male	267	-	-	-	-	-	1	1	1	5	9	15
Female	319	2	1	-	-	1	-	1	-	4	10	9
Asian or Pacific Islander 2/, both sexes	3,586	2	1	4	3	2	7	6	16	38	56	115
Male	1,581	1	-	1	1	1	5	1	11	19	29	66
Female	2,005	1	1	3	2	1	2	5	5	19	27	49
Alabama												
All races, both sexes	2,747	3	-	1	-	3	5	4	15	19	32	70
Male	1,155	2	-	-	-	3	3	3	4	11	14	31
Female	1,592	1	-	1	-	-	2	1	11	8	18	39
White, both sexes	1,985	2	-	1	-	1	4	3	8	8	11	37
Male	803	2	-	-	-	1	2	2	1	5	6	13
Female	1,182	-	-	1	-	-	2	1	7	3	5	24
Black, both sexes	752	1	-	-	-	2	1	1	6	11	20	33
Male	349	-	-	-	-	2	1	1	3	6	8	18
Female	403	1	-	-	-	-	-	-	3	5	12	15
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Alaska												
All races, both sexes	157	-	-	-	-	-	-	-	-	-	3	5
Male	79	-	-	-	-	-	-	-	-	-	-	1
Female	78	-	-	-	-	-	-	-	-	-	3	4
White, both sexes	116	-	-	-	-	-	-	-	-	-	1	2
Male	56	-	-	-	-	-	-	-	-	-	-	1
Female	60	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	26	-	-	-	-	-	-	-	-	-	2	2
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	2	2
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
Arizona												
All races, both sexes	2,207	2	-	1	1	-	3	2	5	13	26	42
Male	866	1	-	1	-	-	1	2	2	7	17	28
Female	1,341	1	-	-	1	-	2	-	3	6	9	14
White, both sexes	2,080	2	-	1	1	-	2	2	4	12	22	37
Male	809	1	-	1	-	-	1	2	1	7	14	23
Female	1,271	1	-	-	1	-	1	-	3	5	8	14
Black, both sexes	52	-	-	-	-	-	1	-	-	-	1	2
Male	24	-	-	-	-	-	-	-	-	-	1	2
Female	28	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	40	-	-	-	-	-	-	-	1	1	3	3
Male	19	-	-	-	-	-	-	-	1	-	2	3
Female	21	-	-	-	-	-	-	-	-	1	1	-
Asian or Pacific Islander 2/, both sexes	35	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Arkansas												
All races, both sexes	1,873	1	-	-	-	1	3	5	5	14	29	32
Male	746	1	-	-	-	-	-	2	2	9	18	17
Female	1,127	-	-	-	-	1	3	3	3	5	11	15
White, both sexes	1,587	1	-	-	-	1	3	5	3	11	20	22
Male	637	1	-	-	-	-	-	2	2	7	12	12
Female	950	-	-	-	-	1	3	3	1	4	8	10
Black, both sexes	274	-	-	-	-	-	-	-	2	3	7	10
Male	105	-	-	-	-	-	-	-	-	2	5	5
Female	169	-	-	-	-	-	-	-	2	1	2	5
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	2	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
California												
All races, both sexes	14,557	16	3	5	5	11	19	24	26	101	160	314
Male	6,029	7	1	2	2	4	15	9	18	66	96	200
Female	8,528	9	2	3	3	7	4	15	8	35	64	114
White, both sexes	11,602	13	1	4	4	7	10	17	18	73	117	205
Male	4,734	4	1	2	1	2	8	7	12	52	73	130
Female	6,868	9	-	2	3	5	2	10	6	21	44	75
Black, both sexes	1,202	2	2	-	-	1	4	4	4	8	18	51
Male	510	2	-	-	-	1	3	1	3	5	7	29
Female	692	-	2	-	-	-	1	3	1	3	11	22
American Indian 1/, both sexes	70	-	-	-	-	1	-	-	-	1	1	4
Male	31	-	-	-	-	-	-	-	-	1	1	3
Female	39	-	-	-	-	1	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1,683	1	-	1	1	2	5	3	4	19	24	54
Male	754	1	-	-	1	1	4	1	3	8	15	38
Female	929	-	-	1	-	1	1	2	1	11	9	16

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Colorado												
All races, both sexes	1,600	5	1	1	-	-	3	1	2	8	19	17
Male	613	3	-	1	-	-	1	-	1	4	8	5
Female	987	2	1	-	-	-	2	1	1	4	11	12
White, both sexes	1,507	4	1	1	-	-	2	1	2	8	15	14
Male	578	2	-	1	-	-	-	-	1	4	6	5
Female	929	2	1	-	-	-	2	1	1	4	9	9
Black, both sexes	62	1	-	-	-	-	1	-	-	-	4	2
Male	24	1	-	-	-	-	1	-	-	-	2	-
Female	38	-	-	-	-	-	-	-	-	-	2	2
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	27	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	1
Connecticut												
All races, both sexes	1,463	2	-	-	-	-	2	4	3	7	10	15
Male	560	1	-	-	-	-	1	2	1	4	5	11
Female	903	1	-	-	-	-	1	2	2	3	5	4
White, both sexes	1,347	2	-	-	-	-	2	4	3	5	8	12
Male	516	1	-	-	-	-	1	2	1	3	5	10
Female	831	1	-	-	-	-	1	2	2	2	3	2
Black, both sexes	99	-	-	-	-	-	-	-	-	2	2	3
Male	35	-	-	-	-	-	-	-	-	1	-	1
Female	64	-	-	-	-	-	-	-	-	1	2	2
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Delaware												
All races, both sexes	374	-	-	-	1	-	1	-	1	1	4	11
Male	142	-	-	-	1	-	1	-	1	-	1	9
Female	232	-	-	-	-	-	-	-	-	1	3	2
White, both sexes	293	-	-	-	1	-	1	-	-	1	3	4
Male	99	-	-	-	1	-	1	-	-	-	-	2
Female	194	-	-	-	-	-	-	-	-	1	3	2
Black, both sexes	75	-	-	-	-	-	-	-	1	-	-	7
Male	40	-	-	-	-	-	-	-	1	-	-	7
Female	35	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	220	-	-	-	-	-	-	-	-	1	1	13
Male	82	-	-	-	-	-	-	-	-	-	-	8
Female	138	-	-	-	-	-	-	-	-	1	1	5
White, both sexes	43	-	-	-	-	-	-	-	-	-	-	1
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	26	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	176	-	-	-	-	-	-	-	-	1	1	12
Male	64	-	-	-	-	-	-	-	-	-	-	8
Female	112	-	-	-	-	-	-	-	-	1	1	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Florida												
All races, both sexes	8,781	8	2	4	1	4	8	15	17	45	93	168
Male	3,715	4	1	2	-	4	6	7	8	26	52	85
Female	5,066	4	1	2	1	-	2	8	9	19	41	83
White, both sexes	7,510	5	2	4	1	3	5	9	8	27	64	113
Male	3,155	3	1	2	-	3	4	4	3	14	34	57
Female	4,355	2	1	2	1	-	1	5	5	13	30	56
Black, both sexes	1,193	3	-	-	-	1	3	6	9	18	29	50
Male	527	1	-	-	-	1	2	3	5	12	18	27
Female	666	2	-	-	-	-	1	3	4	6	11	23
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	67	-	-	-	-	-	-	-	-	-	-	4
Male	29	-	-	-	-	-	-	-	-	-	-	1
Female	38	-	-	-	-	-	-	-	-	-	-	3
Georgia												
All races, both sexes	3,894	4	2	1	5	4	5	3	13	31	61	123
Male	1,609	4	-	1	2	2	4	-	5	16	32	54
Female	2,285	-	2	-	3	2	1	3	8	15	29	69
White, both sexes	2,703	-	1	1	2	3	4	1	7	14	25	57
Male	1,069	-	-	1	1	2	3	-	3	4	17	26
Female	1,634	-	1	-	1	1	1	1	4	10	8	31
Black, both sexes	1,151	4	1	-	2	1	1	2	6	15	36	64
Male	527	4	-	-	1	-	1	-	2	11	15	27
Female	624	-	1	-	1	1	-	2	4	4	21	37
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	2
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	36	-	-	-	1	-	-	-	-	2	-	-
Male	12	-	-	-	-	-	-	-	-	1	-	-
Female	24	-	-	-	1	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Hawaii												
All races, both sexes	643	-	1	-	-	-	-	-	1	6	6	11
Male	295	-	-	-	-	-	-	-	1	2	4	5
Female	348	-	1	-	-	-	-	-	-	4	2	6
White, both sexes	144	-	-	-	-	-	-	-	-	1	1	1
Male	70	-	-	-	-	-	-	-	-	-	1	-
Female	74	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	8	-	1	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	490	-	-	-	-	-	-	-	1	5	5	10
Male	218	-	-	-	-	-	-	-	1	2	3	5
Female	272	-	-	-	-	-	-	-	-	3	2	5
Idaho												
All races, both sexes	640	2	-	-	-	1	1	-	3	3	2	13
Male	243	1	-	-	-	-	-	-	1	-	1	5
Female	397	1	-	-	-	1	1	-	2	3	1	8
White, both sexes	628	2	-	-	-	1	1	-	3	3	2	12
Male	239	1	-	-	-	-	-	-	1	-	1	4
Female	389	1	-	-	-	1	1	-	2	3	1	8
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Illinois												
All races, both sexes	5,864	5	1	1	1	3	4	7	15	24	51	111
Male	2,252	3	1	-	1	2	2	3	7	14	26	50
Female	3,612	2	-	1	-	1	2	4	8	10	25	61
White, both sexes	4,903	3	-	1	1	3	4	6	7	15	28	54
Male	1,836	2	-	-	1	2	2	3	4	8	13	27
Female	3,067	1	-	1	-	1	2	3	3	7	15	27
Black, both sexes	869	2	1	-	-	-	-	1	7	8	22	55
Male	376	1	1	-	-	-	-	-	3	5	13	22
Female	493	1	-	-	-	-	-	1	4	3	9	33
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	84	-	-	-	-	-	-	-	1	1	1	2
Male	37	-	-	-	-	-	-	-	-	1	-	1
Female	47	-	-	-	-	-	-	-	1	-	1	1
Indiana												
All races, both sexes	3,083	1	2	-	1	1	4	5	10	12	29	59
Male	1,177	-	1	-	1	1	1	2	5	8	14	29
Female	1,906	1	1	-	-	-	3	3	5	4	15	30
White, both sexes	2,834	1	2	-	1	1	3	5	8	10	23	46
Male	1,059	-	1	-	1	1	1	2	4	7	12	24
Female	1,775	1	1	-	-	-	2	3	4	3	11	22
Black, both sexes	244	-	-	-	-	-	1	-	2	2	6	13
Male	115	-	-	-	-	-	-	-	1	1	2	5
Female	129	-	-	-	-	-	1	-	1	1	4	8
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Iowa												
All races, both sexes	1,686	-	-	-	-	-	-	-	1	6	11	19
Male	641	-	-	-	-	-	-	-	-	3	5	13
Female	1,045	-	-	-	-	-	-	-	1	3	6	6
White, both sexes	1,641	-	-	-	-	-	-	-	1	5	9	15
Male	620	-	-	-	-	-	-	-	-	2	5	10
Female	1,021	-	-	-	-	-	-	-	1	3	4	5
Black, both sexes	36	-	-	-	-	-	-	-	-	1	2	2
Male	17	-	-	-	-	-	-	-	-	1	-	2
Female	19	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
Kansas												
All races, both sexes	1,498	-	1	-	1	-	1	1	2	6	16	24
Male	576	-	1	-	1	-	1	-	1	4	7	10
Female	922	-	-	-	-	-	-	1	1	2	9	14
White, both sexes	1,404	-	1	-	1	-	1	-	1	4	14	21
Male	531	-	1	-	1	-	1	-	1	2	6	9
Female	873	-	-	-	-	-	-	-	-	2	8	12
Black, both sexes	73	-	-	-	-	-	-	1	1	1	2	3
Male	38	-	-	-	-	-	-	-	-	1	1	1
Female	35	-	-	-	-	-	-	1	1	-	1	2
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Kentucky												
All races, both sexes	2,144	2	1	-	-	-	2	1	5	14	18	34
Male	799	2	-	-	-	-	2	-	1	5	8	17
Female	1,345	-	1	-	-	-	-	1	4	9	10	17
White, both sexes	1,971	1	-	-	-	-	-	1	4	12	15	28
Male	723	1	-	-	-	-	-	-	1	4	6	14
Female	1,248	-	-	-	-	-	-	1	3	8	9	14
Black, both sexes	165	1	1	-	-	-	2	-	1	2	3	6
Male	71	1	-	-	-	-	2	-	-	1	2	3
Female	94	-	1	-	-	-	-	-	1	1	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	2,147	5	2	1	2	2	4	6	5	21	36	67
Male	899	4	1	1	1	2	4	5	2	10	21	36
Female	1,248	1	1	-	1	-	-	1	3	11	15	31
White, both sexes	1,436	1	2	-	2	1	2	3	2	10	16	24
Male	572	1	1	-	1	1	2	2	-	5	9	14
Female	864	-	1	-	1	-	-	1	2	5	7	10
Black, both sexes	696	4	-	-	-	1	2	3	3	10	20	43
Male	321	3	-	-	-	1	2	3	2	5	12	22
Female	375	1	-	-	-	-	-	-	1	5	8	21
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	1	-	-	-	-	-	1	-	-
Male	5	-	-	1	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Maine												
All races, both sexes	664	-	1	-	-	-	-	-	-	2	4	11
Male	240	-	1	-	-	-	-	-	-	2	3	7
Female	424	-	-	-	-	-	-	-	-	-	1	4
White, both sexes	662	-	-	-	-	-	-	-	-	2	4	11
Male	238	-	-	-	-	-	-	-	-	2	3	7
Female	424	-	-	-	-	-	-	-	-	-	1	4
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	2,364	1	1	1	1	2	2	2	4	14	37	49
Male	944	1	1	-	-	1	1	1	2	9	17	25
Female	1,420	-	-	1	1	1	1	1	2	5	20	24
White, both sexes	1,718	1	1	1	1	2	1	1	1	5	11	15
Male	657	1	1	-	-	1	1	-	-	2	6	8
Female	1,061	-	-	1	1	1	-	1	1	3	5	7
Black, both sexes	599	-	-	-	-	-	1	1	2	8	26	33
Male	261	-	-	-	-	-	-	1	1	6	11	17
Female	338	-	-	-	-	-	1	-	1	2	15	16
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	44	-	-	-	-	-	-	-	1	1	-	1
Male	23	-	-	-	-	-	-	-	1	1	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Massachusetts												
All races, both sexes	2,832	2	3	-	-	1	1	3	4	6	9	36
Male	1,034	1	2	-	-	-	-	2	2	4	5	16
Female	1,798	1	1	-	-	1	1	1	2	2	4	20
White, both sexes	2,689	2	2	-	-	-	-	2	3	4	7	28
Male	976	1	1	-	-	-	-	1	1	2	4	12
Female	1,713	1	1	-	-	-	-	1	2	2	3	16
Black, both sexes	91	-	1	-	-	1	1	1	1	2	1	4
Male	34	-	1	-	-	-	-	1	1	2	-	1
Female	57	-	-	-	-	1	1	-	-	-	1	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	51	-	-	-	-	-	-	-	-	-	1	4
Male	23	-	-	-	-	-	-	-	-	-	1	3
Female	28	-	-	-	-	-	-	-	-	-	-	1
Michigan												
All races, both sexes	4,798	6	1	-	-	3	2	6	11	31	46	82
Male	1,927	4	-	-	-	2	2	1	6	16	24	45
Female	2,871	2	1	-	-	1	-	5	5	15	22	37
White, both sexes	4,110	6	1	-	-	1	2	5	6	14	25	59
Male	1,614	4	-	-	-	1	2	-	3	6	14	34
Female	2,496	2	1	-	-	-	-	5	3	8	11	25
Black, both sexes	633	-	-	-	-	2	-	1	5	14	19	22
Male	291	-	-	-	-	1	-	1	3	9	10	10
Female	342	-	-	-	-	1	-	-	2	5	9	12
American Indian 1/, both sexes	18	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	37	-	-	-	-	-	-	-	-	3	2	-
Male	13	-	-	-	-	-	-	-	-	1	-	-
Female	24	-	-	-	-	-	-	-	-	2	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Minnesota												
All races, both sexes	2,193	2	2	-	2	3	2	3	-	9	13	39
Male	853	1	1	-	1	1	1	3	-	6	4	22
Female	1,340	1	1	-	1	2	1	-	-	3	9	17
White, both sexes	2,100	1	1	-	1	3	1	2	-	8	10	30
Male	805	-	1	-	1	1	-	2	-	6	4	16
Female	1,295	1	-	-	-	2	1	-	-	2	6	14
Black, both sexes	48	1	1	-	1	-	1	-	-	-	1	4
Male	24	1	-	-	-	-	1	-	-	-	-	2
Female	24	-	1	-	1	-	-	-	-	-	1	2
American Indian 1/, both sexes	16	-	-	-	-	-	-	1	-	1	2	1
Male	9	-	-	-	-	-	-	1	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	1	2	-
Asian or Pacific Islander 2/, both sexes	29	-	-	-	-	-	-	-	-	-	-	4
Male	15	-	-	-	-	-	-	-	-	-	-	3
Female	14	-	-	-	-	-	-	-	-	-	-	1
Mississippi												
All races, both sexes	1,589	3	1	2	2	2	1	4	6	9	25	48
Male	651	1	-	1	-	1	-	3	4	5	12	23
Female	938	2	1	1	2	1	1	1	2	4	13	25
White, both sexes	1,005	2	-	2	1	1	-	2	2	4	9	13
Male	358	1	-	1	-	-	-	2	2	1	3	6
Female	647	1	-	1	1	1	-	-	-	3	6	7
Black, both sexes	579	1	1	-	1	1	1	2	4	4	16	34
Male	290	-	-	-	-	1	-	1	2	3	9	17
Female	289	1	1	-	1	-	1	1	2	1	7	17
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Missouri												
All races, both sexes	3,229	3	3	4	-	-	1	5	7	13	41	64
Male	1,240	2	1	1	-	-	-	3	3	6	21	23
Female	1,989	1	2	3	-	-	1	2	4	7	20	41
White, both sexes	2,871	2	3	3	-	-	-	3	4	9	30	46
Male	1,098	1	1	1	-	-	-	2	1	2	18	16
Female	1,773	1	2	2	-	-	-	1	3	7	12	30
Black, both sexes	328	1	-	-	-	-	1	2	2	4	11	17
Male	132	1	-	-	-	-	-	1	1	4	3	7
Female	196	-	-	-	-	-	1	1	1	-	8	10
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	25	-	-	1	-	-	-	-	1	-	-	1
Male	8	-	-	-	-	-	-	-	1	-	-	-
Female	17	-	-	1	-	-	-	-	-	-	-	1
Montana												
All races, both sexes	443	1	-	-	-	-	-	-	-	1	5	8
Male	171	-	-	-	-	-	-	-	-	-	1	4
Female	272	1	-	-	-	-	-	-	-	1	4	4
White, both sexes	425	1	-	-	-	-	-	-	-	1	4	8
Male	164	-	-	-	-	-	-	-	-	-	1	4
Female	261	1	-	-	-	-	-	-	-	1	3	4
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	17	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Nebraska												
All races, both sexes	921	1	2	-	-	1	1	2	2	1	5	11
Male	336	-	-	-	-	-	1	1	-	1	4	7
Female	585	1	2	-	-	1	-	1	2	-	1	4
White, both sexes	876	1	1	-	-	1	1	1	2	1	5	8
Male	316	-	-	-	-	-	1	-	-	1	4	5
Female	560	1	1	-	-	1	-	1	2	-	1	3
Black, both sexes	36	-	-	-	-	-	-	1	-	-	-	2
Male	16	-	-	-	-	-	-	1	-	-	-	1
Female	20	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	5	-	1	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	850	4	-	-	-	-	2	3	3	7	25	30
Male	407	3	-	-	-	-	1	1	2	3	11	17
Female	443	1	-	-	-	-	1	2	1	4	14	13
White, both sexes	710	4	-	-	-	-	2	2	1	5	16	20
Male	344	3	-	-	-	-	1	1	1	2	8	13
Female	366	1	-	-	-	-	1	1	-	3	8	7
Black, both sexes	75	-	-	-	-	-	-	-	2	1	4	6
Male	39	-	-	-	-	-	-	-	1	-	1	3
Female	36	-	-	-	-	-	-	-	1	1	3	3
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	1	1	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	58	-	-	-	-	-	-	1	-	-	4	4
Male	21	-	-	-	-	-	-	-	-	-	2	1
Female	37	-	-	-	-	-	-	1	-	-	2	3

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
New Hampshire												
All races, both sexes	489	2	1	-	-	-	-	-	-	1	5	8
Male	181	1	1	-	-	-	-	-	-	1	3	7
Female	308	1	-	-	-	-	-	-	-	-	2	1
White, both sexes	485	2	1	-	-	-	-	-	-	1	4	8
Male	181	1	1	-	-	-	-	-	-	1	3	7
Female	304	1	-	-	-	-	-	-	-	-	1	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	3,492	2	1	2	3	1	2	3	7	14	36	58
Male	1,375	2	-	1	1	-	2	2	5	9	19	36
Female	2,117	-	1	1	2	1	-	1	2	5	17	22
White, both sexes	2,944	1	-	2	2	1	2	2	4	7	15	38
Male	1,140	1	-	1	1	-	2	2	3	4	9	25
Female	1,804	-	-	1	1	1	-	-	1	3	6	13
Black, both sexes	458	1	1	-	1	-	-	1	2	6	20	17
Male	199	1	-	-	-	-	-	-	1	4	10	10
Female	259	-	1	-	1	-	-	1	1	2	10	7
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	85	-	-	-	-	-	-	-	1	1	1	3
Male	34	-	-	-	-	-	-	-	1	1	-	1
Female	51	-	-	-	-	-	-	-	-	-	1	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
New Mexico												
All races, both sexes	804	3	-	-	-	-	2	-	3	3	6	13
Male	333	-	-	-	-	-	-	-	3	1	3	5
Female	471	3	-	-	-	-	2	-	-	2	3	8
White, both sexes	740	2	-	-	-	-	2	-	2	2	5	12
Male	301	-	-	-	-	-	-	-	2	1	2	5
Female	439	2	-	-	-	-	2	-	-	1	3	7
Black, both sexes	17	-	-	-	-	-	-	-	1	-	1	1
Male	10	-	-	-	-	-	-	-	1	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	43	1	-	-	-	-	-	-	-	1	-	-
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	22	1	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	6,160	4	2	2	5	8	3	7	19	38	67	124
Male	2,411	2	1	-	2	4	2	4	10	19	33	66
Female	3,749	2	1	2	3	4	1	3	9	19	34	58
White, both sexes	5,135	2	2	1	3	4	2	5	16	21	42	81
Male	1,979	1	1	-	2	2	1	4	9	11	23	45
Female	3,156	1	1	1	1	2	1	1	7	10	19	36
Black, both sexes	857	1	-	-	1	4	1	2	3	16	22	36
Male	354	1	-	-	-	2	1	-	1	7	8	16
Female	503	-	-	-	1	2	-	2	2	9	14	20
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	163	1	-	1	1	-	-	-	-	1	3	7
Male	76	-	-	-	-	-	-	-	-	1	2	5
Female	87	1	-	1	1	-	-	-	-	-	1	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
North Carolina												
All races, both sexes	4,530	3	-	1	-	1	5	8	11	23	37	86
Male	1,788	-	-	1	-	1	2	4	6	10	18	45
Female	2,742	3	-	-	-	-	3	4	5	13	19	41
White, both sexes	3,422	1	-	1	-	1	3	4	5	9	14	40
Male	1,311	-	-	1	-	1	1	4	2	5	6	19
Female	2,111	1	-	-	-	-	2	-	3	4	8	21
Black, both sexes	1,036	2	-	-	-	-	1	4	6	13	21	44
Male	445	-	-	-	-	-	-	-	4	5	11	24
Female	591	2	-	-	-	-	1	4	2	8	10	20
American Indian 1/, both sexes	41	-	-	-	-	-	-	-	-	1	1	1
Male	17	-	-	-	-	-	-	-	-	-	1	1
Female	24	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	31	-	-	-	-	-	1	-	-	-	1	1
Male	15	-	-	-	-	-	1	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	1	-
North Dakota												
All races, both sexes	330	-	1	-	-	-	-	-	-	2	1	4
Male	126	-	-	-	-	-	-	-	-	1	-	4
Female	204	-	1	-	-	-	-	-	-	1	1	-
White, both sexes	325	-	1	-	-	-	-	-	-	1	1	4
Male	124	-	-	-	-	-	-	-	-	-	-	4
Female	201	-	1	-	-	-	-	-	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Ohio												
All races, both sexes	5,905	3	2	1	3	5	6	8	12	32	56	93
Male	2,299	2	1	1	1	3	3	6	6	17	28	53
Female	3,606	1	1	-	2	2	3	2	6	15	28	40
White, both sexes	5,159	1	1	-	3	4	5	6	8	21	36	71
Male	1,953	1	1	-	1	2	2	4	3	11	18	40
Female	3,206	-	-	-	2	2	3	2	5	10	18	31
Black, both sexes	713	2	1	1	-	1	1	2	4	9	17	22
Male	336	1	-	1	-	1	1	2	3	5	9	13
Female	377	1	1	-	-	-	-	-	1	4	8	9
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	29	-	-	-	-	-	-	-	-	2	2	-
Male	8	-	-	-	-	-	-	-	-	1	1	-
Female	21	-	-	-	-	-	-	-	-	1	1	-
Oklahoma												
All races, both sexes	2,126	3	-	1	1	-	1	5	9	8	24	35
Male	840	3	-	-	1	-	-	2	5	4	8	23
Female	1,286	-	-	1	-	-	1	3	4	4	16	12
White, both sexes	1,872	2	-	1	1	-	1	4	8	8	16	21
Male	728	2	-	-	1	-	-	2	4	4	3	11
Female	1,144	-	-	1	-	-	1	2	4	4	13	10
Black, both sexes	141	1	-	-	-	-	-	1	1	-	6	11
Male	64	1	-	-	-	-	-	-	1	-	4	10
Female	77	-	-	-	-	-	-	1	-	-	2	1
American Indian 1/, both sexes	103	-	-	-	-	-	-	-	-	-	2	3
Male	45	-	-	-	-	-	-	-	-	-	1	2
Female	58	-	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Oregon												
All races, both sexes	1,835	1	-	-	-	-	2	-	2	7	16	40
Male	783	-	-	-	-	-	1	-	2	3	10	29
Female	1,052	1	-	-	-	-	1	-	-	4	6	11
White, both sexes	1,745	1	-	-	-	-	1	-	2	7	15	35
Male	736	-	-	-	-	-	-	-	2	3	9	25
Female	1,009	1	-	-	-	-	1	-	-	4	6	10
Black, both sexes	31	-	-	-	-	-	-	-	-	-	1	1
Male	15	-	-	-	-	-	-	-	-	-	1	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	10	-	-	-	-	-	1	-	-	-	-	-
Male	6	-	-	-	-	-	1	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	49	-	-	-	-	-	-	-	-	-	-	4
Male	26	-	-	-	-	-	-	-	-	-	-	3
Female	23	-	-	-	-	-	-	-	-	-	-	1
Pennsylvania												
All races, both sexes	7,152	4	2	2	-	2	3	7	12	23	48	97
Male	2,742	2	1	-	-	2	1	4	5	9	29	49
Female	4,410	2	1	2	-	-	2	3	7	14	19	48
White, both sexes	6,443	4	2	-	-	2	3	4	8	22	36	65
Male	2,439	2	1	-	-	2	1	3	3	8	24	34
Female	4,004	2	1	-	-	-	2	1	5	14	12	31
Black, both sexes	643	-	-	2	-	-	-	2	3	1	10	30
Male	271	-	-	-	-	-	-	1	1	1	4	13
Female	372	-	-	2	-	-	-	1	2	-	6	17
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	62	-	-	-	-	-	-	1	1	-	2	1
Male	31	-	-	-	-	-	-	-	1	-	1	1
Female	31	-	-	-	-	-	-	1	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Rhode Island												
All races, both sexes	457	-	-	-	-	-	-	-	1	2	4	7
Male	168	-	-	-	-	-	-	-	1	1	1	3
Female	289	-	-	-	-	-	-	-	-	1	3	4
White, both sexes	432	-	-	-	-	-	-	-	1	-	2	6
Male	157	-	-	-	-	-	-	-	1	-	1	3
Female	275	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	18	-	-	-	-	-	-	-	-	2	2	1
Male	10	-	-	-	-	-	-	-	-	1	-	-
Female	8	-	-	-	-	-	-	-	-	1	2	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	2,466	-	2	1	2	-	-	2	13	26	33	60
Male	1,026	-	1	-	-	-	-	-	7	17	21	34
Female	1,440	-	1	1	2	-	-	2	6	9	12	26
White, both sexes	1,667	-	1	-	2	-	-	1	8	7	11	23
Male	650	-	1	-	-	-	-	-	5	4	8	14
Female	1,017	-	-	-	2	-	-	1	3	3	3	9
Black, both sexes	786	-	1	1	-	-	-	1	5	19	21	37
Male	370	-	-	-	-	-	-	-	2	13	12	20
Female	416	-	1	1	-	-	-	1	3	6	9	17
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
South Dakota												
All races, both sexes	410	1	-	-	-	2	-	-	1	-	1	7
Male	161	-	-	-	-	-	-	-	-	-	-	3
Female	249	1	-	-	-	2	-	-	1	-	1	4
White, both sexes	389	-	-	-	-	2	-	-	1	-	1	6
Male	150	-	-	-	-	-	-	-	-	-	-	3
Female	239	-	-	-	-	2	-	-	1	-	1	3
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	18	1	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	9	1	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	3,450	1	2	-	-	1	6	4	7	23	51	74
Male	1,322	1	1	-	-	-	2	3	3	11	29	39
Female	2,128	-	1	-	-	1	4	1	4	12	22	35
White, both sexes	2,893	1	1	-	-	1	3	3	2	14	32	47
Male	1,075	1	1	-	-	-	-	2	-	7	18	25
Female	1,818	-	-	-	-	1	3	1	2	7	14	22
Black, both sexes	535	-	1	-	-	-	3	1	4	9	19	27
Male	239	-	-	-	-	-	2	1	3	4	11	14
Female	296	-	1	-	-	-	1	-	1	5	8	13
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	20	-	-	-	-	-	-	-	1	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Texas												
All races, both sexes	9,796	12	7	3	3	4	6	17	32	70	117	212
Male	3,937	7	2	2	1	2	5	7	14	42	64	118
Female	5,859	5	5	1	2	2	1	10	18	28	53	94
White, both sexes	8,332	10	5	1	3	4	6	14	21	49	82	153
Male	3,321	5	1	1	1	2	5	6	11	31	52	87
Female	5,011	5	4	-	2	2	1	8	10	18	30	66
Black, both sexes	1,322	2	2	2	-	-	-	3	9	20	31	54
Male	556	2	1	1	-	-	-	1	2	10	11	30
Female	766	-	1	1	-	-	-	2	7	10	20	24
American Indian 1/, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	128	-	-	-	-	-	-	-	2	1	4	4
Male	54	-	-	-	-	-	-	-	1	1	1	1
Female	74	-	-	-	-	-	-	-	1	-	3	3
Utah												
All races, both sexes	755	3	-	-	1	-	-	2	2	4	7	9
Male	285	1	-	-	1	-	-	1	1	2	5	3
Female	470	2	-	-	-	-	-	1	1	2	2	6
White, both sexes	730	3	-	-	1	-	-	2	2	4	6	9
Male	271	1	-	-	1	-	-	1	1	2	4	3
Female	459	2	-	-	-	-	-	1	1	2	2	6
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Vermont												
All races, both sexes	269	-	-	-	-	-	-	1	1	-	2	1
Male	97	-	-	-	-	-	-	1	-	-	1	1
Female	172	-	-	-	-	-	-	-	1	-	1	-
White, both sexes	268	-	-	-	-	-	-	1	1	-	2	1
Male	96	-	-	-	-	-	-	1	-	-	1	1
Female	172	-	-	-	-	-	-	-	1	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	3,313	5	1	2	1	3	3	4	9	8	28	57
Male	1,295	5	-	1	-	3	2	2	6	2	11	29
Female	2,018	-	1	1	1	-	1	2	3	6	17	28
White, both sexes	2,498	1	1	2	-	2	2	3	2	3	15	32
Male	935	1	-	1	-	2	1	2	1	1	6	19
Female	1,563	-	1	1	-	-	1	1	1	2	9	13
Black, both sexes	747	4	-	-	1	1	1	-	6	4	13	23
Male	332	4	-	-	-	1	1	-	4	-	5	10
Female	415	-	-	-	1	-	-	-	2	4	8	13
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	67	-	-	-	-	-	-	1	1	1	-	2
Male	27	-	-	-	-	-	-	-	1	1	-	-
Female	40	-	-	-	-	-	-	1	-	-	-	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Washington												
All races, both sexes	2,692	3	1	1	2	2	2	5	3	7	17	32
Male	1,068	3	-	1	2	2	1	4	3	3	10	16
Female	1,624	-	1	-	-	-	1	1	-	4	7	16
White, both sexes	2,474	3	-	1	2	2	2	4	2	7	14	26
Male	974	3	-	1	2	2	1	4	2	3	9	13
Female	1,500	-	-	-	-	-	1	-	-	4	5	13
Black, both sexes	73	-	-	-	-	-	-	-	-	-	-	3
Male	34	-	-	-	-	-	-	-	-	-	-	2
Female	39	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	33	-	-	-	-	-	-	1	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	112	-	1	-	-	-	-	-	1	-	3	3
Male	50	-	-	-	-	-	-	-	1	-	1	1
Female	62	-	1	-	-	-	-	-	-	-	2	2
West Virginia												
All races, both sexes	1,113	1	-	-	-	-	1	-	5	4	9	19
Male	441	1	-	-	-	-	1	-	1	1	7	10
Female	672	-	-	-	-	-	-	-	4	3	2	9
White, both sexes	1,055	1	-	-	-	-	1	-	4	4	8	17
Male	419	1	-	-	-	-	1	-	1	1	6	9
Female	636	-	-	-	-	-	-	-	3	3	2	8
Black, both sexes	54	-	-	-	-	-	-	-	1	-	1	1
Male	19	-	-	-	-	-	-	-	-	-	1	1
Female	35	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Wisconsin												
All races, both sexes	2,738	2	-	-	1	-	5	1	9	10	17	49
Male	1,071	1	-	-	1	-	3	-	5	5	10	29
Female	1,667	1	-	-	-	-	2	1	4	5	7	20
White, both sexes	2,583	2	-	-	1	-	2	1	7	8	13	39
Male	1,004	1	-	-	1	-	2	-	5	4	7	26
Female	1,579	1	-	-	-	-	-	1	2	4	6	13
Black, both sexes	122	-	-	-	-	-	2	-	2	2	2	9
Male	50	-	-	-	-	-	1	-	-	1	1	3
Female	72	-	-	-	-	-	1	-	2	1	1	6
American Indian 1/, both sexes	14	-	-	-	-	-	-	-	-	-	2	1
Male	7	-	-	-	-	-	-	-	-	-	2	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	19	-	-	-	-	-	1	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	1	-	-	-	-	-
Wyoming												
All races, both sexes	209	-	-	-	-	-	-	-	1	3	1	5
Male	81	-	-	-	-	-	-	-	-	2	1	5
Female	128	-	-	-	-	-	-	-	1	1	-	-
White, both sexes	204	-	-	-	-	-	-	-	1	3	1	5
Male	77	-	-	-	-	-	-	-	-	2	1	5
Female	127	-	-	-	-	-	-	-	1	1	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
United States												
All races, both sexes	3,769	4,745	5,755	7,461	10,546	16,973	25,006	28,161	19,235	7,315	1,264	6
Male	2,097	2,757	3,235	3,978	5,373	7,868	9,832	9,423	4,955	1,376	161	4
Female	1,672	1,988	2,520	3,483	5,173	9,105	15,174	18,738	14,280	5,939	1,103	2
White, both sexes	2,457	3,192	4,246	5,660	8,363	14,354	21,986	25,409	17,531	6,688	1,106	4
Male	1,405	1,851	2,391	3,009	4,263	6,685	8,637	8,563	4,515	1,251	128	3
Female	1,052	1,341	1,855	2,651	4,100	7,669	13,349	16,846	13,016	5,437	978	1
Black, both sexes	1,151	1,306	1,285	1,502	1,755	2,068	2,317	2,063	1,302	491	136	2
Male	602	779	721	815	899	921	919	604	300	91	25	1
Female	549	527	564	687	856	1,147	1,398	1,459	1,002	400	111	1
American Indian 1/, both sexes	27	50	33	44	70	77	77	84	45	15	4	-
Male	14	27	22	19	34	44	33	28	11	2	1	-
Female	13	23	11	25	36	33	44	56	34	13	3	-
Asian or Pacific Islander 2/, both sexes	134	197	191	255	358	474	626	605	357	121	18	-
Male	76	100	101	135	177	218	243	228	129	32	7	-
Female	58	97	90	120	181	256	383	377	228	89	11	-
Alabama												
All races, both sexes	110	132	134	190	246	380	513	492	286	98	14	-
Male	64	80	74	107	136	175	195	168	68	13	4	-
Female	46	52	60	83	110	205	318	324	218	85	10	-
White, both sexes	52	67	92	129	167	292	401	399	228	73	10	-
Male	31	42	50	66	88	141	153	137	49	12	2	-
Female	21	25	42	63	79	151	248	262	179	61	8	-
Black, both sexes	57	65	42	61	78	87	109	93	57	24	4	-
Male	32	38	24	41	47	34	42	31	18	1	2	-
Female	25	27	18	20	31	53	67	62	39	23	2	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	1	1	-	-
Male	-	-	-	-	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	1	3	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	3	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Alaska												
All races, both sexes	4	7	2	15	18	23	21	31	23	3	2	-
Male	3	5	2	9	15	13	11	14	6	-	-	-
Female	1	2	-	6	3	10	10	17	17	3	2	-
White, both sexes	2	6	1	13	12	18	16	24	16	3	2	-
Male	1	5	1	8	10	9	7	11	3	-	-	-
Female	1	1	-	5	2	9	9	13	13	3	2	-
Black, both sexes	1	-	-	1	1	-	-	-	2	-	-	-
Male	1	-	-	1	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	1	-	-	1	2	4	5	5	4	-	-	-
Male	1	-	-	-	1	3	4	3	2	-	-	-
Female	-	-	-	1	1	1	1	2	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	3	1	-	2	1	-	-	-
Male	-	-	1	-	3	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	2	1	-	-	-
Arizona												
All races, both sexes	54	83	91	153	175	265	423	470	281	107	10	-
Male	31	43	54	84	89	118	142	153	71	20	2	-
Female	23	40	37	69	86	147	281	317	210	87	8	-
White, both sexes	46	74	84	139	164	251	408	449	271	102	9	-
Male	28	37	51	77	82	112	136	150	67	17	2	-
Female	18	37	33	62	82	139	272	299	204	85	7	-
Black, both sexes	2	4	4	7	4	6	7	7	4	2	1	-
Male	-	3	2	3	3	2	4	-	2	2	-	-
Female	2	1	2	4	1	4	3	7	2	-	1	-
American Indian 1/, both sexes	2	1	1	2	4	5	4	7	4	2	-	-
Male	1	-	-	2	2	2	2	2	2	-	-	-
Female	1	1	1	-	2	3	2	5	2	2	-	-
Asian or Pacific Islander 2/, both sexes	4	4	2	5	3	3	4	7	2	1	-	-
Male	2	3	1	2	2	2	-	1	-	1	-	-
Female	2	1	1	3	1	1	4	6	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Arkansas												
All races, both sexes	63	71	79	114	153	230	336	383	252	84	18	-
Male	40	44	40	70	78	108	127	122	53	13	2	-
Female	23	27	39	44	75	122	209	261	199	71	16	-
White, both sexes	42	50	60	93	128	198	304	346	212	73	15	-
Male	30	33	32	54	68	93	120	114	44	11	2	-
Female	12	17	28	39	60	105	184	232	168	62	13	-
Black, both sexes	21	20	19	20	24	30	30	35	39	11	3	-
Male	10	11	8	15	10	14	6	8	9	2	-	-
Female	11	9	11	5	14	16	24	27	30	9	3	-
American Indian 1/, both sexes	-	1	-	1	-	-	1	1	1	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	2	1	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	1	1	1	-	-	-	-
California												
All races, both sexes	458	514	607	771	1,082	1,741	2,637	3,025	2,055	831	151	1
Male	269	312	331	429	535	777	1,067	1,072	605	187	25	-
Female	189	202	276	342	547	964	1,570	1,953	1,450	644	126	1
White, both sexes	312	357	441	565	774	1,381	2,138	2,530	1,776	729	129	1
Male	189	227	237	308	380	626	870	902	516	167	20	-
Female	123	130	204	257	394	755	1,268	1,628	1,260	562	109	1
Black, both sexes	85	76	92	93	143	123	169	179	96	41	11	-
Male	44	43	50	58	76	44	69	48	19	8	-	-
Female	41	33	42	35	67	79	100	131	77	33	11	-
American Indian 1/, both sexes	3	4	4	5	6	10	12	13	2	4	-	-
Male	3	3	2	3	3	2	3	5	1	1	-	-
Female	-	1	2	2	3	8	9	8	1	3	-	-
Asian or Pacific Islander 2/, both sexes	58	77	70	108	159	227	318	303	181	57	11	-
Male	33	39	42	60	76	105	125	117	69	11	5	-
Female	25	38	28	48	83	122	193	186	112	46	6	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Colorado												
All races, both sexes	44	58	58	79	132	196	298	345	228	84	21	-
Male	27	36	34	42	62	87	103	113	67	17	2	-
Female	17	22	24	37	70	109	195	232	161	67	19	-
White, both sexes	39	50	52	76	124	184	286	329	221	79	19	-
Male	24	32	32	42	58	83	99	106	64	17	2	-
Female	15	18	20	34	66	101	187	223	157	62	17	-
Black, both sexes	5	6	4	-	7	9	5	8	4	4	2	-
Male	3	4	2	-	3	2	2	3	1	-	-	-
Female	2	2	2	-	4	7	3	5	3	4	2	-
American Indian 1/, both sexes	-	-	1	1	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	1	2	-	2	7	8	3	1	-	-
Male	-	-	-	-	-	1	2	4	2	-	-	-
Female	-	2	1	2	-	1	5	4	1	1	-	-
Connecticut												
All races, both sexes	24	40	45	67	89	176	245	388	233	96	17	-
Male	14	24	22	24	40	83	106	139	58	22	3	-
Female	10	16	23	43	49	93	139	249	175	74	14	-
White, both sexes	16	32	37	54	78	152	235	371	226	93	17	-
Male	10	21	18	20	34	72	102	135	57	21	3	-
Female	6	11	19	34	44	80	133	236	169	72	14	-
Black, both sexes	7	7	5	12	7	22	8	14	7	3	-	-
Male	4	2	2	4	4	10	3	2	1	1	-	-
Female	3	5	3	8	3	12	5	12	6	2	-	-
American Indian 1/, both sexes	-	1	1	-	1	1	-	-	-	-	-	-
Male	-	1	1	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	2	1	3	1	2	3	-	-	-	-
Male	-	-	1	-	1	-	1	2	-	-	-	-
Female	1	-	1	1	2	1	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Delaware												
All races, both sexes	6	11	11	28	43	50	70	61	48	20	7	-
Male	1	9	7	16	27	15	23	17	12	2	-	-
Female	5	2	4	12	16	35	47	44	36	18	7	-
White, both sexes	4	6	8	17	33	40	56	51	43	18	7	-
Male	1	5	5	11	20	12	16	14	9	2	-	-
Female	3	1	3	6	13	28	40	37	34	16	7	-
Black, both sexes	2	5	3	11	9	9	12	10	4	2	-	-
Male	-	4	2	5	6	3	7	3	2	-	-	-
Female	2	1	1	6	3	6	5	7	2	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	1	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	18	18	12	18	11	27	33	34	24	5	5	-
Male	11	9	6	8	5	10	8	10	4	1	2	-
Female	7	9	6	10	6	17	25	24	20	4	3	-
White, both sexes	2	-	-	3	3	3	12	8	6	2	3	-
Male	1	-	-	2	-	3	5	4	-	-	2	-
Female	1	-	-	1	3	-	7	4	6	2	1	-
Black, both sexes	16	18	12	15	7	24	21	26	18	3	2	-
Male	10	9	6	6	4	7	3	6	4	1	-	-
Female	6	9	6	9	3	17	18	20	14	2	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Florida												
All races, both sexes	235	302	389	495	681	1,110	1,669	1,810	1,167	482	76	-
Male	139	164	227	258	370	544	691	661	338	118	10	-
Female	96	138	162	237	311	566	978	1,149	829	364	66	-
White, both sexes	144	207	299	365	523	969	1,500	1,675	1,076	441	70	-
Male	93	113	177	192	288	482	631	625	319	104	6	-
Female	51	94	122	173	235	487	869	1,050	757	337	64	-
Black, both sexes	87	91	86	124	143	131	157	125	85	39	6	-
Male	44	50	48	62	74	58	55	34	17	12	4	-
Female	43	41	38	62	69	73	102	91	68	27	2	-
American Indian 1/, both sexes	-	1	-	-	2	3	3	-	1	-	-	-
Male	-	-	-	-	1	1	2	-	-	-	-	-
Female	-	1	-	-	1	2	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	4	3	4	6	13	7	9	10	5	2	-	-
Male	2	1	2	4	7	3	3	2	2	2	-	-
Female	2	2	2	2	6	4	6	8	3	-	-	-
Georgia												
All races, both sexes	172	186	263	291	330	453	644	685	434	157	26	1
Male	104	102	157	189	171	212	230	202	97	22	2	1
Female	68	84	106	102	159	241	414	483	337	135	24	-
White, both sexes	82	91	149	185	214	318	506	542	357	124	19	1
Male	52	49	93	122	111	145	182	164	78	14	1	1
Female	30	42	56	63	103	173	324	378	279	110	18	-
Black, both sexes	88	93	110	104	111	129	130	141	73	33	7	-
Male	51	52	62	67	59	65	45	37	19	8	1	-
Female	37	41	48	37	52	64	85	104	54	25	6	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	4	2	4	6	7	2	4	-	-	-
Male	1	1	2	-	1	2	3	1	-	-	-	-
Female	1	1	2	2	3	4	4	1	4	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Hawaii												
All races, both sexes	14	34	24	34	32	83	121	151	86	32	7	-
Male	11	18	13	17	17	43	58	59	35	10	2	-
Female	3	16	11	17	15	40	63	92	51	22	5	-
White, both sexes	2	6	8	10	8	16	29	38	16	7	1	-
Male	2	2	6	8	6	11	14	13	6	1	-	-
Female	-	4	2	2	2	5	15	25	10	6	1	-
Black, both sexes	-	2	-	-	1	1	1	1	-	-	1	-
Male	-	1	-	-	1	1	1	1	-	-	1	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	25	16	24	23	66	91	112	70	25	5	-
Male	9	14	7	9	10	31	43	45	29	9	1	-
Female	3	11	9	15	13	35	48	67	41	16	4	-
Idaho												
All races, both sexes	13	15	23	22	48	76	103	145	128	38	4	-
Male	6	8	14	10	26	30	45	44	40	12	-	-
Female	7	7	9	12	22	46	58	101	88	26	4	-
White, both sexes	13	14	21	21	46	75	101	144	127	38	4	-
Male	6	8	13	9	25	30	45	44	40	12	-	-
Female	7	6	8	12	21	45	56	100	87	26	4	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	1	-	1	-	1	1	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	1	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Illinois												
All races, both sexes	153	205	207	318	442	705	1,093	1,214	891	362	51	-
Male	80	118	124	181	225	321	435	393	215	46	5	-
Female	73	87	83	137	217	384	658	821	676	316	46	-
White, both sexes	93	129	145	235	338	582	945	1,106	836	326	46	-
Male	52	71	87	134	175	264	380	360	202	44	5	-
Female	41	58	58	101	163	318	565	746	634	282	41	-
Black, both sexes	56	70	58	78	94	114	126	90	49	33	5	-
Male	25	43	36	45	44	53	45	27	11	2	-	-
Female	31	27	22	33	50	61	81	63	38	31	5	-
American Indian 1/, both sexes	2	1	-	-	2	1	1	1	-	-	-	-
Male	1	1	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	1	1	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	5	4	5	8	8	21	17	6	3	-	-
Male	2	3	1	2	5	4	10	6	2	-	-	-
Female	-	2	3	3	3	4	11	11	4	3	-	-
Indiana												
All races, both sexes	76	116	116	149	242	349	611	630	463	172	35	-
Male	44	58	60	80	134	169	229	190	116	33	2	-
Female	32	58	56	69	108	180	382	440	347	139	33	-
White, both sexes	54	98	95	130	216	316	572	604	446	168	35	-
Male	30	50	47	67	117	152	215	180	113	33	2	-
Female	24	48	48	63	99	164	357	424	333	135	33	-
Black, both sexes	22	18	21	18	25	32	38	26	16	4	-	-
Male	14	8	13	12	16	16	14	10	3	-	-	-
Female	8	10	8	6	9	16	24	16	13	4	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	1	1	-	1	-	-	-
Male	-	-	-	1	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Iowa												
All races, both sexes	33	28	51	58	126	207	314	377	298	132	25	-
Male	17	19	35	33	61	107	135	122	62	27	2	-
Female	16	9	16	25	65	100	179	255	236	105	23	-
White, both sexes	28	26	46	54	123	204	308	370	295	132	25	-
Male	16	19	33	29	60	104	132	120	61	27	2	-
Female	12	7	13	25	63	100	176	250	234	105	23	-
Black, both sexes	4	2	2	4	3	3	5	5	3	-	-	-
Male	1	-	1	4	1	3	2	1	1	-	-	-
Female	3	2	1	-	2	-	3	4	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	3	-	-	-	-	2	-	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	1	-	2	-	-	-	-	1	-	-	-	-
Kansas												
All races, both sexes	24	36	57	75	99	168	258	329	266	111	23	-
Male	10	18	27	37	50	82	104	118	81	20	4	-
Female	14	18	30	38	49	86	154	211	185	91	19	-
White, both sexes	17	27	55	64	90	155	250	321	250	109	23	-
Male	7	12	26	32	43	73	100	115	78	20	4	-
Female	10	15	29	32	47	82	150	206	172	89	19	-
Black, both sexes	6	9	2	7	7	8	7	5	13	1	-	-
Male	3	6	1	4	6	6	4	3	2	-	-	-
Female	3	3	1	3	1	2	3	2	11	1	-	-
American Indian 1/, both sexes	-	-	-	-	1	2	-	3	1	1	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	3	1	1	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	4	1	3	1	-	2	-	-	-
Male	-	-	-	1	1	2	-	-	1	-	-	-
Female	1	-	-	3	-	1	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Kentucky												
All races, both sexes	44	81	122	130	178	299	389	424	273	109	18	-
Male	23	41	61	67	86	144	153	112	53	23	1	-
Female	21	40	61	63	92	155	236	312	220	86	17	-
White, both sexes	34	69	112	114	162	277	363	402	255	105	17	-
Male	19	31	57	59	80	134	139	105	50	22	1	-
Female	15	38	55	55	82	143	224	297	205	83	16	-
Black, both sexes	10	11	9	15	16	21	26	20	16	4	1	-
Male	4	9	3	7	6	9	14	6	3	1	-	-
Female	6	2	6	8	10	12	12	14	13	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	1	-	1	-	2	2	-	-	-
Male	-	1	1	1	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	2	-	-	-
Louisiana												
All races, both sexes	88	111	119	144	192	302	382	348	225	74	10	1
Male	50	60	59	72	101	129	159	111	56	13	1	1
Female	38	51	60	72	91	173	223	237	169	61	9	-
White, both sexes	47	43	55	72	126	216	302	271	173	62	6	-
Male	29	21	27	35	69	91	124	83	46	10	1	-
Female	18	22	28	37	57	125	178	188	127	52	5	-
Black, both sexes	39	67	62	71	64	84	80	75	51	12	4	1
Male	19	38	32	36	32	38	35	27	10	3	-	1
Female	20	29	30	35	32	46	45	48	41	9	4	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	2	1	2	2	-	2	1	-	-	-
Male	2	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	2	-	2	2	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Maine												
All races, both sexes	15	11	16	32	47	89	123	147	112	46	8	-
Male	10	5	9	18	21	39	43	39	31	9	3	-
Female	5	6	7	14	26	50	80	108	81	37	5	-
White, both sexes	15	11	16	32	47	88	123	147	112	46	8	-
Male	10	5	9	18	21	38	43	39	31	9	3	-
Female	5	6	7	14	26	50	80	108	81	37	5	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	70	83	102	100	190	304	396	496	345	135	28	1
Male	37	51	54	46	103	132	168	177	82	31	4	1
Female	33	32	48	54	87	172	228	319	263	104	24	-
White, both sexes	33	37	47	61	107	226	299	426	302	118	21	1
Male	19	21	27	28	53	95	131	152	78	30	2	1
Female	14	16	20	33	54	131	168	274	224	88	19	-
Black, both sexes	37	42	51	33	77	71	92	61	42	16	6	-
Male	18	28	25	13	46	32	36	22	3	1	1	-
Female	19	14	26	20	31	39	56	39	39	15	5	-
American Indian 1/, both sexes	-	-	-	-	1	2	-	-	-	-	-	-
Male	-	-	-	-	1	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	4	4	6	5	5	5	9	1	1	1	-
Male	-	2	2	5	3	3	1	3	1	-	1	-
Female	-	2	2	1	2	2	4	6	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Massachusetts												
All races, both sexes	44	53	78	97	195	346	536	670	502	203	43	-
Male	21	32	40	46	113	162	198	212	135	38	5	-
Female	23	21	38	51	82	184	338	458	367	165	38	-
White, both sexes	38	43	71	85	176	325	518	656	488	199	42	-
Male	19	29	35	40	105	157	189	209	131	36	4	-
Female	19	14	36	45	71	168	329	447	357	163	38	-
Black, both sexes	3	6	3	8	10	13	11	12	11	1	1	-
Male	1	1	2	5	3	3	6	2	4	-	1	-
Female	2	5	1	3	7	10	5	10	7	1	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	4	4	4	8	8	7	2	3	3	-	-
Male	1	2	3	1	4	2	3	1	-	2	-	-
Female	2	2	1	3	4	6	4	1	3	1	-	-
Michigan												
All races, both sexes	135	142	188	264	327	600	934	1,051	684	246	39	-
Male	71	84	114	136	159	288	384	357	174	55	5	-
Female	64	58	74	128	168	312	550	694	510	191	34	-
White, both sexes	86	84	139	200	259	533	829	970	631	227	33	-
Male	48	50	86	103	123	253	343	331	159	49	5	-
Female	38	34	53	97	136	280	486	639	472	178	28	-
Black, both sexes	47	55	47	59	64	58	94	73	49	18	6	-
Male	23	31	26	31	34	28	40	24	14	6	-	-
Female	24	24	21	28	30	30	54	49	35	12	6	-
American Indian 1/, both sexes	1	-	1	2	2	4	4	2	1	-	-	-
Male	-	-	1	1	1	4	-	1	-	-	-	-
Female	1	-	-	1	1	-	4	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	3	1	3	2	5	7	6	3	1	-	-
Male	-	3	1	1	1	3	1	1	1	-	-	-
Female	1	-	-	2	1	2	6	5	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Minnesota												
All races, both sexes	43	61	55	99	139	225	390	523	393	159	31	-
Male	23	37	37	58	68	121	164	178	98	26	3	-
Female	20	24	18	41	71	104	226	345	295	133	28	-
White, both sexes	34	47	51	94	129	219	378	514	388	158	31	-
Male	17	27	34	55	66	118	159	173	97	25	3	-
Female	17	20	17	39	63	101	219	341	291	133	28	-
Black, both sexes	7	7	3	1	5	3	7	2	4	-	-	-
Male	5	4	2	1	1	2	3	1	1	-	-	-
Female	2	3	1	-	4	1	4	1	3	-	-	-
American Indian 1/, both sexes	1	3	1	-	2	1	1	-	1	1	-	-
Male	-	3	1	-	1	1	-	-	-	1	-	-
Female	1	-	-	-	1	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	4	-	4	3	2	4	7	-	-	-	-
Male	1	3	-	2	-	-	2	4	-	-	-	-
Female	-	1	-	2	3	2	2	3	-	-	-	-
Mississippi												
All races, both sexes	57	77	92	111	147	205	286	253	184	57	17	-
Male	36	60	53	50	80	94	101	76	38	11	2	-
Female	21	17	39	61	67	111	185	177	146	46	15	-
White, both sexes	27	26	43	63	91	142	198	194	132	44	9	-
Male	15	17	22	28	49	59	65	57	23	7	-	-
Female	12	9	21	35	42	83	133	137	109	37	9	-
Black, both sexes	30	51	49	48	56	62	87	58	52	13	8	-
Male	21	43	31	22	31	35	35	18	15	4	2	-
Female	9	8	18	26	25	27	52	40	37	9	6	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Missouri												
All races, both sexes	65	105	121	175	233	387	621	683	479	192	27	-
Male	33	50	64	98	128	177	250	231	117	32	-	-
Female	32	55	57	77	105	210	371	452	362	160	27	-
White, both sexes	49	77	105	152	197	331	566	647	443	182	22	-
Male	26	39	53	87	108	152	229	220	112	30	-	-
Female	23	38	52	65	89	179	337	427	331	152	22	-
Black, both sexes	15	26	14	22	32	49	51	33	33	10	5	-
Male	7	11	10	11	18	20	21	10	5	2	-	-
Female	8	15	4	11	14	29	30	23	28	8	5	-
American Indian 1/, both sexes	1	-	-	1	-	2	-	1	-	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	2	-	4	5	4	2	3	-	-	-
Male	-	-	1	-	2	3	-	1	-	-	-	-
Female	-	2	1	-	2	2	4	1	3	-	-	-
Montana												
All races, both sexes	9	12	24	21	28	52	72	101	76	32	1	-
Male	5	9	11	12	9	29	31	31	22	7	-	-
Female	4	3	13	9	19	23	41	70	54	25	1	-
White, both sexes	8	10	23	19	28	47	71	98	74	32	1	-
Male	4	8	11	11	9	25	31	31	22	7	-	-
Female	4	2	12	8	19	22	40	67	52	25	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	2	1	2	-	5	1	2	2	-	-	-
Male	1	1	-	1	-	4	-	-	-	-	-	-
Female	-	1	1	1	-	1	1	2	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Nebraska												
All races, both sexes	20	27	21	42	64	108	169	189	160	77	18	-
Male	11	20	12	20	39	51	65	64	30	8	2	-
Female	9	7	9	22	25	57	104	125	130	69	16	-
White, both sexes	16	20	19	39	59	104	167	183	158	74	16	-
Male	7	16	11	20	36	49	64	63	30	8	1	-
Female	9	4	8	19	23	55	103	120	128	66	15	-
Black, both sexes	3	6	1	3	3	3	2	6	1	3	2	-
Male	3	3	1	-	2	2	1	1	-	-	1	-
Female	-	3	-	3	1	1	1	5	1	3	1	-
American Indian 1/, both sexes	1	-	-	-	1	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	1	1	-	-	-	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
Nevada												
All races, both sexes	30	38	54	65	82	115	145	133	87	24	3	-
Male	21	22	31	36	50	54	70	46	32	5	2	-
Female	9	16	23	29	32	61	75	87	55	19	1	-
White, both sexes	25	32	40	43	68	96	130	121	80	22	3	-
Male	17	20	21	27	43	45	64	42	30	4	2	-
Female	8	12	19	16	25	51	66	79	50	18	1	-
Black, both sexes	2	1	9	12	9	9	9	5	4	2	-	-
Male	2	-	8	5	4	5	5	3	1	1	-	-
Female	-	1	1	7	5	4	4	2	3	1	-	-
American Indian 1/, both sexes	-	-	-	1	1	-	2	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	5	5	9	4	10	4	6	3	-	-	-
Male	2	2	2	4	2	4	1	-	1	-	-	-
Female	1	3	3	5	2	6	3	6	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
New Hampshire												
All races, both sexes	4	17	10	18	40	50	88	108	91	42	4	-
Male	1	12	8	9	20	21	32	32	24	9	-	-
Female	3	5	2	9	20	29	56	76	67	33	4	-
White, both sexes	4	17	9	17	40	50	88	108	90	42	4	-
Male	1	12	8	9	20	21	32	32	24	9	-	-
Female	3	5	1	8	20	29	56	76	66	33	4	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-
New Jersey												
All races, both sexes	97	116	116	185	236	456	685	740	496	203	32	1
Male	50	65	64	109	123	211	263	237	133	36	7	-
Female	47	51	52	76	113	245	422	503	363	167	25	1
White, both sexes	64	80	83	130	186	398	579	681	455	187	27	-
Male	37	48	45	77	96	187	217	222	125	32	6	-
Female	27	32	38	53	90	211	362	459	330	155	21	-
Black, both sexes	27	31	27	46	39	49	87	48	34	15	5	1
Male	9	16	15	27	19	22	41	13	6	4	1	-
Female	18	15	12	19	20	27	46	35	28	11	4	1
American Indian 1/, both sexes	-	-	-	1	1	-	2	-	1	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	6	5	6	8	10	9	17	11	6	1	-	-
Male	4	1	4	5	7	2	4	2	2	-	-	-
Female	2	4	2	3	3	7	13	9	4	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
New Mexico												
All races, both sexes	16	30	31	55	75	111	154	169	89	42	2	-
Male	7	16	16	34	46	65	45	59	26	6	1	-
Female	9	14	15	21	29	46	109	110	63	36	1	-
White, both sexes	16	26	30	49	63	104	144	158	83	40	2	-
Male	7	14	16	31	38	59	41	54	24	6	1	-
Female	9	12	14	18	25	45	103	104	59	34	1	-
Black, both sexes	-	1	-	2	3	3	3	2	-	-	-	-
Male	-	1	-	2	2	3	-	-	-	-	-	-
Female	-	-	-	-	1	-	3	2	-	-	-	-
American Indian 1/, both sexes	-	3	1	4	9	4	5	8	5	2	-	-
Male	-	1	-	1	6	3	4	4	2	-	-	-
Female	-	2	1	3	3	1	1	4	3	2	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	2	-	1	-	-	-
New York												
All races, both sexes	176	215	246	307	500	776	1,083	1,306	843	360	69	-
Male	81	127	139	155	248	360	403	457	228	65	5	-
Female	95	88	107	152	252	416	680	849	615	295	64	-
White, both sexes	117	142	180	227	385	629	929	1,182	770	335	60	-
Male	54	82	104	115	190	296	348	417	212	57	5	-
Female	63	60	76	112	195	333	581	765	558	278	55	-
Black, both sexes	53	64	56	70	92	121	120	106	61	19	9	-
Male	27	40	27	34	44	52	40	34	13	7	-	-
Female	26	24	29	36	48	69	80	72	48	12	9	-
American Indian 1/, both sexes	-	1	-	1	2	1	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	1	-	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	8	10	9	21	25	34	18	12	6	-	-
Male	-	5	8	5	13	12	15	6	3	1	-	-
Female	6	3	2	4	8	13	19	12	9	5	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
North Carolina												
All races, both sexes	135	188	216	276	395	600	807	890	608	197	43	-
Male	74	119	121	154	200	275	296	276	145	35	6	-
Female	61	69	95	122	195	325	511	614	463	162	37	-
White, both sexes	66	97	125	181	295	463	668	723	524	167	35	-
Male	39	56	74	105	148	213	243	233	127	30	4	-
Female	27	41	51	76	147	250	425	490	397	137	31	-
Black, both sexes	65	81	84	88	94	127	133	158	79	29	7	-
Male	34	57	44	46	51	56	48	41	18	5	1	-
Female	31	24	40	42	43	71	85	117	61	24	6	-
American Indian 1/, both sexes	2	4	4	4	4	4	4	7	3	1	1	-
Male	-	3	2	2	1	2	3	1	-	-	1	-
Female	2	1	2	2	3	2	1	6	3	1	-	-
Asian or Pacific Islander 2/, both sexes	2	6	3	3	2	6	2	2	2	-	-	-
Male	1	3	1	1	-	4	2	1	-	-	-	-
Female	1	3	2	2	2	2	-	1	2	-	-	-
North Dakota												
All races, both sexes	5	4	12	17	16	35	57	77	62	35	2	-
Male	3	4	3	7	7	19	27	25	19	7	-	-
Female	2	-	9	10	9	16	30	52	43	28	2	-
White, both sexes	5	4	12	16	15	33	57	77	62	35	2	-
Male	3	4	3	7	7	18	27	25	19	7	-	-
Female	2	-	9	9	8	15	30	52	43	28	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	1	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Ohio												
All races, both sexes	147	193	242	313	490	753	1,086	1,224	900	277	59	-
Male	70	121	129	142	235	349	427	425	229	45	6	-
Female	77	72	113	171	255	404	659	799	671	232	53	-
White, both sexes	99	139	191	239	407	655	993	1,126	849	252	53	-
Male	51	88	97	105	199	304	378	389	213	42	4	-
Female	48	51	94	134	208	351	615	737	636	210	49	-
Black, both sexes	48	53	47	73	78	95	91	91	47	24	6	-
Male	19	33	31	37	36	44	48	33	15	2	2	-
Female	29	20	16	36	42	51	43	58	32	22	4	-
American Indian 1/, both sexes	-	-	1	-	-	1	-	1	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	3	1	5	2	2	6	4	1	-	-
Male	-	-	-	-	-	-	1	3	1	1	-	-
Female	-	1	3	1	5	2	1	3	3	-	-	-
Oklahoma												
All races, both sexes	65	60	94	134	188	313	387	417	284	83	14	-
Male	35	35	61	68	93	134	143	137	70	15	3	-
Female	30	25	33	66	95	179	244	280	214	68	11	-
White, both sexes	52	41	75	104	167	276	354	387	266	79	9	-
Male	30	25	45	55	86	122	128	128	67	14	1	-
Female	22	16	30	49	81	154	226	259	199	65	8	-
Black, both sexes	7	7	11	17	8	24	22	13	7	2	3	-
Male	2	5	9	6	3	6	10	3	1	1	2	-
Female	5	2	2	11	5	18	12	10	6	1	1	-
American Indian 1/, both sexes	5	9	7	13	12	12	9	17	10	2	2	-
Male	3	4	7	7	4	6	4	6	1	-	-	-
Female	2	5	-	6	8	6	5	11	9	2	2	-
Asian or Pacific Islander 2/, both sexes	1	3	1	-	1	1	2	-	1	-	-	-
Male	-	1	-	-	-	-	1	-	1	-	-	-
Female	1	2	1	-	1	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Oregon												
All races, both sexes	43	58	74	96	142	234	324	402	281	95	18	-
Male	25	28	41	49	75	124	141	148	78	24	5	-
Female	18	30	33	47	67	110	183	254	203	71	13	-
White, both sexes	42	53	67	87	126	223	312	390	277	89	18	-
Male	24	24	37	45	66	115	137	146	76	22	5	-
Female	18	29	30	42	60	108	175	244	201	67	13	-
Black, both sexes	-	2	2	4	6	2	4	5	2	2	-	-
Male	-	2	2	3	4	1	-	-	1	-	-	-
Female	-	-	-	1	2	1	4	5	1	2	-	-
American Indian 1/, both sexes	-	1	1	-	2	1	2	2	-	-	-	-
Male	-	1	1	-	1	1	1	-	-	-	-	-
Female	-	-	-	-	1	-	1	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	4	5	8	8	6	5	2	4	-	-
Male	1	1	1	1	4	7	3	2	1	2	-	-
Female	-	1	3	4	4	1	3	3	1	2	-	-
Pennsylvania												
All races, both sexes	159	193	253	318	454	922	1,431	1,628	1,139	407	48	-
Male	84	110	135	163	226	431	576	569	270	71	5	-
Female	75	83	118	155	228	491	855	1,059	869	336	43	-
White, both sexes	109	138	196	262	385	829	1,344	1,520	1,085	385	44	-
Male	56	80	103	133	193	386	544	537	257	67	5	-
Female	53	58	93	129	192	443	800	983	828	318	39	-
Black, both sexes	47	50	53	49	60	84	81	97	51	20	3	-
Male	26	27	29	23	30	41	30	29	12	4	-	-
Female	21	23	24	26	30	43	51	68	39	16	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	1	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	1	-
Asian or Pacific Islander 2/, both sexes	3	5	4	7	9	9	6	10	2	2	-	-
Male	2	3	3	7	3	4	2	3	1	-	-	-
Female	1	2	1	-	6	5	4	7	1	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Rhode Island												
All races, both sexes	11	9	19	16	20	46	93	107	85	34	3	-
Male	4	3	15	6	14	18	38	35	23	6	-	-
Female	7	6	4	10	6	28	55	72	62	28	3	-
White, both sexes	10	9	15	15	17	43	91	105	83	32	3	-
Male	4	3	11	6	11	17	37	35	22	6	-	-
Female	6	6	4	9	6	26	54	70	61	26	3	-
Black, both sexes	-	-	4	-	2	3	2	-	1	1	-	-
Male	-	-	4	-	2	1	1	-	1	-	-	-
Female	-	-	-	-	-	2	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	1	-	-	2	-	1	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	2	-	1	-	-
South Carolina												
All races, both sexes	108	122	145	187	216	296	427	440	272	99	14	1
Male	54	81	90	112	106	128	179	132	50	12	1	1
Female	54	41	55	75	110	168	248	308	222	87	13	-
White, both sexes	50	63	77	107	131	204	315	354	225	78	9	1
Male	22	39	47	63	65	88	131	110	40	12	-	1
Female	28	24	30	44	66	116	184	244	185	66	9	-
Black, both sexes	58	59	65	78	83	91	110	84	47	21	5	-
Male	32	42	41	49	39	40	47	22	10	-	1	-
Female	26	17	24	29	44	51	63	62	37	21	4	-
American Indian 1/, both sexes	-	-	-	1	-	-	2	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	3	1	2	1	-	1	-	-	-	-
Male	-	-	2	-	2	-	-	-	-	-	-	-
Female	-	-	1	1	-	1	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
South Dakota												
All races, both sexes	10	8	13	13	19	54	64	101	76	32	8	-
Male	6	4	9	7	10	26	29	39	20	8	-	-
Female	4	4	4	6	9	28	35	62	56	24	8	-
White, both sexes	8	5	11	12	16	50	62	100	75	32	8	-
Male	5	1	8	6	9	23	28	39	20	8	-	-
Female	3	4	3	6	7	27	34	61	55	24	8	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	2	1	-	3	4	2	1	1	-	-	-
Male	1	2	1	-	1	3	1	-	-	-	-	-
Female	1	-	-	-	2	1	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	1	-	-	-	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	113	138	150	235	290	450	619	668	444	145	29	-
Male	63	80	77	122	129	210	236	195	102	14	5	-
Female	50	58	73	113	161	240	383	473	342	131	24	-
White, both sexes	67	91	113	184	238	390	545	602	400	132	27	-
Male	39	54	57	93	113	182	203	174	89	13	4	-
Female	28	37	56	91	125	208	342	428	311	119	23	-
Black, both sexes	44	43	35	49	50	58	72	63	43	12	2	-
Male	23	24	20	27	16	26	33	20	13	1	1	-
Female	21	19	15	22	34	32	39	43	30	11	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	2	4	2	2	2	2	1	3	1	-	-	-
Male	1	2	-	2	-	2	-	1	-	-	-	-
Female	1	2	2	-	2	-	1	2	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Texas												
All races, both sexes	310	400	516	572	827	1,242	1,780	1,906	1,220	451	89	-
Male	188	226	279	281	405	552	691	645	316	82	8	-
Female	122	174	237	291	422	690	1,089	1,261	904	369	81	-
White, both sexes	209	297	380	449	675	1,062	1,586	1,741	1,098	412	75	-
Male	130	169	214	222	330	479	615	599	282	74	5	-
Female	79	128	166	227	345	583	971	1,142	816	338	70	-
Black, both sexes	95	94	122	113	137	162	170	146	111	35	14	-
Male	55	53	60	53	68	64	68	39	28	7	3	-
Female	40	41	62	60	69	98	102	107	83	28	11	-
American Indian 1/, both sexes	1	1	1	1	1	2	2	3	1	-	-	-
Male	-	-	1	-	-	2	-	2	1	-	-	-
Female	1	1	-	1	1	-	2	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	8	13	9	14	16	22	16	10	4	-	-
Male	3	4	4	6	7	7	8	5	5	1	-	-
Female	2	4	9	3	7	9	14	11	5	3	-	-
Utah												
All races, both sexes	19	20	29	35	62	94	152	173	105	31	7	-
Male	9	10	17	18	29	38	56	62	28	4	-	-
Female	10	10	12	17	33	56	96	111	77	27	7	-
White, both sexes	16	19	27	33	57	90	151	169	104	31	6	-
Male	7	9	15	17	27	35	55	61	28	4	-	-
Female	9	10	12	16	30	55	96	108	76	27	6	-
Black, both sexes	-	-	-	1	-	1	-	2	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	2	1	-	-	-
American Indian 1/, both sexes	1	1	1	-	-	-	-	-	-	-	-	-
Male	1	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	1	1	5	3	1	2	-	-	1	-
Male	1	-	1	1	2	2	1	1	-	-	-	-
Female	1	-	-	-	3	1	-	1	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Vermont												
All races, both sexes	3	2	6	13	15	25	60	64	50	26	-	-
Male	2	1	1	8	8	12	22	20	12	8	-	-
Female	1	1	5	5	7	13	38	44	38	18	-	-
White, both sexes	3	2	6	12	15	25	60	64	50	26	-	-
Male	2	1	1	7	8	12	22	20	12	8	-	-
Female	1	1	5	5	7	13	38	44	38	18	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	84	123	160	190	282	422	580	680	474	171	26	-
Male	47	71	99	105	147	192	217	206	117	29	4	-
Female	37	52	61	85	135	230	363	474	357	142	22	-
White, both sexes	47	75	103	118	192	300	465	568	405	141	21	-
Male	25	43	65	62	97	137	170	169	106	23	4	-
Female	22	32	38	56	95	163	295	399	299	118	17	-
Black, both sexes	36	44	53	63	82	112	106	104	63	26	5	-
Male	22	27	31	38	44	52	46	33	9	5	-	-
Female	14	17	22	25	38	60	60	71	54	21	5	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	4	4	9	8	10	9	8	5	4	-	-
Male	-	1	3	5	6	3	1	4	1	1	-	-
Female	1	3	1	4	2	7	8	4	4	3	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Washington												
All races, both sexes	62	75	90	124	205	330	504	601	435	169	22	-
Male	29	32	55	63	104	155	205	207	124	43	6	-
Female	33	43	35	61	101	175	299	394	311	126	16	-
White, both sexes	45	58	77	107	178	307	468	571	413	165	22	-
Male	21	28	46	53	91	143	192	194	118	42	6	-
Female	24	30	31	54	87	164	276	377	295	123	16	-
Black, both sexes	9	4	6	7	5	10	9	12	7	1	-	-
Male	3	1	4	5	3	6	3	4	3	-	-	-
Female	6	3	2	2	2	4	6	8	4	1	-	-
American Indian 1/, both sexes	2	7	3	1	4	3	8	4	-	-	-	-
Male	-	1	2	-	1	1	3	2	-	-	-	-
Female	2	6	1	1	3	2	5	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	6	4	9	18	10	19	14	15	3	-	-
Male	5	2	3	5	9	5	7	7	3	1	-	-
Female	1	4	1	4	9	5	12	7	12	2	-	-
West Virginia												
All races, both sexes	29	41	52	65	101	155	222	229	121	52	7	-
Male	19	25	25	32	50	70	91	73	23	10	2	-
Female	10	16	27	33	51	85	131	156	98	42	5	-
White, both sexes	26	35	48	64	100	146	214	217	115	50	5	-
Male	16	20	23	32	50	67	89	70	22	10	1	-
Female	10	15	25	32	50	79	125	147	93	40	4	-
Black, both sexes	3	5	3	1	1	9	8	12	5	2	2	-
Male	3	4	1	-	-	3	2	3	-	-	1	-
Female	-	1	2	1	1	6	6	9	5	2	1	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Wisconsin												
All races, both sexes	50	70	108	127	191	307	534	597	447	183	30	-
Male	32	47	72	76	94	147	208	199	116	24	2	-
Female	18	23	36	51	97	160	326	398	331	159	28	-
White, both sexes	40	57	96	112	174	290	513	584	435	179	30	-
Male	27	40	65	69	86	138	200	196	111	24	2	-
Female	13	17	31	43	88	152	313	388	324	155	28	-
Black, both sexes	7	10	9	12	15	14	15	10	10	3	-	-
Male	2	5	6	6	6	9	4	2	4	-	-	-
Female	5	5	3	6	9	5	11	8	6	3	-	-
American Indian 1/, both sexes	1	2	1	1	1	2	1	2	-	-	-	-
Male	1	1	-	-	1	-	1	1	-	-	-	-
Female	-	1	1	1	-	2	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	2	2	1	1	5	1	2	1	-	-
Male	2	1	1	1	1	-	3	-	1	-	-	-
Female	-	-	1	1	-	1	2	1	1	1	-	-
Wyoming												
All races, both sexes	2	6	12	8	11	31	37	46	30	13	3	-
Male	2	2	7	5	6	17	12	14	6	2	-	-
Female	-	4	5	3	5	14	25	32	24	11	3	-
White, both sexes	2	5	11	8	11	29	37	46	29	13	3	-
Male	2	1	6	5	6	15	12	14	6	2	-	-
Female	-	4	5	3	5	14	25	32	23	11	3	-
Black, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	1	-	-	-	-	-	1	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
United States												
All races, both sexes	5,708	1	1	10	17	26	39	77	102	218	354	540
Male	2,107	-	-	8	8	14	23	38	47	98	152	212
Female	3,601	1	1	2	9	12	16	39	55	120	202	328
White, both sexes	4,553	1	-	8	14	21	28	59	67	145	251	389
Male	1,701	-	-	7	7	11	14	33	31	66	115	168
Female	2,852	1	-	1	7	10	14	26	36	79	136	221
Black, both sexes	828	-	1	2	2	5	6	16	29	55	79	124
Male	295	-	-	1	-	3	5	5	13	24	29	33
Female	533	-	1	1	2	2	1	11	16	31	50	91
American Indian 1/, both sexes	48	-	-	-	-	-	1	-	1	3	6	5
Male	22	-	-	-	-	-	1	-	1	1	1	3
Female	26	-	-	-	-	-	-	-	-	2	5	2
Asian or Pacific Islander 2/, both sexes	279	-	-	-	1	-	4	2	5	15	18	22
Male	89	-	-	-	1	-	3	-	2	7	7	8
Female	190	-	-	-	-	-	1	2	3	8	11	14
Alabama												
All races, both sexes	104	-	-	-	-	-	1	2	3	4	4	10
Male	40	-	-	-	-	-	-	1	1	3	1	1
Female	64	-	-	-	-	-	1	1	2	1	3	9
White, both sexes	81	-	-	-	-	-	1	2	3	3	3	8
Male	28	-	-	-	-	-	-	1	1	2	1	-
Female	53	-	-	-	-	-	1	1	2	1	2	8
Black, both sexes	23	-	-	-	-	-	-	-	-	1	1	2
Male	12	-	-	-	-	-	-	-	-	1	-	1
Female	11	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Alaska												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	123	-	-	-	1	-	-	1	3	5	11	6
Male	48	-	-	-	-	-	-	1	1	2	6	3
Female	75	-	-	-	1	-	-	-	2	3	5	3
White, both sexes	111	-	-	-	1	-	-	1	2	4	9	6
Male	43	-	-	-	-	-	-	1	-	2	5	3
Female	68	-	-	-	1	-	-	-	2	2	4	3
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	1	1	2	-
Male	3	-	-	-	-	-	-	-	1	-	1	-
Female	4	-	-	-	-	-	-	-	-	1	1	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Arkansas												
All races, both sexes	52	-	-	-	-	-	1	1	2	2	4	5
Male	20	-	-	-	-	-	-	-	1	1	3	1
Female	32	-	-	-	-	-	1	1	1	1	1	4
White, both sexes	44	-	-	-	-	-	1	1	2	2	2	4
Male	18	-	-	-	-	-	-	-	1	1	2	1
Female	26	-	-	-	-	-	1	1	1	1	-	3
Black, both sexes	8	-	-	-	-	-	-	-	-	-	2	1
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	636	-	-	1	3	3	10	11	11	32	36	58
Male	255	-	-	1	2	1	8	3	7	22	17	30
Female	381	-	-	-	1	2	2	8	4	10	19	28
White, both sexes	473	-	-	1	2	3	5	8	7	24	28	41
Male	196	-	-	1	1	1	4	2	4	17	14	21
Female	277	-	-	-	1	2	1	6	3	7	14	20
Black, both sexes	50	-	-	-	-	-	1	2	2	2	3	10
Male	20	-	-	-	-	-	1	1	2	2	-	5
Female	30	-	-	-	-	-	-	1	-	-	3	5
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	111	-	-	-	1	-	4	1	2	6	5	6
Male	38	-	-	-	1	-	3	-	1	3	3	3
Female	73	-	-	-	-	-	1	1	1	3	2	3

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Colorado												
All races, both sexes	80	-	-	1	-	-	1	-	1	3	6	7
Male	27	-	-	1	-	-	-	-	1	-	1	2
Female	53	-	-	-	-	-	1	-	-	3	5	5
White, both sexes	68	-	-	1	-	-	1	-	1	3	3	7
Male	23	-	-	1	-	-	-	-	1	-	-	2
Female	45	-	-	-	-	-	1	-	-	3	3	5
Black, both sexes	9	-	-	-	-	-	-	-	-	-	3	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	71	-	-	-	-	-	-	2	1	4	4	2
Male	29	-	-	-	-	-	-	1	-	2	3	-
Female	42	-	-	-	-	-	-	1	1	2	1	2
White, both sexes	65	-	-	-	-	-	-	2	1	4	3	1
Male	27	-	-	-	-	-	-	1	-	2	3	-
Female	38	-	-	-	-	-	-	1	1	2	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Delaware												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	3
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	3
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Florida												
All races, both sexes	359	-	-	1	-	1	2	11	2	15	11	32
Male	132	-	-	1	-	1	2	6	-	7	4	9
Female	227	-	-	-	-	-	-	5	2	8	7	23
White, both sexes	304	-	-	1	-	1	2	6	2	14	5	23
Male	116	-	-	1	-	1	2	3	-	7	2	7
Female	188	-	-	-	-	-	-	3	2	7	3	16
Black, both sexes	45	-	-	-	-	-	-	5	-	1	6	7
Male	14	-	-	-	-	-	-	3	-	-	2	2
Female	31	-	-	-	-	-	-	2	-	1	4	5
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
Georgia												
All races, both sexes	166	-	-	1	1	-	2	1	7	6	14	26
Male	63	-	-	1	-	-	2	-	3	4	6	7
Female	103	-	-	-	1	-	-	1	4	2	8	19
White, both sexes	107	-	-	1	-	-	2	-	5	2	8	14
Male	39	-	-	1	-	-	2	-	2	1	2	5
Female	68	-	-	-	-	-	-	-	3	1	6	9
Black, both sexes	58	-	-	-	1	-	-	1	2	4	6	11
Male	24	-	-	-	-	-	-	-	1	3	4	2
Female	34	-	-	-	1	-	-	1	1	1	2	9
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Hawaii												
All races, both sexes	37	-	-	-	-	-	-	-	1	3	2	-
Male	12	-	-	-	-	-	-	-	1	1	1	-
Female	25	-	-	-	-	-	-	-	-	2	1	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	33	-	-	-	-	-	-	-	1	3	2	-
Male	11	-	-	-	-	-	-	-	1	1	1	-
Female	22	-	-	-	-	-	-	-	-	2	1	-
Idaho												
All races, both sexes	15	-	-	-	-	-	-	-	1	1	1	1
Male	6	-	-	-	-	-	-	-	-	-	1	1
Female	9	-	-	-	-	-	-	-	1	1	-	-
White, both sexes	14	-	-	-	-	-	-	-	1	1	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	9	-	-	-	-	-	-	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Illinois												
All races, both sexes	233	-	-	-	-	3	4	3	6	6	13	17
Male	83	-	-	-	-	2	2	1	3	2	6	4
Female	150	-	-	-	-	1	2	2	3	4	7	13
White, both sexes	192	-	-	-	-	3	4	2	5	3	13	12
Male	71	-	-	-	-	2	2	1	3	-	6	4
Female	121	-	-	-	-	1	2	1	2	3	7	8
Black, both sexes	34	-	-	-	-	-	-	1	-	3	-	5
Male	11	-	-	-	-	-	-	-	-	2	-	-
Female	23	-	-	-	-	-	-	1	-	1	-	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	1	-	-	-
Indiana												
All races, both sexes	119	-	-	-	-	-	2	3	3	3	4	10
Male	49	-	-	-	-	-	-	2	2	2	3	3
Female	70	-	-	-	-	-	2	1	1	1	1	7
White, both sexes	103	-	-	-	-	-	2	3	3	2	4	8
Male	43	-	-	-	-	-	-	2	2	2	3	3
Female	60	-	-	-	-	-	2	1	1	-	1	5
Black, both sexes	15	-	-	-	-	-	-	-	-	1	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	1	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Iowa												
All races, both sexes	56	-	-	-	-	-	-	-	-	2	4	6
Male	22	-	-	-	-	-	-	-	-	-	2	3
Female	34	-	-	-	-	-	-	-	-	2	2	3
White, both sexes	53	-	-	-	-	-	-	-	-	2	4	6
Male	21	-	-	-	-	-	-	-	-	-	2	3
Female	32	-	-	-	-	-	-	-	-	2	2	3
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	60	-	-	-	-	-	-	-	2	-	8	7
Male	13	-	-	-	-	-	-	-	1	-	2	3
Female	47	-	-	-	-	-	-	-	1	-	6	4
White, both sexes	54	-	-	-	-	-	-	-	1	-	8	6
Male	12	-	-	-	-	-	-	-	1	-	2	3
Female	42	-	-	-	-	-	-	-	-	-	6	3
Black, both sexes	5	-	-	-	-	-	-	-	1	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Kentucky												
All races, both sexes	76	-	-	-	-	-	-	1	1	4	5	3
Male	34	-	-	-	-	-	-	-	-	2	1	-
Female	42	-	-	-	-	-	-	1	1	2	4	3
White, both sexes	69	-	-	-	-	-	-	1	-	3	4	2
Male	31	-	-	-	-	-	-	-	-	1	1	-
Female	38	-	-	-	-	-	-	1	-	2	3	2
Black, both sexes	7	-	-	-	-	-	-	-	1	1	1	1
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	-	1	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	87	-	-	-	2	-	1	-	2	3	3	12
Male	27	-	-	-	1	-	1	-	-	-	1	2
Female	60	-	-	-	1	-	-	-	2	3	2	10
White, both sexes	56	-	-	-	2	-	1	-	1	1	2	4
Male	19	-	-	-	1	-	1	-	-	-	1	1
Female	37	-	-	-	1	-	-	-	1	1	1	3
Black, both sexes	28	-	-	-	-	-	-	-	1	1	1	8
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	21	-	-	-	-	-	-	-	1	1	1	7
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Maine												
All races, both sexes	26	-	-	-	-	-	-	-	-	-	1	3
Male	9	-	-	-	-	-	-	-	-	-	1	2
Female	17	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	26	-	-	-	-	-	-	-	-	-	1	3
Male	9	-	-	-	-	-	-	-	-	-	1	2
Female	17	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	100	-	-	-	-	1	-	1	2	5	17	11
Male	34	-	-	-	-	-	-	-	1	2	7	4
Female	66	-	-	-	-	1	-	1	1	3	10	7
White, both sexes	53	-	-	-	-	1	-	1	-	1	8	1
Male	16	-	-	-	-	-	-	-	-	-	3	-
Female	37	-	-	-	-	1	-	1	-	1	5	1
Black, both sexes	45	-	-	-	-	-	-	-	2	4	9	10
Male	18	-	-	-	-	-	-	-	1	2	4	4
Female	27	-	-	-	-	-	-	-	1	2	5	6
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Massachusetts												
All races, both sexes	104	-	-	-	-	1	-	1	3	2	5	12
Male	39	-	-	-	-	-	-	-	1	1	2	4
Female	65	-	-	-	-	1	-	1	2	1	3	8
White, both sexes	92	-	-	-	-	-	-	1	3	1	4	10
Male	33	-	-	-	-	-	-	-	1	-	2	3
Female	59	-	-	-	-	-	-	1	2	1	2	7
Black, both sexes	8	-	-	-	-	1	-	-	-	1	1	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	1	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	2
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	1
Michigan												
All races, both sexes	202	-	-	-	-	1	-	1	4	14	7	15
Male	74	-	-	-	-	1	-	-	1	5	2	5
Female	128	-	-	-	-	-	-	1	3	9	5	10
White, both sexes	164	-	-	-	-	1	-	1	2	7	4	12
Male	62	-	-	-	-	1	-	-	1	3	2	5
Female	102	-	-	-	-	-	-	1	1	4	2	7
Black, both sexes	33	-	-	-	-	-	-	-	2	6	2	3
Male	12	-	-	-	-	-	-	-	-	2	-	-
Female	21	-	-	-	-	-	-	-	2	4	2	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	1	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Minnesota												
All races, both sexes	96	-	-	-	1	-	1	2	-	3	6	12
Male	35	-	-	-	1	-	1	2	-	2	1	2
Female	61	-	-	-	-	-	-	-	-	1	5	10
White, both sexes	89	-	-	-	1	-	-	2	-	3	6	10
Male	31	-	-	-	1	-	-	2	-	2	1	1
Female	58	-	-	-	-	-	-	-	-	1	5	9
Black, both sexes	4	-	-	-	-	-	1	-	-	-	-	1
Male	2	-	-	-	-	-	1	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	45	-	-	-	1	-	-	2	-	3	6	5
Male	17	-	-	-	-	-	-	1	-	1	2	-
Female	28	-	-	-	1	-	-	1	-	2	4	5
White, both sexes	31	-	-	-	1	-	-	1	-	2	3	1
Male	10	-	-	-	-	-	-	1	-	-	1	-
Female	21	-	-	-	1	-	-	-	-	2	2	1
Black, both sexes	12	-	-	-	-	-	-	1	-	-	3	3
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	1	-	-	2	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Missouri												
All races, both sexes	122	-	-	-	-	-	-	1	2	5	13	16
Male	45	-	-	-	-	-	-	1	-	1	6	4
Female	77	-	-	-	-	-	-	-	2	4	7	12
White, both sexes	103	-	-	-	-	-	-	1	1	5	10	12
Male	40	-	-	-	-	-	-	1	-	1	5	4
Female	63	-	-	-	-	-	-	-	1	4	5	8
Black, both sexes	14	-	-	-	-	-	-	-	1	-	3	3
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	1	-	2	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Montana												
All races, both sexes	21	-	-	-	-	-	-	-	-	-	3	1
Male	11	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	19	-	-	-	-	-	-	-	-	-	2	1
Male	10	-	-	-	-	-	-	-	-	-	1	-
Female	9	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Nebraska												
All races, both sexes	33	-	-	-	-	1	-	1	1	1	3	2
Male	10	-	-	-	-	-	-	-	-	1	2	-
Female	23	-	-	-	-	1	-	1	1	-	1	2
White, both sexes	33	-	-	-	-	1	-	1	1	1	3	2
Male	10	-	-	-	-	-	-	-	-	1	2	-
Female	23	-	-	-	-	1	-	1	1	-	1	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	65	-	-	-	-	-	2	2	2	4	10	10
Male	28	-	-	-	-	-	1	1	1	1	5	6
Female	37	-	-	-	-	-	1	1	1	3	5	4
White, both sexes	46	-	-	-	-	-	2	1	-	3	6	8
Male	23	-	-	-	-	-	1	1	-	1	3	6
Female	23	-	-	-	-	-	1	-	-	2	3	2
Black, both sexes	9	-	-	-	-	-	-	-	2	1	2	1
Male	3	-	-	-	-	-	-	-	1	-	1	-
Female	6	-	-	-	-	-	-	-	1	1	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	1	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	1	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
New Hampshire												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	2	3
Male	10	-	-	-	-	-	-	-	-	-	1	3
Female	9	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	18	-	-	-	-	-	-	-	-	-	1	3
Male	10	-	-	-	-	-	-	-	-	-	1	3
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	169	-	-	1	2	-	-	2	1	4	10	14
Male	58	-	-	1	-	-	-	2	1	3	5	5
Female	111	-	-	-	2	-	-	-	-	1	5	9
White, both sexes	127	-	-	1	1	-	-	2	1	1	3	10
Male	48	-	-	1	-	-	-	2	1	1	2	5
Female	79	-	-	-	1	-	-	-	-	-	1	5
Black, both sexes	32	-	-	-	1	-	-	-	-	2	6	3
Male	7	-	-	-	-	-	-	-	-	1	3	-
Female	25	-	-	-	1	-	-	-	-	1	3	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	1	1	1
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	7	-	-	-	-	-	-	-	-	-	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
New Mexico												
All races, both sexes	44	1	-	-	-	-	-	-	2	1	1	4
Male	17	-	-	-	-	-	-	-	2	-	1	2
Female	27	1	-	-	-	-	-	-	-	1	-	2
White, both sexes	41	1	-	-	-	-	-	-	1	1	1	4
Male	16	-	-	-	-	-	-	-	1	-	1	2
Female	25	1	-	-	-	-	-	-	-	1	-	2
Black, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	380	-	-	1	2	6	2	2	5	12	24	35
Male	133	-	-	-	1	4	1	1	2	5	10	19
Female	247	-	-	1	1	2	1	1	3	7	14	16
White, both sexes	296	-	-	1	2	3	1	1	5	5	14	25
Male	106	-	-	-	1	2	-	1	2	1	7	14
Female	190	-	-	1	1	1	1	-	3	4	7	11
Black, both sexes	60	-	-	-	-	3	1	1	-	6	8	6
Male	18	-	-	-	-	2	1	-	-	3	2	2
Female	42	-	-	-	-	1	-	1	-	3	6	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	24	-	-	-	-	-	-	-	-	1	2	4
Male	9	-	-	-	-	-	-	-	-	1	1	3
Female	15	-	-	-	-	-	-	-	-	-	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
North Carolina												
All races, both sexes	175	-	-	1	-	-	1	4	5	10	4	20
Male	59	-	-	1	-	-	-	4	3	3	-	8
Female	116	-	-	-	-	-	1	-	2	7	4	12
White, both sexes	119	-	-	1	-	-	-	4	2	6	3	11
Male	41	-	-	1	-	-	-	4	1	3	-	4
Female	78	-	-	-	-	-	-	-	1	3	3	7
Black, both sexes	48	-	-	-	-	-	1	-	3	3	-	8
Male	16	-	-	-	-	-	-	-	2	-	-	3
Female	32	-	-	-	-	-	1	-	1	3	-	5
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	1	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
North Dakota												
All races, both sexes	12	-	-	-	-	-	-	-	-	1	1	2
Male	3	-	-	-	-	-	-	-	-	-	-	2
Female	9	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	12	-	-	-	-	-	-	-	-	1	1	2
Male	3	-	-	-	-	-	-	-	-	-	-	2
Female	9	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Ohio												
All races, both sexes	255	-	-	-	1	3	3	4	3	10	20	22
Male	91	-	-	-	1	1	2	2	2	4	7	8
Female	164	-	-	-	-	2	1	2	1	6	13	14
White, both sexes	207	-	-	-	1	2	2	3	2	5	14	17
Male	69	-	-	-	1	-	1	1	1	2	4	6
Female	138	-	-	-	-	2	1	2	1	3	10	11
Black, both sexes	44	-	-	-	-	1	1	1	1	4	4	5
Male	20	-	-	-	-	1	1	1	1	2	2	2
Female	24	-	-	-	-	-	-	-	-	2	2	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	1	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Oklahoma												
All races, both sexes	61	-	-	-	-	-	-	1	1	1	7	6
Male	27	-	-	-	-	-	-	-	1	-	-	5
Female	34	-	-	-	-	-	-	1	-	1	7	1
White, both sexes	50	-	-	-	-	-	-	-	1	1	6	5
Male	22	-	-	-	-	-	-	-	1	-	-	4
Female	28	-	-	-	-	-	-	-	-	1	6	1
Black, both sexes	7	-	-	-	-	-	-	1	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	1	-	-	1	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Oregon												
All races, both sexes	76	-	-	-	-	-	1	-	1	2	2	10
Male	34	-	-	-	-	-	1	-	1	2	2	4
Female	42	-	-	-	-	-	-	-	-	-	-	6
White, both sexes	69	-	-	-	-	-	-	-	1	2	2	10
Male	30	-	-	-	-	-	-	-	1	2	2	4
Female	39	-	-	-	-	-	-	-	-	-	-	6
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	274	-	-	-	-	-	-	2	2	4	11	25
Male	99	-	-	-	-	-	-	1	-	1	4	11
Female	175	-	-	-	-	-	-	1	2	3	7	14
White, both sexes	220	-	-	-	-	-	-	1	-	4	9	19
Male	80	-	-	-	-	-	-	1	-	1	4	9
Female	140	-	-	-	-	-	-	-	-	3	5	10
Black, both sexes	52	-	-	-	-	-	-	1	2	-	2	6
Male	18	-	-	-	-	-	-	-	-	-	-	2
Female	34	-	-	-	-	-	-	1	2	-	2	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Rhode Island												
All races, both sexes	20	-	-	-	-	-	-	-	-	1	3	2
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	15	-	-	-	-	-	-	-	-	1	2	2
White, both sexes	15	-	-	-	-	-	-	-	-	-	2	1
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	4	-	-	-	-	-	-	-	-	1	1	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	112	-	-	1	1	-	-	-	5	8	6	9
Male	46	-	-	-	-	-	-	-	4	2	4	6
Female	66	-	-	1	1	-	-	-	1	6	2	3
White, both sexes	77	-	-	-	1	-	-	-	2	1	4	7
Male	30	-	-	-	-	-	-	-	2	-	2	5
Female	47	-	-	-	1	-	-	-	-	1	2	2
Black, both sexes	35	-	-	1	-	-	-	-	3	7	2	2
Male	16	-	-	-	-	-	-	-	2	2	2	1
Female	19	-	-	1	-	-	-	-	1	5	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
South Dakota												
All races, both sexes	13	-	-	-	-	1	-	-	-	-	1	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	1	-	-	-	-	1	-
White, both sexes	11	-	-	-	-	1	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	1	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	142	-	-	-	-	-	2	2	1	6	13	15
Male	65	-	-	-	-	-	-	2	1	4	8	7
Female	77	-	-	-	-	-	2	-	-	2	5	8
White, both sexes	113	-	-	-	-	-	2	2	-	3	9	9
Male	49	-	-	-	-	-	-	2	-	2	5	4
Female	64	-	-	-	-	-	2	-	-	1	4	5
Black, both sexes	27	-	-	-	-	-	-	-	1	3	4	6
Male	15	-	-	-	-	-	-	-	1	2	3	3
Female	12	-	-	-	-	-	-	-	-	1	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Texas												
All races, both sexes	394	-	1	1	1	2	2	7	9	16	23	42
Male	126	-	-	1	1	1	1	1	2	8	11	19
Female	268	-	1	-	-	1	1	6	7	8	12	23
White, both sexes	328	-	-	-	1	2	2	5	5	13	17	32
Male	109	-	-	-	1	1	1	1	1	8	10	17
Female	219	-	-	-	-	1	1	4	4	5	7	15
Black, both sexes	53	-	1	1	-	-	-	2	3	3	4	9
Male	16	-	-	1	-	-	-	-	1	-	1	2
Female	37	-	1	-	-	-	-	2	2	3	3	7
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	1	-	2	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	1	-	2	1
Utah												
All races, both sexes	32	-	-	-	-	-	-	2	-	2	1	3
Male	11	-	-	-	-	-	-	1	-	1	1	1
Female	21	-	-	-	-	-	-	1	-	1	-	2
White, both sexes	31	-	-	-	-	-	-	2	-	2	1	3
Male	10	-	-	-	-	-	-	1	-	1	1	1
Female	21	-	-	-	-	-	-	1	-	1	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Vermont												
All races, both sexes	10	-	-	-	-	-	-	1	-	-	1	-
Male	5	-	-	-	-	-	-	1	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	10	-	-	-	-	-	-	1	-	-	1	-
Male	5	-	-	-	-	-	-	1	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	121	-	-	1	-	1	-	2	-	1	6	10
Male	36	-	-	1	-	1	-	2	-	1	1	4
Female	85	-	-	-	-	-	-	-	-	-	5	6
White, both sexes	92	-	-	1	-	1	-	2	-	-	4	8
Male	30	-	-	1	-	1	-	2	-	-	-	4
Female	62	-	-	-	-	-	-	-	-	-	4	4
Black, both sexes	19	-	-	-	-	-	-	-	-	-	2	1
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	16	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	1	-	1
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Washington												
All races, both sexes	108	-	-	-	1	2	-	1	1	4	7	8
Male	33	-	-	-	1	2	-	1	1	1	3	1
Female	75	-	-	-	-	-	-	-	-	3	4	7
White, both sexes	90	-	-	-	1	2	-	1	1	4	5	6
Male	28	-	-	-	1	2	-	1	1	1	3	1
Female	62	-	-	-	-	-	-	-	-	3	2	5
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	2	2
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	2	2
West Virginia												
All races, both sexes	55	-	-	-	-	-	-	-	3	1	2	4
Male	29	-	-	-	-	-	-	-	1	1	2	2
Female	26	-	-	-	-	-	-	-	2	-	-	2
White, both sexes	51	-	-	-	-	-	-	-	2	1	2	3
Male	28	-	-	-	-	-	-	-	1	1	2	1
Female	23	-	-	-	-	-	-	-	1	-	-	2
Black, both sexes	4	-	-	-	-	-	-	-	1	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Wisconsin												
All races, both sexes	104	-	-	-	-	-	1	-	2	3	5	11
Male	38	-	-	-	-	-	1	-	1	-	3	6
Female	66	-	-	-	-	-	-	-	1	3	2	5
White, both sexes	96	-	-	-	-	-	-	-	2	3	5	9
Male	34	-	-	-	-	-	-	-	1	-	3	5
Female	62	-	-	-	-	-	-	-	1	3	2	4
Black, both sexes	6	-	-	-	-	-	1	-	-	-	-	2
Male	3	-	-	-	-	-	1	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	12	-	-	-	-	-	-	-	1	1	1	2
Male	7	-	-	-	-	-	-	-	-	-	1	2
Female	5	-	-	-	-	-	-	-	1	1	-	-
White, both sexes	11	-	-	-	-	-	-	-	1	1	1	2
Male	6	-	-	-	-	-	-	-	-	-	1	2
Female	5	-	-	-	-	-	-	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
United States												
All races, both sexes	638	589	550	480	433	481	533	368	192	50	9	-
Male	256	251	214	165	150	165	142	107	48	6	3	-
Female	382	338	336	315	283	316	391	261	144	44	6	-
White, both sexes	481	456	458	404	356	412	466	317	170	42	8	-
Male	202	194	181	133	127	141	123	99	42	4	3	-
Female	279	262	277	271	229	271	343	218	128	38	5	-
Black, both sexes	130	88	62	54	49	40	37	31	10	7	1	-
Male	44	45	21	26	15	12	11	5	1	2	-	-
Female	86	43	41	28	34	28	26	26	9	5	1	-
American Indian 1/, both sexes	5	13	3	2	3	4	-	2	-	-	-	-
Male	2	7	1	1	-	3	-	1	-	-	-	-
Female	3	6	2	1	3	1	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	22	32	27	20	25	25	30	18	12	1	-	-
Male	8	5	11	5	8	9	8	2	5	-	-	-
Female	14	27	16	15	17	16	22	16	7	1	-	-
Alabama												
All races, both sexes	15	8	9	9	7	11	14	5	2	-	-	-
Male	8	3	4	5	4	3	4	-	2	-	-	-
Female	7	5	5	4	3	8	10	5	-	-	-	-
White, both sexes	9	7	7	5	5	10	11	5	2	-	-	-
Male	6	3	3	2	3	2	2	-	2	-	-	-
Female	3	4	4	3	2	8	9	5	-	-	-	-
Black, both sexes	6	1	2	4	2	1	3	-	-	-	-	-
Male	2	-	1	3	1	1	2	-	-	-	-	-
Female	4	1	1	1	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Alaska												
All races, both sexes	1	-	-	1	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	1	-	-	-
White, both sexes	1	-	-	1	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	8	18	15	14	8	11	6	11	5	-	-	-
Male	6	7	9	5	3	2	1	2	-	-	-	-
Female	2	11	6	9	5	9	5	9	5	-	-	-
White, both sexes	8	16	14	14	6	10	6	10	4	-	-	-
Male	6	6	9	5	2	2	1	1	-	-	-	-
Female	2	10	5	9	4	8	5	9	4	-	-	-
Black, both sexes	-	1	-	-	1	1	-	-	-	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Arkansas												
All races, both sexes	8	4	5	4	3	3	6	3	1	-	-	-
Male	5	3	-	1	1	1	1	2	-	-	-	-
Female	3	1	5	3	2	2	5	1	1	-	-	-
White, both sexes	6	3	4	4	3	2	6	3	1	-	-	-
Male	5	2	-	1	1	1	1	2	-	-	-	-
Female	1	1	4	3	2	1	5	1	1	-	-	-
Black, both sexes	2	1	1	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	2	-	1	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	69	71	69	49	40	55	52	45	20	1	-	-
Male	26	25	25	19	16	23	13	13	4	-	-	-
Female	43	46	44	30	24	32	39	32	16	1	-	-
White, both sexes	51	52	58	39	29	41	37	30	16	1	-	-
Male	17	22	22	16	11	18	11	11	3	-	-	-
Female	34	30	36	23	18	23	26	19	13	1	-	-
Black, both sexes	8	6	4	-	2	1	3	5	1	-	-	-
Male	4	3	1	-	1	-	-	-	-	-	-	-
Female	4	3	3	-	1	1	3	5	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	13	7	10	9	13	12	9	3	-	-	-
Male	5	-	2	3	4	5	2	2	1	-	-	-
Female	5	13	5	7	5	8	10	7	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Colorado												
All races, both sexes	11	13	6	3	8	7	5	5	2	-	1	-
Male	6	5	1	1	2	3	1	2	1	-	-	-
Female	5	8	5	2	6	4	4	3	1	-	1	-
White, both sexes	10	12	4	3	7	5	5	4	1	-	1	-
Male	5	5	1	1	2	2	1	2	-	-	-	-
Female	5	7	3	2	5	3	4	2	1	-	1	-
Black, both sexes	1	1	1	-	1	1	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	1	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	2	5	7	6	7	8	13	7	3	-	-	-
Male	2	1	4	-	1	5	6	2	2	-	-	-
Female	-	4	3	6	6	3	7	5	1	-	-	-
White, both sexes	2	4	6	6	6	7	13	7	3	-	-	-
Male	2	1	3	-	1	4	6	2	2	-	-	-
Female	-	3	3	6	5	3	7	5	1	-	-	-
Black, both sexes	-	1	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Delaware												
All races, both sexes	3	2	1	-	2	1	3	-	1	-	-	-
Male	-	2	-	-	-	1	-	-	-	-	-	-
Female	3	-	1	-	2	-	3	-	1	-	-	-
White, both sexes	2	2	1	-	2	1	3	-	1	-	-	-
Male	-	2	-	-	-	1	-	-	-	-	-	-
Female	2	-	1	-	2	-	3	-	1	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	3	3	1	1	-	1	-	2	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	-	2	1	1	-	1	-	2	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	3	1	1	-	1	-	2	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	-	2	1	1	-	1	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Florida												
All races, both sexes	29	33	32	39	23	36	45	27	16	4	-	-
Male	13	13	12	16	7	14	14	6	7	-	-	-
Female	16	20	20	23	16	22	31	21	9	4	-	-
White, both sexes	23	30	29	33	18	32	44	24	14	3	-	-
Male	12	12	10	12	6	14	14	6	7	-	-	-
Female	11	18	19	21	12	18	30	18	7	3	-	-
Black, both sexes	6	3	1	5	3	2	1	2	2	1	-	-
Male	1	1	1	3	1	-	-	-	-	-	-	-
Female	5	2	-	2	2	2	1	2	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	1	2	1	-	1	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	2	1	-	1	-	-	-	-
Georgia												
All races, both sexes	18	14	19	13	12	7	13	7	2	3	-	-
Male	8	7	8	7	3	3	2	2	-	-	-	-
Female	10	7	11	6	9	4	11	5	2	3	-	-
White, both sexes	10	9	13	9	10	3	12	6	1	2	-	-
Male	4	5	6	4	3	-	2	2	-	-	-	-
Female	6	4	7	5	7	3	10	4	1	2	-	-
Black, both sexes	8	5	6	4	2	4	1	1	1	1	-	-
Male	4	2	2	3	-	3	-	-	-	-	-	-
Female	4	3	4	1	2	1	1	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Hawaii												
All races, both sexes	-	6	2	1	3	5	9	4	1	-	-	-
Male	-	3	1	-	-	1	3	-	1	-	-	-
Female	-	3	1	1	3	4	6	4	-	-	-	-
White, both sexes	-	2	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	2	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	3	2	-	3	5	9	4	1	-	-	-
Male	-	2	1	-	-	1	3	-	1	-	-	-
Female	-	1	1	-	3	4	6	4	-	-	-	-
Idaho												
All races, both sexes	2	1	-	2	1	4	-	-	1	-	-	-
Male	1	-	-	-	1	2	-	-	-	-	-	-
Female	1	1	-	2	-	2	-	-	1	-	-	-
White, both sexes	2	1	-	2	1	4	-	-	1	-	-	-
Male	1	-	-	-	1	2	-	-	-	-	-	-
Female	1	1	-	2	-	2	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Illinois												
All races, both sexes	32	23	19	23	18	22	19	15	7	3	-	-
Male	10	12	6	11	5	7	3	8	-	1	-	-
Female	22	11	13	12	13	15	16	7	7	2	-	-
White, both sexes	24	18	19	17	13	20	15	14	7	3	-	-
Male	8	9	6	8	4	7	2	8	-	1	-	-
Female	16	9	13	9	9	13	13	6	7	2	-	-
Black, both sexes	8	3	-	4	5	1	3	1	-	-	-	-
Male	2	2	-	3	1	-	1	-	-	-	-	-
Female	6	1	-	1	4	1	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	2	-	1	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	2	-	1	1	-	-	-	-	-
Indiana												
All races, both sexes	19	11	15	7	10	7	8	11	4	2	-	-
Male	9	4	8	1	4	5	1	5	-	-	-	-
Female	10	7	7	6	6	2	7	6	4	2	-	-
White, both sexes	15	10	12	7	9	7	7	9	3	2	-	-
Male	7	4	6	1	3	5	1	4	-	-	-	-
Female	8	6	6	6	6	2	6	5	3	2	-	-
Black, both sexes	4	1	3	-	-	-	1	2	1	-	-	-
Male	2	-	2	-	-	-	-	1	-	-	-	-
Female	2	1	1	-	-	-	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Iowa												
All races, both sexes	9	4	5	6	3	4	3	7	3	-	-	-
Male	2	3	2	4	-	2	1	2	1	-	-	-
Female	7	1	3	2	3	2	2	5	2	-	-	-
White, both sexes	8	3	4	6	3	4	3	7	3	-	-	-
Male	2	3	1	4	-	2	1	2	1	-	-	-
Female	6	-	3	2	3	2	2	5	2	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	2	11	7	5	6	4	1	5	2	-	-	-
Male	-	4	1	1	1	-	-	-	-	-	-	-
Female	2	7	6	4	5	4	1	5	2	-	-	-
White, both sexes	2	10	6	4	5	4	1	5	2	-	-	-
Male	-	3	1	1	1	-	-	-	-	-	-	-
Female	2	7	5	3	4	4	1	5	2	-	-	-
Black, both sexes	-	1	1	1	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Kentucky												
All races, both sexes	5	10	9	10	7	9	8	2	1	1	-	-
Male	3	6	6	6	4	3	3	-	-	-	-	-
Female	2	4	3	4	3	6	5	2	1	1	-	-
White, both sexes	4	10	9	10	6	9	7	2	1	1	-	-
Male	2	6	6	6	4	3	2	-	-	-	-	-
Female	2	4	3	4	2	6	5	2	1	1	-	-
Black, both sexes	1	-	-	-	1	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	5	11	13	4	10	11	5	1	3	1	-	-
Male	1	6	6	-	3	3	2	-	1	-	-	-
Female	4	5	7	4	7	8	3	1	2	1	-	-
White, both sexes	3	5	10	3	8	8	4	1	2	1	-	-
Male	1	2	5	-	3	2	1	-	1	-	-	-
Female	2	3	5	3	5	6	3	1	1	1	-	-
Black, both sexes	2	5	3	1	2	3	1	-	-	-	-	-
Male	-	3	1	-	-	1	1	-	-	-	-	-
Female	2	2	2	1	2	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Maine												
All races, both sexes	4	2	1	4	3	3	3	-	2	-	-	-
Male	1	1	-	1	1	1	-	-	1	-	-	-
Female	3	1	1	3	2	2	3	-	1	-	-	-
White, both sexes	4	2	1	4	3	3	3	-	2	-	-	-
Male	1	1	-	1	1	1	-	-	1	-	-	-
Female	3	1	1	3	2	2	3	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	8	10	8	7	8	7	8	3	3	-	1	-
Male	4	3	2	1	4	4	1	1	-	-	-	-
Female	4	7	6	6	4	3	7	2	3	-	1	-
White, both sexes	5	5	5	6	5	4	6	2	3	-	-	-
Male	3	1	2	1	3	2	-	1	-	-	-	-
Female	2	4	3	5	2	2	6	1	3	-	-	-
Black, both sexes	3	4	2	1	3	3	2	1	-	-	1	-
Male	1	2	-	-	1	2	1	-	-	-	-	-
Female	2	2	2	1	2	1	1	1	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Massachusetts												
All races, both sexes	10	5	8	5	12	11	9	11	5	3	1	-
Male	3	3	2	3	6	4	3	4	3	-	-	-
Female	7	2	6	2	6	7	6	7	2	3	1	-
White, both sexes	8	5	8	3	10	10	9	11	5	3	1	-
Male	2	3	2	1	5	4	3	4	3	-	-	-
Female	6	2	6	2	5	6	6	7	2	3	1	-
Black, both sexes	-	-	-	2	2	1	-	-	-	-	-	-
Male	-	-	-	2	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	28	22	22	24	10	13	20	11	9	1	-	-
Male	9	10	11	10	5	3	6	3	2	1	-	-
Female	19	12	11	14	5	10	14	8	7	-	-	-
White, both sexes	22	17	19	23	9	12	18	10	7	-	-	-
Male	8	6	10	9	5	3	5	3	1	-	-	-
Female	14	11	9	14	4	9	13	7	6	-	-	-
Black, both sexes	4	5	3	1	1	1	2	1	1	1	-	-
Male	1	4	1	1	-	-	1	-	1	1	-	-
Female	3	1	2	-	1	1	1	1	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Minnesota												
All races, both sexes	12	10	5	6	4	6	12	8	8	-	-	-
Male	7	5	2	3	1	2	3	1	2	-	-	-
Female	5	5	3	3	3	4	9	7	6	-	-	-
White, both sexes	10	9	5	6	4	5	12	8	8	-	-	-
Male	6	4	2	3	1	2	3	1	2	-	-	-
Female	4	5	3	3	3	3	9	7	6	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Mississippi												
All races, both sexes	7	4	3	2	-	3	6	1	2	-	-	-
Male	4	4	1	1	-	2	1	-	-	-	-	-
Female	3	-	2	1	-	1	5	1	2	-	-	-
White, both sexes	6	2	3	1	-	2	6	1	2	-	-	-
Male	3	2	1	-	-	1	1	-	-	-	-	-
Female	3	-	2	1	-	1	5	1	2	-	-	-
Black, both sexes	1	2	-	1	-	1	-	-	-	-	-	-
Male	1	2	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Missouri												
All races, both sexes	12	8	11	8	10	8	15	4	5	4	-	-
Male	7	4	5	2	6	1	5	1	1	1	-	-
Female	5	4	6	6	4	7	10	3	4	3	-	-
White, both sexes	10	6	11	6	8	6	14	4	5	4	-	-
Male	6	4	5	1	5	-	5	1	1	1	-	-
Female	4	2	6	5	3	6	9	3	4	3	-	-
Black, both sexes	2	-	-	2	2	1	-	-	-	-	-	-
Male	1	-	-	1	1	-	-	-	-	-	-	-
Female	1	-	-	1	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	2	-	-	-	-	1	-	-	-	-	-
Montana												
All races, both sexes	3	2	1	5	2	1	-	2	1	-	-	-
Male	-	2	1	4	1	-	-	2	-	-	-	-
Female	3	-	-	1	1	1	-	-	1	-	-	-
White, both sexes	3	2	1	4	2	1	-	2	1	-	-	-
Male	-	2	1	3	1	-	-	2	-	-	-	-
Female	3	-	-	1	1	1	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Nebraska												
All races, both sexes	4	6	2	3	2	2	1	2	-	1	1	-
Male	2	3	-	-	-	1	-	1	-	-	-	-
Female	2	3	2	3	2	1	1	1	-	1	1	-
White, both sexes	4	6	2	3	2	2	1	2	-	1	1	-
Male	2	3	-	-	-	1	-	1	-	-	-	-
Female	2	3	2	3	2	1	1	1	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	4	5	8	2	3	5	5	2	-	1	-	-
Male	2	1	2	1	1	2	3	1	-	-	-	-
Female	2	4	6	1	2	3	2	1	-	1	-	-
White, both sexes	3	4	6	1	3	4	3	1	-	1	-	-
Male	2	1	2	1	1	1	2	1	-	-	-	-
Female	1	3	4	-	2	3	1	-	-	1	-	-
Black, both sexes	-	1	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	2	1	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	2	1	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
New Hampshire												
All races, both sexes	1	4	1	2	1	2	1	-	-	2	-	-
Male	-	2	1	1	1	1	-	-	-	-	-	-
Female	1	2	-	1	-	1	1	-	-	2	-	-
White, both sexes	1	4	1	2	1	2	1	-	-	2	-	-
Male	-	2	1	1	1	1	-	-	-	-	-	-
Female	1	2	-	1	-	1	1	-	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	25	15	19	9	18	17	18	8	2	3	1	-
Male	11	5	7	1	8	4	2	2	-	1	-	-
Female	14	10	12	8	10	13	16	6	2	2	1	-
White, both sexes	18	11	12	4	17	17	15	8	2	3	1	-
Male	10	4	4	1	8	4	2	2	-	1	-	-
Female	8	7	8	3	9	13	13	6	2	2	1	-
Black, both sexes	6	3	4	3	1	-	3	-	-	-	-	-
Male	1	1	1	-	-	-	-	-	-	-	-	-
Female	5	2	3	3	1	-	3	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	3	2	-	-	-	-	-	-	-	-
Male	-	-	2	-	-	-	-	-	-	-	-	-
Female	1	1	1	2	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
New Mexico												
All races, both sexes	5	4	3	10	2	5	2	3	1	-	-	-
Male	1	1	2	3	1	2	-	2	-	-	-	-
Female	4	3	1	7	1	3	2	1	1	-	-	-
White, both sexes	5	3	3	9	2	5	2	3	1	-	-	-
Male	1	1	2	3	1	2	-	2	-	-	-	-
Female	4	2	1	6	1	3	2	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	42	34	34	33	29	31	34	33	15	5	1	-
Male	12	13	13	8	10	11	9	7	5	1	1	-
Female	30	21	21	25	19	20	25	26	10	4	-	-
White, both sexes	32	25	28	27	21	27	29	31	14	4	1	-
Male	11	9	12	7	7	10	8	7	5	1	1	-
Female	21	16	16	20	14	17	21	24	9	3	-	-
Black, both sexes	7	6	4	5	4	4	3	1	1	-	-	-
Male	1	3	-	1	2	1	-	-	-	-	-	-
Female	6	3	4	4	2	3	3	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	3	2	1	4	-	2	1	-	1	-	-
Male	-	1	1	-	1	-	1	-	-	-	-	-
Female	3	2	1	1	3	-	1	1	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
North Carolina												
All races, both sexes	22	21	19	17	12	11	10	16	2	-	-	-
Male	7	11	7	7	3	1	-	4	-	-	-	-
Female	15	10	12	10	9	10	10	12	2	-	-	-
White, both sexes	12	13	12	12	10	7	10	14	2	-	-	-
Male	4	6	6	5	3	1	-	3	-	-	-	-
Female	8	7	6	7	7	6	10	11	2	-	-	-
Black, both sexes	9	6	6	5	2	3	-	2	-	-	-	-
Male	3	4	1	2	-	-	-	1	-	-	-	-
Female	6	2	5	3	2	3	-	1	-	-	-	-
American Indian 1/, both sexes	1	1	1	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
North Dakota												
All races, both sexes	3	-	1	1	1	-	1	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	1	1	1	-	1	1	-	-	-	-
White, both sexes	3	-	1	1	1	-	1	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	1	1	1	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Ohio												
All races, both sexes	26	25	21	19	21	16	25	23	10	3	-	-
Male	8	13	11	5	7	4	7	5	4	-	-	-
Female	18	12	10	14	14	12	18	18	6	3	-	-
White, both sexes	19	22	16	17	19	15	23	20	9	1	-	-
Male	7	11	7	4	6	4	7	4	3	-	-	-
Female	12	11	9	13	13	11	16	16	6	1	-	-
Black, both sexes	7	3	5	2	2	1	2	3	-	2	-	-
Male	1	2	4	1	1	-	-	1	-	-	-	-
Female	6	1	1	1	1	1	2	2	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	6	3	9	6	5	11	2	1	2	-	-	-
Male	5	1	3	4	2	6	-	-	-	-	-	-
Female	1	2	6	2	3	5	2	1	2	-	-	-
White, both sexes	5	2	8	4	5	8	2	1	2	-	-	-
Male	4	1	3	3	2	4	-	-	-	-	-	-
Female	1	1	5	1	3	4	2	1	2	-	-	-
Black, both sexes	-	-	1	2	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	2	-	-	-	-	-	-
Male	1	-	-	-	-	2	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Oregon												
All races, both sexes	11	8	10	4	6	10	7	1	3	-	-	-
Male	6	2	2	2	3	7	1	1	-	-	-	-
Female	5	6	8	2	3	3	6	-	3	-	-	-
White, both sexes	10	7	9	3	5	10	6	1	3	-	-	-
Male	5	1	2	2	3	7	-	1	-	-	-	-
Female	5	6	7	1	2	3	6	-	3	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	1	1	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	1	1	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	35	31	19	18	24	27	38	24	11	2	1	-
Male	14	14	7	4	11	6	11	10	4	-	1	-
Female	21	17	12	14	13	21	27	14	7	2	-	-
White, both sexes	23	20	16	15	20	25	36	20	9	2	1	-
Male	10	9	5	2	9	6	11	9	3	-	1	-
Female	13	11	11	13	11	19	25	11	6	2	-	-
Black, both sexes	12	10	3	3	4	2	2	4	1	-	-	-
Male	4	5	2	2	2	-	-	1	-	-	-	-
Female	8	5	1	1	2	2	2	3	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Rhode Island												
All races, both sexes	3	2	2	1	1	-	3	-	2	-	-	-
Male	1	-	2	-	1	-	-	-	-	-	-	-
Female	2	2	-	1	-	-	3	-	2	-	-	-
White, both sexes	2	2	1	1	1	-	3	-	2	-	-	-
Male	1	-	1	-	1	-	-	-	-	-	-	-
Female	1	2	-	1	-	-	3	-	2	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	16	7	8	11	7	8	10	8	5	2	-	-
Male	3	3	5	5	2	1	6	4	1	-	-	-
Female	13	4	3	6	5	7	4	4	4	2	-	-
White, both sexes	10	4	6	9	5	7	8	6	5	2	-	-
Male	1	1	4	4	2	1	4	3	1	-	-	-
Female	9	3	2	5	3	6	4	3	4	2	-	-
Black, both sexes	6	3	2	2	2	1	2	2	-	-	-	-
Male	2	2	1	1	-	-	2	1	-	-	-	-
Female	4	1	1	1	2	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
South Dakota												
All races, both sexes	2	2	-	1	-	-	2	1	2	-	-	-
Male	2	1	-	-	-	-	1	1	-	-	-	-
Female	-	1	-	1	-	-	1	-	2	-	-	-
White, both sexes	1	1	-	1	-	-	2	1	2	-	-	-
Male	1	-	-	-	-	-	1	1	-	-	-	-
Female	-	1	-	1	-	-	1	-	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	19	14	10	20	8	7	10	7	5	1	2	-
Male	8	8	5	7	5	3	3	1	2	-	1	-
Female	11	6	5	13	3	4	7	6	3	1	1	-
White, both sexes	13	12	9	18	6	7	9	6	5	1	2	-
Male	6	6	5	6	3	3	3	1	2	-	1	-
Female	7	6	4	12	3	4	6	5	3	1	1	-
Black, both sexes	5	1	1	2	2	-	1	1	-	-	-	-
Male	2	1	-	1	2	-	-	-	-	-	-	-
Female	3	-	1	1	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Texas												
All races, both sexes	47	47	42	26	33	37	30	14	10	4	-	-
Male	19	17	10	5	5	12	6	6	1	-	-	-
Female	28	30	32	21	28	25	24	8	9	4	-	-
White, both sexes	33	40	36	24	31	35	27	13	8	4	-	-
Male	13	15	9	5	5	10	5	6	1	-	-	-
Female	20	25	27	19	26	25	22	7	7	4	-	-
Black, both sexes	13	5	5	2	-	2	1	1	1	-	-	-
Male	5	2	1	-	-	2	1	-	-	-	-	-
Female	8	3	4	2	-	-	-	1	1	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	-	2	-	2	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	2	-	2	-	1	-	-	-
Utah												
All races, both sexes	3	2	1	6	4	3	2	2	1	-	-	-
Male	2	1	-	-	1	1	1	1	-	-	-	-
Female	1	1	1	6	3	2	1	1	1	-	-	-
White, both sexes	3	1	1	6	4	3	2	2	1	-	-	-
Male	2	-	-	-	1	1	1	1	-	-	-	-
Female	1	1	1	6	3	2	1	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Vermont												
All races, both sexes	1	1	-	-	2	-	3	-	1	-	-	-
Male	1	-	-	-	1	-	2	-	-	-	-	-
Female	-	1	-	-	1	-	1	-	1	-	-	-
White, both sexes	1	1	-	-	2	-	3	-	1	-	-	-
Male	1	-	-	-	1	-	2	-	-	-	-	-
Female	-	1	-	-	1	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	13	18	17	8	11	8	12	9	3	1	-	-
Male	5	6	6	2	2	2	1	1	-	1	-	-
Female	8	12	11	6	9	6	11	8	3	-	-	-
White, both sexes	12	14	14	6	7	5	10	7	1	-	-	-
Male	5	6	5	2	1	1	1	1	-	-	-	-
Female	7	8	9	4	6	4	9	6	1	-	-	-
Black, both sexes	1	3	1	1	3	3	1	1	1	1	-	-
Male	-	-	-	-	-	1	-	-	-	1	-	-
Female	1	3	1	1	3	2	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	2	1	1	-	1	1	1	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	1	1	1	-	-	1	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Washington												
All races, both sexes	15	13	10	10	7	12	8	3	5	1	-	-
Male	2	3	5	2	2	3	4	-	2	-	-	-
Female	13	10	5	8	5	9	4	3	3	1	-	-
White, both sexes	13	8	8	9	4	12	7	3	5	1	-	-
Male	2	3	3	1	1	3	3	-	2	-	-	-
Female	11	5	5	8	3	9	4	3	3	1	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	3	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	1	3	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	1	2	-	1	-	-	-	-	-
Male	-	-	1	1	1	-	1	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	3	3	8	7	4	4	12	4	-	-	-	-
Male	3	2	4	3	1	2	6	2	-	-	-	-
Female	-	1	4	4	3	2	6	2	-	-	-	-
White, both sexes	3	3	8	7	4	4	10	4	-	-	-	-
Male	3	2	4	3	1	2	6	2	-	-	-	-
Female	-	1	4	4	3	2	4	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Wisconsin												
All races, both sexes	7	12	12	6	12	7	14	9	2	1	-	-
Male	3	6	4	2	3	1	5	2	1	-	-	-
Female	4	6	8	4	9	6	9	7	1	1	-	-
White, both sexes	7	11	11	6	11	7	14	8	2	-	-	-
Male	3	5	3	2	3	1	5	2	1	-	-	-
Female	4	6	8	4	8	6	9	6	1	-	-	-
Black, both sexes	-	1	-	-	1	-	-	-	-	1	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Wyoming												
All races, both sexes	-	1	1	2	3	-	-	-	-	-	-	-
Male	-	1	1	1	1	-	-	-	-	-	-	-
Female	-	-	-	1	2	-	-	-	-	-	-	-
White, both sexes	-	-	1	2	3	-	-	-	-	-	-	-
Male	-	-	1	1	1	-	-	-	-	-	-	-
Female	-	-	-	1	2	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
United States												
All races, both sexes	27,453	52	13	11	14	23	37	53	120	306	580	1,149
Male	12,784	29	6	3	5	15	21	30	65	183	330	673
Female	14,669	23	7	8	9	8	16	23	55	123	250	476
White, both sexes	22,305	37	11	6	9	15	22	34	73	192	344	693
Male	10,215	19	5	2	5	10	14	18	38	111	208	415
Female	12,090	18	6	4	4	5	8	16	35	81	136	278
Black, both sexes	3,942	13	1	3	3	7	14	15	40	93	204	386
Male	1,967	10	1	-	-	5	7	11	22	59	102	212
Female	1,975	3	-	3	3	2	7	4	18	34	102	174
American Indian 1/, both sexes	129	2	-	-	-	-	-	2	-	4	6	12
Male	57	-	-	-	-	-	-	1	-	2	4	8
Female	72	2	-	-	-	-	-	1	-	2	2	4
Asian or Pacific Islander 2/, both sexes	1,077	-	1	2	2	1	1	2	7	17	26	58
Male	545	-	-	1	-	-	-	-	5	11	16	38
Female	532	-	1	1	2	1	1	2	2	6	10	20
Alabama												
All races, both sexes	614	-	-	-	-	1	2	1	6	10	17	30
Male	295	-	-	-	-	1	1	1	-	4	8	13
Female	319	-	-	-	-	-	1	-	6	6	9	17
White, both sexes	428	-	-	-	-	-	1	-	4	3	4	10
Male	211	-	-	-	-	-	-	-	-	1	2	3
Female	217	-	-	-	-	-	1	-	4	2	2	7
Black, both sexes	182	-	-	-	-	1	1	1	1	7	13	20
Male	83	-	-	-	-	1	1	1	-	3	6	10
Female	99	-	-	-	-	-	-	-	1	4	7	10
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Alaska												
All races, both sexes	31	-	-	-	-	-	-	-	-	-	1	5
Male	19	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	1	4
White, both sexes	20	-	-	-	-	-	-	-	-	-	-	2
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	1	2
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	2
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Arizona												
All races, both sexes	385	1	-	-	-	-	2	-	1	3	5	13
Male	199	1	-	-	-	-	-	-	1	2	3	11
Female	186	-	-	-	-	-	2	-	-	1	2	2
White, both sexes	349	1	-	-	-	-	1	-	1	3	5	9
Male	179	1	-	-	-	-	-	-	1	2	3	7
Female	170	-	-	-	-	-	1	-	-	1	2	2
Black, both sexes	11	-	-	-	-	-	1	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	3
Male	5	-	-	-	-	-	-	-	-	-	-	3
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Arkansas												
All races, both sexes	340	1	-	-	-	-	-	1	2	5	13	13
Male	151	1	-	-	-	-	-	1	1	4	9	8
Female	189	-	-	-	-	-	-	-	1	1	4	5
White, both sexes	275	1	-	-	-	-	-	1	1	3	7	6
Male	122	1	-	-	-	-	-	1	1	2	4	4
Female	153	-	-	-	-	-	-	-	-	1	3	2
Black, both sexes	61	-	-	-	-	-	-	-	1	2	4	7
Male	28	-	-	-	-	-	-	-	-	2	4	4
Female	33	-	-	-	-	-	-	-	1	-	-	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	2	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
California												
All races, both sexes	2,864	6	-	2	2	2	3	6	9	44	75	143
Male	1,379	3	-	1	-	1	1	3	7	31	51	103
Female	1,485	3	-	1	2	1	2	3	2	13	24	40
White, both sexes	2,134	5	-	1	2	1	2	6	7	31	51	90
Male	1,003	2	-	1	-	1	1	3	5	24	36	66
Female	1,131	3	-	-	2	-	1	3	2	7	15	24
Black, both sexes	256	1	-	-	-	-	1	-	1	3	10	21
Male	132	1	-	-	-	-	-	-	1	2	5	14
Female	124	-	-	-	-	-	1	-	-	1	5	7
American Indian 1/, both sexes	14	-	-	-	-	-	-	-	-	-	1	2
Male	8	-	-	-	-	-	-	-	-	-	1	1
Female	6	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	460	-	-	1	-	1	-	-	1	10	13	30
Male	236	-	-	-	-	-	-	-	1	5	9	22
Female	224	-	-	1	-	1	-	-	-	5	4	8

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Colorado												
All races, both sexes	333	4	1	-	-	-	-	-	-	3	8	5
Male	157	2	-	-	-	-	-	-	-	2	5	1
Female	176	2	1	-	-	-	-	-	-	1	3	4
White, both sexes	306	3	1	-	-	-	-	-	-	3	7	3
Male	145	1	-	-	-	-	-	-	-	2	4	1
Female	161	2	1	-	-	-	-	-	-	1	3	2
Black, both sexes	14	1	-	-	-	-	-	-	-	-	1	1
Male	8	1	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	1
Connecticut												
All races, both sexes	295	-	-	-	-	-	1	-	2	3	2	5
Male	131	-	-	-	-	-	1	-	1	2	-	4
Female	164	-	-	-	-	-	-	-	1	1	2	1
White, both sexes	266	-	-	-	-	-	1	-	2	1	2	5
Male	120	-	-	-	-	-	1	-	1	1	-	4
Female	146	-	-	-	-	-	-	-	1	-	2	1
Black, both sexes	23	-	-	-	-	-	-	-	-	2	-	-
Male	9	-	-	-	-	-	-	-	-	1	-	-
Female	14	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Delaware												
All races, both sexes	78	-	-	-	-	-	1	-	1	1	1	4
Male	29	-	-	-	-	-	1	-	1	-	1	4
Female	49	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	60	-	-	-	-	-	1	-	-	1	-	-
Male	18	-	-	-	-	-	1	-	-	-	-	-
Female	42	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	16	-	-	-	-	-	-	-	1	-	-	4
Male	9	-	-	-	-	-	-	-	1	-	-	4
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	64	-	-	-	-	-	-	-	-	1	1	8
Male	21	-	-	-	-	-	-	-	-	-	-	6
Female	43	-	-	-	-	-	-	-	-	1	1	2
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	47	-	-	-	-	-	-	-	-	1	1	7
Male	18	-	-	-	-	-	-	-	-	-	-	6
Female	29	-	-	-	-	-	-	-	-	1	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Florida												
All races, both sexes	1,800	5	-	1	-	2	4	3	9	22	46	86
Male	858	3	-	-	-	2	3	1	6	15	27	43
Female	942	2	-	1	-	-	1	2	3	7	19	43
White, both sexes	1,493	4	-	1	-	1	2	2	3	9	35	57
Male	705	2	-	-	-	1	2	1	2	5	19	28
Female	788	2	-	1	-	-	-	1	1	4	16	29
Black, both sexes	288	1	-	-	-	1	2	1	6	13	11	28
Male	144	1	-	-	-	1	1	-	4	10	8	15
Female	144	-	-	-	-	-	1	1	2	3	3	13
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
Georgia												
All races, both sexes	830	-	1	-	3	2	1	2	4	14	23	58
Male	397	-	-	-	1	1	-	-	1	7	10	24
Female	433	-	1	-	2	1	1	2	3	7	13	34
White, both sexes	532	-	1	-	2	1	1	1	1	5	10	24
Male	252	-	-	-	1	1	-	-	1	1	8	11
Female	280	-	1	-	1	-	1	1	-	4	2	13
Black, both sexes	280	-	-	-	-	1	-	1	3	8	13	33
Male	138	-	-	-	-	-	-	-	-	6	2	12
Female	142	-	-	-	-	1	-	1	3	2	11	21
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	-	-	-	1	-	-	-	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	1	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Hawaii												
All races, both sexes	168	-	-	-	-	-	-	-	-	2	3	4
Male	84	-	-	-	-	-	-	-	-	1	2	4
Female	84	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	33	-	-	-	-	-	-	-	-	1	-	-
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	133	-	-	-	-	-	-	-	-	1	3	4
Male	62	-	-	-	-	-	-	-	-	1	2	4
Female	71	-	-	-	-	-	-	-	-	-	1	-
Idaho												
All races, both sexes	103	1	-	-	-	1	-	-	-	1	-	8
Male	51	1	-	-	-	-	-	-	-	-	-	2
Female	52	-	-	-	-	1	-	-	-	1	-	6
White, both sexes	101	1	-	-	-	1	-	-	-	1	-	8
Male	50	1	-	-	-	-	-	-	-	-	-	2
Female	51	-	-	-	-	1	-	-	-	1	-	6
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Illinois												
All races, both sexes	1,063	3	1	-	1	-	-	2	5	12	23	49
Male	484	1	1	-	1	-	-	1	2	10	15	22
Female	579	2	-	-	-	-	-	1	3	2	8	27
White, both sexes	869	1	-	-	1	-	-	2	2	8	7	24
Male	386	-	-	-	1	-	-	1	1	7	3	12
Female	483	1	-	-	-	-	-	1	1	1	4	12
Black, both sexes	172	2	1	-	-	-	-	-	3	3	15	23
Male	86	1	1	-	-	-	-	-	1	2	12	9
Female	86	1	-	-	-	-	-	-	2	1	3	14
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	19	-	-	-	-	-	-	-	-	1	1	2
Male	11	-	-	-	-	-	-	-	-	1	-	1
Female	8	-	-	-	-	-	-	-	-	-	1	1
Indiana												
All races, both sexes	573	-	1	-	-	1	-	-	2	6	14	22
Male	228	-	1	-	-	1	-	-	-	4	7	11
Female	345	-	-	-	-	-	-	-	2	2	7	11
White, both sexes	520	-	1	-	-	1	-	-	2	5	10	17
Male	203	-	1	-	-	1	-	-	-	3	6	10
Female	317	-	-	-	-	-	-	-	2	2	4	7
Black, both sexes	51	-	-	-	-	-	-	-	-	1	4	5
Male	24	-	-	-	-	-	-	-	-	1	1	1
Female	27	-	-	-	-	-	-	-	-	-	3	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Iowa												
All races, both sexes	310	-	-	-	-	-	-	-	-	3	4	7
Male	138	-	-	-	-	-	-	-	-	3	1	5
Female	172	-	-	-	-	-	-	-	-	-	3	2
White, both sexes	298	-	-	-	-	-	-	-	-	2	2	5
Male	135	-	-	-	-	-	-	-	-	2	1	3
Female	163	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	8	-	-	-	-	-	-	-	-	1	2	1
Male	2	-	-	-	-	-	-	-	-	1	-	1
Female	6	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	266	-	-	-	-	-	-	-	-	4	4	11
Male	123	-	-	-	-	-	-	-	-	2	2	4
Female	143	-	-	-	-	-	-	-	-	2	2	7
White, both sexes	249	-	-	-	-	-	-	-	-	2	4	10
Male	116	-	-	-	-	-	-	-	-	-	2	3
Female	133	-	-	-	-	-	-	-	-	2	2	7
Black, both sexes	12	-	-	-	-	-	-	-	-	1	-	1
Male	5	-	-	-	-	-	-	-	-	1	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Kentucky												
All races, both sexes	474	-	-	-	-	-	2	-	2	8	5	18
Male	220	-	-	-	-	-	2	-	1	2	3	11
Female	254	-	-	-	-	-	-	-	1	6	2	7
White, both sexes	428	-	-	-	-	-	-	-	2	7	3	14
Male	198	-	-	-	-	-	-	-	1	2	1	9
Female	230	-	-	-	-	-	-	-	1	5	2	5
Black, both sexes	42	-	-	-	-	-	2	-	-	1	2	4
Male	19	-	-	-	-	-	2	-	-	-	2	2
Female	23	-	-	-	-	-	-	-	-	1	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	462	3	1	1	-	2	1	4	2	5	16	29
Male	244	2	-	1	-	2	1	4	1	2	12	20
Female	218	1	1	-	-	-	-	-	1	3	4	9
White, both sexes	293	1	1	-	-	1	-	1	1	2	7	13
Male	150	1	-	-	-	1	-	1	-	1	5	9
Female	143	-	1	-	-	-	-	-	1	1	2	4
Black, both sexes	162	2	-	-	-	1	1	3	1	3	9	16
Male	90	1	-	-	-	1	1	3	1	1	7	11
Female	72	1	-	-	-	-	-	-	-	2	2	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	1	-	-	-	-	-	-	-	-
Male	4	-	-	1	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Maine												
All races, both sexes	143	-	-	-	-	-	-	-	-	1	1	4
Male	63	-	-	-	-	-	-	-	-	1	1	3
Female	80	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	143	-	-	-	-	-	-	-	-	1	1	4
Male	63	-	-	-	-	-	-	-	-	1	1	3
Female	80	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	494	-	-	1	-	1	-	1	1	8	14	22
Male	243	-	-	-	-	1	-	1	1	6	7	11
Female	251	-	-	1	-	-	-	-	-	2	7	11
White, both sexes	318	-	-	1	-	1	-	-	-	4	2	5
Male	149	-	-	-	-	1	-	-	-	2	2	1
Female	169	-	-	1	-	-	-	-	-	2	-	4
Black, both sexes	157	-	-	-	-	-	-	1	-	3	12	16
Male	79	-	-	-	-	-	-	1	-	3	5	10
Female	78	-	-	-	-	-	-	-	-	-	7	6
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	18	-	-	-	-	-	-	-	1	1	-	1
Male	14	-	-	-	-	-	-	-	1	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Massachusetts												
All races, both sexes	599	1	1	-	-	-	1	2	1	1	1	19
Male	249	-	1	-	-	-	-	2	1	1	1	10
Female	350	1	-	-	-	-	1	-	-	-	-	9
White, both sexes	558	1	1	-	-	-	-	1	-	-	1	14
Male	228	-	1	-	-	-	-	1	-	-	1	8
Female	330	1	-	-	-	-	-	-	-	-	-	6
Black, both sexes	26	-	-	-	-	-	1	1	1	1	-	4
Male	11	-	-	-	-	-	-	1	1	1	-	1
Female	15	-	-	-	-	-	1	-	-	-	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	959	3	-	-	-	2	-	5	3	9	21	32
Male	450	2	-	-	-	1	-	1	3	5	11	18
Female	509	1	-	-	-	1	-	4	-	4	10	14
White, both sexes	773	3	-	-	-	-	-	4	-	6	14	24
Male	350	2	-	-	-	-	-	-	-	2	7	13
Female	423	1	-	-	-	-	-	4	-	4	7	11
Black, both sexes	170	-	-	-	-	2	-	1	3	2	7	8
Male	94	-	-	-	-	1	-	1	3	2	4	5
Female	76	-	-	-	-	1	-	-	-	-	3	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Minnesota												
All races, both sexes	453	1	-	-	1	1	1	1	-	5	2	16
Male	199	1	-	-	-	-	-	1	-	3	-	13
Female	254	-	-	-	1	1	1	-	-	2	2	3
White, both sexes	416	-	-	-	-	1	1	-	-	4	1	10
Male	178	-	-	-	-	-	-	-	-	3	-	9
Female	238	-	-	-	-	1	1	-	-	1	1	1
Black, both sexes	19	1	-	-	1	-	-	-	-	-	1	3
Male	10	1	-	-	-	-	-	-	-	-	-	2
Female	9	-	-	-	1	-	-	-	-	-	1	1
American Indian 1/, both sexes	4	-	-	-	-	-	-	1	-	1	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	-	-	3
Male	9	-	-	-	-	-	-	-	-	-	-	2
Female	5	-	-	-	-	-	-	-	-	-	-	1
Mississippi												
All races, both sexes	305	-	-	-	1	1	1	-	1	4	8	18
Male	126	-	-	-	-	-	-	-	1	2	3	7
Female	179	-	-	-	1	1	1	-	-	2	5	11
White, both sexes	190	-	-	-	-	1	-	-	1	2	2	6
Male	73	-	-	-	-	-	-	-	1	1	-	3
Female	117	-	-	-	-	1	-	-	-	1	2	3
Black, both sexes	115	-	-	-	1	-	1	-	-	2	6	12
Male	53	-	-	-	-	-	-	-	-	1	3	4
Female	62	-	-	-	1	-	1	-	-	1	3	8
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Missouri												
All races, both sexes	659	1	2	1	-	-	1	2	3	2	14	28
Male	298	-	1	1	-	-	-	1	1	-	6	12
Female	361	1	1	-	-	-	1	1	2	2	8	16
White, both sexes	572	1	2	1	-	-	-	-	2	2	10	17
Male	260	-	1	1	-	-	-	-	-	-	6	7
Female	312	1	1	-	-	-	-	-	2	2	4	10
Black, both sexes	81	-	-	-	-	-	1	2	-	-	4	11
Male	34	-	-	-	-	-	-	1	-	-	-	5
Female	47	-	-	-	-	-	1	1	-	-	4	6
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	77	1	-	-	-	-	-	-	-	-	1	4
Male	25	-	-	-	-	-	-	-	-	-	-	2
Female	52	1	-	-	-	-	-	-	-	-	1	2
White, both sexes	74	1	-	-	-	-	-	-	-	-	1	4
Male	24	-	-	-	-	-	-	-	-	-	-	2
Female	50	1	-	-	-	-	-	-	-	-	1	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Nebraska												
All races, both sexes	167	1	-	-	-	-	-	-	1	-	1	6
Male	72	-	-	-	-	-	-	-	-	-	1	4
Female	95	1	-	-	-	-	-	-	1	-	-	2
White, both sexes	156	1	-	-	-	-	-	-	1	-	1	3
Male	66	-	-	-	-	-	-	-	-	-	1	2
Female	90	1	-	-	-	-	-	-	1	-	-	1
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	229	3	-	-	-	-	-	-	1	2	12	13
Male	137	2	-	-	-	-	-	-	1	1	6	8
Female	92	1	-	-	-	-	-	-	-	1	6	5
White, both sexes	184	3	-	-	-	-	-	-	1	1	9	8
Male	114	2	-	-	-	-	-	-	1	-	5	5
Female	70	1	-	-	-	-	-	-	-	1	4	3
Black, both sexes	21	-	-	-	-	-	-	-	-	-	1	2
Male	12	-	-	-	-	-	-	-	-	-	-	2
Female	9	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	22	-	-	-	-	-	-	-	-	-	2	3
Male	10	-	-	-	-	-	-	-	-	-	1	1
Female	12	-	-	-	-	-	-	-	-	-	1	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
New Hampshire												
All races, both sexes	96	-	-	-	-	-	-	-	-	-	1	2
Male	43	-	-	-	-	-	-	-	-	-	-	2
Female	53	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	95	-	-	-	-	-	-	-	-	-	1	2
Male	43	-	-	-	-	-	-	-	-	-	-	2
Female	52	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	830	-	-	1	-	1	2	1	4	7	17	26
Male	390	-	-	-	-	-	2	-	2	4	10	17
Female	440	-	-	1	-	1	-	1	2	3	7	9
White, both sexes	691	-	-	1	-	1	2	-	2	4	9	17
Male	317	-	-	-	-	-	2	-	1	2	5	12
Female	374	-	-	1	-	1	-	-	1	2	4	5
Black, both sexes	108	-	-	-	-	-	-	1	2	3	8	7
Male	56	-	-	-	-	-	-	-	1	2	5	4
Female	52	-	-	-	-	-	-	1	1	1	3	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	30	-	-	-	-	-	-	-	-	-	-	2
Male	16	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
New Mexico												
All races, both sexes	125	2	-	-	-	-	-	-	-	2	1	2
Male	59	-	-	-	-	-	-	-	-	1	1	1
Female	66	2	-	-	-	-	-	-	-	1	-	1
White, both sexes	111	1	-	-	-	-	-	-	-	1	-	2
Male	51	-	-	-	-	-	-	-	-	1	-	1
Female	60	1	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	10	1	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	1	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1,492	2	-	-	2	1	-	2	6	18	22	45
Male	655	1	-	-	-	-	-	1	3	10	11	26
Female	837	1	-	-	2	1	-	1	3	8	11	19
White, both sexes	1,216	2	-	-	-	1	-	2	5	10	12	28
Male	527	1	-	-	-	-	-	1	3	6	6	16
Female	689	1	-	-	-	1	-	1	2	4	6	12
Black, both sexes	221	-	-	-	1	-	-	-	1	8	9	16
Male	100	-	-	-	-	-	-	-	-	4	4	9
Female	121	-	-	-	1	-	-	-	1	4	5	7
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	54	-	-	-	1	-	-	-	-	-	1	1
Male	28	-	-	-	-	-	-	-	-	-	1	1
Female	26	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
North Carolina												
All races, both sexes	898	1	-	-	-	-	-	-	3	6	17	39
Male	409	-	-	-	-	-	-	-	1	4	9	22
Female	489	1	-	-	-	-	-	-	2	2	8	17
White, both sexes	652	-	-	-	-	-	-	-	2	3	5	15
Male	294	-	-	-	-	-	-	-	-	2	3	8
Female	358	-	-	-	-	-	-	-	2	1	2	7
Black, both sexes	224	1	-	-	-	-	-	-	1	3	11	23
Male	105	-	-	-	-	-	-	-	1	2	5	13
Female	119	1	-	-	-	-	-	-	-	1	6	10
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	70	-	-	-	-	-	-	-	-	-	-	-
Male	24	-	-	-	-	-	-	-	-	-	-	-
Female	46	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	68	-	-	-	-	-	-	-	-	-	-	-
Male	24	-	-	-	-	-	-	-	-	-	-	-
Female	44	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Ohio												
All races, both sexes	1,203	1	1	-	-	2	1	4	4	11	24	37
Male	546	1	1	-	-	2	-	4	2	7	12	26
Female	657	-	-	-	-	-	1	-	2	4	12	11
White, both sexes	1,029	1	1	-	-	2	1	3	2	7	14	27
Male	456	1	1	-	-	2	-	3	1	5	9	18
Female	573	-	-	-	-	-	1	-	1	2	5	9
Black, both sexes	165	-	-	-	-	-	-	1	2	3	9	10
Male	88	-	-	-	-	-	-	1	1	1	3	8
Female	77	-	-	-	-	-	-	-	1	2	6	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	1	1	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	7	-	-	-	-	-	-	-	-	-	1	-
Oklahoma												
All races, both sexes	310	1	-	1	1	-	1	2	4	3	7	8
Male	151	1	-	-	1	-	-	1	3	2	3	7
Female	159	-	-	1	-	-	1	1	1	1	4	1
White, both sexes	256	1	-	1	1	-	1	2	4	3	4	4
Male	120	1	-	-	1	-	-	1	3	2	1	3
Female	136	-	-	1	-	-	1	1	1	1	3	1
Black, both sexes	24	-	-	-	-	-	-	-	-	-	2	3
Male	17	-	-	-	-	-	-	-	-	-	2	3
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	26	-	-	-	-	-	-	-	-	-	1	1
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Oregon												
All races, both sexes	345	-	-	-	-	-	-	-	-	2	11	15
Male	167	-	-	-	-	-	-	-	-	1	7	14
Female	178	-	-	-	-	-	-	-	-	1	4	1
White, both sexes	325	-	-	-	-	-	-	-	-	2	10	13
Male	152	-	-	-	-	-	-	-	-	1	6	12
Female	173	-	-	-	-	-	-	-	-	1	4	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	-	-	-	2
Male	10	-	-	-	-	-	-	-	-	-	-	2
Female	3	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1,518	2	-	2	-	1	2	4	7	14	26	37
Male	667	1	-	-	-	1	1	3	4	5	18	21
Female	851	1	-	2	-	-	1	1	3	9	8	16
White, both sexes	1,342	2	-	-	-	1	2	2	5	14	18	22
Male	582	1	-	-	-	1	1	2	2	5	14	12
Female	760	1	-	-	-	-	1	-	3	9	4	10
Black, both sexes	149	-	-	2	-	-	-	1	1	-	6	14
Male	71	-	-	-	-	-	-	1	1	-	3	8
Female	78	-	-	2	-	-	-	-	-	-	3	6
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	26	-	-	-	-	-	-	1	1	-	2	1
Male	14	-	-	-	-	-	-	-	1	-	1	1
Female	12	-	-	-	-	-	-	1	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Rhode Island												
All races, both sexes	100	-	-	-	-	-	-	-	-	-	1	3
Male	46	-	-	-	-	-	-	-	-	-	-	2
Female	54	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	95	-	-	-	-	-	-	-	-	-	-	3
Male	44	-	-	-	-	-	-	-	-	-	-	2
Female	51	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	582	-	-	-	-	-	-	-	4	12	17	35
Male	298	-	-	-	-	-	-	-	1	10	12	19
Female	284	-	-	-	-	-	-	-	3	2	5	16
White, both sexes	379	-	-	-	-	-	-	-	3	4	6	12
Male	186	-	-	-	-	-	-	-	1	3	5	8
Female	193	-	-	-	-	-	-	-	2	1	1	4
Black, both sexes	198	-	-	-	-	-	-	-	1	8	10	23
Male	109	-	-	-	-	-	-	-	-	7	6	11
Female	89	-	-	-	-	-	-	-	1	1	4	12
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
South Dakota												
All races, both sexes	77	1	-	-	-	-	-	-	-	-	-	3
Male	32	-	-	-	-	-	-	-	-	-	-	-
Female	45	1	-	-	-	-	-	-	-	-	-	3
White, both sexes	70	-	-	-	-	-	-	-	-	-	-	3
Male	30	-	-	-	-	-	-	-	-	-	-	-
Female	40	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	652	1	-	-	-	-	2	2	3	9	19	33
Male	305	1	-	-	-	-	2	1	2	5	12	17
Female	347	-	-	-	-	-	-	1	1	4	7	16
White, both sexes	522	1	-	-	-	-	-	1	-	7	12	19
Male	234	1	-	-	-	-	-	-	-	4	7	10
Female	288	-	-	-	-	-	-	1	-	3	5	9
Black, both sexes	122	-	-	-	-	-	2	1	2	2	7	14
Male	66	-	-	-	-	-	2	1	2	1	5	7
Female	56	-	-	-	-	-	-	-	-	1	2	7
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	1	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Texas												
All races, both sexes	1,932	3	3	1	1	1	2	3	15	30	50	109
Male	954	2	1	-	-	1	2	2	8	20	27	70
Female	978	1	2	1	1	-	-	1	7	10	23	39
White, both sexes	1,612	1	3	-	1	1	2	3	11	21	36	78
Male	801	-	1	-	-	1	2	2	7	14	23	51
Female	811	1	2	-	1	-	-	1	4	7	13	27
Black, both sexes	278	2	-	1	-	-	-	-	4	8	14	27
Male	134	2	-	-	-	-	-	-	1	5	4	18
Female	144	-	-	1	-	-	-	-	3	3	10	9
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	40	-	-	-	-	-	-	-	-	1	-	3
Male	19	-	-	-	-	-	-	-	-	1	-	1
Female	21	-	-	-	-	-	-	-	-	-	-	2
Utah												
All races, both sexes	136	1	-	-	1	-	-	-	1	1	3	6
Male	70	1	-	-	1	-	-	-	-	1	2	2
Female	66	-	-	-	-	-	-	-	1	-	1	4
White, both sexes	131	1	-	-	1	-	-	-	1	1	3	6
Male	67	1	-	-	1	-	-	-	-	1	2	2
Female	64	-	-	-	-	-	-	-	1	-	1	4
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Vermont												
All races, both sexes	27	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	26	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	672	2	-	-	-	1	1	2	4	5	12	29
Male	302	2	-	-	-	1	1	-	4	1	4	16
Female	370	-	-	-	-	-	-	2	-	4	8	13
White, both sexes	499	-	-	-	-	-	1	1	1	2	6	14
Male	212	-	-	-	-	-	1	-	1	1	3	9
Female	287	-	-	-	-	-	-	1	-	1	3	5
Black, both sexes	157	2	-	-	-	1	-	-	3	3	6	15
Male	82	2	-	-	-	1	-	-	3	-	1	7
Female	75	-	-	-	-	-	-	-	-	3	5	8
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	1	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Washington												
All races, both sexes	488	-	1	-	1	-	1	2	2	1	4	14
Male	219	-	-	-	1	-	1	1	2	-	2	7
Female	269	-	1	-	-	-	-	1	-	1	2	7
White, both sexes	424	-	-	-	1	-	1	1	1	1	4	12
Male	191	-	-	-	1	-	1	1	1	-	2	5
Female	233	-	-	-	-	-	-	-	-	1	2	7
Black, both sexes	19	-	-	-	-	-	-	-	-	-	-	2
Male	10	-	-	-	-	-	-	-	-	-	-	2
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	9	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	36	-	1	-	-	-	-	-	1	-	-	-
Male	16	-	-	-	-	-	-	-	1	-	-	-
Female	20	-	1	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	229	-	-	-	-	-	1	-	1	2	5	7
Male	105	-	-	-	-	-	1	-	-	-	4	4
Female	124	-	-	-	-	-	-	-	1	2	1	3
White, both sexes	215	-	-	-	-	-	1	-	1	2	4	6
Male	97	-	-	-	-	-	1	-	-	-	3	4
Female	118	-	-	-	-	-	-	-	1	2	1	2
Black, both sexes	12	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Wisconsin												
All races, both sexes	496	-	-	-	-	-	3	1	6	3	7	16
Male	243	-	-	-	-	-	1	-	3	1	4	12
Female	253	-	-	-	-	-	2	1	3	2	3	4
White, both sexes	457	-	-	-	-	-	1	1	4	2	4	14
Male	225	-	-	-	-	-	1	-	3	1	2	11
Female	232	-	-	-	-	-	-	1	1	1	2	3
Black, both sexes	29	-	-	-	-	-	1	-	2	1	2	2
Male	13	-	-	-	-	-	-	-	-	-	1	1
Female	16	-	-	-	-	-	1	-	2	1	1	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	1	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	1	-	-	-	-	-
Wyoming												
All races, both sexes	37	-	-	-	-	-	-	-	-	1	-	2
Male	18	-	-	-	-	-	-	-	-	1	-	2
Female	19	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	36	-	-	-	-	-	-	-	-	1	-	2
Male	17	-	-	-	-	-	-	-	-	1	-	2
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
United States												
All races, both sexes	1,575	1,731	1,781	2,003	2,653	3,814	4,894	4,015	2,073	498	55	3
Male	909	1,037	1,033	1,057	1,343	1,802	2,038	1,471	623	97	11	3
Female	666	694	748	946	1,310	2,012	2,856	2,544	1,450	401	44	-
White, both sexes	936	1,108	1,319	1,552	2,189	3,318	4,397	3,671	1,884	449	44	2
Male	560	654	758	823	1,109	1,604	1,830	1,355	579	88	8	2
Female	376	454	561	729	1,080	1,714	2,567	2,316	1,305	361	36	-
Black, both sexes	551	510	368	343	324	349	330	217	130	34	6	1
Male	292	317	220	171	166	129	143	63	28	6	2	1
Female	259	193	148	172	158	220	187	154	102	28	4	-
American Indian 1/, both sexes	12	19	11	10	14	12	11	12	2	-	-	-
Male	6	8	8	1	5	6	5	3	-	-	-	-
Female	6	11	3	9	9	6	6	9	2	-	-	-
Asian or Pacific Islander 2/, both sexes	76	94	83	98	126	135	156	115	57	15	5	-
Male	51	58	47	62	63	63	60	50	16	3	1	-
Female	25	36	36	36	63	72	96	65	41	12	4	-
Alabama												
All races, both sexes	50	51	44	45	57	92	94	78	28	6	2	-
Male	26	31	23	24	35	41	40	35	8	2	2	-
Female	24	20	21	21	22	51	54	43	20	4	-	-
White, both sexes	23	23	27	35	46	76	79	67	25	4	1	-
Male	14	17	14	17	25	39	37	32	7	2	1	-
Female	9	6	13	18	21	37	42	35	18	2	-	-
Black, both sexes	26	28	17	10	11	15	14	11	3	2	1	-
Male	11	14	9	7	10	2	3	3	1	-	1	-
Female	15	14	8	3	1	13	11	8	2	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	1	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Alaska												
All races, both sexes	1	3	1	4	4	5	2	1	4	-	-	-
Male	1	3	1	1	4	4	2	1	1	-	-	-
Female	-	-	-	3	-	1	-	-	3	-	-	-
White, both sexes	-	3	-	3	3	4	1	1	3	-	-	-
Male	-	3	-	1	3	3	1	1	1	-	-	-
Female	-	-	-	2	-	1	-	-	2	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	1	-	-	1	-	-	-
Male	-	-	1	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Arizona												
All races, both sexes	23	30	24	39	43	51	72	48	23	7	-	-
Male	13	16	15	23	23	28	29	25	7	2	-	-
Female	10	14	9	16	20	23	43	23	16	5	-	-
White, both sexes	19	26	21	33	40	46	69	47	21	7	-	-
Male	12	13	13	19	21	26	27	25	7	2	-	-
Female	7	13	8	14	19	20	42	22	14	5	-	-
Black, both sexes	-	1	2	3	1	1	1	-	-	-	-	-
Male	-	1	1	2	1	-	1	-	-	-	-	-
Female	-	-	1	1	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	3	1	-	1	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	1	1	-	-	-	2	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	3	2	1	3	2	1	1	1	1	-	-	-
Male	1	2	1	2	1	1	-	-	-	-	-	-
Female	2	-	-	1	1	-	1	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Arkansas												
All races, both sexes	18	23	19	23	38	53	45	51	30	3	2	-
Male	11	13	11	12	23	17	18	14	8	-	-	-
Female	7	10	8	11	15	36	27	37	22	3	2	-
White, both sexes	12	13	16	18	31	48	40	48	26	2	2	-
Male	9	7	10	9	21	17	18	12	6	-	-	-
Female	3	6	6	9	10	31	22	36	20	2	2	-
Black, both sexes	6	9	3	5	7	4	5	3	4	1	-	-
Male	2	6	1	3	2	-	-	2	2	-	-	-
Female	4	3	2	2	5	4	5	1	2	1	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
California												
All races, both sexes	193	200	203	219	268	349	474	415	199	45	7	-
Male	125	124	121	131	113	146	188	151	65	12	2	-
Female	68	76	82	88	155	203	286	264	134	33	5	-
White, both sexes	121	134	141	160	192	269	375	342	159	41	4	-
Male	79	85	83	92	79	113	145	122	53	11	2	-
Female	42	49	58	68	113	156	230	220	106	30	2	-
Black, both sexes	39	32	32	21	26	15	26	15	12	1	-	-
Male	23	18	18	13	12	2	15	5	2	1	-	-
Female	16	14	14	8	14	13	11	10	10	-	-	-
American Indian 1/, both sexes	3	2	3	1	-	-	1	1	-	-	-	-
Male	3	1	2	-	-	-	-	-	-	-	-	-
Female	-	1	1	1	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	30	32	27	37	50	65	72	57	28	3	3	-
Male	20	20	18	26	22	31	28	24	10	-	-	-
Female	10	12	9	11	28	34	44	33	18	3	3	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Colorado												
All races, both sexes	20	26	22	26	37	44	58	53	20	6	-	-
Male	12	17	14	10	19	21	24	23	7	-	-	-
Female	8	9	8	16	18	23	34	30	13	6	-	-
White, both sexes	18	21	21	24	36	40	55	50	19	5	-	-
Male	10	15	13	10	18	20	22	22	7	-	-	-
Female	8	6	8	14	18	20	33	28	12	5	-	-
Black, both sexes	2	3	1	-	1	2	-	-	1	1	-	-
Male	2	2	1	-	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	2	-	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	2	-	1	3	3	-	-	-	-
Male	-	-	-	-	-	-	2	1	-	-	-	-
Female	-	2	-	2	-	1	1	2	-	-	-	-
Connecticut												
All races, both sexes	12	17	18	18	21	40	58	66	25	5	2	-
Male	6	11	10	10	6	24	24	23	7	2	-	-
Female	6	6	8	8	15	16	34	43	18	3	2	-
White, both sexes	6	13	16	14	18	38	55	64	24	5	2	-
Male	4	10	9	8	5	23	23	23	6	2	-	-
Female	2	3	7	6	13	15	32	41	18	3	2	-
Black, both sexes	5	4	-	4	2	2	1	2	1	-	-	-
Male	2	1	-	2	1	1	-	-	1	-	-	-
Female	3	3	-	2	1	1	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	2	-	1	-	2	-	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	1	-	1	-	1	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Delaware												
All races, both sexes	3	2	1	10	15	9	12	9	7	2	-	-
Male	1	2	1	3	7	2	3	2	1	-	-	-
Female	2	-	-	7	8	7	9	7	6	2	-	-
White, both sexes	2	2	1	5	13	8	10	9	6	2	-	-
Male	1	2	1	2	6	2	1	2	-	-	-	-
Female	1	-	-	3	7	6	9	7	6	2	-	-
Black, both sexes	1	-	-	5	1	1	2	-	1	-	-	-
Male	-	-	-	1	-	-	2	-	1	-	-	-
Female	1	-	-	4	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	9	6	4	4	5	4	7	8	5	1	1	-
Male	2	5	2	2	1	1	1	-	1	-	-	-
Female	7	1	2	2	4	3	6	8	4	1	1	-
White, both sexes	1	-	-	-	3	1	5	2	2	-	1	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	1	-	-	-	3	-	4	2	2	-	1	-
Black, both sexes	8	6	4	4	1	3	2	6	3	1	-	-
Male	2	5	2	2	-	-	-	-	1	-	-	-
Female	6	1	2	2	1	3	2	6	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Florida												
All races, both sexes	120	113	109	123	153	240	329	269	132	34	-	-
Male	67	67	68	66	85	127	143	90	37	8	-	-
Female	53	46	41	57	68	113	186	179	95	26	-	-
White, both sexes	69	68	82	95	126	223	303	255	124	34	-	-
Male	43	43	49	54	72	122	133	86	35	8	-	-
Female	26	25	33	41	54	101	170	169	89	26	-	-
Black, both sexes	49	42	26	28	24	14	23	13	6	-	-	-
Male	23	23	18	12	10	3	10	4	1	-	-	-
Female	26	19	8	16	14	11	13	9	5	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	1	-	3	3	2	1	2	-	-	-
Male	1	1	1	-	3	2	-	-	1	-	-	-
Female	1	1	-	-	-	1	2	1	1	-	-	-
Georgia												
All races, both sexes	81	72	79	75	78	95	104	85	45	6	1	1
Male	53	43	47	50	37	42	43	27	9	-	1	1
Female	28	29	32	25	41	53	61	58	36	6	-	-
White, both sexes	34	30	43	52	55	67	87	71	40	6	-	1
Male	23	20	24	36	27	31	35	24	8	-	-	1
Female	11	10	19	16	28	36	52	47	32	6	-	-
Black, both sexes	46	40	32	23	20	25	17	13	4	-	1	-
Male	29	22	21	14	10	10	8	2	1	-	1	-
Female	17	18	11	9	10	15	9	11	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	4	-	2	3	-	1	1	-	-	-
Male	1	1	2	-	-	1	-	1	-	-	-	-
Female	-	1	2	-	2	2	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Hawaii												
All races, both sexes	11	19	11	9	12	25	33	20	12	7	-	-
Male	10	11	4	4	6	14	14	9	3	2	-	-
Female	1	8	7	5	6	11	19	11	9	5	-	-
White, both sexes	2	3	4	3	3	4	8	3	2	-	-	-
Male	2	1	2	2	3	4	4	2	1	-	-	-
Female	-	2	2	1	-	-	4	1	1	-	-	-
Black, both sexes	-	2	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	14	7	6	9	21	25	17	10	7	-	-
Male	8	9	2	2	3	10	10	7	2	2	-	-
Female	1	5	5	4	6	11	15	10	8	5	-	-
Idaho												
All races, both sexes	6	4	9	6	12	11	20	13	9	2	-	-
Male	2	3	7	5	6	6	9	5	5	-	-	-
Female	4	1	2	1	6	5	11	8	4	2	-	-
White, both sexes	6	4	8	6	11	11	20	13	9	2	-	-
Male	2	3	6	5	6	6	9	5	5	-	-	-
Female	4	1	2	1	5	5	11	8	4	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Illinois												
All races, both sexes	55	77	44	91	96	147	204	143	85	25	-	-
Male	29	39	35	52	57	64	81	40	33	1	-	-
Female	26	38	9	39	39	83	123	103	52	24	-	-
White, both sexes	33	50	32	74	81	128	190	134	80	22	-	-
Male	20	23	26	44	45	54	78	40	30	1	-	-
Female	13	27	6	30	36	74	112	94	50	21	-	-
Black, both sexes	19	25	11	16	12	16	10	8	5	3	-	-
Male	7	15	9	7	9	8	2	-	3	-	-	-
Female	12	10	2	9	3	8	8	8	2	3	-	-
American Indian 1/, both sexes	2	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	1	1	3	2	4	1	-	-	-	-
Male	1	1	-	1	3	2	1	-	-	-	-	-
Female	-	1	1	-	-	-	3	1	-	-	-	-
Indiana												
All races, both sexes	24	43	35	35	53	72	112	85	49	17	2	-
Male	14	17	16	18	33	31	35	28	10	2	-	-
Female	10	26	19	17	20	41	77	57	39	15	2	-
White, both sexes	12	35	28	33	50	68	108	82	49	17	2	-
Male	7	14	12	16	31	29	34	27	10	2	-	-
Female	5	21	16	17	19	39	74	55	39	15	2	-
Black, both sexes	12	8	7	1	3	4	3	3	-	-	-	-
Male	7	3	4	1	2	2	1	1	-	-	-	-
Female	5	5	3	-	1	2	2	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Iowa												
All races, both sexes	16	11	22	17	33	52	66	38	30	11	-	-
Male	8	7	16	6	16	28	31	9	7	1	-	-
Female	8	4	6	11	17	24	35	29	23	10	-	-
White, both sexes	13	11	19	17	33	52	66	37	30	11	-	-
Male	8	7	16	6	16	28	31	9	7	1	-	-
Female	5	4	3	11	17	24	35	28	23	10	-	-
Black, both sexes	2	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	1	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	2	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	2	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	9	9	17	27	22	41	42	53	21	6	-	-
Male	4	5	11	13	11	22	22	17	7	3	-	-
Female	5	4	6	14	11	19	20	36	14	3	-	-
White, both sexes	8	5	17	23	21	39	41	52	21	6	-	-
Male	3	3	11	13	11	21	22	17	7	3	-	-
Female	5	2	6	10	10	18	19	35	14	3	-	-
Black, both sexes	1	4	-	2	1	1	-	1	-	-	-	-
Male	1	2	-	-	-	-	-	-	-	-	-	-
Female	-	2	-	2	1	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	2	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Kentucky												
All races, both sexes	21	20	48	35	52	64	89	54	48	8	-	-
Male	11	11	22	24	22	34	41	19	15	2	-	-
Female	10	9	26	11	30	30	48	35	33	6	-	-
White, both sexes	15	15	42	30	48	58	89	52	45	8	-	-
Male	11	7	19	20	22	32	41	17	14	2	-	-
Female	4	8	23	10	26	26	48	35	31	6	-	-
Black, both sexes	6	5	5	4	4	6	-	1	2	-	-	-
Male	-	4	2	3	-	2	-	1	1	-	-	-
Female	6	1	3	1	4	4	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	-	-	1	1	-	-	-
Male	-	-	1	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Louisiana												
All races, both sexes	38	36	29	35	43	58	82	45	25	6	-	1
Male	24	19	14	19	25	26	39	20	11	1	-	1
Female	14	17	15	16	18	32	43	25	14	5	-	-
White, both sexes	17	15	17	15	33	39	67	38	20	5	-	-
Male	10	8	7	10	20	18	32	17	9	1	-	-
Female	7	7	10	5	13	21	35	21	11	4	-	-
Black, both sexes	19	21	11	19	9	18	15	7	5	1	-	1
Male	12	11	7	8	5	8	7	3	2	-	-	1
Female	7	10	4	11	4	10	8	4	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	1	1	1	1	-	-	-	-	-	-
Male	2	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Maine												
All races, both sexes	3	4	10	9	20	28	28	22	9	4	-	-
Male	2	2	5	6	11	13	10	3	5	1	-	-
Female	1	2	5	3	9	15	18	19	4	3	-	-
White, both sexes	3	4	10	9	20	28	28	22	9	4	-	-
Male	2	2	5	6	11	13	10	3	5	1	-	-
Female	1	2	5	3	9	15	18	19	4	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	30	26	25	34	57	91	70	69	38	5	1	-
Male	17	22	17	12	27	46	35	23	14	2	1	-
Female	13	4	8	22	30	45	35	46	24	3	-	-
White, both sexes	10	11	11	23	32	68	56	54	35	5	-	-
Male	7	10	7	7	13	36	28	19	14	2	-	-
Female	3	1	4	16	19	32	28	35	21	3	-	-
Black, both sexes	20	15	13	8	21	18	14	13	3	-	-	-
Male	10	12	9	2	11	7	7	2	-	-	-	-
Female	10	3	4	6	10	11	7	11	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	3	4	4	-	2	-	-	1	-
Male	-	-	1	3	3	2	-	2	-	-	1	-
Female	-	-	-	-	1	2	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Massachusetts												
All races, both sexes	18	17	33	41	60	100	124	109	52	18	-	-
Male	10	11	15	15	34	52	42	35	15	4	-	-
Female	8	6	18	26	26	48	82	74	37	14	-	-
White, both sexes	14	15	30	37	56	95	120	105	50	18	-	-
Male	9	10	13	12	32	51	38	33	15	4	-	-
Female	5	5	17	25	24	44	82	72	35	14	-	-
Black, both sexes	3	1	1	3	1	3	2	3	1	-	-	-
Male	1	-	-	2	-	1	2	1	-	-	-	-
Female	2	1	1	1	1	2	-	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	2	1	3	2	2	1	1	-	-	-
Male	-	1	2	1	2	-	2	1	-	-	-	-
Female	1	-	-	-	1	2	-	-	1	-	-	-
Michigan												
All races, both sexes	44	44	66	75	86	140	173	148	84	20	4	-
Male	24	31	42	28	45	74	73	62	24	5	1	-
Female	20	13	24	47	41	66	100	86	60	15	3	-
White, both sexes	24	21	47	58	73	122	147	139	72	15	4	-
Male	14	15	30	21	36	64	63	58	22	2	1	-
Female	10	6	17	37	37	58	84	81	50	13	3	-
Black, both sexes	20	21	19	15	12	15	21	8	12	4	-	-
Male	10	14	12	7	8	9	9	4	2	3	-	-
Female	10	7	7	8	4	6	12	4	10	1	-	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	1	1	2	5	1	-	1	-	-
Male	-	2	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	1	-	2	4	1	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Minnesota												
All races, both sexes	13	27	20	28	38	57	98	75	61	7	1	-
Male	7	15	13	17	20	25	41	28	14	1	-	-
Female	6	12	7	11	18	32	57	47	47	6	1	-
White, both sexes	10	18	18	26	34	56	96	72	61	7	1	-
Male	5	8	12	16	19	25	41	25	14	1	-	-
Female	5	10	6	10	15	31	55	47	47	6	1	-
Black, both sexes	2	6	2	-	1	-	1	1	-	-	-	-
Male	1	4	1	-	-	-	-	1	-	-	-	-
Female	1	2	1	-	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	3	-	2	2	1	-	2	-	-	-	-
Male	1	3	-	1	-	-	-	2	-	-	-	-
Female	-	-	-	1	2	1	-	-	-	-	-	-
Mississippi												
All races, both sexes	17	30	24	25	46	36	44	28	16	4	1	-
Male	10	24	10	8	24	13	13	8	3	-	-	-
Female	7	6	14	17	22	23	31	20	13	4	1	-
White, both sexes	9	10	16	12	36	26	29	23	12	4	1	-
Male	5	7	7	6	18	8	8	6	3	-	-	-
Female	4	3	9	6	18	18	21	17	9	4	1	-
Black, both sexes	8	20	8	13	10	10	15	5	4	-	-	-
Male	5	17	3	2	6	5	5	2	-	-	-	-
Female	3	3	5	11	4	5	10	3	4	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Missouri												
All races, both sexes	25	34	42	51	55	86	133	107	58	14	-	-
Male	12	14	24	35	30	32	62	41	23	3	-	-
Female	13	20	18	16	25	54	71	66	35	11	-	-
White, both sexes	21	22	36	48	47	76	126	98	51	12	-	-
Male	10	9	20	32	25	31	57	37	21	3	-	-
Female	11	13	16	16	22	45	69	61	30	9	-	-
Black, both sexes	4	12	6	3	6	9	7	8	6	2	-	-
Male	2	5	4	3	3	1	5	3	2	-	-	-
Female	2	7	2	-	3	8	2	5	4	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	1	-	1	1	-	-	-
Male	-	-	-	-	2	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
Montana												
All races, both sexes	1	5	4	3	5	12	9	21	9	2	-	-
Male	-	2	1	1	1	4	5	4	4	1	-	-
Female	1	3	3	2	4	8	4	17	5	1	-	-
White, both sexes	1	3	4	2	5	12	9	21	9	2	-	-
Male	-	1	1	1	1	4	5	4	4	1	-	-
Female	1	2	3	1	4	8	4	17	5	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	2	-	1	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Nebraska												
All races, both sexes	9	12	8	13	18	20	32	26	17	3	-	-
Male	5	10	5	5	13	6	10	9	4	-	-	-
Female	4	2	3	8	5	14	22	17	13	3	-	-
White, both sexes	8	7	7	13	18	20	32	26	17	2	-	-
Male	4	7	5	5	13	6	10	9	4	-	-	-
Female	4	-	2	8	5	14	22	17	13	2	-	-
Black, both sexes	1	4	-	-	-	-	-	-	-	1	-	-
Male	1	2	-	-	-	-	-	-	-	-	-	-
Female	-	2	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	16	18	19	24	28	30	37	16	8	1	1	-
Male	12	11	12	14	18	17	20	10	4	-	1	-
Female	4	7	7	10	10	13	17	6	4	1	-	-
White, both sexes	12	13	14	14	24	27	33	16	7	1	1	-
Male	8	9	8	10	17	16	18	10	4	-	1	-
Female	4	4	6	4	7	11	15	6	3	1	-	-
Black, both sexes	2	-	3	5	3	1	3	-	1	-	-	-
Male	2	-	3	2	1	-	2	-	-	-	-	-
Female	-	-	-	3	2	1	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	5	2	5	1	2	-	-	-	-	-	-
Male	2	2	1	2	-	1	-	-	-	-	-	-
Female	-	3	1	3	1	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
New Hampshire												
All races, both sexes	3	3	3	5	12	17	21	17	11	1	-	-
Male	1	2	2	2	6	8	8	9	3	-	-	-
Female	2	1	1	3	6	9	13	8	8	1	-	-
White, both sexes	3	3	2	5	12	17	21	17	11	1	-	-
Male	1	2	2	2	6	8	8	9	3	-	-	-
Female	2	1	-	3	6	9	13	8	8	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	45	45	31	70	62	133	159	144	62	16	4	-
Male	23	28	19	43	36	60	77	45	21	3	-	-
Female	22	17	12	27	26	73	82	99	41	13	4	-
White, both sexes	29	31	25	54	44	122	142	134	57	15	2	-
Male	17	19	14	33	22	57	68	42	20	3	-	-
Female	12	12	11	21	22	65	74	92	37	12	2	-
Black, both sexes	11	13	5	13	11	8	13	7	3	1	2	-
Male	2	8	4	7	8	3	9	3	-	-	-	-
Female	9	5	1	6	3	5	4	4	3	1	2	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	1	1	3	6	3	4	3	2	-	-	-
Male	4	1	1	3	5	-	-	-	1	-	-	-
Female	1	-	-	-	1	3	4	3	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
New Mexico												
All races, both sexes	6	9	9	7	17	22	19	18	9	2	-	-
Male	4	6	1	5	10	14	7	7	2	-	-	-
Female	2	3	8	2	7	8	12	11	7	2	-	-
White, both sexes	6	8	8	6	15	19	18	17	8	2	-	-
Male	4	5	1	5	9	11	6	6	2	-	-	-
Female	2	3	7	1	6	8	12	11	6	2	-	-
Black, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	1	1	2	1	1	-	1	-	-	-
Male	-	1	-	-	1	1	1	-	-	-	-	-
Female	-	-	1	1	1	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	64	92	86	100	144	227	261	252	136	27	5	-
Male	31	55	42	47	70	107	104	104	39	4	-	-
Female	33	37	44	53	74	120	157	148	97	23	5	-
White, both sexes	40	64	67	70	114	194	231	226	121	24	5	-
Male	20	36	34	32	55	91	93	93	36	4	-	-
Female	20	28	33	38	59	103	138	133	85	20	5	-
Black, both sexes	21	22	15	25	24	28	18	20	11	2	-	-
Male	11	15	5	12	12	13	6	7	2	-	-	-
Female	10	7	10	13	12	15	12	13	9	2	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	5	4	5	6	5	12	6	4	1	-	-
Male	-	4	3	3	3	3	5	4	1	-	-	-
Female	3	1	1	2	3	2	7	2	3	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
North Carolina												
All races, both sexes	56	68	68	56	110	112	173	117	60	11	1	-
Male	30	44	43	28	53	55	63	36	19	2	-	-
Female	26	24	25	28	57	57	110	81	41	9	1	-
White, both sexes	26	37	33	40	91	89	152	97	52	9	1	-
Male	17	22	22	20	44	45	57	34	18	2	-	-
Female	9	15	11	20	47	44	95	63	34	7	1	-
Black, both sexes	28	27	32	14	16	22	20	17	7	2	-	-
Male	12	20	19	8	8	9	5	2	1	-	-	-
Female	16	7	13	6	8	13	15	15	6	2	-	-
American Indian 1/, both sexes	1	1	1	2	2	-	-	2	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	1	1	-	2	1	-	-	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	3	2	-	1	1	1	1	1	-	-	-
Male	1	2	1	-	-	1	1	-	-	-	-	-
Female	-	1	1	-	1	-	-	1	1	-	-	-
North Dakota												
All races, both sexes	2	2	7	7	4	5	11	12	13	6	1	-
Male	2	2	2	2	2	2	5	4	2	1	-	-
Female	-	-	5	5	2	3	6	8	11	5	1	-
White, both sexes	2	2	7	6	3	5	11	12	13	6	1	-
Male	2	2	2	2	2	2	5	4	2	1	-	-
Female	-	-	5	4	1	3	6	8	11	5	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Ohio												
All races, both sexes	61	67	81	87	117	198	212	177	96	20	2	-
Male	29	43	41	39	53	92	86	70	33	5	-	-
Female	32	24	40	48	64	106	126	107	63	15	2	-
White, both sexes	31	48	67	67	101	177	203	168	91	16	2	-
Male	13	29	34	26	48	82	82	66	31	5	-	-
Female	18	19	33	41	53	95	121	102	60	11	2	-
Black, both sexes	30	19	12	19	15	20	7	9	5	4	-	-
Male	16	14	7	13	5	10	3	4	2	-	-	-
Female	14	5	5	6	10	10	4	5	3	4	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	1	1	1	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	2	1	1	1	1	-	-	-	-	-
Oklahoma												
All races, both sexes	18	18	19	15	33	44	64	40	25	6	-	-
Male	9	13	13	5	17	22	29	15	8	2	-	-
Female	9	5	6	10	16	22	35	25	17	4	-	-
White, both sexes	12	10	12	10	26	40	57	37	25	6	-	-
Male	6	8	7	4	15	22	23	13	8	2	-	-
Female	6	2	5	6	11	18	34	24	17	4	-	-
Black, both sexes	2	2	4	3	1	2	5	-	-	-	-	-
Male	1	2	4	-	1	-	4	-	-	-	-	-
Female	1	-	-	3	-	2	1	-	-	-	-	-
American Indian 1/, both sexes	4	3	2	2	6	2	2	3	-	-	-	-
Male	2	2	2	1	1	-	2	2	-	-	-	-
Female	2	1	-	1	5	2	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	3	1	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	2	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Oregon												
All races, both sexes	14	22	22	25	36	43	65	49	32	8	1	-
Male	8	11	13	11	18	23	29	15	12	4	1	-
Female	6	11	9	14	18	20	36	34	20	4	-	-
White, both sexes	14	20	19	23	32	40	65	48	32	6	1	-
Male	8	9	10	10	15	21	29	15	12	3	1	-
Female	6	11	9	13	17	19	36	33	20	3	-	-
Black, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	1	-	-	1	-	1	-	-	-	-
Male	-	1	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	2	3	2	-	-	-	2	-	-
Male	-	1	1	1	3	1	-	-	-	1	-	-
Female	-	-	-	1	-	1	-	-	-	1	-	-
Pennsylvania												
All races, both sexes	63	71	95	94	122	230	327	270	122	25	4	-
Male	37	38	52	46	54	110	143	99	32	2	-	-
Female	26	33	43	48	68	120	184	171	90	23	4	-
White, both sexes	40	53	74	79	103	212	310	261	118	22	4	-
Male	21	28	39	40	48	103	136	96	31	2	-	-
Female	19	25	35	39	55	109	174	165	87	20	4	-
Black, both sexes	22	16	18	12	13	17	15	5	4	3	-	-
Male	15	8	11	3	4	7	7	2	1	-	-	-
Female	7	8	7	9	9	10	8	3	3	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	3	3	6	1	2	3	-	-	-	-
Male	1	2	2	3	2	-	-	1	-	-	-	-
Female	-	-	1	-	4	1	2	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Rhode Island												
All races, both sexes	2	3	10	7	3	11	26	21	13	-	-	-
Male	1	-	7	3	2	5	12	10	4	-	-	-
Female	1	3	3	4	1	6	14	11	9	-	-	-
White, both sexes	2	3	8	6	3	10	26	21	13	-	-	-
Male	1	-	5	3	2	5	12	10	4	-	-	-
Female	1	3	3	3	1	5	14	11	9	-	-	-
Black, both sexes	-	-	2	-	-	1	-	-	-	-	-	-
Male	-	-	2	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	53	55	33	52	56	66	85	67	34	10	2	1
Male	29	36	20	33	33	30	42	21	7	4	-	1
Female	24	19	13	19	23	36	43	46	27	6	2	-
White, both sexes	22	30	17	31	40	52	69	57	24	10	1	1
Male	10	16	12	19	24	23	34	20	6	4	-	1
Female	12	14	5	12	16	29	35	37	18	6	1	-
Black, both sexes	31	25	15	20	15	14	16	9	10	-	1	-
Male	19	20	7	14	8	7	8	1	1	-	-	-
Female	12	5	8	6	7	7	8	8	9	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	1	-	-	1	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
South Dakota												
All races, both sexes	4	3	6	1	6	15	14	19	4	1	-	-
Male	2	1	4	1	2	8	5	7	1	1	-	-
Female	2	2	2	-	4	7	9	12	3	-	-	-
White, both sexes	3	2	4	1	6	15	13	18	4	1	-	-
Male	2	-	3	1	2	8	5	7	1	1	-	-
Female	1	2	1	-	4	7	8	11	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	-	-	-	1	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	48	50	40	56	67	87	107	84	33	11	-	-
Male	25	32	21	29	30	45	43	32	6	2	-	-
Female	23	18	19	27	37	42	64	52	27	9	-	-
White, both sexes	23	31	31	42	58	78	99	78	32	10	-	-
Male	12	20	16	21	26	42	39	29	6	1	-	-
Female	11	11	15	21	32	36	60	49	26	9	-	-
Black, both sexes	24	19	8	12	9	8	8	5	-	1	-	-
Male	12	12	5	6	4	2	4	2	-	1	-	-
Female	12	7	3	6	5	6	4	3	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	2	-	1	-	1	1	-	-	-
Male	1	-	-	2	-	1	-	1	-	-	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Texas												
All races, both sexes	139	135	150	140	179	257	300	263	112	34	5	-
Male	90	78	90	74	89	116	124	112	40	7	1	-
Female	49	57	60	66	90	141	176	151	72	27	4	-
White, both sexes	88	94	117	115	153	230	272	248	104	30	4	-
Male	61	56	74	63	74	106	114	106	39	6	1	-
Female	27	38	43	52	79	124	158	142	65	24	3	-
Black, both sexes	48	36	27	20	23	23	23	10	8	3	1	-
Male	28	20	13	7	14	7	8	5	1	1	-	-
Female	20	16	14	13	9	16	15	5	7	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	5	6	5	3	4	5	4	-	1	-	-
Male	1	2	3	4	1	3	2	1	-	-	-	-
Female	2	3	3	1	2	1	3	3	-	1	-	-
Utah												
All races, both sexes	11	8	7	8	17	12	28	15	14	2	1	-
Male	4	5	5	7	9	7	10	7	9	-	-	-
Female	7	3	2	1	8	5	18	8	5	2	1	-
White, both sexes	9	8	6	8	16	12	28	15	14	2	-	-
Male	3	5	4	7	8	7	10	7	9	-	-	-
Female	6	3	2	1	8	5	18	8	5	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	1	-	-	-	-	-	1	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Vermont												
All races, both sexes	1	-	2	2	4	1	4	7	1	4	-	-
Male	-	-	1	1	2	1	1	3	-	1	-	-
Female	1	-	1	1	2	-	3	4	1	3	-	-
White, both sexes	1	-	2	1	4	1	4	7	1	4	-	-
Male	-	-	1	-	2	1	1	3	-	1	-	-
Female	1	-	1	1	2	-	3	4	1	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	40	39	50	39	71	93	127	92	57	8	-	-
Male	28	23	32	19	39	41	44	31	15	1	-	-
Female	12	16	18	20	32	52	83	61	42	7	-	-
White, both sexes	19	20	30	28	55	77	107	82	49	7	-	-
Male	11	11	18	12	31	35	35	29	14	1	-	-
Female	8	9	12	16	24	42	72	53	35	6	-	-
Black, both sexes	20	17	18	9	14	15	19	7	7	1	-	-
Male	17	12	12	5	7	5	9	1	-	-	-	-
Female	3	5	6	4	7	10	10	6	7	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	2	2	2	1	1	3	1	-	-	-
Male	-	-	2	2	1	1	-	1	1	-	-	-
Female	1	2	-	-	1	-	1	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Washington												
All races, both sexes	25	23	29	38	49	70	77	90	45	15	1	-
Male	14	8	16	17	24	36	33	41	13	2	1	-
Female	11	15	13	21	25	34	44	49	32	13	-	-
White, both sexes	17	16	24	31	40	66	67	82	44	15	1	-
Male	10	6	12	14	21	35	28	38	13	2	1	-
Female	7	10	12	17	19	31	39	44	31	13	-	-
Black, both sexes	5	-	4	2	1	-	2	3	-	-	-	-
Male	1	-	3	2	-	-	1	1	-	-	-	-
Female	4	-	1	-	1	-	1	2	-	-	-	-
American Indian 1/, both sexes	-	4	-	-	-	1	1	2	-	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	3	-	-	-	1	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	3	1	5	8	3	7	3	1	-	-	-
Male	3	1	1	1	3	1	4	1	-	-	-	-
Female	-	2	-	4	5	2	3	2	1	-	-	-
West Virginia												
All races, both sexes	11	12	10	17	26	40	61	23	8	3	2	-
Male	9	8	2	7	14	18	28	9	-	1	-	-
Female	2	4	8	10	12	22	33	14	8	2	2	-
White, both sexes	9	9	9	17	26	37	60	21	8	3	2	-
Male	7	5	2	7	14	17	28	8	-	1	-	-
Female	2	4	7	10	12	20	32	13	8	2	2	-
Black, both sexes	2	2	1	-	-	3	1	2	-	-	-	-
Male	2	2	-	-	-	1	-	1	-	-	-	-
Female	-	-	1	-	-	2	1	1	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Wisconsin												
All races, both sexes	22	23	31	32	62	75	94	77	34	9	1	-
Male	14	17	23	23	28	39	42	31	5	-	-	-
Female	8	6	8	9	34	36	52	46	29	9	1	-
White, both sexes	16	20	27	29	58	73	90	76	32	9	1	-
Male	11	15	21	21	25	38	40	31	5	-	-	-
Female	5	5	6	8	33	35	50	45	27	9	1	-
Black, both sexes	5	3	2	2	4	2	1	1	1	-	-	-
Male	2	2	2	1	3	1	-	-	-	-	-	-
Female	3	1	-	1	1	1	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	1	-	-	3	-	1	-	-	-
Male	1	-	-	1	-	-	2	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	1	-	-	-
Wyoming												
All races, both sexes	1	4	2	1	1	4	8	6	3	4	-	-
Male	1	1	2	1	-	3	5	2	-	-	-	-
Female	-	3	-	-	1	1	3	4	3	4	-	-
White, both sexes	1	4	2	1	1	3	8	6	3	4	-	-
Male	1	1	2	1	-	2	5	2	-	-	-	-
Female	-	3	-	-	1	1	3	4	3	4	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
United States												
All races, both sexes	6,718	13	1	3	3	8	4	9	25	39	90	152
Male	2,770	10	-	-	1	5	2	3	15	25	54	92
Female	3,948	3	1	3	2	3	2	6	10	14	36	60
White, both sexes	5,757	8	1	2	3	6	3	7	19	25	57	106
Male	2,344	6	-	-	1	4	2	2	11	14	37	64
Female	3,413	2	1	2	2	2	1	5	8	11	20	42
Black, both sexes	791	5	-	-	-	2	1	2	5	14	29	40
Male	353	4	-	-	-	1	-	1	3	11	15	23
Female	438	1	-	-	-	1	1	1	2	3	14	17
American Indian 1/, both sexes	23	-	-	-	-	-	-	-	-	-	1	1
Male	10	-	-	-	-	-	-	-	-	-	1	1
Female	13	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	147	-	-	1	-	-	-	-	1	-	3	5
Male	63	-	-	-	-	-	-	-	1	-	1	4
Female	84	-	-	1	-	-	-	-	-	-	2	1
Alabama												
All races, both sexes	111	-	-	-	-	-	-	-	1	1	1	2
Male	44	-	-	-	-	-	-	-	-	1	-	2
Female	67	-	-	-	-	-	-	-	1	-	1	-
White, both sexes	86	-	-	-	-	-	-	-	1	-	-	1
Male	30	-	-	-	-	-	-	-	-	-	-	1
Female	56	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	25	-	-	-	-	-	-	-	-	1	1	1
Male	14	-	-	-	-	-	-	-	-	1	-	1
Female	11	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Alaska												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	122	-	-	-	-	-	-	-	-	3	2	3
Male	52	-	-	-	-	-	-	-	-	1	2	3
Female	70	-	-	-	-	-	-	-	-	2	-	-
White, both sexes	117	-	-	-	-	-	-	-	-	3	1	3
Male	49	-	-	-	-	-	-	-	-	1	1	3
Female	68	-	-	-	-	-	-	-	-	2	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Arkansas												
All races, both sexes	78	-	-	-	-	-	1	-	-	2	4	1
Male	34	-	-	-	-	-	-	-	-	2	2	1
Female	44	-	-	-	-	-	1	-	-	-	2	-
White, both sexes	66	-	-	-	-	-	1	-	-	2	3	1
Male	30	-	-	-	-	-	-	-	-	2	2	1
Female	36	-	-	-	-	-	1	-	-	-	1	-
Black, both sexes	12	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	596	-	-	-	-	1	-	2	2	2	9	13
Male	263	-	-	-	-	-	-	1	2	1	3	11
Female	333	-	-	-	-	1	-	1	-	1	6	2
White, both sexes	496	-	-	-	-	1	-	2	2	1	6	11
Male	215	-	-	-	-	-	-	1	2	-	3	9
Female	281	-	-	-	-	1	-	1	-	1	3	2
Black, both sexes	42	-	-	-	-	-	-	-	-	1	1	-
Male	19	-	-	-	-	-	-	-	-	1	-	-
Female	23	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	57	-	-	-	-	-	-	-	-	-	2	2
Male	29	-	-	-	-	-	-	-	-	-	-	2
Female	28	-	-	-	-	-	-	-	-	-	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Colorado												
All races, both sexes	80	-	-	-	-	-	-	-	-	-	1	3
Male	31	-	-	-	-	-	-	-	-	-	1	1
Female	49	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	78	-	-	-	-	-	-	-	-	-	1	2
Male	31	-	-	-	-	-	-	-	-	-	1	1
Female	47	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	83	-	-	-	-	-	-	-	-	-	-	-
Male	34	-	-	-	-	-	-	-	-	-	-	-
Female	49	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	72	-	-	-	-	-	-	-	-	-	-	-
Male	31	-	-	-	-	-	-	-	-	-	-	-
Female	41	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Delaware												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	-	3
Male	7	-	-	-	-	-	-	-	-	-	-	2
Female	12	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	2
Male	2	-	-	-	-	-	-	-	-	-	-	2
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Florida												
All races, both sexes	494	1	-	-	1	1	-	1	2	3	10	12
Male	212	1	-	-	-	1	-	-	-	2	7	6
Female	282	-	-	-	1	-	-	1	2	1	3	6
White, both sexes	432	1	-	-	1	1	-	1	1	2	6	9
Male	179	1	-	-	-	1	-	-	-	1	4	5
Female	253	-	-	-	1	-	-	1	1	1	2	4
Black, both sexes	61	-	-	-	-	-	-	-	1	1	4	3
Male	33	-	-	-	-	-	-	-	-	1	3	1
Female	28	-	-	-	-	-	-	-	1	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	143	2	-	-	-	1	-	-	-	1	4	7
Male	71	2	-	-	-	1	-	-	-	-	3	5
Female	72	-	-	-	-	-	-	-	-	1	1	2
White, both sexes	105	-	-	-	-	1	-	-	-	-	2	2
Male	48	-	-	-	-	1	-	-	-	-	2	1
Female	57	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	38	2	-	-	-	-	-	-	-	1	2	5
Male	23	2	-	-	-	-	-	-	-	-	1	4
Female	15	-	-	-	-	-	-	-	-	1	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Hawaii												
All races, both sexes	42	-	-	-	-	-	-	-	-	-	-	1
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	33	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	1
Idaho												
All races, both sexes	26	-	-	-	-	-	-	-	1	-	-	1
Male	14	-	-	-	-	-	-	-	1	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	25	-	-	-	-	-	-	-	1	-	-	1
Male	14	-	-	-	-	-	-	-	1	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Illinois												
All races, both sexes	300	1	-	-	-	-	-	-	-	2	3	7
Male	114	1	-	-	-	-	-	-	-	1	1	4
Female	186	-	-	-	-	-	-	-	-	1	2	3
White, both sexes	257	1	-	-	-	-	-	-	-	2	3	4
Male	96	1	-	-	-	-	-	-	-	1	1	3
Female	161	-	-	-	-	-	-	-	-	1	2	1
Black, both sexes	35	-	-	-	-	-	-	-	-	-	-	3
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	20	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	142	-	-	-	-	-	-	-	-	-	3	7
Male	59	-	-	-	-	-	-	-	-	-	1	6
Female	83	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	133	-	-	-	-	-	-	-	-	-	3	6
Male	55	-	-	-	-	-	-	-	-	-	1	5
Female	78	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Iowa												
All races, both sexes	83	-	-	-	-	-	-	-	1	-	-	1
Male	35	-	-	-	-	-	-	-	-	-	-	1
Female	48	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	82	-	-	-	-	-	-	-	1	-	-	1
Male	35	-	-	-	-	-	-	-	-	-	-	1
Female	47	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	49	-	-	-	-	-	-	-	-	1	1	1
Male	19	-	-	-	-	-	-	-	-	1	1	1
Female	30	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	45	-	-	-	-	-	-	-	-	1	1	1
Male	17	-	-	-	-	-	-	-	-	1	1	1
Female	28	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Kentucky												
All races, both sexes	101	1	-	-	-	-	-	-	-	-	1	5
Male	41	1	-	-	-	-	-	-	-	-	1	2
Female	60	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	98	1	-	-	-	-	-	-	-	-	1	5
Male	40	1	-	-	-	-	-	-	-	-	1	2
Female	58	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	92	-	-	-	-	-	-	-	-	1	2	1
Male	35	-	-	-	-	-	-	-	-	1	1	-
Female	57	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	66	-	-	-	-	-	-	-	-	-	1	-
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	44	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	25	-	-	-	-	-	-	-	-	1	1	1
Male	13	-	-	-	-	-	-	-	-	1	1	-
Female	12	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Maine												
All races, both sexes	26	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	26	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	159	-	-	-	1	-	1	-	-	-	1	2
Male	54	-	-	-	-	-	-	-	-	-	1	2
Female	105	-	-	-	1	-	1	-	-	-	-	-
White, both sexes	123	-	-	-	1	-	-	-	-	-	-	2
Male	38	-	-	-	-	-	-	-	-	-	-	2
Female	85	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	34	-	-	-	-	-	1	-	-	-	1	-
Male	15	-	-	-	-	-	-	-	-	-	1	-
Female	19	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Massachusetts												
All races, both sexes	134	-	-	-	-	-	-	-	-	1	-	1
Male	47	-	-	-	-	-	-	-	-	1	-	-
Female	87	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	131	-	-	-	-	-	-	-	-	1	-	1
Male	46	-	-	-	-	-	-	-	-	1	-	-
Female	85	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	332	1	-	-	-	-	-	-	-	3	6	6
Male	135	1	-	-	-	-	-	-	-	2	3	4
Female	197	-	-	-	-	-	-	-	-	1	3	2
White, both sexes	289	1	-	-	-	-	-	-	-	1	3	3
Male	120	1	-	-	-	-	-	-	-	1	2	2
Female	169	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	40	-	-	-	-	-	-	-	-	2	3	3
Male	15	-	-	-	-	-	-	-	-	1	1	2
Female	25	-	-	-	-	-	-	-	-	1	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Minnesota												
All races, both sexes	114	-	-	-	-	2	-	-	-	-	-	1
Male	37	-	-	-	-	1	-	-	-	-	-	-
Female	77	-	-	-	-	1	-	-	-	-	-	1
White, both sexes	113	-	-	-	-	2	-	-	-	-	-	1
Male	37	-	-	-	-	1	-	-	-	-	-	-
Female	76	-	-	-	-	1	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	70	-	-	1	-	1	-	1	-	-	2	4
Male	43	-	-	-	-	1	-	1	-	-	1	2
Female	27	-	-	1	-	-	-	-	-	-	1	2
White, both sexes	47	-	-	1	-	-	-	1	-	-	1	1
Male	28	-	-	-	-	-	-	1	-	-	-	-
Female	19	-	-	1	-	-	-	-	-	-	1	1
Black, both sexes	23	-	-	-	-	1	-	-	-	-	1	3
Male	15	-	-	-	-	1	-	-	-	-	1	2
Female	8	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Missouri												
All races, both sexes	165	-	-	-	-	-	-	-	1	3	4	8
Male	65	-	-	-	-	-	-	-	1	3	3	3
Female	100	-	-	-	-	-	-	-	-	-	1	5
White, both sexes	141	-	-	-	-	-	-	-	-	1	3	7
Male	54	-	-	-	-	-	-	-	-	1	2	2
Female	87	-	-	-	-	-	-	-	-	-	1	5
Black, both sexes	23	-	-	-	-	-	-	-	1	2	1	1
Male	11	-	-	-	-	-	-	-	1	2	1	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Nebraska												
All races, both sexes	35	-	-	-	-	-	-	1	-	-	-	-
Male	12	-	-	-	-	-	-	1	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	30	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	1	-	-	-	-
Male	3	-	-	-	-	-	-	1	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	26	-	-	-	-	-	-	1	-	1	-	-
Male	17	-	-	-	-	-	-	-	-	1	-	-
Female	9	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	23	-	-	-	-	-	-	1	-	1	-	-
Male	16	-	-	-	-	-	-	-	-	1	-	-
Female	7	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
New Hampshire												
All races, both sexes	22	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	131	1	-	-	-	-	-	-	1	-	-	1
Male	51	1	-	-	-	-	-	-	1	-	-	1
Female	80	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	117	-	-	-	-	-	-	-	1	-	-	1
Male	46	-	-	-	-	-	-	-	1	-	-	1
Female	71	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	12	1	-	-	-	-	-	-	-	-	-	-
Male	4	1	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
New Mexico												
All races, both sexes	44	-	-	-	-	-	-	-	-	-	-	1
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	42	-	-	-	-	-	-	-	-	-	-	1
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	312	-	-	1	1	1	1	-	3	-	6	7
Male	130	-	-	-	1	-	1	-	3	-	4	4
Female	182	-	-	1	-	1	-	-	-	-	2	3
White, both sexes	258	-	-	-	1	-	1	-	2	-	4	6
Male	112	-	-	-	1	-	1	-	2	-	4	4
Female	146	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	42	-	-	-	-	1	-	-	1	-	2	1
Male	14	-	-	-	-	-	-	-	1	-	-	-
Female	28	-	-	-	-	1	-	-	-	-	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	1	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
North Carolina												
All races, both sexes	260	1	-	-	-	1	-	-	1	-	3	5
Male	108	-	-	-	-	1	-	-	1	-	2	3
Female	152	1	-	-	-	-	-	-	-	-	1	2
White, both sexes	203	1	-	-	-	1	-	-	1	-	1	3
Male	79	-	-	-	-	1	-	-	1	-	1	2
Female	124	1	-	-	-	-	-	-	-	-	-	1
Black, both sexes	56	-	-	-	-	-	-	-	-	-	2	2
Male	29	-	-	-	-	-	-	-	-	-	1	1
Female	27	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Ohio												
All races, both sexes	371	1	-	-	-	-	-	-	1	5	2	10
Male	147	-	-	-	-	-	-	-	-	2	1	4
Female	224	1	-	-	-	-	-	-	1	3	1	6
White, both sexes	335	-	-	-	-	-	-	-	1	3	1	7
Male	131	-	-	-	-	-	-	-	-	-	-	2
Female	204	-	-	-	-	-	-	-	1	3	1	5
Black, both sexes	36	1	-	-	-	-	-	-	-	2	1	3
Male	16	-	-	-	-	-	-	-	-	2	1	2
Female	20	1	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	102	-	-	-	-	-	-	-	2	-	3	2
Male	48	-	-	-	-	-	-	-	1	-	1	1
Female	54	-	-	-	-	-	-	-	1	-	2	1
White, both sexes	88	-	-	-	-	-	-	-	1	-	1	1
Male	41	-	-	-	-	-	-	-	-	-	-	-
Female	47	-	-	-	-	-	-	-	1	-	1	1
Black, both sexes	8	-	-	-	-	-	-	-	1	-	2	1
Male	4	-	-	-	-	-	-	-	1	-	1	1
Female	4	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Oregon												
All races, both sexes	83	-	-	-	-	-	-	-	-	-	1	3
Male	37	-	-	-	-	-	-	-	-	-	-	3
Female	46	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	77	-	-	-	-	-	-	-	-	-	1	1
Male	35	-	-	-	-	-	-	-	-	-	-	1
Female	42	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	411	-	-	-	-	-	-	-	1	1	3	9
Male	161	-	-	-	-	-	-	-	-	1	3	3
Female	250	-	-	-	-	-	-	-	1	-	-	6
White, both sexes	362	-	-	-	-	-	-	-	1	1	3	5
Male	142	-	-	-	-	-	-	-	-	1	3	2
Female	220	-	-	-	-	-	-	-	1	-	-	3
Black, both sexes	43	-	-	-	-	-	-	-	-	-	-	3
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Rhode Island												
All races, both sexes	16	-	-	-	-	-	-	-	1	-	-	-
Male	10	-	-	-	-	-	-	-	1	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	1	-	-	-
Male	9	-	-	-	-	-	-	-	1	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	156	-	-	-	-	-	-	1	2	2	-	2
Male	61	-	-	-	-	-	-	-	1	2	-	1
Female	95	-	-	-	-	-	-	1	1	-	-	1
White, both sexes	116	-	-	-	-	-	-	-	1	-	-	-
Male	46	-	-	-	-	-	-	-	1	-	-	-
Female	70	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	40	-	-	-	-	-	-	1	1	2	-	2
Male	15	-	-	-	-	-	-	-	-	2	-	1
Female	25	-	-	-	-	-	-	1	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
South Dakota												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	182	-	-	-	-	-	-	-	-	2	1	5
Male	79	-	-	-	-	-	-	-	-	-	-	3
Female	103	-	-	-	-	-	-	-	-	2	1	2
White, both sexes	157	-	-	-	-	-	-	-	-	1	1	4
Male	68	-	-	-	-	-	-	-	-	-	-	3
Female	89	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	25	-	-	-	-	-	-	-	-	1	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Texas												
All races, both sexes	406	1	-	-	-	-	-	2	5	2	11	6
Male	163	-	-	-	-	-	-	-	3	1	7	4
Female	243	1	-	-	-	-	-	2	2	1	4	2
White, both sexes	346	1	-	-	-	-	-	2	4	2	5	4
Male	135	-	-	-	-	-	-	-	2	1	4	2
Female	211	1	-	-	-	-	-	2	2	1	1	2
Black, both sexes	55	-	-	-	-	-	-	-	-	-	5	2
Male	25	-	-	-	-	-	-	-	-	-	2	2
Female	30	-	-	-	-	-	-	-	-	-	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	1	-	1	-
Male	3	-	-	-	-	-	-	-	1	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	22	-	-	-	-	-	-	-	-	1	1	-
Male	10	-	-	-	-	-	-	-	-	-	1	-
Female	12	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	21	-	-	-	-	-	-	-	-	1	1	-
Male	9	-	-	-	-	-	-	-	-	-	1	-
Female	12	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Vermont												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	167	1	1	1	-	-	-	-	-	-	2	1
Male	63	1	-	-	-	-	-	-	-	-	2	1
Female	104	-	1	1	-	-	-	-	-	-	-	-
White, both sexes	124	-	1	1	-	-	-	-	-	-	1	1
Male	48	-	-	-	-	-	-	-	-	-	1	1
Female	76	-	1	1	-	-	-	-	-	-	-	-
Black, both sexes	41	1	-	-	-	-	-	-	-	-	1	-
Male	14	1	-	-	-	-	-	-	-	-	1	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Washington												
All races, both sexes	140	-	-	-	-	-	-	-	-	-	1	1
Male	51	-	-	-	-	-	-	-	-	-	1	1
Female	89	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	132	-	-	-	-	-	-	-	-	-	1	-
Male	48	-	-	-	-	-	-	-	-	-	1	-
Female	84	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	49	1	-	-	-	-	-	-	-	-	-	1
Male	19	1	-	-	-	-	-	-	-	-	-	1
Female	30	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	46	1	-	-	-	-	-	-	-	-	-	1
Male	19	1	-	-	-	-	-	-	-	-	-	1
Female	27	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Wisconsin												
All races, both sexes	139	1	-	-	-	-	1	-	-	2	2	5
Male	63	1	-	-	-	-	1	-	-	2	1	4
Female	76	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	132	1	-	-	-	-	1	-	-	2	2	5
Male	61	1	-	-	-	-	1	-	-	2	1	4
Female	71	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
United States												
All races, both sexes	202	268	353	401	555	820	1,167	1,244	952	343	65	1
Male	127	164	211	244	281	389	430	407	249	54	6	1
Female	75	104	142	157	274	431	737	837	703	289	59	-
White, both sexes	150	201	278	312	467	705	1,038	1,127	870	312	59	1
Male	97	123	169	187	242	337	389	376	226	50	6	1
Female	53	78	109	125	225	368	649	751	644	262	53	-
Black, both sexes	49	58	70	75	68	90	105	86	64	24	4	-
Male	29	35	38	50	27	42	34	20	17	3	-	-
Female	20	23	32	25	41	48	71	66	47	21	4	-
American Indian 1/, both sexes	-	1	-	-	6	3	3	4	3	1	-	-
Male	-	1	-	-	3	1	-	2	1	-	-	-
Female	-	-	-	-	3	2	3	2	2	1	-	-
Asian or Pacific Islander 2/, both sexes	3	8	5	14	14	22	21	27	15	6	2	-
Male	1	5	4	7	9	9	7	9	5	1	-	-
Female	2	3	1	7	5	13	14	18	10	5	2	-
Alabama												
All races, both sexes	4	4	6	12	11	8	26	23	9	3	-	-
Male	2	2	3	8	5	4	7	9	1	-	-	-
Female	2	2	3	4	6	4	19	14	8	3	-	-
White, both sexes	1	3	5	9	9	6	22	20	6	3	-	-
Male	-	1	3	6	3	2	6	8	-	-	-	-
Female	1	2	2	3	6	4	16	12	6	3	-	-
Black, both sexes	3	1	1	3	2	2	4	3	3	-	-	-
Male	2	1	-	2	2	2	1	1	1	-	-	-
Female	1	-	1	1	-	-	3	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Alaska												
All races, both sexes	-	-	-	-	-	-	2	3	1	1	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	2	2	1	1	-	-
White, both sexes	-	-	-	-	-	-	2	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	2	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2	7	9	14	7	10	30	19	15	1	-	-
Male	2	5	5	9	5	5	3	5	6	1	-	-
Female	-	2	4	5	2	5	27	14	9	-	-	-
White, both sexes	2	7	9	13	6	10	28	19	15	1	-	-
Male	2	5	5	9	4	5	2	5	6	1	-	-
Female	-	2	4	4	2	5	26	14	9	-	-	-
Black, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Arkansas												
All races, both sexes	5	-	5	5	6	11	13	12	8	5	-	-
Male	4	-	3	5	-	5	5	5	-	2	-	-
Female	1	-	2	-	6	6	8	7	8	3	-	-
White, both sexes	5	-	5	3	5	9	11	11	7	3	-	-
Male	4	-	3	3	-	4	5	5	-	1	-	-
Female	1	-	2	-	5	5	6	6	7	2	-	-
Black, both sexes	-	-	-	2	1	2	2	1	1	2	-	-
Male	-	-	-	2	-	1	-	-	-	1	-	-
Female	-	-	-	-	1	1	2	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	25	22	30	34	56	66	94	123	77	37	3	-
Male	15	16	18	16	37	32	38	41	27	5	-	-
Female	10	6	12	18	19	34	56	82	50	32	3	-
White, both sexes	19	14	24	24	48	58	78	103	70	33	2	-
Male	12	11	14	12	32	26	32	33	23	5	-	-
Female	7	3	10	12	16	32	46	70	47	28	2	-
Black, both sexes	4	5	3	6	2	1	8	6	4	1	-	-
Male	2	3	1	2	1	1	4	3	1	-	-	-
Female	2	2	2	4	1	-	4	3	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	2	3	3	4	6	7	8	14	3	2	1	-
Male	1	2	3	2	4	5	2	5	3	-	-	-
Female	1	1	-	2	2	2	6	9	-	2	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Colorado												
All races, both sexes	3	1	6	5	7	7	18	20	6	3	-	-
Male	2	1	4	4	3	2	4	8	1	-	-	-
Female	1	-	2	1	4	5	14	12	5	3	-	-
White, both sexes	2	1	6	5	7	7	18	20	6	3	-	-
Male	2	1	4	4	3	2	4	8	1	-	-	-
Female	-	-	2	1	4	5	14	12	5	3	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	2	-	4	8	6	16	7	17	14	8	1	-
Male	1	-	1	3	5	8	3	9	3	1	-	-
Female	1	-	3	5	1	8	4	8	11	7	1	-
White, both sexes	1	-	4	8	6	10	6	16	13	7	1	-
Male	-	-	1	3	5	6	3	9	3	1	-	-
Female	1	-	3	5	1	4	3	7	10	6	1	-
Black, both sexes	1	-	-	-	-	5	1	1	1	1	-	-
Male	1	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	3	1	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Delaware												
All races, both sexes	-	-	2	-	3	1	4	3	-	2	1	-
Male	-	-	-	-	2	-	2	1	-	-	-	-
Female	-	-	2	-	1	1	2	2	-	2	1	-
White, both sexes	-	-	1	-	3	1	3	3	-	1	1	-
Male	-	-	-	-	2	-	2	1	-	-	-	-
Female	-	-	1	-	1	1	1	2	-	1	1	-
Black, both sexes	-	-	1	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	2	2	2	2	-	4	3	-	1	-	-	-
Male	2	1	1	-	-	2	1	-	-	-	-	-
Female	-	1	1	2	-	2	2	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	2	2	2	2	-	4	3	-	-	-	-	-
Male	2	1	1	-	-	2	1	-	-	-	-	-
Female	-	1	1	2	-	2	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Florida												
All races, both sexes	13	19	24	23	49	57	84	88	74	27	5	-
Male	11	10	16	12	23	31	34	29	23	6	-	-
Female	2	9	8	11	26	26	50	59	51	21	5	-
White, both sexes	8	16	18	17	40	51	76	82	71	26	5	-
Male	7	7	12	9	20	29	31	26	21	5	-	-
Female	1	9	6	8	20	22	45	56	50	21	5	-
Black, both sexes	4	3	6	6	9	6	8	6	3	1	-	-
Male	4	3	4	3	3	2	3	3	2	1	-	-
Female	-	-	2	3	6	4	5	3	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	5	11	8	13	19	17	19	22	10	4	-	-
Male	2	9	7	10	9	7	4	6	6	-	-	-
Female	3	2	1	3	10	10	15	16	4	4	-	-
White, both sexes	3	7	5	9	12	13	18	20	10	3	-	-
Male	1	5	5	7	6	4	4	6	6	-	-	-
Female	2	2	-	2	6	9	14	14	4	3	-	-
Black, both sexes	2	4	3	4	7	4	1	2	-	1	-	-
Male	1	4	2	3	3	3	-	-	-	-	-	-
Female	1	-	1	1	4	1	1	2	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Hawaii												
All races, both sexes	-	-	2	4	1	4	6	14	7	2	1	-
Male	-	-	2	3	1	2	3	7	2	1	-	-
Female	-	-	-	1	-	2	3	7	5	1	1	-
White, both sexes	-	-	1	2	-	1	2	3	-	-	-	-
Male	-	-	1	2	-	1	2	3	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	2	1	3	4	11	7	2	1	-
Male	-	-	1	1	1	1	1	4	2	1	-	-
Female	-	-	-	1	-	2	3	7	5	1	1	-
Idaho												
All races, both sexes	1	1	2	1	2	4	5	7	1	-	-	-
Male	-	1	1	1	1	3	2	4	-	-	-	-
Female	1	-	1	-	1	1	3	3	1	-	-	-
White, both sexes	1	1	2	1	2	3	5	7	1	-	-	-
Male	-	1	1	1	1	3	2	4	-	-	-	-
Female	1	-	1	-	1	-	3	3	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Illinois												
All races, both sexes	6	8	15	16	22	37	61	53	44	23	2	-
Male	4	6	7	9	8	18	28	15	7	5	-	-
Female	2	2	8	7	14	19	33	38	37	18	2	-
White, both sexes	6	4	12	9	18	33	54	50	39	20	2	-
Male	4	4	6	4	7	16	24	14	6	5	-	-
Female	2	-	6	5	11	17	30	36	33	15	2	-
Black, both sexes	-	3	2	5	4	4	5	3	5	1	-	-
Male	-	1	1	4	1	2	3	1	1	-	-	-
Female	-	2	1	1	3	2	2	2	4	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	2	-	-	2	-	-	2	-	-
Male	-	1	-	1	-	-	1	-	-	-	-	-
Female	-	-	1	1	-	-	1	-	-	2	-	-
Indiana												
All races, both sexes	8	12	4	7	14	12	18	26	22	6	3	-
Male	6	6	3	5	9	7	3	6	6	1	-	-
Female	2	6	1	2	5	5	15	20	16	5	3	-
White, both sexes	7	11	4	6	12	12	17	24	22	6	3	-
Male	5	5	3	5	8	7	3	6	6	1	-	-
Female	2	6	1	1	4	5	14	18	16	5	3	-
Black, both sexes	1	1	-	1	2	-	1	2	-	-	-	-
Male	1	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Iowa												
All races, both sexes	1	1	4	3	9	12	19	14	10	4	4	-
Male	1	1	2	2	7	6	7	5	2	1	-	-
Female	-	-	2	1	2	6	12	9	8	3	4	-
White, both sexes	1	1	4	3	8	12	19	14	10	4	4	-
Male	1	1	2	2	7	6	7	5	2	1	-	-
Female	-	-	2	1	1	6	12	9	8	3	4	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	4	3	3	4	13	4	7	6	1	-	-
Male	-	2	-	2	1	4	2	3	2	-	-	-
Female	1	2	3	1	3	9	2	4	4	1	-	-
White, both sexes	1	3	3	3	4	11	4	7	5	1	-	-
Male	-	2	-	2	1	2	2	3	2	-	-	-
Female	1	1	3	1	3	9	2	4	3	1	-	-
Black, both sexes	-	1	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Kentucky												
All races, both sexes	1	6	9	6	8	13	19	17	7	7	1	-
Male	1	4	5	4	4	4	5	5	2	3	-	-
Female	-	2	4	2	4	9	14	12	5	4	1	-
White, both sexes	1	5	9	6	8	12	19	16	7	7	1	-
Male	1	3	5	4	4	4	5	5	2	3	-	-
Female	-	2	4	2	4	8	14	11	5	4	1	-
Black, both sexes	-	1	-	-	-	1	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	2	4	10	8	6	14	13	14	11	3	3	-
Male	1	4	6	6	3	6	5	2	-	-	-	-
Female	1	-	4	2	3	8	8	12	11	3	3	-
White, both sexes	2	2	5	6	4	12	9	10	10	3	2	-
Male	1	2	3	4	2	5	4	1	-	-	-	-
Female	1	-	2	2	2	7	5	9	10	3	2	-
Black, both sexes	-	2	5	2	2	1	4	4	1	-	1	-
Male	-	2	3	2	1	1	1	1	-	-	-	-
Female	-	-	2	-	1	-	3	3	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Maine												
All races, both sexes	2	-	-	2	1	4	3	9	4	1	-	-
Male	2	-	-	1	-	2	3	3	-	-	-	-
Female	-	-	-	1	1	2	-	6	4	1	-	-
White, both sexes	2	-	-	2	1	4	3	9	4	1	-	-
Male	2	-	-	1	-	2	3	3	-	-	-	-
Female	-	-	-	1	1	2	-	6	4	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	4	6	8	-	10	19	27	29	33	14	3	1
Male	3	4	4	-	6	4	8	11	7	3	-	1
Female	1	2	4	-	4	15	19	18	26	11	3	-
White, both sexes	1	3	5	-	7	14	22	24	28	13	2	1
Male	-	1	2	-	4	3	7	9	6	3	-	1
Female	1	2	3	-	3	11	15	15	22	10	2	-
Black, both sexes	3	2	3	-	3	5	5	4	5	1	1	-
Male	3	2	2	-	2	1	1	2	1	-	-	-
Female	-	-	1	-	1	4	4	2	4	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Massachusetts												
All races, both sexes	2	3	4	5	8	16	28	31	23	8	4	-
Male	-	-	2	4	4	6	13	13	2	2	-	-
Female	2	3	2	1	4	10	15	18	21	6	4	-
White, both sexes	2	3	4	5	8	15	26	31	23	8	4	-
Male	-	-	2	4	4	6	12	13	2	2	-	-
Female	2	3	2	1	4	9	14	18	21	6	4	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Michigan												
All races, both sexes	10	14	12	18	24	40	54	80	43	19	2	-
Male	8	10	10	9	14	19	19	22	11	2	1	-
Female	2	4	2	9	10	21	35	58	32	17	1	-
White, both sexes	8	9	10	15	20	36	54	73	38	17	1	-
Male	7	7	8	9	13	17	19	21	10	2	1	-
Female	1	2	2	6	7	19	35	52	28	15	-	-
Black, both sexes	2	5	2	2	3	4	-	7	4	2	1	-
Male	1	3	2	-	1	2	-	1	1	-	-	-
Female	1	2	-	2	2	2	-	6	3	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Minnesota												
All races, both sexes	4	3	5	4	11	10	19	27	17	11	-	-
Male	1	2	3	4	7	5	2	9	3	-	-	-
Female	3	1	2	-	4	5	17	18	14	11	-	-
White, both sexes	4	3	5	4	11	10	19	27	16	11	-	-
Male	1	2	3	4	7	5	2	9	3	-	-	-
Female	3	1	2	-	4	5	17	18	13	11	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	4	5	5	5	9	11	9	2	7	3	1	-
Male	4	3	3	4	6	6	7	2	2	1	-	-
Female	-	2	2	1	3	5	2	-	5	2	1	-
White, both sexes	3	2	2	4	8	7	6	2	5	3	1	-
Male	3	-	2	3	5	4	5	2	2	1	-	-
Female	-	2	-	1	3	3	1	-	3	2	1	-
Black, both sexes	1	3	3	1	1	4	3	-	2	-	-	-
Male	1	3	1	1	1	2	2	-	-	-	-	-
Female	-	-	2	-	-	2	1	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Missouri												
All races, both sexes	4	6	9	8	6	21	28	36	24	6	1	-
Male	1	3	6	3	3	10	10	12	7	-	-	-
Female	3	3	3	5	3	11	18	24	17	6	1	-
White, both sexes	2	4	8	5	5	16	25	35	23	6	1	-
Male	1	3	5	1	2	8	10	12	7	-	-	-
Female	1	1	3	4	3	8	15	23	16	6	1	-
Black, both sexes	2	2	1	3	1	5	2	1	1	-	-	-
Male	-	-	1	2	1	2	-	-	-	-	-	-
Female	2	2	-	1	-	3	2	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Montana												
All races, both sexes	-	1	1	-	1	2	2	1	2	-	-	-
Male	-	1	-	-	-	2	2	-	-	-	-	-
Female	-	-	1	-	1	-	-	1	2	-	-	-
White, both sexes	-	1	1	-	1	1	2	1	2	-	-	-
Male	-	1	-	-	-	1	2	-	-	-	-	-
Female	-	-	1	-	1	-	-	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Nebraska												
All races, both sexes	1	2	2	1	2	4	5	9	5	3	-	-
Male	-	1	1	1	2	3	1	1	-	1	-	-
Female	1	1	1	-	-	1	4	8	5	2	-	-
White, both sexes	1	2	1	1	1	4	5	7	5	3	-	-
Male	-	1	-	1	1	3	1	1	-	1	-	-
Female	1	1	1	-	-	1	4	6	5	2	-	-
Black, both sexes	-	-	1	-	1	-	-	2	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	3	3	2	3	1	4	3	5	-	-	-
Male	-	3	2	2	2	1	3	-	3	-	-	-
Female	-	-	1	-	1	-	1	3	2	-	-	-
White, both sexes	-	3	3	1	3	1	4	2	4	-	-	-
Male	-	3	2	1	2	1	3	-	3	-	-	-
Female	-	-	1	-	1	-	1	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
New Hampshire												
All races, both sexes	-	2	-	-	3	-	3	3	7	3	-	-
Male	-	1	-	-	2	-	1	-	3	1	-	-
Female	-	1	-	-	1	-	2	3	4	2	-	-
White, both sexes	-	2	-	-	3	-	3	3	7	3	-	-
Male	-	1	-	-	2	-	1	-	3	1	-	-
Female	-	1	-	-	1	-	2	3	4	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	5	8	3	6	10	22	23	26	14	11	-	-
Male	4	4	3	4	3	8	5	11	5	1	-	-
Female	1	4	-	2	7	14	18	15	9	10	-	-
White, both sexes	3	6	3	4	9	22	20	26	12	10	-	-
Male	3	4	3	2	3	8	4	11	5	1	-	-
Female	-	2	-	2	6	14	16	15	7	9	-	-
Black, both sexes	2	1	-	2	1	-	2	-	2	1	-	-
Male	1	-	-	2	-	-	-	-	-	-	-	-
Female	1	1	-	-	1	-	2	-	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
New Mexico												
All races, both sexes	-	4	4	2	3	5	11	8	5	1	-	-
Male	-	3	2	1	3	4	2	4	2	-	-	-
Female	-	1	2	1	-	1	9	4	3	1	-	-
White, both sexes	-	4	4	2	1	5	11	8	5	1	-	-
Male	-	3	2	1	1	4	2	4	2	-	-	-
Female	-	1	2	1	-	1	9	4	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	2	-	-	-	-	-	-	-
Male	-	-	-	-	2	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	9	10	18	14	25	47	46	56	46	15	6	-
Male	7	6	9	11	13	19	16	18	15	3	-	-
Female	2	4	9	3	12	28	30	38	31	12	6	-
White, both sexes	5	8	13	12	19	36	39	50	44	13	5	-
Male	4	5	6	9	10	14	16	18	15	3	-	-
Female	1	3	7	3	9	22	23	32	29	10	5	-
Black, both sexes	4	2	5	2	2	6	6	5	2	2	1	-
Male	3	1	3	2	-	4	-	-	-	-	-	-
Female	1	1	2	-	2	2	6	5	2	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	4	5	1	1	-	-	-	-
Male	-	-	-	-	3	1	-	-	-	-	-	-
Female	-	-	-	-	1	4	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
North Carolina												
All races, both sexes	7	8	21	25	16	31	51	37	40	11	2	-
Male	3	7	10	16	5	15	18	17	8	1	1	-
Female	4	1	11	9	11	16	33	20	32	10	1	-
White, both sexes	6	5	14	16	14	26	40	32	35	6	2	-
Male	3	4	8	9	4	13	12	14	6	-	1	-
Female	3	1	6	7	10	13	28	18	29	6	1	-
Black, both sexes	1	3	7	8	2	5	11	5	5	5	-	-
Male	-	3	2	7	1	2	6	3	2	1	-	-
Female	1	-	5	1	1	3	5	2	3	4	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	2	-	2	3	3	2	-	-	-
Male	-	-	-	1	-	1	1	1	-	-	-	-
Female	-	-	-	1	-	1	2	2	2	-	-	-
White, both sexes	-	-	-	2	-	1	3	3	2	-	-	-
Male	-	-	-	1	-	1	1	1	-	-	-	-
Female	-	-	-	1	-	-	2	2	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Ohio												
All races, both sexes	13	12	18	22	34	47	74	49	61	17	5	-
Male	8	6	10	13	16	29	33	10	14	-	1	-
Female	5	6	8	9	18	18	41	39	47	17	4	-
White, both sexes	12	8	17	18	32	45	66	46	58	16	5	-
Male	8	5	10	11	15	28	29	9	13	-	1	-
Female	4	3	7	7	17	17	37	37	45	16	4	-
Black, both sexes	1	4	1	4	2	2	8	3	3	1	-	-
Male	-	1	-	2	1	1	4	1	1	-	-	-
Female	1	3	1	2	1	1	4	2	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	6	6	1	3	8	14	23	19	12	2	1	-
Male	5	4	1	1	3	3	10	11	6	-	1	-
Female	1	2	-	2	5	11	13	8	6	2	-	-
White, both sexes	5	5	1	3	7	13	20	17	11	2	1	-
Male	4	3	1	1	3	3	10	10	5	-	1	-
Female	1	2	-	2	4	10	10	7	6	2	-	-
Black, both sexes	1	-	-	-	-	1	2	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	2	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	1	-	1	2	1	-	-	-
Male	-	1	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	1	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Oregon												
All races, both sexes	1	3	-	7	6	8	17	13	17	7	-	-
Male	1	1	-	3	3	5	9	6	5	1	-	-
Female	-	2	-	4	3	3	8	7	12	6	-	-
White, both sexes	1	2	-	6	6	8	17	12	16	7	-	-
Male	1	1	-	3	3	5	9	6	5	1	-	-
Female	-	1	-	3	3	3	8	6	11	6	-	-
Black, both sexes	-	-	-	1	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	13	13	18	20	26	62	82	81	60	17	5	-
Male	6	8	10	14	10	26	34	24	18	3	1	-
Female	7	5	8	6	16	36	48	57	42	14	4	-
White, both sexes	9	10	13	13	23	54	74	78	58	15	5	-
Male	4	5	7	8	10	24	32	24	18	3	1	-
Female	5	5	6	5	13	30	42	54	40	12	4	-
Black, both sexes	4	2	5	5	3	7	7	3	2	2	-	-
Male	2	2	3	4	-	2	2	-	-	-	-	-
Female	2	-	2	1	3	5	5	3	2	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	2	-	1	1	-	-	-	-	-
Male	-	1	-	2	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Rhode Island												
All races, both sexes	-	-	1	1	2	3	2	2	3	1	-	-
Male	-	-	1	1	1	2	1	1	2	-	-	-
Female	-	-	-	-	1	1	1	1	1	1	-	-
White, both sexes	-	-	1	1	2	3	2	2	2	1	-	-
Male	-	-	1	1	1	2	1	1	1	-	-	-
Female	-	-	-	-	1	1	1	1	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	3	3	8	11	5	22	30	33	26	8	-	-
Male	1	2	4	6	2	10	9	14	8	1	-	-
Female	2	1	4	5	3	12	21	19	18	7	-	-
White, both sexes	1	2	4	8	3	16	25	26	22	8	-	-
Male	-	2	2	5	1	8	8	13	5	1	-	-
Female	1	-	2	3	2	8	17	13	17	7	-	-
Black, both sexes	2	1	4	3	2	6	5	7	4	-	-	-
Male	1	-	2	1	1	2	1	1	3	-	-	-
Female	1	1	2	2	1	4	4	6	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
South Dakota												
All races, both sexes	-	1	-	2	3	1	2	4	2	-	-	-
Male	-	-	-	-	1	1	2	3	-	-	-	-
Female	-	1	-	2	2	-	-	1	2	-	-	-
White, both sexes	-	1	-	2	2	1	2	4	1	-	-	-
Male	-	-	-	-	1	1	2	3	-	-	-	-
Female	-	1	-	2	1	-	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	7	11	12	14	19	21	24	38	24	3	1	-
Male	4	6	8	10	11	13	9	7	8	-	-	-
Female	3	5	4	4	8	8	15	31	16	3	1	-
White, both sexes	5	8	9	13	17	20	21	33	21	3	1	-
Male	3	4	6	9	10	12	7	7	7	-	-	-
Female	2	4	3	4	7	8	14	26	14	3	1	-
Black, both sexes	2	3	3	1	2	1	3	5	3	-	-	-
Male	1	2	2	1	1	1	2	-	1	-	-	-
Female	1	1	1	-	1	-	1	5	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Texas												
All races, both sexes	13	20	24	31	39	41	65	63	57	20	6	-
Male	7	8	17	18	17	20	26	21	9	5	-	-
Female	6	12	7	13	22	21	39	42	48	15	6	-
White, both sexes	9	16	18	24	30	35	59	57	54	20	6	-
Male	5	8	12	12	14	18	24	19	9	5	-	-
Female	4	8	6	12	16	17	35	38	45	15	6	-
Black, both sexes	4	4	6	6	9	5	5	6	3	-	-	-
Male	2	-	5	6	3	1	2	2	-	-	-	-
Female	2	4	1	-	6	4	3	4	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Utah												
All races, both sexes	-	-	1	1	2	6	3	3	4	-	-	-
Male	-	-	1	1	1	3	1	1	1	-	-	-
Female	-	-	-	-	1	3	2	2	3	-	-	-
White, both sexes	-	-	1	1	1	6	3	3	4	-	-	-
Male	-	-	1	1	-	3	1	1	1	-	-	-
Female	-	-	-	-	1	3	2	2	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Vermont												
All races, both sexes	-	-	-	1	1	2	1	4	2	1	-	-
Male	-	-	-	1	-	2	1	-	1	-	-	-
Female	-	-	-	-	1	-	-	4	1	1	-	-
White, both sexes	-	-	-	1	1	2	1	4	2	1	-	-
Male	-	-	-	1	-	2	1	-	1	-	-	-
Female	-	-	-	-	1	-	-	4	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	2	7	11	11	17	20	25	29	30	7	2	-
Male	1	4	7	6	6	11	9	5	10	-	-	-
Female	1	3	4	5	11	9	16	24	20	7	2	-
White, both sexes	1	4	5	8	13	14	19	26	23	5	2	-
Male	1	3	5	4	4	8	8	5	8	-	-	-
Female	-	1	-	4	9	6	11	21	15	5	2	-
Black, both sexes	1	3	6	3	4	6	5	3	6	2	-	-
Male	-	1	2	2	2	3	-	-	2	-	-	-
Female	1	2	4	1	2	3	5	3	4	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Washington												
All races, both sexes	4	7	7	8	13	18	23	21	27	10	-	-
Male	2	2	5	5	7	8	7	7	5	1	-	-
Female	2	5	2	3	6	10	16	14	22	9	-	-
White, both sexes	4	7	7	7	11	17	23	20	25	10	-	-
Male	2	2	5	4	7	7	7	7	5	1	-	-
Female	2	5	2	3	4	10	16	13	20	9	-	-
Black, both sexes	-	-	-	1	-	1	-	1	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	2	-	-	-
West Virginia												
All races, both sexes	1	3	2	1	6	3	12	14	4	1	-	-
Male	1	2	1	-	3	2	4	3	1	-	-	-
Female	-	1	1	1	3	1	8	11	3	1	-	-
White, both sexes	1	3	2	1	6	3	11	12	4	1	-	-
Male	1	2	1	-	3	2	4	3	1	-	-	-
Female	-	1	1	1	3	1	7	9	3	1	-	-
Black, both sexes	-	-	-	-	-	-	1	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Wisconsin												
All races, both sexes	6	5	10	9	11	8	22	26	23	6	2	-
Male	4	5	7	5	6	2	8	9	5	2	1	-
Female	2	-	3	4	5	6	14	17	18	4	1	-
White, both sexes	5	5	10	8	9	7	21	25	23	6	2	-
Male	4	5	7	4	6	2	8	8	5	2	1	-
Female	1	-	3	4	3	5	13	17	18	4	1	-
Black, both sexes	1	-	-	1	2	-	1	1	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	1	-	-	-	2	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	1	1	3	1	3	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	1	-	2	1	3	-	-	-	-
White, both sexes	-	-	-	1	1	2	1	3	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	2	1	3	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
United States												
All races, both sexes	71,117	4	4	3	2	8	28	23	56	116	290	599
Male	26,805	1	1	2	-	3	16	11	23	62	160	340
Female	44,312	3	3	1	2	5	12	12	33	54	130	259
White, both sexes	60,745	3	3	2	2	6	20	16	32	70	173	378
Male	22,333	1	1	1	-	1	10	8	12	33	103	217
Female	38,412	2	2	1	2	5	10	8	20	37	70	161
Black, both sexes	8,641	1	1	1	-	1	7	6	23	42	107	197
Male	3,739	-	-	1	-	1	5	2	10	27	51	110
Female	4,902	1	1	-	-	-	2	4	13	15	56	87
American Indian 1/, both sexes	277	-	-	-	-	-	-	-	-	1	4	6
Male	125	-	-	-	-	-	-	-	-	1	2	3
Female	152	-	-	-	-	-	-	-	-	-	2	3
Asian or Pacific Islander 2/, both sexes	1,454	-	-	-	-	1	1	1	1	3	6	18
Male	608	-	-	-	-	1	1	1	1	1	4	10
Female	846	-	-	-	-	-	-	-	-	2	2	8
Alabama												
All races, both sexes	1,582	-	-	-	-	1	2	1	5	4	8	24
Male	629	-	-	-	-	1	2	1	3	3	4	12
Female	953	-	-	-	-	-	-	-	2	1	4	12
White, both sexes	1,144	-	-	-	-	-	2	1	-	2	3	17
Male	431	-	-	-	-	-	2	1	-	2	2	8
Female	713	-	-	-	-	-	-	-	-	-	1	9
Black, both sexes	433	-	-	-	-	1	-	-	5	2	5	7
Male	196	-	-	-	-	1	-	-	3	1	2	4
Female	237	-	-	-	-	-	-	-	2	1	3	3
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Alaska												
All races, both sexes	85	-	-	-	-	-	-	-	-	-	-	-
Male	44	-	-	-	-	-	-	-	-	-	-	-
Female	41	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	68	-	-	-	-	-	-	-	-	-	-	-
Male	34	-	-	-	-	-	-	-	-	-	-	-
Female	34	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1,267	-	-	-	-	-	1	1	1	2	5	16
Male	456	-	-	-	-	-	1	1	-	2	3	9
Female	811	-	-	-	-	-	-	-	1	-	2	7
White, both sexes	1,206	-	-	-	-	-	1	1	1	2	5	15
Male	432	-	-	-	-	-	1	1	-	2	3	8
Female	774	-	-	-	-	-	-	-	1	-	2	7
Black, both sexes	29	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	18	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Arkansas												
All races, both sexes	1,049	-	-	-	-	1	1	3	1	5	7	12
Male	390	-	-	-	-	-	-	1	-	2	4	7
Female	659	-	-	-	-	1	1	2	1	3	3	5
White, both sexes	904	-	-	-	-	1	1	3	-	4	7	10
Male	337	-	-	-	-	-	-	1	-	2	4	6
Female	567	-	-	-	-	1	1	2	-	2	3	4
Black, both sexes	140	-	-	-	-	-	-	-	1	1	-	2
Male	52	-	-	-	-	-	-	-	-	-	-	1
Female	88	-	-	-	-	-	-	-	1	1	-	1
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	6,755	-	-	1	-	2	1	2	1	7	25	62
Male	2,678	-	-	-	-	1	1	2	1	2	16	35
Female	4,077	-	-	1	-	1	-	-	-	5	9	27
White, both sexes	5,496	-	-	1	-	1	-	1	1	5	18	37
Male	2,161	-	-	-	-	-	-	1	1	2	12	22
Female	3,335	-	-	1	-	1	-	-	-	3	6	15
Black, both sexes	545	-	-	-	-	-	1	-	-	1	4	15
Male	224	-	-	-	-	-	1	-	-	-	2	6
Female	321	-	-	-	-	-	-	-	-	1	2	9
American Indian 1/, both sexes	40	-	-	-	-	-	-	-	-	-	-	1
Male	17	-	-	-	-	-	-	-	-	-	-	1
Female	23	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	674	-	-	-	-	1	-	1	-	1	3	9
Male	276	-	-	-	-	1	-	1	-	-	2	6
Female	398	-	-	-	-	-	-	-	-	1	1	3

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Colorado												
All races, both sexes	791	1	-	-	-	-	2	-	1	2	4	2
Male	285	1	-	-	-	-	1	-	-	2	1	1
Female	506	-	-	-	-	-	1	-	1	-	3	1
White, both sexes	754	1	-	-	-	-	1	-	1	2	4	2
Male	271	1	-	-	-	-	-	-	-	2	1	1
Female	483	-	-	-	-	-	1	-	1	-	3	1
Black, both sexes	27	-	-	-	-	-	1	-	-	-	-	-
Male	11	-	-	-	-	-	1	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	778	-	-	-	-	-	1	2	-	-	4	7
Male	292	-	-	-	-	-	-	1	-	-	2	6
Female	486	-	-	-	-	-	1	1	-	-	2	1
White, both sexes	722	-	-	-	-	-	1	2	-	-	3	5
Male	268	-	-	-	-	-	-	1	-	-	2	5
Female	454	-	-	-	-	-	1	1	-	-	1	-
Black, both sexes	48	-	-	-	-	-	-	-	-	-	1	2
Male	19	-	-	-	-	-	-	-	-	-	-	1
Female	29	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Delaware												
All races, both sexes	190	-	-	-	-	-	-	-	-	-	2	4
Male	75	-	-	-	-	-	-	-	-	-	-	3
Female	115	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	154	-	-	-	-	-	-	-	-	-	2	3
Male	57	-	-	-	-	-	-	-	-	-	-	2
Female	97	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	34	-	-	-	-	-	-	-	-	-	-	1
Male	18	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	94	-	-	-	-	-	-	-	-	-	-	2
Male	37	-	-	-	-	-	-	-	-	-	-	2
Female	57	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	77	-	-	-	-	-	-	-	-	-	-	2
Male	25	-	-	-	-	-	-	-	-	-	-	2
Female	52	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Florida												
All races, both sexes	4,548	1	1	-	-	-	1	-	3	2	22	31
Male	1,888	-	-	-	-	-	-	-	1	2	11	24
Female	2,660	1	1	-	-	-	1	-	2	-	11	7
White, both sexes	3,902	-	1	-	-	-	1	-	2	1	14	19
Male	1,608	-	-	-	-	-	-	-	1	1	6	15
Female	2,294	-	1	-	-	-	1	-	1	-	8	4
Black, both sexes	605	1	-	-	-	-	-	-	1	1	8	10
Male	261	-	-	-	-	-	-	-	-	1	5	8
Female	344	1	-	-	-	-	-	-	1	-	3	2
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	35	-	-	-	-	-	-	-	-	-	-	2
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	20	-	-	-	-	-	-	-	-	-	-	1
Georgia												
All races, both sexes	2,235	-	-	-	-	1	2	-	1	7	16	24
Male	870	-	-	-	-	-	2	-	-	4	10	14
Female	1,365	-	-	-	-	1	-	-	1	3	6	10
White, both sexes	1,595	-	-	-	-	1	1	-	1	5	4	11
Male	595	-	-	-	-	-	1	-	-	2	4	6
Female	1,000	-	-	-	-	1	-	-	1	3	-	5
Black, both sexes	622	-	-	-	-	-	1	-	-	1	12	13
Male	270	-	-	-	-	-	1	-	-	1	6	8
Female	352	-	-	-	-	-	-	-	-	-	6	5
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Hawaii												
All races, both sexes	310	-	-	-	-	-	-	-	-	1	1	5
Male	137	-	-	-	-	-	-	-	-	-	1	1
Female	173	-	-	-	-	-	-	-	-	1	-	4
White, both sexes	81	-	-	-	-	-	-	-	-	-	1	1
Male	32	-	-	-	-	-	-	-	-	-	1	-
Female	49	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	228	-	-	-	-	-	-	-	-	1	-	4
Male	104	-	-	-	-	-	-	-	-	-	-	1
Female	124	-	-	-	-	-	-	-	-	1	-	3
Idaho												
All races, both sexes	346	1	-	-	-	-	1	-	1	-	-	2
Male	117	-	-	-	-	-	-	-	-	-	-	1
Female	229	1	-	-	-	-	1	-	1	-	-	1
White, both sexes	342	1	-	-	-	-	1	-	1	-	-	2
Male	115	-	-	-	-	-	-	-	-	-	-	1
Female	227	1	-	-	-	-	1	-	1	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Illinois												
All races, both sexes	3,247	-	-	-	-	-	-	2	4	3	10	30
Male	1,188	-	-	-	-	-	-	1	2	-	3	18
Female	2,059	-	-	-	-	-	-	1	2	3	7	12
White, both sexes	2,745	-	-	-	-	-	-	2	-	2	3	11
Male	976	-	-	-	-	-	-	1	-	-	2	7
Female	1,769	-	-	-	-	-	-	1	-	2	1	4
Black, both sexes	462	-	-	-	-	-	-	-	4	1	7	19
Male	194	-	-	-	-	-	-	-	2	-	1	11
Female	268	-	-	-	-	-	-	-	2	1	6	8
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	35	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1,776	-	-	-	-	-	2	1	3	-	7	16
Male	658	-	-	-	-	-	1	-	2	-	3	6
Female	1,118	-	-	-	-	-	1	1	1	-	4	10
White, both sexes	1,645	-	-	-	-	-	1	1	2	-	5	13
Male	590	-	-	-	-	-	1	-	1	-	2	4
Female	1,055	-	-	-	-	-	-	1	1	-	3	9
Black, both sexes	130	-	-	-	-	-	1	-	1	-	2	3
Male	68	-	-	-	-	-	-	-	1	-	1	2
Female	62	-	-	-	-	-	1	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Iowa												
All races, both sexes	982	-	-	-	-	-	-	-	-	-	2	3
Male	347	-	-	-	-	-	-	-	-	-	2	3
Female	635	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	960	-	-	-	-	-	-	-	-	-	2	2
Male	333	-	-	-	-	-	-	-	-	-	2	2
Female	627	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	20	-	-	-	-	-	-	-	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	844	-	-	-	-	-	1	1	-	-	2	4
Male	315	-	-	-	-	-	1	-	-	-	1	2
Female	529	-	-	-	-	-	-	1	-	-	1	2
White, both sexes	797	-	-	-	-	-	1	-	-	-	-	4
Male	289	-	-	-	-	-	1	-	-	-	-	2
Female	508	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	36	-	-	-	-	-	-	1	-	-	2	-
Male	23	-	-	-	-	-	-	-	-	-	1	-
Female	13	-	-	-	-	-	-	1	-	-	1	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Kentucky												
All races, both sexes	1,222	-	-	-	-	-	-	-	2	2	5	5
Male	404	-	-	-	-	-	-	-	-	1	1	3
Female	818	-	-	-	-	-	-	-	2	1	4	2
White, both sexes	1,123	-	-	-	-	-	-	-	2	2	5	4
Male	363	-	-	-	-	-	-	-	-	1	1	2
Female	760	-	-	-	-	-	-	-	2	1	4	2
Black, both sexes	96	-	-	-	-	-	-	-	-	-	-	1
Male	40	-	-	-	-	-	-	-	-	-	-	1
Female	56	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1,204	-	-	-	-	-	2	1	-	8	14	23
Male	474	-	-	-	-	-	2	-	-	4	7	13
Female	730	-	-	-	-	-	-	1	-	4	7	10
White, both sexes	793	-	-	-	-	-	1	1	-	5	6	7
Male	293	-	-	-	-	-	1	-	-	3	3	4
Female	500	-	-	-	-	-	-	1	-	2	3	3
Black, both sexes	407	-	-	-	-	-	1	-	-	3	8	16
Male	180	-	-	-	-	-	1	-	-	1	4	9
Female	227	-	-	-	-	-	-	-	-	2	4	7
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Maine												
All races, both sexes	362	-	-	-	-	-	-	-	-	1	1	1
Male	117	-	-	-	-	-	-	-	-	1	-	1
Female	245	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	361	-	-	-	-	-	-	-	-	1	1	1
Male	116	-	-	-	-	-	-	-	-	1	-	1
Female	245	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1,237	-	-	-	-	-	-	-	1	1	4	12
Male	484	-	-	-	-	-	-	-	-	1	2	6
Female	753	-	-	-	-	-	-	-	1	-	2	6
White, both sexes	946	-	-	-	-	-	-	-	1	-	1	6
Male	354	-	-	-	-	-	-	-	-	-	1	4
Female	592	-	-	-	-	-	-	-	1	-	-	2
Black, both sexes	274	-	-	-	-	-	-	-	-	1	3	6
Male	122	-	-	-	-	-	-	-	-	1	1	2
Female	152	-	-	-	-	-	-	-	-	-	2	4
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Massachusetts												
All races, both sexes	1,384	-	-	-	-	-	-	-	-	2	3	1
Male	469	-	-	-	-	-	-	-	-	1	2	1
Female	915	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	1,328	-	-	-	-	-	-	-	-	2	2	1
Male	452	-	-	-	-	-	-	-	-	1	1	1
Female	876	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	34	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	22	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	2,156	-	-	-	-	-	1	-	2	4	5	20
Male	820	-	-	-	-	-	1	-	1	4	3	11
Female	1,336	-	-	-	-	-	-	-	1	-	2	9
White, both sexes	1,895	-	-	-	-	-	1	-	2	-	2	13
Male	700	-	-	-	-	-	1	-	1	-	2	8
Female	1,195	-	-	-	-	-	-	-	1	-	-	5
Black, both sexes	240	-	-	-	-	-	-	-	-	4	3	6
Male	107	-	-	-	-	-	-	-	-	4	1	2
Female	133	-	-	-	-	-	-	-	-	-	2	4
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Minnesota												
All races, both sexes	1,126	-	1	-	-	-	-	-	-	-	4	7
Male	426	-	1	-	-	-	-	-	-	-	2	6
Female	700	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	1,093	-	1	-	-	-	-	-	-	-	2	6
Male	411	-	1	-	-	-	-	-	-	-	2	5
Female	682	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	2	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	2	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	999	-	-	-	-	-	-	1	3	2	7	17
Male	396	-	-	-	-	-	-	1	1	2	4	12
Female	603	-	-	-	-	-	-	-	2	-	3	5
White, both sexes	631	-	-	-	-	-	-	-	-	-	2	4
Male	212	-	-	-	-	-	-	-	-	-	1	3
Female	419	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	365	-	-	-	-	-	-	1	3	2	5	13
Male	182	-	-	-	-	-	-	1	1	2	3	9
Female	183	-	-	-	-	-	-	-	2	-	2	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Missouri												
All races, both sexes	1,717	-	-	-	-	-	-	1	1	3	10	10
Male	635	-	-	-	-	-	-	-	1	2	6	4
Female	1,082	-	-	-	-	-	-	1	-	1	4	6
White, both sexes	1,537	-	-	-	-	-	-	1	1	1	7	8
Male	567	-	-	-	-	-	-	-	1	-	5	3
Female	970	-	-	-	-	-	-	1	-	1	2	5
Black, both sexes	166	-	-	-	-	-	-	-	-	2	3	2
Male	63	-	-	-	-	-	-	-	-	2	1	1
Female	103	-	-	-	-	-	-	-	-	-	2	1
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	248	-	-	-	-	-	-	-	-	1	1	1
Male	89	-	-	-	-	-	-	-	-	-	-	-
Female	159	-	-	-	-	-	-	-	-	1	1	1
White, both sexes	240	-	-	-	-	-	-	-	-	1	1	1
Male	86	-	-	-	-	-	-	-	-	-	-	-
Female	154	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Nebraska												
All races, both sexes	508	-	-	-	-	-	-	-	-	-	-	3
Male	168	-	-	-	-	-	-	-	-	-	-	3
Female	340	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	491	-	-	-	-	-	-	-	-	-	-	3
Male	162	-	-	-	-	-	-	-	-	-	-	3
Female	329	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	421	-	-	-	-	-	-	-	-	-	2	4
Male	192	-	-	-	-	-	-	-	-	-	-	2
Female	229	-	-	-	-	-	-	-	-	-	2	2
White, both sexes	360	-	-	-	-	-	-	-	-	-	1	1
Male	162	-	-	-	-	-	-	-	-	-	-	1
Female	198	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	39	-	-	-	-	-	-	-	-	-	1	3
Male	22	-	-	-	-	-	-	-	-	-	-	1
Female	17	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
New Hampshire												
All races, both sexes	240	-	-	-	-	-	-	-	-	-	1	1
Male	82	-	-	-	-	-	-	-	-	-	1	1
Female	158	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	238	-	-	-	-	-	-	-	-	-	1	1
Male	82	-	-	-	-	-	-	-	-	-	1	1
Female	156	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1,928	-	-	-	-	-	-	-	1	2	7	16
Male	717	-	-	-	-	-	-	-	1	1	4	12
Female	1,211	-	-	-	-	-	-	-	-	1	3	4
White, both sexes	1,632	-	-	-	-	-	-	-	-	1	2	10
Male	592	-	-	-	-	-	-	-	-	-	2	7
Female	1,040	-	-	-	-	-	-	-	-	1	-	3
Black, both sexes	258	-	-	-	-	-	-	-	-	1	5	6
Male	113	-	-	-	-	-	-	-	-	1	2	5
Female	145	-	-	-	-	-	-	-	-	-	3	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	36	-	-	-	-	-	-	-	1	-	-	-
Male	12	-	-	-	-	-	-	-	1	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
New Mexico												
All races, both sexes	409	-	-	-	-	-	1	-	-	-	1	4
Male	162	-	-	-	-	-	-	-	-	-	1	2
Female	247	-	-	-	-	-	1	-	-	-	-	2
White, both sexes	380	-	-	-	-	-	1	-	-	-	1	3
Male	150	-	-	-	-	-	-	-	-	-	1	2
Female	230	-	-	-	-	-	1	-	-	-	-	1
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	3,134	-	-	-	-	-	-	1	3	6	12	29
Male	1,192	-	-	-	-	-	-	-	2	2	7	12
Female	1,942	-	-	-	-	-	-	1	1	4	5	17
White, both sexes	2,622	-	-	-	-	-	-	-	3	4	9	16
Male	969	-	-	-	-	-	-	-	2	2	5	7
Female	1,653	-	-	-	-	-	-	-	1	2	4	9
Black, both sexes	449	-	-	-	-	-	-	1	-	2	3	12
Male	191	-	-	-	-	-	-	-	-	-	2	4
Female	258	-	-	-	-	-	-	1	-	2	1	8
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	60	-	-	-	-	-	-	-	-	-	-	1
Male	30	-	-	-	-	-	-	-	-	-	-	1
Female	30	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
North Carolina												
All races, both sexes	2,447	-	-	-	-	-	3	2	2	6	11	15
Male	917	-	-	-	-	-	2	-	1	3	7	7
Female	1,530	-	-	-	-	-	1	2	1	3	4	8
White, both sexes	1,883	-	-	-	-	-	2	-	-	-	3	10
Male	673	-	-	-	-	-	1	-	-	-	2	5
Female	1,210	-	-	-	-	-	1	-	-	-	1	5
Black, both sexes	532	-	-	-	-	-	-	2	2	6	8	5
Male	227	-	-	-	-	-	-	-	1	3	5	2
Female	305	-	-	-	-	-	-	2	1	3	3	3
American Indian 1/, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	1	-	-	-	-	-
Male	8	-	-	-	-	-	1	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	182	-	-	-	-	-	-	-	-	1	-	1
Male	77	-	-	-	-	-	-	-	-	1	-	1
Female	105	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	180	-	-	-	-	-	-	-	-	-	-	1
Male	75	-	-	-	-	-	-	-	-	-	-	1
Female	105	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Ohio												
All races, both sexes	2,571	-	-	1	1	-	1	-	3	3	5	18
Male	910	-	-	1	-	-	1	-	1	1	4	10
Female	1,661	-	-	-	1	-	-	-	2	2	1	8
White, both sexes	2,280	-	-	-	1	-	1	-	2	3	3	15
Male	775	-	-	-	-	-	1	-	-	1	2	9
Female	1,505	-	-	-	1	-	-	-	2	2	1	6
Black, both sexes	280	-	-	1	-	-	-	-	1	-	2	3
Male	132	-	-	1	-	-	-	-	1	-	2	1
Female	148	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1,324	-	-	-	-	-	-	1	2	4	7	17
Male	501	-	-	-	-	-	-	1	-	2	4	8
Female	823	-	-	-	-	-	-	-	2	2	3	9
White, both sexes	1,182	-	-	-	-	-	-	1	2	4	5	10
Male	444	-	-	-	-	-	-	1	-	2	2	3
Female	738	-	-	-	-	-	-	-	2	2	3	7
Black, both sexes	86	-	-	-	-	-	-	-	-	-	1	5
Male	35	-	-	-	-	-	-	-	-	-	1	4
Female	51	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	50	-	-	-	-	-	-	-	-	-	1	2
Male	20	-	-	-	-	-	-	-	-	-	1	1
Female	30	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Oregon												
All races, both sexes	907	-	-	-	-	-	1	-	-	3	1	12
Male	359	-	-	-	-	-	-	-	-	-	1	8
Female	548	-	-	-	-	-	1	-	-	3	-	4
White, both sexes	873	-	-	-	-	-	1	-	-	3	1	11
Male	346	-	-	-	-	-	-	-	-	-	1	8
Female	527	-	-	-	-	-	1	-	-	3	-	3
Black, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
Pennsylvania												
All races, both sexes	3,855	-	-	-	-	-	-	-	2	2	6	21
Male	1,396	-	-	-	-	-	-	-	1	1	4	11
Female	2,459	-	-	-	-	-	-	-	1	1	2	10
White, both sexes	3,530	-	-	-	-	-	-	-	2	2	5	15
Male	1,268	-	-	-	-	-	-	-	1	1	3	9
Female	2,262	-	-	-	-	-	-	-	1	1	2	6
Black, both sexes	303	-	-	-	-	-	-	-	-	-	1	6
Male	122	-	-	-	-	-	-	-	-	-	1	2
Female	181	-	-	-	-	-	-	-	-	-	-	4
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Rhode Island												
All races, both sexes	238	-	-	-	-	-	-	-	-	1	-	1
Male	86	-	-	-	-	-	-	-	-	1	-	1
Female	152	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	225	-	-	-	-	-	-	-	-	-	-	1
Male	79	-	-	-	-	-	-	-	-	-	-	1
Female	146	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1,221	-	1	-	1	-	-	-	2	3	8	14
Male	472	-	-	-	-	-	-	-	1	2	4	8
Female	749	-	1	-	1	-	-	-	1	1	4	6
White, both sexes	839	-	-	-	1	-	-	-	2	1	1	4
Male	302	-	-	-	-	-	-	-	1	-	1	1
Female	537	-	-	-	1	-	-	-	1	1	-	3
Black, both sexes	377	-	1	-	-	-	-	-	-	2	7	10
Male	168	-	-	-	-	-	-	-	-	2	3	7
Female	209	-	1	-	-	-	-	-	-	-	4	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
South Dakota												
All races, both sexes	232	-	-	-	-	-	-	-	-	-	-	2
Male	83	-	-	-	-	-	-	-	-	-	-	1
Female	149	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	224	-	-	-	-	-	-	-	-	-	-	1
Male	78	-	-	-	-	-	-	-	-	-	-	1
Female	146	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	1,966	-	-	-	-	1	2	-	2	4	16	19
Male	694	-	-	-	-	-	-	-	-	2	9	11
Female	1,272	-	-	-	-	1	2	-	2	2	7	8
White, both sexes	1,671	-	-	-	-	1	1	-	1	3	10	13
Male	575	-	-	-	-	-	-	-	-	1	6	7
Female	1,096	-	-	-	-	1	1	-	1	2	4	6
Black, both sexes	283	-	-	-	-	-	1	-	1	1	6	6
Male	117	-	-	-	-	-	-	-	-	1	3	4
Female	166	-	-	-	-	-	1	-	1	-	3	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Texas												
All races, both sexes	5,106	-	1	-	-	1	-	3	2	17	27	42
Male	1,938	-	-	-	-	-	-	2	1	9	15	21
Female	3,168	-	1	-	-	1	-	1	1	8	12	21
White, both sexes	4,406	-	1	-	-	1	-	2	1	9	20	31
Male	1,640	-	-	-	-	-	-	1	1	4	13	14
Female	2,766	-	1	-	-	1	-	1	-	5	7	17
Black, both sexes	650	-	-	-	-	-	-	1	1	8	6	11
Male	276	-	-	-	-	-	-	1	-	5	2	7
Female	374	-	-	-	-	-	-	-	1	3	4	4
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	43	-	-	-	-	-	-	-	-	-	1	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	1	-
Utah												
All races, both sexes	402	-	-	-	-	-	-	-	1	-	-	-
Male	136	-	-	-	-	-	-	-	1	-	-	-
Female	266	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	389	-	-	-	-	-	-	-	1	-	-	-
Male	130	-	-	-	-	-	-	-	1	-	-	-
Female	259	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Vermont												
All races, both sexes	137	-	-	-	-	-	-	-	-	-	1	-
Male	44	-	-	-	-	-	-	-	-	-	1	-
Female	93	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	137	-	-	-	-	-	-	-	-	-	1	-
Male	44	-	-	-	-	-	-	-	-	-	1	-
Female	93	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1,808	-	-	-	-	1	2	-	4	2	7	12
Male	675	-	-	-	-	1	1	-	1	-	4	6
Female	1,133	-	-	-	-	-	1	-	3	2	3	6
White, both sexes	1,364	-	-	-	-	1	1	-	1	1	3	7
Male	489	-	-	-	-	1	-	-	-	-	2	4
Female	875	-	-	-	-	-	1	-	1	1	1	3
Black, both sexes	411	-	-	-	-	-	1	-	3	1	4	5
Male	175	-	-	-	-	-	1	-	1	-	2	2
Female	236	-	-	-	-	-	-	-	2	1	2	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	32	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Washington												
All races, both sexes	1,403	-	-	1	-	-	-	-	-	1	5	7
Male	521	-	-	1	-	-	-	-	-	1	4	5
Female	882	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	1,314	-	-	1	-	-	-	-	-	1	4	6
Male	480	-	-	1	-	-	-	-	-	1	3	5
Female	834	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	30	-	-	-	-	-	-	-	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	48	-	-	-	-	-	-	-	-	-	1	-
Male	24	-	-	-	-	-	-	-	-	-	1	-
Female	24	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	595	-	-	-	-	-	-	-	1	1	2	6
Male	231	-	-	-	-	-	-	-	-	-	1	2
Female	364	-	-	-	-	-	-	-	1	1	1	4
White, both sexes	572	-	-	-	-	-	-	-	1	1	2	6
Male	222	-	-	-	-	-	-	-	-	-	1	2
Female	350	-	-	-	-	-	-	-	1	1	1	4
Black, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Wisconsin												
All races, both sexes	1,419	1	-	-	-	-	-	-	1	2	2	13
Male	486	-	-	-	-	-	-	-	1	2	1	6
Female	933	1	-	-	-	-	-	-	-	-	1	7
White, both sexes	1,345	1	-	-	-	-	-	-	1	1	1	9
Male	456	-	-	-	-	-	-	-	1	1	-	5
Female	889	1	-	-	-	-	-	-	-	-	1	4
Black, both sexes	60	-	-	-	-	-	-	-	-	1	-	3
Male	22	-	-	-	-	-	-	-	-	1	-	1
Female	38	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	130	-	-	-	-	-	-	-	-	1	-	1
Male	46	-	-	-	-	-	-	-	-	1	-	1
Female	84	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	129	-	-	-	-	-	-	-	-	1	-	1
Male	45	-	-	-	-	-	-	-	-	1	-	1
Female	84	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
United States												
All races, both sexes	1,073	1,701	2,353	3,549	5,227	8,873	13,676	16,564	11,637	4,553	776	2
Male	648	1,034	1,373	1,936	2,687	4,088	5,299	5,333	2,833	857	98	-
Female	425	667	980	1,613	2,540	4,785	8,377	11,231	8,804	3,696	678	2
White, both sexes	713	1,102	1,676	2,634	4,053	7,413	11,992	14,962	10,632	4,178	684	1
Male	443	684	987	1,439	2,081	3,398	4,636	4,844	2,572	784	78	-
Female	270	418	689	1,195	1,972	4,015	7,356	10,118	8,060	3,394	606	1
Black, both sexes	325	543	607	802	1,001	1,211	1,336	1,261	789	300	79	1
Male	187	320	347	438	520	568	522	376	187	53	14	-
Female	138	223	260	364	481	643	814	885	602	247	65	1
American Indian 1/, both sexes	9	12	13	26	33	42	47	45	24	11	4	-
Male	5	8	11	13	17	24	20	12	6	2	1	-
Female	4	4	2	13	16	18	27	33	18	9	3	-
Asian or Pacific Islander 2/, both sexes	26	44	57	87	140	207	301	296	192	64	9	-
Male	13	22	28	46	69	98	121	101	68	18	5	-
Female	13	22	29	41	71	109	180	195	124	46	4	-
Alabama												
All races, both sexes	36	60	65	111	141	219	314	307	205	74	5	-
Male	25	38	37	64	69	102	116	97	47	7	1	-
Female	11	22	28	47	72	117	198	210	158	67	4	-
White, both sexes	16	28	45	70	86	166	241	245	162	55	5	-
Male	10	18	24	37	41	78	91	77	32	7	1	-
Female	6	10	21	33	45	88	150	168	130	48	4	-
Black, both sexes	20	32	20	41	54	53	71	62	42	18	-	-
Male	15	20	13	27	27	24	25	20	14	-	-	-
Female	5	12	7	14	27	29	46	42	28	18	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	1	1	-	-
Male	-	-	-	-	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Alaska												
All races, both sexes	2	4	1	5	12	15	14	19	9	2	2	-
Male	2	2	1	5	9	7	8	9	1	-	-	-
Female	-	2	-	-	3	8	6	10	8	2	2	-
White, both sexes	1	3	1	4	8	12	11	18	6	2	2	-
Male	1	2	1	4	6	5	6	9	-	-	-	-
Female	-	1	-	-	2	7	5	9	6	2	2	-
Black, both sexes	-	-	-	1	1	-	-	-	1	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	1	-	-	-	1	3	3	1	2	-	-	-
Male	1	-	-	-	-	2	2	-	1	-	-	-
Female	-	-	-	-	1	1	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	2	-	-	-	-	-	-	-
Male	-	-	-	-	2	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	18	21	32	73	97	165	254	314	184	75	8	-
Male	9	13	15	39	52	74	87	98	38	13	2	-
Female	9	8	17	34	45	91	167	216	146	62	6	-
White, both sexes	14	18	29	67	93	158	246	299	179	71	7	-
Male	7	11	14	36	50	70	85	96	36	10	2	-
Female	7	7	15	31	43	88	161	203	143	61	5	-
Black, both sexes	2	2	2	2	2	4	3	5	3	2	1	-
Male	-	1	1	1	1	2	1	-	1	2	-	-
Female	2	1	1	1	1	2	2	5	2	-	1	-
American Indian 1/, both sexes	1	-	-	2	1	2	2	4	2	1	-	-
Male	1	-	-	2	-	1	1	1	1	-	-	-
Female	-	-	-	-	1	1	1	3	1	1	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	2	1	1	3	6	-	1	-	-
Male	1	1	-	-	1	1	-	1	-	1	-	-
Female	-	-	1	2	-	-	3	5	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Arkansas												
All races, both sexes	22	38	34	68	74	123	203	239	150	57	11	-
Male	13	23	17	44	35	65	71	70	30	7	1	-
Female	9	15	17	24	39	58	132	169	120	50	10	-
White, both sexes	15	29	25	57	61	105	186	215	124	52	9	-
Male	8	19	13	34	30	54	67	66	25	7	1	-
Female	7	10	12	23	31	51	119	149	99	45	8	-
Black, both sexes	7	9	9	10	12	18	16	23	25	5	2	-
Male	5	4	4	9	5	11	4	4	5	-	-	-
Female	2	5	5	1	7	7	12	19	20	5	2	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	1	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
California												
All races, both sexes	116	159	210	325	499	814	1,332	1,556	1,126	423	93	1
Male	73	107	118	178	257	355	552	535	328	102	15	-
Female	43	52	92	147	242	459	780	1,021	798	321	78	1
White, both sexes	84	115	154	234	351	640	1,095	1,324	980	373	81	1
Male	60	79	85	125	183	281	458	466	283	92	11	-
Female	24	36	69	109	168	359	637	858	697	281	70	1
Black, both sexes	20	25	33	51	82	67	85	92	43	20	6	-
Male	8	15	21	31	42	28	34	22	10	4	-	-
Female	12	10	12	20	40	39	51	70	33	16	6	-
American Indian 1/, both sexes	-	2	-	4	6	7	8	8	1	3	-	-
Male	-	2	-	3	3	1	2	3	1	1	-	-
Female	-	-	-	1	3	6	6	5	-	2	-	-
Asian or Pacific Islander 2/, both sexes	12	17	23	36	60	100	144	132	102	27	6	-
Male	5	11	12	19	29	45	58	44	34	5	4	-
Female	7	6	11	17	31	55	86	88	68	22	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Colorado												
All races, both sexes	7	12	16	36	53	93	155	206	141	51	9	-
Male	4	9	12	22	25	38	56	62	38	11	2	-
Female	3	3	4	14	28	55	99	144	103	40	7	-
White, both sexes	6	10	14	35	48	90	147	198	138	48	9	-
Male	4	7	11	22	23	38	54	57	37	11	2	-
Female	2	3	3	13	25	52	93	141	101	37	7	-
Black, both sexes	1	2	1	-	4	3	5	5	3	2	-	-
Male	-	2	1	-	1	-	2	3	1	-	-	-
Female	1	-	-	-	3	3	3	2	2	2	-	-
American Indian 1/, both sexes	-	-	1	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	3	3	-	1	-	-
Male	-	-	-	-	-	-	-	2	-	-	-	-
Female	-	-	-	-	-	-	3	1	-	1	-	-
Connecticut												
All races, both sexes	8	13	15	28	44	98	136	214	143	56	9	-
Male	5	9	6	10	24	41	59	78	35	13	3	-
Female	3	4	9	18	20	57	77	136	108	43	6	-
White, both sexes	7	11	10	21	37	85	133	205	139	54	9	-
Male	4	8	4	8	19	35	58	74	35	12	3	-
Female	3	3	6	13	18	50	75	131	104	42	6	-
Black, both sexes	1	2	4	6	4	13	3	6	4	2	-	-
Male	1	1	1	2	3	6	1	2	-	1	-	-
Female	-	1	3	4	1	7	2	4	4	1	-	-
American Indian 1/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	2	-	-	3	-	-	-	-
Male	-	-	-	-	1	-	-	2	-	-	-	-
Female	-	-	-	1	1	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Delaware												
All races, both sexes	-	6	5	13	15	27	39	34	29	11	5	-
Male	-	5	5	10	13	8	14	8	7	2	-	-
Female	-	1	-	3	2	19	25	26	22	9	5	-
White, both sexes	-	1	4	9	12	21	31	27	28	11	5	-
Male	-	1	4	7	10	7	10	7	7	2	-	-
Female	-	-	-	2	2	14	21	20	21	9	5	-
Black, both sexes	-	5	1	4	3	5	7	7	1	-	-	-
Male	-	4	1	3	3	1	4	1	-	-	-	-
Female	-	1	-	1	-	4	3	6	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	2	5	4	11	3	15	20	16	12	2	2	-
Male	2	1	2	6	2	7	5	7	1	1	1	-
Female	-	4	2	5	1	8	15	9	11	1	1	-
White, both sexes	-	-	-	3	-	2	6	3	1	1	1	-
Male	-	-	-	2	-	2	4	3	-	-	1	-
Female	-	-	-	1	-	-	2	-	1	1	-	-
Black, both sexes	2	5	4	8	3	13	14	13	11	1	1	-
Male	2	1	2	4	2	5	1	4	1	1	-	-
Female	-	4	2	4	1	8	13	9	10	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Florida												
All races, both sexes	59	110	173	247	363	557	922	1,058	658	297	43	-
Male	42	57	105	126	204	275	387	395	177	75	7	-
Female	17	53	68	121	159	282	535	663	481	222	36	-
White, both sexes	38	71	128	179	274	469	822	972	601	269	41	-
Male	27	37	83	91	155	230	354	371	164	67	6	-
Female	11	34	45	88	119	239	468	601	437	202	35	-
Black, both sexes	20	38	45	64	80	83	92	79	54	27	2	-
Male	14	20	22	32	45	43	29	22	12	7	1	-
Female	6	18	23	32	35	40	63	57	42	20	1	-
American Indian 1/, both sexes	-	-	-	-	2	2	2	-	-	-	-	-
Male	-	-	-	-	1	1	2	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	4	7	3	6	7	3	1	-	-
Male	1	-	-	3	3	1	2	2	1	1	-	-
Female	-	1	-	1	4	2	4	5	2	-	-	-
Georgia												
All races, both sexes	61	77	133	156	172	278	400	456	319	111	21	-
Male	37	35	81	102	94	133	134	138	69	17	-	-
Female	24	42	52	54	78	145	266	318	250	94	21	-
White, both sexes	29	39	76	96	106	194	313	354	262	87	16	-
Male	21	15	50	64	60	88	110	109	54	11	-	-
Female	8	24	26	32	46	106	203	245	208	76	16	-
Black, both sexes	31	38	57	58	65	81	79	101	56	24	5	-
Male	16	20	31	38	34	44	21	29	15	6	-	-
Female	15	18	26	20	31	37	58	72	41	18	5	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	2	1	3	7	1	1	-	-	-
Male	-	-	-	-	-	1	3	-	-	-	-	-
Female	1	-	-	2	1	2	4	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Hawaii												
All races, both sexes	1	9	7	15	13	36	57	89	51	21	4	-
Male	1	4	4	8	8	20	29	34	19	7	1	-
Female	-	5	3	7	5	16	28	55	32	14	3	-
White, both sexes	-	1	2	4	4	8	13	28	11	7	1	-
Male	-	1	2	4	3	5	5	7	3	1	-	-
Female	-	-	-	-	1	3	8	21	8	6	1	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	8	5	11	9	28	44	60	40	14	3	-
Male	1	3	2	4	5	15	24	26	16	6	1	-
Female	-	5	3	7	4	13	20	34	24	8	2	-
Idaho												
All races, both sexes	3	4	6	12	26	44	57	89	74	23	3	-
Male	3	2	3	4	14	13	26	26	18	7	-	-
Female	-	2	3	8	12	31	31	63	56	16	3	-
White, both sexes	3	4	6	11	25	44	56	88	74	23	3	-
Male	3	2	3	3	13	13	26	26	18	7	-	-
Female	-	2	3	8	12	31	30	62	56	16	3	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Illinois												
All races, both sexes	53	79	102	147	227	375	617	768	569	229	32	-
Male	34	54	61	87	112	167	243	241	131	33	1	-
Female	19	25	41	60	115	208	374	527	438	196	31	-
White, both sexes	28	47	66	105	170	301	536	701	538	207	28	-
Male	20	32	40	63	87	134	213	221	124	31	1	-
Female	8	15	26	42	83	167	323	480	414	176	27	-
Black, both sexes	24	31	35	42	50	71	72	53	28	21	4	-
Male	13	21	21	24	22	33	24	15	5	2	-	-
Female	11	10	14	18	28	38	48	38	23	19	4	-
American Indian 1/, both sexes	-	1	-	-	2	-	1	1	-	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	5	3	8	13	3	1	-	-
Male	1	-	-	-	2	-	6	5	2	-	-	-
Female	-	-	1	-	3	3	2	8	1	1	-	-
Indiana												
All races, both sexes	23	45	46	77	129	212	374	398	310	113	20	-
Male	13	28	25	40	69	104	154	112	78	23	-	-
Female	10	17	21	37	60	108	220	286	232	90	20	-
White, both sexes	19	39	38	65	112	186	352	385	298	109	20	-
Male	10	25	19	31	57	91	145	105	76	23	-	-
Female	9	14	19	34	55	95	207	280	222	86	20	-
Black, both sexes	4	6	8	12	17	26	22	13	11	4	-	-
Male	3	3	6	9	12	13	9	7	2	-	-	-
Female	1	3	2	3	5	13	13	6	9	4	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Iowa												
All races, both sexes	6	12	16	26	66	106	174	249	210	94	18	-
Male	5	8	11	18	27	58	72	79	44	18	2	-
Female	1	4	5	8	39	48	102	170	166	76	16	-
White, both sexes	5	11	16	22	64	103	170	245	208	94	18	-
Male	4	8	11	14	26	55	70	78	43	18	2	-
Female	1	3	5	8	38	48	100	167	165	76	16	-
Black, both sexes	1	1	-	4	2	3	3	3	2	-	-	-
Male	1	-	-	4	1	3	1	1	1	-	-	-
Female	-	1	-	-	1	-	2	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Kansas												
All races, both sexes	10	8	25	30	49	85	169	198	178	71	13	-
Male	5	5	13	15	24	47	63	76	48	12	3	-
Female	5	3	12	15	25	38	106	122	130	59	10	-
White, both sexes	5	7	25	28	42	79	164	193	167	69	13	-
Male	3	4	13	13	17	42	59	74	46	12	3	-
Female	2	3	12	15	25	37	105	119	121	57	10	-
Black, both sexes	4	1	-	2	6	4	5	2	8	1	-	-
Male	2	1	-	2	6	4	4	2	1	-	-	-
Female	2	-	-	-	-	-	1	-	7	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	2	-	3	1	1	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	3	1	1	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	-	-	2	-	-	-
Male	-	-	-	-	1	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Kentucky												
All races, both sexes	15	35	47	66	94	181	220	279	182	77	12	-
Male	8	16	21	28	48	90	79	64	31	13	1	-
Female	7	19	26	38	46	91	141	215	151	64	11	-
White, both sexes	12	30	44	56	83	167	199	264	170	74	11	-
Male	5	12	21	25	42	82	69	60	29	13	1	-
Female	7	18	23	31	41	85	130	204	141	61	10	-
Black, both sexes	3	5	3	10	11	13	21	14	11	3	1	-
Male	3	4	-	3	6	7	10	4	2	-	-	-
Female	-	1	3	7	5	6	11	10	9	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Louisiana												
All races, both sexes	37	51	58	78	111	165	224	225	150	52	5	-
Male	22	26	29	39	59	72	89	70	32	9	1	-
Female	15	25	29	39	52	93	135	155	118	43	4	-
White, both sexes	22	17	19	36	63	115	171	174	110	44	2	-
Male	15	6	11	16	35	51	67	49	24	7	1	-
Female	7	11	8	20	28	64	104	125	86	37	1	-
Black, both sexes	15	34	38	42	47	50	53	49	40	8	3	-
Male	7	20	18	23	24	21	22	20	8	2	-	-
Female	8	14	20	19	23	29	31	29	32	6	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	1	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Maine												
All races, both sexes	3	4	2	13	16	34	75	99	79	27	7	-
Male	2	2	1	8	6	15	25	29	19	5	3	-
Female	1	2	1	5	10	19	50	70	60	22	4	-
White, both sexes	3	4	2	13	16	33	75	99	79	27	7	-
Male	2	2	1	8	6	14	25	29	19	5	3	-
Female	1	2	1	5	10	19	50	70	60	22	4	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	26	38	50	48	88	150	228	292	200	85	14	-
Male	13	22	27	27	53	62	97	104	47	20	3	-
Female	13	16	23	21	35	88	131	188	153	65	11	-
White, both sexes	16	17	22	25	45	115	174	261	178	74	11	-
Male	9	9	13	16	24	45	77	91	44	19	2	-
Female	7	8	9	9	21	70	97	170	134	55	9	-
Black, both sexes	10	19	26	21	41	33	52	28	21	10	3	-
Male	4	12	13	9	28	15	20	13	2	1	1	-
Female	6	7	13	12	13	18	32	15	19	9	2	-
American Indian 1/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	2	2	1	1	2	3	1	1	-	-
Male	-	1	1	2	-	1	-	-	1	-	-	-
Female	-	1	1	-	1	-	2	3	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Massachusetts												
All races, both sexes	13	18	24	34	79	153	260	356	296	113	32	-
Male	7	11	17	17	45	67	98	105	72	21	5	-
Female	6	7	7	17	34	86	162	251	224	92	27	-
White, both sexes	13	13	21	28	71	146	251	348	290	111	31	-
Male	7	10	15	16	42	65	95	104	71	20	4	-
Female	6	3	6	12	29	81	156	244	219	91	27	-
Black, both sexes	-	4	1	3	4	2	5	8	5	1	1	-
Male	-	1	1	1	2	-	3	1	1	-	1	-
Female	-	3	-	2	2	2	2	7	4	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	2	3	4	5	4	-	1	1	-	-
Male	-	-	1	-	1	2	-	-	-	1	-	-
Female	-	1	1	3	3	3	4	-	1	-	-	-
Michigan												
All races, both sexes	41	50	49	92	134	268	433	540	360	134	23	-
Male	21	29	31	55	59	139	171	171	91	32	1	-
Female	20	21	18	37	75	129	262	369	269	102	22	-
White, both sexes	25	27	35	65	103	240	388	503	343	128	20	-
Male	13	19	22	39	44	121	153	161	84	31	1	-
Female	12	8	13	26	59	119	235	342	259	97	19	-
Black, both sexes	16	22	12	25	30	23	42	32	16	6	3	-
Male	8	9	7	14	15	13	18	9	6	1	-	-
Female	8	13	5	11	15	10	24	23	10	5	3	-
American Indian 1/, both sexes	-	-	1	1	1	2	3	1	-	-	-	-
Male	-	-	1	1	-	2	-	-	-	-	-	-
Female	-	-	-	-	1	-	3	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	1	-	3	-	4	1	-	-	-
Male	-	1	1	1	-	3	-	1	1	-	-	-
Female	-	-	-	-	-	-	-	3	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Minnesota												
All races, both sexes	9	14	22	44	64	108	196	307	224	104	22	-
Male	6	12	17	26	27	64	87	101	56	19	2	-
Female	3	2	5	18	37	44	109	206	168	85	20	-
White, both sexes	6	12	20	42	59	106	189	302	223	103	22	-
Male	4	11	15	24	26	63	84	100	56	18	2	-
Female	2	1	5	18	33	43	105	202	167	85	20	-
Black, both sexes	2	1	1	1	3	2	3	1	-	-	-	-
Male	2	-	1	1	1	1	1	-	-	-	-	-
Female	-	1	-	-	2	1	2	1	-	-	-	-
American Indian 1/, both sexes	1	1	1	-	1	-	-	-	1	1	-	-
Male	-	1	1	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	1	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	4	4	-	-	-	-
Male	-	-	-	1	-	-	2	1	-	-	-	-
Female	-	-	-	-	1	-	2	3	-	-	-	-
Mississippi												
All races, both sexes	25	31	48	69	76	138	199	190	140	43	10	-
Male	15	24	31	31	42	65	70	58	29	9	2	-
Female	10	7	17	38	34	73	129	132	111	34	8	-
White, both sexes	8	9	17	41	40	97	137	145	96	30	5	-
Male	4	6	9	16	22	41	46	44	15	5	-	-
Female	4	3	8	25	18	56	91	101	81	25	5	-
Black, both sexes	17	22	31	28	36	40	61	44	44	13	5	-
Male	11	18	22	15	20	24	23	13	14	4	2	-
Female	6	4	9	13	16	16	38	31	30	9	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Missouri												
All races, both sexes	17	43	42	93	133	207	325	410	280	125	17	-
Male	9	21	21	52	73	100	121	146	59	20	-	-
Female	8	22	21	41	60	107	204	264	221	105	17	-
White, both sexes	12	33	35	78	113	175	291	391	259	118	14	-
Male	7	17	16	47	63	82	109	140	58	19	-	-
Female	5	16	19	31	50	93	182	251	201	99	14	-
Black, both sexes	4	10	5	14	19	28	32	17	20	7	3	-
Male	2	4	4	5	10	14	12	6	1	1	-	-
Female	2	6	1	9	9	14	20	11	19	6	3	-
American Indian 1/, both sexes	1	-	-	1	-	2	-	1	-	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	1	2	2	1	1	-	-	-
Male	-	-	1	-	-	2	-	-	-	-	-	-
Female	-	-	1	-	1	-	2	1	1	-	-	-
Montana												
All races, both sexes	4	4	14	7	15	27	46	54	49	24	1	-
Male	4	4	8	3	5	16	17	16	13	3	-	-
Female	-	-	6	4	10	11	29	38	36	21	1	-
White, both sexes	3	4	14	7	15	24	45	53	47	24	1	-
Male	3	4	8	3	5	14	17	16	13	3	-	-
Female	-	-	6	4	10	10	28	37	34	21	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	3	1	1	2	-	-	-
Male	1	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Nebraska												
All races, both sexes	4	3	7	15	34	65	94	116	100	52	15	-
Male	2	3	4	8	18	34	35	38	16	5	2	-
Female	2	-	3	7	16	31	59	78	84	47	13	-
White, both sexes	3	3	7	14	30	62	93	114	99	50	13	-
Male	1	3	4	8	16	32	35	38	16	5	1	-
Female	2	-	3	6	14	30	58	76	83	45	12	-
Black, both sexes	-	-	-	1	2	3	1	2	1	2	2	-
Male	-	-	-	-	1	2	-	-	-	-	1	-
Female	-	-	-	1	1	1	1	2	1	2	1	-
American Indian 1/, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	10	10	20	30	42	64	74	89	61	13	2	-
Male	7	6	12	17	26	29	39	28	22	3	1	-
Female	3	4	8	13	16	35	35	61	39	10	1	-
White, both sexes	10	10	14	22	33	53	65	81	56	12	2	-
Male	7	6	7	13	20	23	36	25	20	3	1	-
Female	3	4	7	9	13	30	29	56	36	9	1	-
Black, both sexes	-	-	5	6	6	6	4	4	3	1	-	-
Male	-	-	4	3	3	5	2	3	1	-	-	-
Female	-	-	1	3	3	1	2	1	2	1	-	-
American Indian 1/, both sexes	-	-	-	1	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	2	5	4	4	2	-	-	-
Male	-	-	1	1	2	1	1	-	1	-	-	-
Female	-	-	-	-	-	4	3	4	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
New Hampshire												
All races, both sexes	-	5	3	9	16	22	39	67	52	22	3	-
Male	-	5	2	4	7	10	16	17	14	5	-	-
Female	-	-	1	5	9	12	23	50	38	17	3	-
White, both sexes	-	5	3	8	16	22	39	67	51	22	3	-
Male	-	5	2	4	7	10	16	17	14	5	-	-
Female	-	-	1	4	9	12	23	50	37	17	3	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
New Jersey												
All races, both sexes	19	44	49	90	124	232	409	453	334	127	20	1
Male	10	25	29	54	64	115	151	147	81	18	5	-
Female	9	19	20	36	60	117	258	306	253	109	15	1
White, both sexes	11	29	32	59	97	194	338	413	306	121	19	-
Male	5	18	18	35	52	98	122	137	76	17	5	-
Female	6	11	14	24	45	96	216	276	230	104	14	-
Black, both sexes	8	13	15	27	23	35	59	34	25	5	1	1
Male	5	7	10	17	10	16	26	8	5	1	-	-
Female	3	6	5	10	13	19	33	26	20	4	1	1
American Indian 1/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	2	3	4	3	11	6	3	1	-	-
Male	-	-	1	2	2	1	3	2	-	-	-	-
Female	-	2	1	1	2	2	8	4	3	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
New Mexico												
All races, both sexes	4	12	10	25	42	55	88	89	52	25	1	-
Male	2	6	8	19	23	29	26	29	12	5	-	-
Female	2	6	2	6	19	26	62	60	40	20	1	-
White, both sexes	4	10	10	24	37	53	81	82	49	24	1	-
Male	2	5	8	18	20	28	23	27	11	5	-	-
Female	2	5	2	6	17	25	58	55	38	19	1	-
Black, both sexes	-	1	-	-	2	-	3	2	-	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	3	2	-	-	-	-
American Indian 1/, both sexes	-	1	-	1	3	2	3	5	3	1	-	-
Male	-	-	-	1	2	1	3	2	1	-	-	-
Female	-	1	-	-	1	1	-	3	2	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
New York												
All races, both sexes	38	71	93	137	245	381	592	746	499	238	43	-
Male	21	46	66	76	117	186	217	256	139	44	1	-
Female	17	25	27	61	128	195	375	490	360	194	42	-
White, both sexes	25	38	60	102	182	293	499	677	456	222	36	-
Male	14	26	44	59	86	150	178	232	126	37	1	-
Female	11	12	16	43	96	143	321	445	330	185	35	-
Black, both sexes	13	33	29	32	57	75	75	61	37	12	7	-
Male	7	20	18	15	26	29	30	23	11	6	-	-
Female	6	13	11	17	31	46	45	38	26	6	7	-
American Indian 1/, both sexes	-	-	-	1	1	1	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	4	2	5	12	18	8	6	4	-	-
Male	-	-	4	1	4	7	9	1	2	1	-	-
Female	-	-	-	1	1	5	9	7	4	3	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
North Carolina												
All races, both sexes	41	72	80	141	199	349	429	544	394	133	26	-
Male	26	45	48	86	108	154	156	156	91	23	4	-
Female	15	27	32	55	91	195	273	388	303	110	22	-
White, both sexes	18	32	46	94	141	265	354	442	338	117	21	-
Male	11	18	28	61	75	117	125	130	78	20	2	-
Female	7	14	18	33	66	148	229	312	260	97	19	-
Black, both sexes	22	37	32	44	57	78	70	97	53	15	4	-
Male	15	25	19	23	33	33	27	24	13	3	1	-
Female	7	12	13	21	24	45	43	73	40	12	3	-
American Indian 1/, both sexes	-	1	1	1	1	2	4	4	2	1	1	-
Male	-	1	1	1	-	1	3	1	-	-	1	-
Female	-	-	-	-	1	1	1	3	2	1	-	-
Asian or Pacific Islander 2/, both sexes	1	2	1	2	-	4	1	1	1	-	-	-
Male	-	1	-	1	-	3	1	1	-	-	-	-
Female	1	1	1	1	-	1	-	-	1	-	-	-
North Dakota												
All races, both sexes	-	2	3	6	10	23	34	50	31	20	1	-
Male	-	2	1	3	5	13	18	15	14	4	-	-
Female	-	-	2	3	5	10	16	35	17	16	1	-
White, both sexes	-	2	3	6	10	22	34	50	31	20	1	-
Male	-	2	1	3	5	12	18	15	14	4	-	-
Female	-	-	2	3	5	10	16	35	17	16	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Ohio												
All races, both sexes	31	58	79	118	184	309	502	631	446	152	29	-
Male	16	40	39	55	86	136	179	205	104	28	4	-
Female	15	18	40	63	98	173	323	426	342	124	25	-
White, both sexes	24	39	60	92	151	264	460	578	424	137	26	-
Male	14	28	26	44	73	111	153	189	97	25	2	-
Female	10	11	34	48	78	153	307	389	327	112	24	-
Black, both sexes	7	19	19	26	29	44	42	50	20	14	3	-
Male	2	12	13	11	13	24	26	15	7	2	2	-
Female	5	7	6	15	16	20	16	35	13	12	1	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	4	-	-	3	2	1	-	-
Male	-	-	-	-	-	-	-	1	-	1	-	-
Female	-	-	-	-	4	-	-	2	2	-	-	-
Oklahoma												
All races, both sexes	32	26	61	88	125	206	231	278	179	56	11	-
Male	15	13	43	46	65	85	78	89	41	10	1	-
Female	17	13	18	42	60	121	153	189	138	46	10	-
White, both sexes	28	18	50	70	116	179	216	256	167	53	7	-
Male	15	9	33	39	61	75	72	84	39	9	-	-
Female	13	9	17	31	55	104	144	172	128	44	7	-
Black, both sexes	3	5	6	11	5	20	9	12	5	2	2	-
Male	-	3	5	4	2	6	4	3	1	1	1	-
Female	3	2	1	7	3	14	5	9	4	1	1	-
American Indian 1/, both sexes	-	3	5	7	3	6	4	10	6	1	2	-
Male	-	1	5	3	2	4	1	2	-	-	-	-
Female	-	2	-	4	1	2	3	8	6	1	2	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	1	2	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	1	-	-	-	1	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Oregon												
All races, both sexes	13	17	31	45	70	125	174	211	140	53	11	-
Male	8	10	20	20	35	56	80	71	34	12	4	-
Female	5	7	11	25	35	69	94	140	106	41	7	-
White, both sexes	13	17	29	42	65	121	165	205	139	50	11	-
Male	8	10	19	18	32	53	77	70	34	12	4	-
Female	5	7	10	24	33	68	88	135	105	38	7	-
Black, both sexes	-	-	1	2	3	1	4	3	-	2	-	-
Male	-	-	1	2	2	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	4	3	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	2	-	2	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	3	3	3	1	1	-	-
Male	-	-	-	-	-	3	2	1	-	-	-	-
Female	-	-	1	1	-	-	1	2	1	1	-	-
Pennsylvania												
All races, both sexes	42	61	94	147	218	492	760	983	723	279	25	-
Male	25	41	49	74	120	229	293	328	166	52	2	-
Female	17	20	45	73	98	263	467	655	557	227	23	-
White, both sexes	32	44	74	122	186	440	720	910	690	266	22	-
Male	19	31	41	64	101	202	280	305	159	50	2	-
Female	13	13	33	58	85	238	440	605	531	216	20	-
Black, both sexes	8	16	20	24	30	48	39	67	31	11	2	-
Male	5	10	8	9	18	25	13	22	7	2	-	-
Female	3	6	12	15	12	23	26	45	24	9	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	1	-
Asian or Pacific Islander 2/, both sexes	2	1	-	1	2	4	1	6	1	2	-	-
Male	1	-	-	1	1	2	-	1	-	-	-	-
Female	1	1	-	-	1	2	1	5	1	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Rhode Island												
All races, both sexes	5	2	5	4	10	22	46	64	49	27	2	-
Male	2	2	5	2	7	9	19	19	14	5	-	-
Female	3	-	-	2	3	13	27	45	35	22	2	-
White, both sexes	5	2	4	4	7	20	44	62	49	25	2	-
Male	2	2	4	2	4	8	18	19	14	5	-	-
Female	3	-	-	2	3	12	26	43	35	20	2	-
Black, both sexes	-	-	1	-	2	2	2	-	-	1	-	-
Male	-	-	1	-	2	1	1	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	2	-	1	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	1	-	-
South Carolina												
All races, both sexes	26	44	76	83	109	162	230	253	150	51	8	-
Male	15	31	49	48	53	70	92	69	24	6	-	-
Female	11	13	27	35	56	92	138	184	126	45	8	-
White, both sexes	12	22	40	43	61	106	171	205	126	37	7	-
Male	7	17	25	25	30	46	68	55	20	6	-	-
Female	5	5	15	18	31	60	103	150	106	31	7	-
Black, both sexes	14	22	34	39	47	56	58	48	24	14	1	-
Male	8	14	23	23	22	24	24	14	4	-	-	-
Female	6	8	11	16	25	32	34	34	20	14	1	-
American Indian 1/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
South Dakota												
All races, both sexes	2	1	6	9	8	35	34	57	47	25	6	-
Male	1	1	4	6	6	16	12	17	14	5	-	-
Female	1	-	2	3	2	19	22	40	33	20	6	-
White, both sexes	2	-	6	8	7	31	34	57	47	25	6	-
Male	1	-	4	5	6	13	12	17	14	5	-	-
Female	1	-	2	3	1	18	22	40	33	20	6	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	1	4	-	-	-	-	-	-
Male	-	1	-	-	-	3	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	33	55	71	115	158	273	384	424	294	96	19	-
Male	24	33	37	59	63	118	146	115	65	8	4	-
Female	9	22	34	56	95	155	238	309	229	88	15	-
White, both sexes	25	33	50	85	129	233	335	383	262	90	17	-
Male	18	23	25	43	56	100	126	104	55	8	3	-
Female	7	10	25	42	73	133	209	279	207	82	14	-
Black, both sexes	8	19	20	30	27	39	47	39	32	5	2	-
Male	6	9	12	16	7	17	20	11	10	-	1	-
Female	2	10	8	14	20	22	27	28	22	5	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	3	1	-	2	1	1	2	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	2	1	-	2	-	1	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Texas												
All races, both sexes	89	152	224	293	431	649	986	1,132	734	272	51	-
Male	59	92	116	144	218	282	384	355	183	52	5	-
Female	30	60	108	149	213	367	602	777	551	220	46	-
White, both sexes	67	112	164	230	343	552	877	1,035	669	250	42	-
Male	44	66	89	113	173	243	336	327	164	49	3	-
Female	23	46	75	117	170	309	541	708	505	201	39	-
Black, both sexes	20	39	55	60	80	92	97	89	61	21	9	-
Male	14	25	25	29	41	38	42	26	16	3	2	-
Female	6	14	30	31	39	54	55	63	45	18	7	-
American Indian 1/, both sexes	1	-	1	1	1	-	2	1	-	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	1	-	-	1	1	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	4	2	7	5	10	7	4	1	-	-
Male	1	1	1	2	4	1	6	1	3	-	-	-
Female	-	-	3	-	3	4	4	6	1	1	-	-
Utah												
All races, both sexes	4	5	14	14	30	56	80	111	58	23	6	-
Male	3	3	7	8	14	17	27	39	13	4	-	-
Female	1	2	7	6	16	39	53	72	45	19	6	-
White, both sexes	3	5	13	12	28	53	80	107	58	23	6	-
Male	2	3	6	7	14	15	27	38	13	4	-	-
Female	1	2	7	5	14	38	53	69	45	19	6	-
Black, both sexes	-	-	-	1	-	1	-	2	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	2	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	2	2	-	2	-	-	-	-
Male	-	-	1	1	-	1	-	1	-	-	-	-
Female	-	-	-	-	2	1	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Vermont												
All races, both sexes	1	1	4	9	4	16	27	30	30	14	-	-
Male	1	1	-	5	3	8	9	8	5	3	-	-
Female	-	-	4	4	1	8	18	22	25	11	-	-
White, both sexes	1	1	4	9	4	16	27	30	30	14	-	-
Male	1	1	-	5	3	8	9	8	5	3	-	-
Female	-	-	4	4	1	8	18	22	25	11	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	26	45	72	106	145	217	316	419	301	118	15	-
Male	12	28	47	61	78	92	128	131	66	18	1	-
Female	14	17	25	45	67	125	188	288	235	100	14	-
White, both sexes	13	29	47	61	92	151	244	342	262	98	11	-
Male	7	18	33	34	49	65	96	105	60	14	1	-
Female	6	11	14	27	43	86	148	237	202	84	10	-
Black, both sexes	13	16	25	39	49	58	66	75	36	16	4	-
Male	5	10	14	24	26	26	32	24	5	3	-	-
Female	8	6	11	15	23	32	34	51	31	13	4	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	6	4	8	6	2	2	4	-	-
Male	-	-	-	3	3	1	-	2	-	1	-	-
Female	-	-	-	3	1	7	6	-	2	3	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Washington												
All races, both sexes	11	25	28	50	104	168	296	331	254	106	16	-
Male	4	13	18	27	53	72	121	104	66	28	4	-
Female	7	12	10	23	51	96	175	227	188	78	12	-
White, both sexes	5	21	25	44	96	157	279	314	242	103	16	-
Male	1	11	18	24	47	66	114	97	61	27	4	-
Female	4	10	7	20	49	91	165	217	181	76	12	-
Black, both sexes	3	2	1	2	2	5	3	7	3	1	-	-
Male	2	1	-	-	2	2	2	2	2	-	-	-
Female	1	1	1	2	-	3	1	5	1	1	-	-
American Indian 1/, both sexes	1	-	1	1	-	1	5	2	-	-	-	-
Male	-	-	-	-	-	-	3	1	-	-	-	-
Female	1	-	1	1	-	1	2	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	1	3	6	5	9	8	9	2	-	-
Male	1	1	-	3	4	4	2	4	3	1	-	-
Female	1	1	1	-	2	1	7	4	6	1	-	-
West Virginia												
All races, both sexes	13	17	25	34	52	82	104	143	79	32	4	-
Male	6	9	16	19	27	37	43	45	18	6	2	-
Female	7	8	9	15	25	45	61	98	61	26	2	-
White, both sexes	12	15	22	33	52	79	103	138	76	30	2	-
Male	5	8	14	19	27	37	42	43	17	6	1	-
Female	7	7	8	14	25	42	61	95	59	24	1	-
Black, both sexes	1	2	2	1	-	3	1	5	2	2	2	-
Male	1	1	1	-	-	-	1	2	-	-	1	-
Female	-	1	1	1	-	3	-	3	2	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Wisconsin												
All races, both sexes	12	22	43	53	71	155	280	347	279	117	21	-
Male	9	12	30	28	34	76	95	112	67	12	1	-
Female	3	10	13	25	37	79	185	235	212	105	20	-
White, both sexes	10	14	37	45	66	144	268	340	271	116	21	-
Male	8	9	27	25	33	69	91	111	63	12	1	-
Female	2	5	10	20	33	75	177	229	208	104	20	-
Black, both sexes	1	5	6	6	4	10	10	6	7	1	-	-
Male	-	1	3	3	-	7	3	-	3	-	-	-
Female	1	4	3	3	4	3	7	6	4	1	-	-
American Indian 1/, both sexes	-	2	-	1	-	-	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	1	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	1	1	1	2	-	1	-	-	-
Male	1	1	-	-	1	-	1	-	1	-	-	-
Female	-	-	-	1	-	1	1	-	-	-	-	-
Wyoming												
All races, both sexes	-	1	9	4	3	22	24	34	23	8	-	-
Male	-	-	4	3	2	11	5	11	6	2	-	-
Female	-	1	5	1	1	11	19	23	17	6	-	-
White, both sexes	-	1	8	4	3	22	24	34	23	8	-	-
Male	-	-	3	3	2	11	5	11	6	2	-	-
Female	-	1	5	1	1	11	19	23	17	6	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
United States												
All races, both sexes	24,956	62	33	11	9	6	16	18	22	54	86	176
Male	9,645	37	13	4	6	3	11	8	12	33	46	93
Female	15,311	25	20	7	3	3	5	10	10	21	40	83
White, both sexes	21,335	45	19	10	7	4	13	13	12	38	61	116
Male	8,121	25	10	4	5	2	8	8	6	27	32	61
Female	13,214	20	9	6	2	2	5	5	6	11	29	55
Black, both sexes	2,883	15	13	-	2	1	2	4	8	12	20	48
Male	1,195	11	3	-	1	1	2	-	4	5	12	26
Female	1,688	4	10	-	1	-	-	4	4	7	8	22
American Indian 1/, both sexes	109	-	1	-	-	1	-	-	-	1	2	-
Male	53	-	-	-	-	-	-	-	-	1	1	-
Female	56	-	1	-	-	1	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	629	2	-	1	-	-	1	1	2	3	3	12
Male	276	1	-	-	-	-	1	-	2	-	1	6
Female	353	1	-	1	-	-	-	1	-	3	2	6
Alabama												
All races, both sexes	336	3	-	1	-	1	-	-	-	-	2	4
Male	147	2	-	-	-	1	-	-	-	-	1	3
Female	189	1	-	1	-	-	-	-	-	-	1	1
White, both sexes	246	2	-	1	-	1	-	-	-	-	1	1
Male	103	2	-	-	-	1	-	-	-	-	1	1
Female	143	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	89	1	-	-	-	-	-	-	-	-	-	3
Male	44	-	-	-	-	-	-	-	-	-	-	2
Female	45	1	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Alaska												
All races, both sexes	30	-	-	-	-	-	-	-	-	-	1	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	21	-	-	-	-	-	-	-	-	-	1	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	310	1	-	1	-	-	-	-	-	-	3	4
Male	111	-	-	1	-	-	-	-	-	-	3	2
Female	199	1	-	-	-	-	-	-	-	-	-	2
White, both sexes	297	1	-	1	-	-	-	-	-	-	2	4
Male	106	-	-	1	-	-	-	-	-	-	2	2
Female	191	1	-	-	-	-	-	-	-	-	-	2
Black, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Arkansas												
All races, both sexes	354	-	-	-	-	-	-	-	-	-	1	1
Male	151	-	-	-	-	-	-	-	-	-	-	-
Female	203	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	298	-	-	-	-	-	-	-	-	-	1	1
Male	130	-	-	-	-	-	-	-	-	-	-	-
Female	168	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	53	-	-	-	-	-	-	-	-	-	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	34	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	3,706	10	3	1	-	3	5	3	3	16	15	38
Male	1,454	4	1	-	-	1	5	-	1	10	9	21
Female	2,252	6	2	1	-	2	-	3	2	6	6	17
White, both sexes	3,003	8	1	1	-	1	3	-	1	12	14	26
Male	1,159	2	1	-	-	-	3	-	-	9	8	12
Female	1,844	6	-	1	-	1	-	-	1	3	6	14
Black, both sexes	309	1	2	-	-	1	1	2	1	1	-	5
Male	115	1	-	-	-	1	1	-	-	-	-	4
Female	194	-	2	-	-	-	-	2	1	1	-	1
American Indian 1/, both sexes	13	-	-	-	-	1	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	8	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	381	1	-	-	-	-	1	1	1	2	1	7
Male	175	1	-	-	-	-	1	-	1	-	1	5
Female	206	-	-	-	-	-	-	1	-	2	-	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Colorado												
All races, both sexes	316	-	-	-	-	-	-	1	-	-	-	-
Male	113	-	-	-	-	-	-	-	-	-	-	-
Female	203	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	301	-	-	-	-	-	-	1	-	-	-	-
Male	108	-	-	-	-	-	-	-	-	-	-	-
Female	193	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	236	2	-	-	-	-	-	-	-	-	-	1
Male	74	1	-	-	-	-	-	-	-	-	-	1
Female	162	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	222	2	-	-	-	-	-	-	-	-	-	1
Male	70	1	-	-	-	-	-	-	-	-	-	1
Female	152	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Delaware												
All races, both sexes	73	-	-	-	1	-	-	-	-	-	-	-
Male	28	-	-	-	1	-	-	-	-	-	-	-
Female	45	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	52	-	-	-	1	-	-	-	-	-	-	-
Male	16	-	-	-	1	-	-	-	-	-	-	-
Female	36	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	32	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Florida												
All races, both sexes	1,580	1	1	2	-	-	1	-	1	3	4	7
Male	625	-	1	1	-	-	1	-	1	-	3	3
Female	955	1	-	1	-	-	-	-	-	3	1	4
White, both sexes	1,379	-	1	2	-	-	-	-	-	1	4	5
Male	547	-	1	1	-	-	-	-	-	-	3	2
Female	832	-	-	1	-	-	-	-	-	1	1	3
Black, both sexes	194	1	-	-	-	-	1	-	1	2	-	2
Male	75	-	-	-	-	-	1	-	1	-	-	1
Female	119	1	-	-	-	-	-	-	-	2	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	520	2	1	-	1	-	-	-	1	3	4	8
Male	208	2	-	-	1	-	-	-	1	1	3	4
Female	312	-	1	-	-	-	-	-	-	2	1	4
White, both sexes	364	-	-	-	-	-	-	-	-	2	1	6
Male	135	-	-	-	-	-	-	-	-	-	1	3
Female	229	-	-	-	-	-	-	-	-	2	-	3
Black, both sexes	153	2	1	-	1	-	-	-	1	1	3	2
Male	72	2	-	-	1	-	-	-	1	1	2	1
Female	81	-	1	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Hawaii												
All races, both sexes	86	-	1	-	-	-	-	-	-	-	-	1
Male	41	-	-	-	-	-	-	-	-	-	-	-
Female	45	-	1	-	-	-	-	-	-	-	-	1
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	1	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	63	-	-	-	-	-	-	-	-	-	-	1
Male	29	-	-	-	-	-	-	-	-	-	-	-
Female	34	-	-	-	-	-	-	-	-	-	-	1
Idaho												
All races, both sexes	150	-	-	-	-	-	-	-	-	1	1	1
Male	55	-	-	-	-	-	-	-	-	-	-	1
Female	95	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	146	-	-	-	-	-	-	-	-	1	1	1
Male	55	-	-	-	-	-	-	-	-	-	-	1
Female	91	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Illinois												
All races, both sexes	1,021	1	-	1	-	-	-	-	-	1	2	8
Male	383	1	-	-	-	-	-	-	-	1	1	2
Female	638	-	-	1	-	-	-	-	-	-	1	6
White, both sexes	840	1	-	1	-	-	-	-	-	-	2	3
Male	307	1	-	-	-	-	-	-	-	-	1	1
Female	533	-	-	1	-	-	-	-	-	-	1	2
Black, both sexes	166	-	-	-	-	-	-	-	-	1	-	5
Male	70	-	-	-	-	-	-	-	-	1	-	1
Female	96	-	-	-	-	-	-	-	-	-	-	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	473	1	1	-	1	-	-	1	2	3	1	4
Male	183	-	-	-	1	-	-	-	1	2	-	3
Female	290	1	1	-	-	-	-	1	1	1	1	1
White, both sexes	433	1	1	-	1	-	-	1	1	3	1	2
Male	168	-	-	-	1	-	-	-	1	2	-	2
Female	265	1	1	-	-	-	-	1	-	1	1	-
Black, both sexes	39	-	-	-	-	-	-	-	1	-	-	2
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	25	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Iowa												
All races, both sexes	255	-	-	-	-	-	-	-	-	1	1	2
Male	99	-	-	-	-	-	-	-	-	-	-	1
Female	156	-	-	-	-	-	-	-	-	1	1	1
White, both sexes	248	-	-	-	-	-	-	-	-	1	1	1
Male	96	-	-	-	-	-	-	-	-	-	-	1
Female	152	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Kansas												
All races, both sexes	279	-	1	-	1	-	-	-	-	1	1	1
Male	106	-	1	-	1	-	-	-	-	1	1	-
Female	173	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	259	-	1	-	1	-	-	-	-	1	1	-
Male	97	-	1	-	1	-	-	-	-	1	1	-
Female	162	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Kentucky												
All races, both sexes	271	1	1	-	-	-	-	-	-	-	2	3
Male	100	1	-	-	-	-	-	-	-	-	2	1
Female	171	-	1	-	-	-	-	-	-	-	-	2
White, both sexes	253	-	-	-	-	-	-	-	-	-	2	3
Male	91	-	-	-	-	-	-	-	-	-	2	1
Female	162	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	17	1	1	-	-	-	-	-	-	-	-	-
Male	8	1	-	-	-	-	-	-	-	-	-	-
Female	9	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	302	2	1	-	-	-	-	1	1	4	1	2
Male	119	2	1	-	-	-	-	1	1	3	-	1
Female	183	-	-	-	-	-	-	-	-	1	1	1
White, both sexes	228	-	1	-	-	-	-	1	-	2	-	-
Male	88	-	1	-	-	-	-	1	-	1	-	-
Female	140	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	74	2	-	-	-	-	-	-	1	2	1	2
Male	31	2	-	-	-	-	-	-	1	2	-	1
Female	43	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Maine												
All races, both sexes	107	-	1	-	-	-	-	-	-	-	1	3
Male	40	-	1	-	-	-	-	-	-	-	1	1
Female	67	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	106	-	-	-	-	-	-	-	-	-	1	3
Male	39	-	-	-	-	-	-	-	-	-	1	1
Female	67	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	374	1	1	-	-	-	1	-	-	-	1	2
Male	129	1	1	-	-	-	1	-	-	-	-	2
Female	245	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	278	1	1	-	-	-	1	-	-	-	-	1
Male	100	1	1	-	-	-	1	-	-	-	-	1
Female	178	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	89	-	-	-	-	-	-	-	-	-	1	1
Male	27	-	-	-	-	-	-	-	-	-	-	1
Female	62	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Massachusetts												
All races, both sexes	611	1	2	-	-	-	-	-	-	-	-	3
Male	230	1	1	-	-	-	-	-	-	-	-	1
Female	381	-	1	-	-	-	-	-	-	-	-	2
White, both sexes	580	1	1	-	-	-	-	-	-	-	-	2
Male	217	1	-	-	-	-	-	-	-	-	-	-
Female	363	-	1	-	-	-	-	-	-	-	-	2
Black, both sexes	22	-	1	-	-	-	-	-	-	-	-	-
Male	8	-	1	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1,149	2	1	-	-	-	1	-	2	1	7	9
Male	448	1	-	-	-	-	1	-	1	-	5	7
Female	701	1	1	-	-	-	-	-	1	1	2	2
White, both sexes	989	2	1	-	-	-	1	-	2	-	2	7
Male	382	1	-	-	-	-	1	-	1	-	1	6
Female	607	1	1	-	-	-	-	-	1	-	1	1
Black, both sexes	150	-	-	-	-	-	-	-	-	-	4	2
Male	63	-	-	-	-	-	-	-	-	-	4	1
Female	87	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	1	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Minnesota												
All races, both sexes	404	1	1	-	-	-	-	-	-	1	1	3
Male	156	-	-	-	-	-	-	-	-	1	1	1
Female	248	1	1	-	-	-	-	-	-	-	-	2
White, both sexes	389	1	-	-	-	-	-	-	-	1	1	3
Male	148	-	-	-	-	-	-	-	-	1	1	1
Female	241	1	-	-	-	-	-	-	-	-	-	2
Black, both sexes	10	-	1	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	170	3	1	1	-	-	-	-	2	-	2	4
Male	69	1	-	1	-	-	-	-	2	-	2	2
Female	101	2	1	-	-	-	-	-	-	-	-	2
White, both sexes	106	2	-	1	-	-	-	-	1	-	1	1
Male	35	1	-	1	-	-	-	-	1	-	1	-
Female	71	1	-	-	-	-	-	-	-	-	-	1
Black, both sexes	64	1	1	-	-	-	-	-	1	-	1	3
Male	34	-	-	-	-	-	-	-	1	-	1	2
Female	30	1	1	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Missouri												
All races, both sexes	566	2	1	3	-	-	-	1	-	-	-	2
Male	197	2	-	-	-	-	-	1	-	-	-	-
Female	369	-	1	3	-	-	-	-	-	-	-	2
White, both sexes	518	1	1	2	-	-	-	1	-	-	-	2
Male	177	1	-	-	-	-	-	1	-	-	-	-
Female	341	-	1	2	-	-	-	-	-	-	-	2
Black, both sexes	44	1	-	-	-	-	-	-	-	-	-	-
Male	20	1	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	1	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	86	-	-	-	-	-	-	-	-	-	-	1
Male	40	-	-	-	-	-	-	-	-	-	-	1
Female	46	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	82	-	-	-	-	-	-	-	-	-	-	1
Male	39	-	-	-	-	-	-	-	-	-	-	1
Female	43	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Nebraska												
All races, both sexes	178	-	2	-	-	-	1	-	-	-	1	-
Male	74	-	-	-	-	-	1	-	-	-	1	-
Female	104	-	2	-	-	-	-	-	-	-	-	-
White, both sexes	166	-	1	-	-	-	1	-	-	-	1	-
Male	69	-	-	-	-	-	1	-	-	-	1	-
Female	97	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	109	1	-	-	-	-	-	-	-	-	1	3
Male	33	1	-	-	-	-	-	-	-	-	-	1
Female	76	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	97	1	-	-	-	-	-	-	-	-	-	3
Male	29	1	-	-	-	-	-	-	-	-	-	1
Female	68	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
New Hampshire												
All races, both sexes	112	2	1	-	-	-	-	-	-	1	1	1
Male	38	1	1	-	-	-	-	-	-	1	1	1
Female	74	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	112	2	1	-	-	-	-	-	-	1	1	1
Male	38	1	1	-	-	-	-	-	-	1	1	1
Female	74	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	434	1	1	-	1	-	-	-	-	1	2	1
Male	159	1	-	-	1	-	-	-	-	1	-	1
Female	275	-	1	-	-	-	-	-	-	-	2	-
White, both sexes	377	1	-	-	1	-	-	-	-	1	1	-
Male	137	1	-	-	1	-	-	-	-	1	-	-
Female	240	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	48	-	1	-	-	-	-	-	-	-	1	1
Male	19	-	-	-	-	-	-	-	-	-	-	1
Female	29	-	1	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
New Mexico												
All races, both sexes	182	-	-	-	-	-	1	-	1	-	3	2
Male	74	-	-	-	-	-	-	-	1	-	-	-
Female	108	-	-	-	-	-	1	-	-	-	3	2
White, both sexes	166	-	-	-	-	-	1	-	1	-	3	2
Male	65	-	-	-	-	-	-	-	1	-	-	-
Female	101	-	-	-	-	-	1	-	-	-	3	2
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	842	2	2	-	-	-	-	2	2	2	3	8
Male	301	1	1	-	-	-	-	2	-	2	1	5
Female	541	1	1	-	-	-	-	-	2	-	2	3
White, both sexes	743	-	2	-	-	-	-	2	1	2	3	6
Male	265	-	1	-	-	-	-	2	-	2	1	4
Female	478	-	1	-	-	-	-	-	1	-	2	2
Black, both sexes	85	1	-	-	-	-	-	-	1	-	-	1
Male	31	1	-	-	-	-	-	-	-	-	-	1
Female	54	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	1	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	1	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
North Carolina												
All races, both sexes	750	1	-	-	-	-	1	2	-	1	2	7
Male	295	-	-	-	-	-	-	-	-	-	-	5
Female	455	1	-	-	-	-	1	2	-	1	2	2
White, both sexes	565	-	-	-	-	-	1	-	-	-	2	1
Male	224	-	-	-	-	-	-	-	-	-	-	-
Female	341	-	-	-	-	-	1	-	-	-	2	1
Black, both sexes	176	1	-	-	-	-	-	2	-	1	-	6
Male	68	-	-	-	-	-	-	-	-	-	-	5
Female	108	1	-	-	-	-	-	2	-	1	-	1
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	53	-	1	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	36	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	53	-	1	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	36	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Ohio												
All races, both sexes	1,505	1	1	-	1	-	1	-	1	3	5	6
Male	605	1	-	-	-	-	-	-	1	3	4	5
Female	900	-	1	-	1	-	1	-	-	-	1	1
White, both sexes	1,308	-	-	-	1	-	1	-	1	3	4	5
Male	522	-	-	-	-	-	-	-	1	3	3	5
Female	786	-	-	-	1	-	1	-	-	-	1	-
Black, both sexes	188	1	1	-	-	-	-	-	-	-	1	1
Male	80	1	-	-	-	-	-	-	-	-	1	-
Female	108	-	1	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	329	2	-	-	-	-	-	1	-	-	-	2
Male	113	2	-	-	-	-	-	-	-	-	-	2
Female	216	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	296	1	-	-	-	-	-	1	-	-	-	1
Male	101	1	-	-	-	-	-	-	-	-	-	1
Female	195	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	16	1	-	-	-	-	-	-	-	-	-	1
Male	6	1	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Oregon												
All races, both sexes	424	1	-	-	-	-	-	-	1	-	1	-
Male	186	-	-	-	-	-	-	-	1	-	-	-
Female	238	1	-	-	-	-	-	-	-	-	1	-
White, both sexes	401	1	-	-	-	-	-	-	1	-	1	-
Male	173	-	-	-	-	-	-	-	1	-	-	-
Female	228	1	-	-	-	-	-	-	-	-	1	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1,094	2	2	-	-	1	1	1	-	2	2	5
Male	419	1	1	-	-	1	-	-	-	1	-	3
Female	675	1	1	-	-	-	1	1	-	1	2	2
White, both sexes	989	2	2	-	-	1	1	1	-	1	1	4
Male	367	1	1	-	-	1	-	-	-	-	-	2
Female	622	1	1	-	-	-	1	1	-	1	1	2
Black, both sexes	96	-	-	-	-	-	-	-	-	1	1	1
Male	45	-	-	-	-	-	-	-	-	1	-	1
Female	51	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Rhode Island												
All races, both sexes	83	-	-	-	-	-	-	-	-	-	-	1
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	62	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	82	-	-	-	-	-	-	-	-	-	-	1
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	61	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	395	-	1	-	-	-	-	1	-	1	2	-
Male	149	-	1	-	-	-	-	-	-	1	1	-
Female	246	-	-	-	-	-	-	1	-	-	1	-
White, both sexes	256	-	1	-	-	-	-	1	-	1	-	-
Male	86	-	1	-	-	-	-	-	-	1	-	-
Female	170	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	136	-	-	-	-	-	-	-	-	-	2	-
Male	62	-	-	-	-	-	-	-	-	-	1	-
Female	74	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
South Dakota												
All races, both sexes	73	-	-	-	-	1	-	-	1	-	-	1
Male	33	-	-	-	-	-	-	-	-	-	-	1
Female	40	-	-	-	-	1	-	-	1	-	-	-
White, both sexes	71	-	-	-	-	1	-	-	1	-	-	1
Male	31	-	-	-	-	-	-	-	-	-	-	1
Female	40	-	-	-	-	1	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	508	-	2	-	-	-	-	-	1	2	2	2
Male	179	-	1	-	-	-	-	-	-	-	-	1
Female	329	-	1	-	-	-	-	-	1	2	2	1
White, both sexes	430	-	1	-	-	-	-	-	1	-	-	2
Male	149	-	1	-	-	-	-	-	-	-	-	1
Female	281	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	78	-	1	-	-	-	-	-	-	2	2	-
Male	30	-	-	-	-	-	-	-	-	-	-	-
Female	48	-	1	-	-	-	-	-	-	2	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Texas												
All races, both sexes	1,958	8	2	1	1	-	2	2	1	5	6	13
Male	756	5	1	1	-	-	2	2	-	4	4	4
Female	1,202	3	1	-	1	-	-	-	1	1	2	9
White, both sexes	1,640	8	1	1	1	-	2	2	-	4	4	8
Male	636	5	-	1	-	-	2	2	-	4	2	3
Female	1,004	3	1	-	1	-	-	-	-	-	2	5
Black, both sexes	286	-	1	-	-	-	-	-	1	1	2	5
Male	105	-	1	-	-	-	-	-	-	-	2	1
Female	181	-	-	-	-	-	-	-	1	1	-	4
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	163	2	-	-	-	-	-	-	-	-	2	-
Male	58	-	-	-	-	-	-	-	-	-	1	-
Female	105	2	-	-	-	-	-	-	-	-	1	-
White, both sexes	158	2	-	-	-	-	-	-	-	-	1	-
Male	55	-	-	-	-	-	-	-	-	-	-	-
Female	103	2	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Vermont												
All races, both sexes	83	-	-	-	-	-	-	-	1	-	-	-
Male	32	-	-	-	-	-	-	-	-	-	-	-
Female	51	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	83	-	-	-	-	-	-	-	1	-	-	-
Male	32	-	-	-	-	-	-	-	-	-	-	-
Female	51	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	545	2	-	-	1	-	-	-	1	-	1	5
Male	219	2	-	-	-	-	-	-	1	-	-	2
Female	326	-	-	-	1	-	-	-	-	-	1	3
White, both sexes	419	1	-	-	-	-	-	-	-	-	1	2
Male	156	1	-	-	-	-	-	-	-	-	-	1
Female	263	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	119	1	-	-	1	-	-	-	-	-	-	2
Male	58	1	-	-	-	-	-	-	-	-	-	1
Female	61	-	-	-	1	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	1	-	-	1
Male	5	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Washington												
All races, both sexes	553	3	-	-	-	-	1	2	-	1	-	2
Male	244	3	-	-	-	-	-	2	-	1	-	2
Female	309	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	514	3	-	-	-	-	1	2	-	1	-	2
Male	227	3	-	-	-	-	-	2	-	1	-	2
Female	287	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	185	-	-	-	-	-	-	-	-	-	-	1
Male	57	-	-	-	-	-	-	-	-	-	-	1
Female	128	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	171	-	-	-	-	-	-	-	-	-	-	1
Male	53	-	-	-	-	-	-	-	-	-	-	1
Female	118	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Wisconsin												
All races, both sexes	580	-	-	-	1	-	-	-	-	-	1	4
Male	241	-	-	-	1	-	-	-	-	-	1	1
Female	339	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	553	-	-	-	1	-	-	-	-	-	1	2
Male	228	-	-	-	1	-	-	-	-	-	1	1
Female	325	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	21	-	-	-	-	-	-	-	-	-	-	2
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
United States												
All races, both sexes	281	456	718	1,028	1,678	2,985	4,736	5,970	4,381	1,871	359	-
Male	157	271	404	576	912	1,424	1,923	2,105	1,202	362	43	-
Female	124	185	314	452	766	1,561	2,813	3,865	3,179	1,509	316	-
White, both sexes	177	325	515	758	1,298	2,506	4,093	5,332	3,975	1,707	311	-
Male	103	196	296	427	704	1,205	1,659	1,889	1,096	325	33	-
Female	74	129	219	331	594	1,301	2,434	3,443	2,879	1,382	278	-
Black, both sexes	96	107	178	228	313	378	509	468	309	126	46	-
Male	50	62	95	130	171	170	209	140	67	27	9	-
Female	46	45	83	98	142	208	300	328	242	99	37	-
American Indian 1/, both sexes	1	5	6	6	14	16	16	21	16	3	-	-
Male	1	3	2	4	9	10	8	10	4	-	-	-
Female	-	2	4	2	5	6	8	11	12	3	-	-
Asian or Pacific Islander 2/, both sexes	7	19	19	36	53	85	118	149	81	35	2	-
Male	3	10	11	15	28	39	47	66	35	10	1	-
Female	4	9	8	21	25	46	71	83	46	25	1	-
Alabama												
All races, both sexes	5	9	10	13	30	50	65	79	42	15	7	-
Male	3	6	7	6	23	25	28	27	10	4	1	-
Female	2	3	3	7	7	25	37	52	32	11	6	-
White, both sexes	3	6	8	10	21	34	48	62	33	11	4	-
Male	1	3	6	4	16	20	17	20	8	3	-	-
Female	2	3	2	6	5	14	31	42	25	8	4	-
Black, both sexes	2	3	2	3	9	16	17	17	9	4	3	-
Male	2	3	1	2	7	5	11	7	2	1	1	-
Female	-	-	1	1	2	11	6	10	7	3	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Alaska												
All races, both sexes	-	-	-	5	2	3	3	8	8	-	-	-
Male	-	-	-	3	2	2	1	3	4	-	-	-
Female	-	-	-	2	-	1	2	5	4	-	-	-
White, both sexes	-	-	-	5	1	2	2	4	6	-	-	-
Male	-	-	-	3	1	1	-	1	2	-	-	-
Female	-	-	-	2	-	1	2	3	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	1	1	2	1	-	-	-
Male	-	-	-	-	1	1	1	2	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Arizona												
All races, both sexes	3	7	11	13	20	28	61	78	54	24	2	-
Male	1	2	10	8	6	9	22	23	20	4	-	-
Female	2	5	1	5	14	19	39	55	34	20	2	-
White, both sexes	3	7	11	12	19	27	59	74	52	23	2	-
Male	1	2	10	8	5	9	21	23	18	4	-	-
Female	2	5	1	4	14	18	38	51	34	19	2	-
Black, both sexes	-	-	-	1	-	-	2	2	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	1	-	-	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	2	1	1	-	-
Male	-	-	-	-	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	2	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Arkansas												
All races, both sexes	10	6	16	14	32	40	69	78	63	19	5	-
Male	7	5	9	8	19	20	32	31	15	4	1	-
Female	3	1	7	6	13	20	37	47	48	15	4	-
White, both sexes	4	5	10	11	28	34	61	69	54	16	4	-
Male	4	5	6	7	16	17	29	29	13	3	1	-
Female	-	-	4	4	12	17	32	40	41	13	3	-
Black, both sexes	6	1	6	3	4	5	7	8	9	3	1	-
Male	3	-	3	1	3	2	2	2	2	1	-	-
Female	3	1	3	2	1	3	5	6	7	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
California												
All races, both sexes	55	62	95	144	219	457	685	886	633	325	48	-
Male	30	40	49	85	112	221	276	332	181	68	8	-
Female	25	22	46	59	107	236	409	554	452	257	40	-
White, both sexes	37	42	64	108	154	373	553	731	551	281	42	-
Male	21	30	33	63	75	188	224	270	154	59	7	-
Female	16	12	31	45	79	185	329	461	397	222	35	-
Black, both sexes	14	8	20	15	31	39	47	61	36	19	5	-
Male	7	4	9	12	20	13	16	18	6	3	-	-
Female	7	4	11	3	11	26	31	43	30	16	5	-
American Indian 1/, both sexes	-	-	1	-	-	3	3	3	1	-	-	-
Male	-	-	-	-	-	1	1	2	-	-	-	-
Female	-	-	1	-	-	2	2	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	4	12	10	21	34	42	82	91	45	25	1	-
Male	2	6	7	10	17	19	35	42	21	6	1	-
Female	2	6	3	11	17	23	47	49	24	19	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Colorado												
All races, both sexes	3	6	8	9	27	45	62	61	59	24	11	-
Male	3	4	3	5	13	23	18	18	20	6	-	-
Female	-	2	5	4	14	22	44	43	39	18	11	-
White, both sexes	3	6	7	9	26	42	61	57	57	23	9	-
Male	3	4	3	5	12	21	18	17	19	6	-	-
Female	-	2	4	4	14	21	43	40	38	17	9	-
Black, both sexes	-	-	1	-	1	3	-	2	-	1	2	-
Male	-	-	-	-	1	2	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	2	-	1	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	2	2	-	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	-	1	1	1	-	-	-
Connecticut												
All races, both sexes	-	5	1	7	11	14	31	84	48	27	5	-
Male	-	3	1	1	4	5	14	27	11	6	-	-
Female	-	2	-	6	7	9	17	57	37	21	5	-
White, both sexes	-	4	1	5	11	12	28	79	47	27	5	-
Male	-	2	1	1	4	4	12	27	11	6	-	-
Female	-	2	-	4	7	8	16	52	36	21	5	-
Black, both sexes	-	-	-	2	-	2	3	5	1	-	-	-
Male	-	-	-	-	-	1	2	-	-	-	-	-
Female	-	-	-	2	-	1	1	5	1	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Delaware												
All races, both sexes	-	1	2	5	8	12	12	15	11	5	1	-
Male	-	-	1	3	5	4	4	6	4	-	-	-
Female	-	1	1	2	3	8	8	9	7	5	1	-
White, both sexes	-	1	1	3	3	9	9	12	8	4	1	-
Male	-	-	-	2	2	2	3	4	2	-	-	-
Female	-	1	1	1	1	7	6	8	6	4	1	-
Black, both sexes	-	-	1	2	5	3	3	3	2	1	-	-
Male	-	-	1	1	3	2	1	2	1	-	-	-
Female	-	-	-	1	2	1	2	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	2	2	1	-	3	3	3	8	6	2	2	-
Male	2	1	1	-	2	-	1	3	2	-	1	-
Female	-	1	-	-	1	3	2	5	4	2	1	-
White, both sexes	1	-	-	-	-	-	1	3	2	1	1	-
Male	1	-	-	-	-	-	-	1	-	-	1	-
Female	-	-	-	-	-	-	1	2	2	1	-	-
Black, both sexes	1	2	1	-	3	3	2	5	4	1	1	-
Male	1	1	1	-	2	-	1	2	2	-	-	-
Female	-	1	-	-	1	3	1	3	2	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Florida												
All races, both sexes	14	27	51	63	93	220	289	368	287	120	28	-
Male	6	17	26	38	51	97	113	141	94	29	3	-
Female	8	10	25	25	42	123	176	227	193	91	25	-
White, both sexes	6	22	42	41	65	194	255	342	266	109	24	-
Male	4	14	23	26	35	87	99	136	92	24	-	-
Female	2	8	19	15	30	107	156	206	174	85	24	-
Black, both sexes	8	5	8	21	27	26	33	25	20	10	4	-
Male	2	3	3	12	15	10	13	5	2	4	3	-
Female	6	2	5	9	12	16	20	20	18	6	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	1	-	1	1	-	1	-	-
Male	-	-	-	-	1	-	1	-	-	1	-	-
Female	-	-	1	1	-	-	-	1	-	-	-	-
Georgia												
All races, both sexes	7	12	24	34	49	56	108	115	58	33	4	-
Male	4	8	14	20	28	27	47	29	13	5	1	-
Female	3	4	10	14	21	29	61	86	45	28	3	-
White, both sexes	6	6	12	19	31	41	76	91	44	26	3	-
Male	3	4	8	11	15	22	31	23	10	3	1	-
Female	3	2	4	8	16	19	45	68	34	23	2	-
Black, both sexes	1	6	12	15	17	15	32	24	12	7	1	-
Male	1	4	6	9	12	5	16	6	3	2	-	-
Female	-	2	6	6	5	10	16	18	9	5	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	2	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Hawaii												
All races, both sexes	2	-	2	5	3	13	16	24	15	2	2	-
Male	-	-	2	2	2	6	9	9	10	-	1	-
Female	2	-	-	3	1	7	7	15	5	2	1	-
White, both sexes	-	-	1	-	1	3	6	4	3	-	-	-
Male	-	-	1	-	-	1	3	1	2	-	-	-
Female	-	-	-	-	1	2	3	3	1	-	-	-
Black, both sexes	-	-	-	-	1	1	1	-	-	-	1	-
Male	-	-	-	-	1	1	1	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	1	5	1	9	9	20	12	2	1	-
Male	-	-	1	2	1	4	5	8	8	-	-	-
Female	2	-	-	3	-	5	4	12	4	2	1	-
Idaho												
All races, both sexes	1	5	6	1	7	13	21	36	43	13	1	-
Male	-	2	3	-	4	6	8	9	17	5	-	-
Female	1	3	3	1	3	7	13	27	26	8	1	-
White, both sexes	1	4	5	1	7	13	20	36	42	13	1	-
Male	-	2	3	-	4	6	8	9	17	5	-	-
Female	1	2	2	1	3	7	12	27	25	8	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	1	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Illinois												
All races, both sexes	7	18	27	41	79	124	192	235	186	82	17	-
Male	3	7	15	22	43	65	80	89	44	6	4	-
Female	4	11	12	19	36	59	112	146	142	76	13	-
White, both sexes	2	10	16	30	56	100	150	207	172	74	16	-
Male	-	3	9	15	32	53	63	77	42	6	4	-
Female	2	7	7	15	24	47	87	130	130	68	12	-
Black, both sexes	5	8	10	11	23	22	36	25	11	8	1	-
Male	3	4	5	7	11	10	15	11	2	-	-	-
Female	2	4	5	4	12	12	21	14	9	8	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	2	6	3	3	-	-	-
Male	-	-	1	-	-	2	2	1	-	-	-	-
Female	-	-	-	-	-	-	4	2	3	-	-	-
Indiana												
All races, both sexes	2	5	16	23	36	46	99	110	78	34	10	-
Male	2	3	8	16	19	22	36	39	22	7	2	-
Female	-	2	8	7	17	24	63	71	56	27	8	-
White, both sexes	1	3	13	19	33	43	88	104	74	34	10	-
Male	1	2	7	14	18	20	32	38	21	7	2	-
Female	-	1	6	5	15	23	56	66	53	27	8	-
Black, both sexes	1	2	3	4	3	2	11	6	4	-	-	-
Male	1	1	1	2	1	1	4	1	1	-	-	-
Female	-	1	2	2	2	1	7	5	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Iowa												
All races, both sexes	1	-	4	6	15	33	52	69	45	23	3	-
Male	1	-	4	3	11	13	24	27	8	7	-	-
Female	-	-	-	3	4	20	28	42	37	16	3	-
White, both sexes	1	-	3	6	15	33	50	67	44	23	3	-
Male	1	-	3	3	11	13	23	26	8	7	-	-
Female	-	-	-	3	4	20	27	41	36	16	3	-
Black, both sexes	-	-	1	-	-	-	2	1	1	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	2	4	5	10	18	25	42	66	59	33	10	-
Male	1	2	2	6	13	9	17	22	24	5	1	-
Female	1	2	3	4	5	16	25	44	35	28	9	-
White, both sexes	1	2	4	6	18	22	40	64	55	33	10	-
Male	1	-	1	3	13	8	17	21	23	5	1	-
Female	-	2	3	3	5	14	23	43	32	28	9	-
Black, both sexes	1	2	1	2	-	2	2	2	4	-	-	-
Male	-	2	1	2	-	1	-	1	1	-	-	-
Female	1	-	-	-	-	1	2	1	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Kentucky												
All races, both sexes	2	10	9	13	17	32	53	72	35	16	5	-
Male	-	4	7	5	8	13	25	24	5	5	-	-
Female	2	6	2	8	9	19	28	48	30	11	5	-
White, both sexes	2	9	8	12	17	31	49	68	32	15	5	-
Male	-	3	6	4	8	13	22	23	5	4	-	-
Female	2	6	2	8	9	18	27	45	27	11	5	-
Black, both sexes	-	-	1	1	-	1	4	4	3	1	-	-
Male	-	-	1	1	-	-	3	1	-	1	-	-
Female	-	-	-	-	-	1	1	3	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	6	9	9	19	22	54	58	63	36	12	2	-
Male	2	5	4	8	11	22	24	19	12	3	-	-
Female	4	4	5	11	11	32	34	44	24	9	2	-
White, both sexes	3	4	4	12	18	42	51	48	31	9	2	-
Male	2	3	1	5	9	15	20	16	12	2	-	-
Female	1	1	3	7	9	27	31	32	19	7	2	-
Black, both sexes	3	5	5	7	4	12	7	15	5	3	-	-
Male	-	2	3	3	2	7	4	3	-	1	-	-
Female	3	3	2	4	2	5	3	12	5	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Maine												
All races, both sexes	3	1	3	4	7	20	14	17	18	14	1	-
Male	3	-	3	2	3	8	5	4	6	3	-	-
Female	-	1	-	2	4	12	9	13	12	11	1	-
White, both sexes	3	1	3	4	7	20	14	17	18	14	1	-
Male	3	-	3	2	3	8	5	4	6	3	-	-
Female	-	1	-	2	4	12	9	13	12	11	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	2	3	11	11	27	37	63	103	71	31	9	-
Male	-	-	4	6	13	16	27	38	14	6	-	-
Female	2	3	7	5	14	21	36	65	57	25	9	-
White, both sexes	1	1	4	7	18	25	41	85	58	26	8	-
Male	-	-	3	4	9	9	19	32	14	6	-	-
Female	1	1	1	3	9	16	22	53	44	20	8	-
Black, both sexes	1	2	7	3	9	12	19	15	13	5	1	-
Male	-	-	1	2	4	7	7	5	-	-	-	-
Female	1	2	6	1	5	5	12	10	13	5	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	3	3	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	1	-	-	2	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Massachusetts												
All races, both sexes	1	10	9	12	36	66	115	163	126	61	6	-
Male	1	7	4	7	24	33	42	55	43	11	-	-
Female	-	3	5	5	12	33	73	108	83	50	6	-
White, both sexes	1	7	8	12	31	59	112	161	120	59	6	-
Male	1	6	3	7	22	31	41	55	40	10	-	-
Female	-	1	5	5	9	28	71	106	80	49	6	-
Black, both sexes	-	1	1	-	3	7	3	1	5	-	-	-
Male	-	-	1	-	-	2	1	-	3	-	-	-
Female	-	1	-	-	3	5	2	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	-	1	-	-	1	1	2	-	-
Male	-	1	-	-	1	-	-	-	-	1	-	-
Female	-	1	-	-	-	-	-	1	1	1	-	-
Michigan												
All races, both sexes	12	12	39	55	73	139	254	272	188	72	10	-
Male	9	4	20	34	36	53	115	99	46	15	2	-
Female	3	8	19	21	37	86	139	173	142	57	8	-
White, both sexes	7	10	28	39	54	123	222	245	171	67	8	-
Male	6	3	16	25	25	48	103	88	42	14	2	-
Female	1	7	12	14	29	75	119	157	129	53	6	-
Black, both sexes	5	2	11	16	18	15	29	25	16	5	2	-
Male	3	1	4	9	10	4	12	10	4	1	-	-
Female	2	1	7	7	8	11	17	15	12	4	2	-
American Indian 1/, both sexes	-	-	-	-	1	1	1	1	1	-	-	-
Male	-	-	-	-	1	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Minnesota												
All races, both sexes	5	7	3	17	22	44	65	106	83	37	8	-
Male	2	3	2	8	13	25	31	39	23	6	1	-
Female	3	4	1	9	9	19	34	67	60	31	7	-
White, both sexes	4	5	3	16	21	42	62	105	80	37	8	-
Male	1	2	2	8	13	23	29	38	22	6	1	-
Female	3	3	1	8	8	19	33	67	58	31	7	-
Black, both sexes	1	-	-	-	1	1	3	-	3	-	-	-
Male	1	-	-	-	-	1	2	-	1	-	-	-
Female	-	-	-	-	1	-	1	-	2	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	1	-	1	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	4	7	12	10	16	17	28	32	19	7	5	-
Male	3	5	8	6	8	8	10	8	4	1	-	-
Female	1	2	4	4	8	9	18	24	15	6	5	-
White, both sexes	1	3	5	5	7	10	20	23	17	7	2	-
Male	-	2	3	3	4	5	5	5	3	1	-	-
Female	1	1	2	2	3	5	15	18	14	6	2	-
Black, both sexes	3	4	7	5	9	7	8	9	2	-	3	-
Male	3	3	5	3	4	3	5	3	1	-	-	-
Female	-	1	2	2	5	4	3	6	1	-	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Missouri												
All races, both sexes	7	14	17	15	29	65	120	126	112	43	9	-
Male	4	8	8	6	16	34	52	31	27	8	-	-
Female	3	6	9	9	13	31	68	95	85	35	9	-
White, both sexes	4	12	15	15	24	58	110	119	105	42	7	-
Male	2	6	7	6	13	31	48	30	25	7	-	-
Female	2	6	8	9	11	27	62	89	80	35	7	-
Black, both sexes	3	2	2	-	4	6	10	7	6	1	2	-
Male	2	2	1	-	3	3	4	1	2	1	-	-
Female	1	-	1	-	1	3	6	6	4	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	1	-	-	-
Montana												
All races, both sexes	1	-	4	6	5	10	15	23	15	6	-	-
Male	1	-	1	4	2	7	7	9	5	3	-	-
Female	-	-	3	2	3	3	8	14	10	3	-	-
White, both sexes	1	-	3	6	5	9	15	21	15	6	-	-
Male	1	-	1	4	2	6	7	9	5	3	-	-
Female	-	-	2	2	3	3	8	12	10	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Nebraska												
All races, both sexes	2	4	2	10	8	17	37	36	38	18	2	-
Male	2	3	2	6	6	7	19	15	10	2	-	-
Female	-	1	-	4	2	10	18	21	28	16	2	-
White, both sexes	-	2	2	8	8	16	36	34	37	18	2	-
Male	-	2	2	6	6	7	18	14	10	2	-	-
Female	-	-	-	2	2	9	18	20	27	16	2	-
Black, both sexes	2	2	-	2	-	-	1	2	-	-	-	-
Male	2	1	-	-	-	-	1	1	-	-	-	-
Female	-	1	-	2	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Nevada												
All races, both sexes	-	2	4	7	6	15	25	23	13	9	-	-
Male	-	1	3	2	3	5	5	7	3	2	-	-
Female	-	1	1	5	3	10	20	16	10	7	-	-
White, both sexes	-	2	3	5	5	11	25	21	13	8	-	-
Male	-	1	2	2	3	4	5	6	3	1	-	-
Female	-	1	1	3	2	7	20	15	10	7	-	-
Black, both sexes	-	-	1	1	-	2	-	-	-	1	-	-
Male	-	-	1	-	-	-	-	-	-	1	-	-
Female	-	-	-	1	-	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	2	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	1	1	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
New Hampshire												
All races, both sexes	-	3	3	2	8	9	24	21	21	14	1	-
Male	-	2	3	2	4	2	7	6	4	3	-	-
Female	-	1	-	-	4	7	17	15	17	11	1	-
White, both sexes	-	3	3	2	8	9	24	21	21	14	1	-
Male	-	2	3	2	4	2	7	6	4	3	-	-
Female	-	1	-	-	4	7	17	15	17	11	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	3	4	14	10	22	52	76	109	84	46	7	-
Male	2	3	6	7	12	24	28	32	26	13	2	-
Female	1	1	8	3	10	28	48	77	58	33	5	-
White, both sexes	3	3	11	9	19	43	64	100	78	38	5	-
Male	2	3	6	6	11	20	21	30	24	10	1	-
Female	1	-	5	3	8	23	43	70	54	28	4	-
Black, both sexes	-	1	3	1	3	6	10	7	4	8	2	-
Male	-	-	-	1	1	3	6	2	1	3	1	-
Female	-	1	3	-	2	3	4	5	3	5	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	3	1	2	1	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	-	-	2	1	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
New Mexico												
All races, both sexes	1	1	5	11	11	24	34	51	22	14	1	-
Male	-	-	3	6	9	16	10	17	10	1	1	-
Female	1	1	2	5	2	8	24	34	12	13	-	-
White, both sexes	1	1	5	8	8	22	32	48	20	13	1	-
Male	-	-	3	4	7	14	10	15	9	1	1	-
Female	1	1	2	4	1	8	22	33	11	12	-	-
Black, both sexes	-	-	-	2	1	1	-	-	-	-	-	-
Male	-	-	-	2	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	2	1	1	3	1	1	-	-
Male	-	-	-	-	1	1	-	2	1	-	-	-
Female	-	-	-	1	1	-	1	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
New York												
All races, both sexes	23	8	15	23	57	90	150	219	147	75	14	-
Male	10	7	9	13	38	37	57	72	30	13	3	-
Female	13	1	6	10	19	53	93	147	117	62	11	-
White, both sexes	15	7	12	16	49	79	131	198	135	72	13	-
Male	5	6	8	8	32	31	53	67	30	12	3	-
Female	10	1	4	8	17	48	78	131	105	60	10	-
Black, both sexes	8	1	3	6	5	8	18	19	10	3	1	-
Male	5	1	1	4	4	5	4	4	-	1	-	-
Female	3	-	2	2	1	3	14	15	10	2	1	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	2	3	1	2	2	-	-	-
Male	-	-	-	1	2	1	-	1	-	-	-	-
Female	-	-	-	-	-	2	1	1	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
North Carolina												
All races, both sexes	9	19	28	37	58	97	144	176	112	42	14	-
Male	8	12	13	17	31	50	59	63	27	9	1	-
Female	1	7	15	20	27	47	85	113	85	33	13	-
White, both sexes	4	10	20	19	39	76	112	138	97	35	11	-
Male	4	6	10	10	22	37	49	52	25	8	1	-
Female	-	4	10	9	17	39	63	86	72	27	10	-
Black, both sexes	5	8	7	17	17	19	32	37	14	7	3	-
Male	4	5	3	6	9	12	10	11	2	1	-	-
Female	1	3	4	11	8	7	22	26	12	6	3	-
American Indian 1/, both sexes	-	1	1	1	1	2	-	1	1	-	-	-
Male	-	1	-	1	-	1	-	-	-	-	-	-
Female	-	-	1	-	1	1	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	1	1	1	5	8	11	16	9	-	-
Male	-	-	-	1	-	3	3	5	3	2	-	-
Female	-	-	1	-	1	2	5	6	13	7	-	-
White, both sexes	-	-	1	1	1	5	8	11	16	9	-	-
Male	-	-	-	1	-	3	3	5	3	2	-	-
Female	-	-	1	-	1	2	5	6	13	7	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Ohio												
All races, both sexes	16	31	43	67	134	183	273	344	287	85	23	-
Male	9	19	28	30	73	88	122	135	74	12	1	-
Female	7	12	15	37	61	95	151	209	213	73	22	-
White, both sexes	13	22	31	45	104	154	241	314	267	82	20	-
Male	9	15	20	20	57	79	107	121	69	12	1	-
Female	4	7	11	25	47	75	134	193	198	70	19	-
Black, both sexes	3	8	10	22	30	28	32	26	19	3	3	-
Male	-	4	7	10	16	9	15	12	5	-	-	-
Female	3	4	3	12	14	19	17	14	14	3	3	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	-	1	-	3	1	-	-	-
Male	-	-	-	-	-	-	-	2	-	-	-	-
Female	-	1	1	-	-	1	-	1	1	-	-	-
Oklahoma												
All races, both sexes	3	7	4	22	17	38	67	79	66	19	2	-
Male	1	4	1	12	6	18	26	22	15	3	1	-
Female	2	3	3	10	11	20	41	57	51	16	1	-
White, both sexes	2	6	4	17	13	36	59	76	61	18	1	-
Male	1	4	1	8	5	18	23	21	15	3	-	-
Female	1	2	3	9	8	18	36	55	46	15	1	-
Black, both sexes	1	-	-	1	2	-	6	1	2	-	1	-
Male	-	-	-	1	-	-	2	-	-	-	1	-
Female	1	-	-	-	2	-	4	1	2	-	-	-
American Indian 1/, both sexes	-	1	-	4	2	2	2	2	3	1	-	-
Male	-	-	-	3	1	-	1	1	-	-	-	-
Female	-	1	-	1	1	2	1	1	3	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Oregon												
All races, both sexes	4	8	11	15	24	48	61	128	89	27	6	-
Male	2	4	6	13	16	33	22	55	27	7	-	-
Female	2	4	5	2	8	15	39	73	62	20	6	-
White, both sexes	4	7	10	13	18	44	59	124	87	26	6	-
Male	2	3	6	12	13	29	22	54	25	6	-	-
Female	2	4	4	1	5	15	37	70	62	20	6	-
Black, both sexes	-	1	-	1	2	1	-	1	1	-	-	-
Male	-	1	-	1	2	1	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	4	3	2	2	1	1	-	-
Male	-	-	-	-	1	3	-	1	1	1	-	-
Female	-	-	1	1	3	-	2	1	-	-	-	-
Pennsylvania												
All races, both sexes	6	17	27	39	64	111	224	270	223	84	13	-
Male	2	9	17	25	31	60	95	108	50	14	1	-
Female	4	8	10	14	33	51	129	162	173	70	12	-
White, both sexes	5	11	19	33	53	98	204	251	210	80	12	-
Male	2	7	11	19	25	51	85	103	46	12	1	-
Female	3	4	8	14	28	47	119	148	164	68	11	-
Black, both sexes	1	6	7	5	10	10	18	18	13	4	1	-
Male	-	2	5	5	6	7	8	4	4	2	-	-
Female	1	4	2	-	4	3	10	14	9	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	1	3	2	1	-	-	-	-
Male	-	-	1	1	-	2	2	1	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Rhode Island												
All races, both sexes	1	2	1	3	4	10	16	20	18	6	1	-
Male	-	1	-	-	3	2	6	5	3	1	-	-
Female	1	1	1	3	1	8	10	15	15	5	1	-
White, both sexes	1	2	1	3	4	10	16	20	17	6	1	-
Male	-	1	-	-	3	2	6	5	3	1	-	-
Female	1	1	1	3	1	8	10	15	14	5	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	10	13	20	30	39	38	72	79	57	28	4	-
Male	6	9	12	20	16	17	30	24	10	1	1	-
Female	4	4	8	10	23	21	42	55	47	27	3	-
White, both sexes	5	5	10	16	22	23	42	60	48	21	1	-
Male	4	3	4	10	8	10	17	19	8	1	-	-
Female	1	2	6	6	14	13	25	41	40	20	1	-
Black, both sexes	5	8	10	14	17	14	29	18	9	7	3	-
Male	2	6	8	10	8	7	12	5	2	-	1	-
Female	3	2	2	4	9	7	17	13	7	7	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
South Dakota												
All races, both sexes	2	1	1	-	2	3	12	20	21	6	2	-
Male	1	1	1	-	1	1	9	11	5	2	-	-
Female	1	-	-	-	1	2	3	9	16	4	2	-
White, both sexes	2	1	1	-	1	3	11	20	21	6	2	-
Male	1	1	1	-	-	1	8	11	5	2	-	-
Female	1	-	-	-	1	2	3	9	16	4	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	6	8	17	30	38	62	94	115	88	34	7	-
Male	2	1	6	17	20	31	35	40	21	4	-	-
Female	4	7	11	13	18	31	59	75	67	30	7	-
White, both sexes	1	7	14	26	28	52	81	102	80	28	7	-
Male	-	1	5	14	18	25	28	33	19	4	-	-
Female	1	6	9	12	10	27	53	69	61	24	7	-
Black, both sexes	5	1	3	4	10	10	13	13	8	6	-	-
Male	2	-	1	3	2	6	7	7	2	-	-	-
Female	3	1	2	1	8	4	6	6	6	6	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Texas												
All races, both sexes	22	46	76	82	145	258	399	434	307	121	27	-
Male	13	31	46	40	76	122	151	151	83	18	2	-
Female	9	15	30	42	69	136	248	283	224	103	25	-
White, both sexes	12	35	45	56	118	210	351	388	263	108	23	-
Male	7	24	30	29	64	102	136	141	69	14	1	-
Female	5	11	15	27	54	108	215	247	194	94	22	-
Black, both sexes	10	10	29	25	25	40	44	40	38	11	4	-
Male	6	6	16	11	10	16	15	6	11	3	1	-
Female	4	4	13	14	15	24	29	34	27	8	3	-
American Indian 1/, both sexes	-	-	-	-	-	2	-	1	1	-	-	-
Male	-	-	-	-	-	2	-	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	2	1	2	6	4	5	5	2	-	-
Male	-	1	-	-	2	2	-	3	2	1	-	-
Female	-	-	2	1	-	4	4	2	3	1	-	-
Utah												
All races, both sexes	1	5	6	6	9	17	39	42	28	6	-	-
Male	-	1	4	2	4	10	17	14	5	-	-	-
Female	1	4	2	4	5	7	22	28	23	6	-	-
White, both sexes	1	5	6	6	8	16	38	42	27	6	-	-
Male	-	1	4	2	4	9	16	14	5	-	-	-
Female	1	4	2	4	4	7	22	28	22	6	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Vermont												
All races, both sexes	-	-	-	1	4	6	25	23	16	7	-	-
Male	-	-	-	1	2	1	9	9	6	4	-	-
Female	-	-	-	-	2	5	16	14	10	3	-	-
White, both sexes	-	-	-	1	4	6	25	23	16	7	-	-
Male	-	-	-	1	2	1	9	9	6	4	-	-
Female	-	-	-	-	2	5	16	14	10	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	3	14	10	26	38	84	100	131	83	37	9	-
Male	1	10	7	17	22	46	35	38	26	9	3	-
Female	2	4	3	9	16	38	65	93	57	28	6	-
White, both sexes	2	8	7	15	25	53	85	111	70	31	8	-
Male	1	5	4	10	12	28	30	29	24	8	3	-
Female	1	3	3	5	13	25	55	82	46	23	5	-
Black, both sexes	1	5	3	11	12	30	15	18	13	6	1	-
Male	-	4	3	7	9	17	5	8	2	1	-	-
Female	1	1	-	4	3	13	10	10	11	5	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	1	-	2	-	-	-	-
Male	-	1	-	-	1	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Washington												
All races, both sexes	7	7	16	18	32	62	100	156	104	37	5	-
Male	7	6	11	12	18	36	40	55	38	12	1	-
Female	-	1	5	6	14	26	60	101	66	25	4	-
White, both sexes	6	6	13	16	27	55	92	152	97	36	5	-
Male	6	6	8	10	15	32	40	52	37	12	1	-
Female	-	-	5	6	12	23	52	100	60	24	4	-
Black, both sexes	-	1	1	2	2	4	4	1	4	-	-	-
Male	-	-	1	2	1	3	-	1	1	-	-	-
Female	-	1	-	-	1	1	4	-	3	-	-	-
American Indian 1/, both sexes	-	-	1	-	2	1	2	-	-	-	-	-
Male	-	-	1	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	1	2	2	3	3	1	-	-
Male	1	-	1	-	1	-	-	2	-	-	-	-
Female	-	-	-	-	-	2	2	1	3	1	-	-
West Virginia												
All races, both sexes	1	6	7	6	13	26	33	45	30	16	1	-
Male	-	4	2	3	5	11	10	14	4	3	-	-
Female	1	2	5	3	8	15	23	31	26	13	1	-
White, both sexes	1	5	7	6	12	23	30	42	27	16	1	-
Male	-	3	2	3	5	9	9	14	4	3	-	-
Female	1	2	5	3	7	14	21	28	23	13	1	-
Black, both sexes	-	1	-	-	1	3	3	3	3	-	-	-
Male	-	1	-	-	-	2	1	-	-	-	-	-
Female	-	-	-	-	1	1	2	3	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Wisconsin												
All races, both sexes	3	8	12	27	35	62	124	138	109	50	6	-
Male	2	7	8	18	23	29	58	45	38	10	-	-
Female	1	1	4	9	12	33	66	93	71	40	6	-
White, both sexes	2	7	11	24	30	59	120	135	107	48	6	-
Male	1	6	7	17	19	28	56	44	37	10	-	-
Female	1	1	4	7	11	31	64	91	70	38	6	-
Black, both sexes	-	1	1	3	4	2	3	2	2	1	-	-
Male	-	1	1	1	3	1	1	1	1	-	-	-
Female	-	-	-	2	1	1	2	1	1	1	-	-
American Indian 1/, both sexes	1	-	-	-	1	1	1	1	-	-	-	-
Male	1	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Wyoming												
All races, both sexes	1	-	-	-	3	2	4	3	4	1	3	-
Male	1	-	-	-	2	2	2	1	-	-	-	-
Female	-	-	-	-	1	-	2	2	4	1	3	-
White, both sexes	1	-	-	-	3	2	4	3	3	1	3	-
Male	1	-	-	-	2	2	2	1	-	-	-	-
Female	-	-	-	-	1	-	2	2	3	1	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
United States												
All races, both sexes	30,172	14	4	4	10	23	44	43	94	193	318	472
Male	14,700	8	1	4	7	16	32	30	62	143	226	335
Female	15,472	6	3	-	3	7	12	13	32	50	92	137
White, both sexes	26,369	10	4	3	8	16	32	34	68	133	215	325
Male	12,832	6	1	3	6	11	22	25	44	102	152	236
Female	13,537	4	3	-	2	5	10	9	24	31	63	89
Black, both sexes	3,117	2	-	1	1	6	8	5	21	53	94	132
Male	1,501	2	-	1	1	4	7	3	14	35	67	87
Female	1,616	-	-	-	-	2	1	2	7	18	27	45
American Indian 1/, both sexes	140	1	-	-	-	1	1	1	1	2	3	5
Male	80	-	-	-	-	1	1	1	1	2	2	5
Female	60	1	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	546	1	-	-	1	-	3	3	4	5	6	10
Male	287	-	-	-	-	-	2	1	3	4	5	7
Female	259	1	-	-	1	-	1	2	1	1	1	3
Alabama												
All races, both sexes	572	1	-	-	-	-	1	3	-	6	12	17
Male	284	1	-	-	-	-	1	1	-	3	10	9
Female	288	-	-	-	-	-	-	2	-	3	2	8
White, both sexes	445	1	-	-	-	-	1	2	-	5	9	10
Male	225	1	-	-	-	-	1	1	-	3	7	6
Female	220	-	-	-	-	-	-	1	-	2	2	4
Black, both sexes	123	-	-	-	-	-	-	-	-	1	3	7
Male	56	-	-	-	-	-	-	-	-	-	3	3
Female	67	-	-	-	-	-	-	-	-	1	-	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Alaska												
All races, both sexes	31	-	-	1	-	-	-	-	-	1	1	1
Male	16	-	-	1	-	-	-	-	-	1	-	1
Female	15	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	27	-	-	1	-	-	-	-	-	1	1	1
Male	15	-	-	1	-	-	-	-	-	1	-	1
Female	12	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	592	-	1	-	1	2	2	2	5	5	8	8
Male	316	-	-	-	-	2	2	2	5	5	8	5
Female	276	-	1	-	1	-	-	-	-	-	-	3
White, both sexes	560	-	1	-	1	1	1	2	3	3	7	6
Male	299	-	-	-	-	1	1	2	3	3	7	4
Female	261	-	1	-	1	-	-	-	-	-	-	2
Black, both sexes	15	-	-	-	-	-	-	-	2	1	1	1
Male	7	-	-	-	-	-	-	-	2	1	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	13	-	-	-	-	1	1	-	-	1	-	1
Male	8	-	-	-	-	1	1	-	-	1	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Arkansas												
All races, both sexes	295	-	-	-	-	-	-	-	1	3	3	4
Male	144	-	-	-	-	-	-	-	1	3	1	3
Female	151	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	260	-	-	-	-	-	-	-	-	2	3	3
Male	128	-	-	-	-	-	-	-	-	2	1	2
Female	132	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	32	-	-	-	-	-	-	-	1	1	-	1
Male	16	-	-	-	-	-	-	-	1	1	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	2,970	1	-	-	1	2	6	7	15	20	35	44
Male	1,474	1	-	-	1	2	4	6	11	17	25	33
Female	1,496	-	-	-	-	-	2	1	4	3	10	11
White, both sexes	2,482	1	-	-	1	2	5	6	11	14	27	34
Male	1,203	1	-	-	1	2	3	6	7	12	17	24
Female	1,279	-	-	-	-	-	2	-	4	2	10	10
Black, both sexes	217	-	-	-	-	-	-	-	1	4	6	4
Male	114	-	-	-	-	-	-	-	1	3	6	3
Female	103	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	16	-	-	-	-	-	-	-	-	-	-	2
Male	11	-	-	-	-	-	-	-	-	-	-	2
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	255	-	-	-	-	-	1	1	3	2	2	4
Male	146	-	-	-	-	-	1	-	3	2	2	4
Female	109	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Colorado												
All races, both sexes	535	1	-	-	-	-	-	-	3	2	3	5
Male	228	1	-	-	-	-	-	-	2	2	2	3
Female	307	-	-	-	-	-	-	-	1	-	1	2
White, both sexes	506	1	-	-	-	-	-	-	3	2	2	5
Male	215	1	-	-	-	-	-	-	2	2	2	3
Female	291	-	-	-	-	-	-	-	1	-	-	2
Black, both sexes	19	-	-	-	-	-	-	-	-	-	1	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	368	-	1	1	-	-	-	-	3	5	4	4
Male	152	-	-	1	-	-	-	-	2	4	3	2
Female	216	-	1	-	-	-	-	-	1	1	1	2
White, both sexes	336	-	1	-	-	-	-	-	2	3	2	3
Male	135	-	-	-	-	-	-	-	1	3	1	2
Female	201	-	1	-	-	-	-	-	1	-	1	1
Black, both sexes	27	-	-	1	-	-	-	-	1	2	1	1
Male	15	-	-	1	-	-	-	-	1	1	1	-
Female	12	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Delaware												
All races, both sexes	67	-	-	-	-	-	-	-	-	3	-	2
Male	29	-	-	-	-	-	-	-	-	2	-	2
Female	38	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	54	-	-	-	-	-	-	-	-	2	-	2
Male	26	-	-	-	-	-	-	-	-	2	-	2
Female	28	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	11	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	58	1	-	-	-	-	-	-	1	-	2	3
Male	26	1	-	-	-	-	-	-	1	-	2	2
Female	32	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	10	1	-	-	-	-	-	-	-	-	-	-
Male	4	1	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	46	-	-	-	-	-	-	-	1	-	2	3
Male	22	-	-	-	-	-	-	-	1	-	2	2
Female	24	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Florida												
All races, both sexes	2,181	-	-	1	-	-	3	1	5	14	19	31
Male	1,128	-	-	1	-	-	2	1	2	10	16	21
Female	1,053	-	-	-	-	-	1	-	3	4	3	10
White, both sexes	1,987	-	-	1	-	-	2	1	2	10	12	23
Male	1,036	-	-	1	-	-	1	1	1	8	11	15
Female	951	-	-	-	-	-	1	-	1	2	1	8
Black, both sexes	188	-	-	-	-	-	1	-	3	4	6	8
Male	89	-	-	-	-	-	1	-	1	2	4	6
Female	99	-	-	-	-	-	-	-	2	2	2	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	644	1	-	-	2	-	1	1	2	12	10	18
Male	336	1	-	-	2	-	1	1	2	5	8	10
Female	308	-	-	-	-	-	-	-	-	7	2	8
White, both sexes	453	1	-	-	1	-	-	1	2	6	3	8
Male	233	1	-	-	1	-	-	1	2	2	3	5
Female	220	-	-	-	-	-	-	-	-	4	-	3
Black, both sexes	187	-	-	-	1	-	1	-	-	6	7	10
Male	100	-	-	-	1	-	1	-	-	3	5	5
Female	87	-	-	-	-	-	-	-	-	3	2	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Hawaii												
All races, both sexes	102	-	-	-	-	-	-	-	-	1	2	4
Male	53	-	-	-	-	-	-	-	-	-	2	4
Female	49	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	34	-	-	-	-	-	-	-	-	-	1	3
Male	16	-	-	-	-	-	-	-	-	-	1	3
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	66	-	-	-	-	-	-	-	-	1	1	1
Male	35	-	-	-	-	-	-	-	-	-	1	1
Female	31	-	-	-	-	-	-	-	-	1	-	-
Idaho												
All races, both sexes	146	-	-	-	-	-	-	-	-	1	-	3
Male	72	-	-	-	-	-	-	-	-	-	-	2
Female	74	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	142	-	-	-	-	-	-	-	-	1	-	3
Male	71	-	-	-	-	-	-	-	-	-	-	2
Female	71	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Illinois												
All races, both sexes	1,243	-	-	-	2	-	1	-	1	7	15	18
Male	556	-	-	-	2	-	1	-	1	7	9	10
Female	687	-	-	-	-	-	-	-	-	-	6	8
White, both sexes	1,056	-	-	-	2	-	1	-	1	2	9	10
Male	472	-	-	-	2	-	1	-	1	2	6	7
Female	584	-	-	-	-	-	-	-	-	-	3	3
Black, both sexes	173	-	-	-	-	-	-	-	-	5	6	8
Male	75	-	-	-	-	-	-	-	-	5	3	3
Female	98	-	-	-	-	-	-	-	-	-	3	5
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	795	-	-	-	1	-	1	-	3	3	6	9
Male	357	-	-	-	-	-	1	-	2	1	4	8
Female	438	-	-	-	1	-	-	-	1	2	2	1
White, both sexes	730	-	-	-	-	-	-	-	3	3	4	6
Male	327	-	-	-	-	-	-	-	2	1	3	5
Female	403	-	-	-	-	-	-	-	1	2	1	1
Black, both sexes	62	-	-	-	-	-	1	-	-	-	2	3
Male	29	-	-	-	-	-	1	-	-	-	1	3
Female	33	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Iowa												
All races, both sexes	427	-	-	-	-	-	-	-	1	1	4	7
Male	195	-	-	-	-	-	-	-	-	1	2	5
Female	232	-	-	-	-	-	-	-	1	-	2	2
White, both sexes	421	-	-	-	-	-	-	-	1	1	4	6
Male	191	-	-	-	-	-	-	-	-	1	2	4
Female	230	-	-	-	-	-	-	-	1	-	2	2
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	564	-	-	-	-	1	-	-	1	4	-	6
Male	254	-	-	-	-	1	-	-	1	4	-	5
Female	310	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	536	-	-	-	-	1	-	-	1	4	-	5
Male	245	-	-	-	-	1	-	-	1	4	-	4
Female	291	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Kentucky												
All races, both sexes	484	-	-	-	-	1	2	1	-	2	5	8
Male	238	-	-	-	-	-	2	1	-	1	2	7
Female	246	-	-	-	-	1	-	-	-	1	3	1
White, both sexes	454	-	-	-	-	1	2	1	-	2	5	8
Male	222	-	-	-	-	-	2	1	-	1	2	7
Female	232	-	-	-	-	1	-	-	-	1	3	1
Black, both sexes	29	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	460	-	1	-	1	-	2	1	1	-	5	8
Male	206	-	-	-	-	-	2	1	1	-	3	5
Female	254	-	1	-	1	-	-	-	-	-	2	3
White, both sexes	340	-	1	-	1	-	1	1	1	-	1	4
Male	150	-	-	-	-	-	1	1	1	-	1	3
Female	190	-	1	-	1	-	-	-	-	-	-	1
Black, both sexes	118	-	-	-	-	-	1	-	-	-	4	4
Male	55	-	-	-	-	-	1	-	-	-	2	2
Female	63	-	-	-	-	-	-	-	-	-	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Maine												
All races, both sexes	159	-	-	-	-	-	-	1	-	2	-	1
Male	79	-	-	-	-	-	-	1	-	1	-	1
Female	80	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	157	-	-	-	-	-	-	1	-	2	-	1
Male	79	-	-	-	-	-	-	1	-	1	-	1
Female	78	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	487	-	-	-	-	2	1	1	4	6	8	10
Male	218	-	-	-	-	-	-	1	3	5	6	5
Female	269	-	-	-	-	2	1	-	1	1	2	5
White, both sexes	374	-	-	-	-	1	1	-	3	2	3	4
Male	168	-	-	-	-	-	-	-	2	2	3	2
Female	206	-	-	-	-	1	1	-	1	-	-	2
Black, both sexes	109	-	-	-	-	1	-	1	1	4	5	5
Male	48	-	-	-	-	-	-	1	1	3	3	2
Female	61	-	-	-	-	1	-	-	-	1	2	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Massachusetts												
All races, both sexes	743	-	-	-	-	-	-	-	1	3	2	13
Male	349	-	-	-	-	-	-	-	-	2	1	10
Female	394	-	-	-	-	-	-	-	1	1	1	3
White, both sexes	717	-	-	-	-	-	-	-	1	3	2	13
Male	335	-	-	-	-	-	-	-	-	2	1	10
Female	382	-	-	-	-	-	-	-	1	1	1	3
Black, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1,235	1	-	-	-	2	-	-	1	10	16	19
Male	602	-	-	-	-	2	-	-	-	6	13	16
Female	633	1	-	-	-	-	-	-	1	4	3	3
White, both sexes	1,055	1	-	-	-	1	-	-	1	4	8	8
Male	523	-	-	-	-	1	-	-	-	1	7	8
Female	532	1	-	-	-	-	-	-	1	3	1	-
Black, both sexes	167	-	-	-	-	1	-	-	-	6	8	11
Male	70	-	-	-	-	1	-	-	-	5	6	8
Female	97	-	-	-	-	-	-	-	-	1	2	3
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Minnesota												
All races, both sexes	491	-	1	1	-	-	3	1	2	2	5	9
Male	253	-	1	1	-	-	1	-	1	1	2	9
Female	238	-	-	-	-	-	2	1	1	1	3	-
White, both sexes	466	-	1	1	-	-	2	1	1	2	4	6
Male	242	-	1	1	-	-	1	-	1	1	1	6
Female	224	-	-	-	-	-	1	1	-	1	3	-
Black, both sexes	15	-	-	-	-	-	-	-	-	-	1	2
Male	7	-	-	-	-	-	-	-	-	-	1	2
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	1	-	1	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	1	-	1	-	-	-
Mississippi												
All races, both sexes	212	-	-	-	-	1	-	-	1	2	3	4
Male	111	-	-	-	-	1	-	-	1	1	3	2
Female	101	-	-	-	-	-	-	-	-	1	-	2
White, both sexes	156	-	-	-	-	-	-	-	1	2	1	1
Male	79	-	-	-	-	-	-	-	1	1	1	-
Female	77	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	55	-	-	-	-	1	-	-	-	-	2	3
Male	32	-	-	-	-	1	-	-	-	-	2	2
Female	23	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Missouri												
All races, both sexes	727	-	-	-	-	1	1	-	2	4	6	7
Male	329	-	-	-	-	-	1	-	2	4	4	7
Female	398	-	-	-	-	1	-	-	-	-	2	-
White, both sexes	630	-	-	-	-	1	1	-	2	3	6	6
Male	289	-	-	-	-	-	1	-	2	3	4	6
Female	341	-	-	-	-	1	-	-	-	-	2	-
Black, both sexes	92	-	-	-	-	-	-	-	-	1	-	1
Male	36	-	-	-	-	-	-	-	-	1	-	1
Female	56	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	135	-	-	-	-	-	-	-	1	1	2	2
Male	72	-	-	-	-	-	-	-	-	-	1	1
Female	63	-	-	-	-	-	-	-	1	1	1	1
White, both sexes	128	-	-	-	-	-	-	-	1	1	1	2
Male	69	-	-	-	-	-	-	-	-	-	1	1
Female	59	-	-	-	-	-	-	-	1	1	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Nebraska												
All races, both sexes	230	-	-	-	-	-	-	-	-	2	2	-
Male	118	-	-	-	-	-	-	-	-	2	-	-
Female	112	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	225	-	-	-	-	-	-	-	-	2	2	-
Male	115	-	-	-	-	-	-	-	-	2	-	-
Female	110	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	273	-	-	-	-	-	2	-	1	3	1	3
Male	139	-	-	-	-	-	-	-	-	3	1	1
Female	134	-	-	-	-	-	2	-	1	-	-	2
White, both sexes	244	-	-	-	-	-	2	-	1	3	-	1
Male	128	-	-	-	-	-	-	-	-	3	-	-
Female	116	-	-	-	-	-	2	-	1	-	-	1
Black, both sexes	15	-	-	-	-	-	-	-	-	-	1	1
Male	6	-	-	-	-	-	-	-	-	-	1	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
New Hampshire												
All races, both sexes	136	1	-	-	-	-	-	-	-	1	1	2
Male	73	-	-	-	-	-	-	-	-	1	1	2
Female	63	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	135	1	-	-	-	-	-	-	-	1	1	2
Male	72	-	-	-	-	-	-	-	-	1	1	2
Female	63	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	818	-	-	-	-	2	2	1	-	4	12	9
Male	373	-	-	-	-	2	1	-	-	3	9	7
Female	445	-	-	-	-	-	1	1	-	1	3	2
White, both sexes	720	-	-	-	-	2	2	1	-	3	6	7
Male	332	-	-	-	-	2	1	-	-	3	4	6
Female	388	-	-	-	-	-	1	1	-	-	2	1
Black, both sexes	83	-	-	-	-	-	-	-	-	1	6	2
Male	35	-	-	-	-	-	-	-	-	-	5	1
Female	48	-	-	-	-	-	-	-	-	1	1	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
New Mexico												
All races, both sexes	166	-	-	-	-	-	-	1	1	1	1	2
Male	90	-	-	-	-	-	-	1	-	1	1	1
Female	76	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	156	-	-	-	-	-	-	-	1	1	1	2
Male	82	-	-	-	-	-	-	-	-	1	1	1
Female	74	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	1	-	-	-	-
Male	4	-	-	-	-	-	-	1	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1,676	1	-	-	-	2	3	3	4	9	16	25
Male	864	1	-	-	-	1	2	2	3	9	14	19
Female	812	-	-	-	-	1	1	1	1	-	2	6
White, both sexes	1,430	-	-	-	-	-	2	2	3	6	11	19
Male	740	-	-	-	-	-	1	2	3	6	9	14
Female	690	-	-	-	-	-	1	-	-	-	2	5
Black, both sexes	213	1	-	-	-	2	-	1	1	3	5	6
Male	109	1	-	-	-	1	-	-	-	3	5	5
Female	104	-	-	-	-	1	-	1	1	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	31	-	-	-	-	-	1	-	-	-	-	-
Male	14	-	-	-	-	-	1	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
North Carolina												
All races, both sexes	918	-	-	-	-	-	3	7	5	5	8	12
Male	467	-	-	-	-	-	3	5	3	3	7	9
Female	451	-	-	-	-	-	-	2	2	2	1	3
White, both sexes	720	-	-	-	-	-	3	5	1	3	5	6
Male	373	-	-	-	-	-	3	4	-	1	5	4
Female	347	-	-	-	-	-	-	1	1	2	-	2
Black, both sexes	188	-	-	-	-	-	-	2	3	2	3	6
Male	87	-	-	-	-	-	-	1	2	2	2	5
Female	101	-	-	-	-	-	-	1	1	-	1	1
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	1	-	-	-
Male	5	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	69	-	-	-	-	-	-	-	-	1	-	1
Male	35	-	-	-	-	-	-	-	-	1	-	1
Female	34	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	65	-	-	-	-	-	-	-	-	-	-	1
Male	32	-	-	-	-	-	-	-	-	-	-	1
Female	33	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Ohio												
All races, both sexes	1,415	-	-	-	1	1	1	1	3	6	14	28
Male	681	-	-	-	1	1	1	1	2	6	10	20
Female	734	-	-	-	-	-	-	-	1	-	4	8
White, both sexes	1,272	-	-	-	1	1	-	-	2	6	13	18
Male	598	-	-	-	1	1	-	-	2	6	9	13
Female	674	-	-	-	-	-	-	-	-	-	4	5
Black, both sexes	138	-	-	-	-	-	1	1	1	-	1	10
Male	79	-	-	-	-	-	1	1	-	-	1	7
Female	59	-	-	-	-	-	-	-	1	-	-	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	389	-	-	-	-	-	-	-	4	1	5	6
Male	184	-	-	-	-	-	-	-	3	1	1	5
Female	205	-	-	-	-	-	-	-	1	-	4	1
White, both sexes	350	-	-	-	-	-	-	-	2	1	5	4
Male	159	-	-	-	-	-	-	-	1	1	1	3
Female	191	-	-	-	-	-	-	-	1	-	4	1
Black, both sexes	16	-	-	-	-	-	-	-	2	-	-	2
Male	13	-	-	-	-	-	-	-	2	-	-	2
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Oregon												
All races, both sexes	386	-	-	-	-	1	-	1	1	-	6	6
Male	202	-	-	-	-	-	-	1	-	-	5	3
Female	184	-	-	-	-	1	-	-	1	-	1	3
White, both sexes	371	-	-	-	-	1	-	1	1	-	5	6
Male	193	-	-	-	-	-	-	1	-	-	4	3
Female	178	-	-	-	-	1	-	-	1	-	1	3
Black, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1,538	1	-	-	1	1	1	2	3	4	12	19
Male	699	-	-	-	1	1	1	2	3	3	8	15
Female	839	1	-	-	-	-	-	-	-	1	4	4
White, both sexes	1,419	-	-	-	1	1	1	1	3	3	7	11
Male	640	-	-	-	1	1	1	1	3	3	6	8
Female	779	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	107	-	-	-	-	-	-	-	-	1	4	8
Male	52	-	-	-	-	-	-	-	-	-	1	7
Female	55	-	-	-	-	-	-	-	-	1	3	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	1	-	-	-	-	-	1	-	-	-	-
Male	6	-	-	-	-	-	-	1	-	-	-	-
Female	5	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Rhode Island												
All races, both sexes	137	1	-	-	-	-	-	-	-	-	-	-
Male	46	1	-	-	-	-	-	-	-	-	-	-
Female	91	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	132	1	-	-	-	-	-	-	-	-	-	-
Male	44	1	-	-	-	-	-	-	-	-	-	-
Female	88	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	404	-	-	-	-	1	-	-	2	6	10	7
Male	208	-	-	-	-	-	-	-	2	1	3	5
Female	196	-	-	-	-	1	-	-	-	5	7	2
White, both sexes	293	-	-	-	-	1	-	-	2	2	3	3
Male	153	-	-	-	-	-	-	-	2	-	-	2
Female	140	-	-	-	-	1	-	-	-	2	3	1
Black, both sexes	110	-	-	-	-	-	-	-	-	4	7	4
Male	54	-	-	-	-	-	-	-	-	1	3	3
Female	56	-	-	-	-	-	-	-	-	3	4	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
South Dakota												
All races, both sexes	100	-	-	-	-	-	1	-	-	-	-	1
Male	51	-	-	-	-	-	1	-	-	-	-	1
Female	49	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	91	-	-	-	-	-	1	-	-	-	-	1
Male	50	-	-	-	-	-	1	-	-	-	-	1
Female	41	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	636	1	-	-	-	-	-	2	2	3	7	7
Male	306	1	-	-	-	-	-	1	-	3	6	5
Female	330	-	-	-	-	-	-	1	2	-	1	2
White, both sexes	539	-	-	-	-	-	-	2	1	3	4	5
Male	263	-	-	-	-	-	-	1	-	3	4	4
Female	276	-	-	-	-	-	-	1	1	-	-	1
Black, both sexes	92	1	-	-	-	-	-	-	1	-	3	2
Male	42	1	-	-	-	-	-	-	-	-	2	1
Female	50	-	-	-	-	-	-	-	1	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Texas												
All races, both sexes	1,564	1	-	-	-	-	4	4	7	13	21	41
Male	804	-	-	-	-	-	3	1	3	7	18	26
Female	760	1	-	-	-	-	1	3	4	6	3	15
White, both sexes	1,342	1	-	-	-	-	2	4	7	11	14	29
Male	692	-	-	-	-	-	1	1	3	7	12	20
Female	650	1	-	-	-	-	1	3	4	4	2	9
Black, both sexes	203	-	-	-	-	-	2	-	-	2	7	11
Male	102	-	-	-	-	-	2	-	-	-	6	6
Female	101	-	-	-	-	-	-	-	-	2	1	5
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	18	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	1
Utah												
All races, both sexes	164	1	-	-	-	-	-	-	-	1	4	2
Male	90	-	-	-	-	-	-	-	-	1	4	2
Female	74	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	158	1	-	-	-	-	-	-	-	1	3	1
Male	86	-	-	-	-	-	-	-	-	1	3	1
Female	72	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Vermont												
All races, both sexes	86	-	-	-	-	-	-	-	-	-	1	1
Male	44	-	-	-	-	-	-	-	-	-	1	1
Female	42	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	85	-	-	-	-	-	-	-	-	-	1	1
Male	44	-	-	-	-	-	-	-	-	-	1	1
Female	41	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	661	-	-	-	-	2	1	-	3	5	5	9
Male	311	-	-	-	-	2	1	-	3	5	1	7
Female	350	-	-	-	-	-	-	-	-	-	4	2
White, both sexes	504	-	-	-	-	1	1	-	1	2	4	6
Male	239	-	-	-	-	1	1	-	1	2	1	5
Female	265	-	-	-	-	-	-	-	-	-	3	1
Black, both sexes	147	-	-	-	-	1	-	-	2	3	1	3
Male	68	-	-	-	-	1	-	-	2	3	-	2
Female	79	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Washington												
All races, both sexes	718	1	-	-	-	-	-	1	-	5	7	14
Male	362	-	-	-	-	-	-	-	-	5	4	9
Female	356	1	-	-	-	-	-	1	-	-	3	5
White, both sexes	673	-	-	-	-	-	-	1	-	4	6	10
Male	342	-	-	-	-	-	-	-	-	4	4	7
Female	331	-	-	-	-	-	-	1	-	-	2	3
Black, both sexes	15	-	-	-	-	-	-	-	-	-	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	5	1	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	3	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	25	-	-	-	-	-	-	-	-	1	1	1
Male	13	-	-	-	-	-	-	-	-	1	-	-
Female	12	-	-	-	-	-	-	-	-	-	1	1
West Virginia												
All races, both sexes	259	-	-	-	-	-	1	1	1	1	1	2
Male	116	-	-	-	-	-	-	1	-	-	-	1
Female	143	-	-	-	-	-	1	-	1	1	1	1
White, both sexes	254	-	-	-	-	-	-	1	1	1	1	2
Male	115	-	-	-	-	-	-	1	-	-	-	1
Female	139	-	-	-	-	-	-	-	1	1	1	1
Black, both sexes	5	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Wisconsin												
All races, both sexes	637	-	-	-	-	1	1	-	2	2	6	9
Male	320	-	-	-	-	1	1	-	1	2	4	6
Female	317	-	-	-	-	-	-	-	1	-	2	3
White, both sexes	607	-	-	-	-	1	1	-	1	-	6	8
Male	304	-	-	-	-	1	1	-	1	-	4	5
Female	303	-	-	-	-	-	-	-	-	-	2	3
Black, both sexes	25	-	-	-	-	-	-	-	1	1	-	1
Male	13	-	-	-	-	-	-	-	-	1	-	1
Female	12	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	69	-	-	-	-	-	-	-	1	-	2	1
Male	40	-	-	-	-	-	-	-	1	-	1	1
Female	29	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	68	-	-	-	-	-	-	-	1	-	2	1
Male	39	-	-	-	-	-	-	-	1	-	1	1
Female	29	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
United States												
All races, both sexes	754	1,089	1,557	2,020	2,874	4,067	5,305	5,442	3,824	1,617	401	3
Male	525	755	1,068	1,292	1,656	2,129	2,511	2,226	1,257	359	55	3
Female	229	334	489	728	1,218	1,938	2,794	3,216	2,567	1,258	346	-
White, both sexes	533	828	1,275	1,686	2,496	3,604	4,765	4,977	3,540	1,459	356	2
Male	377	584	896	1,094	1,461	1,911	2,292	2,041	1,192	325	49	2
Female	156	244	379	592	1,035	1,693	2,473	2,936	2,348	1,134	307	-
Black, both sexes	193	228	226	267	307	371	416	381	229	134	41	1
Male	134	150	137	152	155	177	151	142	49	27	5	1
Female	59	78	89	115	152	194	265	239	180	107	36	-
American Indian 1/, both sexes	5	8	15	20	19	15	16	13	8	6	-	-
Male	2	7	7	14	12	6	9	5	2	3	-	-
Female	3	1	8	6	7	9	7	8	6	3	-	-
Asian or Pacific Islander 2/, both sexes	23	25	41	47	52	77	108	71	47	18	4	-
Male	12	14	28	32	28	35	59	38	14	4	1	-
Female	11	11	13	15	24	42	49	33	33	14	3	-
Alabama												
All races, both sexes	20	31	32	42	56	89	79	93	57	25	8	-
Male	12	24	23	30	29	42	37	38	17	5	2	-
Female	8	7	9	12	27	47	42	55	40	20	6	-
White, both sexes	12	25	27	35	45	69	63	76	44	16	5	-
Male	6	19	20	25	25	33	30	33	12	2	1	-
Female	6	6	7	10	20	36	33	43	32	14	4	-
Black, both sexes	8	4	5	7	11	20	16	17	12	9	3	-
Male	6	3	3	5	4	9	7	5	4	3	1	-
Female	2	1	2	2	7	11	9	12	8	6	2	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	1	-	-	-
Male	-	1	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Alaska												
All races, both sexes	4	3	1	1	3	3	2	5	5	-	-	-
Male	2	1	1	1	2	1	2	2	1	-	-	-
Female	2	2	-	-	1	2	-	3	4	-	-	-
White, both sexes	3	3	1	1	2	3	2	4	4	-	-	-
Male	2	1	1	1	1	1	2	2	1	-	-	-
Female	1	2	-	-	1	2	-	2	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	1	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	15	29	41	31	56	75	102	102	71	30	6	-
Male	11	22	26	17	30	46	56	35	29	12	3	-
Female	4	7	15	14	26	29	46	67	42	18	3	-
White, both sexes	14	27	39	29	54	71	99	100	66	30	6	-
Male	10	21	25	15	30	44	56	34	28	12	3	-
Female	4	6	14	14	24	27	43	66	38	18	3	-
Black, both sexes	1	1	1	1	-	1	2	-	3	-	-	-
Male	1	-	-	1	-	1	-	-	-	-	-	-
Female	-	1	1	-	-	-	2	-	3	-	-	-
American Indian 1/, both sexes	-	1	1	1	1	2	1	1	1	-	-	-
Male	-	1	1	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	1	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Arkansas												
All races, both sexes	9	11	18	27	30	49	45	51	30	11	3	-
Male	4	9	9	15	15	27	22	21	10	4	-	-
Female	5	2	9	12	15	22	23	30	20	7	3	-
White, both sexes	7	9	15	25	26	45	38	45	30	10	2	-
Male	3	8	8	15	14	25	20	16	10	4	-	-
Female	4	1	7	10	12	20	18	29	20	6	2	-
Black, both sexes	2	2	2	2	4	3	6	6	-	1	1	-
Male	1	1	1	-	1	2	2	5	-	-	-	-
Female	1	1	1	2	3	1	4	1	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	75	119	156	180	258	393	540	518	375	172	52	1
Male	53	86	116	121	156	209	236	217	122	45	12	1
Female	22	33	40	59	102	184	304	301	253	127	40	-
White, both sexes	47	89	125	143	214	322	450	456	338	153	43	1
Male	34	67	93	98	130	170	186	190	110	40	11	1
Female	13	22	32	45	84	152	264	266	228	113	32	-
Black, both sexes	15	20	18	17	17	27	37	30	11	5	5	-
Male	11	13	13	9	10	16	14	10	4	1	-	-
Female	4	7	5	8	7	11	23	20	7	4	5	-
American Indian 1/, both sexes	-	1	1	1	4	3	2	2	-	-	-	-
Male	-	1	1	1	3	1	1	1	-	-	-	-
Female	-	-	-	-	1	2	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	9	12	19	23	41	51	30	26	14	4	-
Male	8	5	9	13	13	22	35	16	8	4	1	-
Female	5	4	3	6	10	19	16	14	18	10	3	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Colorado												
All races, both sexes	8	18	15	30	49	56	97	111	82	44	11	-
Male	6	12	7	20	27	29	44	36	24	11	2	-
Female	2	6	8	10	22	27	53	75	58	33	9	-
White, both sexes	5	15	14	29	45	55	90	107	80	43	10	-
Male	4	11	7	20	25	28	41	33	24	11	1	-
Female	1	4	7	9	20	27	49	74	56	32	9	-
Black, both sexes	3	3	1	-	2	1	4	2	-	1	1	-
Male	2	1	-	-	2	1	1	1	-	-	1	-
Female	1	2	1	-	-	-	3	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	3	1	1	-	-	-
Male	-	-	-	-	-	-	2	1	-	-	-	-
Female	-	-	-	1	1	-	1	-	1	-	-	-
Connecticut												
All races, both sexes	5	3	16	16	24	56	60	69	65	26	10	-
Male	3	1	9	12	13	31	25	25	17	3	1	-
Female	2	2	7	4	11	25	35	44	48	23	9	-
White, both sexes	4	2	12	14	23	51	60	62	63	25	9	-
Male	3	-	7	11	13	28	25	22	17	2	-	-
Female	1	2	5	3	10	23	35	40	46	23	9	-
Black, both sexes	1	1	3	2	-	5	-	5	2	1	1	-
Male	-	1	2	1	-	3	-	2	-	1	1	-
Female	1	-	1	1	-	2	-	3	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	1	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Delaware												
All races, both sexes	1	1	1	6	5	12	14	12	8	1	1	-
Male	1	-	-	2	2	4	7	4	5	-	-	-
Female	-	1	1	4	3	8	7	8	3	1	1	-
White, both sexes	1	1	1	5	3	9	11	11	6	1	1	-
Male	1	-	-	2	1	4	6	4	4	-	-	-
Female	-	1	1	3	2	5	5	7	2	1	1	-
Black, both sexes	-	-	-	1	2	2	3	1	1	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	1	1	2	2	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	4	5	4	5	10	5	6	9	2	1	-	-
Male	4	3	1	2	5	1	-	4	-	-	-	-
Female	-	2	3	3	5	4	6	5	2	1	-	-
White, both sexes	-	-	-	1	4	-	2	2	-	-	-	-
Male	-	-	-	-	2	-	-	1	-	-	-	-
Female	-	-	-	1	2	-	2	1	-	-	-	-
Black, both sexes	4	4	4	4	6	4	4	7	2	1	-	-
Male	4	3	1	2	3	1	-	3	-	-	-	-
Female	-	1	3	2	3	3	4	4	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Florida												
All races, both sexes	64	77	113	139	202	301	423	357	283	114	34	-
Male	44	60	77	91	119	160	213	173	103	34	1	-
Female	20	17	36	48	83	141	210	184	180	80	33	-
White, both sexes	48	66	101	121	180	277	398	338	273	104	30	-
Male	34	52	72	80	103	149	208	165	102	32	1	-
Female	14	14	29	41	77	128	190	173	171	72	29	-
Black, both sexes	16	11	11	17	22	23	24	19	10	9	4	-
Male	10	8	4	11	16	11	5	8	1	1	-	-
Female	6	3	7	6	6	12	19	11	9	8	4	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	1	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	1	-	-	-	-	-
Georgia												
All races, both sexes	22	34	40	51	70	90	100	89	56	34	11	-
Male	15	21	28	35	43	52	56	32	21	3	-	-
Female	7	13	12	16	27	38	44	57	35	31	11	-
White, both sexes	12	16	23	34	52	65	89	69	42	21	8	-
Male	9	9	18	22	32	38	49	25	15	1	-	-
Female	3	7	5	12	20	27	40	44	27	20	8	-
Black, both sexes	10	17	16	16	18	25	11	19	14	13	3	-
Male	6	11	9	12	11	14	7	7	6	2	-	-
Female	4	6	7	4	7	11	4	12	8	11	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	1	-	-	-	1	-	-	-	-
Male	-	1	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Hawaii												
All races, both sexes	2	8	9	5	8	13	18	15	11	6	-	-
Male	1	5	7	3	7	4	7	8	4	1	-	-
Female	1	3	2	2	1	9	11	7	7	5	-	-
White, both sexes	-	2	4	1	2	4	5	4	5	3	-	-
Male	-	1	3	1	2	1	1	1	1	1	-	-
Female	-	1	1	-	-	3	4	3	4	2	-	-
Black, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	5	4	4	6	9	13	11	6	3	-	-
Male	1	3	3	2	5	3	6	7	3	-	-	-
Female	1	2	1	2	1	6	7	4	3	3	-	-
Idaho												
All races, both sexes	3	5	3	7	15	34	20	28	18	9	-	-
Male	2	3	1	6	9	19	12	10	7	1	-	-
Female	1	2	2	1	6	15	8	18	11	8	-	-
White, both sexes	2	5	3	7	14	32	20	28	18	9	-	-
Male	2	3	1	6	8	19	12	10	7	1	-	-
Female	-	2	2	1	6	13	8	18	11	8	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Illinois												
All races, both sexes	24	51	56	83	114	165	217	217	163	89	20	-
Male	19	34	40	51	64	80	92	82	46	17	1	-
Female	5	17	16	32	50	85	125	135	117	72	19	-
White, both sexes	21	41	45	67	96	140	188	190	151	74	18	-
Male	16	28	31	40	55	70	82	72	44	14	1	-
Female	5	13	14	27	41	70	106	118	107	60	17	-
Black, both sexes	2	10	11	14	16	20	29	24	11	15	2	-
Male	2	6	9	9	7	8	10	8	2	3	-	-
Female	-	4	2	5	9	12	19	16	9	12	2	-
American Indian 1/, both sexes	-	-	-	-	-	3	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	2	2	2	-	3	1	-	-	-
Male	1	-	-	2	2	1	-	2	-	-	-	-
Female	-	-	-	-	-	1	-	1	1	-	-	-
Indiana												
All races, both sexes	17	25	38	45	82	105	132	156	107	56	9	-
Male	15	13	22	23	54	52	55	67	31	8	1	-
Female	2	12	16	22	28	53	77	89	76	48	8	-
White, both sexes	13	21	33	40	78	95	122	147	105	51	9	-
Male	12	10	20	21	50	47	53	63	31	8	1	-
Female	1	11	13	19	28	48	69	84	74	43	8	-
Black, both sexes	3	4	5	5	4	10	9	9	2	5	-	-
Male	3	3	2	2	4	5	1	4	-	-	-	-
Female	-	1	3	3	-	5	8	5	2	5	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Iowa												
All races, both sexes	9	6	23	23	42	45	67	86	74	32	7	-
Male	9	5	12	16	23	24	34	35	27	2	-	-
Female	-	1	11	7	19	21	33	51	47	30	7	-
White, both sexes	7	6	22	23	42	45	66	85	74	32	7	-
Male	7	5	11	16	23	24	34	35	27	2	-	-
Female	-	1	11	7	19	21	32	50	47	30	7	-
Black, both sexes	1	-	1	-	-	-	-	1	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Kansas												
All races, both sexes	11	6	20	30	46	75	92	124	94	47	7	-
Male	9	5	13	14	30	35	41	50	33	12	1	-
Female	2	1	7	16	16	40	51	74	61	35	6	-
White, both sexes	9	6	18	28	42	73	89	119	89	45	7	-
Male	8	5	13	14	26	35	41	49	32	11	1	-
Female	1	1	5	14	16	38	48	70	57	34	6	-
Black, both sexes	2	-	1	2	4	1	3	4	3	2	-	-
Male	1	-	-	-	4	-	-	1	1	1	-	-
Female	1	-	1	2	-	1	3	3	2	1	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Kentucky												
All races, both sexes	14	21	27	48	52	61	83	87	47	19	6	-
Male	10	14	17	30	28	30	42	42	10	2	-	-
Female	4	7	10	18	24	31	41	45	37	17	6	-
White, both sexes	10	19	26	46	47	57	78	85	45	17	5	-
Male	6	12	17	29	25	28	39	42	9	2	-	-
Female	4	7	9	17	22	29	39	43	36	15	5	-
Black, both sexes	4	2	1	1	5	4	5	2	2	2	1	-
Male	4	2	-	-	3	2	3	-	1	-	-	-
Female	-	-	1	1	2	2	2	2	1	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	14	18	31	41	54	47	69	86	55	19	7	-
Male	8	13	21	26	24	23	26	28	18	6	1	-
Female	6	5	10	15	30	24	43	58	37	13	6	-
White, both sexes	6	9	20	28	39	36	55	72	46	13	6	-
Male	4	7	15	17	18	19	20	24	13	5	1	-
Female	2	2	5	11	21	17	35	48	33	8	5	-
Black, both sexes	7	9	11	13	14	11	14	14	9	6	1	-
Male	4	6	6	9	5	4	6	4	5	1	-	-
Female	3	3	5	4	9	7	8	10	4	5	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Maine												
All races, both sexes	2	7	11	11	17	24	24	37	16	5	1	-
Male	2	4	7	6	11	11	13	14	5	2	1	-
Female	-	3	4	5	6	13	11	23	11	3	-	-
White, both sexes	2	7	10	11	17	24	24	37	15	5	1	-
Male	2	4	7	6	11	11	13	14	5	2	1	-
Female	-	3	3	5	6	13	11	23	10	3	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Maryland												
All races, both sexes	17	15	23	29	46	75	84	82	58	21	5	-
Male	11	9	17	16	20	29	39	30	22	5	-	-
Female	6	6	6	13	26	46	45	52	36	16	5	-
White, both sexes	7	6	14	19	34	66	69	70	52	18	5	-
Male	4	3	12	11	17	27	35	26	21	3	-	-
Female	3	3	2	8	17	39	34	44	31	15	5	-
Black, both sexes	10	9	9	9	12	9	15	10	6	3	-	-
Male	7	6	5	5	3	2	4	3	1	2	-	-
Female	3	3	4	4	9	7	11	7	5	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Massachusetts												
All races, both sexes	11	20	26	38	65	103	128	170	101	49	13	-
Male	9	15	20	28	39	54	55	72	30	13	1	-
Female	2	5	6	10	26	49	73	98	71	36	12	-
White, both sexes	11	19	22	36	60	99	124	167	98	49	13	-
Male	9	14	16	26	36	51	54	72	30	13	1	-
Female	2	5	6	10	24	48	70	95	68	36	12	-
Black, both sexes	-	1	4	2	3	3	4	2	3	-	-	-
Male	-	1	4	2	1	3	1	-	-	-	-	-
Female	-	-	-	-	2	-	3	2	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Michigan												
All races, both sexes	39	43	68	86	118	174	214	210	175	46	13	-
Male	24	29	51	47	69	95	104	79	59	6	2	-
Female	15	14	17	39	49	79	110	131	116	40	11	-
White, both sexes	26	30	52	70	100	157	193	186	162	44	12	-
Male	18	20	41	42	64	87	97	73	56	6	2	-
Female	8	10	11	28	36	70	96	113	106	38	10	-
Black, both sexes	13	12	15	13	16	15	20	22	12	2	1	-
Male	6	9	10	3	3	6	6	4	3	-	-	-
Female	7	3	5	10	13	9	14	18	9	2	1	-
American Indian 1/, both sexes	-	1	1	2	1	1	1	2	-	-	-	-
Male	-	-	-	1	1	1	1	2	-	-	-	-
Female	-	1	1	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	1	-	-	1	-	-	-
Male	-	-	-	1	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Minnesota												
All races, both sexes	10	15	30	22	43	68	82	93	63	33	8	-
Male	8	10	19	17	28	43	42	41	23	5	1	-
Female	2	5	11	5	15	25	40	52	40	28	7	-
White, both sexes	7	13	24	21	43	67	79	92	61	33	8	-
Male	6	8	17	16	28	43	42	41	23	5	1	-
Female	1	5	7	5	15	24	37	51	38	28	7	-
Black, both sexes	2	1	3	-	-	1	3	-	2	-	-	-
Male	2	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	2	-	-	1	3	-	2	-	-	-
American Indian 1/, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	2	-	-	-	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	2	-	-	-	-	1	-	-	-	-
Mississippi												
All races, both sexes	11	16	14	15	26	28	32	27	18	10	2	2
Male	7	10	10	10	17	21	10	14	2	-	-	2
Female	4	6	4	5	9	7	22	13	16	10	2	-
White, both sexes	7	9	9	11	22	20	27	23	13	8	1	1
Male	5	6	7	7	15	15	8	11	1	-	-	1
Female	2	3	2	4	7	5	19	12	12	8	1	-
Black, both sexes	4	7	5	4	4	8	5	3	5	2	1	1
Male	2	4	3	3	2	6	2	3	1	-	-	1
Female	2	3	2	1	2	2	3	-	4	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Missouri												
All races, both sexes	19	20	24	45	68	91	131	147	102	46	13	-
Male	11	14	18	33	40	45	55	57	30	6	2	-
Female	8	6	6	12	28	46	76	90	72	40	11	-
White, both sexes	17	15	18	39	60	71	113	128	97	42	11	-
Male	10	11	15	27	38	35	49	51	30	5	2	-
Female	7	4	3	12	22	36	64	77	67	37	9	-
Black, both sexes	2	5	5	5	8	17	18	19	5	4	2	-
Male	1	3	2	5	2	8	6	6	-	1	-	-
Female	1	2	3	-	6	9	12	13	5	3	2	-
American Indian 1/, both sexes	-	-	-	1	-	2	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	5	6	13	5	18	13	29	16	15	7	2	-
Male	5	5	7	5	8	12	13	6	6	3	-	-
Female	-	1	6	-	10	1	16	10	9	4	2	-
White, both sexes	4	6	10	4	18	13	28	16	15	7	2	-
Male	4	5	7	4	8	12	12	6	6	3	-	-
Female	-	1	3	-	10	1	16	10	9	4	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	3	1	-	-	1	-	-	-	-	-
Male	1	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	3	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Nebraska												
All races, both sexes	5	6	8	13	19	19	44	59	34	15	4	-
Male	5	4	6	10	9	10	25	30	11	4	2	-
Female	-	2	2	3	10	9	19	29	23	11	2	-
White, both sexes	4	5	8	11	19	19	43	59	34	15	4	-
Male	4	3	6	9	9	10	25	30	11	4	2	-
Female	-	2	2	2	10	9	18	29	23	11	2	-
Black, both sexes	1	1	-	1	-	-	1	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	6	6	20	25	23	40	56	46	29	9	3	-
Male	4	5	14	16	12	21	28	22	8	3	1	-
Female	2	1	6	9	11	19	28	24	21	6	2	-
White, both sexes	4	5	15	20	20	35	53	46	27	9	3	-
Male	4	4	11	16	12	19	26	22	7	3	1	-
Female	-	1	4	4	8	16	27	24	20	6	2	-
Black, both sexes	-	1	1	4	1	4	1	-	1	-	-	-
Male	-	1	-	-	-	1	1	-	1	-	-	-
Female	-	-	1	4	1	3	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	4	1	1	1	1	-	-	-	-	-
Male	-	-	3	-	-	1	-	-	-	-	-	-
Female	2	-	1	1	1	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
New Hampshire												
All races, both sexes	-	7	11	4	9	19	23	33	16	8	1	-
Male	-	3	10	3	8	12	9	19	5	-	-	-
Female	-	4	1	1	1	7	14	14	11	8	1	-
White, both sexes	-	7	10	4	9	19	23	33	16	8	1	-
Male	-	3	9	3	8	12	9	19	5	-	-	-
Female	-	4	1	1	1	7	14	14	11	8	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	13	27	34	38	75	108	159	154	120	50	10	-
Male	11	20	20	22	37	54	78	53	41	14	1	-
Female	2	7	14	16	38	54	81	101	79	36	9	-
White, both sexes	11	20	30	25	67	101	138	139	113	46	9	-
Male	9	15	19	15	35	50	72	47	40	13	1	-
Female	2	5	11	10	32	51	66	92	73	33	8	-
Black, both sexes	2	7	3	10	6	6	16	13	6	4	1	-
Male	2	5	1	5	1	4	5	5	-	1	-	-
Female	-	2	2	5	5	2	11	8	6	3	1	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	2	2	1	5	1	1	-	-	-
Male	-	-	-	2	1	-	1	1	1	-	-	-
Female	-	-	1	-	1	1	4	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
New Mexico												
All races, both sexes	4	2	9	9	17	33	22	30	28	4	2	-
Male	4	1	8	4	11	16	15	11	14	1	1	-
Female	-	1	1	5	6	17	7	19	14	3	1	-
White, both sexes	3	2	8	9	15	33	20	29	27	3	2	-
Male	3	1	8	4	9	16	13	11	13	-	1	-
Female	-	1	-	5	6	17	7	18	14	3	1	-
Black, both sexes	1	-	-	-	1	-	1	-	1	-	-	-
Male	1	-	-	-	1	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	1	-	1	1	-	1	-	-
Male	-	-	-	-	1	-	1	-	-	1	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	53	69	80	116	158	231	292	314	209	74	17	-
Male	37	50	55	80	89	137	147	130	69	17	2	-
Female	16	19	25	36	69	94	145	184	140	57	15	-
White, both sexes	32	47	52	96	140	200	263	286	187	71	13	-
Male	26	36	35	67	78	123	137	119	65	17	2	-
Female	6	11	17	29	62	77	126	167	122	54	11	-
Black, both sexes	21	20	21	16	16	27	22	26	19	2	4	-
Male	11	13	14	10	11	14	7	10	4	-	-	-
Female	10	7	7	6	5	13	15	16	15	2	4	-
American Indian 1/, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	6	4	2	4	7	2	3	1	-	-
Male	-	-	6	3	-	-	3	1	-	-	-	-
Female	-	1	-	1	2	4	4	1	3	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
North Carolina												
All races, both sexes	28	37	48	73	109	124	156	150	102	40	11	-
Male	19	25	37	49	54	65	81	63	36	5	3	-
Female	9	12	11	24	55	59	75	87	66	35	8	-
White, both sexes	20	21	32	57	78	108	136	115	87	33	10	-
Male	14	15	24	42	40	59	75	49	31	4	3	-
Female	6	6	8	15	38	49	61	66	56	29	7	-
Black, both sexes	8	16	14	14	30	15	19	33	15	7	1	-
Male	5	10	11	6	13	5	6	13	5	1	-	-
Female	3	6	3	8	17	10	13	20	10	6	1	-
American Indian 1/, both sexes	-	-	-	2	1	1	-	1	-	-	-	-
Male	-	-	-	1	1	1	-	1	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	-	-	1	1	-	-	-	-
Male	-	-	2	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
North Dakota												
All races, both sexes	2	1	2	5	7	11	10	16	7	4	2	-
Male	2	1	2	2	5	6	7	6	-	2	-	-
Female	-	-	-	3	2	5	3	10	7	2	2	-
White, both sexes	2	1	-	4	7	11	10	16	7	4	2	-
Male	2	1	-	2	5	6	7	6	-	2	-	-
Female	-	-	-	2	2	5	3	10	7	2	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	2	1	-	-	-	-	-	-	-	-
Male	-	-	2	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Ohio												
All races, both sexes	36	41	81	93	115	195	282	266	170	70	11	-
Male	23	30	54	65	58	103	129	106	53	17	1	-
Female	13	11	27	28	57	92	153	160	117	53	10	-
White, both sexes	30	28	73	77	100	175	257	255	160	65	11	-
Male	17	20	47	53	51	89	119	102	51	16	1	-
Female	13	8	26	24	49	86	138	153	109	49	10	-
Black, both sexes	6	13	8	15	15	17	25	11	10	4	-	-
Male	6	10	7	11	7	12	10	4	2	-	-	-
Female	-	3	1	4	8	5	15	7	8	4	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	3	-	-	-	-	-	-
Male	-	-	-	1	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Oklahoma												
All races, both sexes	13	10	24	34	40	46	73	69	42	19	3	-
Male	9	8	16	26	25	19	32	25	9	4	1	-
Female	4	2	8	8	15	27	41	44	33	15	2	-
White, both sexes	9	9	19	29	36	43	66	68	40	16	3	-
Male	8	7	12	21	22	19	28	24	8	3	1	-
Female	1	2	7	8	14	24	38	44	32	13	2	-
Black, both sexes	3	-	4	-	1	1	1	1	-	1	-	-
Male	1	-	4	-	1	-	1	1	-	1	-	-
Female	2	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	1	5	3	-	6	-	2	2	-	-
Male	-	1	-	5	2	-	3	-	1	-	-	-
Female	1	-	1	-	1	-	3	-	1	2	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Oregon												
All races, both sexes	8	14	22	31	34	48	71	69	56	15	3	-
Male	5	9	17	20	21	29	40	20	27	4	1	-
Female	3	5	5	11	13	19	31	49	29	11	2	-
White, both sexes	7	13	21	27	33	48	68	66	56	15	3	-
Male	5	9	16	17	21	29	38	18	27	4	1	-
Female	2	4	5	10	12	19	30	48	29	11	2	-
Black, both sexes	1	-	1	3	1	-	-	-	-	-	-	-
Male	-	-	1	3	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	3	2	-	-	-	-
Male	-	-	-	-	-	-	2	2	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
Pennsylvania												
All races, both sexes	27	29	66	69	153	209	284	318	229	90	20	-
Male	19	19	51	38	86	84	134	131	76	24	3	-
Female	8	10	15	31	67	125	150	187	153	66	17	-
White, both sexes	16	24	59	64	145	194	261	301	221	88	18	-
Male	10	17	46	35	81	78	123	125	76	23	2	-
Female	6	7	13	29	64	116	138	176	145	65	16	-
Black, both sexes	11	4	6	5	7	13	19	17	8	2	2	-
Male	9	2	4	3	5	5	8	6	-	1	1	-
Female	2	2	2	2	2	8	11	11	8	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	1	2	4	-	-	-	-	-
Male	-	-	1	-	-	1	3	-	-	-	-	-
Female	-	1	-	-	1	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Rhode Island												
All races, both sexes	1	6	3	7	13	12	33	29	20	11	1	-
Male	-	4	2	3	5	4	11	10	4	2	-	-
Female	1	2	1	4	8	8	22	19	16	9	1	-
White, both sexes	1	3	3	6	13	12	33	28	20	11	1	-
Male	-	2	2	3	5	4	11	10	4	2	-	-
Female	1	1	1	3	8	8	22	18	16	9	1	-
Black, both sexes	-	2	-	1	-	-	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	12	15	32	49	52	42	61	64	35	15	1	-
Male	8	11	18	31	37	26	29	25	9	3	-	-
Female	4	4	14	18	15	16	32	39	26	12	1	-
White, both sexes	6	8	22	29	41	38	45	53	28	12	-	-
Male	4	6	15	18	31	24	20	20	9	2	-	-
Female	2	2	7	11	10	14	25	33	19	10	-	-
Black, both sexes	5	7	10	20	11	4	16	11	7	3	1	-
Male	3	5	3	13	6	2	9	5	-	1	-	-
Female	2	2	7	7	5	2	7	6	7	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
South Dakota												
All races, both sexes	2	2	5	2	11	9	22	24	11	9	1	-
Male	-	1	3	1	5	4	13	14	4	3	1	-
Female	2	1	2	1	6	5	9	10	7	6	-	-
White, both sexes	2	-	4	2	9	9	21	23	10	8	1	-
Male	-	-	3	1	5	4	13	14	4	3	1	-
Female	2	-	1	1	4	5	8	9	6	5	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	2	-	-	1	1	1	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	-	1	1	1	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
Tennessee												
All races, both sexes	20	25	46	63	58	107	105	80	74	30	6	-
Male	16	21	32	34	34	52	49	31	17	4	-	-
Female	4	4	14	29	24	55	56	49	57	26	6	-
White, both sexes	16	17	39	50	50	95	95	71	62	25	4	-
Male	12	14	28	27	30	49	45	28	14	4	-	-
Female	4	3	11	23	20	46	50	43	48	21	4	-
Black, both sexes	4	8	7	12	8	11	8	9	11	5	2	-
Male	4	7	4	7	4	2	4	3	3	-	-	-
Female	-	1	3	5	4	9	4	6	8	5	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	2	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	2	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Texas												
All races, both sexes	41	66	89	135	161	221	268	241	162	72	17	-
Male	27	49	57	82	103	122	126	106	58	13	3	-
Female	14	17	32	53	58	99	142	135	104	59	14	-
White, both sexes	28	55	80	114	134	189	241	218	144	57	14	-
Male	16	43	52	69	85	106	117	92	57	9	2	-
Female	12	12	28	45	49	83	124	126	87	48	12	-
Black, both sexes	12	10	8	19	23	31	21	22	17	15	3	-
Male	10	5	5	11	15	16	7	13	1	4	1	-
Female	2	5	3	8	8	15	14	9	16	11	2	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	2	3	1	6	1	1	-	-	-
Male	1	1	-	2	2	-	2	1	-	-	-	-
Female	-	-	1	-	1	1	4	-	1	-	-	-
Utah												
All races, both sexes	6	13	3	11	14	16	24	34	27	6	2	-
Male	4	10	1	10	9	8	12	17	9	2	1	-
Female	2	3	2	1	5	8	12	17	18	4	1	-
White, both sexes	6	11	2	11	14	16	23	34	27	6	2	-
Male	4	8	1	10	9	8	12	17	9	2	1	-
Female	2	3	1	1	5	8	11	17	18	4	1	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Vermont												
All races, both sexes	3	2	5	11	10	7	15	13	9	8	1	-
Male	2	1	4	7	6	6	5	6	2	2	1	-
Female	1	1	1	4	4	1	10	7	7	6	-	-
White, both sexes	3	2	5	11	10	7	14	13	9	8	1	-
Male	2	1	4	7	6	6	5	6	2	2	1	-
Female	1	1	1	4	4	1	9	7	7	6	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	11	32	33	45	61	87	130	115	84	26	12	-
Male	6	17	27	31	34	39	64	43	27	4	-	-
Female	5	15	6	14	27	48	66	72	57	22	12	-
White, both sexes	5	21	25	38	46	63	103	94	66	17	11	-
Male	1	11	21	28	27	29	50	35	24	2	-	-
Female	4	10	4	10	19	34	53	59	42	15	11	-
Black, both sexes	6	11	8	7	12	24	24	19	16	9	1	-
Male	5	6	6	3	5	10	12	8	3	2	-	-
Female	1	5	2	4	7	14	12	11	13	7	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	3	-	3	2	2	-	-	-
Male	-	-	-	-	2	-	2	-	-	-	-	-
Female	-	-	-	-	1	-	1	2	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Washington												
All races, both sexes	12	26	37	56	79	100	103	140	89	37	11	-
Male	8	14	28	35	44	56	55	61	31	11	1	-
Female	4	12	9	21	35	44	48	79	58	26	10	-
White, both sexes	12	26	35	49	72	94	94	134	89	36	11	-
Male	8	14	27	31	42	53	51	58	31	11	1	-
Female	4	12	8	18	30	41	43	76	58	25	10	-
Black, both sexes	-	-	-	1	4	3	3	1	-	1	-	-
Male	-	-	-	-	2	1	1	-	-	-	-	-
Female	-	-	-	1	2	2	2	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	2	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	4	3	2	6	5	-	-	-	-
Male	-	-	1	3	-	2	3	3	-	-	-	-
Female	-	-	1	1	3	-	3	2	-	-	-	-
West Virginia												
All races, both sexes	5	13	18	22	24	37	49	40	29	13	2	-
Male	3	6	12	12	19	20	17	11	10	4	-	-
Female	2	7	6	10	5	17	32	29	19	9	2	-
White, both sexes	5	11	18	22	24	37	48	39	29	13	2	-
Male	3	6	12	12	19	20	16	11	10	4	-	-
Female	2	5	6	10	5	17	32	28	19	9	2	-
Black, both sexes	-	2	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	2	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Wisconsin												
All races, both sexes	10	24	25	39	44	87	114	137	88	37	11	-
Male	7	17	14	25	27	53	57	62	35	8	-	-
Female	3	7	11	14	17	34	57	75	53	29	11	-
White, both sexes	8	22	25	34	41	83	109	134	86	37	11	-
Male	5	16	14	23	25	50	55	61	35	8	-	-
Female	3	6	11	11	16	33	54	73	51	29	11	-
Black, both sexes	2	2	-	4	3	4	5	-	2	-	-	-
Male	2	1	-	1	2	3	2	-	-	-	-	-
Female	-	1	-	3	1	1	3	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	2	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Wyoming												
All races, both sexes	1	3	3	9	11	9	19	4	3	3	-	-
Male	1	2	1	9	6	4	10	2	-	2	-	-
Female	-	1	2	-	5	5	9	2	3	1	-	-
White, both sexes	1	3	2	9	11	9	19	4	3	3	-	-
Male	1	2	-	9	6	4	10	2	-	2	-	-
Female	-	1	2	-	5	5	9	2	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
United States												
All races, both sexes	8,232	1	-	-	-	2	-	1	-	11	16	43
Male	3,220	1	-	-	-	-	-	1	-	8	11	33
Female	5,012	-	-	-	-	2	-	-	-	3	5	10
White, both sexes	7,444	1	-	-	-	1	-	1	-	8	12	27
Male	2,898	1	-	-	-	-	-	1	-	6	8	21
Female	4,546	-	-	-	-	1	-	-	-	2	4	6
Black, both sexes	647	-	-	-	-	1	-	-	-	1	4	15
Male	259	-	-	-	-	-	-	-	-	1	3	11
Female	388	-	-	-	-	1	-	-	-	-	1	4
American Indian 1/, both sexes	30	-	-	-	-	-	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	15	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	111	-	-	-	-	-	-	-	-	2	-	-
Male	48	-	-	-	-	-	-	-	-	1	-	-
Female	63	-	-	-	-	-	-	-	-	1	-	-
Alabama												
All races, both sexes	199	-	-	-	-	-	-	-	-	1	2	8
Male	78	-	-	-	-	-	-	-	-	-	2	4
Female	121	-	-	-	-	-	-	-	-	1	-	4
White, both sexes	155	-	-	-	-	-	-	-	-	1	2	5
Male	59	-	-	-	-	-	-	-	-	-	2	2
Female	96	-	-	-	-	-	-	-	-	1	-	3
Black, both sexes	44	-	-	-	-	-	-	-	-	-	-	3
Male	19	-	-	-	-	-	-	-	-	-	-	2
Female	25	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Alaska												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	195	-	-	-	-	-	-	-	-	-	-	1
Male	94	-	-	-	-	-	-	-	-	-	-	1
Female	101	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	185	-	-	-	-	-	-	-	-	-	-	1
Male	91	-	-	-	-	-	-	-	-	-	-	1
Female	94	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Arkansas												
All races, both sexes	33	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	29	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	956	-	-	-	-	-	-	1	-	1	1	2
Male	380	-	-	-	-	-	-	1	-	1	-	2
Female	576	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	838	-	-	-	-	-	-	1	-	-	1	2
Male	330	-	-	-	-	-	-	1	-	-	-	2
Female	508	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	49	-	-	-	-	-	-	-	-	1	-	-
Male	23	-	-	-	-	-	-	-	-	1	-	-
Female	26	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	65	-	-	-	-	-	-	-	-	-	-	-
Male	25	-	-	-	-	-	-	-	-	-	-	-
Female	40	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Colorado												
All races, both sexes	242	-	-	-	-	-	-	-	-	-	1	-
Male	77	-	-	-	-	-	-	-	-	-	1	-
Female	165	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	233	-	-	-	-	-	-	-	-	-	1	-
Male	74	-	-	-	-	-	-	-	-	-	1	-
Female	159	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	97	-	-	-	-	-	-	-	-	-	-	2
Male	27	-	-	-	-	-	-	-	-	-	-	-
Female	70	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	93	-	-	-	-	-	-	-	-	-	-	1
Male	25	-	-	-	-	-	-	-	-	-	-	-
Female	68	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Delaware												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Florida												
All races, both sexes	655	-	-	-	-	-	-	-	-	-	3	3
Male	276	-	-	-	-	-	-	-	-	-	3	1
Female	379	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	617	-	-	-	-	-	-	-	-	-	2	3
Male	265	-	-	-	-	-	-	-	-	-	2	1
Female	352	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	37	-	-	-	-	-	-	-	-	-	1	-
Male	10	-	-	-	-	-	-	-	-	-	1	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	152	1	-	-	-	-	-	-	-	1	-	2
Male	68	1	-	-	-	-	-	-	-	1	-	2
Female	84	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	126	1	-	-	-	-	-	-	-	1	-	1
Male	56	1	-	-	-	-	-	-	-	1	-	1
Female	70	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	26	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Hawaii												
All races, both sexes	15	-	-	-	-	-	-	-	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	1	-	-
Idaho												
All races, both sexes	30	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	30	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Illinois												
All races, both sexes	307	-	-	-	-	-	-	-	-	-	-	1
Male	103	-	-	-	-	-	-	-	-	-	-	1
Female	204	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	269	-	-	-	-	-	-	-	-	-	-	1
Male	90	-	-	-	-	-	-	-	-	-	-	1
Female	179	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	36	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	266	-	-	-	-	-	-	-	-	-	-	-
Male	82	-	-	-	-	-	-	-	-	-	-	-
Female	184	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	249	-	-	-	-	-	-	-	-	-	-	-
Male	77	-	-	-	-	-	-	-	-	-	-	-
Female	172	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Iowa												
All races, both sexes	176	-	-	-	-	-	-	-	-	-	1	-
Male	69	-	-	-	-	-	-	-	-	-	1	-
Female	107	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	174	-	-	-	-	-	-	-	-	-	1	-
Male	68	-	-	-	-	-	-	-	-	-	1	-
Female	106	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	353	-	-	-	-	-	-	-	-	1	-	1
Male	128	-	-	-	-	-	-	-	-	1	-	1
Female	225	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	337	-	-	-	-	-	-	-	-	1	-	1
Male	124	-	-	-	-	-	-	-	-	1	-	1
Female	213	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Kentucky												
All races, both sexes	106	-	-	-	-	-	-	-	-	-	1	1
Male	38	-	-	-	-	-	-	-	-	-	-	1
Female	68	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	102	-	-	-	-	-	-	-	-	-	1	1
Male	37	-	-	-	-	-	-	-	-	-	-	1
Female	65	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	147	-	-	-	-	-	-	-	-	-	1	1
Male	55	-	-	-	-	-	-	-	-	-	-	1
Female	92	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	116	-	-	-	-	-	-	-	-	-	-	-
Male	40	-	-	-	-	-	-	-	-	-	-	-
Female	76	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	31	-	-	-	-	-	-	-	-	-	1	1
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Maine												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	20	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	74	-	-	-	-	2	-	-	-	-	1	-
Male	25	-	-	-	-	-	-	-	-	-	1	-
Female	49	-	-	-	-	2	-	-	-	-	-	-
White, both sexes	62	-	-	-	-	1	-	-	-	-	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	42	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	11	-	-	-	-	1	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Massachusetts												
All races, both sexes	229	-	-	-	-	-	-	-	-	-	-	2
Male	93	-	-	-	-	-	-	-	-	-	-	2
Female	136	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	223	-	-	-	-	-	-	-	-	-	-	2
Male	91	-	-	-	-	-	-	-	-	-	-	2
Female	132	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	372	-	-	-	-	-	-	-	-	1	-	3
Male	142	-	-	-	-	-	-	-	-	-	-	3
Female	230	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	329	-	-	-	-	-	-	-	-	1	-	-
Male	124	-	-	-	-	-	-	-	-	-	-	-
Female	205	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	38	-	-	-	-	-	-	-	-	-	-	3
Male	15	-	-	-	-	-	-	-	-	-	-	3
Female	23	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Minnesota												
All races, both sexes	73	-	-	-	-	-	-	-	-	-	-	1
Male	40	-	-	-	-	-	-	-	-	-	-	1
Female	33	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	72	-	-	-	-	-	-	-	-	-	-	-
Male	39	-	-	-	-	-	-	-	-	-	-	-
Female	33	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Missouri												
All races, both sexes	247	-	-	-	-	-	-	-	-	1	-	1
Male	74	-	-	-	-	-	-	-	-	1	-	1
Female	173	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	211	-	-	-	-	-	-	-	-	1	-	1
Male	65	-	-	-	-	-	-	-	-	1	-	1
Female	146	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	36	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Nebraska												
All races, both sexes	64	-	-	-	-	-	-	-	-	-	-	-
Male	27	-	-	-	-	-	-	-	-	-	-	-
Female	37	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	63	-	-	-	-	-	-	-	-	-	-	-
Male	26	-	-	-	-	-	-	-	-	-	-	-
Female	37	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	120	-	-	-	-	-	-	-	-	-	-	1
Male	53	-	-	-	-	-	-	-	-	-	-	1
Female	67	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	117	-	-	-	-	-	-	-	-	-	-	-
Male	51	-	-	-	-	-	-	-	-	-	-	-
Female	66	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
New Hampshire												
All races, both sexes	27	-	-	-	-	-	-	-	-	-	-	1
Male	17	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	27	-	-	-	-	-	-	-	-	-	-	1
Male	17	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	178	-	-	-	-	-	-	-	-	-	-	-
Male	60	-	-	-	-	-	-	-	-	-	-	-
Female	118	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	167	-	-	-	-	-	-	-	-	-	-	-
Male	56	-	-	-	-	-	-	-	-	-	-	-
Female	111	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
New Mexico												
All races, both sexes	32	-	-	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	31	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	458	-	-	-	-	-	-	-	-	-	2	1
Male	220	-	-	-	-	-	-	-	-	-	1	1
Female	238	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	373	-	-	-	-	-	-	-	-	-	1	1
Male	180	-	-	-	-	-	-	-	-	-	-	1
Female	193	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	74	-	-	-	-	-	-	-	-	-	1	-
Male	35	-	-	-	-	-	-	-	-	-	1	-
Female	39	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
North Carolina												
All races, both sexes	232	-	-	-	-	-	-	-	-	-	-	1
Male	95	-	-	-	-	-	-	-	-	-	-	-
Female	137	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	172	-	-	-	-	-	-	-	-	-	-	-
Male	69	-	-	-	-	-	-	-	-	-	-	-
Female	103	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	59	-	-	-	-	-	-	-	-	-	-	1
Male	25	-	-	-	-	-	-	-	-	-	-	-
Female	34	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Ohio												
All races, both sexes	328	-	-	-	-	-	-	-	-	1	-	3
Male	134	-	-	-	-	-	-	-	-	1	-	3
Female	194	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	309	-	-	-	-	-	-	-	-	1	-	3
Male	126	-	-	-	-	-	-	-	-	1	-	3
Female	183	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	124	-	-	-	-	-	-	-	-	-	-	-
Male	40	-	-	-	-	-	-	-	-	-	-	-
Female	84	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	116	-	-	-	-	-	-	-	-	-	-	-
Male	36	-	-	-	-	-	-	-	-	-	-	-
Female	80	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Oregon												
All races, both sexes	116	-	-	-	-	-	-	-	-	-	-	-
Male	59	-	-	-	-	-	-	-	-	-	-	-
Female	57	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	116	-	-	-	-	-	-	-	-	-	-	-
Male	59	-	-	-	-	-	-	-	-	-	-	-
Female	57	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	333	-	-	-	-	-	-	-	-	1	-	2
Male	111	-	-	-	-	-	-	-	-	1	-	2
Female	222	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	321	-	-	-	-	-	-	-	-	1	-	1
Male	108	-	-	-	-	-	-	-	-	1	-	1
Female	213	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Rhode Island												
All races, both sexes	51	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	39	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	51	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	39	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	35	-	-	-	-	-	-	-	-	-	1	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	29	-	-	-	-	-	-	-	-	-	1	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
South Dakota												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	137	-	-	-	-	-	-	-	-	1	-	-
Male	47	-	-	-	-	-	-	-	-	1	-	-
Female	90	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	121	-	-	-	-	-	-	-	-	1	-	-
Male	44	-	-	-	-	-	-	-	-	1	-	-
Female	77	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Texas												
All races, both sexes	331	-	-	-	-	-	-	-	-	-	-	3
Male	133	-	-	-	-	-	-	-	-	-	-	2
Female	198	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	296	-	-	-	-	-	-	-	-	-	-	1
Male	119	-	-	-	-	-	-	-	-	-	-	1
Female	177	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	33	-	-	-	-	-	-	-	-	-	-	2
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	21	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	42	-	-	-	-	-	-	-	-	-	1	-
Male	22	-	-	-	-	-	-	-	-	-	1	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	41	-	-	-	-	-	-	-	-	-	1	-
Male	21	-	-	-	-	-	-	-	-	-	1	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Vermont												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	175	-	-	-	-	-	-	-	-	-	-	-
Male	64	-	-	-	-	-	-	-	-	-	-	-
Female	111	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	142	-	-	-	-	-	-	-	-	-	-	-
Male	49	-	-	-	-	-	-	-	-	-	-	-
Female	93	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	31	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Washington												
All races, both sexes	192	-	-	-	-	-	-	-	-	1	-	1
Male	81	-	-	-	-	-	-	-	-	1	-	1
Female	111	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	179	-	-	-	-	-	-	-	-	-	-	-
Male	71	-	-	-	-	-	-	-	-	-	-	-
Female	108	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	81	-	-	-	-	-	-	-	-	-	-	-
Male	31	-	-	-	-	-	-	-	-	-	-	-
Female	50	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	81	-	-	-	-	-	-	-	-	-	-	-
Male	31	-	-	-	-	-	-	-	-	-	-	-
Female	50	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Wisconsin												
All races, both sexes	104	-	-	-	-	-	-	-	-	-	1	-
Male	35	-	-	-	-	-	-	-	-	-	1	-
Female	69	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	101	-	-	-	-	-	-	-	-	-	1	-
Male	34	-	-	-	-	-	-	-	-	-	1	-
Female	67	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
United States												
All races, both sexes	91	153	197	303	526	895	1,403	1,856	1,608	848	278	-
Male	62	101	142	188	286	426	601	674	483	165	38	-
Female	29	52	55	115	240	469	802	1,182	1,125	683	240	-
White, both sexes	75	127	158	260	446	796	1,279	1,722	1,492	785	254	-
Male	51	83	109	160	246	389	557	622	455	154	35	-
Female	24	44	49	100	200	407	722	1,100	1,037	631	219	-
Black, both sexes	15	21	33	33	67	78	99	112	92	55	21	-
Male	11	15	27	20	34	30	33	41	22	9	2	-
Female	4	6	6	13	33	48	66	71	70	46	19	-
American Indian 1/, both sexes	-	4	-	4	5	4	3	3	4	2	-	-
Male	-	3	-	4	3	1	-	2	-	1	-	-
Female	-	1	-	-	2	3	3	1	4	1	-	-
Asian or Pacific Islander 2/, both sexes	1	1	6	6	8	17	22	19	20	6	3	-
Male	-	-	6	4	3	6	11	9	6	1	1	-
Female	1	1	-	2	5	11	11	10	14	5	2	-
Alabama												
All races, both sexes	7	8	9	10	13	25	28	42	27	15	4	-
Male	3	6	7	7	6	8	10	15	8	2	-	-
Female	4	2	2	3	7	17	18	27	19	13	4	-
White, both sexes	4	8	9	10	7	19	21	34	21	11	3	-
Male	-	6	7	7	4	6	7	12	5	1	-	-
Female	4	2	2	3	3	13	14	22	16	10	3	-
Black, both sexes	3	-	-	-	6	6	7	8	6	4	1	-
Male	3	-	-	-	2	2	3	3	3	1	-	-
Female	-	-	-	-	4	4	4	5	3	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Alaska												
All races, both sexes	-	1	1	-	1	-	1	1	1	-	-	-
Male	-	1	1	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
White, both sexes	-	1	1	-	-	-	1	1	1	-	-	-
Male	-	1	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	5	5	4	7	11	20	32	50	38	18	4	-
Male	2	5	2	4	8	11	13	18	19	8	3	-
Female	3	-	2	3	3	9	19	32	19	10	1	-
White, both sexes	5	4	4	7	10	18	31	49	34	18	4	-
Male	2	4	2	4	8	11	13	17	18	8	3	-
Female	3	-	2	3	2	7	18	32	16	10	1	-
Black, both sexes	-	-	-	-	-	-	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-
American Indian 1/, both sexes	-	1	-	-	1	1	1	-	1	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Arkansas												
All races, both sexes	-	-	-	-	6	3	6	5	8	4	1	-
Male	-	-	-	-	2	-	3	2	2	1	-	-
Female	-	-	-	-	4	3	3	3	6	3	1	-
White, both sexes	-	-	-	-	6	3	3	4	8	4	1	-
Male	-	-	-	-	2	-	2	1	2	1	-	-
Female	-	-	-	-	4	3	1	3	6	3	1	-
Black, both sexes	-	-	-	-	-	-	3	1	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	8	15	19	31	52	104	173	210	198	96	45	-
Male	6	8	15	23	29	50	69	77	67	21	11	-
Female	2	7	4	8	23	54	104	133	131	75	34	-
White, both sexes	4	13	14	25	43	86	152	190	179	89	39	-
Male	3	7	10	19	26	42	60	70	60	20	10	-
Female	1	6	4	6	17	44	92	120	119	69	29	-
Black, both sexes	3	1	3	3	3	8	7	11	4	2	3	-
Male	3	1	3	1	1	5	2	4	2	-	-	-
Female	-	-	-	2	2	3	5	7	2	2	3	-
American Indian 1/, both sexes	-	-	-	1	1	1	-	1	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	2	2	5	9	14	8	15	5	3	-
Male	-	-	2	2	1	3	7	3	5	1	1	-
Female	1	1	-	-	4	6	7	5	10	4	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Colorado												
All races, both sexes	-	2	3	6	15	15	46	55	55	34	10	-
Male	-	2	2	3	5	8	15	18	13	8	2	-
Female	-	-	1	3	10	7	31	37	42	26	8	-
White, both sexes	-	2	3	6	13	15	44	53	54	33	9	-
Male	-	2	2	3	4	8	15	17	13	8	1	-
Female	-	-	1	3	9	7	29	36	41	25	8	-
Black, both sexes	-	-	-	-	1	-	1	1	-	1	1	-
Male	-	-	-	-	1	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	1	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	1	2	3	12	16	17	25	12	7	-
Male	-	-	1	1	-	5	5	5	9	1	-	-
Female	-	-	-	1	3	7	11	12	16	11	7	-
White, both sexes	-	-	1	2	3	11	16	16	25	11	7	-
Male	-	-	1	1	-	4	5	5	9	-	-	-
Female	-	-	-	1	3	7	11	11	16	11	7	-
Black, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Delaware												
All races, both sexes	-	-	-	1	-	-	2	-	2	-	1	-
Male	-	-	-	-	-	-	2	-	2	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	1	-
White, both sexes	-	-	-	1	-	-	2	-	2	-	1	-
Male	-	-	-	-	-	-	2	-	2	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	1	2	6	1	1	1	-	1	-	-
Male	-	-	1	1	2	1	-	1	-	-	-	-
Female	-	-	-	1	4	-	1	-	-	1	-	-
White, both sexes	-	-	-	1	3	-	1	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	1	2	-	1	-	-	-	-	-
Black, both sexes	-	-	1	1	3	1	-	-	-	1	-	-
Male	-	-	1	1	1	1	-	-	-	-	-	-
Female	-	-	-	-	2	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Florida												
All races, both sexes	8	14	8	19	46	72	124	131	130	75	22	-
Male	5	11	4	10	34	35	57	59	38	18	1	-
Female	3	3	4	9	12	37	67	72	92	57	21	-
White, both sexes	8	14	8	18	44	65	120	125	124	65	21	-
Male	5	11	4	9	34	32	56	57	37	16	1	-
Female	3	3	4	9	10	33	64	68	87	49	20	-
Black, both sexes	-	-	-	1	2	7	4	6	6	9	1	-
Male	-	-	-	1	-	3	1	2	1	1	-	-
Female	-	-	-	-	2	4	3	4	5	8	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	3	7	5	7	15	22	22	29	21	13	4	-
Male	2	4	4	6	7	11	11	12	7	-	-	-
Female	1	3	1	1	8	11	11	17	14	13	4	-
White, both sexes	3	4	5	6	12	20	21	25	15	9	3	-
Male	2	1	4	5	5	11	10	10	5	-	-	-
Female	1	3	1	1	7	9	11	15	10	9	3	-
Black, both sexes	-	3	-	1	3	2	1	4	6	4	1	-
Male	-	3	-	1	2	-	1	2	2	-	-	-
Female	-	-	-	-	1	2	-	2	4	4	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Hawaii												
All races, both sexes	-	-	2	-	-	4	3	2	2	1	-	-
Male	-	-	1	-	-	1	1	1	1	1	-	-
Female	-	-	1	-	-	3	2	1	1	-	-	-
White, both sexes	-	-	1	-	-	1	-	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	1	1	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	3	3	2	1	-	-	-
Male	-	-	1	-	-	1	1	1	-	-	-	-
Female	-	-	-	-	-	2	2	1	1	-	-	-
Idaho												
All races, both sexes	-	2	1	1	2	5	1	8	5	5	-	-
Male	-	-	-	1	2	4	1	2	1	1	-	-
Female	-	2	1	-	-	1	-	6	4	4	-	-
White, both sexes	-	2	1	1	2	5	1	8	5	5	-	-
Male	-	-	-	1	2	4	1	2	1	1	-	-
Female	-	2	1	-	-	1	-	6	4	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Illinois												
All races, both sexes	-	3	7	13	16	34	60	52	64	43	14	-
Male	-	1	5	7	9	14	25	19	14	7	1	-
Female	-	2	2	6	7	20	35	33	50	36	13	-
White, both sexes	-	2	2	11	11	32	51	46	60	40	13	-
Male	-	1	1	5	6	14	24	17	13	7	1	-
Female	-	1	1	6	5	18	27	29	47	33	12	-
Black, both sexes	-	1	5	2	5	1	9	5	4	3	1	-
Male	-	-	4	2	3	-	1	1	1	-	-	-
Female	-	1	1	-	2	1	8	4	3	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1	2	4	5	14	33	40	65	62	33	7	-
Male	-	-	1	1	8	14	16	22	13	6	1	-
Female	1	2	3	4	6	19	24	43	49	27	6	-
White, both sexes	1	2	4	5	13	28	37	61	61	30	7	-
Male	-	-	1	1	7	12	15	21	13	6	1	-
Female	1	2	3	4	6	16	22	40	48	24	6	-
Black, both sexes	-	-	-	-	1	5	3	4	1	3	-	-
Male	-	-	-	-	1	2	1	1	-	-	-	-
Female	-	-	-	-	-	3	2	3	1	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Iowa												
All races, both sexes	5	2	7	4	15	16	23	37	37	23	6	-
Male	5	2	5	4	5	8	10	13	15	1	-	-
Female	-	-	2	-	10	8	13	24	22	22	6	-
White, both sexes	4	2	7	4	15	16	23	36	37	23	6	-
Male	4	2	5	4	5	8	10	13	15	1	-	-
Female	-	-	2	-	10	8	13	23	22	22	6	-
Black, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	3	3	9	10	25	40	56	93	73	34	5	-
Male	2	3	6	3	14	14	24	30	25	5	-	-
Female	1	-	3	7	11	26	32	63	48	29	5	-
White, both sexes	3	3	8	9	22	39	55	89	69	33	5	-
Male	2	3	6	3	11	14	24	30	24	5	-	-
Female	1	-	2	6	11	25	31	59	45	28	5	-
Black, both sexes	-	-	1	1	3	1	1	3	3	1	-	-
Male	-	-	-	-	3	-	-	-	1	-	-	-
Female	-	-	1	1	-	1	1	3	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Kentucky												
All races, both sexes	4	-	3	7	5	12	17	30	13	11	2	-
Male	2	-	3	3	3	4	5	14	2	1	-	-
Female	2	-	-	4	2	8	12	16	11	10	2	-
White, both sexes	4	-	3	7	3	12	17	30	12	10	2	-
Male	2	-	3	3	2	4	5	14	2	1	-	-
Female	2	-	-	4	1	8	12	16	10	9	2	-
Black, both sexes	-	-	-	-	2	-	-	-	1	1	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	1	1	6	11	20	20	45	24	11	5	-
Male	1	1	1	2	7	9	8	13	6	5	1	-
Female	-	-	-	4	4	11	12	32	18	6	4	-
White, both sexes	-	1	-	3	8	16	15	39	21	9	4	-
Male	-	1	-	1	5	8	4	12	4	4	1	-
Female	-	-	-	2	3	8	11	27	17	5	3	-
Black, both sexes	1	-	1	3	3	4	5	6	3	2	1	-
Male	1	-	1	1	2	1	4	1	2	1	-	-
Female	-	-	-	2	1	3	1	5	1	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Maine												
All races, both sexes	-	-	-	2	3	-	-	5	4	4	1	-
Male	-	-	-	1	1	-	-	3	2	1	1	-
Female	-	-	-	1	2	-	-	2	2	3	-	-
White, both sexes	-	-	-	2	3	-	-	5	4	4	1	-
Male	-	-	-	1	1	-	-	3	2	1	1	-
Female	-	-	-	1	2	-	-	2	2	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	2	-	2	5	4	12	19	14	10	3	-
Male	-	-	-	1	1	2	5	9	5	1	-	-
Female	-	2	-	1	4	2	7	10	9	9	3	-
White, both sexes	-	1	-	1	2	4	11	17	12	10	3	-
Male	-	-	-	1	-	2	5	7	4	1	-	-
Female	-	1	-	-	2	2	6	10	8	9	3	-
Black, both sexes	-	1	-	-	3	-	1	2	2	-	-	-
Male	-	-	-	-	1	-	-	2	1	-	-	-
Female	-	1	-	-	2	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Massachusetts												
All races, both sexes	2	4	5	9	10	24	28	62	43	29	11	-
Male	2	3	5	6	7	16	13	21	12	6	-	-
Female	-	1	-	3	3	8	15	41	31	23	11	-
White, both sexes	2	4	4	9	10	22	28	61	41	29	11	-
Male	2	3	4	6	7	15	13	21	12	6	-	-
Female	-	1	-	3	3	7	15	40	29	23	11	-
Black, both sexes	-	-	1	-	-	1	-	1	2	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Michigan												
All races, both sexes	3	11	11	19	24	40	67	83	75	24	11	-
Male	3	7	9	12	11	20	25	31	18	1	2	-
Female	-	4	2	7	13	20	42	52	57	23	9	-
White, both sexes	3	6	9	17	21	35	62	77	65	23	10	-
Male	3	3	7	11	11	17	24	28	17	1	2	-
Female	-	3	2	6	10	18	38	49	48	22	8	-
Black, both sexes	-	4	2	2	3	3	5	5	9	1	1	-
Male	-	4	2	1	-	1	1	2	1	-	-	-
Female	-	-	-	1	3	2	4	3	8	1	1	-
American Indian 1/, both sexes	-	1	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Minnesota												
All races, both sexes	1	2	2	-	2	13	13	17	9	9	4	-
Male	1	1	1	-	1	9	10	10	5	-	1	-
Female	-	1	1	-	1	4	3	7	4	9	3	-
White, both sexes	1	2	2	-	2	13	13	17	9	9	4	-
Male	1	1	1	-	1	9	10	10	5	-	1	-
Female	-	1	1	-	1	4	3	7	4	9	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	2	1	2	5	1	7	3	4	-	-
Male	-	-	2	-	1	3	-	4	1	-	-	-
Female	-	-	-	1	1	2	1	3	2	4	-	-
White, both sexes	-	-	2	1	2	4	1	7	3	2	-	-
Male	-	-	2	-	1	2	-	4	1	-	-	-
Female	-	-	-	1	1	2	1	3	2	2	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	2	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Missouri												
All races, both sexes	2	1	2	6	16	26	42	65	49	26	10	-
Male	-	1	2	4	11	9	14	20	9	1	1	-
Female	2	-	-	2	5	17	28	45	40	25	9	-
White, both sexes	2	1	2	5	14	20	32	55	45	25	8	-
Male	-	1	2	3	10	8	11	17	9	1	1	-
Female	2	-	-	2	4	12	21	38	36	24	7	-
Black, both sexes	-	-	-	1	2	6	10	10	4	1	2	-
Male	-	-	-	1	1	1	3	3	-	-	-	-
Female	-	-	-	-	1	5	7	7	4	1	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	2	-	4	-	3	4	3	3	2	2	1	-
Male	2	-	2	-	1	4	-	2	-	1	-	-
Female	-	-	2	-	2	-	3	1	2	1	1	-
White, both sexes	2	-	4	-	3	4	3	3	2	2	1	-
Male	2	-	2	-	1	4	-	2	-	1	-	-
Female	-	-	2	-	2	-	3	1	2	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Nebraska												
All races, both sexes	-	1	1	1	4	6	8	21	12	8	2	-
Male	-	1	-	1	2	5	4	6	3	4	1	-
Female	-	-	1	-	2	1	4	15	9	4	1	-
White, both sexes	-	1	1	-	4	6	8	21	12	8	2	-
Male	-	1	-	-	2	5	4	6	3	4	1	-
Female	-	-	1	-	2	1	4	15	9	4	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	4	9	5	15	28	27	21	7	3	-
Male	-	-	3	7	2	6	13	12	6	2	1	-
Female	-	-	1	2	3	9	15	15	15	5	2	-
White, both sexes	-	-	4	9	4	15	28	27	20	7	3	-
Male	-	-	3	7	2	6	13	12	5	2	1	-
Female	-	-	1	2	2	9	15	15	15	5	2	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
New Hampshire												
All races, both sexes	-	1	1	-	1	3	4	9	5	2	-	-
Male	-	1	1	-	1	2	2	6	3	-	-	-
Female	-	-	-	-	-	1	2	3	2	2	-	-
White, both sexes	-	1	1	-	1	3	4	9	5	2	-	-
Male	-	1	1	-	1	2	2	6	3	-	-	-
Female	-	-	-	-	-	1	2	3	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	4	4	2	6	13	19	26	33	40	23	8	-
Male	4	3	2	2	7	6	12	8	9	6	1	-
Female	-	1	-	4	6	13	14	25	31	17	7	-
White, both sexes	4	4	2	6	13	18	25	29	37	22	7	-
Male	4	3	2	2	7	5	12	6	9	5	1	-
Female	-	1	-	4	6	13	13	23	28	17	6	-
Black, both sexes	-	-	-	-	-	1	1	4	3	1	1	-
Male	-	-	-	-	-	1	-	2	-	1	-	-
Female	-	-	-	-	-	-	1	2	3	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
New Mexico												
All races, both sexes	-	1	1	1	2	5	3	6	11	1	1	-
Male	-	-	1	1	1	2	2	3	7	-	-	-
Female	-	1	-	-	1	3	1	3	4	1	1	-
White, both sexes	-	1	1	1	1	5	3	6	11	1	1	-
Male	-	-	1	1	-	2	2	3	7	-	-	-
Female	-	1	-	-	1	3	1	3	4	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	9	10	20	23	32	49	72	99	87	41	13	-
Male	6	8	16	17	20	25	43	44	28	9	2	-
Female	3	2	4	6	12	24	29	55	59	32	11	-
White, both sexes	5	3	11	17	27	39	61	86	73	39	10	-
Male	5	3	9	13	16	21	39	37	25	9	2	-
Female	-	-	2	4	11	18	22	49	48	30	8	-
Black, both sexes	4	7	7	4	5	9	9	12	12	1	3	-
Male	1	5	5	3	4	4	3	6	3	-	-	-
Female	3	2	2	1	1	5	6	6	9	1	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	2	-	1	2	1	2	1	-	-
Male	-	-	2	1	-	-	1	1	-	-	-	-
Female	-	-	-	1	-	1	1	-	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
North Carolina												
All races, both sexes	2	5	9	10	17	27	48	49	43	16	5	-
Male	1	3	7	7	8	10	25	20	12	1	1	-
Female	1	2	2	3	9	17	23	29	31	15	4	-
White, both sexes	-	3	2	8	8	20	42	37	36	12	4	-
Male	-	2	1	6	4	9	22	13	10	1	1	-
Female	-	1	1	2	4	11	20	24	26	11	3	-
Black, both sexes	2	2	6	2	9	7	6	12	7	4	1	-
Male	1	1	5	1	4	1	3	7	2	-	-	-
Female	1	1	1	1	5	6	3	5	5	4	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	1	-	-	1	-	1	1	7	2	2	2	-
Male	1	-	-	1	-	1	-	1	-	1	-	-
Female	-	-	-	-	-	-	1	6	2	1	2	-
White, both sexes	1	-	-	1	-	1	1	7	2	2	2	-
Male	1	-	-	1	-	1	-	1	-	1	-	-
Female	-	-	-	-	-	-	1	6	2	1	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Ohio												
All races, both sexes	4	7	12	8	22	38	62	72	58	33	8	-
Male	1	4	9	7	9	22	30	22	18	8	-	-
Female	3	3	3	1	13	16	32	50	40	25	8	-
White, both sexes	4	7	10	5	20	38	55	68	58	32	8	-
Male	1	4	7	4	8	22	28	22	18	8	-	-
Female	3	3	3	1	12	16	27	46	40	24	8	-
Black, both sexes	-	-	2	3	2	-	7	4	-	1	-	-
Male	-	-	2	3	1	-	2	-	-	-	-	-
Female	-	-	-	-	1	-	5	4	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2	2	3	5	6	8	27	30	25	13	3	-
Male	2	1	1	3	3	3	9	9	5	3	1	-
Female	-	1	2	2	3	5	18	21	20	10	2	-
White, both sexes	2	1	2	4	6	8	25	30	24	11	3	-
Male	2	-	-	2	3	3	9	9	5	2	1	-
Female	-	1	2	2	3	5	16	21	19	9	2	-
Black, both sexes	-	-	1	-	-	-	-	-	-	1	-	-
Male	-	-	1	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	1	-	-	2	-	1	1	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	1	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Oregon												
All races, both sexes	1	4	4	6	4	14	24	23	26	7	3	-
Male	1	3	3	4	4	8	14	6	14	1	1	-
Female	-	1	1	2	-	6	10	17	12	6	2	-
White, both sexes	1	4	4	6	4	14	24	23	26	7	3	-
Male	1	3	3	4	4	8	14	6	14	1	1	-
Female	-	1	1	2	-	6	10	17	12	6	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	6	2	4	5	22	34	62	81	73	30	11	-
Male	5	-	4	2	14	12	21	27	18	4	1	-
Female	1	2	-	3	8	22	41	54	55	26	10	-
White, both sexes	6	2	4	5	21	32	59	78	72	29	11	-
Male	5	-	4	2	13	12	21	27	18	3	1	-
Female	1	2	-	3	8	20	38	51	54	26	10	-
Black, both sexes	-	-	-	-	1	2	3	3	1	1	-	-
Male	-	-	-	-	1	-	-	-	-	1	-	-
Female	-	-	-	-	-	2	3	3	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Rhode Island												
All races, both sexes	-	-	-	-	-	1	11	13	15	10	1	-
Male	-	-	-	-	-	-	4	3	3	2	-	-
Female	-	-	-	-	-	1	7	10	12	8	1	-
White, both sexes	-	-	-	-	-	1	11	13	15	10	1	-
Male	-	-	-	-	-	-	4	3	3	2	-	-
Female	-	-	-	-	-	1	7	10	12	8	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	1	3	4	4	7	8	4	3	-	-
Male	-	-	-	2	2	2	4	5	1	-	-	-
Female	-	-	1	1	2	2	3	3	3	3	-	-
White, both sexes	-	-	-	2	2	4	6	8	3	3	-	-
Male	-	-	-	1	1	2	3	5	1	-	-	-
Female	-	-	-	1	1	2	3	3	2	3	-	-
Black, both sexes	-	-	1	1	2	-	1	-	1	-	-	-
Male	-	-	-	1	1	-	1	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
South Dakota												
All races, both sexes	-	-	-	-	1	1	1	7	2	2	1	-
Male	-	-	-	-	1	-	1	4	1	1	1	-
Female	-	-	-	-	-	1	-	3	1	1	-	-
White, both sexes	-	-	-	-	1	1	1	7	2	2	1	-
Male	-	-	-	-	1	-	1	4	1	1	1	-
Female	-	-	-	-	-	1	-	3	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	1	3	3	10	11	18	20	23	28	15	4	-
Male	1	3	2	5	5	8	6	8	5	3	-	-
Female	-	-	1	5	6	10	14	15	23	12	4	-
White, both sexes	1	3	3	7	9	14	20	23	26	12	2	-
Male	1	3	2	4	4	7	6	8	5	3	-	-
Female	-	-	1	3	5	7	14	15	21	9	2	-
Black, both sexes	-	-	-	3	2	4	-	-	2	3	2	-
Male	-	-	-	1	1	1	-	-	-	-	-	-
Female	-	-	-	2	1	3	-	-	2	3	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Texas												
All races, both sexes	3	7	10	17	30	36	59	70	52	35	9	-
Male	1	6	6	8	18	17	24	24	17	7	3	-
Female	2	1	4	9	12	19	35	46	35	28	6	-
White, both sexes	2	6	8	15	26	34	52	64	49	31	8	-
Male	-	6	4	7	16	17	22	22	17	5	2	-
Female	2	-	4	8	10	17	30	42	32	26	6	-
Black, both sexes	1	1	2	2	3	2	6	6	3	4	1	-
Male	1	-	2	1	1	-	1	2	-	2	1	-
Female	-	1	-	1	2	2	5	4	3	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	4	-	2	2	1	6	11	10	4	1	-
Male	-	4	-	2	1	1	2	6	4	1	-	-
Female	-	-	-	-	1	-	4	5	6	3	1	-
White, both sexes	-	3	-	2	2	1	6	11	10	4	1	-
Male	-	3	-	2	1	1	2	6	4	1	-	-
Female	-	-	-	-	1	-	4	5	6	3	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Vermont												
All races, both sexes	-	-	-	1	1	-	1	2	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	1	2	1	1	-	-
White, both sexes	-	-	-	1	1	-	1	2	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	1	2	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	6	5	9	7	14	37	44	33	13	7	-
Male	-	4	4	6	4	8	15	13	10	-	-	-
Female	-	2	1	3	3	6	22	31	23	13	7	-
White, both sexes	-	5	5	6	5	10	31	40	24	10	6	-
Male	-	3	4	5	2	5	11	11	8	-	-	-
Female	-	2	1	1	3	5	20	29	16	10	6	-
Black, both sexes	-	1	-	3	1	4	6	3	9	3	1	-
Male	-	1	-	1	1	3	4	2	2	-	-	-
Female	-	-	-	2	-	1	2	1	7	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Washington												
All races, both sexes	2	6	1	6	15	17	22	51	39	23	8	-
Male	2	1	-	4	7	10	14	17	16	8	-	-
Female	-	5	1	2	8	7	8	34	23	15	8	-
White, both sexes	2	6	1	4	13	16	19	48	39	23	8	-
Male	2	1	-	2	6	9	12	15	16	8	-	-
Female	-	5	1	2	7	7	7	33	23	15	8	-
Black, both sexes	-	-	-	-	2	-	2	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	1	3	-	-	-	-
Male	-	-	-	1	-	1	1	2	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
West Virginia												
All races, both sexes	-	3	4	3	3	16	15	13	13	9	2	-
Male	-	1	3	2	1	10	4	4	2	4	-	-
Female	-	2	1	1	2	6	11	9	11	5	2	-
White, both sexes	-	3	4	3	3	16	15	13	13	9	2	-
Male	-	1	3	2	1	10	4	4	2	4	-	-
Female	-	2	1	1	2	6	11	9	11	5	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Wisconsin												
All races, both sexes	1	2	1	5	3	12	17	20	22	12	8	-
Male	1	2	-	5	-	7	6	4	7	2	-	-
Female	-	-	1	-	3	5	11	16	15	10	8	-
White, both sexes	1	2	1	5	3	10	16	20	22	12	8	-
Male	1	2	-	5	-	6	6	4	7	2	-	-
Female	-	-	1	-	3	4	10	16	15	10	8	-
Black, both sexes	-	-	-	-	-	2	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	2	-	2	5	3	2	1	-	-
Male	-	-	-	2	-	1	3	1	-	1	-	-
Female	-	-	-	-	-	1	2	2	2	-	-	-
White, both sexes	-	-	-	2	-	2	5	3	2	1	-	-
Male	-	-	-	2	-	1	3	1	-	1	-	-
Female	-	-	-	-	-	1	2	2	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
United States												
All races, both sexes	12,986	1	-	-	7	12	26	29	70	140	224	304
Male	7,543	-	-	-	6	11	21	22	52	110	168	226
Female	5,443	1	-	-	1	1	5	7	18	30	56	78
White, both sexes	11,348	1	-	-	6	7	19	24	49	89	144	206
Male	6,610	-	-	-	5	6	15	18	36	72	107	157
Female	4,738	1	-	-	1	1	4	6	13	17	37	49
Black, both sexes	1,257	-	-	-	1	4	4	2	16	46	72	88
Male	705	-	-	-	1	4	4	2	12	33	54	61
Female	552	-	-	-	-	-	-	-	4	13	18	27
American Indian 1/, both sexes	60	-	-	-	-	1	1	1	1	2	3	3
Male	41	-	-	-	-	1	1	1	1	2	2	3
Female	19	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	321	-	-	-	-	-	2	2	4	3	5	7
Male	187	-	-	-	-	-	1	1	3	3	5	5
Female	134	-	-	-	-	-	1	1	1	-	-	2
Alabama												
All races, both sexes	215	-	-	-	-	-	-	3	-	4	7	6
Male	134	-	-	-	-	-	-	1	-	2	6	5
Female	81	-	-	-	-	-	-	2	-	2	1	1
White, both sexes	169	-	-	-	-	-	-	2	-	3	4	4
Male	107	-	-	-	-	-	-	1	-	2	3	4
Female	62	-	-	-	-	-	-	1	-	1	1	-
Black, both sexes	43	-	-	-	-	-	-	-	-	1	3	2
Male	25	-	-	-	-	-	-	-	-	-	3	1
Female	18	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Alaska												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	247	-	-	-	1	2	2	-	5	5	7	5
Male	153	-	-	-	-	2	2	-	5	5	7	3
Female	94	-	-	-	1	-	-	-	-	-	-	2
White, both sexes	232	-	-	-	1	1	1	-	3	3	6	3
Male	142	-	-	-	-	1	1	-	3	3	6	2
Female	90	-	-	-	1	-	-	-	-	-	-	1
Black, both sexes	9	-	-	-	-	-	-	-	2	1	1	1
Male	5	-	-	-	-	-	-	-	2	1	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	6	-	-	-	-	1	1	-	-	1	-	1
Male	6	-	-	-	-	1	1	-	-	1	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Arkansas												
All races, both sexes	144	-	-	-	-	-	-	-	1	3	2	1
Male	82	-	-	-	-	-	-	-	1	3	1	1
Female	62	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	128	-	-	-	-	-	-	-	-	2	2	-
Male	75	-	-	-	-	-	-	-	-	2	1	-
Female	53	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	13	-	-	-	-	-	-	-	1	1	-	1
Male	7	-	-	-	-	-	-	-	1	1	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1,246	-	-	-	-	-	5	4	10	15	27	35
Male	746	-	-	-	-	-	3	4	9	12	20	28
Female	500	-	-	-	-	-	2	-	1	3	7	7
White, both sexes	1,002	-	-	-	-	-	4	4	6	10	20	28
Male	589	-	-	-	-	-	2	4	5	8	13	21
Female	413	-	-	-	-	-	2	-	1	2	7	7
Black, both sexes	94	-	-	-	-	-	-	-	1	3	5	3
Male	55	-	-	-	-	-	-	-	1	2	5	3
Female	39	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	144	-	-	-	-	-	1	-	3	2	2	3
Male	96	-	-	-	-	-	1	-	3	2	2	3
Female	48	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Colorado												
All races, both sexes	199	-	-	-	-	-	-	-	3	2	2	5
Male	113	-	-	-	-	-	-	-	2	2	1	3
Female	86	-	-	-	-	-	-	-	1	-	1	2
White, both sexes	189	-	-	-	-	-	-	-	3	2	1	5
Male	110	-	-	-	-	-	-	-	2	2	1	3
Female	79	-	-	-	-	-	-	-	1	-	-	2
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	151	-	-	-	-	-	-	-	3	5	2	2
Male	80	-	-	-	-	-	-	-	2	4	2	2
Female	71	-	-	-	-	-	-	-	1	1	-	-
White, both sexes	132	-	-	-	-	-	-	-	2	3	-	2
Male	68	-	-	-	-	-	-	-	1	3	-	2
Female	64	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	15	-	-	-	-	-	-	-	1	2	1	-
Male	10	-	-	-	-	-	-	-	1	1	1	-
Female	5	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Delaware												
All races, both sexes	29	-	-	-	-	-	-	-	-	2	-	1
Male	15	-	-	-	-	-	-	-	-	2	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	23	-	-	-	-	-	-	-	-	2	-	1
Male	13	-	-	-	-	-	-	-	-	2	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	28	-	-	-	-	-	-	-	1	-	2	3
Male	15	-	-	-	-	-	-	-	1	-	2	2
Female	13	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	24	-	-	-	-	-	-	-	1	-	2	3
Male	15	-	-	-	-	-	-	-	1	-	2	2
Female	9	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Florida												
All races, both sexes	927	-	-	-	-	-	-	1	3	9	13	18
Male	562	-	-	-	-	-	-	1	2	6	11	14
Female	365	-	-	-	-	-	-	-	1	3	2	4
White, both sexes	845	-	-	-	-	-	-	1	1	5	8	12
Male	515	-	-	-	-	-	-	1	1	4	8	9
Female	330	-	-	-	-	-	-	-	-	1	-	3
Black, both sexes	78	-	-	-	-	-	-	-	2	4	4	6
Male	45	-	-	-	-	-	-	-	1	2	2	5
Female	33	-	-	-	-	-	-	-	1	2	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	257	-	-	-	2	-	-	-	2	7	7	12
Male	165	-	-	-	2	-	-	-	2	3	5	6
Female	92	-	-	-	-	-	-	-	-	4	2	6
White, both sexes	180	-	-	-	1	-	-	-	2	3	1	6
Male	117	-	-	-	1	-	-	-	2	-	1	3
Female	63	-	-	-	-	-	-	-	-	3	-	3
Black, both sexes	74	-	-	-	1	-	-	-	-	4	6	6
Male	46	-	-	-	1	-	-	-	-	3	4	3
Female	28	-	-	-	-	-	-	-	-	1	2	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Hawaii												
All races, both sexes	64	-	-	-	-	-	-	-	-	-	2	3
Male	35	-	-	-	-	-	-	-	-	-	2	3
Female	29	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	24	-	-	-	-	-	-	-	-	-	1	3
Male	11	-	-	-	-	-	-	-	-	-	1	3
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	38	-	-	-	-	-	-	-	-	-	1	-
Male	22	-	-	-	-	-	-	-	-	-	1	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	72	-	-	-	-	-	-	-	-	-	-	1
Male	43	-	-	-	-	-	-	-	-	-	-	1
Female	29	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	70	-	-	-	-	-	-	-	-	-	-	1
Male	42	-	-	-	-	-	-	-	-	-	-	1
Female	28	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Illinois												
All races, both sexes	578	-	-	-	2	-	-	-	1	6	11	15
Male	317	-	-	-	2	-	-	-	1	6	7	8
Female	261	-	-	-	-	-	-	-	-	-	4	7
White, both sexes	490	-	-	-	2	-	-	-	1	1	5	8
Male	269	-	-	-	2	-	-	-	1	1	4	5
Female	221	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	80	-	-	-	-	-	-	-	-	5	6	7
Male	41	-	-	-	-	-	-	-	-	5	3	3
Female	39	-	-	-	-	-	-	-	-	-	3	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	325	-	-	-	-	-	1	-	3	1	2	4
Male	195	-	-	-	-	-	1	-	2	1	1	4
Female	130	-	-	-	-	-	-	-	1	-	1	-
White, both sexes	301	-	-	-	-	-	-	-	3	1	1	3
Male	181	-	-	-	-	-	-	-	2	1	1	3
Female	120	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	22	-	-	-	-	-	1	-	-	-	1	1
Male	13	-	-	-	-	-	1	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Iowa												
All races, both sexes	148	-	-	-	-	-	-	-	-	1	3	5
Male	83	-	-	-	-	-	-	-	-	1	1	3
Female	65	-	-	-	-	-	-	-	-	-	2	2
White, both sexes	146	-	-	-	-	-	-	-	-	1	3	5
Male	81	-	-	-	-	-	-	-	-	1	1	3
Female	65	-	-	-	-	-	-	-	-	-	2	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	130	-	-	-	-	1	-	-	-	2	-	3
Male	84	-	-	-	-	1	-	-	-	2	-	3
Female	46	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	125	-	-	-	-	1	-	-	-	2	-	2
Male	82	-	-	-	-	1	-	-	-	2	-	2
Female	43	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Kentucky												
All races, both sexes	206	-	-	-	-	-	2	1	-	1	3	3
Male	127	-	-	-	-	-	2	1	-	1	2	3
Female	79	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	193	-	-	-	-	-	2	1	-	1	3	3
Male	118	-	-	-	-	-	2	1	-	1	2	3
Female	75	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	171	-	-	-	-	-	2	1	-	-	4	5
Male	92	-	-	-	-	-	2	1	-	-	3	2
Female	79	-	-	-	-	-	-	-	-	-	1	3
White, both sexes	130	-	-	-	-	-	1	1	-	-	1	2
Male	73	-	-	-	-	-	1	1	-	-	1	1
Female	57	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	41	-	-	-	-	-	1	-	-	-	3	3
Male	19	-	-	-	-	-	1	-	-	-	2	1
Female	22	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Maine												
All races, both sexes	78	-	-	-	-	-	-	1	-	2	-	-
Male	42	-	-	-	-	-	-	1	-	1	-	-
Female	36	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	76	-	-	-	-	-	-	1	-	2	-	-
Male	42	-	-	-	-	-	-	1	-	1	-	-
Female	34	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	228	-	-	-	-	-	-	-	2	5	6	7
Male	116	-	-	-	-	-	-	-	2	4	5	5
Female	112	-	-	-	-	-	-	-	-	1	1	2
White, both sexes	176	-	-	-	-	-	-	-	2	1	3	2
Male	90	-	-	-	-	-	-	-	2	1	3	2
Female	86	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	49	-	-	-	-	-	-	-	-	4	3	4
Male	24	-	-	-	-	-	-	-	-	3	2	2
Female	25	-	-	-	-	-	-	-	-	1	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Massachusetts												
All races, both sexes	294	-	-	-	-	-	-	-	-	3	2	8
Male	154	-	-	-	-	-	-	-	-	2	1	6
Female	140	-	-	-	-	-	-	-	-	1	1	2
White, both sexes	285	-	-	-	-	-	-	-	-	3	2	8
Male	148	-	-	-	-	-	-	-	-	2	1	6
Female	137	-	-	-	-	-	-	-	-	1	1	2
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	550	-	-	-	-	1	-	-	-	9	12	13
Male	325	-	-	-	-	1	-	-	-	6	9	10
Female	225	-	-	-	-	-	-	-	-	3	3	3
White, both sexes	475	-	-	-	-	-	-	-	-	3	5	6
Male	282	-	-	-	-	-	-	-	-	1	4	6
Female	193	-	-	-	-	-	-	-	-	2	1	-
Black, both sexes	69	-	-	-	-	1	-	-	-	6	7	7
Male	37	-	-	-	-	1	-	-	-	5	5	4
Female	32	-	-	-	-	-	-	-	-	1	2	3
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Minnesota												
All races, both sexes	255	-	-	-	-	-	2	-	2	1	3	7
Male	143	-	-	-	-	-	-	-	1	1	-	7
Female	112	-	-	-	-	-	2	-	1	-	3	-
White, both sexes	240	-	-	-	-	-	1	-	1	1	3	5
Male	135	-	-	-	-	-	-	-	1	1	-	5
Female	105	-	-	-	-	-	1	-	-	-	3	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	1	-	1	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	1	-	1	-	-	-
Mississippi												
All races, both sexes	112	-	-	-	-	1	-	-	-	2	3	2
Male	70	-	-	-	-	1	-	-	-	1	3	-
Female	42	-	-	-	-	-	-	-	-	1	-	2
White, both sexes	83	-	-	-	-	-	-	-	-	2	1	1
Male	50	-	-	-	-	-	-	-	-	1	1	-
Female	33	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	29	-	-	-	-	1	-	-	-	-	2	1
Male	20	-	-	-	-	1	-	-	-	-	2	-
Female	9	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Missouri												
All races, both sexes	258	-	-	-	-	-	1	-	1	2	5	3
Male	162	-	-	-	-	-	1	-	1	2	4	3
Female	96	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	233	-	-	-	-	-	1	-	1	1	5	2
Male	149	-	-	-	-	-	1	-	1	1	4	2
Female	84	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	24	-	-	-	-	-	-	-	-	1	-	1
Male	12	-	-	-	-	-	-	-	-	1	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	72	-	-	-	-	-	-	-	-	-	2	2
Male	43	-	-	-	-	-	-	-	-	-	1	1
Female	29	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	67	-	-	-	-	-	-	-	-	-	1	2
Male	41	-	-	-	-	-	-	-	-	-	1	1
Female	26	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Nebraska												
All races, both sexes	101	-	-	-	-	-	-	-	-	2	2	-
Male	59	-	-	-	-	-	-	-	-	2	-	-
Female	42	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	97	-	-	-	-	-	-	-	-	2	2	-
Male	57	-	-	-	-	-	-	-	-	2	-	-
Female	40	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	98	-	-	-	-	-	1	-	1	3	1	1
Male	58	-	-	-	-	-	-	-	-	3	1	-
Female	40	-	-	-	-	-	1	-	1	-	-	1
White, both sexes	78	-	-	-	-	-	1	-	1	3	-	-
Male	52	-	-	-	-	-	-	-	-	3	-	-
Female	26	-	-	-	-	-	1	-	1	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
New Hampshire												
All races, both sexes	56	1	-	-	-	-	-	-	-	1	1	1
Male	30	-	-	-	-	-	-	-	-	1	1	1
Female	26	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	56	1	-	-	-	-	-	-	-	1	1	1
Male	30	-	-	-	-	-	-	-	-	1	1	1
Female	26	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	345	-	-	-	-	1	1	-	-	3	8	6
Male	183	-	-	-	-	1	1	-	-	2	7	5
Female	162	-	-	-	-	-	-	-	-	1	1	1
White, both sexes	297	-	-	-	-	1	1	-	-	2	4	5
Male	159	-	-	-	-	1	1	-	-	2	4	4
Female	138	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	36	-	-	-	-	-	-	-	-	1	4	1
Male	20	-	-	-	-	-	-	-	-	-	3	1
Female	16	-	-	-	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
New Mexico												
All races, both sexes	71	-	-	-	-	-	-	1	-	1	-	1
Male	41	-	-	-	-	-	-	1	-	1	-	1
Female	30	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	65	-	-	-	-	-	-	-	-	1	-	1
Male	36	-	-	-	-	-	-	-	-	1	-	1
Female	29	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	771	-	-	-	-	1	1	1	3	6	13	22
Male	442	-	-	-	-	1	1	1	3	6	12	16
Female	329	-	-	-	-	-	-	-	-	-	1	6
White, both sexes	665	-	-	-	-	-	1	1	3	3	9	16
Male	382	-	-	-	-	-	1	1	3	3	8	11
Female	283	-	-	-	-	-	-	-	-	-	1	5
Black, both sexes	91	-	-	-	-	1	-	-	-	3	4	6
Male	54	-	-	-	-	1	-	-	-	3	4	5
Female	37	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
North Carolina												
All races, both sexes	393	-	-	-	-	-	2	6	4	3	5	6
Male	230	-	-	-	-	-	2	5	3	2	5	6
Female	163	-	-	-	-	-	-	1	1	1	-	-
White, both sexes	339	-	-	-	-	-	2	5	1	2	4	2
Male	200	-	-	-	-	-	2	4	-	1	4	2
Female	139	-	-	-	-	-	-	1	1	1	-	-
Black, both sexes	52	-	-	-	-	-	-	1	2	1	1	4
Male	29	-	-	-	-	-	-	1	2	1	1	4
Female	23	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	39	-	-	-	-	-	-	-	-	1	-	1
Male	25	-	-	-	-	-	-	-	-	1	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	36	-	-	-	-	-	-	-	-	-	-	1
Male	23	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Ohio												
All races, both sexes	622	-	-	-	1	1	1	1	3	4	10	15
Male	362	-	-	-	1	1	1	1	2	4	6	11
Female	260	-	-	-	-	-	-	-	1	-	4	4
White, both sexes	556	-	-	-	1	1	-	-	2	4	9	8
Male	320	-	-	-	1	1	-	-	2	4	5	6
Female	236	-	-	-	-	-	-	-	-	-	4	2
Black, both sexes	61	-	-	-	-	-	1	1	1	-	1	7
Male	38	-	-	-	-	-	1	1	-	-	1	5
Female	23	-	-	-	-	-	-	-	1	-	-	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	179	-	-	-	-	-	-	-	4	1	2	6
Male	107	-	-	-	-	-	-	-	3	1	1	5
Female	72	-	-	-	-	-	-	-	1	-	1	1
White, both sexes	158	-	-	-	-	-	-	-	2	1	2	4
Male	93	-	-	-	-	-	-	-	1	1	1	3
Female	65	-	-	-	-	-	-	-	1	-	1	1
Black, both sexes	9	-	-	-	-	-	-	-	2	-	-	2
Male	7	-	-	-	-	-	-	-	2	-	-	2
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Oregon												
All races, both sexes	170	-	-	-	-	-	-	-	1	-	6	4
Male	100	-	-	-	-	-	-	-	-	-	5	2
Female	70	-	-	-	-	-	-	-	1	-	1	2
White, both sexes	160	-	-	-	-	-	-	-	1	-	5	4
Male	95	-	-	-	-	-	-	-	-	-	4	2
Female	65	-	-	-	-	-	-	-	1	-	1	2
Black, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	722	-	-	-	1	1	1	2	3	1	11	10
Male	374	-	-	-	1	1	1	2	3	1	8	8
Female	348	-	-	-	-	-	-	-	-	-	3	2
White, both sexes	664	-	-	-	1	1	1	1	3	1	7	6
Male	339	-	-	-	1	1	1	1	3	1	6	4
Female	325	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	50	-	-	-	-	-	-	-	-	-	3	4
Male	29	-	-	-	-	-	-	-	-	-	1	4
Female	21	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	1	-	-	-	-
Male	5	-	-	-	-	-	-	1	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Rhode Island												
All races, both sexes	57	-	-	-	-	-	-	-	-	-	-	-
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	36	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	53	-	-	-	-	-	-	-	-	-	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	33	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	208	-	-	-	-	1	-	-	2	5	6	7
Male	114	-	-	-	-	-	-	-	2	1	3	5
Female	94	-	-	-	-	1	-	-	-	4	3	2
White, both sexes	154	-	-	-	-	1	-	-	2	1	1	3
Male	87	-	-	-	-	-	-	-	2	-	-	2
Female	67	-	-	-	-	1	-	-	-	1	1	1
Black, both sexes	53	-	-	-	-	-	-	-	-	4	5	4
Male	26	-	-	-	-	-	-	-	-	1	3	3
Female	27	-	-	-	-	-	-	-	-	3	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
South Dakota												
All races, both sexes	47	-	-	-	-	-	1	-	-	-	-	-
Male	26	-	-	-	-	-	1	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	44	-	-	-	-	-	1	-	-	-	-	-
Male	26	-	-	-	-	-	1	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	253	-	-	-	-	-	-	2	2	2	5	3
Male	149	-	-	-	-	-	-	1	-	2	5	2
Female	104	-	-	-	-	-	-	1	2	-	-	1
White, both sexes	217	-	-	-	-	-	-	2	1	2	3	1
Male	133	-	-	-	-	-	-	1	-	2	3	1
Female	84	-	-	-	-	-	-	1	1	-	-	-
Black, both sexes	33	-	-	-	-	-	-	-	1	-	2	2
Male	16	-	-	-	-	-	-	-	-	-	2	1
Female	17	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Texas												
All races, both sexes	634	-	-	-	-	-	2	3	5	10	13	27
Male	404	-	-	-	-	-	2	1	2	6	12	18
Female	230	-	-	-	-	-	-	2	3	4	1	9
White, both sexes	552	-	-	-	-	-	1	3	5	9	8	20
Male	357	-	-	-	-	-	1	1	2	6	7	14
Female	195	-	-	-	-	-	-	2	3	3	1	6
Black, both sexes	71	-	-	-	-	-	1	-	-	1	5	7
Male	42	-	-	-	-	-	1	-	-	-	5	4
Female	29	-	-	-	-	-	-	-	-	1	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	68	-	-	-	-	-	-	-	-	-	1	1
Male	41	-	-	-	-	-	-	-	-	-	1	1
Female	27	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	63	-	-	-	-	-	-	-	-	-	-	-
Male	38	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Vermont												
All races, both sexes	57	-	-	-	-	-	-	-	-	-	1	1
Male	34	-	-	-	-	-	-	-	-	-	1	1
Female	23	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	57	-	-	-	-	-	-	-	-	-	1	1
Male	34	-	-	-	-	-	-	-	-	-	1	1
Female	23	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	285	-	-	-	-	1	1	-	2	5	2	7
Male	168	-	-	-	-	1	1	-	2	5	1	7
Female	117	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	228	-	-	-	-	-	1	-	1	2	2	5
Male	135	-	-	-	-	-	1	-	1	2	1	5
Female	93	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	52	-	-	-	-	1	-	-	1	3	-	2
Male	30	-	-	-	-	1	-	-	1	3	-	2
Female	22	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Washington												
All races, both sexes	321	-	-	-	-	-	-	1	-	3	5	6
Male	184	-	-	-	-	-	-	-	-	3	4	4
Female	137	-	-	-	-	-	-	1	-	-	1	2
White, both sexes	307	-	-	-	-	-	-	1	-	3	5	5
Male	178	-	-	-	-	-	-	-	-	3	4	4
Female	129	-	-	-	-	-	-	1	-	-	1	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
West Virginia												
All races, both sexes	113	-	-	-	-	-	-	1	1	-	1	-
Male	60	-	-	-	-	-	-	1	-	-	-	-
Female	53	-	-	-	-	-	-	-	1	-	1	-
White, both sexes	112	-	-	-	-	-	-	1	1	-	1	-
Male	60	-	-	-	-	-	-	1	-	-	-	-
Female	52	-	-	-	-	-	-	-	1	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Wisconsin												
All races, both sexes	339	-	-	-	-	1	-	-	1	2	3	8
Male	202	-	-	-	-	1	-	-	-	2	1	6
Female	137	-	-	-	-	-	-	-	1	-	2	2
White, both sexes	322	-	-	-	-	1	-	-	-	-	3	7
Male	193	-	-	-	-	1	-	-	-	-	1	5
Female	129	-	-	-	-	-	-	-	-	-	2	2
Black, both sexes	14	-	-	-	-	-	-	-	1	1	-	1
Male	7	-	-	-	-	-	-	-	-	1	-	1
Female	7	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	41	-	-	-	-	-	-	-	1	-	-	1
Male	25	-	-	-	-	-	-	-	1	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	41	-	-	-	-	-	-	-	1	-	-	1
Male	25	-	-	-	-	-	-	-	1	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
United States												
All races, both sexes	448	610	873	1,094	1,522	1,992	2,354	1,964	1,019	274	22	1
Male	341	453	640	751	947	1,119	1,222	937	430	83	3	1
Female	107	157	233	343	575	873	1,132	1,027	589	191	19	-
White, both sexes	309	462	730	925	1,370	1,807	2,152	1,823	957	249	18	1
Male	241	352	557	652	862	1,022	1,136	876	419	74	2	1
Female	68	110	173	273	508	785	1,016	947	538	175	16	-
Black, both sexes	121	126	106	128	110	139	132	98	43	18	3	-
Male	89	88	61	71	58	74	44	36	7	5	1	-
Female	32	38	45	57	52	65	88	62	36	13	2	-
American Indian 1/, both sexes	2	1	6	7	8	4	8	7	2	3	-	-
Male	1	1	3	5	7	2	6	2	1	2	-	-
Female	1	-	3	2	1	2	2	5	1	1	-	-
Asian or Pacific Islander 2/, both sexes	16	21	31	34	34	42	62	36	17	4	1	-
Male	10	12	19	23	20	21	36	23	3	2	-	-
Female	6	9	12	11	14	21	26	13	14	2	1	-
Alabama												
All races, both sexes	7	12	17	21	29	40	25	25	14	5	-	-
Male	6	11	14	14	18	21	13	14	7	2	-	-
Female	1	1	3	7	11	19	12	11	7	3	-	-
White, both sexes	4	8	14	16	27	32	17	23	12	3	-	-
Male	4	7	11	11	18	17	9	13	6	1	-	-
Female	-	1	3	5	9	15	8	10	6	2	-	-
Black, both sexes	3	3	3	5	2	8	8	2	1	2	-	-
Male	2	3	3	3	-	4	4	1	-	1	-	-
Female	1	-	-	2	2	4	4	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	1	-	-	-
Male	-	1	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Alaska												
All races, both sexes	1	-	-	1	1	1	1	3	2	-	-	-
Male	-	-	-	1	1	1	1	2	1	-	-	-
Female	1	-	-	-	-	-	-	1	1	-	-	-
White, both sexes	1	-	-	1	1	1	1	2	2	-	-	-
Male	-	-	-	1	1	1	1	2	1	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	7	16	22	12	25	36	46	27	21	7	1	-
Male	7	12	18	6	13	24	29	9	8	3	-	-
Female	-	4	4	6	12	12	17	18	13	4	1	-
White, both sexes	6	15	21	12	25	35	44	27	21	7	1	-
Male	6	12	17	6	13	23	29	9	8	3	-	-
Female	-	3	4	6	12	12	15	18	13	4	1	-
Black, both sexes	1	1	-	-	-	-	2	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Arkansas												
All races, both sexes	6	6	11	15	16	27	20	21	11	4	-	-
Male	3	5	8	9	8	17	8	11	5	2	-	-
Female	3	1	3	6	8	10	12	10	6	2	-	-
White, both sexes	4	5	9	14	14	26	18	19	11	4	-	-
Male	2	5	7	9	8	17	8	9	5	2	-	-
Female	2	-	2	5	6	9	10	10	6	2	-	-
Black, both sexes	2	1	1	1	2	-	1	2	-	-	-	-
Male	1	-	1	-	-	-	-	2	-	-	-	-
Female	1	1	-	1	2	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	44	73	93	93	143	191	234	174	77	24	3	1
Male	34	56	69	65	97	111	111	85	29	12	-	1
Female	10	17	24	28	46	80	123	89	48	12	3	-
White, both sexes	27	49	73	76	118	155	188	152	68	21	2	1
Male	21	41	56	54	81	90	82	72	28	10	-	1
Female	6	8	17	22	37	65	106	80	40	11	2	-
Black, both sexes	9	15	12	3	7	12	17	6	1	-	-	-
Male	7	9	7	3	4	6	6	2	-	-	-	-
Female	2	6	5	-	3	6	11	4	1	-	-	-
American Indian 1/, both sexes	-	1	-	-	2	1	-	1	-	-	-	-
Male	-	1	-	-	2	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	8	8	14	16	23	29	15	8	3	1	-
Male	6	5	6	8	10	14	23	10	1	2	-	-
Female	2	3	2	6	6	9	6	5	7	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Colorado												
All races, both sexes	4	11	7	20	20	25	35	41	18	6	-	-
Male	4	7	5	15	13	12	23	13	11	2	-	-
Female	-	4	2	5	7	13	12	28	7	4	-	-
White, both sexes	3	10	6	19	19	25	32	41	17	6	-	-
Male	3	7	5	15	13	12	21	13	11	2	-	-
Female	-	3	1	4	6	13	11	28	6	4	-	-
Black, both sexes	1	1	1	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	2	-	1	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	1	-	-	-
Connecticut												
All races, both sexes	5	2	9	11	10	31	27	26	14	3	1	-
Male	3	1	7	9	5	19	13	10	2	-	1	-
Female	2	1	2	2	5	12	14	16	12	3	-	-
White, both sexes	4	1	6	10	9	28	27	25	12	3	-	-
Male	3	-	5	8	5	17	13	9	2	-	-	-
Female	1	1	1	2	4	11	14	16	10	3	-	-
Black, both sexes	1	1	2	1	-	3	-	-	2	-	1	-
Male	-	1	2	1	-	2	-	-	-	-	1	-
Female	1	-	-	-	-	1	-	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Delaware												
All races, both sexes	-	1	1	2	3	5	5	4	5	-	-	-
Male	-	-	-	1	2	3	3	1	2	-	-	-
Female	-	1	1	1	1	2	2	3	3	-	-	-
White, both sexes	-	1	1	2	2	3	4	4	3	-	-	-
Male	-	-	-	1	1	3	3	1	1	-	-	-
Female	-	1	1	1	1	-	1	3	2	-	-	-
Black, both sexes	-	-	-	-	1	1	1	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	3	5	1	2	2	2	3	4	-	-	-	-
Male	3	3	-	1	2	-	-	1	-	-	-	-
Female	-	2	1	1	-	2	3	3	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	3	4	1	2	2	1	2	3	-	-	-	-
Male	3	3	-	1	2	-	-	1	-	-	-	-
Female	-	1	1	1	-	1	2	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Florida												
All races, both sexes	37	38	65	79	107	153	185	128	73	16	2	-
Male	27	30	46	54	62	86	110	70	36	7	-	-
Female	10	8	19	25	45	67	75	58	37	9	2	-
White, both sexes	28	32	58	71	97	140	177	125	72	16	2	-
Male	21	25	43	50	53	80	109	68	36	7	-	-
Female	7	7	15	21	44	60	68	57	36	9	2	-
Black, both sexes	9	6	6	8	10	12	7	3	1	-	-	-
Male	6	5	2	4	9	6	1	2	-	-	-	-
Female	3	1	4	4	1	6	6	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Georgia												
All races, both sexes	14	16	23	27	35	33	40	25	11	2	1	-
Male	9	11	17	20	26	22	25	11	5	1	-	-
Female	5	5	6	7	9	11	15	14	6	1	1	-
White, both sexes	7	9	13	20	26	24	37	20	10	1	-	-
Male	5	6	11	15	20	14	24	10	5	-	-	-
Female	2	3	2	5	6	10	13	10	5	1	-	-
Black, both sexes	7	6	9	7	9	9	3	4	1	1	1	-
Male	4	4	5	5	6	8	1	1	-	1	-	-
Female	3	2	4	2	3	1	2	3	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	-	-	-	1	-	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Hawaii												
All races, both sexes	1	7	6	3	7	8	7	12	6	2	-	-
Male	1	4	5	2	6	2	3	6	1	-	-	-
Female	-	3	1	1	1	6	4	6	5	2	-	-
White, both sexes	-	2	3	-	2	2	3	3	4	1	-	-
Male	-	1	3	-	2	-	1	-	-	-	-	-
Female	-	1	-	-	-	2	2	3	4	1	-	-
Black, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	4	2	3	5	6	4	9	2	1	-	-
Male	1	2	1	2	4	2	2	6	1	-	-	-
Female	-	2	1	1	1	4	2	3	1	1	-	-
Idaho												
All races, both sexes	2	2	1	3	10	22	12	14	5	-	-	-
Male	2	2	1	3	7	12	6	7	2	-	-	-
Female	-	-	-	-	3	10	6	7	3	-	-	-
White, both sexes	2	2	1	3	9	21	12	14	5	-	-	-
Male	2	2	1	3	6	12	6	7	2	-	-	-
Female	-	-	-	-	3	9	6	7	3	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Illinois												
All races, both sexes	19	35	37	58	62	81	92	89	54	14	2	-
Male	16	26	27	37	34	44	42	40	22	5	-	-
Female	3	9	10	21	28	37	50	49	32	9	2	-
White, both sexes	17	29	33	46	50	70	86	81	50	9	2	-
Male	14	22	24	29	28	38	39	37	22	3	-	-
Female	3	7	9	17	22	32	47	44	28	6	2	-
Black, both sexes	1	6	4	10	10	10	6	7	3	5	-	-
Male	1	4	3	6	4	5	3	2	-	2	-	-
Female	-	2	1	4	6	5	3	5	3	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	2	2	1	-	1	1	-	-	-
Male	1	-	-	2	2	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Indiana												
All races, both sexes	11	18	27	26	47	49	57	49	23	7	-	-
Male	10	10	18	18	33	28	27	29	12	1	-	-
Female	1	8	9	8	14	21	30	20	11	6	-	-
White, both sexes	8	16	24	23	47	45	53	47	23	7	-	-
Male	8	9	16	16	33	26	26	27	12	1	-	-
Female	-	7	8	7	14	19	27	20	11	6	-	-
Black, both sexes	2	2	3	3	-	4	3	2	-	-	-	-
Male	2	1	2	2	-	2	-	2	-	-	-	-
Female	-	1	1	1	-	2	3	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Iowa												
All races, both sexes	3	4	9	14	22	15	28	25	17	2	-	-
Male	3	3	4	10	15	8	17	12	6	-	-	-
Female	-	1	5	4	7	7	11	13	11	2	-	-
White, both sexes	2	4	8	14	22	15	28	25	17	2	-	-
Male	2	3	3	10	15	8	17	12	6	-	-	-
Female	-	1	5	4	7	7	11	13	11	2	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	6	2	8	14	12	23	26	17	11	4	1	-
Male	6	1	6	8	9	15	13	12	6	2	-	-
Female	-	1	2	6	3	8	13	5	5	2	1	-
White, both sexes	5	2	8	13	12	23	25	17	10	4	1	-
Male	5	1	6	8	9	15	13	12	6	2	-	-
Female	-	1	2	5	3	8	12	5	4	2	1	-
Black, both sexes	1	-	-	1	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Kentucky												
All races, both sexes	5	14	11	25	27	31	40	29	12	2	-	-
Male	4	11	7	18	17	17	22	18	3	1	-	-
Female	1	3	4	7	10	14	18	11	9	1	-	-
White, both sexes	2	12	11	24	27	29	37	28	11	2	-	-
Male	1	9	7	17	17	16	21	18	2	1	-	-
Female	1	3	4	7	10	13	16	10	9	1	-	-
Black, both sexes	3	2	-	-	-	2	3	1	1	-	-	-
Male	3	2	-	-	-	1	1	-	1	-	-	-
Female	-	-	-	-	-	1	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	6	8	18	19	24	16	30	19	17	2	-	-
Male	4	7	12	13	13	9	12	7	7	-	-	-
Female	2	1	6	6	11	7	18	12	10	2	-	-
White, both sexes	3	4	14	15	22	13	25	14	14	1	-	-
Male	2	4	10	10	13	8	10	6	6	-	-	-
Female	1	-	4	5	9	5	15	8	8	1	-	-
Black, both sexes	3	4	4	4	2	3	5	5	3	1	-	-
Male	2	3	2	3	-	1	2	1	1	-	-	-
Female	1	1	2	1	2	2	3	4	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Maine												
All races, both sexes	-	-	7	4	11	11	17	20	5	-	-	-
Male	-	-	5	2	9	4	8	9	3	-	-	-
Female	-	-	2	2	2	7	9	11	2	-	-	-
White, both sexes	-	-	6	4	11	11	17	20	4	-	-	-
Male	-	-	5	2	9	4	8	9	3	-	-	-
Female	-	-	1	2	2	7	9	11	1	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Maryland												
All races, both sexes	11	10	9	21	22	36	39	37	19	2	2	-
Male	8	8	7	11	12	13	19	12	10	-	-	-
Female	3	2	2	10	10	23	20	25	9	2	2	-
White, both sexes	3	3	7	14	20	34	33	32	18	2	2	-
Male	2	3	6	7	12	12	19	11	10	-	-	-
Female	1	-	1	7	8	22	14	21	8	2	2	-
Black, both sexes	8	7	2	7	2	2	6	3	1	-	-	-
Male	6	5	1	4	-	1	-	-	-	-	-	-
Female	2	2	1	3	2	1	6	3	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Massachusetts												
All races, both sexes	8	10	15	19	38	48	46	57	31	8	1	-
Male	7	8	12	15	23	21	19	28	9	3	-	-
Female	1	2	3	4	15	27	27	29	22	5	1	-
White, both sexes	8	10	13	17	35	48	46	56	30	8	1	-
Male	7	8	10	13	21	21	19	28	9	3	-	-
Female	1	2	3	4	14	27	27	28	21	5	1	-
Black, both sexes	-	-	2	2	1	-	-	-	1	-	-	-
Male	-	-	2	2	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Michigan												
All races, both sexes	24	26	40	41	62	94	90	74	56	8	-	-
Male	18	17	33	26	39	55	51	30	27	3	-	-
Female	6	9	7	15	23	39	39	44	29	5	-	-
White, both sexes	15	21	32	35	57	89	84	66	54	8	-	-
Male	13	14	28	23	36	52	48	28	26	3	-	-
Female	2	7	4	12	21	37	36	38	28	5	-	-
Black, both sexes	9	5	8	4	3	5	5	7	2	-	-	-
Male	5	3	5	1	1	3	2	1	1	-	-	-
Female	4	2	3	3	2	2	3	6	1	-	-	-
American Indian 1/, both sexes	-	-	-	1	1	-	1	1	-	-	-	-
Male	-	-	-	1	1	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Minnesota												
All races, both sexes	8	9	19	17	26	39	46	38	26	12	-	-
Male	6	6	12	14	19	24	23	16	12	2	-	-
Female	2	3	7	3	7	15	23	22	14	10	-	-
White, both sexes	5	7	14	17	26	38	46	38	26	12	-	-
Male	4	4	10	14	19	24	23	16	12	2	-	-
Female	1	3	4	3	7	14	23	22	14	10	-	-
Black, both sexes	2	1	2	-	-	1	-	-	-	-	-	-
Male	2	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	2	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	2	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	9	5	7	11	18	19	14	13	5	2	1	-
Male	6	4	6	9	13	15	5	7	-	-	-	-
Female	3	1	1	2	5	4	9	6	5	2	1	-
White, both sexes	7	2	4	7	17	13	11	11	4	2	1	-
Male	5	2	4	6	12	10	4	5	-	-	-	-
Female	2	-	-	1	5	3	7	6	4	2	1	-
Black, both sexes	2	3	3	4	1	6	3	2	1	-	-	-
Male	1	2	2	3	1	5	1	2	-	-	-	-
Female	1	1	1	1	-	1	2	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Missouri												
All races, both sexes	10	10	11	19	36	34	53	41	22	9	1	-
Male	7	8	9	16	22	18	30	25	12	3	1	-
Female	3	2	2	3	14	16	23	16	10	6	-	-
White, both sexes	9	8	9	16	33	29	51	37	22	8	1	-
Male	6	6	8	13	22	16	29	24	12	3	1	-
Female	3	2	1	3	11	13	22	13	10	5	-	-
Black, both sexes	1	2	1	3	3	5	2	4	-	1	-	-
Male	1	2	-	3	-	2	1	1	-	-	-	-
Female	-	-	1	-	3	3	1	3	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	6	6	4	13	7	15	10	5	1	-	-
Male	1	5	4	4	7	6	9	4	1	-	-	-
Female	-	1	2	-	6	1	6	6	4	1	-	-
White, both sexes	1	6	4	3	13	7	14	10	5	1	-	-
Male	1	5	4	3	7	6	8	4	1	-	-	-
Female	-	1	-	-	6	1	6	6	4	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	2	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	2	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Nebraska												
All races, both sexes	5	4	3	11	8	8	27	21	8	2	-	-
Male	5	2	3	8	4	3	15	14	3	-	-	-
Female	-	2	-	3	4	5	12	7	5	2	-	-
White, both sexes	4	3	3	10	8	8	26	21	8	2	-	-
Male	4	1	3	8	4	3	15	14	3	-	-	-
Female	-	2	-	2	4	5	11	7	5	2	-	-
Black, both sexes	1	1	-	1	-	-	1	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	5	3	12	11	13	16	16	11	3	1	-	-
Male	3	2	9	6	6	10	10	7	1	-	-	-
Female	2	1	3	5	7	6	6	4	2	1	-	-
White, both sexes	3	3	7	7	11	13	14	11	3	1	-	-
Male	3	2	6	6	6	9	9	7	1	-	-	-
Female	-	1	1	1	5	4	5	4	2	1	-	-
Black, both sexes	-	-	1	3	1	2	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	3	1	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	4	1	1	1	1	-	-	-	-	-
Male	-	-	3	-	-	1	-	-	-	-	-	-
Female	2	-	1	1	1	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
New Hampshire												
All races, both sexes	-	4	6	1	3	11	10	11	3	3	-	-
Male	-	2	5	-	3	7	5	5	-	-	-	-
Female	-	2	1	1	-	4	5	6	3	3	-	-
White, both sexes	-	4	6	1	3	11	10	11	3	3	-	-
Male	-	2	5	-	3	7	5	5	-	-	-	-
Female	-	2	1	1	-	4	5	6	3	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	5	15	20	25	34	51	80	61	31	4	-	-
Male	4	14	12	19	18	28	34	22	15	1	-	-
Female	1	1	8	6	16	23	46	39	16	3	-	-
White, both sexes	4	11	17	14	29	50	70	55	31	3	-	-
Male	3	10	11	12	16	28	30	21	15	1	-	-
Female	1	1	6	2	13	22	40	34	16	2	-	-
Black, both sexes	1	4	2	8	3	-	6	5	-	1	-	-
Male	1	4	1	5	1	-	3	1	-	-	-	-
Female	-	-	1	3	2	-	3	4	-	1	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	2	2	1	4	-	-	-	-	-
Male	-	-	-	2	1	-	1	-	-	-	-	-
Female	-	-	1	-	1	1	3	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
New Mexico												
All races, both sexes	3	1	3	2	10	19	9	12	8	1	-	-
Male	3	1	3	-	8	9	6	4	3	1	-	-
Female	-	-	-	2	2	10	3	8	5	-	-	-
White, both sexes	2	1	3	2	9	19	9	11	7	-	-	-
Male	2	1	3	-	7	9	6	4	2	-	-	-
Female	-	-	-	2	2	10	3	7	5	-	-	-
Black, both sexes	1	-	-	-	1	-	-	-	1	-	-	-
Male	1	-	-	-	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	35	41	43	66	79	123	137	128	63	9	-	-
Male	25	31	30	46	45	76	68	57	24	-	-	-
Female	10	10	13	20	34	47	69	71	39	9	-	-
White, both sexes	21	30	28	54	71	111	127	120	61	9	-	-
Male	17	23	18	39	41	71	66	56	24	-	-	-
Female	4	7	10	15	30	40	61	64	37	9	-	-
Black, both sexes	14	10	10	10	7	11	7	7	1	-	-	-
Male	8	8	8	5	4	5	2	1	-	-	-	-
Female	6	2	2	5	3	6	5	6	1	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	4	2	1	1	3	1	1	-	-	-
Male	-	-	4	2	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	1	3	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
North Carolina												
All races, both sexes	16	17	17	39	52	64	68	52	30	11	1	-
Male	11	14	10	30	29	38	35	26	12	2	-	-
Female	5	3	7	9	23	26	33	26	18	9	1	-
White, both sexes	13	12	13	33	44	60	65	46	26	10	1	-
Male	9	10	8	27	24	36	35	25	11	2	-	-
Female	4	2	5	6	20	24	30	21	15	8	1	-
Black, both sexes	3	5	4	6	8	4	3	5	4	1	-	-
Male	2	4	2	3	5	2	-	1	1	-	-	-
Female	1	1	2	3	3	2	3	4	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
North Dakota												
All races, both sexes	1	1	1	4	6	9	5	5	4	1	-	-
Male	1	1	1	1	5	4	5	4	-	1	-	-
Female	-	-	-	3	1	5	-	1	4	-	-	-
White, both sexes	1	1	-	3	6	9	5	5	4	1	-	-
Male	1	1	-	1	5	4	5	4	-	1	-	-
Female	-	-	-	2	1	5	-	1	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Ohio												
All races, both sexes	22	21	43	56	55	94	134	98	49	14	-	-
Male	18	16	32	41	31	56	66	49	22	4	-	-
Female	4	5	11	15	24	38	68	49	27	10	-	-
White, both sexes	17	13	41	50	47	85	126	94	45	13	-	-
Male	13	10	31	37	27	49	64	46	21	3	-	-
Female	4	3	10	13	20	36	62	48	24	10	-	-
Black, both sexes	5	8	2	5	8	6	8	4	4	-	-	-
Male	5	6	1	3	4	5	2	3	1	-	-	-
Female	-	2	1	2	4	1	6	1	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	3	-	-	-	-	-	-
Male	-	-	-	1	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Oklahoma												
All races, both sexes	10	7	14	18	27	26	31	23	7	3	-	-
Male	6	6	10	16	17	12	19	11	-	-	-	-
Female	4	1	4	2	10	14	12	12	7	3	-	-
White, both sexes	6	7	14	15	25	24	27	22	7	2	-	-
Male	5	6	10	13	15	12	16	10	-	-	-	-
Female	1	1	4	2	10	12	11	12	7	2	-	-
Black, both sexes	3	-	-	-	-	-	1	1	-	-	-	-
Male	1	-	-	-	-	-	1	1	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	3	2	-	3	-	-	1	-	-
Male	-	-	-	3	2	-	2	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Oregon												
All races, both sexes	3	9	13	17	20	27	30	25	12	3	-	-
Male	2	6	10	11	14	16	15	10	7	2	-	-
Female	1	3	3	6	6	11	15	15	5	1	-	-
White, both sexes	2	8	13	16	19	27	27	23	12	3	-	-
Male	2	6	10	10	14	16	13	9	7	2	-	-
Female	-	2	3	6	5	11	14	14	5	1	-	-
Black, both sexes	1	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	3	1	-	-	-	-
Male	-	-	-	-	-	-	2	1	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
Pennsylvania												
All races, both sexes	16	17	43	36	90	121	139	133	64	32	1	-
Male	9	14	35	19	48	57	73	56	27	11	-	-
Female	7	3	8	17	42	64	66	77	37	21	1	-
White, both sexes	6	14	38	35	88	114	130	127	60	31	-	-
Male	1	12	31	18	47	53	67	54	27	11	-	-
Female	5	2	7	17	41	61	63	73	33	20	-	-
Black, both sexes	10	3	4	1	1	5	7	6	4	1	1	-
Male	8	2	3	1	1	3	4	2	-	-	-	-
Female	2	1	1	-	-	2	3	4	4	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	2	2	-	-	-	-	-
Male	-	-	1	-	-	1	2	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Rhode Island												
All races, both sexes	-	3	3	4	9	5	19	11	2	1	-	-
Male	-	2	2	3	3	2	5	4	-	-	-	-
Female	-	1	1	1	6	3	14	7	2	1	-	-
White, both sexes	-	1	3	3	9	5	19	10	2	1	-	-
Male	-	1	2	3	3	2	5	4	-	-	-	-
Female	-	-	1	-	6	3	14	6	2	1	-	-
Black, both sexes	-	1	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	8	10	18	28	29	24	32	25	12	1	-	-
Male	6	6	13	14	19	14	14	12	5	-	-	-
Female	2	4	5	14	10	10	18	13	7	1	-	-
White, both sexes	5	4	12	18	25	23	27	21	10	1	-	-
Male	4	2	11	9	17	14	12	9	5	-	-	-
Female	1	2	1	9	8	9	15	12	5	1	-	-
Black, both sexes	2	6	6	10	4	1	5	4	2	-	-	-
Male	1	4	2	5	2	-	2	3	-	-	-	-
Female	1	2	4	5	2	1	3	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
South Dakota												
All races, both sexes	-	1	4	2	4	5	12	10	6	2	-	-
Male	-	-	3	1	1	2	9	6	2	1	-	-
Female	-	1	1	1	3	3	3	4	4	1	-	-
White, both sexes	-	-	3	2	4	5	12	10	5	2	-	-
Male	-	-	3	1	1	2	9	6	2	1	-	-
Female	-	-	-	1	3	3	3	4	3	1	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	14	11	22	30	27	48	44	23	19	1	-	-
Male	13	8	17	16	19	27	23	11	5	-	-	-
Female	1	3	5	14	8	21	21	12	14	1	-	-
White, both sexes	11	6	19	24	24	45	40	21	17	1	-	-
Male	10	4	16	14	17	27	22	11	5	-	-	-
Female	1	2	3	10	7	18	18	10	12	1	-	-
Black, both sexes	3	5	3	5	3	3	2	2	2	-	-	-
Male	3	4	1	2	2	-	1	-	-	-	-	-
Female	-	1	2	3	1	3	1	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	2	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Texas												
All races, both sexes	25	35	53	56	72	93	109	80	40	10	1	-
Male	19	29	36	44	52	56	57	45	23	2	-	-
Female	6	6	17	12	20	37	52	35	17	8	1	-
White, both sexes	19	31	48	48	62	80	99	75	37	6	1	-
Male	13	27	34	38	46	49	55	40	22	2	-	-
Female	6	4	14	10	16	31	44	35	15	4	1	-
Black, both sexes	5	4	4	6	9	13	5	4	3	4	-	-
Male	5	2	2	4	6	7	1	4	1	-	-	-
Female	-	2	2	2	3	6	4	-	2	4	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	2	1	-	5	1	-	-	-	-
Male	1	-	-	2	-	-	1	1	-	-	-	-
Female	-	-	1	-	1	-	4	-	-	-	-	-
Utah												
All races, both sexes	4	8	3	6	8	8	8	13	7	1	-	-
Male	3	6	1	5	6	5	6	5	2	-	-	-
Female	1	2	2	1	2	3	2	8	5	1	-	-
White, both sexes	4	7	2	6	8	8	7	13	7	1	-	-
Male	3	5	1	5	6	5	6	5	2	-	-	-
Female	1	2	1	1	2	3	1	8	5	1	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Vermont												
All races, both sexes	2	1	5	10	7	6	9	6	4	5	-	-
Male	1	1	4	7	4	5	4	3	1	2	-	-
Female	1	-	1	3	3	1	5	3	3	3	-	-
White, both sexes	2	1	5	10	7	6	9	6	4	5	-	-
Male	1	1	4	7	4	5	4	3	1	2	-	-
Female	1	-	1	3	3	1	5	3	3	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	4	16	17	22	36	46	52	43	24	6	1	-
Male	2	10	16	16	22	23	32	19	9	2	-	-
Female	2	6	1	6	14	23	20	24	15	4	1	-
White, both sexes	2	11	15	20	30	37	41	35	20	5	1	-
Male	-	7	15	15	19	17	25	17	9	1	-	-
Female	2	4	-	5	11	20	16	18	11	4	1	-
Black, both sexes	2	5	2	2	5	9	8	8	3	1	-	-
Male	2	3	1	1	2	6	5	2	-	1	-	-
Female	-	2	1	1	3	3	3	6	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	3	-	1	-	-	-
Male	-	-	-	-	1	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Washington												
All races, both sexes	6	11	26	29	50	46	49	60	21	6	2	-
Male	5	7	20	20	30	24	28	30	7	1	1	-
Female	1	4	6	9	20	22	21	30	14	5	1	-
White, both sexes	6	11	24	25	47	46	46	59	21	6	2	-
Male	5	7	19	18	29	24	27	29	7	1	1	-
Female	1	4	5	7	18	22	19	30	14	5	1	-
Black, both sexes	-	-	-	1	2	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	3	1	-	3	1	-	-	-	-
Male	-	-	1	2	-	-	1	1	-	-	-	-
Female	-	-	1	1	1	-	2	-	-	-	-	-
West Virginia												
All races, both sexes	4	5	8	14	14	13	25	18	7	2	-	-
Male	3	4	5	7	13	7	8	6	6	-	-	-
Female	1	1	3	7	1	6	17	12	1	2	-	-
White, both sexes	4	4	8	14	14	13	25	18	7	2	-	-
Male	3	4	5	7	13	7	8	6	6	-	-	-
Female	1	-	3	7	1	6	17	12	1	2	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Wisconsin												
All races, both sexes	7	20	15	19	32	47	65	72	35	12	-	-
Male	6	13	11	13	21	29	37	41	17	4	-	-
Female	1	7	4	6	11	18	28	31	18	8	-	-
White, both sexes	5	18	15	16	30	47	63	70	35	12	-	-
Male	4	12	11	13	20	29	36	40	17	4	-	-
Female	1	6	4	3	10	18	27	30	18	8	-	-
Black, both sexes	2	2	-	3	2	-	2	-	-	-	-	-
Male	2	1	-	-	1	-	1	-	-	-	-	-
Female	-	1	-	3	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	3	1	7	9	5	11	1	-	1	-	-
Male	1	2	-	7	4	2	6	1	-	-	-	-
Female	-	1	1	-	5	3	5	-	-	1	-	-
White, both sexes	1	3	1	7	9	5	11	1	-	1	-	-
Male	1	2	-	7	4	2	6	1	-	-	-	-
Female	-	1	1	-	5	3	5	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
United States												
All races, both sexes	8,954	12	4	4	3	9	18	13	24	42	78	125
Male	3,937	7	1	4	1	5	11	7	10	25	47	76
Female	5,017	5	3	-	2	4	7	6	14	17	31	49
White, both sexes	7,577	8	4	3	2	8	13	9	19	36	59	92
Male	3,324	5	1	3	1	5	7	6	8	24	37	58
Female	4,253	3	3	-	1	3	6	3	11	12	22	34
Black, both sexes	1,213	2	-	1	-	1	4	3	5	6	18	29
Male	537	2	-	1	-	-	3	1	2	1	10	15
Female	676	-	-	-	-	1	1	2	3	5	8	14
American Indian 1/, both sexes	50	1	-	-	-	-	-	-	-	-	-	1
Male	24	-	-	-	-	-	-	-	-	-	-	1
Female	26	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	114	1	-	-	1	-	1	1	-	-	1	3
Male	52	-	-	-	-	-	1	-	-	-	-	2
Female	62	1	-	-	1	-	-	1	-	-	1	1
Alabama												
All races, both sexes	158	1	-	-	-	-	1	-	-	1	3	3
Male	72	1	-	-	-	-	1	-	-	1	2	-
Female	86	-	-	-	-	-	-	-	-	-	1	3
White, both sexes	121	1	-	-	-	-	1	-	-	1	3	1
Male	59	1	-	-	-	-	1	-	-	1	2	-
Female	62	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	36	-	-	-	-	-	-	-	-	-	-	2
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Alaska												
All races, both sexes	13	-	-	1	-	-	-	-	-	1	-	-
Male	4	-	-	1	-	-	-	-	-	1	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	1	-	-	-	-	-	1	-	-
Male	4	-	-	1	-	-	-	-	-	1	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	150	-	1	-	-	-	-	2	-	-	1	2
Male	69	-	-	-	-	-	-	2	-	-	1	1
Female	81	-	1	-	-	-	-	-	-	-	-	1
White, both sexes	143	-	1	-	-	-	-	2	-	-	1	2
Male	66	-	-	-	-	-	-	2	-	-	1	1
Female	77	-	1	-	-	-	-	-	-	-	-	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Arkansas												
All races, both sexes	118	-	-	-	-	-	-	-	-	-	1	3
Male	52	-	-	-	-	-	-	-	-	-	-	2
Female	66	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	103	-	-	-	-	-	-	-	-	-	1	3
Male	45	-	-	-	-	-	-	-	-	-	-	2
Female	58	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	768	1	-	-	1	2	1	2	5	4	7	7
Male	348	1	-	-	1	2	1	1	2	4	5	3
Female	420	-	-	-	-	-	-	1	3	-	2	4
White, both sexes	642	1	-	-	1	2	1	1	5	4	6	4
Male	284	1	-	-	1	2	1	1	2	4	4	1
Female	358	-	-	-	-	-	-	-	3	-	2	3
Black, both sexes	74	-	-	-	-	-	-	-	-	-	1	1
Male	36	-	-	-	-	-	-	-	-	-	1	-
Female	38	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	46	-	-	-	-	-	-	1	-	-	-	1
Male	25	-	-	-	-	-	-	-	-	-	-	1
Female	21	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Colorado												
All races, both sexes	94	1	-	-	-	-	-	-	-	-	-	-
Male	38	1	-	-	-	-	-	-	-	-	-	-
Female	56	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	84	1	-	-	-	-	-	-	-	-	-	-
Male	31	1	-	-	-	-	-	-	-	-	-	-
Female	53	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	120	-	1	1	-	-	-	-	-	-	2	-
Male	45	-	-	1	-	-	-	-	-	-	1	-
Female	75	-	1	-	-	-	-	-	-	-	1	-
White, both sexes	111	-	1	-	-	-	-	-	-	-	2	-
Male	42	-	-	-	-	-	-	-	-	-	1	-
Female	69	-	1	-	-	-	-	-	-	-	1	-
Black, both sexes	9	-	-	1	-	-	-	-	-	-	-	-
Male	3	-	-	1	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Delaware												
All races, both sexes	32	-	-	-	-	-	-	-	-	1	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	22	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	25	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	17	1	-	-	-	-	-	-	-	-	-	-
Male	5	1	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Florida												
All races, both sexes	599	-	-	1	-	-	3	-	2	5	3	10
Male	290	-	-	1	-	-	2	-	-	4	2	6
Female	309	-	-	-	-	-	1	-	2	1	1	4
White, both sexes	525	-	-	1	-	-	2	-	1	5	2	8
Male	256	-	-	1	-	-	1	-	-	4	1	5
Female	269	-	-	-	-	-	1	-	1	1	1	3
Black, both sexes	73	-	-	-	-	-	1	-	1	-	1	2
Male	34	-	-	-	-	-	1	-	-	-	1	1
Female	39	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	235	-	-	-	-	-	1	1	-	4	3	4
Male	103	-	-	-	-	-	1	1	-	1	3	2
Female	132	-	-	-	-	-	-	-	-	3	-	2
White, both sexes	147	-	-	-	-	-	-	1	-	2	2	1
Male	60	-	-	-	-	-	-	1	-	1	2	1
Female	87	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	87	-	-	-	-	-	1	-	-	2	1	3
Male	42	-	-	-	-	-	1	-	-	-	1	1
Female	45	-	-	-	-	-	-	-	-	2	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Hawaii												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	44	-	-	-	-	-	-	-	-	1	-	2
Male	17	-	-	-	-	-	-	-	-	-	-	1
Female	27	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	42	-	-	-	-	-	-	-	-	1	-	2
Male	17	-	-	-	-	-	-	-	-	-	-	1
Female	25	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Illinois												
All races, both sexes	358	-	-	-	-	-	1	-	-	1	4	2
Male	136	-	-	-	-	-	1	-	-	1	2	1
Female	222	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	297	-	-	-	-	-	1	-	-	1	4	1
Male	113	-	-	-	-	-	1	-	-	1	2	1
Female	184	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	57	-	-	-	-	-	-	-	-	-	-	1
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	35	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	204	-	-	-	1	-	-	-	-	2	4	5
Male	80	-	-	-	-	-	-	-	-	-	3	4
Female	124	-	-	-	1	-	-	-	-	2	1	1
White, both sexes	180	-	-	-	-	-	-	-	-	2	3	3
Male	69	-	-	-	-	-	-	-	-	-	2	2
Female	111	-	-	-	-	-	-	-	-	2	1	1
Black, both sexes	23	-	-	-	-	-	-	-	-	-	1	2
Male	11	-	-	-	-	-	-	-	-	-	1	2
Female	12	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Iowa												
All races, both sexes	103	-	-	-	-	-	-	-	1	-	-	2
Male	43	-	-	-	-	-	-	-	-	-	-	2
Female	60	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	101	-	-	-	-	-	-	-	1	-	-	1
Male	42	-	-	-	-	-	-	-	-	-	-	1
Female	59	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	81	-	-	-	-	-	-	-	1	1	-	2
Male	42	-	-	-	-	-	-	-	1	1	-	1
Female	39	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	74	-	-	-	-	-	-	-	1	1	-	2
Male	39	-	-	-	-	-	-	-	1	1	-	1
Female	35	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Kentucky												
All races, both sexes	172	-	-	-	-	1	-	-	-	1	1	4
Male	73	-	-	-	-	-	-	-	-	-	-	3
Female	99	-	-	-	-	1	-	-	-	1	1	1
White, both sexes	159	-	-	-	-	1	-	-	-	1	1	4
Male	67	-	-	-	-	-	-	-	-	-	-	3
Female	92	-	-	-	-	1	-	-	-	1	1	1
Black, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	142	-	1	-	1	-	-	-	1	-	-	2
Male	59	-	-	-	-	-	-	-	1	-	-	2
Female	83	-	1	-	1	-	-	-	-	-	-	-
White, both sexes	94	-	1	-	1	-	-	-	1	-	-	2
Male	37	-	-	-	-	-	-	-	1	-	-	2
Female	57	-	1	-	1	-	-	-	-	-	-	-
Black, both sexes	46	-	-	-	-	-	-	-	-	-	-	-
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Maine												
All races, both sexes	61	-	-	-	-	-	-	-	-	-	-	-
Male	27	-	-	-	-	-	-	-	-	-	-	-
Female	34	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	61	-	-	-	-	-	-	-	-	-	-	-
Male	27	-	-	-	-	-	-	-	-	-	-	-
Female	34	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	185	-	-	-	-	-	1	1	2	1	1	3
Male	77	-	-	-	-	-	-	1	1	1	-	-
Female	108	-	-	-	-	-	1	-	1	-	1	3
White, both sexes	136	-	-	-	-	-	1	-	1	1	-	2
Male	58	-	-	-	-	-	-	-	-	1	-	-
Female	78	-	-	-	-	-	1	-	1	-	-	2
Black, both sexes	49	-	-	-	-	-	-	1	1	-	1	1
Male	19	-	-	-	-	-	-	1	1	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Massachusetts												
All races, both sexes	220	-	-	-	-	-	-	-	1	-	-	3
Male	102	-	-	-	-	-	-	-	-	-	-	2
Female	118	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	209	-	-	-	-	-	-	-	1	-	-	3
Male	96	-	-	-	-	-	-	-	-	-	-	2
Female	113	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	313	1	-	-	-	1	-	-	1	-	4	3
Male	135	-	-	-	-	1	-	-	-	-	4	3
Female	178	1	-	-	-	-	-	-	1	-	-	-
White, both sexes	251	1	-	-	-	1	-	-	1	-	3	2
Male	117	-	-	-	-	1	-	-	-	-	3	2
Female	134	1	-	-	-	-	-	-	1	-	-	-
Black, both sexes	60	-	-	-	-	-	-	-	-	-	1	1
Male	18	-	-	-	-	-	-	-	-	-	1	1
Female	42	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Minnesota												
All races, both sexes	163	-	1	1	-	-	1	1	-	1	2	1
Male	70	-	1	1	-	-	1	-	-	-	2	1
Female	93	-	-	-	-	-	-	1	-	1	-	-
White, both sexes	154	-	1	1	-	-	1	1	-	1	1	1
Male	68	-	1	1	-	-	1	-	-	-	1	1
Female	86	-	-	-	-	-	-	1	-	1	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	75	-	-	-	-	-	-	-	1	-	-	2
Male	30	-	-	-	-	-	-	-	1	-	-	2
Female	45	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	51	-	-	-	-	-	-	-	1	-	-	-
Male	19	-	-	-	-	-	-	-	1	-	-	-
Female	32	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	23	-	-	-	-	-	-	-	-	-	-	2
Male	11	-	-	-	-	-	-	-	-	-	-	2
Female	12	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Missouri												
All races, both sexes	222	-	-	-	-	1	-	-	1	1	1	3
Male	93	-	-	-	-	-	-	-	1	1	-	3
Female	129	-	-	-	-	1	-	-	-	-	1	-
White, both sexes	186	-	-	-	-	1	-	-	1	1	1	3
Male	75	-	-	-	-	-	-	-	1	1	-	3
Female	111	-	-	-	-	1	-	-	-	-	1	-
Black, both sexes	32	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	39	-	-	-	-	-	-	-	1	1	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	1	1	-	-
White, both sexes	37	-	-	-	-	-	-	-	1	1	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Nebraska												
All races, both sexes	65	-	-	-	-	-	-	-	-	-	-	-
Male	32	-	-	-	-	-	-	-	-	-	-	-
Female	33	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	65	-	-	-	-	-	-	-	-	-	-	-
Male	32	-	-	-	-	-	-	-	-	-	-	-
Female	33	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	55	-	-	-	-	-	1	-	-	-	-	1
Male	28	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	1	-	-	-	-	1
White, both sexes	49	-	-	-	-	-	1	-	-	-	-	1
Male	25	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	1	-	-	-	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
New Hampshire												
All races, both sexes	53	-	-	-	-	-	-	-	-	-	-	-
Male	26	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	52	-	-	-	-	-	-	-	-	-	-	-
Male	25	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	295	-	-	-	-	1	1	1	-	1	4	3
Male	130	-	-	-	-	1	-	-	-	1	2	2
Female	165	-	-	-	-	-	1	1	-	-	2	1
White, both sexes	256	-	-	-	-	1	1	1	-	1	2	2
Male	117	-	-	-	-	1	-	-	-	1	-	2
Female	139	-	-	-	-	-	1	1	-	-	2	-
Black, both sexes	36	-	-	-	-	-	-	-	-	-	2	1
Male	11	-	-	-	-	-	-	-	-	-	2	-
Female	25	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
New Mexico												
All races, both sexes	63	-	-	-	-	-	-	-	1	-	1	1
Male	32	-	-	-	-	-	-	-	-	-	1	-
Female	31	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	60	-	-	-	-	-	-	-	1	-	1	1
Male	30	-	-	-	-	-	-	-	-	-	1	-
Female	30	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	447	1	-	-	-	1	2	2	1	3	1	2
Male	202	1	-	-	-	-	1	1	-	3	1	2
Female	245	-	-	-	-	1	1	1	1	-	-	-
White, both sexes	392	-	-	-	-	-	1	1	-	3	1	2
Male	178	-	-	-	-	-	-	1	-	3	1	2
Female	214	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	48	1	-	-	-	1	-	1	1	-	-	-
Male	20	1	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	-	1	-	1	1	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	1	-	-	-	-	-
Male	3	-	-	-	-	-	1	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
North Carolina												
All races, both sexes	293	-	-	-	-	-	1	1	1	2	3	5
Male	142	-	-	-	-	-	1	-	-	1	2	3
Female	151	-	-	-	-	-	-	1	1	1	1	2
White, both sexes	209	-	-	-	-	-	1	-	-	1	1	4
Male	104	-	-	-	-	-	1	-	-	-	1	2
Female	105	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	77	-	-	-	-	-	-	1	1	1	2	1
Male	33	-	-	-	-	-	-	-	-	1	1	1
Female	44	-	-	-	-	-	-	1	1	-	1	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Ohio												
All races, both sexes	465	-	-	-	-	-	-	-	-	1	4	10
Male	185	-	-	-	-	-	-	-	-	1	4	6
Female	280	-	-	-	-	-	-	-	-	-	-	4
White, both sexes	407	-	-	-	-	-	-	-	-	1	4	7
Male	152	-	-	-	-	-	-	-	-	1	4	4
Female	255	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes	58	-	-	-	-	-	-	-	-	-	-	3
Male	33	-	-	-	-	-	-	-	-	-	-	2
Female	25	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	86	-	-	-	-	-	-	-	-	-	3	-
Male	37	-	-	-	-	-	-	-	-	-	-	-
Female	49	-	-	-	-	-	-	-	-	-	3	-
White, both sexes	76	-	-	-	-	-	-	-	-	-	3	-
Male	30	-	-	-	-	-	-	-	-	-	-	-
Female	46	-	-	-	-	-	-	-	-	-	3	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Oregon												
All races, both sexes	100	-	-	-	-	1	-	1	-	-	-	2
Male	43	-	-	-	-	-	-	1	-	-	-	1
Female	57	-	-	-	-	1	-	-	-	-	-	1
White, both sexes	95	-	-	-	-	1	-	1	-	-	-	2
Male	39	-	-	-	-	-	-	1	-	-	-	1
Female	56	-	-	-	-	1	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	483	1	-	-	-	-	-	-	-	2	1	7
Male	214	-	-	-	-	-	-	-	-	1	-	5
Female	269	1	-	-	-	-	-	-	-	1	1	2
White, both sexes	434	-	-	-	-	-	-	-	-	1	-	4
Male	193	-	-	-	-	-	-	-	-	1	-	3
Female	241	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	45	-	-	-	-	-	-	-	-	1	1	3
Male	20	-	-	-	-	-	-	-	-	-	-	2
Female	25	-	-	-	-	-	-	-	-	1	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Rhode Island												
All races, both sexes	29	1	-	-	-	-	-	-	-	-	-	-
Male	13	1	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	28	1	-	-	-	-	-	-	-	-	-	-
Male	12	1	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	161	-	-	-	-	-	-	-	-	1	3	-
Male	78	-	-	-	-	-	-	-	-	-	-	-
Female	83	-	-	-	-	-	-	-	-	1	3	-
White, both sexes	110	-	-	-	-	-	-	-	-	1	1	-
Male	53	-	-	-	-	-	-	-	-	-	-	-
Female	57	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	51	-	-	-	-	-	-	-	-	-	2	-
Male	25	-	-	-	-	-	-	-	-	-	-	-
Female	26	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
South Dakota												
All races, both sexes	38	-	-	-	-	-	-	-	-	-	-	1
Male	16	-	-	-	-	-	-	-	-	-	-	1
Female	22	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	32	-	-	-	-	-	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	246	1	-	-	-	-	-	-	-	-	2	4
Male	110	1	-	-	-	-	-	-	-	-	1	3
Female	136	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	201	-	-	-	-	-	-	-	-	-	1	4
Male	86	-	-	-	-	-	-	-	-	-	1	3
Female	115	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	43	1	-	-	-	-	-	-	-	-	1	-
Male	23	1	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Texas												
All races, both sexes	599	1	-	-	-	-	2	1	2	3	8	11
Male	267	-	-	-	-	-	1	-	1	1	6	6
Female	332	1	-	-	-	-	1	1	1	2	2	5
White, both sexes	494	1	-	-	-	-	1	1	2	2	6	8
Male	216	-	-	-	-	-	-	-	1	1	5	5
Female	278	1	-	-	-	-	1	1	1	1	1	3
Black, both sexes	99	-	-	-	-	-	1	-	-	1	2	2
Male	48	-	-	-	-	-	1	-	-	-	1	1
Female	51	-	-	-	-	-	-	-	-	1	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Utah												
All races, both sexes	54	1	-	-	-	-	-	-	-	1	2	1
Male	27	-	-	-	-	-	-	-	-	1	2	1
Female	27	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	54	1	-	-	-	-	-	-	-	1	2	1
Male	27	-	-	-	-	-	-	-	-	1	2	1
Female	27	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Vermont												
All races, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	201	-	-	-	-	1	-	-	1	-	3	2
Male	79	-	-	-	-	1	-	-	1	-	-	-
Female	122	-	-	-	-	-	-	-	-	-	3	2
White, both sexes	134	-	-	-	-	1	-	-	-	-	2	1
Male	55	-	-	-	-	1	-	-	-	-	-	-
Female	79	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	64	-	-	-	-	-	-	-	1	-	1	1
Male	24	-	-	-	-	-	-	-	1	-	-	-
Female	40	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Washington												
All races, both sexes	205	1	-	-	-	-	-	-	-	1	2	7
Male	97	-	-	-	-	-	-	-	-	1	-	4
Female	108	1	-	-	-	-	-	-	-	-	2	3
White, both sexes	187	-	-	-	-	-	-	-	-	1	1	5
Male	93	-	-	-	-	-	-	-	-	1	-	3
Female	94	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	2
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	3	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
West Virginia												
All races, both sexes	65	-	-	-	-	-	1	-	-	1	-	2
Male	25	-	-	-	-	-	-	-	-	-	-	1
Female	40	-	-	-	-	-	1	-	-	1	-	1
White, both sexes	61	-	-	-	-	-	-	-	-	1	-	2
Male	24	-	-	-	-	-	-	-	-	-	-	1
Female	37	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	4	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Wisconsin												
All races, both sexes	194	-	-	-	-	-	1	-	1	-	2	1
Male	83	-	-	-	-	-	1	-	1	-	2	-
Female	111	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	184	-	-	-	-	-	1	-	1	-	2	1
Male	77	-	-	-	-	-	1	-	1	-	2	-
Female	107	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	2	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	2	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
United States												
All races, both sexes	215	326	487	623	826	1,180	1,548	1,622	1,197	495	101	2
Male	122	201	286	353	423	584	688	615	344	111	14	2
Female	93	125	201	270	403	596	860	1,007	853	384	87	-
White, both sexes	149	239	387	501	680	1,001	1,334	1,432	1,091	425	84	1
Male	85	149	230	282	353	500	599	543	318	97	12	1
Female	64	90	157	219	327	501	735	889	773	328	72	-
Black, both sexes	57	81	87	106	130	154	185	171	94	61	17	1
Male	34	47	49	61	63	73	74	65	20	13	2	1
Female	23	34	38	45	67	81	111	106	74	48	15	-
American Indian 1/, both sexes	3	3	9	9	6	7	5	3	2	1	-	-
Male	1	3	4	5	2	3	3	1	1	-	-	-
Female	2	-	5	4	4	4	2	2	1	1	-	-
Asian or Pacific Islander 2/, both sexes	6	3	4	7	10	18	24	16	10	8	-	-
Male	2	2	3	5	5	8	12	6	5	1	-	-
Female	4	1	1	2	5	10	12	10	5	7	-	-
Alabama												
All races, both sexes	6	11	6	11	14	24	26	26	16	5	4	-
Male	3	7	2	9	5	13	14	9	2	1	2	-
Female	3	4	4	2	9	11	12	17	14	4	2	-
White, both sexes	4	9	4	9	11	18	25	19	11	2	2	-
Male	2	6	2	7	3	10	14	8	1	-	1	-
Female	2	3	2	2	8	8	11	11	10	2	1	-
Black, both sexes	2	1	2	2	3	6	1	7	5	3	2	-
Male	1	-	-	2	2	3	-	1	1	1	1	-
Female	1	1	2	-	1	3	1	6	4	2	1	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Alaska												
All races, both sexes	3	2	-	-	1	2	-	1	2	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	2	-	-	1	2	-	1	2	-	-	-
White, both sexes	2	2	-	-	1	2	-	1	1	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	2	-	-	1	2	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	3	8	15	12	20	19	24	25	12	5	1	-
Male	2	5	6	7	9	11	14	8	2	1	-	-
Female	1	3	9	5	11	8	10	17	10	4	1	-
White, both sexes	3	8	14	10	19	18	24	24	11	5	1	-
Male	2	5	6	5	9	10	14	8	2	1	-	-
Female	1	3	8	5	10	8	10	16	9	4	1	-
Black, both sexes	-	-	1	1	-	1	-	-	1	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Arkansas												
All races, both sexes	3	5	7	12	8	19	19	25	11	3	2	-
Male	1	4	1	6	5	10	11	8	3	1	-	-
Female	2	1	6	6	3	9	8	17	8	2	2	-
White, both sexes	3	4	6	11	6	16	17	22	11	2	1	-
Male	1	3	1	6	4	8	10	6	3	1	-	-
Female	2	1	5	5	2	8	7	16	8	1	1	-
Black, both sexes	-	1	1	1	2	3	2	3	-	1	1	-
Male	-	1	-	-	1	2	1	2	-	-	-	-
Female	-	-	1	1	1	1	1	1	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	23	31	44	56	63	98	133	134	100	52	4	-
Male	13	22	32	33	30	48	56	55	26	12	1	-
Female	10	9	12	23	33	50	77	79	74	40	3	-
White, both sexes	16	27	38	42	53	81	110	114	91	43	2	-
Male	10	19	27	25	23	38	44	48	22	10	1	-
Female	6	8	11	17	30	43	66	66	69	33	1	-
Black, both sexes	3	4	3	11	7	7	13	13	6	3	2	-
Male	1	3	3	5	5	5	6	4	2	1	-	-
Female	2	1	-	6	2	2	7	9	4	2	2	-
American Indian 1/, both sexes	-	-	1	-	1	1	2	-	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	2	3	2	9	8	7	3	6	-	-
Male	2	-	1	3	2	5	5	3	2	1	-	-
Female	2	-	1	-	-	4	3	4	1	5	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Colorado												
All races, both sexes	4	5	5	4	14	16	16	15	9	4	1	-
Male	2	3	-	2	9	9	6	5	-	1	-	-
Female	2	2	5	2	5	7	10	10	9	3	1	-
White, both sexes	2	3	5	4	13	15	14	13	9	4	1	-
Male	1	2	-	2	8	8	5	3	-	1	-	-
Female	1	1	5	2	5	7	9	10	9	3	1	-
Black, both sexes	2	2	-	-	1	1	2	1	-	-	-	-
Male	1	1	-	-	1	1	1	1	-	-	-	-
Female	1	1	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	1	6	3	11	13	17	26	26	11	2	-
Male	-	-	1	2	8	7	7	10	6	2	-	-
Female	-	1	5	1	3	6	10	16	20	9	2	-
White, both sexes	-	1	5	2	11	12	17	21	26	11	2	-
Male	-	-	1	2	8	7	7	8	6	2	-	-
Female	-	1	4	-	3	5	10	13	20	9	2	-
Black, both sexes	-	-	1	1	-	1	-	5	-	-	-	-
Male	-	-	-	-	-	-	-	2	-	-	-	-
Female	-	-	1	1	-	1	-	3	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Delaware												
All races, both sexes	1	-	-	3	2	7	7	8	1	1	-	-
Male	1	-	-	1	-	1	2	3	1	-	-	-
Female	-	-	-	2	2	6	5	5	-	1	-	-
White, both sexes	1	-	-	2	1	6	5	7	1	1	-	-
Male	1	-	-	1	-	1	1	3	1	-	-	-
Female	-	-	-	1	1	5	4	4	-	1	-	-
Black, both sexes	-	-	-	1	1	1	2	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	1	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	2	1	2	2	2	4	2	-	-	-
Male	1	-	-	-	1	-	-	2	-	-	-	-
Female	-	-	2	1	1	2	2	2	2	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	2	1	1	2	2	4	2	-	-	-
Male	1	-	-	-	-	-	-	2	-	-	-	-
Female	-	-	2	1	1	2	2	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Florida												
All races, both sexes	19	25	40	41	49	76	114	98	80	23	10	-
Male	12	19	27	27	23	39	46	44	29	9	-	-
Female	7	6	13	14	26	37	68	54	51	14	10	-
White, both sexes	12	20	35	32	39	72	101	88	77	23	7	-
Male	8	16	25	21	16	37	43	40	29	9	-	-
Female	4	4	10	11	23	35	58	48	48	14	7	-
Black, both sexes	7	5	5	8	10	4	13	10	3	-	3	-
Male	4	3	2	6	7	2	3	4	-	-	-	-
Female	3	2	3	2	3	2	10	6	3	-	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	5	11	12	17	20	35	38	35	24	19	6	-
Male	4	6	7	9	10	19	20	9	9	2	-	-
Female	1	5	5	8	10	16	18	26	15	17	6	-
White, both sexes	2	3	5	8	14	21	31	24	17	11	5	-
Male	2	2	3	2	7	13	15	5	5	1	-	-
Female	-	1	2	6	7	8	16	19	12	10	5	-
Black, both sexes	3	8	7	8	6	14	7	11	7	8	1	-
Male	2	4	4	6	3	6	5	4	4	1	-	-
Female	1	4	3	2	3	8	2	7	3	7	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Hawaii												
All races, both sexes	1	1	1	2	1	1	8	1	3	3	-	-
Male	-	1	1	1	1	1	3	1	2	-	-	-
Female	1	-	-	1	-	-	5	-	1	3	-	-
White, both sexes	-	-	-	1	-	1	2	1	-	1	-	-
Male	-	-	-	1	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	1	1	-	6	-	3	2	-	-
Male	-	1	1	-	1	-	3	-	2	-	-	-
Female	1	-	-	1	-	-	3	-	1	2	-	-
Idaho												
All races, both sexes	1	1	1	3	3	7	7	6	8	4	-	-
Male	-	1	-	2	-	3	5	1	4	-	-	-
Female	1	-	1	1	3	4	2	5	4	4	-	-
White, both sexes	-	1	1	3	3	6	7	6	8	4	-	-
Male	-	1	-	2	-	3	5	1	4	-	-	-
Female	-	-	1	1	3	3	2	5	4	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Illinois												
All races, both sexes	5	13	12	12	36	50	65	76	45	32	4	-
Male	3	7	8	7	21	22	25	23	10	5	-	-
Female	2	6	4	5	15	28	40	53	35	27	4	-
White, both sexes	4	10	10	10	35	38	51	63	41	25	3	-
Male	2	5	6	6	21	18	19	18	9	4	-	-
Female	2	5	4	4	14	20	32	45	32	21	3	-
Black, both sexes	1	3	2	2	1	9	14	12	4	7	1	-
Male	1	2	2	1	-	3	6	5	1	1	-	-
Female	-	1	-	1	1	6	8	7	3	6	1	-
American Indian 1/, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Indiana												
All races, both sexes	5	5	7	14	21	23	35	42	22	16	2	-
Male	5	3	3	4	13	10	12	16	6	1	-	-
Female	-	2	4	10	8	13	23	26	16	15	2	-
White, both sexes	4	3	5	12	18	22	32	39	21	14	2	-
Male	4	1	3	4	10	9	12	15	6	1	-	-
Female	-	2	2	8	8	13	20	24	15	13	2	-
Black, both sexes	1	2	2	2	3	1	3	3	1	2	-	-
Male	1	2	-	-	3	1	-	1	-	-	-	-
Female	-	-	2	2	-	-	3	2	1	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Iowa												
All races, both sexes	1	-	7	5	5	14	16	24	20	7	1	-
Male	1	-	3	2	3	8	7	10	6	1	-	-
Female	-	-	4	3	2	6	9	14	14	6	1	-
White, both sexes	1	-	7	5	5	14	15	24	20	7	1	-
Male	1	-	3	2	3	8	7	10	6	1	-	-
Female	-	-	4	3	2	6	8	14	14	6	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Kansas												
All races, both sexes	2	1	3	6	9	12	10	14	10	9	1	-
Male	1	1	1	3	7	6	4	8	2	5	1	-
Female	1	-	2	3	2	6	6	6	8	4	-	-
White, both sexes	1	1	2	6	8	11	9	13	10	8	1	-
Male	1	1	1	3	6	6	4	7	2	4	1	-
Female	-	-	1	3	2	5	5	6	8	4	-	-
Black, both sexes	1	-	-	-	1	-	1	1	-	1	-	-
Male	-	-	-	-	1	-	-	1	-	1	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Kentucky												
All races, both sexes	5	7	13	16	20	18	26	28	22	6	4	-
Male	4	3	7	9	8	9	15	10	5	-	-	-
Female	1	4	6	7	12	9	11	18	17	6	4	-
White, both sexes	4	7	12	15	17	16	24	27	22	5	3	-
Male	3	3	7	9	6	8	13	10	5	-	-	-
Female	1	4	5	6	11	8	11	17	17	5	3	-
Black, both sexes	1	-	1	1	3	2	2	1	-	1	1	-
Male	1	-	-	-	2	1	2	-	-	-	-	-
Female	-	-	1	1	1	1	-	1	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	7	9	12	16	19	11	19	22	14	6	2	-
Male	3	5	8	11	4	5	6	8	5	1	-	-
Female	4	4	4	5	15	6	13	14	9	5	2	-
White, both sexes	3	4	6	10	9	7	15	19	11	3	2	-
Male	2	2	5	6	-	3	6	6	3	1	-	-
Female	1	2	1	4	9	4	9	13	8	2	2	-
Black, both sexes	3	5	6	6	9	4	4	3	3	3	-	-
Male	1	3	3	5	3	2	-	2	2	-	-	-
Female	2	2	3	1	6	2	4	1	1	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Maine												
All races, both sexes	2	7	4	5	3	13	7	12	7	1	-	-
Male	2	4	2	3	1	7	5	2	-	1	-	-
Female	-	3	2	2	2	6	2	10	7	-	-	-
White, both sexes	2	7	4	5	3	13	7	12	7	1	-	-
Male	2	4	2	3	1	7	5	2	-	1	-	-
Female	-	3	2	2	2	6	2	10	7	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	6	3	14	6	19	35	33	26	25	9	-	-
Male	3	1	10	4	7	14	15	9	7	4	-	-
Female	3	2	4	2	12	21	18	17	18	5	-	-
White, both sexes	4	2	7	4	12	28	25	21	22	6	-	-
Male	2	-	6	3	5	13	11	8	7	2	-	-
Female	2	2	1	1	7	15	14	13	15	4	-	-
Black, both sexes	2	1	7	2	7	7	8	5	3	3	-	-
Male	1	1	4	1	2	1	4	1	-	2	-	-
Female	1	-	3	1	5	6	4	4	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Massachusetts												
All races, both sexes	1	6	6	10	17	31	54	51	27	12	1	-
Male	-	4	3	7	9	17	23	23	9	4	1	-
Female	1	2	3	3	8	14	31	28	18	8	-	-
White, both sexes	1	5	5	10	15	29	50	50	27	12	1	-
Male	-	3	2	7	8	15	22	23	9	4	1	-
Female	1	2	3	3	7	14	28	27	18	8	-	-
Black, both sexes	-	1	1	-	2	2	4	1	-	-	-	-
Male	-	1	1	-	1	2	1	-	-	-	-	-
Female	-	-	-	-	1	-	3	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	12	6	17	26	32	40	57	53	44	14	2	-
Male	3	5	9	9	19	20	28	18	14	2	-	-
Female	9	1	8	17	13	20	29	35	30	12	2	-
White, both sexes	8	3	11	18	22	33	47	43	43	13	2	-
Male	2	3	6	8	17	18	25	17	13	2	-	-
Female	6	-	5	10	5	15	22	26	30	11	2	-
Black, both sexes	4	3	5	7	10	7	10	10	1	1	-	-
Male	1	2	3	1	2	2	3	1	1	-	-	-
Female	3	1	2	6	8	5	7	9	-	1	-	-
American Indian 1/, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Minnesota												
All races, both sexes	1	4	9	5	15	16	23	38	28	12	4	-
Male	1	3	6	3	8	10	9	15	6	3	-	-
Female	-	1	3	2	7	6	14	23	22	9	4	-
White, both sexes	1	4	8	4	15	16	20	37	26	12	4	-
Male	1	3	6	2	8	10	9	15	6	3	-	-
Female	-	1	2	2	7	6	11	22	20	9	4	-
Black, both sexes	-	-	1	-	-	-	3	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	3	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Mississippi												
All races, both sexes	2	11	5	3	6	4	17	7	10	4	1	2
Male	1	6	2	1	3	3	5	3	1	-	-	2
Female	1	5	3	2	3	1	12	4	9	4	1	-
White, both sexes	-	7	3	3	3	3	15	5	6	4	-	1
Male	-	4	1	1	2	3	4	2	-	-	-	1
Female	-	3	2	2	1	-	11	3	6	4	-	-
Black, both sexes	2	4	2	-	3	1	2	1	4	-	1	1
Male	1	2	1	-	1	-	1	1	1	-	-	1
Female	1	2	1	-	2	1	1	-	3	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Missouri												
All races, both sexes	7	9	11	20	16	31	36	41	31	11	2	-
Male	4	5	7	13	7	18	11	12	9	2	-	-
Female	3	4	4	7	9	13	25	29	22	9	2	-
White, both sexes	6	6	7	18	13	22	30	36	30	9	2	-
Male	4	4	5	11	6	11	9	10	9	1	-	-
Female	2	2	2	7	7	11	21	26	21	8	2	-
Black, both sexes	1	3	4	1	3	6	6	5	1	2	-	-
Male	-	1	2	1	1	5	2	2	-	1	-	-
Female	1	2	2	-	2	1	4	3	1	1	-	-
American Indian 1/, both sexes	-	-	-	1	-	2	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	2	-	3	1	2	2	11	3	8	4	1	-
Male	2	-	1	1	-	2	4	-	5	2	-	-
Female	-	-	2	-	2	-	7	3	3	2	1	-
White, both sexes	1	-	2	1	2	2	11	3	8	4	1	-
Male	1	-	1	1	-	2	4	-	5	2	-	-
Female	-	-	1	-	2	-	7	3	3	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Nebraska												
All races, both sexes	-	1	4	1	7	5	9	17	14	5	2	-
Male	-	1	3	1	3	2	6	10	5	-	1	-
Female	-	-	1	-	4	3	3	7	9	5	1	-
White, both sexes	-	1	4	1	7	5	9	17	14	5	2	-
Male	-	1	3	1	3	2	6	10	5	-	1	-
Female	-	-	1	-	4	3	3	7	9	5	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1	3	4	5	5	9	12	8	5	1	-	-
Male	1	3	2	3	4	5	5	3	1	1	-	-
Female	-	-	2	2	1	4	7	5	4	-	-	-
White, both sexes	1	2	4	4	5	7	11	8	4	1	-	-
Male	1	2	2	3	4	4	4	3	1	1	-	-
Female	-	-	2	1	1	3	7	5	3	-	-	-
Black, both sexes	-	1	-	1	-	2	1	-	-	-	-	-
Male	-	1	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
New Hampshire												
All races, both sexes	-	2	4	3	5	5	9	13	8	3	1	-
Male	-	-	4	3	4	3	2	8	2	-	-	-
Female	-	2	-	-	1	2	7	5	6	3	1	-
White, both sexes	-	2	3	3	5	5	9	13	8	3	1	-
Male	-	-	3	3	4	3	2	8	2	-	-	-
Female	-	2	-	-	1	2	7	5	6	3	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	4	8	12	7	28	38	53	60	49	23	2	-
Male	3	3	6	1	12	20	32	23	17	7	-	-
Female	1	5	6	6	16	18	21	37	32	16	2	-
White, both sexes	3	5	11	5	25	33	43	55	45	21	2	-
Male	2	2	6	1	12	17	30	20	16	7	-	-
Female	1	3	5	4	13	16	13	35	29	14	2	-
Black, both sexes	1	3	1	2	3	5	9	4	3	2	-	-
Male	1	1	-	-	-	3	2	2	-	-	-	-
Female	-	2	1	2	3	2	7	2	3	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
New Mexico												
All races, both sexes	1	-	5	6	5	9	10	12	9	2	1	-
Male	1	-	4	3	2	5	7	4	4	-	1	-
Female	-	-	1	3	3	4	3	8	5	2	-	-
White, both sexes	1	-	4	6	5	9	8	12	9	2	1	-
Male	1	-	4	3	2	5	5	4	4	-	1	-
Female	-	-	-	3	3	4	3	8	5	2	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	9	18	17	27	47	59	83	87	59	24	4	-
Male	6	11	9	17	24	36	36	29	17	8	-	-
Female	3	7	8	10	23	23	47	58	42	16	4	-
White, both sexes	6	14	13	25	42	50	75	80	53	23	3	-
Male	4	10	8	15	21	31	32	26	16	8	-	-
Female	2	4	5	10	21	19	43	54	37	15	3	-
Black, both sexes	3	3	4	2	4	7	6	7	6	1	1	-
Male	2	-	1	2	3	5	2	3	1	-	-	-
Female	1	3	3	-	1	2	4	4	5	1	1	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	2	2	-	-	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	1	2	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
North Carolina												
All races, both sexes	10	15	22	24	40	33	40	49	29	13	5	-
Male	7	8	20	12	17	17	21	17	12	2	2	-
Female	3	7	2	12	23	16	19	32	17	11	3	-
White, both sexes	7	6	17	16	26	28	29	32	25	11	5	-
Male	5	3	15	9	12	14	18	11	10	1	2	-
Female	2	3	2	7	14	14	11	21	15	10	3	-
Black, both sexes	3	9	4	6	13	4	10	16	4	2	-	-
Male	2	5	4	2	4	2	3	5	2	1	-	-
Female	1	4	-	4	9	2	7	11	2	1	-	-
American Indian 1/, both sexes	-	-	-	2	1	1	-	1	-	-	-	-
Male	-	-	-	1	1	1	-	1	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	1	-	1	1	4	4	1	1	-	-
Male	-	-	1	-	-	1	2	1	-	-	-	-
Female	-	-	-	-	1	-	2	3	1	1	-	-
White, both sexes	-	-	-	-	1	1	4	4	1	1	-	-
Male	-	-	-	-	-	1	2	1	-	-	-	-
Female	-	-	-	-	1	-	2	3	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Ohio												
All races, both sexes	10	13	26	29	38	63	86	96	63	23	3	-
Male	4	10	13	17	18	25	33	35	13	5	1	-
Female	6	3	13	12	20	38	53	61	50	18	2	-
White, both sexes	9	8	22	22	33	52	76	93	57	20	3	-
Male	3	6	9	12	16	18	27	34	12	5	1	-
Female	6	2	13	10	17	34	49	59	45	15	2	-
Black, both sexes	1	5	4	7	5	11	10	3	6	3	-	-
Male	1	4	4	5	2	7	6	1	1	-	-	-
Female	-	1	-	2	3	4	4	2	5	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1	1	7	11	7	12	15	16	10	3	-	-
Male	1	1	5	7	5	4	4	5	4	1	-	-
Female	-	-	2	4	2	8	11	11	6	2	-	-
White, both sexes	1	1	3	10	5	11	14	16	9	3	-	-
Male	1	1	2	6	4	4	3	5	3	1	-	-
Female	-	-	1	4	1	7	11	11	6	2	-	-
Black, both sexes	-	-	3	-	1	1	-	-	-	-	-	-
Male	-	-	3	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	1	1	-	1	-	1	-	-	-
Male	-	-	-	1	-	-	1	-	1	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Oregon												
All races, both sexes	4	1	5	8	10	7	17	21	18	5	-	-
Male	2	-	4	5	3	5	11	4	6	1	-	-
Female	2	1	1	3	7	2	6	17	12	4	-	-
White, both sexes	4	1	4	5	10	7	17	20	18	5	-	-
Male	2	-	3	3	3	5	11	3	6	1	-	-
Female	2	1	1	2	7	2	6	17	12	4	-	-
Black, both sexes	-	-	1	2	-	-	-	-	-	-	-	-
Male	-	-	1	2	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	5	10	19	28	41	54	83	104	92	28	8	-
Male	5	5	12	17	24	15	40	48	31	9	2	-
Female	-	5	7	11	17	39	43	56	61	19	6	-
White, both sexes	4	8	17	24	36	48	72	96	89	28	7	-
Male	4	5	11	15	21	13	35	44	31	9	1	-
Female	-	3	6	9	15	35	37	52	58	19	6	-
Black, both sexes	1	1	2	4	5	6	9	8	3	-	1	-
Male	1	-	1	2	3	2	4	4	-	-	1	-
Female	-	1	1	2	2	4	5	4	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Rhode Island												
All races, both sexes	1	3	-	3	4	6	3	5	3	-	-	-
Male	-	2	-	-	2	2	2	3	1	-	-	-
Female	1	1	-	3	2	4	1	2	2	-	-	-
White, both sexes	1	2	-	3	4	6	3	5	3	-	-	-
Male	-	1	-	-	2	2	2	3	1	-	-	-
Female	1	1	-	3	2	4	1	2	2	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	4	5	13	18	19	14	22	31	19	11	1	-
Male	2	5	5	15	16	10	11	8	3	3	-	-
Female	2	-	8	3	3	4	11	23	16	8	1	-
White, both sexes	1	4	10	9	14	11	12	24	15	8	-	-
Male	-	4	4	8	13	8	5	6	3	2	-	-
Female	1	-	6	1	1	3	7	18	12	6	-	-
Black, both sexes	3	1	3	9	5	3	10	7	4	3	1	-
Male	2	1	1	7	3	2	6	2	-	1	-	-
Female	1	-	2	2	2	1	4	5	4	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
South Dakota												
All races, both sexes	2	1	1	-	6	3	9	7	3	5	-	-
Male	-	1	-	-	3	2	3	4	1	1	-	-
Female	2	-	1	-	3	1	6	3	2	4	-	-
White, both sexes	2	-	1	-	4	3	8	6	3	4	-	-
Male	-	-	-	-	3	2	3	4	1	1	-	-
Female	2	-	1	-	1	1	5	2	2	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	2	-	-	1	-	1	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	-	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Tennessee												
All races, both sexes	5	11	21	23	20	41	41	34	27	14	2	-
Male	2	10	13	13	10	17	20	12	7	1	-	-
Female	3	1	8	10	10	24	21	22	20	13	2	-
White, both sexes	4	8	17	19	17	36	35	27	19	12	2	-
Male	1	7	10	9	9	15	17	9	4	1	-	-
Female	3	1	7	10	8	21	18	18	15	11	2	-
Black, both sexes	1	3	4	4	3	4	6	7	7	2	-	-
Male	1	3	3	4	1	1	3	3	3	-	-	-
Female	-	-	1	-	2	3	3	4	4	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Texas												
All races, both sexes	13	24	26	62	59	92	100	91	70	27	7	-
Male	7	14	15	30	33	49	45	37	18	4	-	-
Female	6	10	11	32	26	43	55	54	52	23	7	-
White, both sexes	7	18	24	51	46	75	90	79	58	20	5	-
Male	3	10	14	24	23	40	40	30	18	2	-	-
Female	4	8	10	27	23	35	50	49	40	18	5	-
Black, both sexes	6	5	2	11	11	16	10	12	11	7	2	-
Male	4	3	1	6	8	9	5	7	-	2	-	-
Female	2	2	1	5	3	7	5	5	11	5	2	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	1	-	-	1	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
Utah												
All races, both sexes	2	1	-	3	4	7	10	10	10	1	1	-
Male	1	-	-	3	2	2	4	6	3	1	1	-
Female	1	1	-	-	2	5	6	4	7	-	-	-
White, both sexes	2	1	-	3	4	7	10	10	10	1	1	-
Male	1	-	-	3	2	2	4	6	3	1	1	-
Female	1	1	-	-	2	5	6	4	7	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Vermont												
All races, both sexes	1	1	-	-	2	1	5	5	4	2	1	-
Male	1	-	-	-	2	1	1	3	1	-	1	-
Female	-	1	-	-	-	-	4	2	3	2	-	-
White, both sexes	1	1	-	-	2	1	4	5	4	2	1	-
Male	1	-	-	-	2	1	1	3	1	-	1	-
Female	-	1	-	-	-	-	3	2	3	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	7	10	11	14	18	27	41	28	27	7	4	-
Male	4	3	7	9	8	8	17	11	8	2	-	-
Female	3	7	4	5	10	19	24	17	19	5	4	-
White, both sexes	3	5	5	12	11	16	31	19	22	2	4	-
Male	1	1	2	8	6	7	14	7	7	1	-	-
Female	2	4	3	4	5	9	17	12	15	1	4	-
Black, both sexes	4	5	6	2	6	11	10	8	4	5	-	-
Male	3	2	5	1	2	1	3	4	1	1	-	-
Female	1	3	1	1	4	10	7	4	3	4	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Washington												
All races, both sexes	4	9	10	21	14	37	32	29	29	8	1	-
Male	1	6	8	11	7	22	13	14	8	2	-	-
Female	3	3	2	10	7	15	19	15	21	6	1	-
White, both sexes	4	9	10	20	12	32	29	27	29	7	1	-
Male	1	6	8	11	7	20	12	14	8	2	-	-
Female	3	3	2	9	5	12	17	13	21	5	1	-
Black, both sexes	-	-	-	-	-	3	1	1	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	2	1	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	1	2	1	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	2	-	1	1	-	-	-	-
West Virginia												
All races, both sexes	1	5	6	5	7	8	9	9	9	2	-	-
Male	-	1	4	3	5	3	5	1	2	-	-	-
Female	1	4	2	2	2	5	4	8	7	2	-	-
White, both sexes	1	4	6	5	7	8	8	8	9	2	-	-
Male	-	1	4	3	5	3	4	1	2	-	-	-
Female	1	3	2	2	2	5	4	7	7	2	-	-
Black, both sexes	-	1	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Wisconsin												
All races, both sexes	2	2	9	15	9	28	32	45	31	13	3	-
Male	-	2	3	7	6	17	14	17	11	2	-	-
Female	2	-	6	8	3	11	18	28	20	11	3	-
White, both sexes	2	2	9	13	8	26	30	44	29	13	3	-
Male	-	2	3	5	5	15	13	17	11	2	-	-
Female	2	-	6	8	3	11	17	27	18	11	3	-
Black, both sexes	-	-	-	1	1	2	2	-	2	-	-	-
Male	-	-	-	1	1	2	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Wyoming												
All races, both sexes	-	-	2	-	2	2	3	-	1	1	-	-
Male	-	-	1	-	2	1	1	-	-	1	-	-
Female	-	-	1	-	-	1	2	-	1	-	-	-
White, both sexes	-	-	1	-	2	2	3	-	1	1	-	-
Male	-	-	-	-	2	1	1	-	-	1	-	-
Female	-	-	1	-	-	1	2	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
United States												
All races, both sexes	4,101	53	3	3	3	16	24	51	69	132	190	281
Male	1,902	31	1	2	2	7	15	22	45	74	114	162
Female	2,199	22	2	1	1	9	9	29	24	58	76	119
White, both sexes	3,336	40	3	2	3	12	15	31	48	84	133	187
Male	1,523	22	1	2	2	4	6	15	32	46	85	110
Female	1,813	18	2	-	1	8	9	16	16	38	48	77
Black, both sexes	701	12	-	-	-	4	7	19	21	48	53	87
Male	348	9	-	-	-	3	7	7	13	28	26	47
Female	353	3	-	-	-	1	-	12	8	20	27	40
American Indian 1/, both sexes	17	-	-	-	-	-	1	-	-	-	2	3
Male	6	-	-	-	-	-	1	-	-	-	2	1
Female	11	-	-	-	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	47	1	-	1	-	-	1	1	-	-	2	4
Male	25	-	-	-	-	-	1	-	-	-	1	4
Female	22	1	-	1	-	-	-	1	-	-	1	-
Alabama												
All races, both sexes	66	1	-	-	-	-	-	1	-	1	3	4
Male	32	1	-	-	-	-	-	1	-	-	2	3
Female	34	-	-	-	-	-	-	-	-	1	1	1
White, both sexes	52	1	-	-	-	-	-	1	-	1	1	3
Male	24	1	-	-	-	-	-	1	-	-	1	2
Female	28	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	13	-	-	-	-	-	-	-	-	-	2	1
Male	7	-	-	-	-	-	-	-	-	-	1	1
Female	6	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Alaska												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	88	2	-	-	-	-	-	1	2	4	6	9
Male	36	1	-	-	-	-	-	1	1	1	3	4
Female	52	1	-	-	-	-	-	-	1	3	3	5
White, both sexes	81	2	-	-	-	-	-	1	2	3	4	8
Male	33	1	-	-	-	-	-	1	1	-	2	4
Female	48	1	-	-	-	-	-	-	1	3	2	4
Black, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	1	1
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Arkansas												
All races, both sexes	48	2	-	-	-	-	-	1	1	1	2	1
Male	24	2	-	-	-	-	-	1	-	1	2	-
Female	24	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	36	1	-	-	-	-	-	1	-	-	2	-
Male	17	1	-	-	-	-	-	1	-	-	2	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	12	1	-	-	-	-	-	-	1	1	-	1
Male	7	1	-	-	-	-	-	-	-	1	-	-
Female	5	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	426	7	-	-	-	2	6	4	9	17	17	42
Male	182	4	-	-	-	2	3	2	7	8	13	21
Female	244	3	-	-	-	-	3	2	2	9	4	21
White, both sexes	346	5	-	-	-	1	4	2	5	12	13	28
Male	139	2	-	-	-	1	1	2	5	5	10	12
Female	207	3	-	-	-	-	3	-	-	7	3	16
Black, both sexes	59	2	-	-	-	1	1	1	4	5	4	11
Male	35	2	-	-	-	1	1	-	2	3	3	6
Female	24	-	-	-	-	-	-	1	2	2	1	5
American Indian 1/, both sexes	3	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	18	-	-	-	-	-	-	1	-	-	-	3
Male	7	-	-	-	-	-	-	-	-	-	-	3
Female	11	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Colorado												
All races, both sexes	54	1	-	-	-	-	-	-	-	3	5	5
Male	29	1	-	-	-	-	-	-	-	1	2	3
Female	25	-	-	-	-	-	-	-	-	2	3	2
White, both sexes	45	1	-	-	-	-	-	-	-	1	3	2
Male	25	1	-	-	-	-	-	-	-	-	1	1
Female	20	-	-	-	-	-	-	-	-	1	2	1
Black, both sexes	9	-	-	-	-	-	-	-	-	2	2	3
Male	4	-	-	-	-	-	-	-	-	1	1	2
Female	5	-	-	-	-	-	-	-	-	1	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	34	-	-	-	-	-	-	1	-	-	-	3
Male	15	-	-	-	-	-	-	-	-	-	-	3
Female	19	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	32	-	-	-	-	-	-	1	-	-	-	3
Male	14	-	-	-	-	-	-	-	-	-	-	3
Female	18	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Delaware												
All races, both sexes	12	3	-	-	-	-	-	-	-	-	-	-
Male	5	1	-	-	-	-	-	-	-	-	-	-
Female	7	2	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	3	-	-	-	-	-	-	-	-	-	-
Male	5	1	-	-	-	-	-	-	-	-	-	-
Female	7	2	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	2
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	2
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Florida												
All races, both sexes	266	2	-	-	-	1	2	1	5	11	12	18
Male	137	2	-	-	-	-	2	1	5	7	9	6
Female	129	-	-	-	-	1	-	-	-	4	3	12
White, both sexes	218	1	-	-	-	1	1	1	1	6	9	9
Male	112	1	-	-	-	-	1	1	1	3	7	4
Female	106	-	-	-	-	1	-	-	-	3	2	5
Black, both sexes	47	1	-	-	-	-	1	-	4	5	3	9
Male	24	1	-	-	-	-	1	-	4	4	2	2
Female	23	-	-	-	-	-	-	-	-	1	1	7
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	123	3	-	-	1	-	2	3	-	4	9	6
Male	54	2	-	-	-	-	2	1	-	3	5	1
Female	69	1	-	-	1	-	-	2	-	1	4	5
White, both sexes	80	-	-	-	1	-	-	1	-	1	6	3
Male	32	-	-	-	-	-	-	-	-	-	4	-
Female	48	-	-	-	1	-	-	1	-	1	2	3
Black, both sexes	43	3	-	-	-	-	2	2	-	3	3	3
Male	22	2	-	-	-	-	2	1	-	3	1	1
Female	21	1	-	-	-	-	-	1	-	-	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Hawaii												
All races, both sexes	19	-	-	-	-	-	-	1	-	1	2	2
Male	10	-	-	-	-	-	-	1	-	1	1	1
Female	9	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	9	-	-	-	-	-	-	-	-	1	2	1
Male	3	-	-	-	-	-	-	-	-	1	1	-
Female	6	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	28	1	1	-	-	-	-	1	1	2	-	2
Male	15	1	-	-	-	-	-	1	1	1	-	1
Female	13	-	1	-	-	-	-	-	-	1	-	1
White, both sexes	27	1	1	-	-	-	-	1	1	2	-	2
Male	14	1	-	-	-	-	-	1	1	1	-	1
Female	13	-	1	-	-	-	-	-	-	1	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Illinois												
All races, both sexes	198	2	-	1	-	-	1	1	5	8	16	18
Male	89	1	-	-	-	-	1	-	4	3	7	15
Female	109	1	-	1	-	-	-	1	1	5	9	3
White, both sexes	140	2	-	-	-	-	-	-	3	2	8	9
Male	60	1	-	-	-	-	-	-	2	1	5	7
Female	80	1	-	-	-	-	-	-	1	1	3	2
Black, both sexes	55	-	-	-	-	-	1	1	2	6	7	9
Male	27	-	-	-	-	-	1	-	2	2	1	8
Female	28	-	-	-	-	-	-	1	-	4	6	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	1	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	64	2	-	-	-	-	-	2	1	-	-	3
Male	28	2	-	-	-	-	-	1	1	-	-	2
Female	36	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	59	2	-	-	-	-	-	2	1	-	-	2
Male	26	2	-	-	-	-	-	1	1	-	-	1
Female	33	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Iowa												
All races, both sexes	42	1	-	-	-	-	-	-	-	1	3	-
Male	18	-	-	-	-	-	-	-	-	-	2	-
Female	24	1	-	-	-	-	-	-	-	1	1	-
White, both sexes	41	1	-	-	-	-	-	-	-	1	3	-
Male	17	-	-	-	-	-	-	-	-	-	2	-
Female	24	1	-	-	-	-	-	-	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	36	-	-	-	-	-	-	1	1	1	1	2
Male	11	-	-	-	-	-	-	-	1	1	-	2
Female	25	-	-	-	-	-	-	1	-	-	1	-
White, both sexes	34	-	-	-	-	-	-	1	1	1	-	2
Male	11	-	-	-	-	-	-	-	1	1	-	2
Female	23	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Kentucky												
All races, both sexes	62	-	-	-	-	1	-	2	2	2	2	8
Male	27	-	-	-	-	-	-	2	2	1	1	5
Female	35	-	-	-	-	1	-	-	-	1	1	3
White, both sexes	55	-	-	-	-	1	-	2	2	2	1	8
Male	24	-	-	-	-	-	-	2	2	1	-	5
Female	31	-	-	-	-	1	-	-	-	1	1	3
Black, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	53	1	-	-	-	-	-	1	-	1	2	2
Male	21	1	-	-	-	-	-	1	-	1	1	1
Female	32	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	39	1	-	-	-	-	-	-	-	-	1	1
Male	13	1	-	-	-	-	-	-	-	-	1	-
Female	26	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	14	-	-	-	-	-	-	1	-	1	1	1
Male	8	-	-	-	-	-	-	1	-	1	-	1
Female	6	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Maine												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	2	-
Male	6	-	-	-	-	-	-	-	-	-	2	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	2	-
Male	6	-	-	-	-	-	-	-	-	-	2	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	79	-	-	-	-	-	1	1	3	8	4	4
Male	44	-	-	-	-	-	1	-	1	4	3	-
Female	35	-	-	-	-	-	-	1	2	4	1	4
White, both sexes	45	-	-	-	-	-	-	-	1	-	2	1
Male	29	-	-	-	-	-	-	-	1	-	1	-
Female	16	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	34	-	-	-	-	-	1	1	2	8	2	3
Male	15	-	-	-	-	-	1	-	-	4	2	-
Female	19	-	-	-	-	-	-	1	2	4	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Massachusetts												
All races, both sexes	79	-	-	-	-	2	-	-	2	5	4	6
Male	41	-	-	-	-	1	-	-	-	3	2	4
Female	38	-	-	-	-	1	-	-	2	2	2	2
White, both sexes	66	-	-	-	-	1	-	-	2	4	2	5
Male	34	-	-	-	-	-	-	-	-	3	1	4
Female	32	-	-	-	-	1	-	-	2	1	1	1
Black, both sexes	12	-	-	-	-	1	-	-	-	1	2	1
Male	6	-	-	-	-	1	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	1	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	183	-	-	-	-	4	1	3	3	7	10	16
Male	89	-	-	-	-	1	1	1	2	4	5	7
Female	94	-	-	-	-	3	-	2	1	3	5	9
White, both sexes	141	-	-	-	-	4	1	1	3	2	6	11
Male	71	-	-	-	-	1	1	-	2	1	4	5
Female	70	-	-	-	-	3	-	1	1	1	2	6
Black, both sexes	42	-	-	-	-	-	-	2	-	5	4	5
Male	18	-	-	-	-	-	-	1	-	3	1	2
Female	24	-	-	-	-	-	-	1	-	2	3	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Minnesota												
All races, both sexes	43	5	-	-	-	-	-	-	1	-	-	3
Male	18	1	-	-	-	-	-	-	1	-	-	2
Female	25	4	-	-	-	-	-	-	-	-	-	1
White, both sexes	40	5	-	-	-	-	-	-	1	-	-	2
Male	16	1	-	-	-	-	-	-	1	-	-	1
Female	24	4	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	40	-	-	-	-	1	-	-	1	2	1	4
Male	19	-	-	-	-	-	-	-	1	1	-	4
Female	21	-	-	-	-	1	-	-	-	1	1	-
White, both sexes	26	-	-	-	-	-	-	-	1	1	1	4
Male	12	-	-	-	-	-	-	-	1	-	-	4
Female	14	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	14	-	-	-	-	1	-	-	-	1	-	-
Male	7	-	-	-	-	-	-	-	-	1	-	-
Female	7	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Missouri												
All races, both sexes	87	-	-	-	-	-	-	2	3	-	2	1
Male	40	-	-	-	-	-	-	1	2	-	1	1
Female	47	-	-	-	-	-	-	1	1	-	1	-
White, both sexes	74	-	-	-	-	-	-	1	1	-	1	1
Male	38	-	-	-	-	-	-	1	1	-	1	1
Female	36	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	13	-	-	-	-	-	-	1	2	-	1	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	11	-	-	-	-	-	-	1	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	15	-	-	-	-	-	-	-	-	1	-	-
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	1	-	-
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Nebraska												
All races, both sexes	25	-	-	-	-	-	-	1	-	1	-	2
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	1	-	1	-	2
White, both sexes	25	-	-	-	-	-	-	1	-	1	-	2
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	1	-	1	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	26	1	-	-	-	-	-	-	-	3	2	2
Male	15	-	-	-	-	-	-	-	-	3	1	1
Female	11	1	-	-	-	-	-	-	-	-	1	1
White, both sexes	23	1	-	-	-	-	-	-	-	3	2	1
Male	14	-	-	-	-	-	-	-	-	3	1	1
Female	9	1	-	-	-	-	-	-	-	-	1	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
New Hampshire												
All races, both sexes	16	-	-	-	-	-	-	-	-	2	-	-
Male	9	-	-	-	-	-	-	-	-	2	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	2	-	-
Male	9	-	-	-	-	-	-	-	-	2	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	130	-	-	-	1	-	-	1	1	2	10	9
Male	55	-	-	-	1	-	-	-	1	1	6	5
Female	75	-	-	-	-	-	-	1	-	1	4	4
White, both sexes	101	-	-	-	1	-	-	-	1	1	8	5
Male	43	-	-	-	1	-	-	-	1	-	6	3
Female	58	-	-	-	-	-	-	-	-	1	2	2
Black, both sexes	26	-	-	-	-	-	-	1	-	1	2	4
Male	11	-	-	-	-	-	-	-	-	1	-	2
Female	15	-	-	-	-	-	-	1	-	-	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
New Mexico												
All races, both sexes	37	-	-	-	-	-	1	-	-	3	2	-
Male	17	-	-	-	-	-	1	-	-	2	2	-
Female	20	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	36	-	-	-	-	-	-	-	-	3	2	-
Male	16	-	-	-	-	-	-	-	-	2	2	-
Female	20	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	259	4	-	-	-	-	-	2	6	8	18	25
Male	129	4	-	-	-	-	-	-	3	5	14	14
Female	130	-	-	-	-	-	-	2	3	3	4	11
White, both sexes	186	3	-	-	-	-	-	1	5	5	12	12
Male	87	3	-	-	-	-	-	-	2	3	9	8
Female	99	-	-	-	-	-	-	1	3	2	3	4
Black, both sexes	72	1	-	-	-	-	-	1	1	3	6	13
Male	41	1	-	-	-	-	-	-	1	2	5	6
Female	31	-	-	-	-	-	-	1	-	1	1	7
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
North Carolina												
All races, both sexes	114	1	-	-	-	-	-	1	2	2	6	9
Male	43	-	-	-	-	-	-	1	1	1	2	6
Female	71	1	-	-	-	-	-	-	1	1	4	3
White, both sexes	77	-	-	-	-	-	-	-	1	1	3	5
Male	28	-	-	-	-	-	-	-	1	1	1	4
Female	49	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	36	-	-	-	-	-	-	1	1	1	3	4
Male	15	-	-	-	-	-	-	1	-	-	1	2
Female	21	-	-	-	-	-	-	-	1	1	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Ohio												
All races, both sexes	209	1	-	-	-	1	1	2	2	6	10	8
Male	95	-	-	-	-	1	1	-	1	3	5	7
Female	114	1	-	-	-	-	-	2	1	3	5	1
White, both sexes	180	1	-	-	-	1	1	1	2	6	9	6
Male	86	-	-	-	-	1	1	-	1	3	5	5
Female	94	1	-	-	-	-	-	1	1	3	4	1
Black, both sexes	29	-	-	-	-	-	-	1	-	-	1	2
Male	9	-	-	-	-	-	-	-	-	-	-	2
Female	20	-	-	-	-	-	-	1	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	61	-	-	-	-	-	2	-	3	4	1	3
Male	33	-	-	-	-	-	1	-	2	3	1	2
Female	28	-	-	-	-	-	1	-	1	1	-	1
White, both sexes	49	-	-	-	-	-	1	-	2	4	-	1
Male	26	-	-	-	-	-	-	-	1	3	-	1
Female	23	-	-	-	-	-	1	-	1	1	-	-
Black, both sexes	6	-	-	-	-	-	1	-	1	-	-	1
Male	4	-	-	-	-	-	1	-	1	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Oregon												
All races, both sexes	55	1	-	-	-	-	-	-	-	1	-	6
Male	25	-	-	-	-	-	-	-	-	-	-	3
Female	30	1	-	-	-	-	-	-	-	1	-	3
White, both sexes	53	1	-	-	-	-	-	-	-	1	-	5
Male	24	-	-	-	-	-	-	-	-	-	-	2
Female	29	1	-	-	-	-	-	-	-	1	-	3
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	187	-	-	-	-	-	-	1	3	2	3	15
Male	90	-	-	-	-	-	-	1	2	2	3	11
Female	97	-	-	-	-	-	-	-	1	-	-	4
White, both sexes	163	-	-	-	-	-	-	1	2	2	3	12
Male	77	-	-	-	-	-	-	1	1	2	3	9
Female	86	-	-	-	-	-	-	-	1	-	-	3
Black, both sexes	24	-	-	-	-	-	-	-	1	-	-	3
Male	13	-	-	-	-	-	-	-	1	-	-	2
Female	11	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Rhode Island												
All races, both sexes	14	-	-	-	-	-	-	-	-	1	2	1
Male	7	-	-	-	-	-	-	-	-	1	2	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	1	1	1
Male	5	-	-	-	-	-	-	-	-	1	1	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	68	-	-	-	-	1	-	-	2	1	5	5
Male	30	-	-	-	-	1	-	-	1	1	2	1
Female	38	-	-	-	-	-	-	-	1	-	3	4
White, both sexes	48	-	-	-	-	-	-	-	2	1	3	4
Male	20	-	-	-	-	-	-	-	1	1	1	1
Female	28	-	-	-	-	-	-	-	1	-	2	3
Black, both sexes	20	-	-	-	-	1	-	-	-	-	2	1
Male	10	-	-	-	-	1	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
South Dakota												
All races, both sexes	8	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	112	1	1	-	-	-	1	4	2	-	9	5
Male	49	-	-	-	-	-	1	2	-	-	6	1
Female	63	1	1	-	-	-	-	2	2	-	3	4
White, both sexes	100	-	1	-	-	-	1	4	1	-	6	3
Male	45	-	-	-	-	-	1	2	-	-	4	1
Female	55	-	1	-	-	-	-	2	1	-	2	2
Black, both sexes	12	1	-	-	-	-	-	-	1	-	3	2
Male	4	-	-	-	-	-	-	-	-	-	2	-
Female	8	1	-	-	-	-	-	-	1	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Texas												
All races, both sexes	235	6	-	-	-	1	4	3	2	5	3	12
Male	114	3	-	-	-	-	1	-	1	3	2	10
Female	121	3	-	-	-	1	3	3	1	2	1	2
White, both sexes	193	4	-	-	-	1	4	2	2	5	2	10
Male	92	2	-	-	-	-	1	-	1	3	1	8
Female	101	2	-	-	-	1	3	2	1	2	1	2
Black, both sexes	40	2	-	-	-	-	-	1	-	-	1	2
Male	22	1	-	-	-	-	-	-	-	-	1	2
Female	18	1	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	1	1
Male	15	-	-	-	-	-	-	-	-	-	1	-
Female	9	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	24	-	-	-	-	-	-	-	-	-	1	1
Male	15	-	-	-	-	-	-	-	-	-	1	-
Female	9	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Vermont												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	111	3	-	-	-	1	1	4	3	2	4	8
Male	50	2	-	-	-	1	-	2	1	-	2	5
Female	61	1	-	-	-	-	1	2	2	2	2	3
White, both sexes	86	2	-	-	-	1	1	1	2	-	4	7
Male	38	1	-	-	-	1	-	1	-	-	2	4
Female	48	1	-	-	-	-	1	-	2	-	2	3
Black, both sexes	24	1	-	-	-	-	-	3	1	2	-	1
Male	11	1	-	-	-	-	-	1	1	-	-	1
Female	13	-	-	-	-	-	-	2	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Washington												
All races, both sexes	95	-	-	1	-	-	-	4	-	2	6	2
Male	42	-	-	1	-	-	-	1	-	-	3	1
Female	53	-	-	-	-	-	-	3	-	2	3	1
White, both sexes	93	-	-	1	-	-	-	3	-	2	6	2
Male	40	-	-	1	-	-	-	-	-	-	3	1
Female	53	-	-	-	-	-	-	3	-	2	3	1
Black, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	43	1	-	1	-	-	1	1	-	2	1	3
Male	19	1	-	1	-	-	-	-	-	2	-	3
Female	24	-	-	-	-	-	1	1	-	-	1	-
White, both sexes	41	1	-	1	-	-	1	1	-	2	1	3
Male	18	1	-	1	-	-	-	-	-	2	-	3
Female	23	-	-	-	-	-	1	1	-	-	1	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Wisconsin												
All races, both sexes	75	1	-	-	1	-	-	-	1	5	4	4
Male	35	1	-	-	1	-	-	-	1	3	1	3
Female	40	-	-	-	-	-	-	-	-	2	3	1
White, both sexes	66	1	-	-	1	-	-	-	1	3	3	2
Male	30	1	-	-	1	-	-	-	1	2	-	1
Female	36	-	-	-	-	-	-	-	-	1	3	1
Black, both sexes	8	-	-	-	-	-	-	-	-	2	1	2
Male	5	-	-	-	-	-	-	-	-	1	1	2
Female	3	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	10	-	1	-	-	1	-	-	1	-	-	-
Male	4	-	1	-	-	-	-	-	1	-	-	-
Female	6	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	9	-	1	-	-	1	-	-	1	-	-	-
Male	3	-	1	-	-	-	-	-	1	-	-	-
Female	6	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
United States												
All races, both sexes	289	296	290	250	302	419	469	495	303	124	38	1
Male	173	172	158	132	147	180	184	162	87	30	2	-
Female	116	124	132	118	155	239	285	333	216	94	36	1
White, both sexes	225	221	228	209	251	368	417	440	275	112	31	1
Male	134	121	125	110	130	160	163	145	79	29	2	-
Female	91	100	103	99	121	208	254	295	196	83	29	1
Black, both sexes	58	66	59	39	44	47	45	53	22	11	6	-
Male	35	45	32	21	16	19	17	17	6	-	-	-
Female	23	21	27	18	28	28	28	36	16	11	6	-
American Indian 1/, both sexes	3	1	1	-	2	2	1	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	1	-	2	2	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	3	8	2	2	5	2	6	2	5	1	1	-
Male	2	6	1	1	1	1	4	-	2	1	-	-
Female	1	2	1	1	4	1	2	2	3	-	1	-
Alabama												
All races, both sexes	7	8	2	5	6	4	7	11	3	3	-	-
Male	4	5	2	3	5	1	1	4	-	-	-	-
Female	3	3	-	2	1	3	6	7	3	3	-	-
White, both sexes	5	6	1	3	6	3	5	10	3	3	-	-
Male	3	3	1	1	5	1	1	4	-	-	-	-
Female	2	3	-	2	1	2	4	6	3	3	-	-
Black, both sexes	2	1	1	2	-	1	2	1	-	-	-	-
Male	1	1	1	2	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Alaska												
All races, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	3	5	10	11	6	6	6	6	10	1	-	-
Male	1	4	5	4	4	2	-	3	2	-	-	-
Female	2	1	5	7	2	4	6	3	8	1	-	-
White, both sexes	3	5	10	9	5	6	6	6	10	1	-	-
Male	1	4	5	3	4	2	-	3	2	-	-	-
Female	2	1	5	6	1	4	6	3	8	1	-	-
Black, both sexes	-	-	-	2	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Arkansas												
All races, both sexes	3	3	4	1	4	6	8	3	5	1	2	-
Male	3	1	1	-	1	2	7	2	1	-	-	-
Female	-	2	3	1	3	4	1	1	4	1	2	-
White, both sexes	2	2	3	1	3	5	5	3	5	1	2	-
Male	2	-	1	-	1	1	5	2	1	-	-	-
Female	-	2	2	1	2	4	-	1	4	1	2	-
Black, both sexes	1	1	1	-	1	1	3	-	-	-	-	-
Male	1	1	-	-	-	1	2	-	-	-	-	-
Female	-	-	1	-	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	27	32	43	32	30	40	45	44	15	10	4	-
Male	15	15	20	17	11	14	14	11	4	1	-	-
Female	12	17	23	15	19	26	31	33	11	9	4	-
White, both sexes	20	28	35	25	25	36	40	41	14	9	3	-
Male	10	13	15	14	10	12	12	10	4	1	-	-
Female	10	15	20	11	15	24	28	31	10	8	3	-
Black, both sexes	6	1	7	5	2	2	4	2	-	1	-	-
Male	4	1	5	2	1	1	2	1	-	-	-	-
Female	2	-	2	3	1	1	2	1	-	1	-	-
American Indian 1/, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	-	2	3	2	1	1	1	-	1	-
Male	1	1	-	1	-	1	-	-	-	-	-	-
Female	-	1	-	1	3	1	1	1	1	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Colorado												
All races, both sexes	3	9	2	2	6	4	3	4	7	-	-	-
Male	2	6	1	2	3	3	-	2	3	-	-	-
Female	1	3	1	-	3	1	3	2	4	-	-	-
White, both sexes	3	9	2	2	5	4	3	3	7	-	-	-
Male	2	6	1	2	3	3	-	2	3	-	-	-
Female	1	3	1	-	2	1	3	1	4	-	-	-
Black, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	2	1	3	2	1	3	5	5	6	2	-	-
Male	2	1	-	2	-	1	2	2	1	1	-	-
Female	-	-	3	-	1	2	3	3	5	1	-	-
White, both sexes	2	1	2	2	1	3	4	5	6	2	-	-
Male	2	1	-	2	-	1	1	2	1	1	-	-
Female	-	-	2	-	1	2	3	3	5	1	-	-
Black, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Delaware												
All races, both sexes	-	1	1	-	1	1	-	3	2	-	-	-
Male	-	-	1	-	1	-	-	2	-	-	-	-
Female	-	1	-	-	-	1	-	1	2	-	-	-
White, both sexes	-	1	1	-	1	1	-	3	2	-	-	-
Male	-	-	1	-	1	-	-	2	-	-	-	-
Female	-	1	-	-	-	1	-	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	1	-	1	1	2	-	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	1	1	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	1	1	-	1	1	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Florida												
All races, both sexes	26	13	14	22	20	31	32	27	17	10	2	-
Male	18	9	7	17	13	11	15	6	7	2	-	-
Female	8	4	7	5	7	20	17	21	10	8	2	-
White, both sexes	21	12	10	18	17	28	30	25	17	10	1	-
Male	15	8	6	15	11	11	14	5	7	2	-	-
Female	6	4	4	3	6	17	16	20	10	8	1	-
Black, both sexes	5	1	4	4	3	3	1	2	-	-	1	-
Male	3	1	1	2	2	-	-	1	-	-	-	-
Female	2	-	3	2	1	3	1	1	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	14	12	4	6	7	11	11	16	10	4	-	-
Male	10	7	2	3	2	5	3	6	2	-	-	-
Female	4	5	2	3	5	6	8	10	8	4	-	-
White, both sexes	12	3	1	5	5	10	11	11	9	1	-	-
Male	8	2	-	2	1	5	3	5	2	-	-	-
Female	4	1	1	3	4	5	8	6	7	1	-	-
Black, both sexes	2	9	3	1	2	1	-	5	1	3	-	-
Male	2	5	2	1	1	-	-	1	-	-	-	-
Female	-	4	1	-	1	1	-	4	1	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Hawaii												
All races, both sexes	-	-	1	-	2	1	3	2	3	1	-	-
Male	-	-	-	-	1	1	2	-	1	1	-	-
Female	-	-	1	-	1	-	1	2	2	-	-	-
White, both sexes	-	-	-	-	1	1	1	1	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	-	2	1	2	1	-	-
Male	-	-	-	-	1	-	2	-	1	1	-	-
Female	-	-	1	-	-	-	-	1	1	-	-	-
Idaho												
All races, both sexes	-	2	1	1	3	4	4	2	3	-	-	-
Male	-	2	1	1	3	2	-	1	-	-	-	-
Female	-	-	-	-	-	2	4	1	3	-	-	-
White, both sexes	-	2	1	1	3	3	4	2	3	-	-	-
Male	-	2	1	1	3	1	-	1	-	-	-	-
Female	-	-	-	-	-	2	4	1	3	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Illinois												
All races, both sexes	7	14	12	14	15	11	26	21	18	6	2	-
Male	3	10	6	9	6	2	11	6	5	-	-	-
Female	4	4	6	5	9	9	15	15	13	6	2	-
White, both sexes	6	7	9	10	11	7	23	21	14	6	2	-
Male	3	3	5	6	4	2	11	6	4	-	-	-
Female	3	4	4	4	7	5	12	15	10	6	2	-
Black, both sexes	1	6	3	4	4	4	3	-	4	-	-	-
Male	-	6	1	3	2	-	-	-	1	-	-	-
Female	1	-	2	1	2	4	3	-	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	3	5	2	4	3	7	11	11	8	2	-	-
Male	3	4	-	2	3	1	2	4	3	-	-	-
Female	-	1	2	2	-	6	9	7	5	2	-	-
White, both sexes	3	4	2	4	3	7	9	11	7	2	-	-
Male	3	3	-	2	3	1	2	4	3	-	-	-
Female	-	1	2	2	-	6	7	7	4	2	-	-
Black, both sexes	-	1	-	-	-	-	2	-	1	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Iowa												
All races, both sexes	2	4	4	4	2	4	3	6	6	2	-	-
Male	-	3	-	1	1	4	2	2	1	2	-	-
Female	2	1	4	3	1	-	1	4	5	-	-	-
White, both sexes	2	3	4	4	2	4	3	6	6	2	-	-
Male	-	2	-	1	1	4	2	2	1	2	-	-
Female	2	1	4	3	1	-	1	4	5	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	3	4	1	2	4	1	7	4	2	2	-	-
Male	2	1	-	-	1	-	2	-	-	1	-	-
Female	1	3	1	2	3	1	5	4	2	1	-	-
White, both sexes	3	4	1	2	4	1	6	4	2	2	-	-
Male	2	1	-	-	1	-	2	-	-	1	-	-
Female	1	3	1	2	3	1	4	4	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Kentucky												
All races, both sexes	3	6	3	2	2	3	7	12	6	1	-	-
Male	3	1	2	-	2	2	2	3	-	1	-	-
Female	-	5	1	2	-	1	5	9	6	-	-	-
White, both sexes	3	3	3	1	2	3	6	12	5	1	-	-
Male	3	-	2	-	2	2	1	3	-	1	-	-
Female	-	3	1	1	-	1	5	9	5	-	-	-
Black, both sexes	-	3	-	1	-	-	1	-	1	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	2	-	1	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	2	5	3	3	5	5	5	14	2	1	1	-
Male	1	3	2	1	1	1	1	4	1	1	-	-
Female	1	2	1	2	4	4	4	10	1	-	1	-
White, both sexes	2	4	2	1	5	4	5	10	1	1	1	-
Male	1	2	1	-	1	1	1	3	-	1	-	-
Female	1	2	1	1	4	3	4	7	1	-	1	-
Black, both sexes	-	1	1	2	-	1	-	4	1	-	-	-
Male	-	1	1	1	-	-	-	1	1	-	-	-
Female	-	-	-	1	-	1	-	3	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Maine												
All races, both sexes	2	-	2	-	1	2	-	4	-	1	-	-
Male	1	-	1	-	1	-	-	1	-	-	-	-
Female	1	-	1	-	-	2	-	3	-	1	-	-
White, both sexes	2	-	2	-	1	2	-	4	-	1	-	-
Male	1	-	1	-	1	-	-	1	-	-	-	-
Female	1	-	1	-	-	2	-	3	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	3	7	9	6	3	7	10	8	4	-	1	-
Male	1	5	6	5	1	6	7	3	1	-	-	-
Female	2	2	3	1	2	1	3	5	3	-	1	-
White, both sexes	2	6	6	4	3	5	8	5	1	-	1	-
Male	1	4	6	3	1	4	5	3	-	-	-	-
Female	1	2	-	1	2	1	3	2	1	-	1	-
Black, both sexes	1	1	3	2	-	2	2	3	3	-	-	-
Male	-	1	-	2	-	2	2	-	1	-	-	-
Female	1	-	3	-	-	-	-	3	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Massachusetts												
All races, both sexes	5	5	4	3	7	6	10	10	6	3	1	-
Male	4	3	2	2	4	2	4	4	4	1	1	-
Female	1	2	2	1	3	4	6	6	2	2	-	-
White, both sexes	4	5	4	3	3	6	9	9	5	3	1	-
Male	3	3	2	2	1	2	4	4	3	1	1	-
Female	1	2	2	1	2	4	5	5	2	2	-	-
Black, both sexes	1	-	-	-	4	-	1	1	-	-	-	-
Male	1	-	-	-	3	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	15	20	17	8	13	17	19	15	9	5	1	-
Male	8	14	11	2	5	9	5	9	2	2	1	-
Female	7	6	6	6	8	8	14	6	7	3	-	-
White, both sexes	12	12	12	7	11	15	17	13	9	4	1	-
Male	6	9	9	2	5	9	5	7	2	2	1	-
Female	6	3	3	5	6	6	12	6	7	2	-	-
Black, both sexes	3	8	5	1	2	2	2	2	-	1	-	-
Male	2	5	2	-	-	-	-	2	-	-	-	-
Female	1	3	3	1	2	2	2	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Minnesota												
All races, both sexes	5	3	2	4	1	1	6	3	5	3	1	-
Male	2	3	1	2	1	-	2	1	2	-	-	-
Female	3	-	1	2	-	1	4	2	3	3	1	-
White, both sexes	4	2	2	4	1	1	6	3	5	3	1	-
Male	2	2	1	2	1	-	2	1	2	-	-	-
Female	2	-	1	2	-	1	4	2	3	3	1	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	2	3	1	1	3	5	4	6	3	3	-	-
Male	2	2	1	-	-	2	2	1	1	2	-	-
Female	-	1	-	1	3	3	2	5	2	1	-	-
White, both sexes	-	1	1	1	1	4	2	4	2	3	-	-
Male	-	1	1	-	-	1	1	1	-	2	-	-
Female	-	-	-	1	1	3	1	3	2	1	-	-
Black, both sexes	2	2	-	-	2	1	2	2	1	-	-	-
Male	2	1	-	-	-	1	1	-	1	-	-	-
Female	-	1	-	-	2	-	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Missouri												
All races, both sexes	7	4	5	3	6	14	12	16	6	5	1	-
Male	4	3	2	2	4	6	2	9	2	1	-	-
Female	3	1	3	1	2	8	10	7	4	4	1	-
White, both sexes	5	3	5	3	6	12	10	14	6	5	1	-
Male	4	2	2	2	4	6	2	9	2	1	-	-
Female	1	1	3	1	2	6	8	5	4	4	1	-
Black, both sexes	2	1	-	-	-	2	2	2	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	2	2	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	2	2	1	-	1	-	2	2	4	-	-	-
Male	-	2	-	-	-	-	2	2	1	-	-	-
Female	2	-	1	-	1	-	-	-	3	-	-	-
White, both sexes	2	2	1	-	1	-	2	2	3	-	-	-
Male	-	2	-	-	-	-	2	2	1	-	-	-
Female	2	-	1	-	1	-	-	-	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Nebraska												
All races, both sexes	-	2	2	-	2	3	1	4	5	2	-	-
Male	-	-	2	-	1	3	1	-	3	-	-	-
Female	-	2	-	-	1	-	-	4	2	2	-	-
White, both sexes	-	2	2	-	2	3	1	4	5	2	-	-
Male	-	-	2	-	1	3	1	-	3	-	-	-
Female	-	2	-	-	1	-	-	4	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	3	3	3	-	1	5	-	3	-	-	-	-
Male	2	-	3	-	-	4	-	1	-	-	-	-
Female	1	3	-	-	1	1	-	2	-	-	-	-
White, both sexes	2	2	3	-	1	5	-	3	-	-	-	-
Male	1	-	3	-	-	4	-	1	-	-	-	-
Female	1	2	-	-	1	1	-	2	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
New Hampshire												
All races, both sexes	1	3	1	-	2	1	1	4	1	-	-	-
Male	-	2	1	-	1	1	1	1	-	-	-	-
Female	1	1	-	-	1	-	-	3	1	-	-	-
White, both sexes	1	3	1	-	2	1	1	4	1	-	-	-
Male	-	2	1	-	1	1	1	1	-	-	-	-
Female	1	1	-	-	1	-	-	3	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	11	4	7	6	17	11	16	14	17	2	-	1
Male	9	1	6	2	7	7	4	2	2	1	-	-
Female	2	3	1	4	10	4	12	12	15	1	-	1
White, both sexes	6	4	4	4	14	10	14	13	13	2	-	1
Male	5	1	3	2	6	6	4	2	2	1	-	-
Female	1	3	1	2	8	4	10	11	11	1	-	1
Black, both sexes	4	-	3	2	2	1	2	1	3	-	-	-
Male	3	-	3	-	1	1	-	-	-	-	-	-
Female	1	-	-	2	1	-	2	1	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
New Mexico												
All races, both sexes	3	2	2	1	2	3	11	4	2	1	-	-
Male	1	1	1	1	-	1	4	2	1	-	-	-
Female	2	1	1	-	2	2	7	2	1	1	-	-
White, both sexes	3	2	2	1	2	3	11	4	2	1	-	-
Male	1	1	1	1	-	1	4	2	1	-	-	-
Female	2	1	1	-	2	2	7	2	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	16	25	17	17	20	28	24	24	13	8	4	-
Male	10	19	8	9	2	16	12	7	4	2	-	-
Female	6	6	9	8	18	12	12	17	9	6	4	-
White, both sexes	9	12	13	13	16	19	21	21	13	8	3	-
Male	5	9	6	7	2	11	10	6	4	2	-	-
Female	4	3	7	6	14	8	11	15	9	6	3	-
Black, both sexes	7	12	4	4	4	9	3	3	-	-	1	-
Male	5	9	2	2	-	5	2	1	-	-	-	-
Female	2	3	2	2	4	4	1	2	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
North Carolina												
All races, both sexes	9	9	5	8	11	15	11	15	7	3	-	-
Male	5	4	3	4	5	4	2	5	-	-	-	-
Female	4	5	2	4	6	11	9	10	7	3	-	-
White, both sexes	9	4	2	6	7	12	10	10	6	1	-	-
Male	5	1	1	2	4	4	2	2	-	-	-	-
Female	4	3	1	4	3	8	8	8	6	1	-	-
Black, both sexes	-	5	3	2	4	3	1	5	1	2	-	-
Male	-	3	2	2	1	-	-	3	-	-	-	-
Female	-	2	1	-	3	3	1	2	1	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	2	1	-	2	2	-	1	5	-	-	-	-
Male	1	1	-	1	1	-	-	1	-	-	-	-
Female	1	-	-	1	1	-	1	4	-	-	-	-
White, both sexes	2	1	-	2	2	-	1	5	-	-	-	-
Male	1	1	-	1	1	-	-	1	-	-	-	-
Female	1	-	-	1	1	-	1	4	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Ohio												
All races, both sexes	11	15	15	18	11	29	26	33	11	6	3	-
Male	5	6	10	9	9	13	8	12	4	1	-	-
Female	6	9	5	9	2	16	18	21	7	5	3	-
White, both sexes	8	13	10	18	8	29	24	27	9	5	2	-
Male	4	6	9	9	7	13	8	9	4	1	-	-
Female	4	7	1	9	1	16	16	18	5	4	2	-
Black, both sexes	3	2	5	-	3	-	2	6	2	1	1	-
Male	1	-	1	-	2	-	-	3	-	-	-	-
Female	2	2	4	-	1	-	2	3	2	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	5	1	4	1	6	10	7	7	4	3	-	-
Male	4	1	3	-	4	4	2	1	3	2	-	-
Female	1	-	1	1	2	6	5	6	1	1	-	-
White, both sexes	3	-	4	1	6	7	6	7	4	3	-	-
Male	2	-	3	-	4	4	2	1	3	2	-	-
Female	1	-	1	1	2	3	4	6	1	1	-	-
Black, both sexes	-	1	-	-	-	1	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	2	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Oregon												
All races, both sexes	2	4	4	3	5	5	8	5	7	3	1	-
Male	1	2	3	-	2	1	5	5	2	1	-	-
Female	1	2	1	3	3	4	3	-	5	2	1	-
White, both sexes	1	4	4	3	5	5	8	5	7	3	1	-
Male	1	2	3	-	2	1	5	5	2	1	-	-
Female	-	2	1	3	3	4	3	-	5	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	15	3	22	10	15	24	19	29	13	10	3	-
Male	8	1	12	4	10	13	9	7	3	4	-	-
Female	7	2	10	6	5	11	10	22	10	6	3	-
White, both sexes	12	3	17	9	14	22	16	27	12	9	2	-
Male	6	1	8	3	10	12	8	7	2	4	-	-
Female	6	2	9	6	4	10	8	20	10	5	2	-
Black, both sexes	3	-	5	1	1	2	3	2	1	1	1	-
Male	2	-	4	1	-	1	1	-	1	-	-	-
Female	1	-	1	-	1	1	2	2	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Rhode Island												
All races, both sexes	-	1	1	1	1	-	2	2	1	-	1	-
Male	-	-	-	1	1	-	1	-	-	-	-	-
Female	-	1	1	-	-	-	1	2	1	-	1	-
White, both sexes	-	1	1	1	1	-	1	2	1	-	1	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	1	2	1	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	4	4	6	7	4	6	8	10	4	-	1	-
Male	3	1	5	4	2	2	5	1	1	-	-	-
Female	1	3	1	3	2	4	3	9	3	-	1	-
White, both sexes	3	4	3	4	3	5	6	5	4	-	1	-
Male	2	1	2	3	1	2	4	-	1	-	-	-
Female	1	3	1	1	2	3	2	5	3	-	1	-
Black, both sexes	1	-	3	3	1	1	2	5	-	-	-	-
Male	1	-	3	1	1	-	1	1	-	-	-	-
Female	-	-	-	2	-	1	1	4	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
South Dakota												
All races, both sexes	1	1	1	-	2	-	-	-	-	2	-	-
Male	1	1	-	-	-	-	-	-	-	1	-	-
Female	-	-	1	-	2	-	-	-	-	1	-	-
White, both sexes	1	-	1	-	2	-	-	-	-	2	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	1	-	2	-	-	-	-	1	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	4	11	8	6	9	12	12	14	7	5	1	-
Male	2	7	4	4	5	2	9	2	4	-	-	-
Female	2	4	4	2	4	10	3	12	3	5	1	-
White, both sexes	4	11	7	4	9	11	12	13	7	5	1	-
Male	2	7	3	3	5	2	9	2	4	-	-	-
Female	2	4	4	1	4	9	3	11	3	5	1	-
Black, both sexes	-	-	1	2	-	1	-	1	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Texas												
All races, both sexes	20	15	17	13	21	31	29	25	21	6	1	-
Male	12	9	11	7	12	15	10	12	5	1	-	-
Female	8	6	6	6	9	16	19	13	16	5	1	-
White, both sexes	13	10	15	13	16	24	24	22	19	6	1	-
Male	9	7	10	7	10	10	7	11	4	1	-	-
Female	4	3	5	6	6	14	17	11	15	5	1	-
Black, both sexes	6	5	2	-	5	7	4	3	2	-	-	-
Male	3	2	1	-	2	5	3	1	1	-	-	-
Female	3	3	1	-	3	2	1	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Utah												
All races, both sexes	4	1	1	2	1	1	3	9	-	-	-	-
Male	3	1	1	1	1	1	1	5	-	-	-	-
Female	1	-	-	1	-	-	2	4	-	-	-	-
White, both sexes	4	1	1	2	1	1	3	9	-	-	-	-
Male	3	1	1	1	1	1	1	5	-	-	-	-
Female	1	-	-	1	-	-	2	4	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Vermont												
All races, both sexes	2	-	1	-	-	-	1	1	-	-	-	-
Male	2	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	2	-	1	-	-	-	1	1	-	-	-	-
Male	2	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	7	10	7	8	5	14	8	12	7	3	4	-
Male	2	5	3	5	3	7	6	4	2	-	-	-
Female	5	5	4	3	2	7	2	8	5	3	4	-
White, both sexes	3	7	5	8	4	13	7	10	6	2	3	-
Male	-	3	3	5	3	6	5	2	2	-	-	-
Female	3	4	2	3	1	7	2	8	4	2	3	-
Black, both sexes	4	2	2	-	1	1	1	2	1	1	1	-
Male	2	1	-	-	-	1	1	2	-	-	-	-
Female	2	1	2	-	1	-	-	-	1	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Washington												
All races, both sexes	10	7	7	5	3	11	10	12	13	1	1	-
Male	7	3	1	2	2	5	5	5	5	1	-	-
Female	3	4	6	3	1	6	5	7	8	-	1	-
White, both sexes	9	7	7	5	3	11	10	12	13	1	1	-
Male	6	3	1	2	2	5	5	5	5	1	-	-
Female	3	4	6	3	1	6	5	7	8	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	4	2	3	2	3	3	11	1	4	-	-	-
Male	1	-	2	1	2	-	4	-	2	-	-	-
Female	3	2	1	1	1	3	7	1	2	-	-	-
White, both sexes	4	2	2	2	2	3	11	1	4	-	-	-
Male	1	-	1	1	2	-	4	-	2	-	-	-
Female	3	2	1	1	-	3	7	1	2	-	-	-
Black, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Wisconsin												
All races, both sexes	6	2	4	3	6	10	13	6	6	1	2	-
Male	4	2	3	2	3	3	5	1	2	-	-	-
Female	2	-	1	1	3	7	8	5	4	1	2	-
White, both sexes	6	2	3	3	5	10	13	6	6	-	1	-
Male	4	2	2	2	3	3	5	1	2	-	-	-
Female	2	-	1	1	2	7	8	5	4	-	1	-
Black, both sexes	-	-	1	-	-	-	-	-	-	1	1	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	1	1	-	-	1	-	1	-	2	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	1	-	2	-	-
White, both sexes	1	1	-	-	-	1	-	1	-	2	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	1	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
United States												
All races, both sexes	3,037	8	1	2	-	15	23	49	62	120	159	221
Male	1,335	5	-	1	-	6	14	21	41	64	93	115
Female	1,702	3	1	1	-	9	9	28	21	56	66	106
White, both sexes	2,450	6	1	1	-	11	14	30	43	76	110	140
Male	1,056	3	-	1	-	3	5	14	28	39	69	72
Female	1,394	3	1	-	-	8	9	16	15	37	41	68
Black, both sexes	553	2	-	-	-	4	7	18	19	44	48	77
Male	266	2	-	-	-	3	7	7	13	25	23	41
Female	287	-	-	-	-	1	-	11	6	19	25	36
American Indian 1/, both sexes	9	-	-	-	-	-	1	-	-	-	1	2
Male	2	-	-	-	-	-	1	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	25	-	-	1	-	-	1	1	-	-	-	2
Male	11	-	-	-	-	-	1	-	-	-	-	2
Female	14	-	-	1	-	-	-	1	-	-	-	-
Alabama												
All races, both sexes	36	1	-	-	-	-	-	1	-	-	1	2
Male	17	1	-	-	-	-	-	1	-	-	1	1
Female	19	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	28	1	-	-	-	-	-	1	-	-	-	1
Male	14	1	-	-	-	-	-	1	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	8	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	1	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Alaska												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	73	-	-	-	-	-	-	1	2	4	4	9
Male	32	-	-	-	-	-	-	1	1	1	2	4
Female	41	-	-	-	-	-	-	-	1	3	2	5
White, both sexes	69	-	-	-	-	-	-	1	2	3	4	8
Male	30	-	-	-	-	-	-	1	1	-	2	4
Female	39	-	-	-	-	-	-	-	1	3	2	4
Black, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Arkansas												
All races, both sexes	38	1	-	-	-	-	-	1	1	1	2	1
Male	20	1	-	-	-	-	-	1	-	1	2	-
Female	18	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	29	-	-	-	-	-	-	1	-	-	2	-
Male	14	-	-	-	-	-	-	1	-	-	2	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	1	-	-	-	-	-	-	1	1	-	1
Male	6	1	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	363	1	-	-	-	1	6	4	8	17	15	35
Male	149	-	-	-	-	1	3	2	6	8	12	15
Female	214	1	-	-	-	-	3	2	2	9	3	20
White, both sexes	291	1	-	-	-	-	4	2	4	12	11	23
Male	112	-	-	-	-	-	1	2	4	5	9	8
Female	179	1	-	-	-	-	3	-	-	7	2	15
Black, both sexes	57	-	-	-	-	1	1	1	4	5	4	11
Male	33	-	-	-	-	1	1	-	2	3	3	6
Female	24	-	-	-	-	-	-	1	2	2	1	5
American Indian 1/, both sexes	3	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	1	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Colorado												
All races, both sexes	44	-	-	-	-	-	-	-	-	3	5	4
Male	21	-	-	-	-	-	-	-	-	1	2	2
Female	23	-	-	-	-	-	-	-	-	2	3	2
White, both sexes	37	-	-	-	-	-	-	-	-	1	3	2
Male	18	-	-	-	-	-	-	-	-	-	1	1
Female	19	-	-	-	-	-	-	-	-	1	2	1
Black, both sexes	7	-	-	-	-	-	-	-	-	2	2	2
Male	3	-	-	-	-	-	-	-	-	1	1	1
Female	4	-	-	-	-	-	-	-	-	1	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	23	-	-	-	-	-	-	1	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	23	-	-	-	-	-	-	1	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Delaware												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	2
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	2
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Florida												
All races, both sexes	209	-	-	-	-	1	2	1	5	10	11	17
Male	101	-	-	-	-	-	2	1	5	6	9	5
Female	108	-	-	-	-	1	-	-	-	4	2	12
White, both sexes	166	-	-	-	-	1	1	1	1	6	8	8
Male	80	-	-	-	-	-	1	1	1	3	7	3
Female	86	-	-	-	-	1	-	-	-	3	1	5
Black, both sexes	42	-	-	-	-	-	1	-	4	4	3	9
Male	20	-	-	-	-	-	1	-	4	3	2	2
Female	22	-	-	-	-	-	-	-	-	1	1	7
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	83	-	-	-	-	-	2	3	-	4	7	6
Male	36	-	-	-	-	-	2	1	-	3	3	1
Female	47	-	-	-	-	-	-	2	-	1	4	5
White, both sexes	56	-	-	-	-	-	-	1	-	1	5	3
Male	21	-	-	-	-	-	-	-	-	-	3	-
Female	35	-	-	-	-	-	-	1	-	1	2	3
Black, both sexes	27	-	-	-	-	-	2	2	-	3	2	3
Male	15	-	-	-	-	-	2	1	-	3	-	1
Female	12	-	-	-	-	-	-	1	-	-	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Hawaii												
All races, both sexes	8	-	-	-	-	-	-	1	-	1	1	2
Male	5	-	-	-	-	-	-	1	-	1	-	1
Female	3	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	4	-	-	-	-	-	-	-	-	1	1	1
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	20	-	-	-	-	-	-	1	1	1	-	2
Male	11	-	-	-	-	-	-	1	1	-	-	1
Female	9	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	19	-	-	-	-	-	-	1	1	1	-	2
Male	10	-	-	-	-	-	-	1	1	-	-	1
Female	9	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Illinois												
All races, both sexes	143	-	-	1	-	-	1	1	5	6	14	15
Male	63	-	-	-	-	-	1	-	4	2	6	12
Female	80	-	-	1	-	-	-	1	1	4	8	3
White, both sexes	100	-	-	-	-	-	-	-	3	2	7	8
Male	45	-	-	-	-	-	-	-	2	1	5	6
Female	55	-	-	-	-	-	-	-	1	1	2	2
Black, both sexes	41	-	-	-	-	-	1	1	2	4	7	7
Male	17	-	-	-	-	-	1	-	2	1	1	6
Female	24	-	-	-	-	-	-	1	-	3	6	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	43	1	-	-	-	-	-	1	1	-	-	2
Male	17	1	-	-	-	-	-	-	1	-	-	2
Female	26	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	39	1	-	-	-	-	-	1	1	-	-	1
Male	15	1	-	-	-	-	-	-	1	-	-	1
Female	24	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Iowa												
All races, both sexes	33	-	-	-	-	-	-	-	-	1	3	-
Male	15	-	-	-	-	-	-	-	-	-	2	-
Female	18	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	32	-	-	-	-	-	-	-	-	1	3	-
Male	14	-	-	-	-	-	-	-	-	-	2	-
Female	18	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	26	-	-	-	-	-	-	1	-	1	1	2
Male	6	-	-	-	-	-	-	-	-	1	-	2
Female	20	-	-	-	-	-	-	1	-	-	1	-
White, both sexes	24	-	-	-	-	-	-	1	-	1	-	2
Male	6	-	-	-	-	-	-	-	-	1	-	2
Female	18	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Kentucky												
All races, both sexes	41	-	-	-	-	1	-	2	2	2	2	6
Male	18	-	-	-	-	-	-	2	2	1	1	3
Female	23	-	-	-	-	1	-	-	-	1	1	3
White, both sexes	36	-	-	-	-	1	-	2	2	2	1	6
Male	16	-	-	-	-	-	-	2	2	1	-	3
Female	20	-	-	-	-	1	-	-	-	1	1	3
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	38	-	-	-	-	-	-	1	-	1	1	2
Male	12	-	-	-	-	-	-	1	-	1	-	1
Female	26	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	27	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	11	-	-	-	-	-	-	1	-	1	1	1
Male	6	-	-	-	-	-	-	1	-	1	-	1
Female	5	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Maine												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	2	-
Male	3	-	-	-	-	-	-	-	-	-	2	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	2	-
Male	3	-	-	-	-	-	-	-	-	-	2	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	57	-	-	-	-	-	1	-	3	8	3	4
Male	33	-	-	-	-	-	1	-	1	4	2	-
Female	24	-	-	-	-	-	-	-	2	4	1	4
White, both sexes	30	-	-	-	-	-	-	-	1	-	2	1
Male	21	-	-	-	-	-	-	-	1	-	1	-
Female	9	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	27	-	-	-	-	-	1	-	2	8	1	3
Male	12	-	-	-	-	-	1	-	-	4	1	-
Female	15	-	-	-	-	-	-	-	2	4	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Massachusetts												
All races, both sexes	57	-	-	-	-	2	-	-	2	5	3	3
Male	30	-	-	-	-	1	-	-	-	3	2	2
Female	27	-	-	-	-	1	-	-	2	2	1	1
White, both sexes	47	-	-	-	-	1	-	-	2	4	2	2
Male	24	-	-	-	-	-	-	-	-	3	1	2
Female	23	-	-	-	-	1	-	-	2	1	1	-
Black, both sexes	9	-	-	-	-	1	-	-	-	1	1	1
Male	5	-	-	-	-	1	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	148	-	-	-	-	4	1	3	3	7	10	15
Male	69	-	-	-	-	1	1	1	2	4	5	7
Female	79	-	-	-	-	3	-	2	1	3	5	8
White, both sexes	109	-	-	-	-	4	1	1	3	2	6	11
Male	53	-	-	-	-	1	1	-	2	1	4	5
Female	56	-	-	-	-	3	-	1	1	1	2	6
Black, both sexes	39	-	-	-	-	-	-	2	-	5	4	4
Male	16	-	-	-	-	-	-	1	-	3	1	2
Female	23	-	-	-	-	-	-	1	-	2	3	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Minnesota												
All races, both sexes	24	1	-	-	-	-	-	-	1	-	-	2
Male	10	-	-	-	-	-	-	-	1	-	-	1
Female	14	1	-	-	-	-	-	-	-	-	-	1
White, both sexes	22	1	-	-	-	-	-	-	1	-	-	2
Male	9	-	-	-	-	-	-	-	1	-	-	1
Female	13	1	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	28	-	-	-	-	1	-	-	1	1	1	2
Male	12	-	-	-	-	-	-	-	1	-	-	2
Female	16	-	-	-	-	1	-	-	-	1	1	-
White, both sexes	20	-	-	-	-	-	-	-	1	1	1	2
Male	9	-	-	-	-	-	-	-	1	-	-	2
Female	11	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	8	-	-	-	-	1	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Missouri												
All races, both sexes	58	-	-	-	-	-	-	2	3	-	2	1
Male	23	-	-	-	-	-	-	1	2	-	1	1
Female	35	-	-	-	-	-	-	1	1	-	1	-
White, both sexes	46	-	-	-	-	-	-	1	1	-	1	1
Male	21	-	-	-	-	-	-	1	1	-	1	1
Female	25	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	12	-	-	-	-	-	-	1	2	-	1	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	10	-	-	-	-	-	-	1	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	10	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Nebraska												
All races, both sexes	14	-	-	-	-	-	-	1	-	1	-	2
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	1	-	1	-	2
White, both sexes	14	-	-	-	-	-	-	1	-	1	-	2
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	1	-	1	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	16	-	-	-	-	-	-	-	-	3	1	-
Male	9	-	-	-	-	-	-	-	-	3	-	-
Female	7	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	15	-	-	-	-	-	-	-	-	3	1	-
Male	8	-	-	-	-	-	-	-	-	3	-	-
Female	7	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
New Hampshire												
All races, both sexes	13	-	-	-	-	-	-	-	-	2	-	-
Male	8	-	-	-	-	-	-	-	-	2	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	2	-	-
Male	8	-	-	-	-	-	-	-	-	2	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	100	-	-	-	-	-	-	1	1	2	7	6
Male	38	-	-	-	-	-	-	-	1	1	4	2
Female	62	-	-	-	-	-	-	1	-	1	3	4
White, both sexes	77	-	-	-	-	-	-	-	1	1	5	4
Male	30	-	-	-	-	-	-	-	1	-	4	2
Female	47	-	-	-	-	-	-	-	-	1	1	2
Black, both sexes	21	-	-	-	-	-	-	1	-	1	2	2
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	13	-	-	-	-	-	-	1	-	-	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
New Mexico												
All races, both sexes	29	-	-	-	-	-	1	-	-	3	2	-
Male	12	-	-	-	-	-	1	-	-	2	2	-
Female	17	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	28	-	-	-	-	-	-	-	-	3	2	-
Male	11	-	-	-	-	-	-	-	-	2	2	-
Female	17	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	195	2	-	-	-	-	-	2	5	6	16	21
Male	92	2	-	-	-	-	-	-	2	3	12	13
Female	103	-	-	-	-	-	-	2	3	3	4	8
White, both sexes	134	1	-	-	-	-	-	1	4	3	10	9
Male	58	1	-	-	-	-	-	-	1	1	7	7
Female	76	-	-	-	-	-	-	1	3	2	3	2
Black, both sexes	61	1	-	-	-	-	-	1	1	3	6	12
Male	34	1	-	-	-	-	-	-	1	2	5	6
Female	27	-	-	-	-	-	-	1	-	1	1	6
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
North Carolina												
All races, both sexes	84	-	-	-	-	-	-	1	1	2	6	7
Male	28	-	-	-	-	-	-	1	1	1	2	5
Female	56	-	-	-	-	-	-	-	-	1	4	2
White, both sexes	56	-	-	-	-	-	-	-	1	1	3	3
Male	17	-	-	-	-	-	-	-	1	1	1	3
Female	39	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	28	-	-	-	-	-	-	1	-	1	3	4
Male	11	-	-	-	-	-	-	1	-	-	1	2
Female	17	-	-	-	-	-	-	-	-	1	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Ohio												
All races, both sexes	168	-	-	-	-	1	1	2	1	4	8	7
Male	75	-	-	-	-	1	1	-	1	1	4	6
Female	93	-	-	-	-	-	-	2	-	3	4	1
White, both sexes	143	-	-	-	-	1	1	1	1	4	7	5
Male	67	-	-	-	-	1	1	-	1	1	4	4
Female	76	-	-	-	-	-	-	1	-	3	3	1
Black, both sexes	25	-	-	-	-	-	-	1	-	-	1	2
Male	8	-	-	-	-	-	-	-	-	-	-	2
Female	17	-	-	-	-	-	-	1	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	43	-	-	-	-	-	2	-	2	4	1	2
Male	22	-	-	-	-	-	1	-	1	3	1	1
Female	21	-	-	-	-	-	1	-	1	1	-	1
White, both sexes	34	-	-	-	-	-	1	-	1	4	-	-
Male	17	-	-	-	-	-	-	-	-	3	-	-
Female	17	-	-	-	-	-	1	-	1	1	-	-
Black, both sexes	6	-	-	-	-	-	1	-	1	-	-	1
Male	4	-	-	-	-	-	1	-	1	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	1	1
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Oregon												
All races, both sexes	38	-	-	-	-	-	-	-	-	1	-	4
Male	16	-	-	-	-	-	-	-	-	-	-	1
Female	22	-	-	-	-	-	-	-	-	1	-	3
White, both sexes	38	-	-	-	-	-	-	-	-	1	-	4
Male	16	-	-	-	-	-	-	-	-	-	-	1
Female	22	-	-	-	-	-	-	-	-	1	-	3
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	121	-	-	-	-	-	-	1	3	1	1	10
Male	56	-	-	-	-	-	-	1	2	1	1	8
Female	65	-	-	-	-	-	-	-	1	-	-	2
White, both sexes	103	-	-	-	-	-	-	1	2	1	1	8
Male	44	-	-	-	-	-	-	1	1	1	1	6
Female	59	-	-	-	-	-	-	-	1	-	-	2
Black, both sexes	18	-	-	-	-	-	-	-	1	-	-	2
Male	12	-	-	-	-	-	-	-	1	-	-	2
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Rhode Island												
All races, both sexes	11	-	-	-	-	-	-	-	-	1	2	1
Male	5	-	-	-	-	-	-	-	-	1	2	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	1	1	1
Male	4	-	-	-	-	-	-	-	-	1	1	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	50	-	-	-	-	1	-	-	2	1	4	4
Male	19	-	-	-	-	1	-	-	1	1	1	-
Female	31	-	-	-	-	-	-	-	1	-	3	4
White, both sexes	37	-	-	-	-	-	-	-	2	1	2	3
Male	13	-	-	-	-	-	-	-	1	1	-	-
Female	24	-	-	-	-	-	-	-	1	-	2	3
Black, both sexes	13	-	-	-	-	1	-	-	-	-	2	1
Male	6	-	-	-	-	1	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
South Dakota												
All races, both sexes	7	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	91	-	1	-	-	-	1	4	1	-	7	3
Male	37	-	-	-	-	-	1	2	-	-	4	-
Female	54	-	1	-	-	-	-	2	1	-	3	3
White, both sexes	84	-	1	-	-	-	1	4	1	-	5	1
Male	35	-	-	-	-	-	1	2	-	-	3	-
Female	49	-	1	-	-	-	-	2	1	-	2	1
Black, both sexes	7	-	-	-	-	-	-	-	-	-	2	2
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Texas												
All races, both sexes	163	1	-	-	-	1	3	3	2	4	3	9
Male	70	-	-	-	-	-	-	-	1	2	2	7
Female	93	1	-	-	-	1	3	3	1	2	1	2
White, both sexes	133	1	-	-	-	1	3	2	2	4	2	7
Male	57	-	-	-	-	-	-	-	1	2	1	5
Female	76	1	-	-	-	1	3	2	1	2	1	2
Black, both sexes	29	-	-	-	-	-	-	1	-	-	1	2
Male	13	-	-	-	-	-	-	-	-	-	1	2
Female	16	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Vermont												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	70	-	-	-	-	1	1	4	3	2	4	5
Male	25	-	-	-	-	1	-	2	1	-	2	2
Female	45	-	-	-	-	-	1	2	2	2	2	3
White, both sexes	57	-	-	-	-	1	1	1	2	-	4	4
Male	19	-	-	-	-	1	-	1	-	-	2	1
Female	38	-	-	-	-	-	1	-	2	-	2	3
Black, both sexes	13	-	-	-	-	-	-	3	1	2	-	1
Male	6	-	-	-	-	-	-	1	1	-	-	1
Female	7	-	-	-	-	-	-	2	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Washington												
All races, both sexes	79	-	-	1	-	-	-	4	-	2	5	-
Male	34	-	-	1	-	-	-	1	-	-	3	-
Female	45	-	-	-	-	-	-	3	-	2	2	-
White, both sexes	78	-	-	1	-	-	-	3	-	2	5	-
Male	33	-	-	1	-	-	-	-	-	-	3	-
Female	45	-	-	-	-	-	-	3	-	2	2	-
Black, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	29	-	-	-	-	-	1	1	-	2	1	2
Male	11	-	-	-	-	-	-	-	-	2	-	2
Female	18	-	-	-	-	-	1	1	-	-	1	-
White, both sexes	29	-	-	-	-	-	1	1	-	2	1	2
Male	11	-	-	-	-	-	-	-	-	2	-	2
Female	18	-	-	-	-	-	1	1	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Wisconsin												
All races, both sexes	55	-	-	-	-	-	-	-	1	5	3	3
Male	25	-	-	-	-	-	-	-	1	3	1	2
Female	30	-	-	-	-	-	-	-	-	2	2	1
White, both sexes	47	-	-	-	-	-	-	-	1	3	2	1
Male	20	-	-	-	-	-	-	-	1	2	-	-
Female	27	-	-	-	-	-	-	-	-	1	2	1
Black, both sexes	8	-	-	-	-	-	-	-	-	2	1	2
Male	5	-	-	-	-	-	-	-	-	1	1	2
Female	3	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	7	-	-	-	-	1	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	5	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	1	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	5	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
United States												
All races, both sexes	218	226	222	185	218	308	339	350	205	82	23	1
Male	118	128	116	88	102	119	124	109	52	18	1	-
Female	100	98	106	97	116	189	215	241	153	64	22	1
White, both sexes	167	174	181	152	184	267	298	313	185	77	19	1
Male	89	95	95	71	94	105	110	97	47	18	1	-
Female	78	79	86	81	90	162	188	216	138	59	18	1
Black, both sexes	50	48	39	31	30	39	37	36	15	5	4	-
Male	29	31	20	16	8	13	12	12	4	-	-	-
Female	21	17	19	15	22	26	25	24	11	5	4	-
American Indian 1/, both sexes	-	1	1	-	-	1	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	1	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	3	1	2	4	1	3	1	4	-	-	-
Male	-	2	1	1	-	1	2	-	1	-	-	-
Female	1	1	-	1	4	-	1	1	3	-	-	-
Alabama												
All races, both sexes	3	3	1	2	5	3	3	7	3	1	-	-
Male	1	1	1	2	5	-	-	3	-	-	-	-
Female	2	2	-	-	-	3	3	4	3	1	-	-
White, both sexes	2	3	1	1	5	2	1	6	3	1	-	-
Male	1	1	1	1	5	-	-	3	-	-	-	-
Female	1	2	-	-	-	2	1	3	3	1	-	-
Black, both sexes	1	-	-	1	-	1	2	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Alaska												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	3	4	8	11	5	5	4	5	7	1	-	-
Male	1	4	4	4	4	1	-	3	2	-	-	-
Female	2	-	4	7	1	4	4	2	5	1	-	-
White, both sexes	3	4	8	9	5	5	4	5	7	1	-	-
Male	1	4	4	3	4	1	-	3	2	-	-	-
Female	2	-	4	6	1	4	4	2	5	1	-	-
Black, both sexes	-	-	-	2	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Arkansas												
All races, both sexes	2	3	3	-	3	3	7	2	5	1	2	-
Male	2	1	1	-	1	1	6	2	1	-	-	-
Female	-	2	2	-	2	2	1	-	4	1	2	-
White, both sexes	1	2	3	-	3	3	4	2	5	1	2	-
Male	1	-	1	-	1	1	4	2	1	-	-	-
Female	-	2	2	-	2	2	-	-	4	1	2	-
Black, both sexes	1	1	-	-	-	-	3	-	-	-	-	-
Male	1	1	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	21	26	40	29	27	35	39	37	12	7	3	-
Male	11	11	18	15	11	13	11	8	3	1	-	-
Female	10	15	22	14	16	22	28	29	9	6	3	-
White, both sexes	15	23	32	22	22	32	34	34	11	6	3	-
Male	7	10	13	12	10	11	9	7	3	1	-	-
Female	8	13	19	10	12	21	25	27	8	5	3	-
Black, both sexes	6	1	7	5	2	2	4	2	-	1	-	-
Male	4	1	5	2	1	1	2	1	-	-	-	-
Female	2	-	2	3	1	1	2	1	-	1	-	-
American Indian 1/, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	2	3	1	1	1	1	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	1	-	1	3	-	1	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Colorado												
All races, both sexes	3	8	2	2	5	3	3	2	4	-	-	-
Male	2	5	1	2	2	2	-	1	1	-	-	-
Female	1	3	1	-	3	1	3	1	3	-	-	-
White, both sexes	3	8	2	2	4	3	3	2	4	-	-	-
Male	2	5	1	2	2	2	-	1	1	-	-	-
Female	1	3	1	-	2	1	3	1	3	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	1	2	1	1	2	4	4	5	1	-	-
Male	1	1	-	1	-	1	1	1	1	1	-	-
Female	-	-	2	-	1	1	3	3	4	-	-	-
White, both sexes	1	1	2	1	1	2	4	4	5	1	-	-
Male	1	1	-	1	-	1	1	1	1	1	-	-
Female	-	-	2	-	1	1	3	3	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Delaware												
All races, both sexes	-	1	1	-	1	1	-	3	2	-	-	-
Male	-	-	1	-	1	-	-	2	-	-	-	-
Female	-	1	-	-	-	1	-	1	2	-	-	-
White, both sexes	-	1	1	-	1	1	-	3	2	-	-	-
Male	-	-	1	-	1	-	-	2	-	-	-	-
Female	-	1	-	-	-	1	-	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	1	-	1	1	2	-	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	1	1	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	1	1	-	1	1	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Florida												
All races, both sexes	21	9	10	16	13	22	27	21	15	7	1	-
Male	14	6	5	12	8	6	12	5	5	-	-	-
Female	7	3	5	4	5	16	15	16	10	7	1	-
White, both sexes	17	8	7	12	10	20	25	19	15	7	-	-
Male	12	5	5	10	6	6	11	4	5	-	-	-
Female	5	3	2	2	4	14	14	15	10	7	-	-
Black, both sexes	4	1	3	4	3	2	1	2	-	-	1	-
Male	2	1	-	2	2	-	-	1	-	-	-	-
Female	2	-	3	2	1	2	1	1	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	9	9	2	3	3	8	7	12	7	1	-	-
Male	6	6	1	1	-	4	1	5	2	-	-	-
Female	3	3	1	2	3	4	6	7	5	1	-	-
White, both sexes	7	2	1	2	3	7	7	10	6	1	-	-
Male	4	2	-	-	-	4	1	5	2	-	-	-
Female	3	-	1	2	3	3	6	5	4	1	-	-
Black, both sexes	2	7	1	1	-	1	-	2	1	-	-	-
Male	2	4	1	1	-	-	-	-	-	-	-	-
Female	-	3	-	-	-	1	-	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Hawaii												
All races, both sexes	-	-	-	-	-	1	1	-	1	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Idaho												
All races, both sexes	-	1	1	-	3	4	3	2	1	-	-	-
Male	-	1	1	-	3	2	-	1	-	-	-	-
Female	-	-	-	-	-	2	3	1	1	-	-	-
White, both sexes	-	1	1	-	3	3	3	2	1	-	-	-
Male	-	1	1	-	3	1	-	1	-	-	-	-
Female	-	-	-	-	-	2	3	1	1	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Illinois												
All races, both sexes	5	11	8	10	9	8	20	16	8	5	-	-
Male	1	8	4	5	4	1	11	3	1	-	-	-
Female	4	3	4	5	5	7	9	13	7	5	-	-
White, both sexes	4	6	7	7	7	5	17	16	6	5	-	-
Male	1	3	4	3	4	1	11	3	1	-	-	-
Female	3	3	3	4	3	4	6	13	5	5	-	-
Black, both sexes	1	4	1	3	2	3	3	-	2	-	-	-
Male	-	4	-	2	-	-	-	-	-	-	-	-
Female	1	-	1	1	2	3	3	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1	4	2	1	2	6	8	8	4	2	-	-
Male	1	3	-	1	2	1	1	2	2	-	-	-
Female	-	1	2	-	-	5	7	6	2	2	-	-
White, both sexes	1	3	2	1	2	6	6	8	4	2	-	-
Male	1	2	-	1	2	1	1	2	2	-	-	-
Female	-	1	2	-	-	5	5	6	2	2	-	-
Black, both sexes	-	1	-	-	-	-	2	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Iowa												
All races, both sexes	2	3	3	4	2	4	2	4	3	2	-	-
Male	-	2	-	1	1	4	2	1	-	2	-	-
Female	2	1	3	3	1	-	-	3	3	-	-	-
White, both sexes	2	2	3	4	2	4	2	4	3	2	-	-
Male	-	1	-	1	1	4	2	1	-	2	-	-
Female	2	1	3	3	1	-	-	3	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	2	3	1	2	1	1	5	3	2	1	-	-
Male	1	1	-	-	-	-	1	-	-	-	-	-
Female	1	2	1	2	1	1	4	3	2	1	-	-
White, both sexes	2	3	1	2	1	1	4	3	2	1	-	-
Male	1	1	-	-	-	-	1	-	-	-	-	-
Female	1	2	1	2	1	1	3	3	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Kentucky												
All races, both sexes	1	6	2	2	2	1	2	7	2	1	-	-
Male	1	1	2	-	2	-	-	2	-	1	-	-
Female	-	5	-	2	-	1	2	5	2	-	-	-
White, both sexes	1	3	2	1	2	1	2	7	2	1	-	-
Male	1	-	2	-	2	-	-	2	-	1	-	-
Female	-	3	-	1	-	1	2	5	2	-	-	-
Black, both sexes	-	3	-	1	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	2	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	4	3	2	3	5	4	9	2	-	-	-
Male	-	2	2	-	-	1	1	2	1	-	-	-
Female	1	2	1	2	3	4	3	7	1	-	-	-
White, both sexes	1	4	2	1	3	4	4	6	1	-	-	-
Male	-	2	1	-	-	1	1	1	-	-	-	-
Female	1	2	1	1	3	3	3	5	1	-	-	-
Black, both sexes	-	-	1	1	-	1	-	3	1	-	-	-
Male	-	-	1	-	-	-	-	1	1	-	-	-
Female	-	-	-	1	-	1	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Maine												
All races, both sexes	1	-	-	-	1	1	-	2	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	2	-	-	-	-
White, both sexes	1	-	-	-	1	1	-	2	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	5	7	4	1	6	6	4	3	-	1	-
Male	1	3	5	4	-	5	4	2	1	-	-	-
Female	-	2	2	-	1	1	2	2	2	-	1	-
White, both sexes	1	5	5	2	1	4	5	2	-	-	1	-
Male	1	3	5	2	-	3	3	2	-	-	-	-
Female	-	2	-	-	1	1	2	-	-	-	1	-
Black, both sexes	-	-	2	2	-	2	1	2	3	-	-	-
Male	-	-	-	2	-	2	1	-	1	-	-	-
Female	-	-	2	-	-	-	-	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Massachusetts												
All races, both sexes	5	3	2	1	6	5	8	6	5	1	-	-
Male	4	3	1	-	3	1	4	1	4	1	-	-
Female	1	-	1	1	3	4	4	5	1	-	-	-
White, both sexes	4	3	2	1	3	5	8	5	4	1	-	-
Male	3	3	1	-	1	1	4	1	3	1	-	-
Female	1	-	1	1	2	4	4	4	1	-	-	-
Black, both sexes	1	-	-	-	3	-	-	1	-	-	-	-
Male	1	-	-	-	2	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	13	18	14	7	10	10	12	12	4	4	1	-
Male	6	12	8	1	4	5	2	8	-	1	1	-
Female	7	6	6	6	6	5	10	4	4	3	-	-
White, both sexes	10	12	9	6	8	8	10	10	4	3	1	-
Male	4	9	6	1	4	5	2	6	-	1	1	-
Female	6	3	3	5	4	3	8	4	4	2	-	-
Black, both sexes	3	6	5	1	2	2	2	2	-	1	-	-
Male	2	3	2	-	-	-	-	2	-	-	-	-
Female	1	3	3	1	2	2	2	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Minnesota												
All races, both sexes	4	1	2	2	1	1	4	2	1	1	1	-
Male	2	1	1	1	1	-	1	1	-	-	-	-
Female	2	-	1	1	-	1	3	1	1	1	1	-
White, both sexes	3	-	2	2	1	1	4	2	1	1	1	-
Male	2	-	1	1	1	-	1	1	-	-	-	-
Female	1	-	1	1	-	1	3	1	1	1	1	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1	2	1	1	2	3	2	6	1	3	-	-
Male	1	2	1	-	-	1	1	1	-	2	-	-
Female	-	-	-	1	2	2	1	5	1	1	-	-
White, both sexes	-	1	1	1	-	2	2	4	1	3	-	-
Male	-	1	1	-	-	-	1	1	-	2	-	-
Female	-	-	-	1	-	2	1	3	1	1	-	-
Black, both sexes	1	1	-	-	2	1	-	2	-	-	-	-
Male	1	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	2	-	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Missouri												
All races, both sexes	5	2	3	1	4	10	12	7	4	2	-	-
Male	3	1	1	1	3	4	2	3	-	-	-	-
Female	2	1	2	-	1	6	10	4	4	2	-	-
White, both sexes	4	1	3	1	4	8	10	5	4	2	-	-
Male	3	-	1	1	3	4	2	3	-	-	-	-
Female	1	1	2	-	1	4	8	2	4	2	-	-
Black, both sexes	1	1	-	-	-	2	2	2	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	2	2	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	2	1	-	1	-	1	-	3	-	-	-
Male	-	2	-	-	-	-	1	-	1	-	-	-
Female	1	-	1	-	1	-	-	-	2	-	-	-
White, both sexes	1	2	1	-	1	-	1	-	2	-	-	-
Male	-	2	-	-	-	-	1	-	1	-	-	-
Female	1	-	1	-	1	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Nebraska												
All races, both sexes	-	1	1	-	1	2	-	2	2	1	-	-
Male	-	-	1	-	1	2	-	-	2	-	-	-
Female	-	1	-	-	-	-	-	2	-	1	-	-
White, both sexes	-	1	1	-	1	2	-	2	2	1	-	-
Male	-	-	1	-	1	2	-	-	2	-	-	-
Female	-	1	-	-	-	-	-	2	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	3	1	2	-	1	3	-	2	-	-	-	-
Male	2	-	2	-	-	2	-	-	-	-	-	-
Female	1	1	-	-	1	1	-	2	-	-	-	-
White, both sexes	2	1	2	-	1	3	-	2	-	-	-	-
Male	1	-	2	-	-	2	-	-	-	-	-	-
Female	1	1	-	-	1	1	-	2	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
New Hampshire												
All races, both sexes	1	3	1	-	1	-	1	3	1	-	-	-
Male	-	2	1	-	1	-	1	1	-	-	-	-
Female	1	1	-	-	-	-	-	2	1	-	-	-
White, both sexes	1	3	1	-	1	-	1	3	1	-	-	-
Male	-	2	1	-	1	-	1	1	-	-	-	-
Female	1	1	-	-	-	-	-	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	9	3	5	3	11	11	14	9	15	2	-	1
Male	7	1	4	1	4	7	3	1	1	1	-	-
Female	2	2	1	2	7	4	11	8	14	1	-	1
White, both sexes	5	3	3	3	8	10	12	8	11	2	-	1
Male	4	1	2	1	3	6	3	1	1	1	-	-
Female	1	2	1	2	5	4	9	7	10	1	-	1
Black, both sexes	4	-	2	-	2	1	2	1	3	-	-	-
Male	3	-	2	-	1	1	-	-	-	-	-	-
Female	1	-	-	-	1	-	2	1	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
New Mexico												
All races, both sexes	3	1	1	1	2	3	9	1	2	-	-	-
Male	1	-	-	1	-	1	3	-	1	-	-	-
Female	2	1	1	-	2	2	6	1	1	-	-	-
White, both sexes	3	1	1	1	2	3	9	1	2	-	-	-
Male	1	-	-	1	-	1	3	-	1	-	-	-
Female	2	1	1	-	2	2	6	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	13	19	13	14	17	20	13	16	11	5	2	-
Male	7	13	5	7	1	11	6	5	3	2	-	-
Female	6	6	8	7	16	9	7	11	8	3	2	-
White, both sexes	7	10	10	10	14	13	10	15	11	5	1	-
Male	3	7	4	5	1	8	4	4	3	2	-	-
Female	4	3	6	5	13	5	6	11	8	3	1	-
Black, both sexes	6	9	3	4	3	7	3	1	-	-	1	-
Male	4	6	1	2	-	3	2	1	-	-	-	-
Female	2	3	2	2	3	4	1	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
North Carolina												
All races, both sexes	5	8	2	6	5	12	10	11	6	2	-	-
Male	1	3	1	2	2	3	1	5	-	-	-	-
Female	4	5	1	4	3	9	9	6	6	2	-	-
White, both sexes	5	4	-	5	3	9	9	7	5	1	-	-
Male	1	1	-	1	2	3	1	2	-	-	-	-
Female	4	3	-	4	1	6	8	5	5	1	-	-
Black, both sexes	-	4	2	1	2	3	1	4	1	1	-	-
Male	-	2	1	1	-	-	-	3	-	-	-	-
Female	-	2	1	-	2	3	1	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	2	1	-	1	2	-	-	1	-	-	-	-
Male	1	1	-	1	1	-	-	1	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	2	1	-	1	2	-	-	1	-	-	-	-
Male	1	1	-	1	1	-	-	1	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Ohio												
All races, both sexes	10	12	12	15	10	23	19	29	8	4	2	-
Male	5	6	8	6	8	9	6	10	3	-	-	-
Female	5	6	4	9	2	14	13	19	5	4	2	-
White, both sexes	7	11	8	15	7	23	18	24	6	3	1	-
Male	4	6	7	6	6	9	6	8	3	-	-	-
Female	3	5	1	9	1	14	12	16	3	3	1	-
Black, both sexes	3	1	4	-	3	-	1	5	2	1	1	-
Male	1	-	1	-	2	-	-	2	-	-	-	-
Female	2	1	3	-	1	-	1	3	2	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	3	1	3	1	5	8	6	2	1	2	-	-
Male	2	1	2	-	3	3	2	-	1	1	-	-
Female	1	-	1	1	2	5	4	2	-	1	-	-
White, both sexes	3	-	3	1	5	6	5	2	1	2	-	-
Male	2	-	2	-	3	3	2	-	1	1	-	-
Female	1	-	1	1	2	3	3	2	-	1	-	-
Black, both sexes	-	1	-	-	-	1	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Oregon												
All races, both sexes	-	4	4	3	2	4	6	4	4	2	-	-
Male	-	2	3	-	-	1	4	4	1	-	-	-
Female	-	2	1	3	2	3	2	-	3	2	-	-
White, both sexes	-	4	4	3	2	4	6	4	4	2	-	-
Male	-	2	3	-	-	1	4	4	1	-	-	-
Female	-	2	1	3	2	3	2	-	3	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	12	3	14	5	11	13	11	20	8	7	1	-
Male	5	1	8	1	7	7	5	4	2	3	-	-
Female	7	2	6	4	4	6	6	16	6	4	1	-
White, both sexes	9	3	11	4	10	11	8	19	7	7	1	-
Male	3	1	5	-	7	6	4	4	1	3	-	-
Female	6	2	6	4	3	5	4	15	6	4	1	-
Black, both sexes	3	-	3	1	1	2	3	1	1	-	-	-
Male	2	-	3	1	-	1	1	-	1	-	-	-
Female	1	-	-	-	1	1	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Rhode Island												
All races, both sexes	-	1	1	1	-	-	1	2	-	-	1	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	1	2	-	-	1	-
White, both sexes	-	1	1	1	-	-	1	2	-	-	1	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	1	2	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	2	3	4	5	3	5	6	7	2	-	1	-
Male	2	1	3	3	1	1	3	-	1	-	-	-
Female	-	2	1	2	2	4	3	7	1	-	1	-
White, both sexes	1	3	3	3	3	4	4	5	2	-	1	-
Male	1	1	2	2	1	1	2	-	1	-	-	-
Female	-	2	1	1	2	3	2	5	1	-	1	-
Black, both sexes	1	-	1	2	-	1	2	2	-	-	-	-
Male	1	-	1	1	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	1	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
South Dakota												
All races, both sexes	1	1	-	-	2	-	-	-	-	2	-	-
Male	1	1	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	2	-	-	-	-	1	-	-
White, both sexes	1	-	-	-	2	-	-	-	-	2	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	2	-	-	-	-	1	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	4	10	7	4	7	9	10	12	5	5	1	-
Male	2	6	3	2	4	2	7	2	2	-	-	-
Female	2	4	4	2	3	7	3	10	3	5	1	-
White, both sexes	4	10	6	3	7	8	10	12	5	5	1	-
Male	2	6	2	2	4	2	7	2	2	-	-	-
Female	2	4	4	1	3	6	3	10	3	5	1	-
Black, both sexes	-	-	1	1	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Texas												
All races, both sexes	13	8	13	9	17	22	17	19	14	4	1	-
Male	7	4	8	4	8	8	8	7	4	-	-	-
Female	6	4	5	5	9	14	9	12	10	4	1	-
White, both sexes	6	4	12	9	14	18	13	17	13	4	1	-
Male	4	3	8	4	8	6	5	7	3	-	-	-
Female	2	1	4	5	6	12	8	10	10	4	1	-
Black, both sexes	6	4	1	-	3	4	4	2	1	-	-	-
Male	3	1	-	-	-	2	3	-	1	-	-	-
Female	3	3	1	-	3	2	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	4	1	1	2	-	-	2	6	-	-	-	-
Male	3	1	1	1	-	-	-	4	-	-	-	-
Female	1	-	-	1	-	-	2	2	-	-	-	-
White, both sexes	4	1	1	2	-	-	2	6	-	-	-	-
Male	3	1	1	1	-	-	-	4	-	-	-	-
Female	1	-	-	1	-	-	2	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Vermont												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	5	6	5	5	3	7	3	9	4	1	2	-
Male	1	3	2	3	2	2	1	3	-	-	-	-
Female	4	3	3	2	1	5	2	6	4	1	2	-
White, both sexes	2	6	4	5	3	6	3	8	4	1	2	-
Male	-	3	2	3	2	1	1	2	-	-	-	-
Female	2	3	2	2	1	5	2	6	4	1	2	-
Black, both sexes	3	-	1	-	-	1	-	1	-	-	-	-
Male	1	-	-	-	-	1	-	1	-	-	-	-
Female	2	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Washington												
All races, both sexes	8	6	7	4	3	9	9	7	12	1	1	-
Male	5	2	1	1	2	4	4	4	5	1	-	-
Female	3	4	6	3	1	5	5	3	7	-	1	-
White, both sexes	8	6	7	4	3	9	9	7	12	1	1	-
Male	5	2	1	1	2	4	4	4	5	1	-	-
Female	3	4	6	3	1	5	5	3	7	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	4	1	2	2	-	2	8	-	3	-	-	-
Male	1	-	1	1	-	-	3	-	1	-	-	-
Female	3	1	1	1	-	2	5	-	2	-	-	-
White, both sexes	4	1	2	2	-	2	8	-	3	-	-	-
Male	1	-	1	1	-	-	3	-	1	-	-	-
Female	3	1	1	1	-	2	5	-	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Wisconsin												
All races, both sexes	6	2	4	2	3	4	10	6	3	1	2	-
Male	4	2	3	2	1	1	4	1	-	-	-	-
Female	2	-	1	-	2	3	6	5	3	1	2	-
White, both sexes	6	2	3	2	3	4	10	6	3	-	1	-
Male	4	2	2	2	1	1	4	1	-	-	-	-
Female	2	-	1	-	2	3	6	5	3	-	1	-
Black, both sexes	-	-	1	-	-	-	-	-	-	1	1	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	1	-	-	1	-	1	-	2	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	2	-	-
White, both sexes	-	-	-	-	-	1	-	1	-	2	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
United States												
All races, both sexes	1,064	45	2	1	3	1	1	2	7	12	31	60
Male	567	26	1	1	2	1	1	1	4	10	21	47
Female	497	19	1	-	1	-	-	1	3	2	10	13
White, both sexes	886	34	2	1	3	1	1	1	5	8	23	47
Male	467	19	1	1	2	1	1	1	4	7	16	38
Female	419	15	1	-	1	-	-	-	1	1	7	9
Black, both sexes	148	10	-	-	-	-	-	1	2	4	5	10
Male	82	7	-	-	-	-	-	-	-	3	3	6
Female	66	3	-	-	-	-	-	1	2	1	2	4
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	1	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	22	1	-	-	-	-	-	-	-	-	2	2
Male	14	-	-	-	-	-	-	-	-	-	1	2
Female	8	1	-	-	-	-	-	-	-	-	1	-
Alabama												
All races, both sexes	30	-	-	-	-	-	-	-	-	1	2	2
Male	15	-	-	-	-	-	-	-	-	-	1	2
Female	15	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	24	-	-	-	-	-	-	-	-	1	1	2
Male	10	-	-	-	-	-	-	-	-	-	1	2
Female	14	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Alaska												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	15	2	-	-	-	-	-	-	-	-	2	-
Male	4	1	-	-	-	-	-	-	-	-	1	-
Female	11	1	-	-	-	-	-	-	-	-	1	-
White, both sexes	12	2	-	-	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	9	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Arkansas												
All races, both sexes	10	1	-	-	-	-	-	-	-	-	-	-
Male	4	1	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	1	-	-	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	63	6	-	-	-	1	-	-	1	-	2	7
Male	33	4	-	-	-	1	-	-	1	-	1	6
Female	30	2	-	-	-	-	-	-	-	-	1	1
White, both sexes	55	4	-	-	-	1	-	-	1	-	2	5
Male	27	2	-	-	-	1	-	-	1	-	1	4
Female	28	2	-	-	-	-	-	-	-	-	1	1
Black, both sexes	2	2	-	-	-	-	-	-	-	-	-	-
Male	2	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	2
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Colorado												
All races, both sexes	10	1	-	-	-	-	-	-	-	-	-	1
Male	8	1	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	1	-	-	-	-	-	-	-	-	-	-
Male	7	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	3
Male	7	-	-	-	-	-	-	-	-	-	-	3
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	3
Male	6	-	-	-	-	-	-	-	-	-	-	3
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Delaware												
All races, both sexes	3	3	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	2	2	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	3	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	2	2	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Florida												
All races, both sexes	57	2	-	-	-	-	-	-	-	1	1	1
Male	36	2	-	-	-	-	-	-	-	1	-	1
Female	21	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	52	1	-	-	-	-	-	-	-	-	1	1
Male	32	1	-	-	-	-	-	-	-	-	-	1
Female	20	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	5	1	-	-	-	-	-	-	-	1	-	-
Male	4	1	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	40	3	-	-	1	-	-	-	-	-	2	-
Male	18	2	-	-	-	-	-	-	-	-	2	-
Female	22	1	-	-	1	-	-	-	-	-	-	-
White, both sexes	24	-	-	-	1	-	-	-	-	-	1	-
Male	11	-	-	-	-	-	-	-	-	-	1	-
Female	13	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	16	3	-	-	-	-	-	-	-	-	1	-
Male	7	2	-	-	-	-	-	-	-	-	1	-
Female	9	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Hawaii												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	8	1	1	-	-	-	-	-	-	1	-	-
Male	4	1	-	-	-	-	-	-	-	1	-	-
Female	4	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	8	1	1	-	-	-	-	-	-	1	-	-
Male	4	1	-	-	-	-	-	-	-	1	-	-
Female	4	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Illinois												
All races, both sexes	55	2	-	-	-	-	-	-	-	2	2	3
Male	26	1	-	-	-	-	-	-	-	1	1	3
Female	29	1	-	-	-	-	-	-	-	1	1	-
White, both sexes	40	2	-	-	-	-	-	-	-	-	1	1
Male	15	1	-	-	-	-	-	-	-	-	-	1
Female	25	1	-	-	-	-	-	-	-	-	1	-
Black, both sexes	14	-	-	-	-	-	-	-	-	2	-	2
Male	10	-	-	-	-	-	-	-	-	1	-	2
Female	4	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	21	1	-	-	-	-	-	1	-	-	-	1
Male	11	1	-	-	-	-	-	1	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	20	1	-	-	-	-	-	1	-	-	-	1
Male	11	1	-	-	-	-	-	1	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Iowa												
All races, both sexes	9	1	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	1	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	10	-	-	-	-	-	-	-	1	-	-	-
Male	5	-	-	-	-	-	-	-	1	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	1	-	-	-
Male	5	-	-	-	-	-	-	-	1	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Kentucky												
All races, both sexes	21	-	-	-	-	-	-	-	-	-	-	2
Male	9	-	-	-	-	-	-	-	-	-	-	2
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	2
Male	8	-	-	-	-	-	-	-	-	-	-	2
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	15	1	-	-	-	-	-	-	-	-	1	-
Male	9	1	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	1	-	-	-	-	-	-	-	-	1	-
Male	7	1	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Maine												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	22	-	-	-	-	-	-	1	-	-	1	-
Male	11	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	1	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Massachusetts												
All races, both sexes	22	-	-	-	-	-	-	-	-	-	1	3
Male	11	-	-	-	-	-	-	-	-	-	-	2
Female	11	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	3
Male	10	-	-	-	-	-	-	-	-	-	-	2
Female	9	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	35	-	-	-	-	-	-	-	-	-	-	1
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	32	-	-	-	-	-	-	-	-	-	-	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Minnesota												
All races, both sexes	19	4	-	-	-	-	-	-	-	-	-	1
Male	8	1	-	-	-	-	-	-	-	-	-	1
Female	11	3	-	-	-	-	-	-	-	-	-	-
White, both sexes	18	4	-	-	-	-	-	-	-	-	-	-
Male	7	1	-	-	-	-	-	-	-	-	-	-
Female	11	3	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	12	-	-	-	-	-	-	-	-	1	-	2
Male	7	-	-	-	-	-	-	-	-	1	-	2
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	2
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Missouri												
All races, both sexes	29	-	-	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Nebraska												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	10	1	-	-	-	-	-	-	-	-	1	2
Male	6	-	-	-	-	-	-	-	-	-	1	1
Female	4	1	-	-	-	-	-	-	-	-	-	1
White, both sexes	8	1	-	-	-	-	-	-	-	-	1	1
Male	6	-	-	-	-	-	-	-	-	-	1	1
Female	2	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
New Hampshire												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	30	-	-	-	1	-	-	-	-	-	3	3
Male	17	-	-	-	1	-	-	-	-	-	2	3
Female	13	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	24	-	-	-	1	-	-	-	-	-	3	1
Male	13	-	-	-	1	-	-	-	-	-	2	1
Female	11	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	2
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
New Mexico												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	64	2	-	-	-	-	-	-	1	2	2	4
Male	37	2	-	-	-	-	-	-	1	2	2	1
Female	27	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	52	2	-	-	-	-	-	-	1	2	2	3
Male	29	2	-	-	-	-	-	-	1	2	2	1
Female	23	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
North Carolina												
All races, both sexes	30	1	-	-	-	-	-	-	1	-	-	2
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	15	1	-	-	-	-	-	-	1	-	-	1
White, both sexes	21	-	-	-	-	-	-	-	-	-	-	2
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	8	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Ohio												
All races, both sexes	41	1	-	-	-	-	-	-	1	2	2	1
Male	20	-	-	-	-	-	-	-	-	2	1	1
Female	21	1	-	-	-	-	-	-	1	-	1	-
White, both sexes	37	1	-	-	-	-	-	-	1	2	2	1
Male	19	-	-	-	-	-	-	-	-	2	1	1
Female	18	1	-	-	-	-	-	-	1	-	1	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	18	-	-	-	-	-	-	-	1	-	-	1
Male	11	-	-	-	-	-	-	-	1	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	1	-	-	1
Male	9	-	-	-	-	-	-	-	1	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Oregon												
All races, both sexes	17	1	-	-	-	-	-	-	-	-	-	2
Male	9	-	-	-	-	-	-	-	-	-	-	2
Female	8	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	1	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	7	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	66	-	-	-	-	-	-	-	-	1	2	5
Male	34	-	-	-	-	-	-	-	-	1	2	3
Female	32	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	60	-	-	-	-	-	-	-	-	1	2	4
Male	33	-	-	-	-	-	-	-	-	1	2	3
Female	27	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Rhode Island												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	1	1
Male	11	-	-	-	-	-	-	-	-	-	1	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	1	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
South Dakota												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	21	1	-	-	-	-	-	-	1	-	2	2
Male	12	-	-	-	-	-	-	-	-	-	2	1
Female	9	1	-	-	-	-	-	-	1	-	-	1
White, both sexes	16	-	-	-	-	-	-	-	-	-	1	2
Male	10	-	-	-	-	-	-	-	-	-	1	1
Female	6	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	5	1	-	-	-	-	-	-	1	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	3	1	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Texas												
All races, both sexes	72	5	-	-	-	-	1	-	-	1	-	3
Male	44	3	-	-	-	-	1	-	-	1	-	3
Female	28	2	-	-	-	-	-	-	-	-	-	-
White, both sexes	60	3	-	-	-	-	1	-	-	1	-	3
Male	35	2	-	-	-	-	1	-	-	1	-	3
Female	25	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	11	2	-	-	-	-	-	-	-	-	-	-
Male	9	1	-	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Vermont												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	41	3	-	-	-	-	-	-	-	-	-	3
Male	25	2	-	-	-	-	-	-	-	-	-	3
Female	16	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	29	2	-	-	-	-	-	-	-	-	-	3
Male	19	1	-	-	-	-	-	-	-	-	-	3
Female	10	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	11	1	-	-	-	-	-	-	-	-	-	-
Male	5	1	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Washington												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	1	2
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	15	-	-	-	-	-	-	-	-	-	1	2
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	14	1	-	1	-	-	-	-	-	-	-	1
Male	8	1	-	1	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	1	-	1	-	-	-	-	-	-	-	1
Male	7	1	-	1	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Wisconsin												
All races, both sexes	20	1	-	-	1	-	-	-	-	-	1	1
Male	10	1	-	-	1	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	19	1	-	-	1	-	-	-	-	-	1	1
Male	10	1	-	-	1	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	3	-	1	-	-	-	-	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	1	-	-	-	-	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
United States												
All races, both sexes	71	70	68	65	84	111	130	145	98	42	15	-
Male	55	44	42	44	45	61	60	53	35	12	1	-
Female	16	26	26	21	39	50	70	92	63	30	14	-
White, both sexes	58	47	47	57	67	101	119	127	90	35	12	-
Male	45	26	30	39	36	55	53	48	32	11	1	-
Female	13	21	17	18	31	46	66	79	58	24	11	-
Black, both sexes	8	18	20	8	14	8	8	17	7	6	2	-
Male	6	14	12	5	8	6	5	5	2	-	-	-
Female	2	4	8	3	6	2	3	12	5	6	2	-
American Indian 1/, both sexes	3	-	-	-	2	1	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	2	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	5	1	-	1	1	3	1	1	1	1	-
Male	2	4	-	-	1	-	2	-	1	1	-	-
Female	-	1	1	-	-	1	1	1	-	-	1	-
Alabama												
All races, both sexes	4	5	1	3	1	1	4	4	-	2	-	-
Male	3	4	1	1	-	1	1	1	-	-	-	-
Female	1	1	-	2	1	-	3	3	-	2	-	-
White, both sexes	3	3	-	2	1	1	4	4	-	2	-	-
Male	2	2	-	-	-	1	1	1	-	-	-	-
Female	1	1	-	2	1	-	3	3	-	2	-	-
Black, both sexes	1	1	1	1	-	-	-	-	-	-	-	-
Male	1	1	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Alaska												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	1	2	-	1	1	2	1	3	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	1	1	-	1	-	2	1	3	-	-	-
White, both sexes	-	1	2	-	-	1	2	1	3	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	1	1	-	-	-	2	1	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Arkansas												
All races, both sexes	1	-	1	1	1	3	1	1	-	-	-	-
Male	1	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	1	1	1	2	-	1	-	-	-	-
White, both sexes	1	-	-	1	-	2	1	1	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	2	-	1	-	-	-	-
Black, both sexes	-	-	1	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	6	6	3	3	3	5	6	7	3	3	1	-
Male	4	4	2	2	-	1	3	3	1	-	-	-
Female	2	2	1	1	3	4	3	4	2	3	1	-
White, both sexes	5	5	3	3	3	4	6	7	3	3	-	-
Male	3	3	2	2	-	1	3	3	1	-	-	-
Female	2	2	1	1	3	3	3	4	2	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	1	-	-	-	-	1	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Colorado												
All races, both sexes	-	1	-	-	1	1	-	2	3	-	-	-
Male	-	1	-	-	1	1	-	1	2	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
White, both sexes	-	1	-	-	1	1	-	1	3	-	-	-
Male	-	1	-	-	1	1	-	1	2	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	-	1	1	-	1	1	1	1	1	-	-
Male	1	-	-	1	-	-	1	1	-	-	-	-
Female	-	-	1	-	-	1	-	-	1	1	-	-
White, both sexes	1	-	-	1	-	1	-	1	1	1	-	-
Male	1	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	1	-	-
Black, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Florida												
All races, both sexes	5	4	4	6	7	9	5	6	2	3	1	-
Male	4	3	2	5	5	5	3	1	2	2	-	-
Female	1	1	2	1	2	4	2	5	-	1	1	-
White, both sexes	4	4	3	6	7	8	5	6	2	3	1	-
Male	3	3	1	5	5	5	3	1	2	2	-	-
Female	1	1	2	1	2	3	2	5	-	1	1	-
Black, both sexes	1	-	1	-	-	1	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	5	3	2	3	4	3	4	4	3	3	-	-
Male	4	1	1	2	2	1	2	1	-	-	-	-
Female	1	2	1	1	2	2	2	3	3	3	-	-
White, both sexes	5	1	-	3	2	3	4	1	3	-	-	-
Male	4	-	-	2	1	1	2	-	-	-	-	-
Female	1	1	-	1	1	2	2	1	3	-	-	-
Black, both sexes	-	2	2	-	2	-	-	3	-	3	-	-
Male	-	1	1	-	1	-	-	1	-	-	-	-
Female	-	1	1	-	1	-	-	2	-	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Hawaii												
All races, both sexes	-	-	1	-	2	-	2	2	2	1	-	-
Male	-	-	-	-	1	-	1	-	1	1	-	-
Female	-	-	1	-	1	-	1	2	1	-	-	-
White, both sexes	-	-	-	-	1	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	-	1	1	1	1	-	-
Male	-	-	-	-	1	-	1	-	1	1	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-
Idaho												
All races, both sexes	-	1	-	1	-	-	1	-	2	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	2	-	-	-
White, both sexes	-	1	-	1	-	-	1	-	2	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Illinois												
All races, both sexes	2	3	4	4	6	3	6	5	10	1	2	-
Male	2	2	2	4	2	1	-	3	4	-	-	-
Female	-	1	2	-	4	2	6	2	6	1	2	-
White, both sexes	2	1	2	3	4	2	6	5	8	1	2	-
Male	2	-	1	3	-	1	-	3	3	-	-	-
Female	-	1	1	-	4	1	6	2	5	1	2	-
Black, both sexes	-	2	2	1	2	1	-	-	2	-	-	-
Male	-	2	1	1	2	-	-	-	1	-	-	-
Female	-	-	1	-	-	1	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	2	1	-	3	1	1	3	3	4	-	-	-
Male	2	1	-	1	1	-	1	2	1	-	-	-
Female	-	-	-	2	-	1	2	1	3	-	-	-
White, both sexes	2	1	-	3	1	1	3	3	3	-	-	-
Male	2	1	-	1	1	-	1	2	1	-	-	-
Female	-	-	-	2	-	1	2	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Iowa												
All races, both sexes	-	1	1	-	-	-	1	2	3	-	-	-
Male	-	1	-	-	-	-	-	1	1	-	-	-
Female	-	-	1	-	-	-	1	1	2	-	-	-
White, both sexes	-	1	1	-	-	-	1	2	3	-	-	-
Male	-	1	-	-	-	-	-	1	1	-	-	-
Female	-	-	1	-	-	-	1	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	1	-	-	3	-	2	1	-	1	-	-
Male	1	-	-	-	1	-	1	-	-	1	-	-
Female	-	1	-	-	2	-	1	1	-	-	-	-
White, both sexes	1	1	-	-	3	-	2	1	-	1	-	-
Male	1	-	-	-	1	-	1	-	-	1	-	-
Female	-	1	-	-	2	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Kentucky												
All races, both sexes	2	-	1	-	-	2	5	5	4	-	-	-
Male	2	-	-	-	-	2	2	1	-	-	-	-
Female	-	-	1	-	-	-	3	4	4	-	-	-
White, both sexes	2	-	1	-	-	2	4	5	3	-	-	-
Male	2	-	-	-	-	2	1	1	-	-	-	-
Female	-	-	1	-	-	-	3	4	3	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	1	-	1	2	-	1	5	-	1	1	-
Male	1	1	-	1	1	-	-	2	-	1	-	-
Female	-	-	-	-	1	-	1	3	-	-	1	-
White, both sexes	1	-	-	-	2	-	1	4	-	1	1	-
Male	1	-	-	-	1	-	-	2	-	1	-	-
Female	-	-	-	-	1	-	1	2	-	-	1	-
Black, both sexes	-	1	-	1	-	-	-	1	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Maine												
All races, both sexes	1	-	2	-	-	1	-	2	-	1	-	-
Male	1	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	1	-	1	-	1	-	-
White, both sexes	1	-	2	-	-	1	-	2	-	1	-	-
Male	1	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	1	-	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	2	2	2	2	2	1	4	4	1	-	-	-
Male	-	2	1	1	1	1	3	1	-	-	-	-
Female	2	-	1	1	1	-	1	3	1	-	-	-
White, both sexes	1	1	1	2	2	1	3	3	1	-	-	-
Male	-	1	1	1	1	1	2	1	-	-	-	-
Female	1	-	-	1	1	-	1	2	1	-	-	-
Black, both sexes	1	1	1	-	-	-	1	1	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	1	-	1	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Massachusetts												
All races, both sexes	-	2	2	2	1	1	2	4	1	2	1	-
Male	-	-	1	2	1	1	-	3	-	-	1	-
Female	-	2	1	-	-	-	2	1	1	2	-	-
White, both sexes	-	2	2	2	-	1	1	4	1	2	1	-
Male	-	-	1	2	-	1	-	3	-	-	1	-
Female	-	2	1	-	-	-	1	1	1	2	-	-
Black, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	2	2	3	1	3	7	7	3	5	1	-	-
Male	2	2	3	1	1	4	3	1	2	1	-	-
Female	-	-	-	-	2	3	4	2	3	-	-	-
White, both sexes	2	-	3	1	3	7	7	3	5	1	-	-
Male	2	-	3	1	1	4	3	1	2	1	-	-
Female	-	-	-	-	2	3	4	2	3	-	-	-
Black, both sexes	-	2	-	-	-	-	-	-	-	-	-	-
Male	-	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Minnesota												
All races, both sexes	1	2	-	2	-	-	2	1	4	2	-	-
Male	-	2	-	1	-	-	1	-	2	-	-	-
Female	1	-	-	1	-	-	1	1	2	2	-	-
White, both sexes	1	2	-	2	-	-	2	1	4	2	-	-
Male	-	2	-	1	-	-	1	-	2	-	-	-
Female	1	-	-	1	-	-	1	1	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1	1	-	-	1	2	2	-	2	-	-	-
Male	1	-	-	-	-	1	1	-	1	-	-	-
Female	-	1	-	-	1	1	1	-	1	-	-	-
White, both sexes	-	-	-	-	1	2	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	1	-	-	-
Black, both sexes	1	1	-	-	-	-	2	-	1	-	-	-
Male	1	-	-	-	-	-	1	-	1	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Missouri												
All races, both sexes	2	2	2	2	2	4	-	9	2	3	1	-
Male	1	2	1	1	1	2	-	6	2	1	-	-
Female	1	-	1	1	1	2	-	3	-	2	1	-
White, both sexes	1	2	2	2	2	4	-	9	2	3	1	-
Male	1	2	1	1	1	2	-	6	2	1	-	-
Female	-	-	1	1	1	2	-	3	-	2	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	-	-	-	-	-	1	2	1	-	-	-
Male	-	-	-	-	-	-	1	2	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	1	-	-	-	-	-	1	2	1	-	-	-
Male	-	-	-	-	-	-	1	2	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Nebraska												
All races, both sexes	-	1	1	-	1	1	1	2	3	1	-	-
Male	-	-	1	-	-	1	1	-	1	-	-	-
Female	-	1	-	-	1	-	-	2	2	1	-	-
White, both sexes	-	1	1	-	1	1	1	2	3	1	-	-
Male	-	-	1	-	-	1	1	-	1	-	-	-
Female	-	1	-	-	1	-	-	2	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	2	1	-	-	2	-	1	-	-	-	-
Male	-	-	1	-	-	2	-	1	-	-	-	-
Female	-	2	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	1	-	-	2	-	1	-	-	-	-
Male	-	-	1	-	-	2	-	1	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
New Hampshire												
All races, both sexes	-	-	-	-	1	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	1	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	2	1	2	3	6	-	2	5	2	-	-	-
Male	2	-	2	1	3	-	1	1	1	-	-	-
Female	-	1	-	2	3	-	1	4	1	-	-	-
White, both sexes	1	1	1	1	6	-	2	5	2	-	-	-
Male	1	-	1	1	3	-	1	1	1	-	-	-
Female	-	1	-	-	3	-	1	4	1	-	-	-
Black, both sexes	-	-	1	2	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	2	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
New Mexico												
All races, both sexes	-	1	1	-	-	-	2	3	-	1	-	-
Male	-	1	1	-	-	-	1	2	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	1	-	-
White, both sexes	-	1	1	-	-	-	2	3	-	1	-	-
Male	-	1	1	-	-	-	1	2	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	3	6	4	3	3	8	11	8	2	3	2	-
Male	3	6	3	2	1	5	6	2	1	-	-	-
Female	-	-	1	1	2	3	5	6	1	3	2	-
White, both sexes	2	2	3	3	2	6	11	6	2	3	2	-
Male	2	2	2	2	1	3	6	2	1	-	-	-
Female	-	-	1	1	1	3	5	4	1	3	2	-
Black, both sexes	1	3	1	-	1	2	-	2	-	-	-	-
Male	1	3	1	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
North Carolina												
All races, both sexes	4	1	3	2	6	3	1	4	1	1	-	-
Male	4	1	2	2	3	1	1	-	-	-	-	-
Female	-	-	1	-	3	2	-	4	1	1	-	-
White, both sexes	4	-	2	1	4	3	1	3	1	-	-	-
Male	4	-	1	1	2	1	1	-	-	-	-	-
Female	-	-	1	-	2	2	-	3	1	-	-	-
Black, both sexes	-	1	1	1	2	-	-	1	-	1	-	-
Male	-	1	1	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	1	-	-	1	4	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	4	-	-	-	-
White, both sexes	-	-	-	1	-	-	1	4	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	4	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Ohio												
All races, both sexes	1	3	3	3	1	6	7	4	3	2	1	-
Male	-	-	2	3	1	4	2	2	1	1	-	-
Female	1	3	1	-	-	2	5	2	2	1	1	-
White, both sexes	1	2	2	3	1	6	6	3	3	2	1	-
Male	-	-	2	3	1	4	2	1	1	1	-	-
Female	1	2	-	-	-	2	4	2	2	1	1	-
Black, both sexes	-	1	1	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	1	1	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2	-	1	-	1	2	1	5	3	1	-	-
Male	2	-	1	-	1	1	-	1	2	1	-	-
Female	-	-	-	-	-	1	1	4	1	-	-	-
White, both sexes	-	-	1	-	1	1	1	5	3	1	-	-
Male	-	-	1	-	1	1	-	1	2	1	-	-
Female	-	-	-	-	-	-	1	4	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	1	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Oregon												
All races, both sexes	2	-	-	-	3	1	2	1	3	1	1	-
Male	1	-	-	-	2	-	1	1	1	1	-	-
Female	1	-	-	-	1	1	1	-	2	-	1	-
White, both sexes	1	-	-	-	3	1	2	1	3	1	1	-
Male	1	-	-	-	2	-	1	1	1	1	-	-
Female	-	-	-	-	1	1	1	-	2	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	3	-	8	5	4	11	8	9	5	3	2	-
Male	3	-	4	3	3	6	4	3	1	1	-	-
Female	-	-	4	2	1	5	4	6	4	2	2	-
White, both sexes	3	-	6	5	4	11	8	8	5	2	1	-
Male	3	-	3	3	3	6	4	3	1	1	-	-
Female	-	-	3	2	1	5	4	5	4	1	1	-
Black, both sexes	-	-	2	-	-	-	-	1	-	1	1	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Rhode Island												
All races, both sexes	-	-	-	-	1	-	1	-	1	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	2	1	2	2	1	1	2	3	2	-	-	-
Male	1	-	2	1	1	1	2	1	-	-	-	-
Female	1	1	-	1	-	-	-	2	2	-	-	-
White, both sexes	2	1	-	1	-	1	2	-	2	-	-	-
Male	1	-	-	1	-	1	2	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	2	-	-	-
Black, both sexes	-	-	2	1	1	-	-	3	-	-	-	-
Male	-	-	2	-	1	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
South Dakota												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	1	1	2	2	3	2	2	2	-	-	-
Male	-	1	1	2	1	-	2	-	2	-	-	-
Female	-	-	-	-	1	3	-	2	-	-	-	-
White, both sexes	-	1	1	1	2	3	2	1	2	-	-	-
Male	-	1	1	1	1	-	2	-	2	-	-	-
Female	-	-	-	-	1	3	-	1	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Texas												
All races, both sexes	7	7	4	4	4	9	12	6	7	2	-	-
Male	5	5	3	3	4	7	2	5	1	1	-	-
Female	2	2	1	1	-	2	10	1	6	1	-	-
White, both sexes	7	6	3	4	2	6	11	5	6	2	-	-
Male	5	4	2	3	2	4	2	4	1	1	-	-
Female	2	2	1	1	-	2	9	1	5	1	-	-
Black, both sexes	-	1	1	-	2	3	-	1	1	-	-	-
Male	-	1	1	-	2	3	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	1	1	1	3	-	-	-	-
Male	-	-	-	-	1	1	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
White, both sexes	-	-	-	-	1	1	1	3	-	-	-	-
Male	-	-	-	-	1	1	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Vermont												
All races, both sexes	-	-	1	-	-	-	1	1	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	1	-	-	-	1	1	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	2	4	2	3	2	7	5	3	3	2	2	-
Male	1	2	1	2	1	5	5	1	2	-	-	-
Female	1	2	1	1	1	2	-	2	1	2	2	-
White, both sexes	1	1	1	3	1	7	4	2	2	1	1	-
Male	-	-	1	2	1	5	4	-	2	-	-	-
Female	1	1	-	1	-	2	-	2	-	1	1	-
Black, both sexes	1	2	1	-	1	-	1	1	1	1	1	-
Male	1	1	-	-	-	-	1	1	-	-	-	-
Female	-	1	1	-	1	-	-	-	1	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Washington												
All races, both sexes	2	1	-	1	-	2	1	5	1	-	-	-
Male	2	1	-	1	-	1	1	1	-	-	-	-
Female	-	-	-	-	-	1	-	4	1	-	-	-
White, both sexes	1	1	-	1	-	2	1	5	1	-	-	-
Male	1	1	-	1	-	1	1	1	-	-	-	-
Female	-	-	-	-	-	1	-	4	1	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	1	1	-	3	1	3	1	1	-	-	-
Male	-	-	1	-	2	-	1	-	1	-	-	-
Female	-	1	-	-	1	1	2	1	-	-	-	-
White, both sexes	-	1	-	-	2	1	3	1	1	-	-	-
Male	-	-	-	-	2	-	1	-	1	-	-	-
Female	-	1	-	-	-	1	2	1	-	-	-	-
Black, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Wisconsin												
All races, both sexes	-	-	-	1	3	6	3	-	3	-	-	-
Male	-	-	-	-	2	2	1	-	2	-	-	-
Female	-	-	-	1	1	4	2	-	1	-	-	-
White, both sexes	-	-	-	1	2	6	3	-	3	-	-	-
Male	-	-	-	-	2	2	1	-	2	-	-	-
Female	-	-	-	1	-	4	2	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-