

Table. Estimated number of male newborn infants discharged from short-stay hospitals, and percentage circumcised during birth hospitalization, by geographic region: United States, 1979-2010

Year	All regions			Northeast			Midwest			South			West		
	Male newborns	Percent circumcised	SE	Male newborns	Percent circumcised	SE	Male newborns	Percent circumcised	SE	Male newborns	Percent circumcised	SE	Male newborns	Percent circumcised	SE
1979	1,906,000	64.5	2.7	375,000	66.2	3.5	534,000	74.3	2.5	642,000	55.8	6.3	355,000	63.9	3.1
1980	1,950,000	64.7	2.5	376,000	67.4	2.7	531,000	75.9	2.9	676,000	56.0	5.4	366,000	61.8	3.6
1981	1,987,000	64.9	2.5	383,000	66.5	2.6	543,000	77.5	2.4	663,000	58.4	6.0	398,000	57.2	3.8
1982	2,020,000	62.5	2.5	380,000	63.4	2.7	515,000	75.9	2.4	707,000	57.3	5.9	417,000	54.3	4.3
1983	1,972,000	63.2	3.0	374,000	69.2	2.7	482,000	74.9	3.4	712,000	58.1	6.8	403,000	52.7	4.7
1984	2,003,000	62.4	2.7	356,000	68.2	3.1	488,000	75.1	2.5	761,000	58.3	6.0	398,000	49.4	4.5
1985	1,953,000	59.5	2.9	339,000	65.2	3.6	469,000	70.5	2.9	701,000	56.0	6.5	445,000	49.0	5.3
1986	1,954,000	59.4	2.2	360,000	67.9	3.1	472,000	71.8	2.6	656,000	57.5	4.4	466,000	43.0	5.6
1987	2,033,000	58.6	2.7	369,000	68.6	2.8	480,000	73.1	2.8	683,000	54.7	6.1	501,000	42.4	5.5
1988	1,882,000	58.0	2.2	373,000	63.2	2.9	443,000	72.6	4.1	614,000	53.8	3.3	452,000	45.2	5.9
1989	1,989,000	58.8	2.5	392,000	63.2	3.6	481,000	74.0	4.0	644,000	57.9	3.2	472,000	41.0	6.3
1990	1,982,000	59.0	2.2	379,000	62.6	3.6	449,000	76.0	2.9	693,000	57.1	3.3	461,000	42.4	5.5
1991	2,003,000	61.1	1.9	366,000	62.4	3.2	465,000	78.2	2.6	685,000	63.6	3.1	487,000	40.4	4.5
1992	1,926,000	60.7	1.8	365,000	67.6	2.4	424,000	78.2	2.8	656,000	62.6	3.3	482,000	37.5	4.0
1993	1,823,000	59.5	2.0	350,000	65.2	2.9	408,000	75.6	2.9	651,000	61.4	3.2	414,000	35.5	5.0
1994	1,935,000	62.0	1.9	377,000	69.6	2.7	431,000	79.9	1.9	681,000	64.7	3.0	445,000	34.2	4.5
1995	1,872,000	64.1	1.9	364,000	68.3	2.2	397,000	79.8	2.2	688,000	66.1	3.5	423,000	42.6	4.7
1996	1,999,000	60.2	2.3	331,000	66.5	3.2	422,000	80.9	2.4	696,000	63.6	3.3	550,000	36.3	5.0
1997	1,931,000	62.8	2.1	296,000	68.3	2.2	440,000	81.6	2.2	740,000	64.5	3.6	454,000	38.0	5.3
1998	1,976,000	63.2	1.9	382,000	68.0	2.6	440,000	82.9	1.6	689,000	64.6	3.6	465,000	38.3	4.7
1999	1,893,000	61.5	1.7	348,000	65.4	2.5	416,000	81.4	1.3	673,000	64.1	2.9	456,000	36.7	4.6
2000	1,869,000	62.4	1.9	396,000	64.6	2.7	405,000	81.4	1.7	686,000	63.9	3.1	382,000	37.3	5.6
2001	1,866,000	63.1	2.0	382,000	66.9	2.8	395,000	81.0	1.5	723,000	62.5	3.6	365,000	40.9	5.9
2002	1,940,000	60.1	2.2	369,000	68.9	2.6	385,000	81.0	1.7	679,000	64.0	3.7	507,000	32.6	5.0
2003	2,018,000	55.9	2.3	364,000	64.7	3.0	397,000	77.8	1.8	719,000	57.7	4.0	538,000	31.4	4.9
2004	2,020,000	57.4	2.2	363,000	66.4	2.1	419,000	79.5	2.2	724,000	58.5	3.9	515,000	31.7	5.1
2005	2,044,000	57.3	2.1	379,000	66.9	2.1	420,000	78.7	2.1	719,000	58.7	3.0	526,000	31.5	4.9
2006	2,041,000	56.1	2.2	352,000	63.6	1.9	424,000	77.9	1.9	760,000	55.3	3.4	505,000	33.8	4.9
2007	2,104,000	55.4	2.0	364,000	60.7	3.0	443,000	79.6	2.3	775,000	54.9	3.2	521,000	32.0	4.3
2008	2,138,000	55.9	4.1	340,000	65.3	4.3	420,000	71.1	5.2	824,000	54.6	4.7	555,000	40.5	11.9
2009	2,100,000	56.2	3.8	345,000	65.4	3.1	450,000	68.8	6.2	801,000	56.0	4.5	504,000	39.0	11.7
2010	2,029,000	58.3	3.6	350,000	66.3	4.0	421,000	71.0	4.1	799,000	58.4	4.8	460,000	40.2	11.7

NOTES: Circumcision is identified by *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM) procedure code 64.0. SE is standard error of the percentage. Estimated numbers are rounded to thousands.

SOURCE: CDC/NCHS, National Hospital Discharge Survey, 1979-2010.