

Table 1. Annual number and percent distribution of ambulatory care visits by setting type, according to selected patient and visit characteristics: United States, 2004

| Characteristic | Combined settings | Primary care offices | Surgical specialty offices | Medical specialty offices | Hospital outpatient departments | Hospital emergency departments |
|---|-------------------------------|----------------------|----------------------------|---------------------------|---------------------------------|--------------------------------|
| | Number of visits in thousands | | | | | |
| All visits | 1,106,067 | 532,420 | 176,431 | 202,006 | 84,994 | 110,216 |
| Patient age | | | | | | |
| Under 15 years | 189,644 | 120,924 | 11,913 | 15,073 | 18,792 | 22,942 |
| Under 1 year | 34,525 | 25,569 | 823 | * | 3,516 | 3,902 |
| 1-4 years | 59,015 | 38,224 | 3,366 | *3,068 | 6,038 | 8,318 |
| 5-14 years | 96,104 | 57,131 | 7,724 | 11,289 | 9,238 | 10,722 |
| 15-24 years | 98,478 | 47,555 | 9,482 | 13,556 | 9,953 | 17,931 |
| 25-44 years | 247,476 | 122,257 | 32,395 | 39,609 | 21,110 | 32,105 |
| 45-64 years | 307,736 | 138,254 | 58,074 | 67,775 | 22,127 | 21,506 |
| 65 years and over | 262,734 | 103,430 | 64,568 | 65,993 | 13,011 | 15,732 |
| 65-74 years | 126,802 | 49,045 | 31,309 | 33,072 | 7,157 | 6,219 |
| 75 years and over | 135,931 | 54,385 | 33,259 | 32,921 | 5,854 | 9,513 |
| Patient sex | | | | | | |
| Female | 647,205 | 328,355 | 96,720 | 110,467 | 51,767 | 59,896 |
| Male | 458,863 | 204,065 | 79,712 | 91,540 | 33,226 | 50,320 |
| Patient race | | | | | | |
| White | 919,628 | 439,306 | 155,925 | 179,788 | 62,847 | 81,762 |
| Black or African American | 141,075 | 67,405 | 14,048 | 16,548 | 18,176 | 24,898 |
| Asian | 34,141 | 19,348 | 5,385 | 4,398 | 2,853 | 2,157 |
| Native Hawaiian or other Pacific Islander | 3,534 | 2,182 | * | * | 270 | 396 |
| American Indian or Alaska Native | 3,834 | *1,429 | * | *757 | *531 | *688 |
| Two or more races reported | 3,856 | 2,749 | * | * | 317 | *316 |
| Patient ethnicity | | | | | | |
| Hispanic or Latino | 120,868 | 63,478 | 15,552 | 13,341 | 14,348 | 14,149 |
| Not Hispanic or Latino | 985,200 | 468,942 | 160,880 | 188,666 | 70,646 | 96,067 |
| Primary expected source of payment | | | | | | |
| Private insurance | 580,097 | 322,829 | 84,231 | 102,712 | 30,980 | 39,344 |
| Medicare | 236,748 | 91,585 | 55,565 | 59,540 | 13,149 | 16,909 |
| Medicaid or SCHIP ¹ | 140,301 | 66,149 | 10,220 | 13,278 | 26,165 | 24,489 |
| Self-pay, no charge, or charity | 69,927 | 20,609 | 11,089 | 11,703 | 7,972 | 18,553 |
| Other | 78,994 | 31,247 | 15,326 | 14,773 | 6,727 | 10,920 |
| Geographic region of provider | | | | | | |
| Northeast | 214,820 | 96,420 | 34,353 | 39,126 | 22,647 | 22,274 |
| Midwest | 250,237 | 117,004 | 42,436 | 38,067 | 25,924 | 26,806 |
| South | 419,914 | 206,833 | 67,034 | 79,985 | 24,912 | 41,150 |
| West | 221,096 | 112,163 | 32,608 | 44,829 | 11,510 | 19,986 |
| MSA ² status of provider | | | | | | |
| MSA | 957,764 | 459,847 | 153,430 | 177,352 | 72,308 | 94,826 |
| Not MSA | 148,304 | 72,573 | 23,001 | 24,654 | 12,686 | 15,391 |

| | | Percent distribution | | | | |
|---|-------|----------------------|------|-------|------|-------|
| All visits | 100.0 | 48.1 | 16.0 | 18.3 | 7.7 | 10.0 |
| Patient age | | | | | | |
| Under 15 years | 100.0 | 63.8 | 6.3 | 8.0 | 9.9 | 12.1 |
| Under 1 year | 100.0 | 74.1 | 2.4 | * | 10.2 | 11.3 |
| 1-4 years | 100.0 | 64.8 | 5.7 | *5.20 | 10.2 | 14.1 |
| 5-14 years | 100.0 | 59.5 | 8.0 | 11.8 | 9.6 | 11.2 |
| 15-24 years | 100.0 | 48.3 | 9.6 | 13.8 | 10.1 | 18.2 |
| 25-44 years | 100.0 | 49.4 | 13.1 | 16.0 | 8.5 | 13.0 |
| 45-64 years | 100.0 | 44.9 | 18.9 | 22.0 | 7.2 | 7.0 |
| 65 years and over | 100.0 | 39.4 | 24.6 | 25.1 | 5.0 | 6.0 |
| 65-74 years | 100.0 | 38.7 | 24.7 | 26.1 | 5.6 | 4.9 |
| 75 years and over | 100.0 | 40.0 | 24.5 | 24.2 | 4.3 | 7.0 |
| Patient sex | | | | | | |
| Female | 100.0 | 50.7 | 14.9 | 17.1 | 8.0 | 9.3 |
| Male | 100.0 | 44.5 | 17.4 | 20.0 | 7.2 | 11.0 |
| Patient race | | | | | | |
| White | 100.0 | 47.8 | 17.0 | 19.6 | 6.8 | 8.9 |
| Black or African American | 100.0 | 47.8 | 10.0 | 11.7 | 12.9 | 17.7 |
| Asian | 100.0 | 56.7 | 15.8 | 12.9 | 8.4 | 6.3 |
| Native Hawaiian or other Pacific Islander | 100.0 | 61.8 | * | * | 7.6 | 11.2 |
| American Indian or Alaska Native | 100.0 | 37.3 | * | 19.8 | 13.9 | 18.0 |
| Two or more races reported | 100.0 | 71.3 | * | * | 8.2 | *8.19 |
| Patient ethnicity | | | | | | |
| Hispanic or Latino | 100.0 | 52.5 | 12.9 | 11.0 | 11.9 | 11.7 |
| Not Hispanic or Latino | 100.0 | 47.6 | 16.3 | 19.2 | 7.2 | 9.8 |
| Primary expected source of payment | | | | | | |
| Private insurance | 100.0 | 55.7 | 14.5 | 17.7 | 5.3 | 6.8 |
| Medicare | 100.0 | 38.7 | 23.5 | 25.2 | 5.6 | 7.1 |
| Medicaid or SCHIP ¹ | 100.0 | 47.2 | 7.3 | 9.5 | 18.7 | 17.5 |
| Self-pay, no charge, or charity | 100.0 | 29.5 | 15.9 | 16.7 | 11.4 | 26.5 |
| Other | 100.0 | 39.6 | 19.4 | 18.7 | 8.5 | 13.8 |
| Geographic region of provider | | | | | | |
| Northeast | 100.0 | 44.9 | 16.0 | 18.2 | 10.5 | 10.4 |
| Midwest | 100.0 | 46.8 | 17.0 | 15.2 | 10.4 | 10.7 |
| South | 100.0 | 49.3 | 16.0 | 19.1 | 5.9 | 9.8 |
| West | 100.0 | 50.7 | 14.8 | 20.3 | 5.2 | 9.0 |
| MSA ² status of provider | | | | | | |
| MSA | 100.0 | 48.0 | 16.0 | 18.5 | 7.6 | 9.9 |
| Not MSA | 100.0 | 48.9 | 15.5 | 16.6 | 8.6 | 10.4 |

* Figure does not meet standard of reliability of precision.

¹SCHIP is State Children's Health Insurance Program.

²MSA is metropolitan statistical area.

Table 2. Annual rate of ambulatory care visits with corresponding standard errors, by setting type and selected patient and provider characteristics: United States, 2004

| Characteristic | Combined settings | Primary care offices | Surgical specialty offices | Medical specialty offices | Hospital outpatient departments | Hospital emergency departments |
|---|---|----------------------|----------------------------|---------------------------|---------------------------------|--------------------------------|
| | Number of visits per 100 persons ^{1,2,3} | | | | | |
| All visits | 383.6 | 184.6 | 61.2 | 70.0 | 29.5 | 38.2 |
| Patient age | | | | | | |
| Under 15 years | 312.1 | 199.0 | 19.6 | 24.8 | 30.9 | 37.8 |
| Under 1 year | 847.5 | 627.7 | 20.2 | * | 86.3 | 95.8 |
| 1-4 years | 369.0 | 239.0 | 21.0 | *19.2 | 37.8 | 52.0 |
| 5-14 years | 236.1 | 140.4 | 19.0 | 27.7 | 22.7 | 26.3 |
| 15-24 years | 242.4 | 117.1 | 23.3 | 33.4 | 24.5 | 44.1 |
| 25-44 years | 301.3 | 148.9 | 39.4 | 48.2 | 25.7 | 39.1 |
| 45-64 years | 438.4 | 196.9 | 82.7 | 96.5 | 31.5 | 30.6 |
| 65 years and over | 758.2 | 298.5 | 186.3 | 190.4 | 37.5 | 45.4 |
| 65-74 years | 696.0 | 269.2 | 171.8 | 181.5 | 39.3 | 34.1 |
| 75 years and over | 827.1 | 330.9 | 202.4 | 200.3 | 35.6 | 57.9 |
| Patient sex | | | | | | |
| Female | 439.0 | 222.7 | 65.6 | 74.9 | 35.1 | 40.6 |
| Male | 325.5 | 144.8 | 56.5 | 64.9 | 23.6 | 35.7 |
| Patient race | | | | | | |
| White | 395.8 | 189.1 | 67.1 | 77.4 | 27.0 | 35.2 |
| Black or African American | 390.5 | 186.6 | 38.9 | 45.8 | 50.3 | 68.9 |
| Asian | 278.6 | 157.9 | 43.9 | 35.9 | 23.3 | 17.6 |
| Native Hawaiian or other Pacific Islander | 711.7 | 439.5 | * | * | 54.3 | 79.7 |
| American Indian or Alaska Native | 138.4 | *51.6 | * | *27.3 | *19.2 | *24.8 |
| Two or more races reported | 87.9 | 62.7 | * | * | 7.2 | *7.2 |
| Patient ethnicity | | | | | | |
| Hispanic or Latino | 296.2 | 155.6 | 38.1 | 32.7 | 35.2 | 34.7 |
| Not Hispanic or Latino | 397.9 | 189.4 | 65.0 | 76.2 | 28.5 | 38.8 |
| Primary expected source of payment | | | | | | |
| Private insurance | 299.7 | 166.8 | 43.5 | 53.1 | 16.0 | 20.3 |
| Medicare | 659.1 | 255.0 | 154.7 | 165.8 | 36.6 | 47.1 |
| Medicaid or SCHIP ⁴ | 460.2 | 217.0 | 33.5 | 43.6 | 85.8 | 80.3 |
| Self-pay, no charge, or charity | 168.1 | 49.5 | 26.7 | 28.1 | 19.2 | 44.6 |
| Geographic region of provider | | | | | | |
| Northeast | 399.7 | 179.4 | 63.9 | 72.8 | 42.1 | 41.4 |
| Midwest | 386.7 | 180.8 | 65.6 | 58.8 | 40.1 | 41.4 |
| South | 405.2 | 199.6 | 64.7 | 77.2 | 24.0 | 39.7 |
| West | 333.5 | 169.2 | 49.2 | 67.6 | 17.4 | 30.1 |
| MSA ⁵ status of provider | | | | | | |
| MSA | 393.8 | 189.1 | 63.1 | 72.9 | 29.7 | 39.0 |
| Not MSA | 328.3 | 160.6 | 50.9 | 54.6 | 28.1 | 34.1 |

| | Standard error of rate | | | | | |
|---|------------------------|-------|------|------|------|------|
| All visits | 12.5 | 8.3 | 3.9 | 6.2 | 2.6 | 1.9 |
| Patient age | | | | | | |
| Under 15 years | 18.6 | 15.3 | 3.1 | 7.0 | 3.9 | 2.8 |
| Under 1 year | 60.0 | 56.5 | 5.9 | ... | 11.7 | 9.7 |
| 1-4 years | 26.7 | 21.5 | 4.3 | 8.9 | 5.4 | 4.1 |
| 5-14 years | 14.5 | 11.8 | 2.8 | 6.7 | 3.0 | 1.8 |
| 15-24 years | 11.5 | 8.8 | 2.5 | 5.9 | 2.3 | 2.5 |
| 25-44 years | 12.6 | 9.9 | 3.3 | 4.6 | 2.3 | 2.0 |
| 45-64 years | 17.4 | 11.8 | 5.3 | 8.8 | 3.2 | 1.5 |
| 65 years and over | 36.4 | 23.0 | 15.4 | 18.0 | 4.2 | 2.3 |
| 65-74 years | 37.1 | 24.1 | 14.8 | 19.2 | 4.4 | 1.9 |
| 75 years and over | 40.2 | 26.0 | 17.4 | 18.5 | 4.5 | 3.1 |
| Patient sex | | | | | | |
| Female | 14.5 | 10.6 | 4.6 | 6.6 | 3.2 | 2.1 |
| Male | 12.0 | 7.6 | 3.6 | 6.1 | 2.2 | 1.7 |
| Patient race | | | | | | |
| White | 14.6 | 9.1 | 4.5 | 7.1 | 2.8 | 1.9 |
| Black or African American | 20.9 | 18.5 | 5.1 | 5.8 | 5.4 | 4.8 |
| Asian | 33.7 | 30.0 | 10.6 | 8.8 | 3.8 | 1.9 |
| Native Hawaiian or other Pacific Islander | 152.2 | 113.0 | ... | ... | 16.1 | 23.8 |
| American Indian or Alaska Native | 28.1 | 21.8 | ... | 8.2 | 7.1 | 8.1 |
| Two or more races reported | 18.8 | 18.2 | ... | ... | 1.7 | 2.3 |
| Patient ethnicity | | | | | | |
| Hispanic or Latino | 26.3 | 16.3 | 9.5 | 7.7 | 5.1 | 2.7 |
| Not Hispanic or Latino | 13.4 | 9.2 | 4.3 | 6.5 | 2.8 | 2.0 |
| Primary expected source of payment | | | | | | |
| Private insurance | 11.8 | 8.7 | 2.7 | 5.6 | 2.0 | 1.1 |
| Medicare | 32.7 | 21.1 | 13.7 | 15.7 | 4.0 | 2.5 |
| Medicaid or SCHIP ⁴ | 28.3 | 21.4 | 5.8 | 7.9 | 9.2 | 5.3 |
| Self-pay, no charge, or charity | 10.6 | 6.3 | 7.1 | 4.0 | 3.1 | 3.3 |
| Geographic region of provider | | | | | | |
| Northeast | 33.0 | 18.2 | 8.2 | 16.9 | 6.2 | 4.1 |
| Midwest | 25.5 | 16.6 | 10.1 | 9.1 | 7.2 | 4.1 |
| South | 23.8 | 16.6 | 7.0 | 12.2 | 4.1 | 3.8 |
| West | 18.1 | 16.7 | 5.5 | 10.2 | 4.2 | 2.8 |
| MSA ⁵ status of provider | | | | | | |
| MSA | 14.7 | 9.0 | 4.4 | 7.1 | 3.0 | 2.2 |
| Not MSA | 42.5 | 26.3 | 12.0 | 13.5 | 7.1 | 5.5 |

* Figure does not meet standard of reliability or precision.

... Category not applicable.

¹ Visit rates for age, sex, race, and region are based on the July 1, 2004 set of estimates of the civilian noninstitutional population of the United States as developed by the Population Division, U.S. Census Bureau. These population estimates reflect Census 2000 data and are available from the U.S. Census Bureau.

² Population estimates of metropolitan statistical area status are based on data from the 2004 National Health Interview Survey, National Center for Health Statistics, adjusted to the U.S. Census Bureau definition of core-based statistical areas as of December 2003. See <http://www.census.gov/population/www/estimates.metrodef.html> for more about metropolitan statistical definitions.

³ Population estimates for expected source of payment are based on the data from the 2004 National Health Interview Survey.

⁴ SCHIP is State Children's Health Insurance Program.

⁵ MSA is metropolitan statistical area.

Table 3. Annual number and percent distribution of ambulatory care visits with corresponding standard errors, by primary diagnosis group, with percent distribution by setting type: United States, 2004

| Primary diagnosis group and ICD-9-CM codes(s) ¹ | Number of visits in thousands | Percent distribution | Total | Primary care offices | Surgical specialty offices | Medical specialty offices | Hospital outpatient departments | Hospital emergency departments | Percent distribution | | | | | |
|--|---|----------------------|-------|----------------------|----------------------------|---------------------------|---------------------------------|--------------------------------|----------------------|--|--|--|--|--|
| | | | | | | | | | | | | | | |
| All visits | 1,106,067 | 100.0 | 100.0 | 48.1 | 16.0 | 18.3 | 7.7 | 10.0 | | | | | | |
| Essential hypertension | 401 | 42,067 | 3.8 | 100.0 | 81.3 | * | 8.0 | 8.3 | 1.8 | | | | | |
| Routine infant or child health check | V20.2 | 34,926 | 3.2 | 100.0 | 89.4 | - | * | 10.0 | * | | | | | |
| Acute upper respiratory infections, excluding pharyngitis | 460-461,463-466 | 33,854 | 3.1 | 100.0 | 76.7 | 1.6 | * | 8.3 | 9.9 | | | | | |
| Malignant neoplasms | 140-208,230-234 | 30,439 | 2.8 | 100.0 | *8.5 | 15.3 | 67.5 | *8.3 | 0.5 | | | | | |
| Diabetes mellitus | 250 | 30,286 | 2.7 | 100.0 | 68.5 | 10.3 | *10.9 | 9.0 | 1.3 | | | | | |
| Spinal disorders | 720-724 | 28,908 | 2.6 | 100.0 | 51.1 | 18.5 | 13.3 | 7.0 | 10.0 | | | | | |
| Arthropathies and related disorders | 710-719 | 28,191 | 2.5 | 100.0 | 43.0 | 39.2 | 5.5 | 7.4 | 5.0 | | | | | |
| Normal pregnancy | V22 | 26,379 | 2.4 | 100.0 | 88.0 | - | - | 10.9 | *1.0 | | | | | |
| Rheumatism, excluding back | 725-729 | 21,679 | 2.0 | 100.0 | 42.9 | 31.1 | *11.6 | 6.3 | 8.1 | | | | | |
| Specific procedures and aftercare | V50-V59.9 | 18,165 | 1.6 | 100.0 | 48.5 | 22.3 | 13.9 | 9.5 | 5.7 | | | | | |
| Asthma | 493 | 16,500 | 1.5 | 100.0 | 54.1 | * | *27.9 | 6.4 | 11.1 | | | | | |
| Gynecological examination | V72.3 | 16,188 | 1.5 | 100.0 | 93.7 | * | - | 6.3 | - | | | | | |
| General medical examination | V70 | 15,311 | 1.4 | 100.0 | 83.3 | * | *7.7 | 6.4 | 1.9 | | | | | |
| Otitis media and eustachian tube disorders | 381-382 | 15,215 | 1.4 | 100.0 | 59.5 | 16.1 | * | 9.1 | 13.8 | | | | | |
| Allergic rhinitis | 477 | 14,885 | 1.3 | 100.0 | 35.8 | *7.9 | 50.7 | *4.8 | 0.8 | | | | | |
| Heart disease, excluding ischemic | 391-392.0,393-398,402,404,415-416,420-429 | 14,586 | 1.3 | 100.0 | 35.6 | * | 43.9 | 7.3 | 10.8 | | | | | |
| Chronic sinusitis | 473 | 14,251 | 1.3 | 100.0 | 74.6 | 10.9 | * | 7.9 | 4.1 | | | | | |
| Follow up examination | V67 | 13,471 | 1.2 | 100.0 | 36.7 | 43.6 | *10.2 | 8.0 | 1.4 | | | | | |
| Psychoses, excluding major depressive disorder | 290-295,296.0-296.1,296.4-299 | 11,653 | 1.1 | 100.0 | * | - | 67.5 | 11.7 | 7.4 | | | | | |
| Potential health hazards related personal and family history | V10-V19 | 11,528 | 1.0 | 100.0 | 40.8 | 30.1 | 18.9 | 7.7 | 2.5 | | | | | |
| Disorders of lipid metabolism | 272 | 11,442 | 1.0 | 100.0 | 86.4 | * | *6.5 | 5.9 | * | | | | | |
| Abdominal pain | 789.0 | 11,318 | 1.0 | 100.0 | 38.9 | 6.5 | * | 5.4 | 39.0 | | | | | |
| Chronic and unspecified bronchitis | 490-491 | 11,213 | 1.0 | 100.0 | 73.5 | * | * | 5.9 | 11.7 | | | | | |
| Ischemic heart disease | 410-414.9 | 10,807 | 1.0 | 100.0 | 24.0 | * | 57.3 | 6.4 | 6.6 | | | | | |
| Acute pharyngitis | 462 | 10,503 | 0.9 | 100.0 | 69.8 | * | * | 10.0 | 15.1 | | | | | |
| Benign neoplasms | 210-229,235-239 | 10,413 | 0.9 | 100.0 | 24.7 | 25.6 | 42.7 | 6.1 | * | | | | | |
| Artificial opening and other postsurgical status | V44-V45 | 10,183 | 0.9 | 100.0 | 16.7 | 65.2 | 7.6 | 9.4 | 1.2 | | | | | |
| Sprains and strains, excluding ankle and back | 840-844,845.1,848 | 9,949 | 0.9 | 100.0 | 30.0 | 36.1 | * | 5.8 | 24.1 | | | | | |
| Depressive disorder, not elsewhere classified | 311 | 9,474 | 0.9 | 100.0 | 55.1 | * | 31.8 | 7.9 | 5.2 | | | | | |
| Contact dermatitis and other eczema | 692 | 9,187 | 0.8 | 100.0 | 54.8 | * | 29.3 | 6.8 | 7.9 | | | | | |
| Open wound, excluding head | 874-897 | 9,133 | 0.8 | 100.0 | 26.5 | 16.2 | * | 6.0 | 46.8 | | | | | |
| Fractures, excluding lower limb | 800-819 | 8,908 | 0.8 | 100.0 | * | 44.2 | * | 7.3 | 29.4 | | | | | |
| Major depressive disorder | 296.2-296.3 | 8,834 | 0.8 | 100.0 | * | - | 85.9 | 9.6 | *1.4 | | | | | |
| Glaucoma | 365 | 8,772 | 0.8 | 100.0 | - | 96.2 | - | *3.7 | * | | | | | |
| Chest pain | 786.5 | 8,646 | 0.8 | 100.0 | 28.7 | * | 18.8 | 4.1 | 48.3 | | | | | |
| All other diagnoses | | 518,805 | 46.9 | 100.0 | 41.3 | 18.7 | 19.5 | 7.6 | 13.0 | | | | | |

| | | | | | Standard error of percent | | | | |
|--|---|--------|-----|-----|---------------------------|-----|------|-----|-----|
| All visits | | 35,927 | | ... | 1.5 | 1.0 | 1.3 | 0.7 | 0.5 |
| Essential hypertension | 401 | 3,727 | 0.3 | ... | 2.6 | ... | 1.6 | 1.4 | 0.4 |
| Routine infant or child health check | V20.2 | 3,155 | 0.3 | ... | 1.8 | ... | ... | 1.7 | ... |
| Acute upper respiratory infections, excluding pharyngitis | 460-461,463-466 | 2,473 | 0.2 | ... | 2.3 | 0.4 | ... | 1.3 | 1.0 |
| Malignant neoplasms | 140-208,230-234 | 5,568 | 0.5 | ... | 2.8 | 2.7 | 5.8 | 2.5 | 0.1 |
| Diabetes mellitus | 250 | 3,019 | 0.3 | ... | 5.3 | 2.2 | 5.8 | 1.8 | 0.2 |
| Spinal disorders | 720-724 | 3,210 | 0.3 | ... | 5.5 | 4.5 | 2.7 | 1.2 | 1.3 |
| Arthropathies and related disorders | 710-719 | 2,324 | 0.2 | ... | 3.9 | 4.4 | 1.4 | 1.4 | 0.6 |
| Normal pregnancy | V22 | 5,013 | 0.4 | ... | 2.7 | ... | ... | 2.6 | 0.3 |
| Rheumatism, excluding back | 725-729 | 2,009 | 0.2 | ... | 4.3 | 4.7 | 3.9 | 1.0 | 1.0 |
| Specific procedures and aftercare | V50-V59.9 | 1,849 | 0.2 | ... | 5.0 | 4.3 | 3.1 | 1.6 | 0.7 |
| Asthma | 493 | 2,275 | 0.2 | ... | 7.6 | ... | 8.7 | 1.2 | 1.6 |
| Gynecological examination | V72.3 | 1,940 | 0.2 | ... | 1.1 | ... | ... | 1.1 | ... |
| General medical examination | V70 | 1,801 | 0.2 | ... | 3.3 | ... | 2.9 | 1.2 | 0.5 |
| Otitis media and eustachian tube disorders | 381-382 | 1,350 | 0.1 | ... | 3.9 | 2.7 | ... | 1.6 | 1.4 |
| Allergic rhinitis | 477 | 3,533 | 0.3 | ... | 9.1 | 2.9 | 11.6 | 1.4 | 0.2 |
| Heart disease, excluding ischemic | 391-392.0,393-398,402,404,415-416,420-429 | 1,281 | 0.1 | ... | 4.5 | ... | 4.6 | 1.3 | 1.3 |
| Chronic sinusitis | 473 | 2,061 | 0.2 | ... | 4.4 | 2.4 | ... | 1.8 | 0.9 |
| Follow up examination | V67 | 1,952 | 0.2 | ... | 7.1 | 6.5 | 4.1 | 1.7 | 0.4 |
| Psychoses, excluding major depressive disorder | 290-295,296.0-296.1,296.4-299 | 1,364 | 0.1 | ... | ... | ... | 4.0 | 2.0 | 1.1 |
| Potential health hazards related personal and family history | V10-V19 | 1,427 | 0.1 | ... | 6.0 | 4.3 | 4.1 | 1.6 | 0.7 |
| Disorders of lipid metabolism | 272 | 1,582 | 0.1 | ... | 2.9 | ... | 2.0 | 1.5 | ... |
| Abdominal pain | 789.0 | 1,055 | 0.1 | ... | 5.8 | 1.3 | ... | 0.9 | 3.8 |
| Chronic and unspecified bronchitis | 490-491 | 1,372 | 0.1 | ... | 4.1 | ... | ... | 1.2 | 1.9 |
| Ischemic heart disease | 410-414.9 | 1,303 | 0.1 | ... | 4.6 | ... | 5.6 | 1.8 | 1.0 |
| Acute pharyngitis | 462 | 1,003 | 0.1 | ... | 4.0 | ... | ... | 2.0 | 2.2 |
| Benign neoplasms | 210-229,235-239 | 1,107 | 0.1 | ... | 3.7 | 5.4 | 5.2 | 1.3 | ... |
| Artificial opening and other postsurgical status | V44-V45 | 1,295 | 0.1 | ... | 4.5 | 5.7 | 2.2 | 2.1 | 0.3 |
| Sprains and strains, excluding ankle and back | 840-844,845.1,848 | 1,007 | 0.1 | ... | 4.2 | 4.8 | ... | 1.1 | 2.6 |
| Depressive disorder, not elsewhere classified | 311 | 1,152 | 0.1 | ... | 5.5 | ... | 5.6 | 1.5 | 0.9 |
| Contact dermatitis and other eczema | 692 | 838 | 0.1 | ... | 4.2 | ... | 3.8 | 1.2 | 1.2 |
| Open wound, excluding head | 874-897 | 743 | 0.1 | ... | 4.6 | 3.9 | ... | 1.1 | 3.8 |
| Fractures, excluding lower limb | 800-819 | 982 | 0.1 | ... | ... | 4.6 | ... | 1.5 | 3.3 |
| Major depressive disorder | 296.2-296.3 | 1,396 | 0.1 | ... | ... | ... | 2.9 | 1.8 | 0.4 |
| Glaucoma | 365 | 1,525 | 0.1 | ... | ... | 1.2 | ... | 1.2 | ... |
| Chest pain | 786.5 | 628 | 0.1 | ... | 4.3 | ... | 3.0 | 0.7 | 3.6 |
| All other diagnoses | | 18,692 | 0.8 | ... | 1.4 | 1.2 | 1.6 | 0.7 | 0.7 |

* Figure does not meet standards of reliability or precision.

... Category not applicable.

-Quantity zero.

¹Based on the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). However, certain codes have been combined in this table to form larger categories that better describe the utilization of ambulatory care services.