

Table 32 (page 1 of 4). Death rates for malignant neoplasms, by sex, race, Hispanic origin, and age: United States, selected years 1950–2007

[Data are based on death certificates]

<i>Sex, race, Hispanic origin, and age</i>	1950 ^{1,2}	1960 ^{1,2}	1970 ²	1980 ²	1990 ²	2000 ³	2006 ³	2007 ³
All persons								
All ages, age-adjusted ⁴	193.9	193.9	198.6	207.9	216.0	199.6	180.7	178.4
All ages, crude	139.8	149.2	162.8	183.9	203.2	196.5	187.0	186.6
Under 1 year	8.7	7.2	4.7	3.2	2.3	2.4	1.8	1.7
1–4 years	11.7	10.9	7.5	4.5	3.5	2.7	2.3	2.2
5–14 years	6.7	6.8	6.0	4.3	3.1	2.5	2.2	2.4
15–24 years	8.6	8.3	8.3	6.3	4.9	4.4	3.9	3.9
25–34 years	20.0	19.5	16.5	13.7	12.6	9.8	9.0	8.5
35–44 years	62.7	59.7	59.5	48.6	43.3	36.6	31.9	30.8
45–54 years	175.1	177.0	182.5	180.0	158.9	127.5	116.3	114.3
55–64 years	390.7	396.8	423.0	436.1	449.6	366.7	321.2	315.4
65–74 years	698.8	713.9	754.2	817.9	872.3	816.3	727.2	715.5
75–84 years	1,153.3	1,127.4	1,169.2	1,232.3	1,348.5	1,335.6	1,263.8	1,256.3
85 years and over	1,451.0	1,450.0	1,320.7	1,594.6	1,752.9	1,819.4	1,606.1	1,590.2
Male								
All ages, age-adjusted ⁴	208.1	225.1	247.6	271.2	280.4	248.9	220.1	217.5
All ages, crude	142.9	162.5	182.1	205.3	221.3	207.2	196.6	197.0
Under 1 year	9.7	7.7	4.4	3.7	2.4	2.6	1.8	1.8
1–4 years	12.5	12.4	8.3	5.2	3.7	3.0	2.5	2.3
5–14 years	7.4	7.6	6.7	4.9	3.5	2.7	2.5	2.4
15–24 years	9.7	10.2	10.4	7.8	5.7	5.1	4.6	4.5
25–34 years	17.7	18.8	16.3	13.4	12.6	9.2	8.6	8.2
35–44 years	45.6	48.9	53.0	44.0	38.5	32.7	27.4	26.4
45–54 years	156.2	170.8	183.5	188.7	162.5	130.9	119.0	117.5
55–64 years	413.1	459.9	511.8	520.8	532.9	415.8	363.6	358.5
65–74 years	791.5	890.5	1,006.8	1,093.2	1,122.2	1,001.9	870.4	854.3
75–84 years	1,332.6	1,389.4	1,588.3	1,790.5	1,914.4	1,760.6	1,631.3	1,617.4
85 years and over	1,668.3	1,741.2	1,720.8	2,369.5	2,739.9	2,710.7	2,248.7	2,249.2
Female								
All ages, age-adjusted ⁴	182.3	168.7	163.2	166.7	175.7	167.6	153.6	151.3
All ages, crude	136.8	136.4	144.4	163.6	186.0	186.2	177.6	176.5
Under 1 year	7.6	6.8	5.0	2.7	2.2	2.3	1.8	1.6
1–4 years	10.8	9.3	6.7	3.7	3.2	2.5	2.1	2.2
5–14 years	6.0	6.0	5.2	3.6	2.8	2.2	2.0	2.3
15–24 years	7.6	6.5	6.2	4.8	4.1	3.6	3.1	3.2
25–34 years	22.2	20.1	16.7	14.0	12.6	10.4	9.5	8.9
35–44 years	79.3	70.0	65.6	53.1	48.1	40.4	36.4	35.2
45–54 years	194.0	183.0	181.5	171.8	155.5	124.2	113.7	111.3
55–64 years	368.2	337.7	343.2	361.7	375.2	321.3	281.8	275.2
65–74 years	612.3	560.2	557.9	607.1	677.4	663.6	605.9	597.6
75–84 years	1,000.7	924.1	891.9	903.1	1,010.3	1,058.5	1,012.5	1,007.4
85 years and over	1,299.7	1,263.9	1,096.7	1,255.7	1,372.1	1,456.4	1,305.5	1,276.7
White male ⁵								
All ages, age-adjusted ⁴	210.0	224.7	244.8	265.1	272.2	243.9	217.9	215.1
All ages, crude	147.2	166.1	185.1	208.7	227.7	218.1	208.7	208.8
25–34 years	17.7	18.8	16.2	13.6	12.3	9.2	8.6	8.1
35–44 years	44.5	46.3	50.1	41.1	35.8	30.9	26.7	25.9
45–54 years	150.8	164.1	172.0	175.4	149.9	123.5	113.6	112.0
55–64 years	409.4	450.9	498.1	497.4	508.2	401.9	352.9	346.7
65–74 years	798.7	887.3	997.0	1,070.7	1,090.7	984.3	862.0	845.4
75–84 years	1,367.6	1,413.7	1,592.7	1,779.7	1,883.2	1,736.0	1,631.3	1,617.4
85 years and over	1,732.7	1,791.4	1,772.2	2,375.6	2,715.1	2,693.7	2,258.3	2,253.2
Black or African American male ⁵								
All ages, age-adjusted ⁴	178.9	227.6	291.9	353.4	397.9	340.3	284.9	282.3
All ages, crude	106.6	136.7	171.6	205.5	221.9	188.5	172.3	172.9
25–34 years	18.0	18.4	18.8	14.1	15.7	10.1	10.0	9.5
35–44 years	55.7	72.9	81.3	73.8	64.3	48.4	36.5	34.0
45–54 years	211.7	244.7	311.2	333.0	302.6	214.2	182.2	178.0
55–64 years	490.8	579.7	689.2	812.5	859.2	626.4	542.9	544.1
65–74 years	636.5	938.5	1,168.9	1,417.2	1,613.9	1,363.8	1,156.5	1,139.5
75–84 years ⁶	853.5	1,053.3	1,624.8	2,029.6	2,478.3	2,351.8	1,979.1	1,936.9
85 years and over	---	1,155.2	1,387.0	2,393.9	3,238.3	3,264.8	2,543.3	2,637.1

See footnotes at end of table.

Table 32 (page 2 of 4). Death rates for malignant neoplasms, by sex, race, Hispanic origin, and age: United States, selected years 1950–2007

[Data are based on death certificates]

<i>Sex, race, Hispanic origin, and age</i>	1950 ^{1,2}	1960 ^{1,2}	1970 ²	1980 ²	1990 ²	2000 ³	2006 ³	2007 ³
Deaths per 100,000 resident population								
American Indian or Alaska Native male⁵								
All ages, age-adjusted ⁴	---	---	---	140.5	145.8	155.8	135.5	139.4
All ages, crude	---	---	---	58.1	61.4	67.0	76.1	83.3
25–34 years	---	---	---	*	*	*	*	0.0
35–44 years	---	---	---	*	22.8	21.4	15.1	16.0
45–54 years	---	---	---	86.9	86.9	70.3	74.5	78.3
55–64 years	---	---	---	213.4	246.2	255.6	222.8	264.5
65–74 years	---	---	---	613.0	530.6	648.0	583.5	565.5
75–84 years	---	---	---	936.4	1,038.4	1,152.5	1,016.8	984.5
85 years and over	---	---	---	1,471.2	1,654.4	1,584.2	1,161.0	1,271.2
Asian or Pacific Islander male⁵								
All ages, age-adjusted ⁴	---	---	---	165.2	172.5	150.8	126.7	130.2
All ages, crude	---	---	---	81.9	82.7	85.2	84.5	89.0
25–34 years	---	---	---	6.3	9.2	7.4	6.9	6.5
35–44 years	---	---	---	29.4	27.7	26.1	19.6	18.8
45–54 years	---	---	---	108.2	92.6	78.5	70.2	73.4
55–64 years	---	---	---	298.5	274.6	229.2	197.2	190.0
65–74 years	---	---	---	581.2	687.2	559.4	459.9	470.7
75–84 years	---	---	---	1,147.6	1,229.9	1,086.1	942.3	1,014.5
85 years and over	---	---	---	1,798.7	1,837.0	1,823.2	1,439.0	1,427.4
Hispanic or Latino male^{5,7}								
All ages, age-adjusted ⁴	---	---	---	---	174.7	171.7	143.4	141.4
All ages, crude	---	---	---	---	65.5	61.3	60.4	61.6
25–34 years	---	---	---	---	8.0	6.9	6.1	6.3
35–44 years	---	---	---	---	22.5	20.1	16.0	17.4
45–54 years	---	---	---	---	96.6	79.4	71.4	74.2
55–64 years	---	---	---	---	294.0	253.1	224.8	221.9
65–74 years	---	---	---	---	655.5	651.2	574.8	560.3
75–84 years	---	---	---	---	1,233.4	1,306.4	1,098.4	1,072.6
85 years and over	---	---	---	---	2,019.4	2,049.7	1,440.1	1,417.9
White, not Hispanic or Latino male⁷								
All ages, age-adjusted ⁴	---	---	---	---	276.7	247.7	223.4	220.8
All ages, crude	---	---	---	---	246.2	244.4	239.9	240.6
25–34 years	---	---	---	---	12.8	9.7	9.3	8.6
35–44 years	---	---	---	---	36.8	32.3	28.9	27.7
45–54 years	---	---	---	---	153.9	127.2	118.7	116.8
55–64 years	---	---	---	---	520.6	412.0	363.4	357.6
65–74 years	---	---	---	---	1,109.0	1,002.1	883.0	867.3
75–84 years	---	---	---	---	1,906.6	1,750.2	1,662.9	1,652.8
85 years and over	---	---	---	---	2,744.4	2,714.1	2,300.2	2,300.4
White female⁵								
All ages, age-adjusted ⁴	182.0	167.7	162.5	165.2	174.0	166.9	153.6	151.2
All ages, crude	139.9	139.8	149.4	170.3	196.1	199.4	190.1	188.8
25–34 years	20.9	18.8	16.3	13.5	11.9	10.1	9.1	8.6
35–44 years	74.5	66.6	62.4	50.9	46.2	38.2	34.9	33.9
45–54 years	185.8	175.7	177.3	166.4	150.9	120.1	109.5	107.1
55–64 years	362.5	329.0	338.6	355.5	368.5	319.7	279.1	271.8
65–74 years	616.5	562.1	554.7	605.2	675.1	665.6	611.5	602.3
75–84 years	1,026.6	939.3	903.5	905.4	1,011.8	1,063.4	1,023.0	1,017.7
85 years and over	1,348.3	1,304.9	1,126.6	1,266.8	1,372.3	1,459.1	1,317.5	1,291.6

See footnotes at end of table.

Table 32 (page 3 of 4). Death rates for malignant neoplasms, by sex, race, Hispanic origin, and age: United States, selected years 1950–2007

[Data are based on death certificates]

Sex, race, Hispanic origin, and age	1950 ^{1,2}	1960 ^{1,2}	1970 ²	1980 ²	1990 ²	2000 ³	2006 ³	2007 ³
Black or African American female ⁵								
Deaths per 100,000 resident population								
All ages, age-adjusted ⁴	174.1	174.3	173.4	189.5	205.9	193.8	176.1	174.9
All ages, crude	111.8	113.8	117.3	136.5	156.1	151.8	147.7	148.2
25–34 years	34.3	31.0	20.9	18.3	18.7	13.5	12.5	12.0
35–44 years	119.8	102.4	94.6	73.5	67.4	58.9	50.8	48.5
45–54 years	277.0	254.8	228.6	230.2	209.9	173.9	158.7	156.1
55–64 years	484.6	442.7	404.8	450.4	482.4	391.0	356.9	352.5
65–74 years	477.3	541.6	615.8	662.4	773.2	753.1	672.9	681.0
75–84 years ⁶	605.3	696.3	763.3	923.9	1,059.9	1,124.0	1,065.3	1,071.7
85 years and over	---	728.9	791.5	1,159.9	1,431.3	1,527.7	1,324.4	1,265.2
American Indian or Alaska Native female ⁵								
All ages, age-adjusted ⁴	---	---	---	94.0	106.9	108.3	108.3	102.1
All ages, crude	---	---	---	50.4	62.1	61.3	76.8	75.0
25–34 years	---	---	---	*	*	*	*	0.0
35–44 years	---	---	---	36.9	31.0	23.7	25.4	20.3
45–54 years	---	---	---	96.9	104.5	59.7	72.8	75.6
55–64 years	---	---	---	198.4	213.3	200.9	193.8	190.3
65–74 years	---	---	---	350.8	438.9	458.3	469.8	444.3
75–84 years	---	---	---	446.4	554.3	714.0	756.8	712.1
85 years and over	---	---	---	786.5	843.7	983.2	684.8	639.5
Asian or Pacific Islander female ⁵								
All ages, age-adjusted ⁴	---	---	---	93.0	103.0	100.7	92.2	90.0
All ages, crude	---	---	---	54.1	60.5	72.1	77.8	78.2
25–34 years	---	---	---	9.5	7.3	8.1	7.3	6.0
35–44 years	---	---	---	38.7	29.8	28.9	24.7	24.0
45–54 years	---	---	---	99.8	93.9	78.2	73.5	70.0
55–64 years	---	---	---	174.7	196.2	176.5	160.2	162.2
65–74 years	---	---	---	301.9	346.2	357.4	330.9	308.8
75–84 years	---	---	---	522.1	641.4	650.1	602.4	601.2
85 years and over	---	---	---	800.0	971.7	988.5	878.4	875.2
Hispanic or Latina female ^{5,7}								
All ages, age-adjusted ⁴	---	---	---	---	111.9	110.8	100.4	98.6
All ages, crude	---	---	---	---	60.7	58.5	59.7	59.9
25–34 years	---	---	---	---	9.7	7.8	8.5	8.9
35–44 years	---	---	---	---	34.8	30.7	27.9	26.5
45–54 years	---	---	---	---	100.5	84.7	78.1	75.5
55–64 years	---	---	---	---	205.4	192.5	174.4	175.6
65–74 years	---	---	---	---	404.8	410.0	370.2	364.4
75–84 years	---	---	---	---	663.0	716.5	665.9	665.7
85 years and over	---	---	---	---	1,022.7	1,056.5	884.9	814.2

See footnotes at end of table.

Table 32 (page 4 of 4). Death rates for malignant neoplasms, by sex, race, Hispanic origin, and age: United States, selected years 1950–2007

[Data are based on death certificates]

Sex, race, Hispanic origin, and age	1950 ^{1,2}	1960 ^{1,2}	1970 ²	1980 ²	1990 ²	2000 ³	2006 ³	2007 ³
White, not Hispanic or Latina female ⁷								
	Deaths per 100,000 resident population							
All ages, age-adjusted ⁴	---	---	---	---	177.5	170.0	157.6	155.3
All ages, crude	---	---	---	---	210.6	220.6	214.7	213.7
25–34 years	---	---	---	---	11.9	10.5	9.2	8.4
35–44 years	---	---	---	---	47.0	38.9	35.9	35.2
45–54 years	---	---	---	---	154.9	123.0	113.0	110.9
55–64 years	---	---	---	---	379.5	328.9	288.5	280.6
65–74 years	---	---	---	---	688.5	681.0	631.3	622.2
75–84 years	---	---	---	---	1,027.2	1,075.3	1,044.4	1,040.1
85 years and over	---	---	---	---	1,385.7	1,468.7	1,336.7	1,315.2

--- Data not available.

* Rates based on fewer than 20 deaths are considered unreliable and are not shown.

0.0 Quantity more than zero but less than 0.05.

¹Includes deaths of persons who were not residents of the 50 states and the District of Columbia (D.C.).

²Underlying cause of death was coded according to the 6th Revision of the *International Classification of Diseases* (ICD) in 1950, 7th Revision in 1960, 8th Revision in 1970, and 9th Revision in 1980–1998. See Appendix II, Cause of death; Tables IV and V.

³Starting with 1999 data, cause of death is coded according to ICD–10. See Appendix II, Cause of death, Table V; Comparability ratio, Table VI.

⁴Age-adjusted rates are calculated using the year 2000 standard population. Prior to 2003, age-adjusted rates were calculated using standard million proportions based on rounded population numbers. Starting with 2003 data, unrounded population numbers are used to calculate age-adjusted rates. See Appendix II, Age adjustment.

⁵The race groups, white, black, Asian or Pacific Islander, and American Indian or Alaska Native, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Death rates for the American Indian or Alaska Native and Asian or Pacific Islander populations are known to be underestimated. See Appendix II, Race, for a discussion of sources of bias in death rates by race and Hispanic origin.

⁶In 1950, rate is for the age group 75 years and over.

⁷Prior to 1997, excludes data from states lacking an Hispanic-origin item on the death certificate. See Appendix II, Hispanic origin.

NOTES: Starting with *Health, United States, 2003*, rates for 1991–1999 were revised using intercensal population estimates based on the 2000 census. Rates for 2000 were revised based on 2000 census counts. Rates for 2001 and later years were computed using 2000-based postcensal estimates. See Appendix I, Population Census and Population Estimates. See Appendix II, Cause of death; Tables IV and V. Age groups were selected to minimize the presentation of unstable age-specific death rates based on small numbers of deaths and for consistency among comparison groups. Starting with 2003 data, some states allowed the reporting of more than one race on the death certificate. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states. See Appendix II, Race. Data for additional years are available. See Appendix III.

SOURCE: CDC/NCHS, National Vital Statistics System; Grove RD, Hetzel AM. Vital statistics rates in the United States, 1940–1960. Washington, DC: U.S. Government Printing Office. 1968; numerator data from National Vital Statistics System, annual mortality files; denominator data from national population estimates for race groups from Table 1 and unpublished Hispanic population estimates for 1985–1996 prepared by the Housing and Household Economic Statistics Division, U.S. Census Bureau; additional mortality tables are available from: http://www.cdc.gov/nchs/nvss/mortality_tables.htm; Xu JQ, Kochanek KD, Murphy SL, Tejada-Vera B. Deaths: Final data for 2007. National vital statistics reports; vol 58 no 19. Hyattsville, MD: NCHS; 2010. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr58/nvsr58_19.pdf.