# Table 46 (page 1 of 5). Health conditions among children under 18 years of age, by selected characteristics: United States, average annual, selected years 1997–1999 through 2007–2009

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic		Current	asthma¹		Asthma attack in the past 12 months <sup>2</sup>				
	1997–1999	2000–2002	2003–2005	2007–2009	1997–1999	2000–2002	2003–2005	2007–2009	
	Percent of children								
Under 18 years <sup>3</sup>			8.7	9.4	5.4	5.7	5.4	5.4	
Age									
0–4 years			6.1 9.6	6.4 10.6	4.3 5.7	4.7 6.1	4.2 5.8	4.3 5.9	
5–9 years			9.1 9.9	10.2 10.8	5.6 5.8	6.3 5.9	6.1 5.7	6.2 5.7	
Sex									
Male			9.9 7.3	10.8 7.9	6.2 4.5	6.6 4.7	6.3 4.4	6.2 4.6	
Race <sup>4</sup>									
White only			7.7 13.0 12.2	8.0 16.0 *10.8	5.0 7.0 6.4	5.2 8.0 *8.7	4.9 7.6 *6.1	4.7 8.6 *	
Asian only			4.8	6.3	4.3	4.7	3.3	4.1	
Islander only			13.5	13.9		7.3	8.8	9.3	
Hispanic origin and race <sup>4</sup>									
Hispanic or Latino.  Not Hispanic or Latino.  White only.  Black or African American only	  		7.6 8.9 7.9 13.0	7.9 9.8 8.2 16.0	4.8 5.5 5.1 7.0	4.2 6.0 5.5 7.9	4.6 5.6 5.0 7.5	4.6 5.7 4.8 8.5	
Percent of poverty level <sup>5</sup>									
Below 100%. 100%-199%. 200%-399%. 400% or more			10.4 8.6 8.3 7.9	12.2 9.8 8.2 8.4	6.1 5.3 5.0 5.2	7.1 5.4 5.3 5.5	6.5 5.2 5.2 4.9	6.9 5.7 4.7 5.0	
Health insurance status at the time of interview <sup>6</sup>									
Insured Private Medicaid Uninsured			9.0 8.0 11.4 5.6	9.7 8.4 12.1 6.3	5.6 5.0 7.7 3.9	5.9 5.3 7.7 4.3	5.6 5.0 7.1 3.3	5.6 5.1 6.7 3.5	

See footnotes at end of table.

Health, United States, 2010 Trend Tables 199

# Table 46 (page 2 of 5). Health conditions among children under 18 years of age, by selected characteristics: United States, average annual, selected years 1997–1999 through 2007–2009

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Attent	ion deficit hyp	peractivity dis	order <sup>7</sup>	Serious emotional or behavioral difficulties8				
	1997–1999	2000–2002	2003–2005	2007–2009	1997–1999	2000–2002	2003–2005	2007–2009	
Age	Percent of children								
5–17 years5–9 years	6.5 4.8 7.6	7.5 5.2 9.0	7.6 5.6 8.9	9.0 5.8 10.9			5.1 4.3 5.6	5.5 4.7 6.0	
Sex									
Male	9.6 3.2	10.8 4.2	10.7 4.4	12.3 5.5			6.1 4.1	6.9 4.0	
Race <sup>4</sup>									
White only . Black or African American only	7.1 5.0 *8.5 *1.7	8.1 7.0 *	7.8 7.7 *9.4 *1.6	9.2 9.3 * *1.8			5.1 5.3 *1.7	5.2 6.6 *	
Native Hawaiian or Other Pacific Islander only. 2 or more races		* 7.4	* 9.7	* 13.1			* 8.2	*3.3 10.3	
Hispanic origin and race <sup>4</sup>									
Hispanic or Latino	3.6 7.0 7.7 5.0	4.2 8.2 9.0 6.8	4.6 8.3 8.8 7.5	4.9 10.0 10.6 9.5			3.8 5.4 5.6 5.2	3.6 5.9 5.8 6.5	
Percent of poverty level <sup>5</sup>									
Below 100%. 100%–199%. 200%–399%. 400% or more	7.2 6.7 6.2 6.1	8.2 7.5 7.7 7.1	8.4 7.8 7.8 6.9	10.3 10.6 8.1 7.8			7.4 5.4 4.9 3.7	8.7 6.9 4.6 3.3	
Health insurance status at the time of interview <sup>6</sup>									
Insured	6.7 5.9 10.5 4.8	7.8 7.0 10.7 5.4	7.8 7.0 10.3 6.1	9.3 7.6 12.7 6.1			5.2 4.1 8.5 4.6	5.6 3.9 9.1 4.2	

See footnotes at end of table.

200 **Trend Tables** Health, United States, 2010

# Table 46 (page 3 of 5). Health conditions among children under 18 years of age, by selected characteristics: United States, average annual, selected years 1997–1999 through 2007–2009

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic		Food a	allergy <sup>9</sup>		Skin allergy <sup>10</sup>				
	1997–1999	2000–2002	2003–2005	2007–2009	1997–1999	2000–2002	2003–2005	2007–2009	
	Percent of children								
Under 18 years	3.4	3.6	3.8	4.6	7.4	8.1	9.6	10.7	
Age									
0–4 years. 5–17 years. 5–9 years. 10–17 years	3.8 3.3 3.1 3.4	4.0 3.4 3.6 3.3	4.3 3.6 3.5 3.6	5.0 4.4 4.6 4.3	8.1 7.2 7.5 7.1	8.7 7.9 8.6 7.5	11.0 9.1 10.0 8.6	12.4 10.1 11.4 9.3	
Sex									
Male	3.4 3.5	3.7 3.4	3.8 3.8	4.6 4.5	7.3 7.6	7.9 8.4	9.5 9.8	10.6 10.8	
Race <sup>4</sup>									
White only	3.5 3.1 *	3.6 3.0 *4.8 4.4	3.8 3.7 * 4.3	4.4 4.4 *7.9 4.3	7.1 9.0 *4.1 8.0	7.6 10.4 *9.1 8.4	9.0 12.4 11.3 7.5	9.7 14.5 *9.8 10.6	
Native Hawaiian or Other Pacific Islander only. 2 or more races		* 5.2	* 4.6	* 5.7		* 10.9	* 14.0	* 15.7	
Hispanic origin and race <sup>4</sup>									
Hispanic or Latino.  Not Hispanic or Latino.  White only.  Black or African American only.	2.1 3.7 3.8 3.1	2.5 3.8 3.9 3.1	2.8 4.0 4.1 3.7	3.7 4.8 4.7 4.5	5.5 7.8 7.5 9.0	5.6 8.7 8.2 10.4	7.2 10.2 9.7 12.4	8.6 11.3 10.2 14.6	
Percent of poverty level <sup>5</sup>									
Below 100% 100%—199% 200%—399% 400% or more	3.3 3.0 3.2 4.2	3.2 3.4 3.4 4.0	3.3 3.8 3.8 4.1	3.9 4.4 4.8 5.0	7.3 7.2 7.3 7.9	7.1 7.6 8.5 8.8	9.0 8.7 10.0 10.5	11.4 10.1 10.2 11.4	
Health insurance status at the time of interview <sup>6</sup>									
Insured Private Medicaid. Uninsured	3.5 3.5 3.6 2.6	3.7 3.7 3.7 2.4	3.9 4.0 3.6 3.0	4.6 4.8 4.2 3.9	7.7 7.4 8.4 5.9	8.5 8.5 8.4 5.3	10.0 10.1 9.5 6.8	11.0 10.7 11.0 7.9	

See footnotes at end of table.

Health, United States, 2010 Trend Tables 201

# Table 46 (page 4 of 5). Health conditions among children under 18 years of age, by selected characteristics: United States, average annual, selected years 1997–1999 through 2007–2009

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Наз	y fever or res	piratory allerg	<i>gy</i> <sup>11</sup>	Three or more ear infections 12				
	1997–1999	2000–2002	2003–2005	2007–2009	1997–1999	2000–2002	2003–2005	2007–2009	
	Percent of children								
Under 18 years	17.5	17.7	17.3	16.6	7.1	6.7	5.8	5.5	
Age									
0–4 years. 5–17 years. 5–9 years . 10–17 years	10.7 19.9 17.3 21.6	10.4 20.3 18.1 21.7	10.1 20.0 17.9 21.2	10.1 19.2 17.3 20.4	13.7 4.8 7.1 3.2	12.8 4.5 6.9 2.9	11.0 3.8 5.7 2.7	10.5 3.5 5.6 2.2	
Sex									
Male	18.6 16.3	18.8 16.5	18.9 15.6	18.1 15.1	7.3 6.9	6.9 6.5	5.9 5.6	5.6 5.4	
Race <sup>4</sup>									
White only Black or African American only American Indian or Alaska Native only Asian only Native Hawaiian or Other Pacific	17.9 16.2 15.2 15.3	18.5 15.6 16.4 12.6	17.8 15.2 16.5 11.3	17.2 14.2 *13.0 13.2	7.4 5.9 *10.8 3.7	7.2 5.0 *6.3 2.6	6.3 4.1 *5.1 3.3	5.9 3.8 *7.6 2.7	
Islander only		20.9	20.8	20.5		7.4	5.0	6.1	
Hispanic origin and race <sup>4</sup>									
Hispanic or Latino	12.4 18.4 19.1 16.3	12.4 18.8 19.9 15.5	12.8 18.3 19.4 15.1	11.8 18.0 19.1 14.2	6.1 7.3 7.7 5.9	6.7 6.7 7.3 4.9	6.2 5.7 6.3 4.0	6.3 5.3 5.8 3.8	
Percent of poverty level <sup>5</sup>									
Below 100%. 100%–199%. 200%–399%. 400% or more	14.3 15.4 18.5 20.3	14.0 15.6 18.1 21.1	14.2 16.0 17.7 19.7	13.7 14.6 16.8 20.3	8.3 7.1 6.8 6.6	7.9 6.8 6.5 6.1	6.7 5.7 5.6 5.5	7.1 5.9 4.9 4.7	
Health insurance status at the time of interview <sup>6</sup>									
Insured	18.0 18.8 15.0 14.3	18.3 19.2 16.0 12.6	17.7 18.5 16.1 13.5	16.9 18.6 13.9 13.9	7.3 6.6 10.2 5.9	6.9 6.4 8.7 4.9	5.8 5.2 7.4 5.4	5.7 4.7 7.4 3.8	

See footnotes at end of table.

Trend Tables Health, United States, 2010

## Table 46 (page 5 of 5). Health conditions among children under 18 years of age, by selected characteristics: United States, average annual, selected years 1997–1999 through 2007–2009

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

<sup>3</sup>Includes all other races not shown separately and unknown health insurance status.

<sup>4</sup>The race groups white, black, American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, and 2 or more races include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity and are not strictly comparable with estimates for earlier years. The five single-race categories plus multiple-race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group; the category 2 or more races includes persons who reported more than one racial group. Prior to 1999, data were tabulated according to the 1977 Standards with four racial groups, and the Asian only category included Native Hawaiian or Other Pacific Islander. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See Appendix II, Hispanic origin; Race.

Epercent of poverty level is based on family income and family size and composition using U.S. Census Bureau poverty thresholds. Missing family income data were imputed for 1997 and beyond. See Appendix II, Family income; Poverty; Table VII.

<sup>6</sup>Health insurance categories are mutually exclusive. Persons who reported both Medicaid and private coverage are classified as having private coverage. Starting with 1997 data, state-sponsored health plan coverage is included as Medicaid coverage. Starting with 1999 data, coverage by the Children's Health Insurance Program (CHIP) is included as Medicaid coverage. In addition to private and Medicaid, the insured category also includes military, other government, and Medicare coverage. Persons not covered by private insurance, Medicaid, CHIP, state-sponsored or other government-sponsored health plans, Medicare, or military plans are considered to have no health insurance coverage. Persons with only Indian Health Service coverage are considered to have no health insurance coverage. See Appendix II, Health insurance coverage.

Based on parents responding "yes" to the question: "Has a doctor or health professional ever told you that your child had attention deficit hyperactivity disorder (ADHD) or attention deficit disorder (ADD)?'

Based on parents responding "yes, definite" or "yes, severe" to the question, "Overall, do you think that [child] has difficulties in any of the following areas: emotions, concentration, behavior, or being able to get along with other people?

Based on parents responding "yes" to the question, "During the past 12 months, has your child had any kind of food or digestive allergy?"

10 Based on parents responding "yes" to the question, "During the past 12 months, has your child had any eczema or any kind of skin allergy?"

11Based on parents responding "yes" to the question "During the past 12 months, has your child had hay fever?" or "yes" to the question, "During the past 12 months, has your child had any kind of respiratory allergy?"

12Based on parents responding "yes" to the question "During the past 12 months, has your child had three or more ear infections?"

NOTES: Answers to questions are supplied by the parents or knowledgable adult in the family. Standard errors are available in the spreadsheet version of this table. Available from: http://www.cdc.gov/nchs/hus.htm. Data for additional years are available. See Appendix III.

SOURCE: CDC/NCHS, National Health Interview Survey, sample child and family core questionnaire.

Health, United States, 2010 **Trend Tables** 203

<sup>- - -</sup> Data not available

<sup>\*</sup>Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%-30%. Data not shown have an RSE greater than 30%. Based on parents responding "yes" to both of the questions, "Has a doctor or other health professional ever told you that your child had asthma?" and "Does your child still have asthma?

<sup>&</sup>lt;sup>2</sup>Based on parents responding "yes" to both questions, "Has a doctor or other health professional ever told you that your child had asthma?" and "During the past 12 months, did your child have an episode of asthma or an asthma attack?'