

**Data table for Figure 7. Joint pain in the past 30 days among adults 18 years of age and over, by age: United States, 2002–2009**

<i>Age</i>	<i>2002</i>	<i>2003</i>	<i>2004</i>	<i>2005</i>	<i>2006</i>	<i>2007</i>	<i>2008</i>	<i>2009</i>
	Percent							
18 years and over, age-adjusted . . . . .	29.5	31.6	30.8	30.7	29.2	27.0	30.5	32.0
18 years and over, crude . . . . .	29.5	31.8	31.1	31.1	29.7	27.6	31.3	33.0
18–44 years. . . . .	19.3	20.7	19.2	19.1	18.0	16.3	20.3	20.7
45–64 years. . . . .	37.5	40.2	39.8	39.9	38.3	36.4	39.3	41.8
65–74 years. . . . .	46.0	49.3	49.3	49.3	47.6	41.6	46.8	47.9
75 years and over . . . . .	48.7	52.4	54.2	52.6	48.8	46.6	48.1	53.8
	Standard error							
18 years and over, age-adjusted . . . . .	0.3	0.3	0.3	0.3	0.4	0.4	0.4	0.4
18 years and over, crude . . . . .	0.3	0.4	0.3	0.3	0.4	0.4	0.4	0.4
18–44 years. . . . .	0.4	0.4	0.4	0.4	0.4	0.4	0.5	0.5
45–64 years. . . . .	0.6	0.6	0.6	0.6	0.7	0.7	0.7	0.7
65–74 years. . . . .	1.0	1.1	1.1	1.0	1.3	1.2	1.3	1.1
75 years and over . . . . .	1.1	1.1	1.1	1.1	1.2	1.3	1.4	1.3

NOTES: Respondents were asked, “During the past 30 days, have you had any symptoms of pain, aching, or stiffness in or around a joint?” Respondents were instructed to not include the back or neck because other questions focused on those areas. To facilitate their response, respondents were shown a card illustrating the body joints. Estimates are age-adjusted to the year 2000 standard population using five age groups: 18–44 years, 45–54 years, 55–64 years, 65–74 years, and 75 years and over. See Appendix II, Age adjustment; Table 53.

SOURCE: CDC/NCHS, National Health Interview Survey, sample adult questionnaire.