

**Data table for Figure 17. Statin drug use in the past 30 days among adults 45 years of age and over, by sex and age: United States, 1988–1994, 1999–2002, and 2005–2008**

Sex and age	1988–1994		1999–2002		2005–2008	
	Percent	SE	Percent	SE	Percent	SE
Total						
45 years and over . . . . .	2.4	0.3	15.6	0.7	25.1	0.7
45–64 years. . . . .	2.2	0.3	12.2	1.0	17.2	0.9
65–74 years. . . . .	3.7	0.7	24.9	1.6	42.0	1.5
75 years and over . . . . .	*1.4	0.4	18.4	1.2	41.4	1.3
Men						
45 years and over . . . . .	2.3	0.3	17.5	1.1	26.9	0.9
45–64 years. . . . .	2.5	0.5	15.1	1.4	18.3	1.2
65–74 years. . . . .	*	*	25.7	1.5	49.7	2.1
75 years and over . . . . .	*	*	18.6	1.7	45.0	1.4
Women						
45 years and over . . . . .	2.6	0.5	13.9	0.8	23.6	1.1
45–64 years. . . . .	1.9	0.5	9.6	1.0	16.1	1.2
65–74 years. . . . .	*4.9	1.0	24.2	2.6	36.0	2.2
75 years and over . . . . .	*1.7	0.5	18.2	1.9	38.9	2.1

SE is standard error.

\* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30%.

NOTES: Respondents reporting use of a prescription drug containing any of the following ingredients—atorvastatin, cerivastatin, fluvastatin, lovastatin, pravastatin, rosuvastatin, or simvastatin—were classified as taking a statin drug. Also see Figure 16, Table 68.

SOURCE: CDC/NCHS, National Health and Nutrition Examination Survey.

**References** (continued from page 26)

- LaRosa JC. Unresolved issues in early trials of cholesterol lowering. *Am J Cardiol* 1995;76(9):5C–9C.
- Evans M, Roberts A, Davies S, Rees A. Medical lipid-regulating therapy: Current evidence, ongoing trials, and future developments. *Drugs* 2004;64(11):1181–96.
- Carroll MD, Lacher DA, Sorlie PD, Cleeman JI, Gordon DJ, Wolz M, et al. Trends in serum lipids and lipoproteins of adults, 1960–2002. *JAMA* 2005;294(14):1773–81.