

Data table for Figure 22. Health insurance coverage among adults 18–64 years of age, by type of coverage and age: United States, 1999–2009

Type of coverage and age	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Percent											
Private											
18–44 years.	72.0	70.5	70.1	68.7	67.7	67.3	66.6	65.0	65.5	64.4	61.7
45–64 years.	79.3	78.7	78.6	77.3	77.3	77.1	76.9	75.2	75.5	74.3	72.6
Medicaid											
18–44 years.	5.7	5.6	6.3	7.1	7.4	7.7	8.3	8.6	8.7	9.2	10.3
45–64 years.	4.4	4.5	4.7	5.3	5.3	5.5	5.5	6.3	5.9	6.4	6.9
Uninsured											
18–44 years.	21.0	22.4	22.2	23.0	23.5	23.5	23.5	24.6	23.9	24.4	25.9
45–64 years.	12.2	12.6	12.2	13.1	12.5	12.8	12.9	13.2	13.5	13.6	14.6
Standard error											
Private											
18–44 years.	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.5	0.5	0.5	0.5
45–64 years.	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.5	0.5	0.5	0.5
Medicaid											
18–44 years.	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.3	0.3
45–64 years.	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2
Uninsured											
18–44 years.	0.3	0.3	0.3	0.3	0.4	0.3	0.4	0.4	0.4	0.4	0.4
45–64 years.	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.3	0.4

NOTES: Data are for the civilian noninstitutionalized population. The category Medicaid includes state-sponsored plans and Children’s Health Insurance Program (CHIP) (starting in 1999). Uninsured adults are not covered by private insurance, Medicaid, CHIP, state-sponsored or other government-sponsored health plans, Medicare, or military plans. Adults with only Indian Health Service coverage are considered to have no coverage. Percents do not add to 100 because the percentage of adults with Medicare, military plans, and other government-sponsored plans is not shown and because adults with both private insurance and Medicaid appear in both categories. Starting with data from the third quarter of 2004, persons under 65 years of age with no reported coverage were asked explicitly about Medicaid coverage. Estimates for Medicaid coverage shown in this table include the additional information. Estimates for 2000–2002 were calculated using 2000-based sample weights and may differ from estimates in other reports that used 1990-based sample weights for 2000–2002 estimates. See Appendix II, Health insurance coverage; Tables 135, 137, and 138.

SOURCE: CDC/NCHS, National Health Interview Survey.