

Data table for Figure 35. Average number of days in ICU/CCU for Medicare decedents in the last 6 months of life, by state of residence: United States, 2005

<i>State of residence</i>	<i>Mean</i>	<i>State of residence</i>	<i>Mean</i>
United States	3.47	Hawaii	2.83
New Jersey	5.71	Arkansas	2.73
Florida	5.42	Connecticut	2.72
California	4.69	Nebraska	2.72
Nevada	4.19	Mississippi	2.61
Illinois	4.09	Rhode Island	2.60
Texas	4.08	New Mexico	2.47
Delaware	3.92	Oklahoma	2.42
Pennsylvania	3.86	Massachusetts	2.30
South Carolina	3.74	Kansas	2.20
Arizona	3.51	Alaska	2.18
District of Columbia	3.51	Washington	2.18
Ohio	3.51	Colorado	2.06
Virginia	3.37	Utah	2.02
Kentucky	3.36	Montana	1.88
Missouri	3.36	Minnesota	1.84
Tennessee	3.36	South Dakota	1.81
Louisiana	3.32	Iowa	1.78
Michigan	3.28	Maine	1.77
West Virginia	3.25	Wyoming	1.77
Alabama	3.22	New Hampshire	1.76
Georgia	3.20	Wisconsin	1.69
New York	3.17	Idaho	1.53
Maryland	3.13	Oregon	1.51
Indiana	3.11	Vermont	1.41
North Carolina	3.02	North Dakota	1.34

NOTES: Includes Medicare enrollees who had full Part A and Part B entitlement throughout the last 6 months of life. Persons enrolled in managed care organizations (HMOs) were excluded. Restricted to those whose age on the date of death was 65–99 years. Geographic location is based on decedent’s residence, not place of care. Estimates were adjusted for differences in age, sex, and race.

SOURCE: The Dartmouth Atlas of Health Care. Available from: <http://www.dartmouthatlas.org/>.