

**Table 90 (page 1 of 3). Use of mammography among women 40 years of age and over, by selected characteristics: United States, selected years 1987–2010**

Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2011.htm#090>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	1987	1990	1993	1994	2000	2003	2005	2008	2010
Percent of women having a mammogram within the past 2 years <sup>1</sup>									
40 years and over, age-adjusted <sup>2,3</sup>	29.0	51.7	59.7	61.0	70.4	69.5	66.6	67.1	66.5
40 years and over, crude <sup>2</sup>	28.7	51.4	59.7	60.9	70.4	69.7	66.8	67.6	67.1
50 years and over, age-adjusted <sup>2,3</sup>	27.3	49.8	59.7	60.9	73.7	72.4	68.2	70.3	68.8
50 years and over, crude <sup>2</sup>	27.4	49.7	59.7	60.6	73.6	72.4	68.4	70.5	69.2
Age									
40–49 years	31.9	55.1	59.9	61.3	64.3	64.4	63.5	61.5	62.3
50–64 years	31.7	56.0	65.1	66.5	78.7	76.2	71.8	74.2	72.6
65 years and over	22.8	43.4	54.2	55.0	67.9	67.7	63.8	65.5	64.4
65–74 years	26.6	48.7	64.2	63.0	74.0	74.6	72.5	72.6	71.9
75 years and over	17.3	35.8	41.0	44.6	61.3	60.6	54.7	57.9	55.7
Race <sup>4</sup>									
40 years and over, crude:									
White only	29.6	52.2	60.0	60.6	71.4	70.1	67.4	67.9	67.4
Black or African American only	24.0	46.4	59.1	64.3	67.8	70.4	64.9	68.0	67.9
American Indian or Alaska Native only	*	43.2	49.8	65.8	47.4	63.1	72.8	62.7	71.2
Asian only	*	46.0	55.1	55.8	53.5	57.6	54.6	66.1	62.4
Native Hawaiian or Other Pacific Islander only	---	---	---	---	*	*	*	*	*
2 or more races	---	---	---	---	69.2	65.3	63.7	55.2	51.4
Hispanic origin and race <sup>4</sup>									
40 years and over, crude:									
Hispanic or Latina	18.3	45.2	50.9	51.9	61.2	65.0	58.8	61.2	64.2
Not Hispanic or Latina	29.4	51.8	60.3	61.5	71.1	70.1	67.5	68.3	67.4
White only	30.3	52.7	60.6	61.3	72.2	70.5	68.3	68.7	67.8
Black or African American only	23.8	46.0	59.2	64.4	67.9	70.5	65.2	68.3	67.4
Age, Hispanic origin, and race <sup>4</sup>									
40–49 years:									
Hispanic or Latina	*15.3	45.1	52.6	47.5	54.1	59.4	54.2	54.1	59.8
Not Hispanic or Latina:									
White only	34.3	57.0	61.6	62.0	67.2	65.2	65.5	64.1	62.6
Black or African American only	27.8	48.4	55.6	67.2	60.9	68.2	62.1	59.5	63.5
50–64 years:									
Hispanic or Latina	23.0	47.5	59.2	60.1	66.5	69.4	61.5	71.3	68.6
Not Hispanic or Latina:									
White only	33.6	58.1	66.2	67.5	80.6	77.2	73.5	74.1	73.5
Black or African American only	26.4	48.4	65.5	63.6	77.7	76.2	71.6	76.7	74.0
65 years and over:									
Hispanic or Latina	*	41.1	*35.7	48.0	68.3	69.5	63.8	59.0	65.2
Not Hispanic or Latina:									
White only	24.0	43.8	54.7	54.9	68.3	68.1	64.7	66.1	65.0
Black or African American only	14.1	39.7	56.3	61.0	65.5	65.4	60.5	66.4	60.9
Age and percent of poverty level <sup>5</sup>									
40 years and over, crude:									
Below 100%	14.6	30.8	41.1	44.2	54.8	55.4	48.5	51.4	51.4
100%–199%	20.9	39.1	47.5	48.6	58.1	60.8	55.3	55.8	53.8
200%–399%	29.7	53.3	63.2	65.0	68.8	69.9	67.2	64.4	66.2
400% or more	42.9	68.7	74.1	74.1	81.5	77.7	76.6	79.0	78.1
40–49 years:									
Below 100%	18.6	32.2	36.1	43.0	47.4	50.6	42.5	46.6	48.1
100%–199%	18.4	39.0	47.8	47.6	43.6	54.0	49.8	46.5	46.2
200%–399%	31.2	55.2	63.0	64.5	60.2	63.0	61.8	56.8	59.2
400% or more	44.1	68.9	69.6	69.9	75.8	71.6	73.6	72.5	73.6
50–64 years:									
Below 100%	14.6	29.9	47.3	46.2	61.7	58.3	50.4	57.5	54.7
100%–199%	24.2	39.8	47.0	49.0	68.3	64.0	58.8	58.9	57.3
200%–399%	29.7	56.2	66.1	69.6	75.1	74.1	70.7	69.8	70.7
400% or more	44.7	71.6	78.7	78.0	86.9	84.9	80.6	84.3	82.8
65 years and over:									
Below 100%	13.1	30.8	40.4	43.9	54.8	57.0	52.3	49.1	50.6
100%–199%	19.9	38.6	47.6	48.8	60.3	62.8	56.1	59.4	55.5
200%–399%	27.7	47.4	60.3	61.0	71.1	72.3	68.6	65.0	67.2
400% or more	34.7	61.2	71.3	73.0	81.9	73.0	72.6	78.3	74.5

See footnotes at end of table.

**Table 90 (page 2 of 3). Use of mammography among women 40 years of age and over, by selected characteristics: United States, selected years 1987–2010**

Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2011.htm#090>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	1987	1990	1993	1994	2000	2003	2005	2008	2010
Health insurance status at the time of interview <sup>6</sup>									
Percent of women having a mammogram within the past 2 years <sup>1</sup>									
40–64 years:									
Insured . . . . .	---	---	66.2	68.3	76.0	75.1	72.5	73.4	74.1
Private . . . . .	---	---	67.1	69.4	77.1	76.3	74.5	74.2	75.6
Medicaid . . . . .	---	---	51.9	54.5	61.7	63.5	55.6	64.2	64.4
Uninsured . . . . .	---	---	36.0	34.0	40.7	41.5	38.1	39.7	36.0
Health insurance status prior to interview <sup>6</sup>									
40–64 years:									
Insured continuously all 12 months . . . . .	---	---	66.6	68.6	76.8	75.6	73.1	74.1	74.7
Uninsured for any period up to 12 months . . . . .	---	---	49.4	49.9	53.0	56.0	51.3	55.3	57.3
Uninsured more than 12 months . . . . .	---	---	28.4	26.6	34.0	37.0	32.9	34.6	30.0
Age and education <sup>7</sup>									
40 years and over, crude:									
No high school diploma or GED . . . . .	17.8	36.4	46.4	48.2	57.7	58.1	52.8	53.8	53.0
High school diploma or GED . . . . .	31.3	52.7	59.0	61.0	69.7	67.8	64.9	65.2	64.4
Some college or more . . . . .	37.7	62.8	69.5	69.7	76.2	75.1	72.7	73.4	72.1
40–49 years:									
No high school diploma or GED . . . . .	15.1	38.5	43.6	50.4	46.8	53.3	51.2	46.9	44.9
High school diploma or GED . . . . .	32.6	53.1	56.6	55.8	59.0	60.8	58.8	57.2	58.4
Some college or more . . . . .	39.2	62.3	66.1	68.7	70.6	68.1	68.3	66.3	66.5
50–64 years:									
No high school diploma or GED . . . . .	21.2	41.0	51.4	51.6	66.5	63.4	56.9	64.9	56.7
High school diploma or GED . . . . .	33.8	56.5	62.4	67.8	76.6	71.8	70.1	70.4	69.9
Some college or more . . . . .	40.5	68.0	78.5	74.7	84.2	82.7	77.0	78.5	77.0
65 years and over:									
No high school diploma or GED . . . . .	16.5	33.0	44.2	45.6	57.4	56.9	50.7	49.2	54.1
High school diploma or GED . . . . .	25.9	47.5	57.4	59.1	71.8	69.7	64.3	65.7	62.5
Some college or more . . . . .	32.3	56.7	64.8	64.3	74.1	75.1	73.0	75.6	70.9
Disability measure <sup>8</sup>									
40 years and over, crude:									
Any basic actions difficulty or complex activity limitation . . . . .	---	---	---	---	67.8	67.2	63.5	63.9	63.3
Any basic actions difficulty . . . . .	---	---	---	---	67.9	67.3	63.5	63.9	63.3
Any complex activity limitation . . . . .	---	---	---	---	64.1	62.3	59.9	60.2	58.2
No disability . . . . .	---	---	---	---	72.6	71.8	69.8	71.1	70.8

See footnotes at end of table.

## Table 90 (page 3 of 3). Use of mammography among women 40 years of age and over, by selected characteristics: United States, selected years 1987–2010

Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2011.htm#090>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

\* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30%.

- - - Data not available.

<sup>1</sup>Questions concerning use of mammography differed slightly on the National Health Interview Survey across the years for which data are shown. See [Appendix II, Mammography](#).

<sup>2</sup>Includes all other races not shown separately, unknown poverty level in 1987, unknown health insurance status, unknown education level, and unknown disability status.

<sup>3</sup>Estimates for women 40 years of age and over are age-adjusted to the year 2000 standard population using four age groups: 40–49 years, 50–64 years, 65–74 years, and 75 years and over. Estimates for women 50 years of age and over are age-adjusted using three age groups. See [Appendix II, Age adjustment](#).

<sup>4</sup>The race groups, white, black, American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, and 2 or more races, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The five single-race categories plus multiple-race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group; the category 2 or more races includes persons who reported more than one racial group. Prior to 1999, data were tabulated according to the 1977 Standards with four racial groups, and the Asian only category included Native Hawaiian or Other Pacific Islander. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See [Appendix II, Hispanic origin; Race](#).

<sup>5</sup>Percent of poverty level is based on family income and family size and composition using U.S. Census Bureau poverty thresholds. Poverty level was unknown for 11% of women 40 years of age and over in 1987. Missing family income data were imputed for 1997 and beyond. See [Appendix II, Family income; Poverty; Table VI](#).

<sup>6</sup>Health insurance categories are mutually exclusive. Persons who reported both Medicaid and private coverage are classified as having private coverage. Starting with 1997 data, state-sponsored health plan coverage is included as Medicaid coverage. Starting with 1999 data, coverage by the Children's Health Insurance Program (CHIP) is included with Medicaid coverage. In addition to private and Medicaid, the insured category also includes military plans, other government-sponsored health plans, and Medicare, not shown separately. Persons not covered by private insurance, Medicaid, CHIP, public assistance (through 1996), state-sponsored or other government-sponsored health plans (starting in 1997), Medicare, or military plans are considered to have no health insurance coverage. Persons with only Indian Health Service coverage are considered to have no health insurance coverage. See [Appendix II, Health insurance coverage](#).

<sup>7</sup>Education categories shown are for 1998 and subsequent years. GED is General Educational Development high school equivalency diploma. In years prior to 1998, the following categories based on number of years of school completed were used: less than 12 years, 12 years, 13 years or more. See [Appendix II, Education](#).

<sup>8</sup>Any basic actions difficulty or complex activity limitation is defined as having one or more of the following limitations or difficulties: movement difficulty, emotional difficulty, sensory (seeing or hearing) difficulty, cognitive difficulty, self-care (activity of daily living or instrumental activities of daily living) limitation, social limitation, or work limitation. For more information, see [Appendix II, Basic actions difficulty; Complex activity limitation](#). Starting with 2007 data, the hearing question, a component of the basic actions difficulty measure, was revised. Consequently, data prior to 2007 are not comparable with 2007 data and beyond. For more information on the impact of the revised hearing question, see [Appendix II, Hearing trouble](#).

NOTES: Standard errors are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data starting in 1997 are not strictly comparable with data for earlier years due to the 1997 questionnaire redesign. See [Appendix I, National Health Interview Survey \(NHIS\)](#). Data for additional years are available. See [Appendix III](#). Data have been revised and differ from previous editions of *Health, United States*.

SOURCE: CDC/NCHS, National Health Interview Survey. Data are from the following supplements: cancer control (1987), health promotion and disease prevention (1990–1991), and year 2000 objectives (1993–1994). Starting in 1998, data are from the family core and sample adult questionnaires. See [Appendix I, National Health Interview Survey \(NHIS\)](#).