

Data table for Figure 28. Vaccinations among adolescents 13–17 years of age, by type of vaccine and percent of poverty level: United States, 2009

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2011.htm#fig28>

Vaccination	Percent of poverty level									
	Total 13–17 years		Below 100%		100%–199%		200%–399%		400% or more	
	Percent	SE	Percent	SE	Percent	SE	Percent	SE	Percent	SE
Measles, mumps, rubella (MMR) (2 doses or more)	89.1	0.4	87.8	1.1	86.9	1.0	88.6	0.8	91.8	0.6
Hepatitis B (Hep B) (3 doses or more)	89.9	0.4	88.3	1.0	88.7	0.9	90.0	0.7	91.5	0.6
History of varicella disease or received varicella vaccine (2 doses or more)	75.7	0.6	74.4	1.6	72.6	1.4	76.0	1.0	78.1	0.8
Tetanus and diphtheria toxoids and acellular pertussis (Tdap) (1 dose or more since age 10).	55.6	0.6	52.8	1.8	49.9	1.6	57.8	1.1	58.4	1.0
Meningococcal conjugate vaccine (MenACWY) (1 dose or more)	53.6	0.6	52.5	1.8	48.6	1.6	50.2	1.1	60.9	1.0
Human papillomavirus (HPV) (3 doses or more, reported among females)	26.7	0.8	25.5	2.2	23.9	1.9	22.8	1.3	32.3	1.3

NOTES: SE is standard error. Data are for the civilian noninstitutionalized population. Varicella is chickenpox. For meningococcal vaccine, includes persons receiving MenACWY or meningococcal-unknown type vaccine. Information on the adolescent vaccination schedule is available from: <http://www.cdc.gov/vaccines/recs/schedules/downloads/child/mmwr-child-schedule.pdf>. Percent of poverty level is based on family income and family size and composition using U.S. Census Bureau poverty thresholds. Missing family income data were imputed. See Appendix II, Family income; Poverty; Vaccination. Also see Table 87.

SOURCE: CDC/NCHS, and National Center for Immunization and Respiratory Diseases, National Immunization Survey (NIS)–Teen. See Appendix I, National Immunization Survey (NIS).

References (continued from Figure 28 text)

2. CDC. State vaccination requirements; 2011. Available from: <http://www.cdc.gov/vaccines/vac-gen/laws/state-reqs.htm>.
3. CDC. Quadrivalent human papillomavirus vaccine: Recommendations of the Advisory Committee on Immunization Practices (ACIP). MMWR 2007;56(RR-2):1–24. Available from: <http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5602a1.htm>.
4. CDC. Prevention and control of meningococcal disease: Recommendations of the Advisory Committee on Immunization Practices (ACIP). MMWR 2005;54(RR-7):1–21. Available from: <http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5407a1.htm>.
5. CDC. Updated recommendations for use of meningococcal conjugate vaccines—Advisory Committee on Immunization Practices (ACIP), 2010. MMWR 2011;60(03):72–6. Available from: http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6003a3.htm?s_cid=mm6003a3_e.
6. CDC. Vaccines for Children Program (VFC); 2011. Available from: <http://www.cdc.gov/vaccines/programs/vfc/default.htm>.
7. CDC. National, state, and local area vaccination coverage among adolescents aged 13–17 years—United States, 2009. MMWR 2010;59(32):1018–23. Available from: <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5932a3.htm>.