

Table 42 (page 1 of 4). Age-adjusted cancer incidence rates for selected cancer sites, by sex, race, and Hispanic origin: United States, selected geographic areas, selected years 1990–2010

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2012.htm#042>.

[Data are based on the Surveillance, Epidemiology, and End Results (SEER) Program's 13 population-based cancer registries]

Site, sex, race, and Hispanic origin	1990	1995	2000	2002	2003	2005	2008	2009	2010	1990–2010 APC ¹
All sites										
Number of new cases per 100,000 population ²										
All persons	475.7	471.1	474.6	473.7	462.7	459.7	459.8	456.8	443.1	†–0.4
White	483.4	477.8	486.1	484.8	474.1	472.6	472.0	468.3	453.8	†–0.3
Black or African American	513.3	535.7	520.2	520.2	509.2	498.8	496.1	493.7	470.8	†–0.6
American Indian or Alaska Native ³	347.0	367.9	362.1	354.6	376.2	401.6	383.2	385.8	390.2	†0.3
Asian or Pacific Islander	334.2	337.3	337.1	342.5	330.5	326.8	325.7	320.7	313.8	†–0.4
Hispanic or Latino ⁴	357.4	360.1	362.0	373.0	360.2	366.9	359.1	355.5	341.1	†–0.2
White, not Hispanic or Latino ⁴	495.1	491.5	503.8	502.1	492.1	490.5	492.0	488.9	474.6	†–0.2
Male	584.0	564.4	564.8	557.8	545.0	533.6	528.5	520.2	502.6	†–0.9
White	591.1	563.8	570.0	563.9	550.7	542.8	535.8	526.9	508.3	†–0.8
Black or African American	686.4	737.1	700.0	683.9	663.0	627.8	631.4	614.5	578.3	†–1.3
American Indian or Alaska Native ³	393.8	420.7	371.7	379.4	439.1	423.9	422.5	429.0	430.7	–0.1
Asian or Pacific Islander	385.1	395.8	394.9	385.6	382.7	366.3	354.6	345.2	334.8	†–0.9
Hispanic or Latino ⁴	416.1	439.1	434.6	441.4	422.7	425.9	410.8	407.6	386.6	†–0.5
White, not Hispanic or Latino ⁴	606.7	577.8	589.1	581.7	569.3	560.8	556.7	547.4	529.3	†–0.7
Female	411.4	410.4	413.6	416.7	406.6	408.8	411.5	412.3	401.1	–0.1
White	421.5	423.6	430.5	431.8	422.4	424.7	427.6	427.4	415.6	0.0
Black or African American	404.7	401.0	399.2	411.1	405.9	411.7	403.4	411.9	396.2	0.0
American Indian or Alaska Native ³	315.6	334.3	361.7	335.3	335.3	386.3	361.4	360.2	367.6	†0.7
Asian or Pacific Islander	294.2	294.3	297.3	315.6	296.4	302.0	308.9	307.8	303.0	0.1
Hispanic or Latino ⁴	326.1	312.1	319.3	331.3	322.2	330.2	327.6	322.9	314.0	0.0
White, not Hispanic or Latino ⁴	430.4	436.8	446.2	448.0	439.4	441.8	446.1	447.5	435.4	0.1
Lung and bronchus										
Male	95.0	86.9	77.8	75.7	75.5	71.5	67.2	66.0	62.0	†–2.0
White	94.2	85.0	76.4	75.0	74.4	70.9	66.1	65.2	61.9	†–1.9
Black or African American	133.9	136.7	110.7	109.1	111.1	97.6	95.4	94.5	80.6	†–2.4
American Indian or Alaska Native ³	74.8	82.8	62.6	46.2	72.6	67.4	69.5	62.0	58.6	–0.8
Asian or Pacific Islander	64.2	60.0	63.3	57.6	58.2	57.7	55.3	51.9	49.3	†–1.1
Hispanic or Latino ⁴	59.3	52.2	45.4	49.0	46.6	44.1	39.9	38.5	33.8	†–1.9
White, not Hispanic or Latino ⁴	97.4	88.5	80.3	78.5	78.0	74.7	70.0	69.2	66.2	†–1.8
Female	47.2	49.3	48.6	49.4	49.8	49.9	47.7	48.0	45.2	–0.1
White	48.5	51.8	50.8	51.6	52.4	51.9	50.3	50.1	47.3	0.0
Black or African American	52.9	49.7	54.7	55.1	54.7	57.9	53.0	54.9	51.4	0.2
American Indian or Alaska Native ³	30.4	46.1	38.5	39.9	41.4	45.5	44.6	34.8	36.4	1.3
Asian or Pacific Islander	28.3	27.2	27.1	29.2	28.8	30.5	27.1	30.4	28.4	0.2
Hispanic or Latino ⁴	26.4	25.0	24.0	25.3	25.4	24.2	25.0	26.0	24.2	–0.3
White, not Hispanic or Latino ⁴	50.8	54.9	54.5	55.5	56.5	56.3	54.6	54.3	51.2	0.1
Colon and rectum										
Male	72.3	63.2	62.6	60.1	58.4	54.6	51.6	48.9	46.5	†–1.9
White	73.0	62.5	62.2	59.0	57.2	54.2	50.6	47.5	44.8	†–2.1
Black or African American	72.7	74.7	72.6	72.3	75.6	66.2	65.7	60.7	57.2	†–1.1
American Indian or Alaska Native ³	61.9	65.2	48.0	49.4	68.6	65.6	48.0	64.7	64.0	–0.3
Asian or Pacific Islander	60.8	58.2	57.3	58.3	52.7	46.9	46.2	45.9	44.7	†–1.5
Hispanic or Latino ⁴	47.3	45.7	50.4	45.9	47.3	47.2	47.3	45.7	41.6	†–0.5
White, not Hispanic or Latino ⁴	75.1	64.0	63.5	60.3	58.3	55.0	50.9	47.5	45.5	†–2.2
Female	50.2	45.9	46.1	45.1	43.5	41.4	39.7	37.9	35.7	†–1.4
White	49.8	45.5	45.6	44.1	43.0	40.3	39.0	36.6	34.2	†–1.5
Black or African American	61.1	54.7	57.9	56.0	55.2	53.8	48.4	50.1	45.3	†–0.9
American Indian or Alaska Native ³	45.8	47.9	39.1	50.3	44.5	48.3	46.2	43.9	41.0	–0.2
Asian or Pacific Islander	37.6	38.5	37.3	41.5	36.3	36.7	35.0	34.1	33.3	†–0.9
Hispanic or Latino ⁴	34.9	32.1	34.1	32.2	34.9	33.9	32.5	31.5	30.6	–0.3
White, not Hispanic or Latino ⁴	50.9	46.7	46.8	45.6	43.8	41.3	40.0	37.3	34.7	†–1.5
Prostate										
Male	166.9	166.4	178.8	177.7	165.2	153.7	152.5	148.7	140.1	†–1.5
White	168.5	161.4	174.9	174.6	161.5	149.7	148.2	143.6	135.0	†–1.6
Black or African American	218.9	276.5	288.3	278.1	251.3	238.6	235.0	231.4	210.9	†–1.5
American Indian or Alaska Native ³	99.5	92.5	69.6	93.0	110.3	92.7	81.9	91.2	81.9	†–1.5
Asian or Pacific Islander	88.3	103.8	106.6	102.4	103.5	94.2	85.9	82.3	75.6	†–1.4
Hispanic or Latino ⁴	118.6	140.2	149.6	151.5	138.7	132.3	125.2	122.0	113.1	†–1.0
White, not Hispanic or Latino ⁴	172.2	163.8	178.8	177.8	165.1	152.3	152.6	147.9	139.1	†–1.6

See footnotes at end of table.

Table 42 (page 2 of 4). Age-adjusted cancer incidence rates for selected cancer sites, by sex, race, and Hispanic origin: United States, selected geographic areas, selected years 1990–2010

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2012.htm#042>.

[Data are based on the Surveillance, Epidemiology, and End Results (SEER) Program's 13 population-based cancer registries]

Site, sex, race, and Hispanic origin	1990	1995	2000	2002	2003	2005	2008	2009	2010	1990–2010 APC ¹
Breast										
Number of new cases per 100,000 population ²										
Female	129.4	130.9	134.1	132.7	124.0	124.2	125.8	127.1	122.4	†–0.3
White	134.3	136.5	140.9	138.9	129.2	129.8	129.8	131.8	126.6	†–0.3
Black or African American	116.9	122.3	120.6	122.5	122.3	118.1	124.2	126.8	121.2	†0.2
American Indian or Alaska Native ³	69.6	94.2	99.1	80.1	93.0	104.5	88.1	97.2	90.5	0.3
Asian or Pacific Islander	87.5	86.8	93.6	99.4	90.6	93.0	100.6	97.5	97.8	†0.6
Hispanic or Latina ⁴	92.0	90.3	97.3	94.4	88.2	93.3	93.2	90.7	85.5	–0.1
White, not Hispanic or Latina ⁴	138.5	141.9	147.2	145.9	135.8	136.5	136.5	139.8	135.0	–0.2
Cervix uteri										
Female	11.9	9.9	8.9	8.4	8.2	7.9	7.6	7.4	7.1	†–2.5
White	11.2	9.2	8.9	8.3	7.9	7.8	7.5	7.4	7.1	†–2.1
Black or African American	16.5	14.8	10.6	10.0	10.7	9.2	9.2	7.9	8.3	†–3.7
American Indian or Alaska Native ³	14.7	*	*	*	*	10.7	9.1	10.6	8.3	†–2.2
Asian or Pacific Islander	12.1	11.0	7.9	8.2	8.1	7.7	6.2	6.6	6.3	†–4.0
Hispanic or Latina ⁴	21.4	17.4	17.1	14.6	14.1	13.7	11.8	10.1	10.3	†–3.7
White, not Hispanic or Latina ⁴	9.7	7.8	7.1	6.9	6.5	6.4	6.4	6.6	6.1	†–2.1
Corpus and uterus, not otherwise specified										
Female	24.7	24.9	23.9	24.0	23.6	24.1	25.2	26.5	26.6	0.2
White	26.4	26.4	25.6	24.8	25.0	25.4	26.3	27.4	27.4	0.0
Black or African American	16.9	17.7	17.2	22.1	20.2	21.4	23.8	25.3	24.4	†2.0
American Indian or Alaska Native ³	19.3	*	18.5	18.6	19.5	14.4	18.9	28.7	27.5	*
Asian or Pacific Islander	13.5	17.7	16.5	18.7	16.5	18.6	19.9	21.0	22.1	†1.7
Hispanic or Latina ⁴	18.1	16.6	16.2	17.8	18.1	19.4	19.3	21.0	20.3	†1.0
White, not Hispanic or Latina ⁴	27.0	27.5	26.8	25.8	25.9	26.2	27.3	28.3	28.3	0.0
Ovary										
Female	15.6	14.5	14.2	13.9	13.5	13.1	12.9	12.6	12.3	†–1.0
White	16.4	15.4	15.1	14.7	14.3	13.9	13.7	13.4	13.2	†–1.0
Black or African American	11.3	10.8	10.7	9.8	11.5	10.6	9.9	10.0	8.9	–0.5
American Indian or Alaska Native ³	21.9	*	18.9	*	13.4	12.2	11.1	16.1	11.8	*
Asian or Pacific Islander	11.2	10.4	10.1	12.0	10.1	10.7	9.9	9.4	9.4	–0.5
Hispanic or Latina ⁴	12.3	11.7	10.9	14.0	12.0	11.9	12.3	10.3	11.7	–0.4
White, not Hispanic or Latina ⁴	16.7	15.9	15.6	14.7	14.6	14.1	13.9	13.8	13.2	†–1.0
Oral cavity and pharynx										
Male	18.5	16.5	15.8	15.7	15.2	15.1	15.8	15.8	15.3	†–0.9
White	18.0	16.3	15.7	15.9	15.3	15.5	16.3	16.4	15.8	†–0.6
Black or African American	25.4	22.3	19.3	17.9	17.4	15.7	14.3	14.8	13.3	†–2.9
American Indian or Alaska Native ³	*	*	*	*	15.7	10.4	*	10.0	21.1	*
Asian or Pacific Islander	14.9	11.8	13.2	12.8	11.6	11.3	12.6	11.5	11.4	†–1.1
Hispanic or Latino ⁴	10.7	12.3	9.0	9.5	8.9	9.7	10.3	11.2	8.6	†–0.9
White, not Hispanic or Latino ⁴	18.8	16.9	16.7	16.9	16.3	16.4	17.5	17.5	17.2	†–0.4
Female	7.3	7.0	6.2	6.5	5.9	6.1	6.3	6.1	6.0	†–1.0
White	7.4	7.1	6.2	6.5	5.9	6.0	6.4	6.1	6.2	†–1.0
Black or African American	6.4	6.6	5.4	6.3	6.7	6.9	5.0	6.1	5.2	†–1.2
American Indian or Alaska Native ³	*	*	*	*	*	*	*	*	*	*
Asian or Pacific Islander	6.1	5.2	6.1	6.0	5.1	5.8	5.7	4.7	5.1	†–1.0
Hispanic or Latina ⁴	4.1	3.7	3.7	3.8	4.0	3.5	4.5	4.2	4.0	–0.4
White, not Hispanic or Latina ⁴	7.8	7.6	6.6	7.1	6.2	6.5	6.7	6.6	6.6	†–0.8
Stomach										
Male	14.6	13.5	12.6	12.0	11.7	11.4	10.6	11.3	10.3	†–1.7
White	12.8	11.9	10.7	10.4	10.1	9.6	9.3	9.9	9.2	†–1.7
Black or African American	21.4	18.6	18.4	15.8	18.5	17.4	16.6	15.3	13.0	†–2.2
American Indian or Alaska Native ³	*	24.1	20.2	25.3	*	20.8	12.7	16.6	19.2	–0.5
Asian or Pacific Islander	26.8	24.5	22.8	20.4	19.1	20.1	15.6	17.2	14.8	†–2.8
Hispanic or Latino ⁴	20.2	19.3	16.1	16.4	16.2	15.5	16.0	16.1	14.7	†–1.5
White, not Hispanic or Latino ⁴	12.1	11.1	10.0	9.6	9.2	8.7	8.2	8.8	8.2	†–2.0

See footnotes at end of table.

Table 42 (page 3 of 4). Age-adjusted cancer incidence rates for selected cancer sites, by sex, race, and Hispanic origin: United States, selected geographic areas, selected years 1990–2010

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2012.htm#042>.

[Data are based on the Surveillance, Epidemiology, and End Results (SEER) Program's 13 population-based cancer registries]

Site, sex, race, and Hispanic origin	1990	1995	2000	2002	2003	2005	2008	2009	2010	1990–2010 APC ¹
Stomach										
Number of new cases per 100,000 population ²										
Female	6.7	6.2	6.1	6.2	6.0	5.7	5.6	5.7	5.6	†–0.9
White	5.7	5.1	5.0	5.1	5.0	4.7	4.6	4.5	4.7	†–1.0
Black or African American	9.9	9.8	8.6	9.9	9.5	8.1	8.1	8.8	8.3	†–1.1
American Indian or Alaska Native ³	*	*	*	17.2	*	*	*	9.9	13.0	*
Asian or Pacific Islander	15.4	13.0	13.0	11.3	11.1	10.3	9.8	10.7	8.7	†–2.6
Hispanic or Latina ⁴	10.8	11.3	10.9	10.8	10.4	10.4	8.6	8.3	9.2	†–1.0
White, not Hispanic or Latina ⁴	5.1	4.4	4.2	4.2	4.1	3.7	3.7	3.7	3.6	†–1.7
Pancreas										
Male	13.0	12.7	12.8	12.8	12.5	13.6	14.0	14.0	13.5	†0.5
White	12.7	12.4	12.6	13.0	12.4	13.4	13.8	14.1	13.3	†0.6
Black or African American	19.3	19.1	18.1	13.7	17.2	18.1	18.5	18.8	18.5	–0.3
American Indian or Alaska Native ³	*	*	*	*	*	21.2	16.0	13.7	17.2	*
Asian or Pacific Islander	11.0	10.3	10.7	9.8	10.1	11.7	11.6	10.4	11.1	–0.1
Hispanic or Latino ⁴	10.7	12.0	12.2	10.9	10.0	12.2	11.6	13.4	11.3	†0.9
White, not Hispanic or Latino ⁴	12.8	12.4	12.7	13.3	12.7	13.5	14.1	14.2	13.5	†0.6
Female	10.0	9.9	9.9	10.5	10.3	10.8	10.9	11.2	11.1	†0.6
White	9.7	9.6	9.6	10.1	10.2	10.6	10.6	11.0	10.7	†0.6
Black or African American	12.9	15.5	12.7	15.8	14.4	16.4	15.2	14.5	14.9	–0.1
American Indian or Alaska Native ³	*	*	20.4	*	*	12.4	12.6	11.4	12.0	*
Asian or Pacific Islander	9.9	8.1	9.2	8.9	8.1	8.0	9.1	9.5	10.2	†0.9
Hispanic or Latina ⁴	9.9	8.9	9.3	11.0	9.0	11.6	9.9	10.1	10.1	0.3
White, not Hispanic or Latina ⁴	9.7	9.7	9.6	10.1	10.4	10.5	10.8	11.3	10.8	†0.7
Urinary bladder										
Male	37.2	35.4	36.8	35.8	36.9	37.0	35.5	34.9	35.5	–0.1
White	40.7	38.9	40.8	39.4	40.8	40.9	39.1	38.3	39.5	–0.1
Black or African American	19.5	19.3	20.2	20.5	22.8	22.8	22.6	21.9	21.8	†0.5
American Indian or Alaska Native ³	*	*	*	*	*	16.8	21.7	20.2	13.0	*
Asian or Pacific Islander	15.4	16.6	16.6	19.4	17.8	17.0	17.8	17.2	16.9	†0.8
Hispanic or Latino ⁴	22.1	17.8	20.4	21.1	20.1	20.0	17.3	19.0	18.0	–0.3
White, not Hispanic or Latino ⁴	42.4	41.0	43.2	41.7	43.4	43.6	42.2	41.2	42.7	0.1
Female	9.5	9.3	9.1	9.1	9.2	9.0	8.8	8.4	8.5	†–0.5
White	10.0	10.1	10.0	10.1	10.0	9.7	9.7	9.1	9.3	†–0.3
Black or African American	8.6	7.2	7.7	8.5	7.7	7.8	6.4	6.8	6.9	–0.4
American Indian or Alaska Native ³	*	*	*	*	*	*	*	*	*	*
Asian or Pacific Islander	5.3	4.4	4.1	3.2	4.9	5.1	4.9	3.8	4.6	–0.2
Hispanic or Latina ⁴	6.0	5.3	5.7	6.5	4.6	6.3	5.6	4.9	4.4	–0.6
White, not Hispanic or Latina ⁴	10.3	10.6	10.5	10.6	10.8	10.2	10.4	9.9	10.2	–0.1
Non-Hodgkin lymphoma										
Male	22.6	25.1	23.5	23.8	24.2	24.6	24.6	24.7	25.0	†0.3
White	23.6	26.2	25.0	25.1	25.7	25.9	26.0	26.1	26.1	†0.3
Black or African American	17.4	21.5	17.5	18.0	19.2	19.5	18.3	18.9	21.0	0.2
American Indian or Alaska Native ³	*	*	15.3	16.6	*	23.2	13.7	20.3	18.1	*
Asian or Pacific Islander	16.7	16.5	15.9	16.3	16.2	17.8	17.6	17.1	16.9	0.1
Hispanic or Latino ⁴	17.3	21.0	20.4	20.5	19.5	19.9	21.0	19.9	22.3	†0.6
White, not Hispanic or Latino ⁴	24.3	26.7	25.5	25.7	26.5	26.9	26.8	27.2	26.7	†0.4
Female	14.5	15.2	16.0	16.5	17.2	16.4	16.5	16.8	16.8	†0.8
White	15.4	16.0	16.9	17.5	18.0	17.7	17.3	17.9	17.8	†0.9
Black or African American	10.3	10.2	11.7	11.8	13.2	13.1	13.1	11.9	12.4	†1.5
American Indian or Alaska Native ³	*	*	13.5	*	*	13.1	12.1	15.6	12.5	*
Asian or Pacific Islander	9.1	11.8	11.4	12.2	12.6	9.5	12.2	11.6	11.7	0.6
Hispanic or Latina ⁴	13.9	13.3	13.7	14.0	15.6	15.3	14.8	17.3	15.5	†1.1
White, not Hispanic or Latina ⁴	15.6	16.2	17.3	18.0	18.4	18.0	17.7	18.0	18.1	†0.9

See footnotes at end of table.

Table 42 (page 4 of 4). Age-adjusted cancer incidence rates for selected cancer sites, by sex, race, and Hispanic origin: United States, selected geographic areas, selected years 1990–2010

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2012.htm#042>.

[Data are based on the Surveillance, Epidemiology, and End Results (SEER) Program's 13 population-based cancer registries]

Site, sex, race, and Hispanic origin	1990	1995	2000	2002	2003	2005	2008	2009	2010	1990–2010 APC [†]
Leukemia										
Number of new cases per 100,000 population ²										
Male	17.1	17.6	17.0	17.0	17.2	17.1	17.0	16.7	17.0	–0.1
White	18.0	19.0	18.1	18.4	18.3	18.6	18.1	17.8	18.3	0.0
Black or African American	16.0	13.3	14.1	12.6	14.4	12.4	13.9	14.1	12.5	–0.2
American Indian or Alaska Native ³	*	*	*	7.9	*	12.3	14.1	*	*	*
Asian or Pacific Islander	8.5	10.0	10.4	9.3	10.3	9.0	9.7	9.4	9.7	–0.3
Hispanic or Latino ⁴	12.2	14.6	12.9	12.3	12.3	13.1	12.3	12.2	12.3	0.2
White, not Hispanic or Latino ⁴	18.3	19.3	18.6	19.0	18.9	19.0	18.6	18.4	18.8	0.1
Female	9.9	10.2	10.3	10.0	10.0	9.9	10.5	9.9	10.1	0.1
White	10.3	10.9	11.0	10.8	10.6	10.4	11.1	10.4	10.7	0.2
Black or African American	8.5	8.2	9.7	7.4	9.0	9.3	7.8	7.6	8.6	–0.3
American Indian or Alaska Native ³	*	*	*	*	*	*	10.7	*	7.7	*
Asian or Pacific Islander	5.7	6.3	6.3	6.3	6.4	6.3	6.9	6.7	5.6	–0.1
Hispanic or Latina ⁴	8.6	8.2	7.8	8.6	7.2	8.4	9.4	8.2	8.6	0.3
White, not Hispanic or Latina ⁴	10.3	11.1	11.0	10.8	11.0	10.5	11.1	10.5	11.0	†0.3

[†] Annual percent change (APC) is significantly different from zero ($p < 0.05$).

0.0 APC is greater than –0.05 but less than 0.05.

* Estimates are considered unreliable. Data not shown if the rate is based on fewer than 16 cases for the time interval. The trend is not shown if it is based on fewer than 10 cases for at least 1 year within the time interval.

¹ APC was calculated by fitting a linear regression model to the natural logarithm of the yearly rates from 1990–2010.

² Age-adjusted by 5-year age groups to the year 2000 U.S. standard population. Age-adjusted rates are based on at least 16 cases. See [Appendix II, Age adjustment](#).

³ Starting with *Health, United States, 2007*, estimates for the American Indian or Alaska Native populations are based on the Contract Health Service Delivery Area (CHSDA) counties within SEER areas.

⁴ Starting with *Health, United States, 2007*, Hispanic data exclude cases from Alaska. The race groups, white, black, Asian or Pacific Islander, and American Indian or Alaska Native, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. The North American Association of Central Cancer Registries (NAACCR) Hispanic Identification Algorithm was used on a combination of variables to classify cases as Hispanic for analytic purposes. See the report, NAACCR Guideline for Enhancing Hispanic-Latino Identification, for more information. Available from: http://seer.cancer.gov/seerstat/variables/seer/yr1973_2006/race_ethnicity/. See [Appendix II, Hispanic origin](#).

NOTES: See [Appendix II, Incidence](#). Estimates are based on 13 SEER areas (November 2012 submission) and differ from published estimates based on 9 SEER areas or other submission dates. See [Appendix I, Surveillance, Epidemiology, and End Results Program \(SEER\)](#). The site variable distinguishes Kaposi Sarcoma and Mesothelioma as individual cancer sites. As a result, Kaposi Sarcoma and Mesothelioma cases do not contribute to other cancer sites. Estimates for 2001–2009 were computed using intercensal population estimates based on the 2000 and 2010 censuses. Data have been revised and differ from previous editions of *Health, United States*. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: National Institutes of Health, National Cancer Institute, Surveillance, Epidemiology, and End Results Program. Available from: <http://www.seer.cancer.gov>. See [Appendix I, Surveillance, Epidemiology, and End Results Program \(SEER\)](#).