

Table 40 (page 1 of 3). Human immunodeficiency virus (HIV) diagnoses, by year of diagnosis and selected characteristics: United States, 2008–2011

Updated data when available, Excel, and PDF: <http://www.cdc.gov/nchs/hus/contents2013.htm#040>.

[Data are based on reporting by 50 states and the District of Columbia]

Sex, race and Hispanic origin, age at diagnosis, and region of residence	Year of diagnosis				
	All years (2008–2011) ¹	2008	2009	2010	2011
		Estimated number of HIV diagnoses ²			
All persons ³	193,451	50,501	47,408	46,268	49,273
Male, 13 years and over.	149,238	38,104	36,392	35,918	38,825
Female, 13 years and over.	43,332	12,146	10,804	10,125	10,257
Children, under 13 years	881	252	213	226	192
Region of residence					
Northeast	40,520	10,711	10,061	9,725	10,024
Midwest	23,846	5,902	5,933	5,774	6,237
South	94,758	25,091	22,992	22,379	24,296
West	34,327	8,797	8,423	8,390	8,717
Male, 13 years and over					
Hispanic origin and race:					
Not Hispanic or Latino:					
White	47,185	12,206	11,583	11,355	12,041
Black or African American	63,508	16,277	15,491	15,293	16,447
Asian ⁴	2,764	685	612	646	821
Native Hawaiian or Other Pacific Islander	253	68	62	53	70
American Indian or Alaska Native	652	179	151	161	161
Hispanic or Latino ⁵	32,133	7,979	7,825	7,725	8,605
Multiple race	2,742	710	667	686	679
Age at diagnosis:					
13–14 years.	72	13	11	20	27
15–24 years.	31,366	7,113	7,480	8,093	8,680
25–34 years.	41,156	10,127	9,990	9,933	11,107
35–44 years.	36,363	10,319	9,146	8,371	8,527
45–54 years.	27,905	7,352	6,787	6,547	7,220
55–64 years.	9,773	2,503	2,363	2,357	2,551
65 years and over	2,603	677	616	597	713
Female, 13 years and over					
Hispanic origin and race:					
Not Hispanic or Latina:					
White	7,274	2,046	1,772	1,680	1,776
Black or African American	28,132	7,967	6,977	6,592	6,595
Asian ⁴	542	123	134	132	153
Native Hawaiian or Other Pacific Islander	46	10	18	9	8
American Indian or Alaska Native	210	44	53	61	51
Hispanic or Latina ⁵	6,309	1,681	1,635	1,463	1,530
Multiple race	820	274	214	188	144
Age at diagnosis:					
13–14 years.	100	30	20	25	26
15–24 years.	6,716	1,835	1,704	1,563	1,615
25–34 years.	11,060	3,206	2,687	2,581	2,586
35–44 years.	11,213	3,242	2,898	2,562	2,511
45–54 years.	9,530	2,618	2,388	2,229	2,295
55–64 years.	3,774	966	874	944	990
65 years and over	939	249	234	221	235
Children, under 13 years					
Hispanic origin and race:					
Not Hispanic or Latino:					
White	106	26	16	35	29
Black or African American	596	176	149	146	125
Asian ⁴	24	7	7	2	8
Native Hawaiian or Other Pacific Islander	0	0	0	0	0
American Indian or Alaska Native	2	2	0	0	0
Hispanic or Latino ⁵	129	31	35	38	25
Multiple race	25	10	5	5	4

See footnotes at end of table.

Table 40 (page 2 of 3). Human immunodeficiency virus (HIV) diagnoses, by year of diagnosis and selected characteristics: United States, 2008–2011

Updated data when available, Excel, and PDF: <http://www.cdc.gov/nchs/hus/contents2013.htm#040>.

[Data are based on reporting by 50 states and the District of Columbia]

Sex, race and Hispanic origin, age at diagnosis, and region of residence	Year of diagnosis				
	All years (2008–2011) ¹	2008	2009	2010	2011
	Percent distribution ⁶				
All persons ³	100.0	100.0	100.0	100.0	100.0
Male, 13 years and over	77.1	75.5	76.8	77.6	78.8
Female, 13 years and over	22.4	24.1	22.8	21.9	20.8
Children, under 13 years	0.5	0.5	0.4	0.5	0.4
Region of residence					
Northeast	20.9	21.2	21.2	21.0	20.3
Midwest	12.3	11.7	12.5	12.5	12.7
South	49.0	49.7	48.5	48.4	49.3
West	17.7	17.4	17.8	18.1	17.7
Male, 13 years and over					
Hispanic origin and race:					
Not Hispanic or Latino:					
White	31.6	32.0	31.8	31.6	31.0
Black or African American	42.6	42.7	42.6	42.6	42.4
Asian ⁴	1.9	1.8	1.7	1.8	2.1
Native Hawaiian or Other Pacific Islander	0.2	0.2	0.2	0.1	0.2
American Indian or Alaska Native	0.4	0.5	0.4	0.4	0.4
Hispanic or Latino ⁵	21.5	20.9	21.5	21.5	22.2
Multiple race	1.8	1.9	1.8	1.9	1.7
Age at diagnosis:					
13–14 years	0.0	0.0	0.0	0.1	0.1
15–24 years	21.0	18.7	20.6	22.5	22.4
25–34 years	27.6	26.6	27.5	27.7	28.6
35–44 years	24.4	27.1	25.1	23.3	22.0
45–54 years	18.7	19.3	18.6	18.2	18.6
55–64 years	6.5	6.6	6.5	6.6	6.6
65 years and over	1.7	1.8	1.7	1.7	1.8
Female, 13 years and over					
Hispanic origin and race:					
Not Hispanic or Latina:					
White	16.8	16.8	16.4	16.6	17.3
Black or African American	64.9	65.6	64.6	65.1	64.3
Asian ⁴	1.3	1.0	1.2	1.3	1.5
Native Hawaiian or Other Pacific Islander	0.1	0.1	0.2	0.1	0.1
American Indian or Alaska Native	0.5	0.4	0.5	0.6	0.5
Hispanic or Latina ⁵	14.6	13.8	15.1	14.4	14.9
Multiple race	1.9	2.3	2.0	1.9	1.4
Age at diagnosis:					
13–14 years	0.2	0.2	0.2	0.2	0.3
15–24 years	15.5	15.1	15.8	15.4	15.7
25–34 years	25.5	26.4	24.9	25.5	25.2
35–44 years	25.9	26.7	26.8	25.3	24.5
45–54 years	22.0	21.6	22.1	22.0	22.4
55–64 years	8.7	8.0	8.1	9.3	9.7
65 years and over	2.2	2.1	2.2	2.2	2.3
Children, under 13 years					
Hispanic origin and race:					
Not Hispanic or Latino:					
White	12.0	10.2	7.7	15.3	15.1
Black or African American	67.6	69.8	70.0	64.6	65.5
Asian ⁴	2.8	2.9	3.5	0.9	4.0
Native Hawaiian or Other Pacific Islander	—	—	—	—	—
American Indian or Alaska Native	0.2	0.9	—	—	—
Hispanic or Latino ⁵	14.6	12.3	16.4	16.8	13.1
Multiple race	2.8	4.1	2.4	2.3	2.3

See footnotes at end of table.

Table 40 (page 3 of 3). Human immunodeficiency virus (HIV) diagnoses, by year of diagnosis and selected characteristics: United States, 2008–2011

Updated data when available, Excel, and PDF: <http://www.cdc.gov/nchs/hus/contents2013.htm#040>.

[Data are based on reporting by 50 states and the District of Columbia]

0.0 Quantity more than zero but less than 0.05.

– Quantity zero.

¹Based on diagnoses that occurred during 2008–2011 that were reported to CDC through June 2012.

²Numbers are point estimates that result from statistical adjustments for reporting delays and missing risk factor information. The estimates do not include adjustments for incomplete reporting. See [Appendix I, National HIV Surveillance System](#).

³All persons totals were calculated independent of values for subpopulations. Consequently, sums of subpopulations may not equal totals for all persons.

⁴Includes Asian and Pacific Islander legacy cases.

⁵Persons of Hispanic origin may be of any race. See [Appendix II, Hispanic origin](#).

⁶Percents may not sum to 100% due to rounding.

NOTES: See [Appendix II, Human immunodeficiency virus \(HIV\); Acquired immunodeficiency syndrome \(AIDS\)](#), for discussion of HIV/AIDS diagnoses reporting definitions and other issues affecting interpretation of trends. Data shown are for the 50 states and the District of Columbia. Starting with *Health, United States, 2013*, this table shows HIV diagnoses data. In previous editions of *Health, United States* AIDS diagnoses data were shown.

SOURCE: CDC, National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention. Division of HIV/AIDS Prevention. HIV Surveillance Report. Diagnoses of HIV infection in the United States and Dependent Areas, 2011 (vol. 23). Atlanta, GA: U.S. Department of Health and Human Services, CDC. 2011 and unpublished data. Available from: http://www.cdc.gov/hiv/library/reports/surveillance/2011/surveillance_Report_vol_23.html. See [Appendix I, National HIV Surveillance System](#).