

Table 58 (page 1 of 3). Current cigarette smoking among adults aged 18 and over, by sex, race, Hispanic origin, age, and education level: United States, average annual, selected years 1990–1992 through 2010–2012

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2013.htm#058>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Male			Female		
	1990–1992 ¹	1999–2001	2010–2012	1990–1992 ¹	1999–2001	2010–2012
18 years and over, age-adjusted ²	Percent of adults who were current cigarette smokers ³					
All persons ⁴	27.9	25.0	21.0	23.7	21.1	16.7
Race ⁵						
White only	27.4	25.1	21.1	24.3	22.2	17.6
Black or African American only	33.9	27.2	22.8	23.1	19.7	15.4
American Indian or Alaska Native only	34.2	30.3	23.2	36.7	34.7	23.2
Asian only	24.8	20.3	14.7	6.3	6.7	5.2
Native Hawaiian or Other Pacific Islander only	---	*	*	---	*	*
2 or more races	---	34.4	26.0	---	30.7	22.9
American Indian or Alaska Native; White	---	38.7	33.6	---	38.9	30.2
Hispanic origin and race ⁵						
Hispanic or Latino	25.7	22.2	16.2	15.8	12.1	8.3
Mexican	26.2	21.9	16.3	14.8	10.6	7.5
Not Hispanic or Latino	28.1	25.5	22.1	24.4	22.3	18.3
White only	27.7	25.5	22.5	25.2	23.5	19.8
Black or African American only	33.9	27.2	23.2	23.2	19.7	15.5
18 years and over, crude						
All persons ⁴	28.4	25.5	21.2	23.6	21.0	16.6
Race ⁵						
White only	27.8	25.4	21.1	24.1	21.7	17.2
Black or African American only	33.2	27.5	23.4	23.3	19.8	15.6
American Indian or Alaska Native only	35.5	31.8	24.6	37.3	36.9	23.8
Asian only	24.9	21.4	15.6	6.3	6.9	5.3
Native Hawaiian or Other Pacific Islander only	---	*	*	---	*	*
2 or more races	---	35.9	26.9	---	31.5	22.5
American Indian or Alaska Native; White	---	41.1	33.6	---	40.1	30.9
Hispanic origin and race ⁵						
Hispanic or Latino	26.5	23.2	16.7	16.6	12.6	8.4
Mexican	27.1	22.8	16.6	15.0	11.0	7.6
Not Hispanic or Latino	28.5	25.8	22.0	24.2	21.9	17.8
White only	28.0	25.5	22.0	24.8	22.7	18.9
Black or African American only	33.3	27.5	23.7	23.3	19.8	15.8
Age and Hispanic origin and race ⁵						
18–24 years:						
Hispanic or Latino	19.3	22.6	14.6	12.8	12.9	7.6
Not Hispanic or Latino:						
White only	28.9	32.7	25.2	28.7	30.8	20.8
Black or African American only	17.7	21.9	16.9	10.8	13.0	10.5
25–34 years:						
Hispanic or Latino	29.9	23.2	18.4	19.2	12.5	8.6
Not Hispanic or Latino:						
White only	32.7	30.8	31.1	30.9	27.4	24.8
Black or African American only	34.6	23.3	25.8	29.2	16.9	18.2
35–44 years:						
Hispanic or Latino	32.1	25.3	17.1	19.9	14.1	8.1
Not Hispanic or Latino:						
White only	32.3	29.6	24.0	27.3	28.3	23.0
Black or African American only	44.1	32.0	24.5	31.3	27.5	15.4
45–64 years:						
Hispanic or Latino	26.6	24.7	17.5	17.1	13.5	10.5
Not Hispanic or Latino:						
White only	28.4	25.1	22.6	26.1	22.1	20.6
Black or African American only	38.0	34.0	28.6	26.1	23.6	19.6
65 years and over:						
Hispanic or Latino	16.1	12.6	11.3	6.6	5.9	4.7
Not Hispanic or Latino:						
White only	14.2	10.0	9.4	12.3	9.8	8.3
Black or African American only	25.2	17.6	13.8	10.7	11.0	9.3

See footnotes at end of table.

Table 58 (page 2 of 3). Current cigarette smoking among adults aged 18 and over, by sex, race, Hispanic origin, age, and education level: United States, average annual, selected years 1990–1992 through 2010–2012

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2013.htm#058>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Male			Female		
	1990–1992 ¹	1999–2001	2010–2012	1990–1992 ¹	1999–2001	2010–2012
Percent of poverty level ^{2,6}		Percent of adults who were current cigarette smokers ³				
Below 100%	40.5	36.5	33.2	30.7	29.1	25.1
100%–199%	35.0	32.8	27.8	26.9	25.6	21.9
200%–399%	26.5	27.3	22.5	22.6	22.3	17.5
400% or more	22.5	18.8	13.7	19.0	15.9	10.1
Hispanic origin and race and percent of poverty level ^{2,5,6}						
Hispanic or Latino:						
Below 100%	29.2	25.3	18.5	16.3	14.4	9.4
100%–199%	29.5	22.0	17.2	16.0	11.8	8.2
200%–399%	23.7	23.6	16.4	15.9	12.0	9.0
400% or more	19.7	18.1	12.4	13.6	9.4	5.9
Not Hispanic or Latino:						
White only:						
Below 100%	44.2	40.7	43.1	37.8	38.3	37.2
100%–199%	36.3	37.5	33.4	31.1	32.0	30.3
200%–399%	26.4	28.5	24.9	23.7	24.8	20.8
400% or more	22.5	19.1	14.4	19.5	17.1	11.3
Black or African American only:						
Below 100%	43.5	40.6	35.1	28.9	27.7	24.4
100%–199%	36.0	33.9	29.7	20.3	21.3	16.8
200%–399%	31.4	24.9	21.0	21.4	17.3	12.7
400% or more	24.3	17.9	12.2	19.2	12.6	6.7
Disability measure ^{2,7}						
Any basic actions difficulty or complex activity limitation	---	33.1	29.3	---	28.1	25.3
Any basic actions difficulty	---	33.2	29.8	---	28.2	25.5
Any complex activity limitation	---	37.6	31.7	---	30.6	29.8
No disability	---	22.8	18.3	---	18.8	13.5
Education, Hispanic origin, and race ^{5,8}						
25 years and over, age-adjusted ⁹						
No high school diploma or GED:						
Hispanic or Latino	30.2	24.3	17.2	15.8	12.1	7.6
Not Hispanic or Latino:						
White only	46.1	43.5	45.2	40.4	39.3	40.8
Black or African American only	45.4	40.0	35.9	31.3	29.4	28.1
High school diploma or GED:						
Hispanic or Latino	29.6	24.1	22.0	18.4	12.5	10.5
Not Hispanic or Latino:						
White only	32.9	31.8	32.2	28.4	29.2	30.0
Black or African American only	38.2	31.4	28.7	25.4	23.0	19.0
Some college or more:						
Hispanic or Latino	20.4	17.1	11.5	14.3	11.1	7.5
Not Hispanic or Latino:						
White only	19.3	17.6	15.3	18.1	16.7	14.6
Black or African American only	25.6	19.2	18.0	22.8	16.9	12.4

See footnotes at end of table.

Table 58 (page 3 of 3). Current cigarette smoking among adults aged 18 and over, by sex, race, Hispanic origin, age, and education level: United States, average annual, selected years 1990–1992 through 2010–2012

Updated data when available, Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2013.htm#058>.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

-- Data not available.

* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30%.

¹Data prior to 1997 are not strictly comparable with data for later years due to the 1997 questionnaire redesign. See [Appendix I, National Health Interview Survey \(NHIS\)](#).

²Estimates are age-adjusted to the year 2000 standard population using five age groups: 18–24 years, 25–34 years, 35–44 years, 45–64 years, and 65 years and over. See [Appendix II, Age adjustment](#). For age groups where smoking is 0% or 100%, the age-adjustment procedure was modified to substitute the percentage smoking from the previous 3-year period.

³Starting with 1993 data, current cigarette smokers were defined as ever smoking 100 cigarettes in their lifetime and smoking now every day or some days. For previous definition, see [Appendix II, Cigarette smoking](#).

⁴Includes all other races not shown separately, unknown education level, and unknown disability measure.

⁵The race groups, white, black, American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, and 2 or more races, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The five single-race categories plus multiple-race categories shown in the table conform to the 1997 Standards. Starting with 1999–2001 data, race-specific estimates are for persons who reported only one racial group; the category 2 or more races includes persons who reported more than one racial group. Prior to 1999, data were tabulated according to the 1977 Standards with four racial groups, and the Asian only category included Native Hawaiian or Other Pacific Islander. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See [Appendix II, Hispanic origin; Race](#).

⁶Percent of poverty level is based on family income and family size and composition using U.S. Census Bureau poverty thresholds. Missing family income data were imputed for 1990 and beyond. See [Appendix II, Family income; Poverty; Table VI](#).

⁷Any basic actions difficulty or complex activity limitation is defined as having one or more of the following limitations or difficulties: movement difficulty, emotional difficulty, sensory (seeing or hearing) difficulty, cognitive difficulty, self-care (activities of daily living or instrumental activities of daily living) limitation, social limitation, or work limitation. For more information, see [Appendix II, Basic actions difficulty; Complex activity limitation](#). Starting with 2007 data, the hearing question, a component of the basic actions difficulty measure, was revised. Consequently, data prior to 2007 are not comparable with data for 2007 and beyond. For more information on the impact of the revised hearing question, see [Appendix II, Hearing trouble](#).

⁸Education categories shown are for 1997 and subsequent years. GED is General Educational Development high school equivalency diploma. In years prior to 1997, the following categories based on number of years of school completed were used: less than 12 years, 12 years, 13 years or more. See [Appendix II, Education](#).

⁹Estimates are age-adjusted to the year 2000 standard using four age groups: 25–34 years, 35–44 years, 45–64 years, and 65 years and over. See [Appendix II, Age adjustment](#).

NOTES: Standard errors for selected years are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Health Interview Survey. Data are from the following questionnaire supplements: health promotion and disease prevention (1990–1991), cancer control and cancer epidemiology (1992), and year 2000 objectives (1993–1995). Starting with 1997, data are from the family core and sample adult questionnaires. See [Appendix I, National Health Interview Survey \(NHIS\)](#).