

Data table for Figure 24. Age-adjusted death rates for human immunodeficiency virus (HIV) disease for all ages, by sex and race and Hispanic origin: United States, 1990–2010

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2013.htm#fig24>

Year	Male	Female	White, not Hispanic male	Black, not Hispanic male	Hispanic male	American Indian or Alaska Native male	Asian or Pacific Islander male	White, not Hispanic female	Black, not Hispanic female	Hispanic female	American Indian or Alaska Native female	Asian or Pacific Islander female
Deaths per 100,000 population												
1990	18.5	2.2	14.1	47.1	28.8	3.3	4.3	0.7	10.3	3.8	*	*
1991	21.0	2.7	15.5	55.6	31.9	6.6	4.3	0.9	12.2	4.9	*	*
1992	23.3	3.2	16.7	65.4	35.1	4.8	4.6	1.0	14.9	5.7	*	0.5
1993	25.1	3.9	17.4	74.5	35.2	7.9	5.3	1.3	17.8	6.8	*	0.8
1994	27.5	4.8	18.5	88.2	41.5	8.9	6.7	1.6	23.0	8.0	*	0.7
1995	27.3	5.3	17.9	90.6	40.8	10.5	6.0	1.7	25.0	8.8	2.5	0.6
1996	19.0	4.2	11.2	71.8	28.0	6.4	4.4	1.3	21.4	6.3	*	0.5
1997	9.6	2.6	4.8	41.5	14.0	3.3	1.6	0.7	13.9	3.3	*	*
1998	7.6	2.2	3.7	33.8	10.2	3.5	1.3	0.5	12.2	2.8	*	*
1999	8.2	2.5	4.0	36.5	10.9	4.2	1.4	0.7	13.4	3.0	*	*
2000	7.9	2.5	3.8	35.5	10.6	3.5	1.2	0.7	13.5	2.9	*	*
2001	7.6	2.5	3.6	34.3	9.8	4.1	1.2	0.6	13.8	2.7	*	*
2002	7.4	2.5	3.5	33.8	9.3	3.3	1.5	0.6	13.8	2.7	*	*
2003	7.1	2.4	3.4	31.8	9.4	3.4	1.1	0.6	13.1	2.6	1.4	*
2004	6.6	2.4	3.1	29.5	8.4	4.1	1.2	0.6	13.3	2.4	1.4	*
2005	6.3	2.3	3.0	28.4	7.7	3.7	1.0	0.6	12.2	1.9	1.3	*
2006	5.9	2.2	2.8	26.6	7.2	2.9	1.1	0.6	12.4	1.9	1.3	*
2007	5.4	2.1	2.5	24.6	6.5	3.3	0.8	0.5	11.7	1.8	1.5	*
2008	4.8	1.9	2.3	21.8	5.5	2.8	1.0	0.5	10.0	1.7	*	0.3
2009	4.4	1.7	2.0	20.1	5.0	2.4	0.7	0.4	9.2	1.4	*	*
2010	3.8	1.4	1.8	17.0	4.6	2.6	0.7	0.4	7.9	1.1	*	*

* Rates based on fewer than 20 deaths are considered unreliable and are not shown.

NOTES: Rates are based on resident population. Rates are age-adjusted using 11 age groups. Age-adjusted rates are calculated using the year 2000 standard population with unrounded population numbers. Highly active antiretroviral therapy (HAART) was introduced in 1996. The lines in the data table around 1995 to 1997 denote the period from pre-HAART to widespread HAART use. Categories for the coding and classification of HIV disease were introduced in the United States in 1987. For the period 1990–1998, underlying cause of death was coded according to the 9th Revision of the *International Classification of Diseases* (ICD–9). Starting with 1999 data, cause of death is coded according to ICD–10. Persons of Hispanic origin may be of any race. See [Appendix II, Age adjustment](#); [Cause of death](#); [Hispanic origin](#); [Human immunodeficiency virus \(HIV\) disease](#); [Table IV](#). See related [Table 31](#).

SOURCE: CDC/NCHS, National Vital Statistics System. See [Appendix I, National Vital Statistics System \(NVSS\)](#).