

**Table 9 (page 1 of 6). Contraceptive use in the past month among women aged 15–44, by age, race and Hispanic origin, and method of contraception: United States, selected years 1982–2013**

Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2014.htm#009>.

[Data are based on household interviews of samples of women of childbearing age]

Race and Hispanic origin and year <sup>1</sup>	Age, in years				
	15–44	15–19	20–24	25–34	35–44
Number of women in population, in thousands					
All women: <sup>2</sup>					
1982 . . . . .	54,099	9,521	10,629	19,644	14,305
1995 . . . . .	60,201	8,961	9,041	20,758	21,440
2002 . . . . .	61,561	9,834	9,840	19,522	22,365
2006–2010 . . . . .	61,755	10,478	10,365	19,722	21,190
2011–2013 . . . . .	60,887	9,547	10,338	20,790	20,212
Not Hispanic or Latina:					
White only:					
1982 . . . . .	41,279	7,010	8,081	14,945	11,243
1995 . . . . .	42,154	5,865	6,020	14,471	15,798
2002 . . . . .	39,498	6,069	5,938	12,073	15,418
2006–2010 . . . . .	37,384	6,034	6,173	11,953	13,224
2011–2013 . . . . .	34,674	4,889	5,606	12,086	12,094
Black or African American only:					
1982 . . . . .	6,825	1,383	1,456	2,392	1,593
1995 . . . . .	8,060	1,334	1,305	2,780	2,641
2002 . . . . .	8,250	1,409	1,396	2,587	2,857
2006–2010 . . . . .	8,451	1,566	1,493	2,621	2,771
2011–2013 . . . . .	8,491	1,432	1,572	2,794	2,693
Hispanic or Latina: <sup>3</sup>					
1982 . . . . .	4,393	886	811	1,677	1,018
1995 . . . . .	6,702	1,150	1,163	2,450	1,940
2002 . . . . .	9,107	1,521	1,632	3,249	2,705
2006–2010 . . . . .	10,474	1,904	1,734	3,611	3,225
2011–2013 . . . . .	12,024	2,144	2,105	4,062	3,713
Percent of women in population using contraception					
All women: <sup>2</sup>					
1982 . . . . .	55.7	24.2	55.8	66.7	61.6
1995 . . . . .	64.2	29.8	63.5	71.1	72.3
2002 . . . . .	61.9	31.5	60.7	68.6	69.9
2006–2010 . . . . .	62.2	30.5	58.3	67.3	74.9
2011–2013 . . . . .	61.7	33.3	60.4	67.4	70.0
Not Hispanic or Latina:					
White only:					
1982 . . . . .	57.3	23.6	58.7	67.8	63.5
1995 . . . . .	66.2	30.5	65.4	72.9	73.6
2002 . . . . .	64.6	35.0	66.3	69.9	71.4
2006–2010 . . . . .	65.6	35.1	62.7	69.7	77.2
2011–2013 . . . . .	65.3	40.1	60.9	69.2	73.7
Black or African American only:					
1982 . . . . .	51.6	29.8	52.3	63.5	52.0
1995 . . . . .	62.3	36.1	67.6	66.8	68.3
2002 . . . . .	57.6	32.9	50.8	67.9	63.8
2006–2010 . . . . .	54.2	25.5	50.0	60.9	66.2
2011–2013 . . . . .	57.9	21.0	64.9	65.2	65.9
Hispanic or Latina: <sup>3</sup>					
1982 . . . . .	50.6	*	*36.8	67.2	59.0
1995 . . . . .	59.0	26.1	50.6	69.2	70.8
2002 . . . . .	59.0	20.4	57.4	66.2	72.9
2006–2010 . . . . .	59.7	22.3	54.0	66.0	77.7
2011–2013 . . . . .	57.3	26.3	52.7	67.8	66.5

See footnotes at end of table.

**Table 9 (page 2 of 6). Contraceptive use in the past month among women aged 15–44, by age, race and Hispanic origin, and method of contraception: United States, selected years 1982–2013**

Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2014.htm#009>.

[Data are based on household interviews of samples of women of childbearing age]

Race and Hispanic origin and year <sup>1</sup>	Age, in years				
	15–44	15–19	20–24	25–34	35–44
Number of sexually active women in population, in thousands <sup>4</sup>					
All women: <sup>2</sup>					
1982	---	---	---	---	---
1995	41,796	3,341	6,272	15,687	16,495
2002	42,683	3,775	6,798	14,857	17,252
2006–2010	43,145	3,896	6,944	14,785	17,520
2011–2013	41,762	3,752	7,136	15,464	15,409
Not Hispanic or Latina:					
White only:					
1982	---	---	---	---	---
1995	29,994	2,202	4,276	11,194	12,322
2002	28,079	2,519	4,329	9,224	12,006
2006–2010	27,105	2,471	4,341	9,105	11,188
2011–2013	24,720	2,168	3,840	9,118	9,593
Black or African American only:					
1982	---	---	---	---	---
1995	5,579	598	967	2,039	1,975
2002	5,611	564	949	1,978	2,121
2006–2010	5,526	517	939	1,946	2,124
2011–2013	5,680	464	1,199	2,040	1,977
Hispanic or Latina: <sup>3</sup>					
1982	---	---	---	---	---
1995	4,330	409	685	1,794	1,442
2002	6,075	405	1,070	2,462	2,138
2006–2010	6,978	563	1,076	2,656	2,683
2011–2013	7,853	723	1,335	3,076	2,720
Percent of sexually active women in population using contraception <sup>4</sup>					
All women: <sup>2</sup>					
1982	---	---	---	---	---
1995	92.5	80.2	91.7	94.0	93.9
2002	89.3	82.0	87.9	90.2	90.7
2006–2010	89.0	82.0	87.0	89.8	90.6
2011–2013	90.0	84.6	87.5	90.6	91.8
Not Hispanic or Latina:					
White only:					
1982	---	---	---	---	---
1995	93.0	81.7	93.0	93.9	94.2
2002	90.9	84.4	90.9	91.5	91.7
2006–2010	90.5	85.7	89.1	91.6	91.2
2011–2013	91.6	90.4	89.0	91.7	92.9
Black or African American only:					
1982	---	---	---	---	---
1995	90.0	80.0	91.3	91.6	90.9
2002	84.7	82.2	74.8	88.9	86.0
2006–2010	82.8	77.3	79.4	82.1	86.3
2011–2013	86.6	*	85.2	89.3	89.8
Hispanic or Latina: <sup>3</sup>					
1982	---	---	---	---	---
1995	91.4	75.5	82.5	95.4	95.2
2002	88.4	76.4	87.5	87.4	92.3
2006–2010	89.6	75.5	87.0	89.7	93.4
2011–2013	87.8	78.0	83.1	89.5	90.8

See footnotes at end of table.

**Table 9 (page 3 of 6). Contraceptive use in the past month among women aged 15–44, by age, race and Hispanic origin, and method of contraception: United States, selected years 1982–2013**

Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2014.htm#009>.

[Data are based on household interviews of samples of women of childbearing age]

Method of contraception and year	Age, in years				
	15–44	15–19	20–24	25–34	35–44
Female sterilization					
Percent of contracepting women					
1982 . . . . .	23.2	—	*4.5	22.1	43.5
1995 . . . . .	27.8	*	4.0	23.8	45.0
2002 . . . . .	27.0	—	3.6	21.6	45.8
2006–2010 . . . . .	26.6	*	*2.6	22.9	44.0
2011–2013 . . . . .	25.1	—	*2.3	21.7	44.2
Male sterilization					
1982 . . . . .	10.9	*	*3.6	10.1	19.9
1995 . . . . .	10.9	—	*	7.8	19.5
2002 . . . . .	10.2	—	*	7.2	18.2
2006–2010 . . . . .	10.8	*	*	7.1	19.8
2011–2013 . . . . .	9.0	—	*	4.2	19.6
Implant and other hormonal contraceptives <sup>5</sup>					
1982 . . . . .	...	...	...	...	...
1995 . . . . .	1.3	*	3.7	*1.3	*
2002 . . . . .	1.0	*	*	*1.7	*
2006–2010 . . . . .	3.4	*4.7	6.4	4.4	*1.1
2011–2013 . . . . .	3.9	*	9.0	4.6	*
Injectable <sup>6</sup>					
1982 . . . . .	...	...	...	...	...
1995 . . . . .	3.0	9.7	6.1	2.9	*0.8
2002 . . . . .	5.5	14.2	10.6	5.5	*1.9
2006–2010 . . . . .	3.9	11.4	5.9	4.2	*1.3
2011–2013 . . . . .	4.6	11.4	7.1	4.8	*
Birth control pill <sup>7</sup>					
1982 . . . . .	28.0	63.9	55.1	25.7	*3.7
1995 . . . . .	27.0	43.8	52.1	33.4	8.7
2002 . . . . .	31.0	53.8	52.5	34.8	15.0
2006–2010 . . . . .	28.4	53.6	47.3	30.5	14.3
2011–2013 . . . . .	26.7	56.5	42.6	25.3	14.4
Intrauterine device					
1982 . . . . .	7.1	*	*4.2	9.7	6.9
1995 . . . . .	0.8	—	*	*0.8	1.1
2002 . . . . .	2.1	*	1.8	3.7	*1.3
2006–2010 . . . . .	5.6	*	5.6	7.2	4.9
2011–2013 . . . . .	10.7	*	10.2	15.1	8.4

See footnotes at end of table.

**Table 9 (page 4 of 6). Contraceptive use in the past month among women aged 15–44, by age, race and Hispanic origin, and method of contraception: United States, selected years 1982–2013**

Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2014.htm#009>.

[Data are based on household interviews of samples of women of childbearing age]

Method of contraception and year	Age, in years				
	15–44	15–19	20–24	25–34	35–44
Diaphragm					
Percent of contracepting women					
1982 . . . . .	8.1	*6.0	10.2	10.3	4.0
1995 . . . . .	1.9	*	*	1.7	2.8
2002 . . . . .	*	–	*	*	*
2006–2010 . . . . .	*	–	–	*	*
2011–2013 . . . . .	*	–	–	*	–
Condom					
1982 . . . . .	12.0	20.8	10.7	11.4	11.3
1995 . . . . .	23.4	45.8	33.7	23.7	15.3
2002 . . . . .	23.8	44.6	36.0	23.1	15.6
2006–2010 . . . . .	23.1	34.7	39.6	25.2	12.8
2011–2013 . . . . .	22.8	34.1	38.5	23.8	12.4
Periodic abstinence-calendar rhythm					
1982 . . . . .	3.3	2.0	3.1	3.3	3.7
1995 . . . . .	3.3	*	*1.5	3.7	3.9
2002 . . . . .	2.0	*	*2.3	*1.7	*2.4
2006–2010 . . . . .	1.7	*	*	2.0	2.1
2011–2013 . . . . .	2.9	*	*	*2.7	*2.5
Periodic abstinence-natural family planning					
1982 . . . . .	0.6	–	*	0.9	*
1995 . . . . .	*0.5	–	*	*0.7	*
2002 . . . . .	*0.4	–	–	*	*
2006–2010 . . . . .	*	–	*	*	*
2011–2013 . . . . .	*	*	*	*	*
Withdrawal					
1982 . . . . .	2.0	2.9	3.0	1.8	1.3
1995 . . . . .	6.1	13.2	7.1	6.0	4.5
2002 . . . . .	8.8	15.0	11.9	10.7	4.7
2006–2010 . . . . .	10.1	14.5	15.1	10.2	7.3
2011–2013 . . . . .	12.6	17.9	16.8	15.3	6.9
Other methods <sup>8</sup>					
1982 . . . . .	4.9	2.6	5.4	4.8	5.3
1995 . . . . .	3.2	*	3.2	3.1	3.4
2002 . . . . .	1.7	*	*0.9	*1.5	*1.8
2006–2010 . . . . .	0.6	*	*	*0.8	*
2011–2013 . . . . .	*	*	*	*	*

See footnotes at end of table.

**Table 9 (page 5 of 6). Contraceptive use in the past month among women aged 15–44, by age, race and Hispanic origin, and method of contraception: United States, selected years 1982–2013**

Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2014.htm#009>.

[Data are based on household interviews of samples of women of childbearing age]

Method of contraception and year	Not Hispanic or Latina <sup>1</sup>		
	White only	Black or African American only	Hispanic or Latina <sup>3</sup>
Female sterilization			
Percent of contracepting women			
1982 . . . . .	22.0	30.0	23.0
1995 . . . . .	24.5	39.9	36.6
2002 . . . . .	23.9	39.2	33.8
2006–2010 . . . . .	23.6	37.3	31.7
2011–2013 . . . . .	21.4	36.8	32.9
Male sterilization			
1982 . . . . .	13.0	*1.5	*
1995 . . . . .	13.7	*1.8	*4.0
2002 . . . . .	12.9	*	4.7
2006–2010 . . . . .	14.2	*	5.8
2011–2013 . . . . .	12.9	*1.9	4.0
Implant and other hormonal contraceptives <sup>5</sup>			
1982 . . . . .	...	...	...
1995 . . . . .	*1.0	*2.4	*2.0
2002 . . . . .	*0.6	*	*2.6
2006–2010 . . . . .	3.0	4.7	3.3
2011–2013 . . . . .	*3.3	*	*4.1
Injectable <sup>6</sup>			
1982 . . . . .	...	...	...
1995 . . . . .	2.4	5.4	4.7
2002 . . . . .	4.3	9.4	7.8
2006–2010 . . . . .	2.5	8.9	6.0
2011–2013 . . . . .	*3.1	10.1	*4.7
Birth control pill <sup>7</sup>			
1982 . . . . .	26.4	37.9	30.2
1995 . . . . .	28.7	23.7	23.0
2002 . . . . .	34.9	23.1	22.0
2006–2010 . . . . .	33.1	18.7	20.2
2011–2013 . . . . .	30.2	17.9	19.2
Intrauterine device			
1982 . . . . .	5.8	9.3	19.2
1995 . . . . .	0.7	*	*
2002 . . . . .	1.7	*	5.3
2006–2010 . . . . .	5.6	5.0	6.8
2011–2013 . . . . .	11.0	6.5	13.5
Diaphragm			
1982 . . . . .	9.2	*3.2	*
1995 . . . . .	2.3	*	*
2002 . . . . .	*	*	—
2006–2010 . . . . .	*	*	*
2011–2013 . . . . .	—	—	—
Condom			
1982 . . . . .	13.1	6.3	*6.9
1995 . . . . .	22.5	24.9	21.2
2002 . . . . .	21.7	29.6	24.1
2006–2010 . . . . .	20.8	29.9	22.2
2011–2013 . . . . .	21.5	25.9	21.0
Periodic abstinence-calendar rhythm			
1982 . . . . .	3.2	2.9	3.9
1995 . . . . .	3.3	*1.7	3.2
2002 . . . . .	2.3	*	*
2006–2010 . . . . .	1.3	*	*2.7
2011–2013 . . . . .	*2.7	*2.2	*

See footnotes at end of table.

**Table 9 (page 6 of 6). Contraceptive use in the past month among women aged 15–44, by age, race and Hispanic origin, and method of contraception: United States, selected years 1982–2013**

Excel, PDF, more data years, and standard errors: <http://www.cdc.gov/nchs/hus/contents2014.htm#009>.

[Data are based on household interviews of samples of women of childbearing age]

Method of contraception and year	Not Hispanic or Latina <sup>1</sup>		
	White only	Black or African American only	Hispanic or Latina <sup>3</sup>
Periodic abstinence-natural family planning			
Percent of contracepting women			
1982 . . . . .	0.7	0.3	—
1995 . . . . .	0.7	*	*
2002 . . . . .	*	*	*
2006–2010 . . . . .	*	*	*
2011–2013 . . . . .	*	—	*
Withdrawal			
1982 . . . . .	2.1	1.3	2.6
1995 . . . . .	6.4	3.3	5.7
2002 . . . . .	9.5	4.8	6.3
2006–2010 . . . . .	10.3	7.1	10.4
2011–2013 . . . . .	12.8	10.1	13.0
Other methods <sup>8</sup>			
1982 . . . . .	4.6	7.3	5.0
1995 . . . . .	3.3	3.8	*2.2
2002 . . . . .	*1.7	*1.9	*1.2
2006–2010 . . . . .	0.6	*	*
2011–2013 . . . . .	*	*	*

\* Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%–30%. Data not shown have an RSE greater than 30% or based on fewer than 100 sample cases.

--- Data not available.

— Quantity zero.

. . . Data not applicable.

<sup>1</sup>Starting with 1995 data, race-specific estimates are tabulated according to 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. Starting with 1995 data, race-specific estimates are for persons who reported only one racial group. Prior to data year 1995, data were tabulated according to the 1977 Standards. Estimates for single-race categories prior to 1995 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. See Appendix II, Race.

<sup>2</sup>Includes women of other or multiple race not shown separately.

<sup>3</sup>Persons of Hispanic origin may be of any race. See Appendix II, Hispanic origin.

<sup>4</sup>Had sexual (vaginal) intercourse in the past 3 months.

<sup>5</sup>Data collected starting with the 1995 survey. Includes data about the contraceptive patch, with data collection starting in the 2002 survey, and the contraceptive ring, with data collection starting in the 2006–2010 survey.

<sup>6</sup>Data collected starting with the 1995 survey.

<sup>7</sup>In 2011–2013, includes the oral contraceptive pill only. In previous surveys includes the oral contraceptive pill and emergency contraception/morning-after pill.

<sup>8</sup>In 2011–2013, includes emergency contraception, female condom/vaginal pouch, foam, cervical cap, Today sponge, suppository or insert, jelly or cream (without diaphragm), and other methods. See Appendix II, Contraception, for the list of other methods reported in previous surveys.

NOTES: Survey collects up to four methods of contraception used in the month of interview. Percents may not add to the total because more than one method could have been used in the month of interview. These data replace estimates of most effective method used and may differ from previous editions of *Health, United States*. Standard errors for selected years are available in the spreadsheet version of this table. Available from: <http://www.cdc.gov/nchs/hus.htm>. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Survey of Family Growth. See Appendix I, National Survey of Family Growth (NSFG).