

Table 30 (page 1 of 3). Death rates for drug poisoning and drug poisoning involving opioid analgesics, by sex, age, race, and Hispanic origin: United States, selected years 1999–2013

Updated data when available, Excel, and PDF: <http://www.cdc.gov/nchs/hus/contents2014.htm#030>.

[Data are based on death certificates]

Sex, age, race, and Hispanic origin	1999	2000	2004	2005	2009	2010	2011	2012	2013
Drug poisoning deaths per 100,000 resident population ¹									
All persons									
All ages, age-adjusted ²	6.1	6.2	9.4	10.1	11.9	12.3	13.2	13.1	13.8
All ages, crude	6.0	6.2	9.4	10.1	12.1	12.4	13.3	13.2	13.9
Under 15 years	0.1	0.1	0.2	0.2	0.2	0.2	0.2	0.2	0.2
15–24 years	3.2	3.7	6.6	6.9	7.7	8.2	8.6	8.0	8.3
25–34 years	8.1	7.9	11.9	13.6	17.2	18.4	20.2	20.1	20.9
35–44 years	14.0	14.3	19.3	19.6	20.5	20.8	22.5	22.1	23.0
45–54 years	11.1	11.6	19.3	21.1	25.4	25.1	26.7	26.9	27.5
55–64 years	4.2	4.2	7.8	9.0	13.7	15.0	15.9	16.6	19.2
65–74 years	2.4	2.0	2.9	3.2	4.7	4.7	5.4	5.8	6.4
75–84 years	2.8	2.4	2.9	3.1	3.8	3.4	3.4	3.4	3.6
85 years and over	3.8	4.4	4.0	4.1	4.4	4.7	4.2	4.3	4.3
Male									
All ages, age-adjusted ²	8.2	8.3	11.8	12.8	14.8	15.0	16.1	16.1	17.0
All ages, crude	8.2	8.4	11.9	12.9	15.0	15.2	16.3	16.3	17.2
Under 15 years	0.1	0.2	0.2	0.2	0.2	0.3	0.2	0.2	0.2
15–24 years	4.5	5.3	9.6	10.0	11.3	11.6	12.4	11.4	11.7
25–34 years	11.5	11.3	16.6	18.7	24.0	25.0	27.5	27.0	28.6
35–44 years	19.2	19.5	23.8	24.4	25.2	24.9	26.8	27.1	28.1
45–54 years	15.2	15.7	23.8	25.8	29.1	28.5	30.4	30.4	31.5
55–64 years	4.9	4.4	8.6	10.6	16.0	17.3	18.5	19.4	22.7
65–74 years	2.7	2.1	2.9	3.3	4.8	4.5	5.4	6.2	6.9
75–84 years	2.5	2.5	2.8	3.4	3.5	3.6	3.4	3.2	3.7
85 years and over	4.4	5.9	4.8	5.2	5.2	5.1	4.3	5.3	5.9
Female									
All ages, age-adjusted ²	3.9	4.1	6.9	7.3	9.1	9.6	10.2	10.2	10.6
All ages, crude	3.9	4.1	6.9	7.4	9.2	9.8	10.3	10.3	10.7
Under 15 years	0.1	0.1	0.2	0.2	0.2	0.2	0.2	0.2	0.2
15–24 years	1.8	1.9	3.3	3.5	4.1	4.6	4.6	4.4	4.8
25–34 years	4.6	4.6	7.2	8.5	10.4	11.9	12.8	13.1	13.0
35–44 years	8.7	9.2	14.8	14.8	16.0	16.8	18.2	17.1	18.0
45–54 years	7.2	7.7	15.0	16.5	21.8	21.8	23.1	23.4	23.6
55–64 years	3.5	3.9	7.0	7.5	11.6	12.9	13.5	14.0	15.9
65–74 years	2.1	2.0	3.0	3.1	4.6	4.8	5.3	5.5	5.9
75–84 years	3.0	2.3	2.9	2.9	3.9	3.3	3.4	3.5	3.4
85 years and over	3.5	3.9	3.7	3.7	3.9	4.5	4.2	3.8	3.5
All ages, age-adjusted ^{2,3}									
Male:									
White	8.1	8.4	12.6	13.6	16.4	16.8	18.1	18.1	19.0
Black or African American	11.5	10.8	11.1	12.8	10.8	10.1	11.0	11.3	12.9
American Indian or Alaska Native	5.7	6.1	11.2	10.8	14.2	11.8	12.9	12.8	12.9
Asian or Pacific Islander	1.5	1.4	2.1	2.2	2.8	2.5	3.2	3.1	3.2
Hispanic or Latino	8.6	7.1	7.5	8.4	8.2	7.6	8.1	8.5	9.2
White, not Hispanic or Latino	8.0	8.6	13.7	14.7	18.3	19.0	20.5	20.4	21.4
Female:									
White	4.0	4.3	7.5	8.0	10.3	10.9	11.7	11.6	12.1
Black or African American	3.9	4.1	5.5	6.0	5.6	5.7	5.9	6.0	6.3
American Indian or Alaska Native	4.6	3.7	7.9	8.6	9.6	9.7	10.7	12.2	11.6
Asian or Pacific Islander	1.0	0.8	1.1	1.3	1.3	1.5	1.6	1.4	1.5
Hispanic or Latina	2.2	2.0	2.9	3.0	3.5	3.6	4.0	4.0	4.1
White, not Hispanic or Latina	4.3	4.5	8.3	8.8	11.6	12.5	13.3	13.2	13.8

See footnotes at end of table.

Table 30 (page 2 of 3). Death rates for drug poisoning and drug poisoning involving opioid analgesics, by sex, age, race, and Hispanic origin: United States, selected years 1999–2013

Updated data when available, Excel, and PDF: <http://www.cdc.gov/nchs/hus/contents2014.htm#030>.

[Data are based on death certificates]

Sex, age, race, and Hispanic origin	1999	2000	2004	2005	2009	2010	2011	2012	2013
Drug poisoning deaths involving opioid analgesics per 100,000 resident population ⁴									
All persons									
All ages, age-adjusted ²	1.4	1.5	3.4	3.7	5.0	5.4	5.4	5.1	5.1
All ages, crude	1.4	1.6	3.4	3.7	5.1	5.4	5.4	5.1	5.1
Under 15 years	*	0.0	0.1	0.1	0.1	0.1	0.1	0.1	0.1
15–24 years	0.7	0.8	2.7	2.7	3.6	3.9	3.6	2.8	2.6
25–34 years	1.9	1.9	4.4	5.3	7.6	8.5	8.5	7.7	7.5
35–44 years	3.5	3.7	6.8	6.9	8.6	9.1	9.3	8.8	8.6
45–54 years	2.9	3.2	7.1	7.9	10.6	10.9	11.2	10.6	10.6
55–64 years	1.0	1.1	2.6	3.1	5.8	6.2	6.3	6.6	7.5
65–74 years	0.4	0.4	0.8	1.0	1.7	1.5	1.8	2.0	2.3
75–84 years	0.3	0.2	0.5	0.6	0.8	0.7	0.7	0.9	0.8
85 years and over	*	*	0.5	0.9	0.7	1.1	0.8	0.8	0.9
Male									
All ages, age-adjusted ²	2.0	2.0	4.2	4.6	6.2	6.5	6.5	6.0	5.9
All ages, crude	2.0	2.1	4.2	4.6	6.2	6.6	6.5	6.0	5.9
Under 15 years	*	*	0.1	0.1	0.1	0.2	0.1	0.1	0.1
15–24 years	1.0	1.2	4.2	4.2	5.3	5.6	5.3	4.2	3.9
25–34 years	2.7	2.7	6.1	7.2	10.6	11.7	11.4	10.0	10.0
35–44 years	5.0	4.9	8.2	8.3	10.4	10.9	10.9	10.3	9.6
45–54 years	3.9	4.3	8.3	9.4	11.6	12.0	12.1	11.1	11.1
55–64 years	1.1	1.0	2.8	3.5	6.3	7.0	6.9	7.3	8.0
65–74 years	0.5	0.3	0.7	0.7	1.6	1.2	1.7	2.0	2.2
75–84 years	*	*	0.4	0.6	0.6	0.7	0.7	0.7	0.9
85 years and over	*	*	*	*	1.2	1.3	*	1.0	1.3
Female									
All ages, age-adjusted ²	0.9	1.1	2.5	2.8	3.9	4.2	4.3	4.2	4.3
All ages, crude	0.9	1.1	2.5	2.8	4.0	4.2	4.4	4.2	4.4
Under 15 years	*	*	0.1	*	0.1	0.1	0.1	0.1	0.1
15–24 years	0.3	0.4	1.1	1.2	1.7	2.1	1.9	1.5	1.4
25–34 years	1.1	1.2	2.8	3.4	4.7	5.3	5.5	5.3	5.0
35–44 years	2.1	2.5	5.4	5.6	6.9	7.3	7.8	7.3	7.6
45–54 years	1.9	2.2	5.9	6.5	9.7	9.8	10.2	10.1	10.1
55–64 years	0.8	1.1	2.4	2.8	5.2	5.5	5.7	6.0	6.9
65–74 years	0.3	0.4	0.9	1.2	1.7	1.7	1.8	2.0	2.4
75–84 years	0.4	*	0.6	0.6	0.9	0.7	0.7	0.9	0.7
85 years and over	*	*	*	0.8	*	1.1	0.8	0.7	0.8
All ages, age-adjusted ^{2,3}									
Male:									
White	2.2	2.3	4.8	5.3	7.2	7.7	7.6	7.0	6.8
Black or African American	1.2	1.2	1.8	2.1	2.4	2.2	2.4	2.3	2.7
American Indian or Alaska Native	*	1.9	4.5	4.4	7.5	5.3	5.5	5.8	4.8
Asian or Pacific Islander	*	*	0.4	0.5	0.7	0.8	1.0	0.7	0.9
Hispanic or Latino	2.9	1.7	2.1	2.2	2.6	2.4	2.6	2.5	2.7
White, not Hispanic or Latino	2.1	2.3	5.3	5.9	8.2	9.0	8.8	8.1	7.9
Female:									
White	1.0	1.2	2.9	3.2	4.5	4.8	5.1	4.9	5.0
Black or African American	0.6	0.6	1.2	1.4	1.8	2.0	2.0	2.0	2.2
American Indian or Alaska Native	*	*	2.7	3.8	4.7	4.9	4.6	5.4	5.4
Asian or Pacific Islander	*	*	*	0.4	0.4	0.5	0.4	0.4	0.3
Hispanic or Latina	0.5	0.5	1.0	1.0	1.3	1.3	1.4	1.5	1.5
White, not Hispanic or Latina	1.1	1.3	3.2	3.5	5.2	5.6	5.8	5.6	5.8

See footnotes at end of table.

Table 30 (page 3 of 3). Death rates for drug poisoning and drug poisoning involving opioid analgesics, by sex, age, race, and Hispanic origin: United States, selected years 1999–2013

Updated data when available, Excel, and PDF: <http://www.cdc.gov/nchs/hus/contents2014.htm#030>.

[Data are based on death certificates]

* Rates based on fewer than 20 deaths are considered unreliable and are not shown.

0.0 Rate more than zero but less than 0.05.

¹Drug poisoning was coded using underlying cause of death according to the 10th Revision of the *International Classification of Diseases* (ICD–10). See Appendix II, Cause of death; Table IV. Drug poisoning deaths include those resulting from accidental or intentional overdoses of a drug, being given the wrong drug, taking the wrong drug in error, taking a drug inadvertently, or other misuses of drugs. These deaths are from all manners and intents, including unintentional, suicide, homicide, undetermined intent, legal intervention, and operations of war.

²Age-adjusted rates are calculated using the year 2000 standard population with unrounded population numbers. See Appendix II, Age adjustment.

³The race groups, white, black, Asian or Pacific Islander, and American Indian or Alaska Native, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Death rates for Hispanic, American Indian or Alaska Native, and Asian or Pacific Islander persons should be interpreted with caution because of inconsistencies in reporting Hispanic origin or race on the death certificate (death rate numerators) compared with population figures (death rate denominators). The net effect of misclassification is an underestimation of deaths and death rates for races other than white and black. See Appendix II, Race, for a detailed discussion of sources of bias in death rates by race and Hispanic origin.

⁴Opioid analgesics include pharmaceutical opioids such as hydrocodone, codeine, and methadone, and synthetic narcotics such as fentanyl and propoxyphene. Drug poisoning deaths involving opioid analgesics include those with an underlying cause of drug poisoning and with opioid analgesics mentioned in the ICD–10 multiple causes of death. See NCHS Health E-Stat for more information on this topic, available from: http://www.cdc.gov/nchs/data/hestat/drug_poisoning/drug_poisoning.htm. See Appendix I, National Vital Statistics System (NVSS), Mortality Multiple Cause-of-Death File, for information about tabulating cause-of-death data in this table. These deaths include all manners and intents. See Appendix II, Cause of death; Table IV. In 1999–2013, 21%–25% of drug poisoning deaths did not include specific information on the death certificate on the type of drug that was involved.

NOTES: Rates for 1999 were computed using intercensal population estimates based on the 1990 and 2000 censuses. Rates for 2000 were computed based on 2000 bridged-race April 1 census counts. Starting with *Health, United States, 2012*, rates for 2001–2009 were revised using intercensal population estimates based on the 2000 and 2010 censuses. Rates for 2010 were based on 2010 bridged-race April 1 census counts. Rates for 2011 and beyond were computed using 2010-based postcensal estimates. See Appendix I, Population Census and Population Estimates. Age groups were selected to minimize the presentation of unstable age-specific death rates based on small numbers of deaths and for consistency among comparison groups. For additional injury-related statistics, see the Web-based Injury Statistics Query and Reporting System, available from: <http://www.cdc.gov/injury/wisqars/index.html>. Starting with 2003 data, some states allowed the reporting of more than one race on the death certificate. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards, for comparability with other states. See Appendix II, Race. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: CDC/NCHS, National Vital Statistics System; numerator data from National Vital Statistics System, annual public-use Mortality Files; denominator data from national population estimates for race groups from Table 1; Xu JQ, Murphy SL, Kochanek KD, et al. Deaths: Final data for 2013. National vital statistics reports; vol 64 no 2. Hyattsville, MD: NCHS; 2015. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr64/nvsr64_02.pdf. See Appendix I, National Vital Statistics System (NVSS).