## Table 76 (page 1 of 3). Use of mammography among women aged 40 and over, by selected characteristics: United States, selected years 1987–2013

Excel, PDF, more data years, and standard errors: http://www.cdc.gov/nchs/hus/contents2014.htm#076.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	1987	1993	1994	2000	2003	2005	2008	2010	2013
		Perce	ent of wom	en having a	a mammogr	am within th	ne past 2 y	ears <sup>1</sup>	
40 years and over, age-adjusted <sup>2,3</sup> 40 years and over, crude <sup>2</sup>	29.0 28.7	59.7 59.7	61.0 60.9	70.4 70.4	69.5 69.7	66.6 66.8	67.1 67.6	66.5 67.1	65.7 66.8
50 years and over, age-adjusted <sup>2,3</sup> 50 years and over, crude <sup>2</sup>	27.3 27.4	59.7 59.7	60.9 60.6	73.7 73.6	72.4 72.4	68.2 68.4	70.3 70.5	68.8 69.2	69.1 69.5
Age 40–49 years	31.9 31.7 22.8 26.6 17.3	59.9 65.1 54.2 64.2 41.0	61.3 66.5 55.0 63.0 44.6	64.3 78.7 67.9 74.0 61.3	64.4 76.2 67.7 74.6 60.6	63.5 71.8 63.8 72.5 54.7	61.5 74.2 65.5 72.6 57.9	62.3 72.6 64.4 71.9 55.7	59.6 71.4 66.9 75.3 56.5
Race <sup>4</sup>									
40 years and over, crude:     White only.     Black or African American only.     American Indian or Alaska Native only.     Asian only.     Native Hawaiian or Other Pacific	29.6 24.0 *	60.0 59.1 49.8 55.1	60.6 64.3 65.8 55.8	71.4 67.8 47.4 53.5	70.1 70.4 63.1 57.6	67.4 64.9 72.8 54.6	67.9 68.0 62.7 66.1	67.4 67.9 71.2 62.4	66.8 67.1 62.6 66.6
Islander only 2 or more races				* 69.2	* 65.3	* 63.7	* 55.2	* 51.4	* 65.4
Hispanic origin and race <sup>4</sup>									
40 years and over, crude: Hispanic or Latina Not Hispanic or Latina White only Black or African American only	18.3 29.4 30.3 23.8	50.9 60.3 60.6 59.2	51.9 61.5 61.3 64.4	61.2 71.1 72.2 67.9	65.0 70.1 70.5 70.5	58.8 67.5 68.3 65.2	61.2 68.3 68.7 68.3	64.2 67.4 67.8 67.4	61.4 67.5 67.6 67.2
Age, Hispanic origin, and race <sup>4</sup>									
40–49 years: Hispanic or Latina	*15.3	52.6	47.5	54.1	59.4	54.2	54.1	59.8	56.4
White only	34.3 27.8	61.6 55.6	62.0 67.2	67.2 60.9	65.2 68.2	65.5 62.1	64.1 59.5	62.6 63.5	60.3 59.4
i0–64 years: Hispanic or Latina	23.0	59.2	60.1	66.5	69.4	61.5	71.3	68.6	65.6
White only Black or African American only	33.6 26.4	66.2 65.5	67.5 63.6	80.6 77.7	77.2 76.2	73.5 71.6	74.1 76.7	73.5 74.0	72.1 71.7
5 years and over: Hispanic or Latina	*	*35.7	48.0	68.3	69.5	63.8	59.0	65.2	63.2
White only	24.0 14.1	54.7 56.3	54.9 61.0	68.3 65.5	68.1 65.4	64.7 60.5	66.1 66.4	65.0 60.9	67.3 68.8
Age and percent of poverty level <sup>5</sup>									
40 years and over, crude:       Below 100%       100%-199%       200%-399%       400% or more	14.6 20.9 29.7 42.9	41.1 47.5 63.2 74.1	44.2 48.6 65.0 74.1	54.8 58.1 68.8 81.5	55.4 60.8 69.9 77.7	48.5 55.3 67.2 76.6	51.4 55.8 64.4 79.0	51.4 53.8 66.2 78.1	49.9 56.7 66.0 77.2
40–49 years: Below 100% 100%–199% 200%–399% 400% or more	18.6 18.4 31.2 44.1	36.1 47.8 63.0 69.6	43.0 47.6 64.5 69.9	47.4 43.6 60.2 75.8	50.6 54.0 63.0 71.6	42.5 49.8 61.8 73.6	46.6 46.5 56.8 72.5	48.1 46.2 59.2 73.6	43.3 52.0 58.5 69.0
50–64 years: Below 100% 100%–199% 200%–399% 400% or more	14.6 24.2 29.7 44.7	47.3 47.0 66.1 78.7	46.2 49.0 69.6 78.0	61.7 68.3 75.1 86.9	58.3 64.0 74.1 84.9	50.4 58.8 70.7 80.6	57.5 58.9 69.8 84.3	54.7 57.3 70.7 82.8	55.0 57.2 69.5 80.9
65 years and over: Below 100% 100%–199% 200%–399% 400% or more	13.1 19.9 27.7 34.7	40.4 47.6 60.3 71.3	43.9 48.8 61.0 73.0	54.8 60.3 71.1 81.9	57.0 62.8 72.3 73.0	52.3 56.1 68.6 72.6	49.1 59.4 65.0 78.3	50.6 55.5 67.2 74.5	49.8 59.3 68.1 79.0

See footnotes at end of table.

## Table 76 (page 2 of 3). Use of mammography among women aged 40 and over, by selected characteristics: United States, selected years 1987–2013

## Excel, PDF, more data years, and standard errors: http://www.cdc.gov/nchs/hus/contents2014.htm#076.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	1987	1993	1994	2000	2003	2005	2008	2010	2013		
Health insurance status at the time of interview <sup>6</sup>	Percent of women having a mammogram within the past 2 years <sup>1</sup>										
40–64 years: Insured Private Medicaid Uninsured		66.2 67.1 51.9 36.0	68.3 69.4 54.5 34.0	76.0 77.1 61.7 40.7	75.1 76.3 63.5 41.5	72.5 74.5 55.6 38.1	73.4 74.2 64.2 39.7	74.1 75.6 64.4 36.0	72.1 73.4 63.5 37.3		
Health insurance status prior to interview <sup>6</sup>											
40–64 years: Insured continuously all 12 months Uninsured for any period up to 12 months Uninsured more than 12 months	 	66.6 49.4 28.4	68.6 49.9 26.6	76.8 53.0 34.0	75.6 56.0 37.0	73.1 51.3 32.9	74.1 55.3 34.6	74.7 57.3 30.0	72.7 54.5 32.8		
Age and education <sup>7</sup>											
40 years and over, crude: No high school diploma or GED High school diploma or GED	17.8 31.3 37.7	46.4 59.0 69.5	48.2 61.0 69.7	57.7 69.7 76.2	58.1 67.8 75.1	52.8 64.9 72.7	53.8 65.2 73.4	53.0 64.4 72.1	53.6 63.4 71.6		
40–49 years: No high school diploma or GED High school diploma or GED Some college or more	15.1 32.6 39.2	43.6 56.6 66.1	50.4 55.8 68.7	46.8 59.0 70.6	53.3 60.8 68.1	51.2 58.8 68.3	46.9 57.2 66.3	44.9 58.4 66.5	46.9 51.8 64.3		
50–64 years: No high school diploma or GED High school diploma or GED Some college or more	21.2 33.8 40.5	51.4 62.4 78.5	51.6 67.8 74.7	66.5 76.6 84.2	63.4 71.8 82.7	56.9 70.1 77.0	64.9 70.4 78.5	56.7 69.9 77.0	58.2 66.9 75.7		
65 years and over: No high school diploma or GED High school diploma or GED Some college or more	16.5 25.9 32.3	44.2 57.4 64.8	45.6 59.1 64.3	57.4 71.8 74.1	56.9 69.7 75.1	50.7 64.3 73.0	49.2 65.7 75.6	54.1 62.5 70.9	53.4 66.5 73.6		
Disability measure <sup>8</sup>											
40 years and over, crude: Any basic actions difficulty or complex activity limitation Any basic actions difficulty Any complex activity limitation No disability	  			67.8 67.9 64.1 72.6	67.2 67.3 62.3 71.8	63.5 63.5 59.9 69.8	63.9 63.9 60.2 71.1	63.3 63.3 58.2 70.8	63.5 63.8 58.4 69.8		

See footnotes at end of table.

## Table 76 (page 3 of 3). Use of mammography among women aged 40 and over, by selected characteristics: United States, selected years 1987–2013

Excel, PDF, more data years, and standard errors: http://www.cdc.gov/nchs/hus/contents2014.htm#076.

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

<sup>4</sup>The race groups, white, black, American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, and 2 or more races, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. The five single-race categories plus multiple-race categories shown in the table conform to the 1997 Standards. Starting with 1999 data, race-specific estimates are tabulated according to the 1977 Standards with four racial group; the category 2 or more races includes persons who reported more than one racial group. Prior to 1999, data were tabulated according to the 1977 Standards with four racial groups, and the Asian only category included Native Hawaiian or Other Pacific Islander. Estimates for single-race categories prior to 1999 included persons who reported one race as best representing their race. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See Appendix II. Hispanic origin: Race.

<sup>5</sup>Percent of poverty level is based on family income and family size and composition using U.S. Census Bureau poverty thresholds. Poverty level was unknown for 11% of women aged 40 and over in 1987. Missing family income data were imputed for 1997 and beyond. See Appendix II, Family income; Poverty; Table VI.

<sup>6</sup>Health insurance categories are mutually exclusive. Persons who reported both Medicaid and private coverage are classified as having private coverage. Starting with 1997 data, state-sponsored health plan coverage is included as Medicaid coverage. Starting with 1999 data, coverage by the Children's Health Insurance Program (CHIP) is included with Medicaid coverage. In addition to private and Medicaid, the insured category also includes military plans, other government-sponsored health plans, and Medicaire, not shown separately. Persons not covered by private insurance, Medicaid, CHIP, public assistance (through 1996), state-sponsored or other government-sponsored health plans, and Medicaid, in 1997). Medicare, or military plans are considered to have no health insurance coverage. Persons with only Indian Health Service coverage are considered to have no health insurance coverage. See Appendix II, Health insurance coverage.

<sup>7</sup>Education categories shown are for 1998 and subsequent years. GED is General Educational Development high school equivalency diploma. In years prior to 1998, the following categories based on number of years of school completed were used: less than 12 years, 12 years, 13 years or more. See Appendix II, Education. <sup>8</sup>Any basic actions difficulty or complex activity limitation is defined as having one or more of the following limitations or difficulties: movement difficulty, emotional difficulty, sensory (seeing or hearing) difficulty, cognitive difficulty; complex activity limitation. For more information, see Appendix II, Basic actions difficulty; Complex activity limitation. Starting with 2007 data, the hearing question, a component of the basic actions difficulty, see Appendix II, Hearing trouble.

NOTES: See Appendix II, Mammography, for a discussion of the U.S. Preventive Services Task Force recommendations for mammography screening. Standard errors are available in the spreadsheet version of this table. Available from: http://www.cdc.gov/nchs/hus.htm. Data starting in 1997 are not strictly comparable with data for earlier years due to the 1997 questionnaire redesign. See Appendix I, National Health Interview Survey (NHIS). Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: http://www.cdc.gov/nchs/hus.htm.

SOURCE: CDC/NCHS, National Health Interview Survey. Data are from the following supplements: cancer control (1987), health promotion and disease prevention (1990–1991), and year 2000 objectives (1993–1994). Starting in 1998, data are from the family core and sample adult questionnaires. See Appendix I, National Health Interview Survey (NHIS).

<sup>\*</sup> Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) of 20%-30%. Data not shown have an RSE greater than 30%. - - - Data not available.

<sup>&</sup>lt;sup>1</sup>Questions concerning use of mammography differed slightly on the National Health Interview Survey across the years for which data are shown. See Appendix II, Mammography.

<sup>&</sup>lt;sup>2</sup>Includes all other races not shown separately, unknown poverty level in 1987, unknown health insurance status, unknown education level, and unknown disability status. <sup>3</sup>Estimates for women aged 40 and over are age-adjusted to the year 2000 standard population using four age groups: 40–49 years, 50–64 years, 65–74 years, and 75 years and over. Estimates for women 50 years of age and over are age-adjusted using three age groups. See Appendix II, Age adjustment.