

Data table for Figure 27. Use of preventive services and screening among noninstitutionalized adults aged 55–64: United States, 2003 and 2013

Excel and Powerpoint: <http://www.cdc.gov/nchs/hus/contents2014.htm#fig27>

Health care utilization	2003		2013	
	Percent	SE	Percent	SE
Influenza vaccination in past 12 months ¹	40.4	0.9	49.9	0.9
Pneumococcal vaccination ever (high-risk group) ²	32.5	1.4	35.2	1.2
Colorectal test or procedure ³	40.0	0.9	58.3	0.9
Mammogram in past 2 years ⁴	76.9	1.0	71.3	1.0
Pap smear in past 3 years ⁵	85.5	1.1	75.9	1.2

SE is standard error.

¹Respondents were asked about influenza vaccination in the past 12 months. See Appendix II, Vaccination.

²Respondents were asked, “Have you ever had a pneumonia shot? This shot is usually given only once or twice in a person’s lifetime and is different from the flu shot. It is also called the pneumococcal vaccine.” High-risk group membership is based on recommendations of CDC’s Advisory Committee on Immunization Practices (ACIP). The high-risk group includes persons who reported diabetes; cancer; heart, lung, liver, or kidney disease. Starting in 2009, this definition was expanded to include persons who reported asthma or cigarette smoking, to be consistent with the revised ACIP recommendation. Starting with data year 2012, the survey questionnaire was changed to ask respondents a new question on chronic obstructive pulmonary disease (COPD), and this information was added to the list of lung diseases used to construct the high-risk category. For more information on high-risk groups, see the 2009 ACIP recommendation available from: <http://www.cdc.gov/mmwr/pdf/wk/mm5934.pdf>.

³Colorectal test or procedure is defined as fecal occult blood test (FOBT) in the past year, sigmoidoscopy in the past 5 years with FOBT in the past 3 years, or colonoscopy in the past 10 years. See Appendix II, Colorectal tests or procedures.

⁴Questions concerning use of mammography differed slightly on the National Health Interview Survey across the years for which data are shown. See Appendix II, Mammography.

⁵Estimates are for women who have not had a hysterectomy. Questions concerning use of Pap smears and hysterectomy status differed slightly on the National Health Interview Survey across the years for which data are shown. Hysterectomy status was not assessed in 2003; therefore, data shown for Pap smear were for 2000 and 2013 when hysterectomy status was available. See Appendix II, Pap smear.

NOTE: Utilization of colorectal, cervical, and breast cancer procedures can occur for routine screening or for diagnostic reasons.

SOURCE: CDC/NCHS, National Health Interview Survey. Family core and sample adult questionnaires. See Appendix I, National Health Interview Survey (NHIS).