

**Table 36 (page 1 of 4). Age-adjusted cancer incidence rates for selected cancer sites, by sex, race, and Hispanic origin: United States, selected geographic areas, selected years 1990–2012**

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#036>.

[Data are based on the Surveillance, Epidemiology, and End Results (SEER) Program's 13 population-based cancer registries]

Site, sex, race, and Hispanic origin	1990	1995	2000	2002	2003	2005	2010	2011	2012	1990–2012 APC <sup>1</sup>
All sites										
Number of new cases per 100,000 population <sup>2</sup>										
All persons . . . . .	475.8	471.3	475.0	474.2	463.2	460.6	449.6	441.7	426.1	†–0.5
White . . . . .	483.2	477.4	485.6	484.5	473.8	472.7	460.5	452.8	435.6	†–0.4
Black or African American . . . . .	515.1	539.1	525.7	524.1	513.9	504.7	482.0	474.3	459.2	†–0.7
American Indian or Alaska Native <sup>3</sup> . . . . .	348.2	370.8	364.8	356.9	379.9	403.5	403.4	375.4	353.6	0.2
Asian or Pacific Islander . . . . .	335.3	340.5	341.5	346.8	334.0	331.7	323.0	321.3	308.5	†–0.4
Hispanic or Latino <sup>4</sup> . . . . .	355.9	359.8	361.5	371.9	357.9	364.8	347.6	349.8	334.2	†–0.3
White, not Hispanic or Latino <sup>4</sup> . . . . .	495.2	491.2	503.5	502.0	492.1	490.9	481.2	472.4	455.2	†–0.3
Male . . . . .	584.1	564.7	565.4	558.3	545.7	534.6	511.1	496.4	464.2	†–1.0
White . . . . .	591.1	563.4	569.4	563.8	550.5	543.1	517.5	503.7	469.8	†–0.9
Black or African American . . . . .	688.3	742.4	707.1	688.5	668.9	636.4	593.5	567.7	540.8	†–1.4
American Indian or Alaska Native <sup>3</sup> . . . . .	396.3	424.0	373.1	381.5	447.2	425.2	451.2	401.6	344.7	–0.3
Asian or Pacific Islander . . . . .	385.9	398.9	400.5	389.9	387.0	371.5	346.1	343.1	314.7	†–1.0
Hispanic or Latino <sup>4</sup> . . . . .	417.4	440.6	435.7	443.5	423.7	426.5	396.9	390.5	362.2	†–0.7
White, not Hispanic or Latino <sup>4</sup> . . . . .	606.7	577.3	588.6	581.4	568.9	561.3	538.4	523.8	489.9	†–0.8
Female . . . . .	411.5	410.6	414.0	417.1	406.9	409.6	405.9	403.7	401.3	–0.1
White . . . . .	421.2	423.2	430.0	431.4	422.1	424.6	420.3	417.7	414.3	0.0
Black or African American . . . . .	406.5	403.1	403.8	414.6	409.8	415.9	404.7	409.9	403.2	0.0
American Indian or Alaska Native <sup>3</sup> . . . . .	315.6	337.0	365.5	337.7	335.5	389.0	376.2	361.3	367.7	†0.7
Asian or Pacific Islander . . . . .	295.7	297.7	300.8	319.9	299.4	306.7	311.0	309.8	308.9	†0.2
Hispanic or Latina <sup>4</sup> . . . . .	322.6	310.5	317.6	328.0	317.4	326.3	317.5	326.1	320.1	0.1
White, not Hispanic or Latina <sup>4</sup> . . . . .	430.6	436.7	446.0	448.1	439.7	442.1	439.9	436.3	433.2	0.1
Lung and bronchus										
Male . . . . .	95.0	86.9	77.8	75.7	75.5	71.5	62.9	60.7	58.3	†–2.1
White . . . . .	94.1	85.0	76.3	74.9	74.3	70.8	62.7	60.1	57.4	†–2.0
Black or African American . . . . .	134.2	137.0	111.7	109.1	112.1	98.0	82.3	82.3	80.6	†–2.5
American Indian or Alaska Native <sup>3</sup> . . . . .	77.3	82.8	62.6	46.2	74.2	67.4	65.3	55.3	44.1	†–1.2
Asian or Pacific Islander . . . . .	64.4	60.1	63.9	57.9	58.7	58.2	50.9	49.9	48.3	†–1.2
Hispanic or Latino <sup>4</sup> . . . . .	59.4	52.4	45.2	49.0	46.5	44.0	34.3	35.6	33.6	†–2.1
White, not Hispanic or Latino <sup>4</sup> . . . . .	97.4	88.4	80.2	78.4	77.9	74.6	67.0	64.0	61.4	†–1.9
Female . . . . .	47.2	49.3	48.6	49.4	49.8	50.0	46.0	44.8	43.9	†–0.3
White . . . . .	48.5	51.7	50.7	51.5	52.4	51.9	48.0	46.5	45.5	–0.2
Black or African American . . . . .	52.9	50.0	55.1	55.4	55.4	58.4	52.8	49.8	51.1	0.0
American Indian or Alaska Native <sup>3</sup> . . . . .	30.4	46.1	38.5	39.9	41.4	45.5	36.6	42.1	34.4	0.9
Asian or Pacific Islander . . . . .	28.4	27.6	27.3	29.7	29.1	30.8	29.1	31.3	28.8	†0.3
Hispanic or Latina <sup>4</sup> . . . . .	26.2	25.1	24.3	24.9	24.4	22.9	24.6	23.9	22.6	†–0.5
White, not Hispanic or Latina <sup>4</sup> . . . . .	50.8	54.8	54.4	55.4	56.6	56.4	51.9	50.6	49.7	0.0
Colon and rectum										
Male . . . . .	72.3	63.2	62.6	60.1	58.4	54.7	46.9	45.4	43.5	†–2.1
White . . . . .	73.0	62.4	62.0	58.9	57.0	54.1	45.4	43.4	41.9	†–2.2
Black or African American . . . . .	73.1	75.1	73.6	72.5	76.3	67.0	57.6	57.0	55.5	†–1.3
American Indian or Alaska Native <sup>3</sup> . . . . .	61.9	66.2	48.7	49.4	70.4	65.6	64.6	64.4	57.8	–0.1
Asian or Pacific Islander . . . . .	60.9	58.8	58.0	59.0	53.3	47.7	45.5	45.7	42.1	†–1.6
Hispanic or Latino <sup>4</sup> . . . . .	47.6	46.0	50.5	46.0	47.4	47.3	42.0	43.0	40.0	†–0.7
White, not Hispanic or Latino <sup>4</sup> . . . . .	75.0	63.9	63.4	60.3	58.1	54.8	46.0	43.5	42.2	†–2.3
Female . . . . .	50.2	45.9	46.1	45.2	43.5	41.4	36.0	34.7	33.4	†–1.5
White . . . . .	49.7	45.4	45.6	44.1	42.9	40.3	34.5	33.7	32.7	†–1.6
Black or African American . . . . .	61.3	54.9	58.1	56.3	55.4	54.1	46.3	42.2	41.0	†–1.2
American Indian or Alaska Native <sup>3</sup> . . . . .	45.8	47.9	39.1	50.9	45.1	48.3	41.9	46.0	45.1	–0.3
Asian or Pacific Islander . . . . .	37.8	38.8	37.5	41.8	36.4	36.9	34.3	31.8	28.7	†–1.2
Hispanic or Latina <sup>4</sup> . . . . .	34.5	32.0	34.1	32.1	33.5	33.6	30.7	29.0	28.1	†–0.6
White, not Hispanic or Latina <sup>4</sup> . . . . .	50.9	46.7	46.7	45.5	44.0	41.3	34.9	34.5	33.5	†–1.6
Prostate										
Male . . . . .	166.9	166.6	178.9	177.9	165.4	154.0	143.1	136.7	110.4	†–1.8
White . . . . .	168.5	161.3	174.8	174.5	161.6	149.9	138.2	131.9	104.0	†–1.9
Black or African American . . . . .	219.4	279.2	291.4	280.3	253.2	242.9	218.3	206.3	178.1	†–1.8
American Indian or Alaska Native <sup>3</sup> . . . . .	99.5	93.5	69.6	93.7	112.4	93.8	83.7	67.1	59.5	†–1.9
Asian or Pacific Islander . . . . .	88.5	104.6	108.2	104.1	104.9	95.7	78.7	79.4	63.3	†–1.8
Hispanic or Latino <sup>4</sup> . . . . .	119.2	141.2	150.2	152.8	139.3	132.4	117.9	109.6	93.5	†–1.5
White, not Hispanic or Latino <sup>4</sup> . . . . .	172.2	163.6	178.6	177.6	165.1	152.5	142.2	136.3	106.7	†–1.9

See footnotes at end of table.

**Table 36 (page 2 of 4). Age-adjusted cancer incidence rates for selected cancer sites, by sex, race, and Hispanic origin: United States, selected geographic areas, selected years 1990–2012**

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#036>.

[Data are based on the Surveillance, Epidemiology, and End Results (SEER) Program's 13 population-based cancer registries]

Site, sex, race, and Hispanic origin	1990	1995	2000	2002	2003	2005	2010	2011	2012	1990–2012 APC <sup>1</sup>
Breast										
Number of new cases per 100,000 population <sup>2</sup>										
Female . . . . .	129.4	130.9	134.2	132.8	124.1	124.4	123.3	126.6	125.9	†–0.3
White . . . . .	134.2	136.3	140.6	138.6	129.0	129.5	127.2	130.3	128.9	†–0.3
Black or African American . . . . .	117.4	123.0	122.6	124.2	123.3	119.3	123.2	128.9	130.0	†0.3
American Indian or Alaska Native <sup>3</sup> . . . . .	69.6	96.0	100.2	81.3	93.5	105.6	91.3	105.2	96.9	0.4
Asian or Pacific Islander . . . . .	88.1	88.4	94.7	101.0	91.9	95.6	100.3	102.7	103.9	†0.7
Hispanic or Latina <sup>4</sup> . . . . .	90.8	89.4	97.0	93.4	87.8	93.1	86.9	95.7	91.8	0.1
White, not Hispanic or Latina <sup>4</sup> . . . . .	138.6	141.8	147.0	145.8	135.7	136.2	135.4	137.2	136.5	–0.2
Cervix uteri										
Female . . . . .	11.9	9.9	8.9	8.4	8.2	7.9	7.2	7.0	6.8	†–2.4
White . . . . .	11.2	9.2	8.9	8.3	7.9	7.8	7.2	7.2	6.7	†–2.1
Black or African American . . . . .	16.6	14.9	10.7	10.1	10.8	9.3	8.5	7.8	7.6	†–3.6
American Indian or Alaska Native <sup>3</sup> . . . . .	14.7	*	*	*	*	10.7	8.3	9.5	*	†–2.0
Asian or Pacific Islander . . . . .	12.1	11.2	8.0	8.2	8.1	7.8	6.6	5.5	6.5	†–3.8
Hispanic or Latina <sup>4</sup> . . . . .	21.3	17.4	17.1	14.5	14.0	13.7	10.4	9.3	9.2	†–3.8
White, not Hispanic or Latina <sup>4</sup> . . . . .	9.7	7.8	7.1	7.0	6.5	6.4	6.2	6.5	5.9	†–1.9
Corpus and uterus, not otherwise specified										
Female . . . . .	24.7	24.9	23.9	24.0	23.6	24.1	26.7	26.5	27.0	†0.4
White . . . . .	26.4	26.4	25.6	24.8	25.0	25.4	27.6	27.2	27.9	0.2
Black or African American . . . . .	17.0	17.9	17.4	22.4	20.3	21.5	24.6	27.7	25.0	†2.2
American Indian or Alaska Native <sup>3</sup> . . . . .	19.3	*	18.5	19.2	19.5	14.4	27.6	19.5	19.3	*
Asian or Pacific Islander . . . . .	13.5	17.8	16.6	18.8	16.6	18.9	22.5	21.8	22.3	†1.8
Hispanic or Latina <sup>4</sup> . . . . .	17.7	16.5	15.8	17.2	17.8	19.3	20.5	22.1	23.7	†1.5
White, not Hispanic or Latina <sup>4</sup> . . . . .	27.1	27.5	26.8	25.8	26.0	26.2	28.6	27.7	28.3	0.1
Ovary										
Female . . . . .	15.6	14.6	14.2	13.9	13.6	13.2	12.5	12.1	12.2	†–1.0
White . . . . .	16.4	15.4	15.0	14.7	14.3	13.8	13.4	12.9	12.9	†–1.0
Black or African American . . . . .	11.3	10.8	10.9	9.9	11.6	10.8	9.4	9.9	10.4	–0.4
American Indian or Alaska Native <sup>3</sup> . . . . .	21.9	*	18.9	*	13.4	12.2	11.9	10.8	15.7	*
Asian or Pacific Islander . . . . .	11.1	10.4	10.4	12.2	10.2	10.9	9.7	9.6	9.5	†–0.5
Hispanic or Latina <sup>4</sup> . . . . .	12.2	11.7	10.6	13.8	11.6	11.7	11.8	11.0	12.1	–0.3
White, not Hispanic or Latina <sup>4</sup> . . . . .	16.8	15.9	15.6	14.7	14.7	14.1	13.4	13.1	12.9	†–1.1
Oral cavity and pharynx										
Male . . . . .	18.5	16.5	15.8	15.7	15.2	15.1	15.5	15.9	15.4	†–0.7
White . . . . .	18.0	16.4	15.6	15.9	15.2	15.5	16.0	16.8	16.2	†–0.4
Black or African American . . . . .	25.4	22.3	19.4	18.1	17.5	15.8	13.8	13.8	14.0	†–2.8
American Indian or Alaska Native <sup>3</sup> . . . . .	*	*	*	*	16.3	10.4	21.3	15.9	15.3	*
Asian or Pacific Islander . . . . .	14.9	11.8	13.7	12.9	11.9	11.5	11.7	10.9	10.6	†–1.1
Hispanic or Latino <sup>4</sup> . . . . .	10.7	12.3	9.0	9.6	8.8	9.7	8.8	10.9	9.9	–0.6
White, not Hispanic or Latino <sup>4</sup> . . . . .	18.8	16.9	16.6	16.9	16.3	16.4	17.5	18.0	17.4	–0.2
Female . . . . .	7.3	7.0	6.2	6.5	5.9	6.1	6.1	6.1	6.3	†–0.8
White . . . . .	7.4	7.1	6.2	6.5	5.8	6.0	6.3	6.3	6.5	†–0.7
Black or African American . . . . .	6.4	6.7	5.4	6.3	6.7	6.9	5.3	5.2	4.5	†–1.5
American Indian or Alaska Native <sup>3</sup> . . . . .	*	*	*	*	*	*	*	*	7.4	*
Asian or Pacific Islander . . . . .	6.1	5.2	6.3	6.0	5.2	5.9	5.2	4.7	6.0	†–0.8
Hispanic or Latina <sup>4</sup> . . . . .	4.0	3.7	3.7	3.7	3.8	3.5	4.2	3.7	3.8	–0.4
White, not Hispanic or Latina <sup>4</sup> . . . . .	7.8	7.5	6.6	7.0	6.2	6.5	6.7	6.9	7.0	†–0.6
Stomach										
Male . . . . .	14.6	13.5	12.6	12.0	11.8	11.4	10.4	10.5	10.1	†–1.7
White . . . . .	12.8	11.9	10.6	10.4	10.1	9.6	9.3	9.5	8.8	†–1.6
Black or African American . . . . .	21.5	18.6	18.6	15.9	18.7	17.4	13.3	14.0	14.6	†–2.1
American Indian or Alaska Native <sup>3</sup> . . . . .	*	24.1	20.2	25.3	*	20.8	20.7	21.0	16.6	–0.5
Asian or Pacific Islander . . . . .	26.8	24.5	22.9	20.5	19.2	20.2	15.2	14.2	14.8	†–2.9
Hispanic or Latino <sup>4</sup> . . . . .	20.2	19.3	16.1	16.4	16.2	15.5	14.9	14.2	13.1	†–1.7
White, not Hispanic or Latino <sup>4</sup> . . . . .	12.1	11.1	10.0	9.6	9.2	8.6	8.2	8.4	7.8	†–1.9

See footnotes at end of table.

**Table 36 (page 3 of 4). Age-adjusted cancer incidence rates for selected cancer sites, by sex, race, and Hispanic origin: United States, selected geographic areas, selected years 1990–2012**

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#036>.

[Data are based on the Surveillance, Epidemiology, and End Results (SEER) Program's 13 population-based cancer registries]

Site, sex, race, and Hispanic origin	1990	1995	2000	2002	2003	2005	2010	2011	2012	1990–2012 APC <sup>1</sup>
Stomach										
Number of new cases per 100,000 population <sup>2</sup>										
Female . . . . .	6.7	6.2	6.1	6.2	6.0	5.7	5.7	5.6	5.8	†–0.8
White . . . . .	5.7	5.1	5.0	5.1	5.0	4.7	4.7	4.8	5.0	†–0.8
Black or African American . . . . .	10.0	9.8	8.7	9.9	9.5	8.1	8.5	9.4	8.0	†–1.0
American Indian or Alaska Native <sup>3</sup> . . . . .	*	*	*	17.2	*	*	13.0	*	12.2	*
Asian or Pacific Islander . . . . .	15.4	13.0	12.9	11.5	11.1	10.3	8.9	7.8	8.7	†–2.7
Hispanic or Latina <sup>4</sup> . . . . .	10.8	11.2	10.6	10.7	10.2	10.2	9.4	8.8	9.4	†–1.0
White, not Hispanic or Latina <sup>4</sup> . . . . .	5.1	4.5	4.2	4.2	4.1	3.8	3.6	3.9	3.9	†–1.5
Pancreas										
Male . . . . .	13.0	12.7	12.8	12.8	12.5	13.6	13.7	14.2	13.9	†0.5
White . . . . .	12.7	12.4	12.6	13.0	12.3	13.4	13.5	14.2	14.3	†0.7
Black or African American . . . . .	19.3	19.1	18.2	13.9	17.4	18.3	18.5	16.8	15.8	–0.4
American Indian or Alaska Native <sup>3</sup> . . . . .	*	*	*	*	*	21.2	17.3	17.6	*	*
Asian or Pacific Islander . . . . .	11.0	10.3	10.7	9.9	10.2	11.8	11.4	12.3	10.3	0.1
Hispanic or Latino <sup>4</sup> . . . . .	10.7	12.0	12.2	10.9	10.1	12.3	11.5	13.4	10.6	†0.8
White, not Hispanic or Latino <sup>4</sup> . . . . .	12.8	12.4	12.7	13.2	12.6	13.5	13.7	14.3	14.8	†0.7
Female . . . . .	10.0	9.9	9.9	10.5	10.4	10.8	11.2	10.6	11.2	†0.6
White . . . . .	9.7	9.6	9.6	10.1	10.2	10.6	10.9	10.3	10.9	†0.6
Black or African American . . . . .	12.9	15.5	12.8	15.9	14.4	16.5	15.1	15.0	15.0	0.0
American Indian or Alaska Native <sup>3</sup> . . . . .	*	*	20.4	*	*	12.4	13.8	*	11.3	*
Asian or Pacific Islander . . . . .	9.9	8.1	9.2	9.1	8.1	8.0	10.4	9.1	10.1	†1.0
Hispanic or Latina <sup>4</sup> . . . . .	9.9	9.0	9.1	11.1	8.6	11.1	9.9	10.1	10.1	0.1
White, not Hispanic or Latina <sup>4</sup> . . . . .	9.7	9.7	9.6	10.0	10.5	10.6	11.1	10.4	11.0	†0.7
Urinary bladder										
Male . . . . .	37.2	35.4	36.8	35.8	36.9	37.0	35.8	33.8	34.0	†–0.2
White . . . . .	40.7	38.8	40.8	39.3	40.7	40.9	39.8	37.9	37.8	†–0.2
Black or African American . . . . .	19.7	19.7	20.2	20.9	23.2	23.1	22.3	20.7	23.1	†0.5
American Indian or Alaska Native <sup>3</sup> . . . . .	*	*	*	*	*	16.8	16.1	17.5	*	*
Asian or Pacific Islander . . . . .	15.4	17.2	16.9	19.7	18.0	17.5	17.2	14.9	16.3	0.3
Hispanic or Latino <sup>4</sup> . . . . .	22.4	17.8	20.4	21.2	20.2	20.1	18.5	19.8	18.9	–0.3
White, not Hispanic or Latino <sup>4</sup> . . . . .	42.4	40.9	43.2	41.6	43.3	43.6	43.0	40.8	40.7	0.0
Female . . . . .	9.5	9.3	9.1	9.1	9.2	9.0	8.6	8.3	8.3	†–0.5
White . . . . .	10.0	10.1	10.0	10.0	10.0	9.7	9.4	9.1	9.1	†–0.4
Black or African American . . . . .	8.7	7.3	7.8	8.5	7.8	7.9	7.0	7.1	6.2	–0.6
American Indian or Alaska Native <sup>3</sup> . . . . .	*	*	*	*	*	*	*	*	*	*
Asian or Pacific Islander . . . . .	5.3	4.5	4.3	3.4	5.0	5.2	4.7	4.3	4.1	–0.2
Hispanic or Latino <sup>4</sup> . . . . .	5.8	5.3	5.8	6.4	4.6	5.9	4.7	5.6	4.9	–0.4
White, not Hispanic or Latino <sup>4</sup> . . . . .	10.3	10.6	10.5	10.5	10.8	10.3	10.2	9.8	9.9	†–0.2
Non-Hodgkin lymphoma										
Male . . . . .	22.6	25.1	23.6	23.8	24.2	24.7	25.4	23.7	23.4	0.2
White . . . . .	23.6	26.2	24.9	25.1	25.7	25.9	26.6	24.9	24.6	†0.2
Black or African American . . . . .	17.7	21.7	17.6	18.1	19.4	19.7	21.3	17.5	17.1	–0.1
American Indian or Alaska Native <sup>3</sup> . . . . .	*	*	15.3	16.6	*	23.2	18.7	10.8	13.6	*
Asian or Pacific Islander . . . . .	16.7	16.6	16.4	16.5	16.3	17.9	17.7	17.9	17.5	0.3
Hispanic or Latino <sup>4</sup> . . . . .	17.3	21.0	20.5	20.6	19.6	20.0	22.9	19.9	19.2	0.4
White, not Hispanic or Latino <sup>4</sup> . . . . .	24.3	26.7	25.5	25.7	26.5	26.9	27.2	25.6	25.7	†0.3
Female . . . . .	14.5	15.2	16.0	16.5	17.2	16.5	17.0	15.8	16.0	†0.6
White . . . . .	15.4	15.9	16.9	17.5	18.0	17.7	18.1	16.5	16.9	†0.7
Black or African American . . . . .	10.4	10.3	11.9	11.8	13.4	13.3	12.6	12.9	13.5	†1.3
American Indian or Alaska Native <sup>3</sup> . . . . .	*	*	13.5	*	*	13.1	13.0	15.1	10.9	*
Asian or Pacific Islander . . . . .	9.5	12.2	11.5	12.4	12.7	9.7	11.9	12.0	11.3	0.5
Hispanic or Latina <sup>4</sup> . . . . .	13.7	13.1	13.8	13.6	15.3	15.1	15.6	14.5	14.9	†0.8
White, not Hispanic or Latina <sup>4</sup> . . . . .	15.6	16.2	17.3	18.1	18.4	18.1	18.4	16.9	17.1	†0.7

See footnotes at end of table.

**Table 36 (page 4 of 4). Age-adjusted cancer incidence rates for selected cancer sites, by sex, race, and Hispanic origin: United States, selected geographic areas, selected years 1990–2012**

Updated data when available, Excel, PDF, and more data years: <http://www.cdc.gov/nchs/hus/contents2015.htm#036>.

[Data are based on the Surveillance, Epidemiology, and End Results (SEER) Program's 13 population-based cancer registries]

Site, sex, race, and Hispanic origin	1990	1995	2000	2002	2003	2005	2010	2011	2012	1990–2012 APC <sup>1</sup>
Leukemia										
Number of new cases per 100,000 population <sup>2</sup>										
Male . . . . .	17.2	17.7	17.1	17.2	17.4	17.4	17.7	17.3	16.9	0.1
White . . . . .	18.0	19.0	18.2	18.6	18.5	18.9	19.2	18.5	17.8	0.1
Black or African American . . . . .	16.1	13.4	14.3	12.7	15.0	12.8	13.4	14.2	14.8	0.1
American Indian or Alaska Native <sup>3</sup> . . . . .	*	*	*	7.9	*	12.3	*	9.7	14.6	*
Asian or Pacific Islander . . . . .	8.4	10.2	10.6	9.6	10.5	9.3	10.0	10.3	10.2	0.0
Hispanic or Latino <sup>4</sup> . . . . .	12.2	14.6	13.2	12.3	12.3	13.2	13.0	13.6	11.2	0.3
White, not Hispanic or Latino <sup>4</sup> . . . . .	18.3	19.3	18.7	19.2	19.0	19.3	19.7	19.1	18.6	†0.2
Female . . . . .	9.9	10.3	10.4	10.1	10.1	10.1	10.6	10.3	10.4	†0.2
White . . . . .	10.3	10.9	11.0	10.9	10.7	10.6	11.3	11.0	11.1	†0.3
Black or African American . . . . .	8.5	8.4	9.8	7.5	9.1	9.5	9.0	9.1	9.1	0.1
American Indian or Alaska Native <sup>3</sup> . . . . .	*	*	*	*	*	*	8.7	*	6.6	*
Asian or Pacific Islander . . . . .	5.7	6.4	6.4	6.4	6.5	6.4	6.0	6.3	6.5	0.0
Hispanic or Latina <sup>4</sup> . . . . .	8.5	8.2	7.8	8.7	7.2	8.2	8.6	8.5	8.8	0.4
White, not Hispanic or Latina <sup>4</sup> . . . . .	10.3	11.1	11.1	10.9	11.1	10.6	11.6	11.3	11.2	†0.4

† Annual percent change (APC) is significantly different from zero ( $p < 0.05$ ).

0.0 APC is greater than -0.05 but less than 0.05.

\* Estimates are considered unreliable. Data not shown if the rate is based on fewer than 16 cases for the time interval. The trend is not shown if it is based on fewer than 10 cases for at least 1 year within the time interval.

<sup>1</sup> APC was calculated by fitting a linear regression model to the natural logarithm of the yearly rates from 1990–2012.

<sup>2</sup> Age-adjusted by 5-year age groups to the year 2000 U.S. standard population. Age-adjusted rates are based on at least 16 cases. See Appendix II, Age adjustment.

<sup>3</sup> Estimates for the American Indian or Alaska Native populations are based on the Contract Health Service Delivery Area (CHSDA) counties within SEER areas.

<sup>4</sup> Hispanic data exclude cases from Alaska. The race groups, white, black, Asian or Pacific Islander, and American Indian or Alaska Native, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. The North American Association of Central Cancer Registries (NAACCR) Hispanic Identification Algorithm was used on a combination of variables to classify cases as Hispanic for analytic purposes. See the report, NAACCR guideline for enhancing Hispanic-Latino identification, for more information. Available from: [http://seer.cancer.gov/seerstat/variables/seer/yr1973\\_2006/race\\_ethnicity/](http://seer.cancer.gov/seerstat/variables/seer/yr1973_2006/race_ethnicity/). See Appendix II, Hispanic origin.

NOTES: See Appendix II, Incidence. Estimates are based on 13 SEER areas (November 2014 submission) and differ from published estimates based on 9 SEER areas or other submission dates. See Appendix I, Surveillance, Epidemiology, and End Results Program (SEER). The site variable distinguishes Kaposi Sarcoma and Mesothelioma as individual cancer sites. As a result, Kaposi Sarcoma and Mesothelioma cases do not contribute to other cancer sites. Estimates for 2001–2009 were computed using intercensal population estimates based on the 2000 and 2010 censuses. Data have been revised and differ from previous editions of *Health, United States*. Data for additional years are available. See the Excel spreadsheet on the *Health, United States* website at: <http://www.cdc.gov/nchs/hus.htm>.

SOURCE: National Institutes of Health, National Cancer Institute, Surveillance, Epidemiology, and End Results Program. Available from: <http://www.seer.cancer.gov>. See Appendix I, Surveillance, Epidemiology, and End Results Program (SEER).