

Data table for Figure 19. Mammography use and colorectal cancer testing use, by race and Hispanic origin: United States, selected years 1987–2015

Excel and PowerPoint: <http://www.cdc.gov/nchs/hus/contents2016.htm#fig19>

Use of mammography in the past 2 years among women aged 40–74: United States, selected years 1987–2015

Year	Race and Hispanic origin ¹			
	Hispanic or Latino	Not Hispanic or Latino		
		White only	Black only	Asian only
Percent of women having a mammogram within the past 2 years ²				
1987	19.3	32.5	25.6	*
1990	45.3	55.7	47.1	47.7
1991	49.9	59.4	48.8	47.0
1993	55.0	64.2	59.6	56.3
1994	53.6	64.4	65.4	58.1
1998	60.9	69.9	67.7	59.0
1999	66.1	73.4	72.0	59.3
2000	61.5	74.1	69.3	53.4
2003	65.5	72.2	72.3	57.8
2005	59.6	70.6	66.9	55.4
2008	61.7	70.5	70.1	69.0
2010	65.2	69.9	69.5	64.1
2013	62.1	69.6	68.1	70.4
2015	62.8	68.2	72.3	62.7
Standard error				
1987	2.4	0.8	2.2	*
1990	2.5	0.7	1.6	4.8
1991	2.5	0.7	1.6	4.2
1993	3.4	0.8	2.3	5.7
1994	3.7	0.8	2.1	5.5
1998	1.7	0.7	1.6	4.2
1999	1.9	0.7	1.6	4.5
2000	1.8	0.7	1.6	4.5
2003	1.7	0.7	1.6	3.9
2005	1.8	0.7	1.6	3.8
2008	2.2	0.9	1.8	3.3
2010	1.7	0.9	1.5	2.7
2013	1.5	0.8	1.7	2.4
2015	1.7	0.8	1.6	3.0

Use of any colorectal test or procedure among adults aged 50–75: United States, selected years 2000–2015

Percent of adults with any colorectal test or procedure ³				
2000 ⁴	21.7	35.7	29.7	20.6
2003	27.2	41.0	35.3	26.3
2005	28.5	47.4	38.0	30.0
2008	34.0	54.8	47.4	47.3
2010	46.5	61.3	55.3	46.6
2013	41.5	60.4	58.2	51.2
2015	47.4	65.6	60.3	52.1
Standard error				
2000 ⁴	1.7	0.7	1.8	3.4
2003	1.7	0.7	1.7	3.7
2005	1.8	0.7	1.7	3.4
2008	2.0	0.8	2.0	3.4
2010	1.9	0.8	1.6	2.9
2013	1.7	0.7	1.5	2.7
2015	1.8	0.8	1.5	2.8

* Estimates are considered unreliable. Data not shown have an RSE greater than 30%.

¹ Starting with 1999 data, race-specific estimates are tabulated according to the 1997 *Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity* and are not strictly comparable with estimates for earlier years. Prior to 1999, data were tabulated according to the 1977 Standards with four racial groups, and the Asian only category included Native Hawaiian or Other Pacific Islander. Estimates for single-race categories prior to 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. See Appendix II, Hispanic origin; Race.

² Questions concerning use of mammography differed slightly on the National Health Interview Survey across the survey years. See Appendix II, Mammography. Data prior to 1997 are not strictly comparable with data for later years due to the 1997 questionnaire redesign.

³ Adults with a colorectal test or procedure are those who reported a home fecal occult blood test (FOBT) in the past year, a sigmoidoscopy procedure in the past five years with FOBT in the past three years, or a colonoscopy in the past 10 years. Questions concerning colorectal testing differed slightly on the National Health Interview Survey across survey years. See Appendix II, Colorectal tests or procedures.

⁴ Colorectal cancer testing data begin in 2000 due to changes in available procedures prior to 2000.

SOURCE: NCHS, National Health Interview Survey. See Appendix I, National Health Interview Survey (NHIS).