Table 30. Death rates for suicide, by sex, race, Hispanic origin, and age: United States, selected years 1950–2016

Sex, race, Hispanic origin, and age	1950 ^{1,2}	1960 ^{1,2}	1970 ²	1980 ²	1990 ²	2000 ³	2010 ³	2015 ³	2016 ³
All persons			De	eaths per 10	00,000 resid	lent populat	ion		
All ages, age-adjusted ⁴	13.2	12.5	13.1	12.2	12.5	10.4	12.1	13.3	13.5
All ages, crude	11.4	10.6	11.6	11.9	12.4	10.4	12.4	13.7	13.9
Under 1 year									
1–4 years					•••	•••			
5–14 years	0.2	0.3	0.3	0.4	8.0	0.7	0.7	1.0	1.1
15–24 years	4.5	5.2	8.8	12.3	13.2	10.2	10.5	12.5	13.2
15–19 years	2.7	3.6	5.9	8.5	11.1	8.0	7.5	9.8	10.0
20–24 years	6.2	7.1	12.2	16.1	15.1	12.5	13.6	15.1	16.1
25-44 years	11.6	12.2	15.4	15.6	15.2	13.4	15.0	16.4	16.9
25–34 years	9.1	10.0	14.1	16.0	15.2	12.0	14.0	15.7	16.5
35–44 years	14.3	14.2	16.9	15.4	15.3	14.5	16.0	17.1	17.4
45–64 years	23.5	22.0	20.6	15.9	15.3	13.5	18.6	19.6	19.2
45–54 years	20.9	20.7	20.0	15.9	14.8	14.4	19.6	20.3	19.7
55–64 years	26.8	23.7	21.4	15.9	16.0	12.1	17.5	18.9	18.7
65 years and over	30.0	24.5	20.8	17.6	20.5	15.2	14.9	16.6	16.7
65–74 years	29.6	23.0	20.8	16.9	17.9	12.5	13.7	15.2	15.4
75–84 years	31.1	27.9	21.2	19.1	24.9	17.6	15.7	17.9	18.2
85 years and over	28.8	26.0	19.0	19.2	22.2	19.6	17.6	19.4	19.0
Male									
All ages, age-adjusted ⁴	21.2	20.0	19.8	19.9	21.5	17.7	19.8	21.1	21.4
All ages, crude	17.8	16.5	16.8	18.6	20.4	17.1	19.9	21.5	21.8
Under 1 year									
1–4 years									
5–14 years	0.3	0.4	0.5	0.6	1.1	1.2	0.9	1.2	1.3
15–24 years	6.5	8.2	13.5	20.2	22.0	17.1	16.9	19.4	20.5
15–19 years	3.5	5.6	8.8	13.8	18.1	13.0	11.7	14.2	14.8
20–24 years	9.3	11.5	19.3	26.8	25.7	21.4	22.2	24.2	25.9
25–44 years	17.2	17.9	20.9	24.0	24.4	21.3	23.6	25.2	26.2
25–34 years	13.4	14.7	19.8	25.0	24.8	19.6	22.5	24.7	26.0
35–44 years	21.3	21.0	22.1	22.5	23.9	22.8	24.6	25.9	26.3
45–64 years	37.1	34.4	30.0	23.7	24.3	21.3	29.2	29.5	29.1
45–54 years	32.0	31.6	27.9	22.9	23.2	22.4	30.4	30.1	29.4
55–64 years	43.6	38.1	32.7	24.5	25.7	19.4	27.7	28.9	28.7
65 years and over	52.8	44.0	38.4	35.0	41.6	31.1	29.0	31.0	31.0
65–74 years	50.5	39.6	36.0	30.4	32.2	22.7	23.9	26.2	25.9
75–84 years	58.3	52.5	42.8	42.3	56.1	38.6	32.3	35.2	36.0
85 years and over	58.3	57.4	42.4	50.6	65.9	57.5	47.3	48.2	48.0
,	55.5	~		00.0	00.0	00	5		

Page 1 of 5 Trend Tables Health, United States, 2017

Table 30. Death rates for suicide, by sex, race, Hispanic origin, and age: United States, selected years 1950–2016

Sex, race, Hispanic origin, and age	1950 ^{1,2}	1960 ^{1,2}	1970 ²	1980 ²	1990 ²	2000 ³	2010 ³	2015 ³	2016 ³
Female			De	eaths per 10	0,000 resid	ent populat	ion		
All ages, age-adjusted ⁴	5.6	5.6	7.4	5.7	4.8	4.0	5.0	6.0	6.0
All ages, crude	5.1	4.9	6.6	5.5	4.8	4.0	5.2	6.2	6.2
Under 1 year									
1–4 years									
5–14 years	0.1	0.1	0.2	0.2	0.4	0.3	0.4	8.0	0.9
15–24 years	2.6	2.2	4.2	4.3	3.9	3.0	3.9	5.3	5.4
15–19 years	1.8	1.6	2.9	3.0	3.7	2.7	3.1	5.1	5.0
20–24 years	3.3	2.9	5.7	5.5	4.1	3.2	4.7	5.5	5.8
25–44 years	6.2	6.6	10.2	7.7	6.2	5.4	6.4	7.5	7.6
25–34 years	4.9	5.5	8.6	7.1	5.6	4.3	5.3	6.6	6.7
35–44 years	7.5	7.7	11.9	8.5	6.8	6.4	7.5	8.4	8.5
45–64 years	9.9	10.2	12.0	8.9	7.1	6.2	8.6	10.2	9.9
45–54 years	9.9	10.2	12.6	9.4	6.9	6.7	9.0	10.7	10.3
55–64 years	9.9	10.2	11.4	8.4	7.3	5.4	8.0	9.7	9.4
65 years and over	9.4	8.4	8.1	6.1	6.4	4.0	4.2	5.1	5.3
65–74 years	10.1	8.4	9.0	6.5	6.7	4.0	4.8	5.7	6.2
75–84 years	8.1	8.9	7.0	5.5	6.3	4.0	3.7	4.6	4.5
85 years and over	8.2	6.0	5.9	5.5	5.4	4.2	3.3	4.2	3.5
White male ⁵									
All ages, age-adjusted ⁴	22.3	21.1	20.8	20.9	22.8	19.1	22.0	23.6	23.9
All ages, crude	19.0	17.6	18.0	19.9	22.0	18.8	22.6	24.6	24.8
15–24 years	6.6	8.6	13.9	21.4	23.2	17.9	18.3	20.9	21.9
25–44 years	17.9	18.5	21.5	24.6	25.4	22.9	26.2	28.6	29.2
45–64 years	39.3	36.5	31.9	25.0	26.0	23.2	33.0	33.8	33.2
65 years and over	55.8	46.7	41.1	37.2	44.2	33.3	31.7	34.3	34.4
65–74 years	53.2	42.0	38.7	32.5	34.2	24.3	26.3	29.1	28.8
75–84 years	61.9	55.7	45.5	45.5	60.2	41.1	34.9	38.6	39.7
85 years and over	61.9	61.3	45.8	52.8	70.3	61.6	50.8	52.5	52.5
Black or African American male ⁵									
All ages, age-adjusted ⁴	7.5	8.4	10.0	11.4	12.8	10.0	9.1	9.6	10.2
All ages, crude	6.3	6.4	8.0	10.3	12.0	9.4	8.7	9.4	10.1
15–24 years	4.9	4.1	10.5	12.3	15.1	14.2	11.1	12.9	14.7
25–44 years	9.8	12.6	16.1	19.2	19.6	14.3	14.5	14.8	16.4
45–64 years	12.7	13.0	12.4	11.8	13.1	9.9	9.5	9.7	9.6
65 years and over	9.0	9.9	8.7	11.4	14.9	11.5	8.3	8.6	8.7
65–74 years	10.0	11.3	8.7	11.1	14.7	11.1	7.6	7.3	8.0
75–84 years ⁶	*	*	*	10.5	14.4	12.1	9.9	11.2	10.6
85 years and over		*	*	*	*	*	*	*	*

Page 2 of 5 Trend Tables Health, United States, 2017

Table 30. Death rates for suicide, by sex, race, Hispanic origin, and age: United States, selected years 1950–2016

Sex, race, Hispanic origin, and age	1950 ^{1,2}	1960 ^{1,2}	1970 ²	1980 ²	1990 ²	2000 ³	2010 ³	2015 ³	2016 ³
American Indian or Alaska Native male ⁵			De	eaths per 10	0,000 resid	ent populat	ion		
All ages, age-adjusted ⁴				19.3	20.1	16.0	15.5	18.8	20.7
All ages, crude				20.9	20.9	15.9	16.1	18.5	20.7
15–24 years				45.3	49.1	26.2	30.6	29.5	32.6
25–44 years				31.2	27.8	24.5	20.9	27.3	33.6
45–64 years				*	*	15.4	17.8	20.8	19.8
65 years and over				*	*	*	*	13.2	11.8
Asian or Pacific Islander male ⁵									
All ages, age-adjusted ⁴				10.7	9.6	8.6	9.5	9.1	10.1
All ages, crude				8.8	8.7	7.9	9.3	9.2	10.3
15–24 years				10.8	13.5	9.1	10.9	15.6	16.3
25–44 years				11.0	10.6	9.9	10.6	9.6	11.9
45–64 years				13.0	9.7	9.7	12.8	10.9	11.8
65 years and over				18.6	16.8	15.4	14.9	12.6	12.1
Hispanic or Latino male ^{5,7}									
All ages, age-adjusted ⁴					13.7	10.3	9.9	9.9	10.9
All ages, crude					11.4	8.4	8.5	9.0	10.1
15–24 years					14.7	10.9	10.7	12.8	13.5
25–44 years					16.2	11.2	11.2	12.2	14.6
45–64 years					16.1	12.0	12.9	11.4	12.7
65 years and over					23.4	19.5	15.7	14.5	13.7
White, not Hispanic or Latino male ⁷									
All ages, age-adjusted ⁴					23.5	20.2	24.2	26.6	26.6
All ages, crude					23.1	20.4	25.7	28.3	28.3
15–24 years					24.4	19.5	20.4	23.4	24.5
25–44 years					26.4	25.1	30.3	33.5	33.5
45–64 years					26.8	24.0	35.4	37.2	36.5
65 years and over					45.4	33.9	32.7	35.9	36.1
Black, not Hispanic or Latino male ⁷									
All ages, age-adjusted ⁴						10.2	9.4	10.0	10.6
All ages, crude						9.6	9.1	9.8	10.6
15–24 years						14.6	11.5	13.1	15.4
25–44 years						14.7	15.1	15.5	17.1
45–64 years						9.8	9.8	10.0	10.0
65 years and over						11.8	8.4	8.8	8.9

Page 3 of 5 Trend Tables Health, United States, 2017

Table 30. Death rates for suicide, by sex, race, Hispanic origin, and age: United States, selected years 1950–2016

Sex, race, Hispanic origin, and age	1950 ^{1,2}	1960 ^{1,2}	1970 ²	1980 ²	1990 ²	2000 ³	2010 ³	2015 ³	2016 ³
White female ⁵			De	eaths per 10	0,000 resid	ent populati	on		
All ages, age-adjusted ⁴	6.0	5.9	7.9	6.1	5.2	4.3	5.6	6.9	6.9
All ages, crude	5.5	5.3	7.1	5.9	5.3	4.4	5.9	7.2	7.2
5–24 years	2.7	2.3	4.2	4.6	4.2	3.1	4.2	5.6	5.5
25–44 years	6.6	7.0	11.0	8.1	6.6	6.0	7.3	8.7	8.8
15–64 years	10.6	10.9	13.0	9.6	7.7	6.9	9.9	11.9	11.6
5 years and over	9.9	8.8	8.5	6.4	6.8	4.3	4.5	5.6	5.8
Black or African American female ⁵									
ll ages, age-adjusted ⁴	1.8	2.0	2.9	2.4	2.4	1.8	1.8	2.0	2.4
ıll ages, crude	1.5	1.6	2.6	2.2	2.3	1.7	1.8	2.1	2.4
5–24 years	1.8	*	3.8	2.3	2.3	2.2	2.0	3.6	4.0
5–44 years	2.3	3.0	4.8	4.3	3.8	2.6	2.8	2.7	3.6
5–64 years	2.7	3.1	2.9	2.5	2.9	2.1	2.1	2.6	2.3
5 years and over	*	*	2.6	*	1.9	1.3	*	0.9	1.4
American Indian or Alaska Native female ⁵									
ıll ages, age-adjusted ⁴				4.7	3.6	3.8	6.1	6.5	6.4
ıll ages, crude				4.7	3.7	4.0	5.9	6.6	6.4
5–24 years				*	*	*	10.4	11.7	11.8
5–44 years				10.7	*	7.2	7.4	8.8	10.7
5–64 years				*	*	*	6.2	6.9	4.1
5 years and over				*	*	*	*	*	*
Asian or Pacific Islander female ⁵									
ıll ages, age-adjusted ⁴				5.5	4.1	2.8	3.4	4.0	3.6
ıll ages, crude				4.7	3.4	2.7	3.4	4.1	3.7
5–24 years				*	3.9	2.7	3.5	4.9	6.3
5–44 years				5.4	3.8	3.3	4.1	4.5	3.7
5–64 years				7.9	5.0	3.2	4.7	5.2	4.5
5 years and over				*	8.5	5.2	4.3	5.2	4.7
Hispanic or Latina female ^{5,7}									
ıll ages, age-adjusted ⁴					2.3	1.7	2.1	2.6	2.6
all ages, crude					2.2	1.5	2.0	2.6	2.5
5–24 years					3.1	2.0	3.1	3.9	3.8
5–44 years					3.1	2.1	2.4	3.2	3.5
5–64 years					2.5	2.5	2.8	3.8	3.3
65 years and over					*	*	2.2	1.9	1.8

Page 4 of 5 Trend Tables Health, United States, 2017

Table 30. Death rates for suicide, by sex, race, Hispanic origin, and age: United States, selected years 1950–2016

Excel version (with more data years and standard errors when available): https://www.cdc.gov/nchs/hus/contents2017.htm#030. [Data are based on death certificates]

Sex, race, Hispanic origin, and age	1950 ^{1,2}	1960 ^{1,2}	1970 ²	1980 ²	1990 ²	2000 ³	2010 ³	2015 ³	2016 ³	
White, not Hispanic or Latina female ⁷	Deaths per 100,000 resident population									
All ages, age-adjusted ⁴					5.4	4.7	6.2	7.8	7.9	
All ages, crude					5.6	4.9	6.7	8.3	8.2	
15–24 years					4.3	3.3	4.4	6.1	6.1	
25–44 years					7.0	6.7	8.6	10.3	10.3	
45–64 years					8.0	7.3	10.7	13.1	13.0	
65 years and over					7.0	4.4	4.7	6.0	6.2	
Black, not Hispanic or Latina female ⁷										
All ages, age-adjusted ⁴						1.8	1.9	2.1	2.4	
All ages, crude						1.7	1.8	2.1	2.5	
15–24 years						2.3	2.1	3.7	4.0	
25–44 years						2.7	2.9	2.9	3.8	
45–64 years						2.2	2.2	2.6	2.3	
65 years and over						1.3	*	0.9	1.4	

^{...} Category not applicable.

NOTES: Starting with Health, United States, 2003, rates for 1991–1999 (shown in spreadsheet version) were revised using intercensal population estimates based on the 1990 and 2000 Censuses. For 2000, population estimates are bridged-race April 1 census counts. Starting with Health, United States, 2012, rates for 2001–2009 (shown in spreadsheet version) were revised using intercensal population estimates based on the 2000 and 2010 Censuses. For 2010, population estimates are bridged-race April 1 census counts. Rates for 2011 and beyond were computed using 2010-based postcensal estimates. See Appendix I, Population Census and Population Estimates. Figures for 2001 include September 11-related deaths for which death certificates were filed as of October 24, 2002. See Appendix II, Cause of death; Table IV for terrorism-related ICD-10 codes. Age groups were selected to minimize the presentation of unstable age-specific death rates based on small numbers of deaths and for consistency among comparison groups. For additional injury-related statistics, see Web-based Injury Statistics Query and Reporting System (WISQARS), available from: https://www.cdc.gov/injury/wisqars/index.html. Starting with 2003 data (shown in spreadsheet version), some states began to collect information on more than one race on the death certificate, according to 1997 Office of Management and Budget (OMB) standards. The multiple-race data for these states were bridged to the single-race categories of the 1977 OMB standards, for comparability with other states. See Appendix II, Race. Data for additional years are available. See the Excel spreadsheet on the Health, United States website at: https://www.cdc.gov/inchs/hus.htm.

SOURCE: NCHS, National Vital Statistics System; Grove RD, Hetzel AM. Vital statistics rates in the United States, 1940–1960. 1968; numerator data from National Vital Statistics System, annual public-use Mortality Files; denominator data from national population estimates for race groups from Table 1 and unpublished Hispanic population estimates for 1985–1996 prepared by the Housing and Household Economic Statistics Division, U.S. Census Bureau; Xu JQ, Murphy SL, Kochanek KD, Bastian B, Arias E. Deaths: Final data for 2016. National Vital Statistics Reports; vol 67. Hyattsville, MD: National Center for Health Statistics. 2018. Available from: https://www.cdc.gov/nchs/products/nvsr.htm. See Appendix I, National Vital Statistics System (NVSS).

Page 5 of 5 Trend Tables Health, United States, 2017

^{*} Rates based on fewer than 20 deaths are considered unreliable and are not shown.

^{- - -} Data not available.

¹Includes deaths of persons who were not residents of the 50 states and the District of Columbia (D.C.).

²Underlying cause of death was coded according to the 6th revision of the *International Classification of Diseases* (ICD) in 1950, 7th revision in 1960, 8th revision in 1970, and 9th revision in 1980–1998. See Appendix II, Cause of death; Table III; Table IV.

³Starting with 1999 data, cause of death is coded according to ICD-10. See Appendix II, Cause of death; Comparability ratio; Table IV; Table V.

⁴Age-adjusted rates are calculated using the year 2000 standard population. Prior to 2001, age-adjusted rates were calculated using standard million proportions based on rounded population numbers. Starting with 2001 data (shown in spreadsheet version), unrounded population numbers are used to calculate age-adjusted rates. See Appendix II, Age adjustment.

⁵The race groups white, black, Asian or Pacific Islander, and American Indian or Alaska Native include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Death rates for Hispanic, American Indian or Alaska Native, and Asian or Pacific Islander persons should be interpreted with caution because of inconsistencies in reporting Hispanic origin or race on the death certificate (death rate numerators) compared with population figures (death rate denominators). The net effect of misclassification is an underestimation of deaths and death rates of the race of the rate and black. See Appendix II, Race, for a detailed discussion of sources of bias in death rates by race and Hispanic origin.

⁷Prior to 1997, data from states that did not report Hispanic origin on the death certificate were excluded. See Appendix II, Hispanic origin.