

**Data table for Figure 12. Vaccination coverage for combined 7-vaccine series among children aged 19–35 months, by state: United States, 2016**

Excel and PowerPoint: [https://www.cdc.gov/nchs/hus/contents2017.htm#Figure\\_012](https://www.cdc.gov/nchs/hus/contents2017.htm#Figure_012)

State	Percent
United States . . . . .	70.7
Alabama . . . . .	77.3
Alaska . . . . .	68.8
Arizona . . . . .	69.9
Arkansas . . . . .	67.8
California . . . . .	65.3
Colorado . . . . .	76.4
Connecticut . . . . .	75.7
Delaware . . . . .	78.1
District of Columbia . . . . .	68.2
Florida . . . . .	67.1
Georgia . . . . .	77.3
Hawaii . . . . .	75.1
Idaho . . . . .	73.9
Illinois . . . . .	71.5
Indiana . . . . .	68.8
Iowa . . . . .	73.5
Kansas . . . . .	76.4
Kentucky . . . . .	74.5
Louisiana . . . . .	66.8
Maine . . . . .	70.6
Maryland . . . . .	74.4
Massachusetts . . . . .	85.3
Michigan . . . . .	70.2
Minnesota . . . . .	73.8
Mississippi . . . . .	70.4
Missouri . . . . .	66.9
Montana . . . . .	63.6
Nebraska . . . . .	80.6
Nevada . . . . .	71.9
New Hampshire . . . . .	78.0
New Jersey . . . . .	70.2
New Mexico . . . . .	68.5
New York . . . . .	72.3
North Carolina . . . . .	77.8
North Dakota . . . . .	68.2
Ohio . . . . .	68.0
Oklahoma . . . . .	67.0
Oregon . . . . .	58.1
Pennsylvania . . . . .	73.7
Rhode Island . . . . .	75.5
South Carolina . . . . .	69.7
South Dakota . . . . .	70.4
Tennessee . . . . .	67.4
Texas . . . . .	69.5
Utah . . . . .	72.2
Vermont . . . . .	76.8
Virginia . . . . .	65.9
Washington . . . . .	75.7
West Virginia . . . . .	64.7
Wisconsin . . . . .	79.4
Wyoming . . . . .	62.8

NOTES: The combined 7-vaccine series consists of 4 or more doses of either the diphtheria, tetanus toxoids, and pertussis vaccine (DTP), the diphtheria and tetanus toxoids vaccine (DT), or the diphtheria, tetanus toxoids, and acellular pertussis vaccine (DTaP); 3 or more doses of any poliovirus vaccine; 1 or more doses of a measles-containing vaccine (MCV); 3 or more doses or 4 or more doses of *Haemophilus influenzae* type b vaccine (Hib) depending on Hib vaccine product type (full series Hib); 3 or more doses of hepatitis B vaccine; 1 or more doses of varicella vaccine; and 4 or more doses of pneumococcal conjugate vaccine (PCV). Data from U.S. territories were not included in the national estimates. Data for the map are displayed by a modified Jenks classification for the 50 U.S. states and D.C., which creates categories that minimize within-group variation and maximize between-group variation. Data are for the civilian noninstitutionalized population.

SOURCE: National Center for Immunization and Respiratory Diseases, National Immunization Survey-Child. See [Appendix I](#), National Immunization Survey (NIS). CDC/NCIRD 2016 childhood combined 7-vaccine series coverage report. Available from: <https://www.cdc.gov/vaccines/imz-managers/coverage/childvaxview/data-reports/7-series/reports/2016.html>.