

Table 30. Health care visits to doctor offices, emergency departments, and home visits within the past 12 months, by selected characteristics: United States, selected years 1997–2018

Excel version (with more data years and standard errors when available): <https://www.cdc.gov/nchs/hus/contents2019.htm#Table-030>

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Number of health care visits ¹											
	None			1–3 visits			4–9 visits			10 or more visits		
	1997	2010	2018	1997	2010	2018	1997	2010	2018	1997	2010	2018
	Percent distribution											
Total, age adjusted ^{2,3}	16.5	15.6	14.5	46.2	45.4	49.2	23.6	25.8	23.4	13.7	13.2	12.9
Total, crude ²	16.5	15.4	14.0	46.5	45.2	48.4	23.5	26.0	24.0	13.5	13.5	13.6
Age group												
Under 18 years	11.8	8.1	7.4	54.1	55.6	60.3	25.2	28.2	25.4	8.9	8.2	6.9
Under 6 years	5.0	3.7	4.9	44.9	48.9	52.7	37.0	36.8	33.9	13.0	10.6	8.4
6–17 years	15.3	10.4	8.6	58.7	59.1	63.9	19.3	23.6	21.4	6.8	6.9	6.2
18–44 years	21.7	24.2	21.9	46.7	43.9	47.6	19.0	20.6	18.7	12.6	11.3	11.7
18–24 years	22.0	25.9	22.2	46.8	43.4	48.9	20.0	21.1	19.1	11.2	9.6	9.8
25–44 years	21.6	23.6	21.8	46.7	44.1	47.2	18.7	20.5	18.6	13.0	11.9	12.4
45–64 years	16.9	14.8	14.2	42.9	42.8	45.5	24.7	26.1	24.7	15.5	16.4	15.7
45–54 years	17.9	17.6	16.5	43.9	43.5	47.6	23.4	23.9	21.6	14.8	15.0	14.3
55–64 years	15.3	11.1	11.9	41.3	41.9	43.4	26.7	28.8	27.6	16.7	18.2	17.1
65 years and over	8.9	5.3	5.5	34.7	33.8	38.0	32.5	36.7	32.5	23.8	24.2	24.0
65–74 years	9.8	6.3	6.4	36.9	36.1	40.1	31.6	35.7	31.8	21.6	21.9	21.7
75 years and over	7.7	4.1	4.1	31.8	31.0	34.8	33.8	38.0	33.6	26.6	27.0	27.6
Sex ³												
Male	21.3	20.4	18.7	47.1	46.4	49.7	20.6	22.7	21.2	11.0	10.5	10.3
Female	11.8	10.9	10.3	45.4	44.4	48.8	26.5	28.8	25.6	16.3	15.9	15.4
Race ^{3,4}												
White only	16.0	15.3	14.0	46.1	44.9	48.5	23.9	26.1	24.0	14.0	13.7	13.4
Black or African American only	16.8	15.7	13.8	46.1	47.2	51.9	23.2	24.7	22.1	13.9	12.4	12.2
American Indian or Alaska Native only	17.1	19.4	16.5	38.0	40.3	41.5	24.2	28.1	25.9	20.7	12.2	16.1
Asian only	22.8	20.4	19.9	49.1	49.9	53.3	19.7	22.1	18.8	8.3	7.6	8.0
Native Hawaiian or Other Pacific Islander only	---	*	20.9	---	*	53.2	---	*	15.7	---	*	*
2 or more races	---	13.9	15.3	---	42.3	47.1	---	25.2	23.5	---	18.6	14.1
Hispanic origin and race ^{3,4}												
Hispanic or Latino	24.9	23.5	20.3	42.3	43.2	48.1	20.3	22.6	21.0	12.5	10.7	10.6
Mexican	28.9	25.2	21.4	40.8	43.3	47.4	18.5	21.4	20.5	11.8	10.1	10.7
Not Hispanic or Latino	15.4	14.0	13.1	46.7	45.8	49.4	24.0	26.5	24.0	13.9	13.7	13.5
White only	14.7	13.2	12.1	46.6	45.3	48.5	24.4	27.1	25.1	14.3	14.4	14.3
Black or African American only	16.9	15.6	13.7	46.1	47.3	51.8	23.1	24.9	22.2	13.8	12.2	12.3
Percent of poverty level ^{3,5}												
Below 100%	20.6	20.4	18.9	37.8	37.5	40.8	22.7	25.1	23.2	18.9	17.0	17.1
100%–199%	20.1	20.8	18.3	43.3	42.1	45.7	21.7	23.1	22.0	14.9	13.9	13.9
200%–399%	16.4	16.2	15.1	47.2	46.3	49.8	23.6	25.4	22.8	12.8	12.1	12.3
400% or more	12.8	10.2	11.0	49.8	49.4	52.2	24.9	27.6	24.6	12.5	12.7	12.1

See footnotes at end of table.

Table 30. Health care visits to doctor offices, emergency departments, and home visits within the past 12 months, by selected characteristics: United States, selected years 1997–2018—Con.

Excel version (with more data years and standard errors when available): <https://www.cdc.gov/nchs/hus/contents2019.htm#Table-030>

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Number of health care visits ¹											
	None			1–3 visits			4–9 visits			10 or more visits		
	1997	2010	2018	1997	2010	2018	1997	2010	2018	1997	2010	2018
Percent distribution												
Hispanic origin and race and percent of poverty level ^{3–5}												
Hispanic or Latino:												
Below 100%	30.2	28.7	27.1	34.8	36.5	42.3	19.9	22.5	19.1	15.0	12.3	11.5
100%–199%	28.7	27.7	21.4	39.7	42.7	46.3	20.4	19.9	21.1	11.2	9.8	11.3
200%–399%	20.7	21.6	19.2	47.4	45.0	49.8	19.8	23.1	20.5	12.1	10.3	10.4
400% or more	15.2	11.3	14.3	50.4	51.1	53.4	22.6	26.1	22.9	11.8	11.5	9.5
Not Hispanic or Latino:												
White only:												
Below 100%	17.0	15.0	15.6	38.3	37.0	37.5	23.9	27.4	25.4	20.9	20.6	21.5
100%–199%	17.3	18.4	15.9	44.1	40.4	43.7	22.2	24.7	23.7	16.3	16.5	16.8
200%–399%	15.4	14.7	13.1	46.9	46.0	48.8	24.3	26.3	24.4	13.4	13.0	13.7
400% or more	12.5	9.9	10.1	49.1	48.2	50.8	25.5	28.4	25.9	13.0	13.5	13.3
Black or African American only:												
Below 100%	17.4	18.4	13.8	38.5	39.8	41.8	23.4	25.0	26.6	20.7	16.8	17.9
100%–199%	18.8	17.6	17.1	43.7	45.7	49.4	22.9	24.3	20.2	14.5	12.5	13.4
200%–399%	16.6	15.1	12.8	49.7	49.0	54.3	22.9	25.7	22.6	10.8	10.2	10.2
400% or more	14.0	10.0	11.5	54.3	58.2	59.4	22.7	22.5	19.6	9.0	9.3	9.5
Health insurance status at time of interview ^{6,7}												
Under 65 years:												
Insured	14.3	12.3	13.0	49.0	48.5	51.8	23.6	26.1	23.1	13.1	13.1	12.1
Private	14.7	12.4	13.2	50.6	51.0	53.6	23.1	25.5	22.7	11.6	11.1	10.4
Medicaid	9.8	10.9	12.2	35.5	38.2	44.5	26.5	28.0	24.3	28.2	23.0	19.0
Uninsured	33.7	37.2	36.8	42.8	42.2	44.9	15.3	15.2	13.2	8.2	5.4	5.1
Health insurance status prior to interview ^{6,7}												
Under 65 years:												
Insured continuously all 12 months	14.1	12.1	12.9	49.2	48.6	51.9	23.6	26.2	23.1	13.0	13.0	12.1
Uninsured for any period up to 12 months	18.9	18.5	21.2	46.0	47.8	48.4	20.8	22.0	20.9	14.4	11.6	9.5
Uninsured more than 12 months	39.0	43.8	45.3	41.4	39.7	41.6	13.2	12.6	9.9	6.4	3.9	3.1
Percent of poverty level and health insurance status prior to interview ^{5–7}												
Under 65 years:												
Below 100%:												
Insured continuously all 12 months	13.8	12.7	14.0	39.7	39.5	43.4	25.2	27.5	24.3	21.4	20.3	18.2
Uninsured for any period up to 12 months	19.7	16.9	24.6	37.6	43.0	35.6	21.9	25.0	24.1	20.9	15.1	15.8
Uninsured more than 12 months	41.2	45.0	53.0	39.9	38.1	35.4	12.2	13.6	8.9	6.6	3.3	2.7
100%–199%:												
Insured continuously all 12 months	16.0	14.8	15.0	46.4	44.4	48.5	21.9	24.8	21.8	15.8	16.0	14.6
Uninsured for any period up to 12 months	18.8	21.0	24.5	45.1	46.0	45.6	21.0	20.6	20.3	15.0	12.4	9.6
Uninsured more than 12 months	38.7	43.2	46.1	41.0	39.4	42.3	14.0	12.4	8.5	6.3	5.0	3.1
200%–399%:												
Insured continuously all 12 months	15.1	13.6	13.7	49.4	49.4	52.1	23.4	25.3	22.5	12.1	11.7	11.7
Uninsured for any period up to 12 months	17.9	18.8	22.0	49.3	49.7	53.3	20.0	19.7	18.3	12.8	11.8	6.4
Uninsured more than 12 months	37.0	43.8	42.7	43.8	40.7	40.9	12.6	13.3	12.6	6.6	*2.2	3.7

See footnotes at end of table.

Table 30. Health care visits to doctor offices, emergency departments, and home visits within the past 12 months, by selected characteristics: United States, selected years 1997–2018—Con.

Excel version (with more data years and standard errors when available): <https://www.cdc.gov/nchs/hus/contents2019.htm#Table-030>

[Data are based on household interviews of a sample of the civilian noninstitutionalized population]

Characteristic	Number of health care visits ¹											
	None			1–3 visits			4–9 visits			10 or more visits		
	1997	2010	2018	1997	2010	2018	1997	2010	2018	1997	2010	2018
Percent of poverty level and health insurance status prior to interview ^{5–7} —Con.	Percent distribution											
400% or more:												
Insured continuously all 12 months.....	12.4	9.7	11.3	52.2	51.8	54.1	23.9	26.8	23.8	11.5	11.6	10.8
Uninsured for any period up to 12 months.....	17.2	16.6	14.7	50.0	53.5	51.6	24.2	23.9	24.8	*8.5	*6.0	8.9
Uninsured more than 12 months.....	35.1	39.2	27.7	44.1	46.0	61.6	15.1	*8.8	7.8	*5.7	*	*
Respondent-assessed health status ³												
Fair or poor.....	7.8	8.4	8.3	23.3	24.0	29.2	29.0	30.2	25.5	39.9	37.3	37.0
Good to excellent.....	17.2	16.3	15.1	48.4	47.5	51.2	23.3	25.5	23.3	11.1	10.7	10.4
Level of difficulty ^{3,8}												
18 years and over:												
A lot of difficulty or cannot do at all.....	---	*9.7	7.6	---	20.5	27.9	---	27.2	24.9	---	42.6	39.5
Some difficulty.....	---	14.2	14.9	---	37.8	41.6	---	28.1	25.2	---	19.9	18.3
No difficulty.....	---	20.2	18.9	---	46.6	50.9	---	23.1	21.2	---	10.1	9.0
Geographic region ³												
Northeast.....	13.2	12.6	10.9	45.9	46.3	50.9	26.0	26.4	24.2	14.9	14.7	13.9
Midwest.....	15.9	13.4	13.5	47.7	46.8	49.8	22.8	26.4	23.3	13.6	13.3	13.4
South.....	17.2	16.1	15.3	46.1	44.2	48.3	23.3	26.6	23.7	13.5	13.2	12.7
West.....	19.1	19.1	16.4	44.8	45.2	49.0	22.8	23.5	22.5	13.3	12.2	12.1
Location of residence ^{3,9}												
Within MSA.....	16.2	15.6	14.4	46.4	45.8	49.5	23.7	25.6	23.3	13.7	13.0	12.8
Outside MSA.....	17.3	15.9	14.8	45.4	42.7	47.5	23.3	27.0	24.2	13.9	14.4	13.6

--- Data not available.

* Estimates are considered unreliable. Starting with 2016 data, the reliability of survey percentage estimates is assessed using new multistep National Center for Health Statistics data presentation standards for proportions. Before 2016 data, the reliability of estimates was evaluated based on relative standard errors (RSEs). Data preceded by an asterisk have an RSE of 20%–30%. Data not shown have an RSE greater than 30%. For more information, see Appendix II, Data presentation standards for proportions; Relative standard error (RSE).

¹Summary measure of the number of visits to hospital emergency departments, home visits by a nurse or other health care professional, and visits to doctor offices, clinics, or some other place during a 12-month period. See Appendix II, Emergency department or emergency room visit; Health care visit; Home visit.

²Includes all other races not shown separately, unknown health insurance status, and unknown disability status.

³Estimates are age adjusted to the year 2000 standard population using six age groups: under 18 years, 18–44 years, 45–54 years, 55–64 years, 65–74 years, and 75 years and over. The disability measure is age adjusted using five adult age groups: 18–44 years, 45–54 years, 55–64 years, 65–74 years, and 75 years and over. Age-adjusted estimates in this table may differ from other age-adjusted estimates based on the same data and presented elsewhere if different age groups are used in the adjustment procedure. See Appendix II, Age adjustment.

⁴The race groups white, black, American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, and 2 or more races include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. Starting with 1999 data, race-specific estimates are tabulated according to the 1997 “Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity” and are not strictly comparable with estimates for earlier years. The five single- and multiple-race categories shown follow the 1997 standards. Starting with 1999 data, race-specific estimates are for persons who reported only one racial group; the category 2 or more races includes persons who reported more than one racial group. Before 1999, data were tabulated with four racial groups according to the 1977 standards, and the Asian only category included Native Hawaiian or Other Pacific Islander. Estimates for single-race categories before 1999 included persons who reported one race or, if they reported more than one race, identified one race as best representing their race. Starting with 2003 data, race responses of other race and unspecified multiple race are treated as missing, and then race is imputed if these were the only race responses. Almost all persons with a race response of other race are of Hispanic origin. See Appendix II, Hispanic origin; Race.

⁵Percent of poverty level is based on family income and family size and composition using U.S. Census Bureau poverty thresholds. Missing family income data are imputed for 1997 and beyond. See Appendix II, Family income; Poverty; Table VII.

⁶Estimates for persons under age 65 are age adjusted to the year 2000 standard population using four age groups: under 18 years, 18–44 years, 45–54 years, and 55–64 years. Age-adjusted estimates in this table may differ from other age-adjusted estimates based on the same data and presented elsewhere if different age groups are used in the adjustment procedure. See Appendix II, Age adjustment.

⁷Health insurance categories are mutually exclusive. Persons who reported both Medicaid and private coverage are classified as having private coverage. Starting with 1997 data, state-sponsored health plan coverage is included as Medicaid coverage. Starting with 1999 data, coverage by the Children’s Health Insurance Program (CHIP) is included as Medicaid coverage. In addition to private and Medicaid, the insured category includes military plans, other government-sponsored health plans, and Medicare. Persons not covered by private insurance, Medicaid, CHIP, state-sponsored or other government-sponsored health plans (starting in 1997), Medicare, or military plans are considered to have no health insurance coverage. Persons with only Indian Health Service coverage are considered to have no health insurance coverage. See Appendix II, Health insurance coverage.

⁸Functional limitation is defined by the reported level of difficulty in six domains of functioning: seeing (even if wearing glasses), hearing (even if wearing hearing aids), mobility (walking or climbing stairs), communication (understanding or being understood by others), cognition (remembering or concentrating), and self-care (such as washing all over or dressing). Respondents with answers to one or more of the six questions are included in one of three mutually exclusive categories. Adults who respond “a lot of difficulty” or “cannot do at all/unable to do” to at least one question are classified in the “a lot of difficulty or cannot do at all” category. Of the remaining respondents, adults who respond “some difficulty” to at least one question are classified in the “some difficulty” category, and adults who respond “no difficulty” to all questions are classified in the “no difficulty” category. Adults who respond “don’t know” or “refused” to all six questions are excluded. During 2010–2018, 1%–8% of respondents were missing data and excluded. See Appendix II, Functional limitation.

⁹MSA is metropolitan statistical area. Starting with 2016 data, MSA status is determined using 2010 Census data and the 2010 standards for defining MSAs. For more information, including the definition applied before 2016, see Appendix II, Metropolitan statistical area (MSA).

NOTE: Standard errors and data for additional years are available in the Excel spreadsheet version of this table on the *Health, United States* website at: <https://www.cdc.gov/nchs/hus/index.htm>.

SOURCE: NCHS, National Health Interview Survey, Family Core and Sample Adult questionnaires. Data for level of difficulty are from the 2010 Quality of Life, 2011–2017 Functioning and Disability, and 2018 Sample Adult questionnaires. See Appendix I, National Health Interview Survey (NHIS).