

Table 39. Maternal mortality for complications of pregnancy, childbirth, and the puerperium, by race, Hispanic origin, and age: United States, selected years 1950-2007

(Data are based on death certificates)

Race, Hispanic origin, and age	1950 ^{1,2}	1960 ^{1,2}	1970 ²	1980 ²	1990 ²	2000 ³	2005 ^{3,4}	2006 ^{3,4}	2007 ^{3,4}
Number of deaths									
All persons.....	2,960	1,579	803	334	343	396	623	569	548
White.....	1,873	936	445	193	177	240	360	313	335
Black or African American.....	1,041	624	342	127	153	137	231	218	179
American Indian or Alaska Native.....	---	---	---	3	4	6	5	9	10
Asian or Pacific Islander.....	---	---	---	11	9	13	27	29	24
Hispanic or Latina ⁵	---	---	---	---	47	81	95	106	95
White, not Hispanic or Latina ⁵ ...	---	---	---	---	125	160	267	210	242
Deaths per 100,000 live births									
All persons									
All ages, age-adjusted ⁶	73.7	32.1	21.5	9.4	7.6	8.2	12.4	11.2	10.2
All ages, crude.....	83.3	37.1	21.5	9.2	8.2	9.8	15.1	13.3	12.7
Under 20 years.....	70.7	22.7	18.9	7.6	7.5	*	7.4	5.0	7.1
20-24 years.....	47.6	20.7	13.0	5.8	6.1	7.4	10.7	10.2	8.1
25-29 years.....	63.5	29.8	17.0	7.7	6.0	7.9	11.8	11.7	9.4
30-34 years.....	107.7	50.3	31.6	13.6	9.5	10.0	12.8	12.6	12.1
35 years and over ⁷	222.0	104.3	81.9	36.3	20.7	22.7	38.0	29.3	32.3
White									
All ages, age-adjusted ⁶	53.1	22.4	14.4	6.7	5.1	6.2	9.1	8.1	7.7
All ages, crude.....	61.1	26.0	14.3	6.6	5.4	7.5	11.1	9.5	10.0
Under 20 years.....	44.9	14.8	13.8	5.8	*	*	*	*	*
20-24 years.....	35.7	15.3	8.4	4.2	3.9	5.6	9.0	8.3	5.9
25-29 years.....	45.0	20.3	11.1	5.4	4.8	5.9	7.2	7.4	6.7
30-34 years.....	75.9	34.3	18.7	9.3	5.0	7.1	9.3	8.2	9.7
35 years and over ⁷	174.1	73.9	59.3	25.5	12.6	18.0	28.9	20.5	27.5
Black or African American									
All ages, age-adjusted ⁶	---	92.0	65.5	24.9	21.7	20.1	31.7	28.7	23.8
All ages, crude.....	---	103.6	60.9	22.4	22.4	22.0	36.5	32.7	26.5
Under 20 years.....	---	54.8	32.3	13.1	*	*	*	*	*
20-24 years.....	---	56.9	41.9	13.9	14.7	15.3	18.2	17.8	18.6
25-29 years.....	---	92.8	65.2	22.4	14.9	21.8	37.1	36.0	24.7
30-34 years.....	---	150.6	117.8	44.0	44.2	34.8	46.6	45.1	31.3
35 years and over ⁷	---	299.5	207.5	100.6	79.7	62.8	112.8	97.0	74.0
Hispanic or Latina ^{5,8}									

All ages, age-adjusted ⁶	---	---	---	---	7.4	9.0	8.2	8.8	7.2
All ages, crude.....	---	---	---	---	7.9	9.9	9.6	10.2	8.9
White, not Hispanic or Latina ⁵									
All ages, age-adjusted ⁶	---	---	---	---	4.4	5.5	9.6	8.0	8.1
All ages, crude.....	---	---	---	---	4.8	6.8	11.7	9.1	10.5

---Data not available.

-Quantity zero.

*Rates based on fewer than 20 deaths are considered unreliable and are not shown.

¹Includes deaths of persons who were not residents of the 50 states and the District of Columbia (D.C.).

²Underlying cause of death was coded according to the 6th Revision of the *International Classification of Diseases* (ICD) in 1950, 7th Revision in 1960, 8th Revision in 1970, and 9th Revision in 1980-1998. See Appendix II, Cause of death; Tables IV and V.

³Starting with 1999 data, cause of death is coded according to ICD-10. Major changes in the classification and coding of maternal deaths account for an increase in the number of maternal deaths under ICD-10.

See Appendix II, Cause of death, Table V; Comparability ratio, Table VI; *International Classification of Diseases* (ICD); Maternal death.

⁴In 2003, states began adopting the 2003 revision of the U.S. Standard Certificate of death that introduced a standard question format for maternal deaths.

Increases are due to methodological changes in reporting and data processing.

See Appendix II, Maternal death.

⁵Prior to 1997, excludes data from states lacking an Hispanic-origin item on the death certificate.

See Appendix II, Hispanic origin.

⁶Rates are age-adjusted to the 1970 distribution of live births by mother's age in the United States. See Appendix II, Age adjustment; Table III.

⁷Rates computed by relating deaths of women 35 years of age and over to live births to women 35-49 years of age.

See Appendix II, Rate: Death and related rates.

⁸Age-specific maternal mortality rates are not calculated because rates based on fewer than 20 deaths are considered unreliable.

NOTES: The race groups, white, black, Asian or Pacific Islander, and American Indian or Alaska Native, include persons of Hispanic and non-Hispanic origin. Persons of Hispanic origin may be of any race. For 1950 and 1960, rates were based on live births by race of child; for all other years, rates are based on live births by race of mother.

Starting with 2003 data, some states allowed the reporting of more than one race on the death certificate.

The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other states.

See Appendix II, Race. Rates are not calculated for American Indian or Alaska Native and Asian or Pacific Islander mothers because rates based on fewer than 20 deaths are considered unreliable.

Data for additional years are available. See Appendix III.

SOURCE: CDC/NCHS,

National Vital Statistics System;

numerator data from annual mortality files; denominator data from annual natality files;

Xu JQ, Kochanek KD, Murphy SL, Tejada-Vera B. Deaths: Final data for 2007. National vital statistics reports;

Vol 58 no 19. Hyattsville, MD: NCHS. 2010. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr58/nvsr58_19.pdf.

Health, United States 2009 Web Update