


Figure 15.1. Percentage of persons of all ages who experienced an asthma episode in the past 12 months: United States, 1997–2005




NOTES: Information on an episode of asthma or asthma attack during the past 12 months is self-reported by adults aged 18 years and over. For children under age 18 years, the information is collected from an adult family member, usually a parent, who is knowledgeable about the child's health. The analyses excluded people with unknown asthma episode status (about 0.3% of respondents each year). Beginning with the 2003 data, the National Health Interview Survey (NHIS) transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–02 were recalculated using weights derived from the 2000 census. See "About This Release" and Table III in the Appendix for more details.

DATA SOURCE: Combined Sample Adult and Sample Child Core components of the 1997–2005 NHIS. The estimate for 2005 was based on data collected from January through June. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ For the period January through June 2005, the percentage of persons of all ages who experienced an asthma episode in the past 12 months was 4.4% (95% confidence interval = 4.0–4.7%), which was not significantly different from the 2004 estimate of 4.1%.

■ The annual percentage of persons of all ages who experienced an asthma episode in the past 12 months was 4.2% in 1997, 4.0% in 1998, 3.9% in 1999, 4.0% in 2000, 4.3% in 2001, 4.3% in 2002, 3.9% in 2003, and 4.1% in 2004.

Figure 15.2. Percentage of persons of all ages who experienced an asthma episode in the past 12 months, by age group and sex: United States, January–June 2005




NOTES: Information on an episode of asthma or asthma attack during the past 12 months is self-reported by adults aged 18 years and over. For children under age 18 years, the information is collected from an adult family member, usually a parent, who is knowledgeable about the child's health. The analyses excluded 35 persons (0.2%) with unknown asthma episode status.

DATA SOURCE: Based on data collected from January through June in the Sample Adult and Sample Child Core components of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- For both sexes combined, the percentage of persons who had an asthma episode in the past 12 months was higher among children under age 15 years than among persons aged 15–34 years and 35 years and over. This age difference was seen among males, but not among females.

- For children under age 15 years, the percentage of persons who had an asthma episode in the past 12 months was higher among boys than among girls. However, for the age groups 15–34 years and 35 years and over, the percentage was higher among females than among males.

Figure 15.3. Sex-adjusted percentage of persons of all ages who experienced an asthma episode in the past 12 months, by age group and race/ethnicity: United States, January–June 2005




NOTES: Information on an episode of asthma or asthma attack during the past 12 months is self-reported by adults aged 18 years and over. For children under age 18 years, the information is collected from an adult family member, usually a parent, who is knowledgeable about the child's health. The analyses excluded 35 persons (0.2%) with unknown asthma episode status.

DATA SOURCE: Based on data collected from January through June in the Sample Adult and Sample Child Core components of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- For children under age 15 years, the sex-adjusted percentage of persons who had an asthma episode in the past 12 months was higher among non-Hispanic black children than among Hispanic or non-Hispanic white children, but this difference was not significant.
- For persons aged 15 years and over, the sex-adjusted percentages of persons who had an asthma episode in the past 12 months were similar across the three race/ethnicity groups.

Figure 15.4. Prevalence of current asthma among persons of all ages: United States, 2001–2005




NOTES: Information on current asthma is self-reported by adults aged 18 years and over. For children under age 18 years, the information is collected from an adult family member, usually a parent, who is knowledgeable about the child's health. The analyses excluded people with unknown current asthma status (about 0.3% of respondents each year). Beginning with the 2003 data, the National Health Interview Survey (NHIS) transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2001–02 were recalculated using weights derived from the 2000 census. See "About This Release" and Table III in the Appendix for more details.

DATA SOURCE: Combined Sample Adult and Sample Child Core components of the 2001–2005 NHIS. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ For the period January through June 2005, 7.9% (95% confidence interval = 7.4–8.3%) of persons of all ages currently had asthma, which was higher than the 2004 estimate of 7.3%.

■ The annual prevalence of current asthma among persons of all ages was 7.6% in 2001, 7.4% in 2002, 7.1% in 2003, and 7.3% in 2004.

Figure 15.5. Prevalence of current asthma among persons of all ages, by age group and sex: United States, January–June 2005




NOTES: Information on current asthma is self-reported by adults aged 18 years and over. For children under age 18 years, the information is collected from an adult family member, usually a parent, who is knowledgeable about the child's health. The analyses excluded 41 persons (0.2%) with unknown current asthma status.

DATA SOURCE: Based on data collected from January through June in the Sample Adult and Sample Child Core components of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ For both sexes combined, the prevalence of current asthma was higher among children 0–14 years old than among persons aged 15–34 years and 35 years and over. This pattern was seen among males, but not females.

■ For children under age 15 years, the prevalence of current asthma was higher among boys than among girls. However, for the age groups 15–34 years and 35 years and over, the prevalence of current asthma was higher among females than among males.

Figure 15.6. Sex-adjusted prevalence of current asthma among persons of all ages, by age group and race/ethnicity: United States, January–June 2005



NOTES: Information on current asthma is self-reported by adults aged 18 years and over. For children under age 18 years, the information is collected from an adult family member, usually a parent, who is knowledgeable about the child's health. The analyses excluded 41 persons (0.2%) with unknown current asthma status.

DATA SOURCE: Based on data collected from January through June in the Sample Adult and Sample Child Core components of the 2005 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

■ For children under age 15 years, the sex-adjusted prevalence of current asthma was higher among non-Hispanic black children than among Hispanic or non-Hispanic white children.

■ Among the six age-specific race/ethnicity groups, the sex-adjusted prevalence of current asthma was highest among non-Hispanic black children under age 15 years (13.2%).

Data tables for figures 15.1–15.6:

Data for figure 15.1. Percentage of persons of all ages who experienced an asthma episode in the past 12 months: United States, 1997–2005

Year	Percent (95% confidence interval)	
	Crude ¹	Age-adjusted ²
1997	4.2 (4.0–4.4)	4.2 (3.9–4.4)
1998	4.0 (3.7–4.2)	3.9 (3.7–4.2)
1999	3.9 (3.6–4.1)	3.9 (3.6–4.1)
2000	4.0 (3.8–4.2)	4.0 (3.8–4.2)
2001	4.3 (4.1–4.6)	4.3 (4.1–4.5)
2002	4.3 (4.0–4.5)	4.3 (4.0–4.5)
2003	3.9 (3.7–4.1)	3.9 (3.7–4.1)
2004	4.1 (3.8–4.3)	4.1 (3.8–4.3)
January–June 2005	4.4 (4.0–4.7)	4.4 (4.0–4.7)

¹Crude estimates are presented in the figure.

²Estimates are age adjusted to the 2000 projected U.S. standard population using three age groups: 0–14 years, 15–34 years, and 35 years and over.

NOTES: Beginning with the 2003 data, the National Health Interview Survey (NHIS) transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–02 were recalculated using weights derived from the 2000 census. See “About This Release” and Table III in the Appendix for more details.

DATA SOURCE: NHIS, 1997–2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

Data table for figure 15.2. Percentage of persons of all ages who experienced an asthma episode in the past 12 months, by age group and sex: United States, January–June 2005

Age and sex	Percent	95% confidence interval
0–14 years		
Total	5.9	5.0–6.7
Male	7.0	5.8–8.3
Female	4.6	3.7–5.6
15–34 years		
Total	4.3	3.6–4.9
Male	2.9	2.1–3.8
Female	5.6	4.6–6.7
35 years and over		
Total	3.8	3.4–4.2
Male	2.7	2.1–3.2
Female	4.9	4.2–5.5
All ages: crude¹		
Total	4.4	4.0–4.7
Male	3.7	3.2–4.1
Female	5.0	4.5–5.5
All ages: age-adjusted²		
Total	4.4	4.0–4.7
Male	3.7	3.2–4.1
Female	5.0	4.5–5.5

¹Crude estimates are presented in the figure.

²Estimates are age adjusted to the 2000 projected U.S. standard population using three age groups: 0–14 years, 15–34 years, and 35 years and over.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

Data table for figure 15.3. Sex-adjusted percentage of persons of all ages who experienced an asthma episode in the past 12 months, by age group and race/ethnicity: United States, January–June 2005

Age and race/ethnicity	Percent ¹	95% confidence interval
0–14 years		
Hispanic or Latino	5.6	4.1–7.0
Not Hispanic or Latino:		
White, single race	5.5	4.5–6.5
Black, single race	6.4	3.9–8.8
15 years and over		
Hispanic or Latino	3.2	2.6–3.9
Not Hispanic or Latino:		
White, single race	4.1	3.7–4.6
Black, single race	3.6	2.7–4.4

¹Estimates are sex adjusted to the 2000 projected U.S. standard population.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

Data for figure 15.4. Prevalence of current asthma among persons of all ages: United States, 2001–2005

Year	Percent (95% confidence interval)	
	Crude ¹	Age-adjusted ²
2001	7.6 (7.3–7.9)	7.6 (7.3–7.9)
2002	7.4 (7.1–7.7)	7.4 (7.1–7.7)
2003	7.1 (6.8–7.4)	7.1 (6.8–7.4)
2004	7.3 (7.0–7.6)	7.3 (7.0–7.6)
January–June 2005	7.9 (7.4–8.3)	7.9 (7.4–8.3)

¹Crude estimates are presented in the figure.

²Estimates are age adjusted to the 2000 projected U.S. standard population using three age groups: 0–14 years, 15–34 years, and 35 years and over.

NOTES: Beginning with the 2003 data, the National Health Interview Survey (NHIS) transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–02 were recalculated using weights derived from the 2000 census. See “About This Release” and Table III in the Appendix for more details.

DATA SOURCE: NHIS, 2001–05. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

Data table for figure 15.5. Prevalence of current asthma among persons of all ages, by age group and sex: United States, January–June 2005

Age and sex	Percent	95% confidence interval
0–14 years		
Total	9.4	8.4–10.4
Male	11.3	9.7–12.8
Female	7.5	6.4–8.7
15–34 years		
Total	7.6	6.7–8.5
Male	5.5	4.4–6.6
Female	9.7	8.4–11.1
35 years and over		
Total	7.4	6.8–8.0
Male	5.2	4.5–6.0
Female	9.4	8.5–10.2
All ages: crude¹		
Total	7.9	7.4–8.3
Male	6.6	6.0–7.2
Female	9.1	8.4–9.7
All ages: age-adjusted²		
Total	7.9	7.4–8.3
Male	6.6	6.0–7.2
Female	9.1	8.4–9.7

¹Crude estimates are presented in the figure.

²Estimates are age adjusted to the 2000 projected U.S. standard population using three age groups: 0–14 years, 15–34 years, and 35 years and over.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

Data table for figure 15.6. Sex-adjusted prevalence of current asthma among persons of all ages, by age group and race/ethnicity: United States, January–June 2005

Age and race/ethnicity	Percent¹	95% confidence interval
0–14 years		
Hispanic or Latino	9.3	7.5–11.0
Not Hispanic or Latino:		
White, single race	8.1	6.8–9.3
Black, single race	13.2	10.1–16.4
15 years and over		
Hispanic or Latino	5.8	4.8–6.8
Not Hispanic or Latino:		
White, single race	7.6	7.0–8.2
Black, single race	8.4	7.1–9.7

¹Estimates are sex adjusted to the 2000 projected U.S. standard population.

DATA SOURCE: National Health Interview Survey, 2005. Data are based on household interviews of a sample of the civilian noninstitutionalized population.