


Figure 5.1. Percentage of adults aged 65 years and over who had ever received a pneumococcal vaccination: United States, 1997–June 2007




NOTES: The analyses excluded those with unknown pneumococcal vaccination status (about 3% of respondents each year). Beginning with the 2003 data, the National Health Interview Survey (NHIS) transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–2002 were recalculated using weights derived from the 2000 census. See “About This Early Release” and Table III in the Appendix for more details.

DATA SOURCE: Sample Adult Core component of the 1997–2007 NHIS. The estimate for 2007 was based on data collected from January through June. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- For the period January through June 2007, the percentage of adults aged 65 years and over who had ever received a pneumococcal vaccination was 57.8% (95% confidence interval = 55.26–60.27%), which was not significantly different from the 2006 estimate of 57.1%.
- The percentage of adults aged 65 years and over who had ever received a pneumococcal vaccination increased from 42.4% in 1997 to 57.8% for the period January through June 2007.

Figure 5.2. Percentage of adults aged 65 years and over who had ever received a pneumococcal vaccination, by age group and sex: United States, January–June 2007




NOTE: The analyses excluded 130 adults (5.2%) aged 65 years and over with unknown pneumococcal vaccination status.

DATA SOURCE: Based on data collected from January through June in the Sample Adult Core component of the 2007 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- The percentage of adults who had ever received a pneumococcal vaccination was 50.8% for persons aged 65–74 years and 65.6% for persons aged 75 years and over.
- For both age groups, there was no significant difference in the percentage of adults who have ever received a pneumococcal vaccination between men and women.

Figure 5.3. Percentage of adults aged 65 years and over who had ever received a pneumococcal vaccination, by race/ethnicity: United States, January–June 2007



NOTE: The analyses excluded 130 adults (5.2%) aged 65 years and over with unknown pneumococcal vaccination status.

DATA SOURCE: Based on data collected from January through June in the Sample Adult Core component of the 2007 National Health Interview Survey. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- The percentage of adults aged 65 years and over who had ever received a pneumococcal vaccination was 34.3% for Hispanic persons, 62.2% for non-Hispanic white persons, and 47.0% for non-Hispanic black persons.
- Hispanic persons were less likely than non-Hispanic white persons and non-Hispanic black persons to have ever received a pneumococcal vaccination.

Data tables for Figures 5.1–5.3:

Data table for Figure 5.1. Percentage of adults aged 65 years and over who had ever received a pneumococcal vaccination: United States, 1997–June 2007

Year	Percent (95% confidence interval)	
	Crude ¹	Age-adjusted ²
1997	42.4 (40.9-43.9)	42.6 (41.1-44.1)
1998	46.0 (44.5-47.5)	46.3 (44.8-47.8)
1999	49.7 (48.1-51.3)	49.9 (48.3-51.5)
2000	53.1 (51.5-54.7)	53.4 (51.8-55.0)
2001	54.0 (52.4-55.6)	54.2 (52.6-55.7)
2002	56.0 (54.5-57.5)	56.2 (54.8-57.7)
2003	55.6 (54.1-57.1)	55.7 (54.2-57.1)
2004	56.8 (55.3-58.3)	57.0 (55.5-58.5)
2005	56.2 (54.82-57.64)	56.3 (54.95-57.72)
2006	57.1 (55.25-58.88)	57.2 (55.43-59.04)
January–June 2007	57.8 (55.26-60.27)	57.9 (55.39-60.33)

¹Crude estimates are presented in the figure.

²Estimates for this Healthy People 2010 Leading Health Indicator are age adjusted using the projected 2000 U.S. population as the standard population and using two age groups: 65–74 years and 75 years and over.

NOTES: Beginning with the 2003 data, the National Health Interview Survey (NHIS) transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–2002 were recalculated using weights derived from the 2000 census. See “About This Early Release” and Table III in the Appendix for more details.

DATA SOURCE: NHIS, 1997–June 2007. Data are based on household interviews of a sample of the civilian noninstitutionalized population.



Data table for Figure 5.2. Percentage of adults aged 65 years and over who had ever received a pneumococcal vaccination, by age group and sex: United States, January–June 2007

Age and sex	Percent	95% confidence interval
65–74 years		
Total	50.8	47.41-54.18
Male	47.2	42.55-51.83
Female	53.8	49.01-58.57
75 years and over		
Total	65.6	62.35-68.84
Male	67.0	61.88-72.06
Female	64.7	60.53-68.89
65 years and over: crude¹		
Total	57.8	55.26-60.27
Male	55.8	52.09-59.50
Female	59.2	55.79-62.66
65 years and over: age-adjusted²		
Total	57.9	55.39-60.33
Male	56.6	53.06-60.21
Female	59.0	55.57-62.44

¹Crude estimates are presented in the figure.

²Estimates for this Healthy People 2010 Leading Health Indicator are age adjusted using the projected 2000 U.S. population as the standard population and using two age groups: 65–74 years and 75 years and over.

DATA SOURCE: National Health Interview Survey, January–June 2007. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

Data table for Figure 5.3. Percentage of adults aged 65 years and over who had ever received a pneumococcal vaccination, by race/ethnicity: United States, January–June 2007

Race/ethnicity	Percent (95% confidence interval)	
	Crude ¹	Age-adjusted ²
Hispanic or Latino	34.3 (27.06-41.50)	34.8 (27.16-42.34)
Not Hispanic or Latino:		
White, single race	62.2 (59.45-64.96)	62.0 (59.29-64.73)
Black, single race	47.0 (40.69-53.26)	47.8 (41.15-54.40)

¹Crude estimates are presented in the figure.

²Estimates for this Healthy People 2010 Leading Health Indicator are age adjusted using the projected 2000 U.S. population as the standard population and using two age groups: 65–74 years and 75 years and over.

DATA SOURCE: National Health Interview Survey, January–June 2007. Data are based on household interviews of a sample of the civilian noninstitutionalized population.