


Figure 12.1. Percentage of adults aged 65 years and over who need help with personal care from other persons: United States, 1997-June 2009




NOTES: Personal care needs, or activities of daily living (ADLs), include eating, bathing, dressing, or getting around inside the person’s home. The analyses excluded persons with unknown information on personal care needs (about 0.0% of respondents each year). Beginning with the 2003 data, the National Health Interview Survey transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000-2002 were recalculated using weights derived from the 2000 census. See “About This Early Release” for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, 1997-June 2009, Family Core component. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- For the period January through June 2009, 6.3% (95% confidence interval = 5.47%-7.21%) of adults aged 65 years and over needed help with personal care from other persons. This estimate was not significantly different from the 2008 estimate of 6.4%.
- The annual percentage of adults who needed help with personal care from other persons ranged from 6.1% in 2006 to 6.9% in 2007.

Figure 12.2. Percentage of adults aged 65 years and over who need help with personal care from other persons, by age group and sex: United States, January-June 2009




NOTES: Personal care needs, or activities of daily living (ADLs), include eating, bathing, dressing, or getting around inside the person’s home. The analyses excluded 4 persons (0.1%) with unknown information on personal care needs.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January-June 2009, Family Core component. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- For both sexes combined, adults aged 85 years and over (19.8%) were more than five times as likely as adults aged 65-74 (3.5%) to need help with personal care from other persons.
- For adults aged 75-84 years and 85 and over, women were more likely than men to need help with personal care.

Figure 12.3. Age-sex-adjusted percentage of adults aged 65 years and over who need help with personal care from other persons, by race/ethnicity: United States, January-June 2009



NOTES: Personal care needs, or activities of daily living (ADLs), include eating, bathing, dressing, or getting around inside the person’s home. The analyses excluded 4 persons (0.1%) with unknown information on personal care needs. Estimates are age-sex adjusted using the projected 2000 U.S. population as the standard population and using three age groups: 65-74 years, 75-84 years, and 85 years and over.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January-June 2009, Family Core component. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- The age-sex-adjusted percentage of persons who needed help with personal care from other persons was 10.0% for Hispanic persons, 5.6% for non-Hispanic white persons, and 10.8% for non-Hispanic black persons.
- Non-Hispanic white persons were less likely to need help with personal care from other persons than Hispanic persons or non-Hispanic black persons.

Data tables for Figures 12.1-12.3:

Data table for Figure 12.1. Percentage of adults aged 65 years and over who need help with personal care from other persons: United States, 1997-June 2009

Year	Percent	95% confidence interval
1997	6.6	6.1-7.1
1998	6.3	5.9-6.8
1999	6.4	5.8-6.9
2000	6.4	5.9-6.9
2001	6.5	6.0-7.0
2002	6.2	5.7-6.7
2003	6.5	6.0-7.1
2004	6.3	5.8-6.8
2005	6.3	5.83-6.76
2006	6.1	5.44-6.69
2007	6.9	6.21-7.58
2008	6.4	5.84-7.03
Jan.-June 2009	6.3	5.47-7.21

NOTES: Beginning with the 2003 data, the National Health Interview Survey transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000-2002 were recalculated using weights derived from the 2000 census. See "About This Early Release" for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, 1997-June 2009, Family Core component. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

Data table for Figure 12.2. Percentage of adults aged 65 years and over who need help with personal care from other persons, by age group and sex: United States, January-June 2009

Age and sex	Percent	95% confidence interval
65-74 years, total	3.5	2.48-4.55
65-74 years, male	3.5	1.78-5.23
65-74 years, female	3.5	2.32-4.74
75-84 years, total	6.5	4.92-8.06
75-84 years, male	4.8	3.06-6.55
75-84 years, female	7.7	5.55-9.77
85 years and over, total	19.8	14.86-24.81
85 years and over, male	13.8	7.64-20.05
85 years and over, female	23.4	17.09-29.75
65 years and over (crude ¹), total	6.3	5.47-7.21
65 years and over (crude ¹), male	4.9	3.72-6.11
65 years and over (crude ¹), female	7.4	6.14-8.69
65 years and over (age-adjusted ²), total	6.6	5.67-7.48
65 years and over (age-adjusted ²), male	5.2	4.00-6.46
65 years and over (age-adjusted ²), female	7.4	6.18-8.69

¹Crude estimates are presented in the figure.

²Estimates are age adjusted using the projected 2000 U.S. population as the standard population and using three age groups: 65-74 years, 75-84 years, and 85 years and over.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January-June 2009, Family Core component. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

Data table for Figure 12.3. Age-sex-adjusted percentage of adults aged 65 years and over who need help with personal care from other persons, by race/ethnicity: United States, January-June 2009

Race/ethnicity	Percent ¹	95% confidence interval
Hispanic or Latino	10.0	6.62-13.40
Not Hispanic or Latino, single race, white	5.6	4.66-6.60
Not Hispanic or Latino, single race, black	10.8	6.83-14.73

¹Estimates are age-sex adjusted using the projected 2000 U.S. population as the standard population and using three age groups: 65-74 years, 75-84 years, and 85 years and over.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January-June 2009, Family Core component. Data are based on household interviews of a sample of the civilian noninstitutionalized population.