


## Receipt of pneumococcal vaccination

**Figure 5.1. Percentage of adults aged 65 and over who had ever received a pneumococcal vaccination: United States, 1997–June 2014**




NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. The analyses excluded those with unknown pneumococcal vaccination status (about 5% of respondents each year). See [Technical Notes](#) for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, 1997–June 2014, Sample Adult Core component.

- For January–June 2014, the percentage of adults aged 65 and over who had ever received a pneumococcal vaccination was 60.5% (95% confidence interval = 58.43%–62.57%), which was higher than, but not significantly different from the 2013 estimate of 59.7%.
- The percentage of adults aged 65 and over who had ever received a pneumococcal vaccination increased rapidly from 42.4% in 1997 to 53.1% in 2000, then increased more slowly to 62.3% in 2011.

**Figure 5.2. Percentage of adults aged 65 and over who had ever received a pneumococcal vaccination, by age group and sex: United States, January–June 2014**




NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. The analyses excluded the 4.0% of adults aged 65 and over with unknown pneumococcal vaccination status. See [Technical Notes](#) for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January–June 2014, Sample Adult Core component.

- For both sexes combined, the percentage of adults who had ever received a pneumococcal vaccination was 54.9% for persons aged 65–74 and 68.5% for persons aged 75 and over.
- For both sexes combined, adults aged 75 and over were more likely than those aged 65–74 to have ever received a pneumococcal vaccination. This pattern was seen in both men and women.
- For adults aged 65 and over, and for adults aged 65–74 and 75 and over, there were no differences between men and women for having ever received a pneumococcal vaccination.

**Figure 5.3. Percentage of adults aged 65 and over who had ever received a pneumococcal vaccination, by race/ethnicity: United States, January–June 2014**



NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. The analyses excluded the 4.0% of adults aged 65 and over with unknown pneumococcal vaccination status. See [Technical Notes](#) for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January–June 2014, Sample Adult Core component.

- The percentage of adults aged 65 and over who had ever received a pneumococcal vaccination was 43.6% for Hispanic persons, 64.3% for non-Hispanic white persons, and 47.5% for non-Hispanic black persons.
- Of the three race/ethnicity groups, non-Hispanic white adults were more likely to have ever received a pneumococcal vaccination compared to Hispanic and non-Hispanic black adults.

## Data tables for Figures 5.1–5.3:

## Data table for Figure 5.1. Percentage of adults aged 65 and over who had ever received a pneumococcal vaccination: United States, 1997–June 2014

Year	Crude <sup>1</sup> percent (95% confidence interval)	Age-adjusted <sup>2</sup> percent (95% confidence interval)
1997	42.4 (40.9-43.9)	42.6 (41.1-44.1)
1998	46.0 (44.5-47.5)	46.3 (44.8-47.8)
1999	49.7 (48.1-51.3)	49.9 (48.3-51.5)
2000	53.1 (51.5-54.7)	53.4 (51.8-55.0)
2001	54.0 (52.4-55.6)	54.2 (52.6-55.7)
2002	56.0 (54.5-57.5)	56.2 (54.8-57.7)
2003	55.6 (54.1-57.1)	55.7 (54.2-57.1)
2004	56.8 (55.3-58.3)	57.0 (55.5-58.5)
2005	56.2 (54.82-57.64)	56.3 (54.95-57.72)
2006	57.1 (55.25-58.88)	57.2 (55.43-59.04)
2007	57.7 (55.68-59.69)	57.8 (55.78-59.80)
2008	59.6 (57.68-61.48)	59.8 (57.98-61.64)
2009	60.6 (58.74-62.51)	61.0 (59.15-62.79)
2010	59.4 (57.61-61.17)	59.8 (58.02-61.51)
2011	62.3 (60.55-63.95)	62.7 (61.04-64.31)
2012	59.9 (58.29-61.49)	60.4 (58.86-62.01)
2013	59.7 (58.02-61.36)	60.4 (58.77-62.05)
January–June 2014	60.5 (58.43-62.57)	61.4 (59.35-63.50)

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates are age adjusted using the projected 2000 U.S. population as the standard population and using two age groups: 65–74 and 75 and over.

NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. Beginning with the 2012 data, the National Health Interview Survey transitioned to weights derived from the 2010 census. For 2003–2011 data, weights were derived from the 2000 census. In this Early Release, estimates for 2000–2002 were recalculated using weights derived from the 2000 census. For 1997–1999 data, weights were derived from the 1990 census. See [Technical Notes](#) for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, 1997–June 2014, Sample Adult Core component.

**Data table for Figure 5.2. Percentage of adults aged 65 and over who had ever received a pneumococcal vaccination, by age group and sex: United States, January–June 2014**

Age (years) and sex	Percent	95% confidence interval
65–74 , total	54.9	52.32-57.50
65–74 , male	52.7	48.41-56.94
65–74 , female	56.8	53.18-60.45
75 and over, total	68.5	65.38-71.71
75 and over, male	67.2	62.18-72.13
75 and over, female	69.5	65.58-73.43
65 and over (crude <sup>1</sup> ), total	60.5	58.43-62.57
65 and over (crude <sup>1</sup> ), male	58.2	54.85-61.53
65 and over (crude <sup>1</sup> ), female	62.3	59.68-64.94
65 and over (age-adjusted <sup>2</sup> ), total	61.4	59.35-63.50
65 and over (age-adjusted <sup>2</sup> ), male	59.6	56.33-62.85
65 and over (age-adjusted <sup>2</sup> ), female	62.9	60.26-65.49

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates are age adjusted using the projected 2000 U.S. population as the standard population and using two age groups: 65–74 and 75 and over.

NOTE: Data are based on household interviews of a sample of the civilian noninstitutionalized population.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January–June 2014, Sample Adult Core component.

**Data table for Figure 5.3. Percentage of adults aged 65 and over who had ever received a pneumococcal vaccination, by race/ethnicity: United States, January–June 2014**

Race/ethnicity	Crude <sup>1</sup> percent (95% confidence interval)	Age-adjusted <sup>2</sup> percent (95% confidence interval)
Hispanic or Latino	43.6 (35.87-51.32)	44.2 (36.19-52.24)
Not Hispanic or Latino, single race, white	64.3 (61.88-66.67)	65.0 (62.64-67.42)
Not Hispanic or Latino, single race, black	47.5 (41.98-52.98)	49.8 (44.41-55.28)

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates are age adjusted using the projected 2000 U.S. population as the standard population and using two age groups: 65–74 and 75 and over.

NOTE: Data are based on household interviews of a sample of the civilian noninstitutionalized population.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January–June 2014, Sample Adult Core component.