


## Human immunodeficiency virus testing

**Figure 10.1. Percentage of adults aged 18 and over who had ever been tested for HIV: United States, 1997–June 2017**



† The AIDS Knowledge and Attitudes section of the National Health Interview Survey (NHIS) was dropped in 2011; only the human immunodeficiency virus (HIV) testing question was retained, and it was moved to the Adult Access to Health Care and Utilization section of the Sample Adult questionnaire. HIV testing estimates based on 2011–2012 NHIS (dashed line) are not comparable with those from 2010 and earlier, or 2013 and later. Differences observed in estimates based on 2010 and earlier NHIS, 2011–2012 NHIS, and 2013 and later NHIS may be partially or fully attributable to these changes in placement of the HIV testing question in the NHIS questionnaire.


†† In 2013, the HIV testing question was moved from the Adult Access to Care and Utilization section of the Sample Adult questionnaire to the Adult Selected Items section of the Sample Adult questionnaire. HIV testing estimates based on 2013 and later NHIS (dotted line) are not comparable with those from 2012 and earlier. Differences observed in estimates based on 2012 and earlier NHIS and on 2013 and later NHIS may be partially or fully attributable to these changes in placement of the HIV testing question in the NHIS questionnaire.

NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. Persons who received HIV testing solely as a result of blood donation were considered not to have been tested for HIV. The analyses exclude those with unknown HIV test status (about 5% of respondents each year). See [Technical Notes](#) for more details.

DATA SOURCE: NCHS, National Health Interview Survey, 1997–June 2017, Sample Adult Core component.

- For January–June 2017, the percentage of adults who had ever been tested for HIV was 40.8% (95% confidence interval = 39.56%–42.09%), which was not statistically different from the 2016 estimate of 41.1%.
- The percentage of adults who had ever had an HIV test increased, from 31.8% in 1997 to 39.5% in 2010.
- From 2013 through January–June 2017, the percentage of adults who had ever been tested for HIV increased, from 37.3% to 40.8%.

**Figure 10.2. Percentage of adults aged 18 and over who had ever been tested for HIV, by age group and sex: United States, January–June 2017**




NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. Persons who received human immunodeficiency virus (HIV) testing solely as a result of blood donation were considered not to have been tested for HIV. The AIDS Knowledge and Attitudes section of the National Health Interview Survey (NHIS) was dropped in 2011; only the HIV testing question was retained, and it was moved to the Adult Access to Health Care and Utilization section of the Sample Adult questionnaire. In 2013, the HIV testing question was moved again to the Adult Selected Items section of the Sample Adult questionnaire and is not comparable with 2011–2012. Differences observed in estimates based on 2010 and earlier NHIS, 2011–2012 NHIS, and 2013 and later NHIS may be partially or fully attributable to these changes in placement of the HIV testing question in the NHIS questionnaire. The analyses exclude the 5.4% of adults with unknown HIV test status. See [Technical Notes](#) for more details.

DATA SOURCE: NCHS, National Health Interview Survey, January–June 2017, Sample Adult Core component.

- For both sexes combined, the percentage of adults by age group who ever had an HIV test was highest among adults aged 25–34 (53.1%) and 35–44 (56.8%) and lowest among adults aged 65 and over (19.2%). This pattern held for males and females.
- For adults aged 18 and over, and for age groups 18–24, 25–34, and 35–44, women were more likely than men to have ever had an HIV test.

**Figure 10.3. Age-sex-adjusted percentage of adults aged 18 and over who had ever been tested for HIV, by race and ethnicity: United States, January–June 2017**



NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. Persons who received human immunodeficiency virus (HIV) testing solely as a result of blood donation were considered not to have been tested for HIV. The AIDS Knowledge and Attitudes section of the National Health Interview Survey (NHIS) was dropped in 2011; only the HIV testing question was retained, and it was moved to the Adult Access to Health Care and Utilization section of the Sample Adult questionnaire. In 2013, the HIV testing question was moved again to the Adult Selected Items section of the Sample Adult questionnaire and is not comparable with 2011–2012. Differences observed in estimates based on 2010 and earlier NHIS, 2011–2012 NHIS, and 2013 and later NHIS may be partially or fully attributable to these changes in placement of the HIV testing question in the NHIS questionnaire. The analyses exclude the 5.4% of adults with unknown HIV test status. See [Technical Notes](#) for more details.

DATA SOURCE: NCHS, National Health Interview Survey, January–June 2017, Sample Adult Core component.

- The age-sex-adjusted percentage of adults by race and ethnicity who ever had an HIV test was 47.8% for Hispanic adults, 38.5% for non-Hispanic white adults, and 60.8% for non-Hispanic black adults.
- Of the three race and ethnicity groups, non-Hispanic black adults were more likely to have ever had an HIV test compared with Hispanic and non-Hispanic white adults. Non-Hispanic white adults were less likely to have ever had an HIV test compared with Hispanic adults.

## Data tables for Figures 10.1–10.3:

## Data table for Figure 10.1. Percentage of adults aged 18 and over who had ever been tested for HIV: United States, 1997– June 2017

| Year              | Crude <sup>1</sup> percent (95% confidence interval) | Age-adjusted <sup>2</sup> percent (95% confidence interval) |
|-------------------|--|---|
| 1997              | 31.8 (31.1–32.5)                                     | 31.0 (30.4–31.6)  |
| 1998              | 30.2 (29.5–30.9)                                     | 29.6 (28.9–30.2)  |
| 1999              | 32.3 (31.6–33.0)                                     | 31.8 (31.1–32.5)  |
| 2000              | 32.1 (31.4–32.8)                                     | 31.8 (31.2–32.5)  |
| 2001              | 33.2 (32.5–34.0)                                     | 33.1 (32.4–33.7)  |
| 2002              | 34.5 (33.8–35.2)                                     | 34.5 (33.8–35.1)  |
| 2003              | 35.4 (34.7–36.1)                                     | 35.3 (34.7–36.0)  |
| 2004              | 34.6 (33.9–35.3)                                     | 34.8 (34.1–35.5)  |
| 2005              | 35.3 (34.58–36.01)                                   | 35.4 (34.73–36.10)  |
| 2006              | 35.8 (34.94–36.70)                                   | 36.1 (35.25–36.92)  |
| 2007              | 36.6 (35.59–37.62)                                   | 37.0 (35.28–38.79)  |
| 2008              | 38.7 (37.73–39.59)                                   | 39.2 (38.40–40.09)  |
| 2009              | 39.8 (38.70–40.84)                                   | 40.4 (39.36–41.40)  |
| 2010              | 39.5 (38.65–40.41)                                   | 40.3 (39.52–41.15)  |
| 2011 <sup>3</sup> | 35.9 (35.14–36.72)                                   | 36.7 (35.97–37.51)  |
| 2012              | 34.8 (34.04–35.64)                                   | 35.8 (34.98–36.62)  |
| 2013 <sup>3</sup> | 37.3 (36.33–38.20)                                   | 38.3 (37.42–39.25)  |
| 2014              | 37.5 (36.66–38.41)                                   | 38.8 (37.92–39.71)  |
| 2015              | 38.6 (37.66–39.45)                                   | 39.8 (38.95–40.73)  |
| 2016              | 41.1 (40.18–42.10)                                   | 42.4 (41.47–43.32)  |
| January–June 2017 | 40.8 (39.56–42.09)                                   | 42.2 (40.92–43.53)  |

<sup>1</sup>Crude estimates are presented.

<sup>2</sup>Estimates are age-adjusted using the projected 2000 U.S. population as the standard population and five age groups: 18–24, 25–34, 35–44, 45–64, and 65 and over.

<sup>3</sup>The AIDS Knowledge and Attitudes section of the National Health Interview Survey (NHIS) was dropped in 2011; only the HIV testing question was retained, and it was moved to the Adult Access to Health Care and Utilization section of the Sample Adult questionnaire. In 2013, the HIV testing question was moved again to the Adult Selected Items section of the Sample Adult questionnaire. HIV testing estimates from these three periods (2010 and earlier, 2011–2012, and 2013 and later) are not comparable; differences observed in estimates may be partially or fully attributable to these changes in placement of the HIV testing question in the NHIS questionnaire.

NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. Beginning with 2012 data, NHIS transitioned to weights derived from the 2010 census. For 2003–2011 data, weights were derived from the 2000 census. In this Early Release, estimates for 2000–2002 were recalculated using weights derived from the 2000 census. For 1997–1999 data, weights were derived from the 1990 census. See [Technical Notes](#) for more details.

DATA SOURCE: NCHS, National Health Interview Survey, 1997–June 2017, Sample Adult Core component.

**Data table for Figure 10.2. Percentage of adults aged 18 and over who had ever been tested for HIV, by age group and sex: United States, January–June 2017**

| Age (years) and sex                              | Percent | 95% confidence interval |
|--|---------|-------------------------|
| 18–24, total                                     | 32.5    | 28.55–36.74             |
| 18–24, male                                      | 24.6    | 18.47–31.56             |
| 18–24, female                                    | 40.7    | 36.68–44.85             |
| 25–34, total                                     | 53.1    | 50.28–55.82             |
| 25–34, male                                      | 45.6    | 40.85–50.36             |
| 25–34, female                                    | 60.5    | 56.72–64.18             |
| 35–44, total                                     | 56.8    | 53.87–59.62             |
| 35–44, male                                      | 51.6    | 46.84–56.25             |
| 35–44, female                                    | 61.8    | 57.29–66.26             |
| 45–64, total                                     | 42.2    | 40.30–44.16             |
| 45–64, male                                      | 40.8    | 38.21–43.48             |
| 45–64, female                                    | 43.5    | 40.58–46.50             |
| 65 and over, total                               | 19.2    | 17.13–21.41             |
| 65 and over, male                                | 21.0    | 18.25–24.03             |
| 65 and over, female                              | 17.7    | 15.60–19.97             |
| 18–64, total                                     | 46.1    | 44.75–47.52             |
| 18–64, male                                      | 41.6    | 39.27–43.89             |
| 18–64, female                                    | 50.6    | 48.28–52.83             |
| 18 and over (crude <sup>1</sup> ), total         | 40.8    | 39.56–42.09             |
| 18 and over (crude <sup>1</sup> ), male          | 37.8    | 35.78–39.85             |
| 18 and over (crude <sup>1</sup> ), female        | 43.7    | 41.77–45.54             |
| 18 and over (age-adjusted <sup>2</sup> ), total  | 42.2    | 40.92–43.53             |
| 18 and over (age-adjusted <sup>2</sup> ), male   | 38.6    | 36.55–40.64             |
| 18 and over (age-adjusted <sup>2</sup> ), female | 45.9    | 43.83–47.94             |

<sup>1</sup>Crude estimates are presented.

<sup>2</sup>Estimates are age-adjusted using the projected 2000 U.S. population as the standard population and five age groups: 18–24, 25–34, 35–44, 45–64, and 65 and over.

NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population.

DATA SOURCE: NCHS, National Health Interview Survey, January–June 2017, Sample Adult Core component.

**Data table for Figure 10.3. Age-sex-adjusted percentage of adults aged 18 and over who had ever been tested for HIV, by race and ethnicity: United States, January–June 2017**

| Race and ethnicity                         | Percent <sup>1</sup> | 95% confidence interval |
|--|----------------------|-------------------------|
| Hispanic or Latino                         | 47.8                 | 44.84–50.72             |
| Not Hispanic or Latino, single race, white | 38.5                 | 36.86–40.20             |
| Not Hispanic or Latino, single race, black | 60.8                 | 57.31–64.26             |

<sup>1</sup>Estimates are age-sex-adjusted using the projected 2000 U.S. population as the standard population and five age groups: 18–24, 25–34, 35–44, 45–64, and 65 and over.

NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population.

DATA SOURCE: NCHS, National Health Interview Survey, January–June 2017, Sample Adult Core component.