



NATIONAL CENTER FOR HEALTH STATISTICS

Preferred Reporting Items for Complex Sample Survey Analysis

Round 8: Data collected February–March 2026



Last revised August 27, 2026

Persons with disabilities experiencing problems accessing PDF files
should contact nchsed@cdc.gov or call 301-458-4688



Table of contents

Overview.....3

PRICSSA-related Content, in Brief3

Table 1: PRICSSA-related Content, in Detail4

Other Resources.....7

Suggested citation7

Overview

In the *Journal of Survey Statistics and Methodology*, Seidenberg, Moser, and West (2023) proposed an itemized checklist to guide researchers publishing analyses using complex sample survey data. This checklist—the Preferred Reporting Items for Complex Sample Survey Analysis (PRICSSA)—is intended to help eliminate analytic and reporting errors and increase transparency and reproducibility. NCHS is providing the following information to support researchers publishing analyses using Rapid Surveys System Round 8 (RSS-8) data.

Additional information about data collection methods, round-specific content, and other useful resources for the RSS-8 is available in the *RSS-8 Survey Description* document, at: www.cdc.gov/nchs/data/rss/round8/RSS8-SurveyDescription.pdf and the *RSS-8 Quality Profile* at www.cdc.gov/nchs/data/rss/round8/rss8-quality-profile.pdf.

PRICSSA-related Content, in Brief

- Name and wave of survey: NCHS Rapid Surveys System Round 8 (RSS-8)
- Data collection mode: Two commercially available probability-based online panels—NORC at the University of Chicago’s (NORC’s) AmeriSpeak panel and Ipsos’s KnowledgePanel—with some telephone supplementation for the AmeriSpeak panel
- Dates of data collection: February 26–March 23, 2026 (February 26–March 23, 2026, for the AmeriSpeak Panel and February 27–March 14, 2026, for the KnowledgePanel).
- Target population: Adult civilian noninstitutionalized population age 18 and older residing in the 50 U.S. states and Washington, DC
- Populations excluded: Persons in long-term care and correctional facilities, persons with no fixed household address
- Design: Stratified cluster sample (AmeriSpeak Panel) and probability proportional to size sampling (KnowledgePanel)
- Variance estimation: Taylor Series Linearization
- Weight and design variables
 - Weight: WEIGHT
 - PSU: P_PSU_R
 - Stratum: P_STRATA_R

- Number of completed interviews: 8,026 adults (2,022 adults for the AmeriSpeak Panel and 6,004 adults for the KnowledgePanel)
- Unweighted survey completion rate: 64.0% (51.0% for the AmeriSpeak Panel and 70.0% for the KnowledgePanel)
- Unweighted final cumulative response rates, incorporating panel recruitment and retention rates: 13.8% (AmeriSpeak Panel) and 3.9% (KnowledgePanel). For more information on calculation of final response rates, please see www.cdc.gov/nchs/data/rss/round8/rss8-quality-profile.pdf.

Unlike the first two rounds of RSS, only one sample was drawn from each panel for RSS-8.

Table 1: PRICSSA-related Content, in Detail

Rapid Surveys System Round 8 (RSS-8)	
1.1 Data collection dates	The RSS-8 interviews were conducted between February 26–March 23, 2026 (February 26–March 23, 2026, for the AmeriSpeak Panel and February 27–March 14, 2026, for the KnowledgePanel).
1.2 Data collection mode(s)	Interviews for the KnowledgePanel were conducted solely by web. Interviews for AmeriSpeak were also primarily web-based; however, telephone interviews were used for panelists known to prefer completing surveys over the phone.
1.3 Target population	The target population for RSS-8 is all civilian noninstitutionalized U.S. adults age 18 and older residing in the 50 U.S. states and Washington, DC. Persons in long-term care and correctional facilities and persons with no fixed household address are excluded.
1.4 Sample design	RSS uses two online panels: Ipsos KnowledgePanel and NORC AmeriSpeak. These panels are based on probability samples of the population of U.S. households and are designed to serve as sampling frames for sample selection and production of national estimates. All members of both panels complete questions about demographics and household composition before participating in any surveys. RSS-8 only includes panelists age 18 or older in the sample.

	8,582 KnowledgePanel panelists were sampled for an expected 6,000 completes, and 3,964 AmeriSpeak panelists were sampled for an expected 2,000 completes.
1.5 Survey response rate(s)	8,582 KnowledgePanel panelists were sampled for final yield of 6,004 completes, and 3,964 AmeriSpeak panelists were sampled for a final yield of 2,022 completes, leading to completion rates of 70.0% and 51.0% respectively. Final response rates that reflect not only the survey completion rate but also the panel recruitment and retention rates are 3.9% for KnowledgePanel and 13.8% for AmeriSpeak.
2.1 Missingness rates	For most measures in RSS-8, percentages with unknown values are typically small. Of the 241 survey items, nearly half had an item nonresponse rate of less than 1% in the combined file (49.0%). The combined file had 19 questions with item nonresponse greater than or equal to 10%. Variables were imputed by the panel providers for their internal weighting procedures and in post-processing for weighting to the National Health Interview Survey (NHIS). The percent of values imputed ranged from 0.0% to 4.9%. Imputation flags can be used to identify imputed values in the data file.
2.2 Observation deletion	Cases were removed if the survey was completed in less than a minimum time or skipped more than half of the eligible questions. Overall, 1.5% of RSS-8 records were removed due to speeders or respondents with high refusal rates.
2.3 Sample sizes	Numbers of completed interviews for RSS-8 are as follows: 8,026 adults (2,022 adults for the AmeriSpeak Panel and 6,004 adults for the KnowledgePanel).
2.4 Confidence intervals and standard errors	NCHS recommends using two-sided 95% confidence intervals calculated using the Clopper-Pearson method adapted for complex surveys by Korn and Graubard. Standard errors used in this calculation should be obtained using software (e.g., SUDAAN) that takes into account the complex sampling design of RSS. NCHS uses the Taylor series linearization method for variance estimation.
2.5 Weighting	The composite calibrated weight, WEIGHT, should be used to generate national estimates based on both panel providers. This

	weight accounts for sampling probabilities and nonresponse and is calibrated to control totals from the NHIS.
2.6 Variance estimation	Users of the public-use and restricted-use files should use the Taylor series linearization method to estimate variances. Sample design variables are provided in the data files for this purpose. This method requires the use of statistical software with this functionality. Analysts who instead apply simple random sampling techniques to RSS-8 data generally will produce standard error estimates that are, on average, too small and are likely to produce results that are subject to excessive Type I error.
2.7 Subpopulation analysis	To compute accurate standard errors, NCHS recommends that subpopulation analyses be carried out using the full data file instead of subsetting the data file to the subpopulation of interest. This can be done with the SUBPOPN statement in SUDAAN or an equivalent procedure with another software package that supports complex design variance estimation.
2.8 Suppression rules	NCHS recommends suppressing percentages that do not meet the criteria specified in National Center for Health Statistics Data Presentation Standards for Proportions . These standards include thresholds for sample size, absolute and relative confidence interval width, and degrees of freedom.
2.9 Software and code	To appropriately analyze RSS-8 data, it is necessary to utilize weights and variance estimation variables and software that can appropriately analyze complex samples. This includes, but is not limited to, SUDAAN, Stata, SPSS, SAS, and R software packages.
2.10 Singleton problem	Taylor Series Linearization requires at least two PSUs per stratum for variance estimation. NCHS uses the MISSUNIT option in SUDAAN to account for the presence of strata with only one PSU.
2.11 Public/restricted data	For data users and researchers throughout the world, public-use data files and codebooks are freely available on the internet. Analysts interested in working with RSS data that were suppressed or edited to protect confidentiality may apply to access unmodified restricted-use data files through the NCHS Research Data Center (RDC). Codebooks listing variables (with no frequencies) available on restricted-use files are available on the RSS website.
2.12 Embedded experiments	There were no embedded experiments in RSS-8.

Other Resources

All information in this document has been summarized from the *NCHS Rapid Surveys System Round 8 Survey Description* and the *Quality Profile*:

National Center for Health Statistics. Quality Profile, Rapid Surveys System Round 8. Hyattsville, Maryland. September 2026. Available from:
www.cdc.gov/nchs/data/rss/round8/rss8-quality-profile.pdf.

National Center for Health Statistics. Survey Description, Rapid Surveys System Round 8. Hyattsville, Maryland. September 2026. Available from:
www.cdc.gov/nchs/data/rss/round8/RSS8-SurveyDescription.pdf.

Additional information may also be found in the following publication:

Parker JD, Talih M, Malec DJ, et al. National Center for Health Statistics data presentation standards for proportions. National Center for Health Statistics. Vital Health Stat 2(175). 2017. Available from:
https://www.cdc.gov/nchs/data/series/sr_02/sr02_175.pdf.

Suggested citation

Preferred Reporting Items for Complex Sample Survey. NCHS Rapid Surveys System. Round 8. September 2026. National Center for Health Statistics. Available from: www.cdc.gov/nchs/data/rss/round7/pricssa.pdf.