

Acute Conditions

incidence and associated disability

United States-July 1963 - June 1964

U. S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE Public Health Service





VITAL and HEALTH STATISTICS DATA FROM THE NATIONAL HEALTH SURVEY

Acute Conditions

incidence and associated disability United States-July 1963 - June 1964

Statistics on the incidence of acute conditions and the associated days of restricted activity, bed disability, and time lost from work and school, by age, sex, calendar quarter, residence, and geographic region. Based on data collected in household interviews during the period July 1963–June 1964.

Washington, D.C.

April 1965

U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE Anthony J. Celebrezze Secretary

Public Health Service Luther L. Terry Surgeon General

NATIONAL CENTER FOR HEALTH STATISTICS

FORREST E. LINDER, Ph. D., Director
THEODORE D. WOOLSEY, Deputy Director
O. K. SAGEN, Ph. D., Assistant Director
WALT R. SIMMONS, M.A., Statistical Advisor
ALICE M. WATERHOUSE, M.D., Medical Advisor
JAMES E. KELLY, D.D.S., Dental Advisor
LOUIS R. STOLCIS, M.A., Executive Officer

DIVISION OF HEALTH INTERVIEW STATISTICS

PHILIP S. LAWRENCE, Sc. D., Chief ELIJAH L. WHITE, Assistant Chief

COOPERATION OF THE BUREAU OF THE CENSUS

Under the legislation establishing the National Health Survey, the Public Health Service is authorized to use, insofar as possible, the services or facilities of other Federal, State, or private agencies.

In accordance with specifications established by the National Health Survey, the Bureau of the Census, under a contractual arrangement, participates in most aspects of survey planning, selects the sample, collects the data, and carries out certain parts of the statistical processing.

CONTENTS

	Page
Selected Findings	1
Source of Data	2
Incidence of Acute Conditions and Associated Days of Disability	
Condition Group, Sex, and Age	3
Geographic Distribution	5
Seasonal Variation: Calendar Quarter	8
Detailed Tables	9
Appendix I. Technical Notes on Methods	45
Background of This Report	45
Statistical Design of the Health Interview Survey	45
General Qualifications	46
Reliability of Estimates	46
Appendix II. Definitions of Certain Terms Used in This Report	48
Terms Relating to Conditions	48
Terms Relating to Disability	49
Demographic Terms	49

ACUTE CONDITIONS

INCIDENCE AND ASSOCIATED DISABILITY

Charles S. Wilder, Division of Health Interview Statistics

SELECTED FINDINGS

The incidence of acute respiratory illnesses which required medical attention or reduction of daily activity was substantially lower during the period July 1963-June 1964 than during the preceding year July 1962-June 1963. An estimated 204.3 million acute respiratory illnesses were reported in household interviews conducted by the Health Interview Survey, National Center for Health Statistics, during the more recent 12 months, or 1.1 cases per person per year. This estimate represents a reduction of about 28.6 million cases from that reported in the earlier period. A decline in incidence of illnesses reported as influenza, amounting to 21.5 million cases, was the principal cause of the drop.

During the year ending June 1964 the civilian, noninstitutional population of the United States experienced a total incidence of 387.4 million acute illnesses or injuries causing the persons concerned to seek medical attention or to restrict usual daily activities, a rate of 2.1 conditions per person per year. This estimate represents less of a decline in incidence than that noted above for respiratory illnesses alone because of an increased number of injuries and common childhood diseases. There were approximately 4.5 million more injuries and 9.9 million more common childhood diseases during the current year than had been reported in the previous period. Epidemics of measles and rubella (German

measles) occurring during the spring of 1964 contributed substantially to the rise in incidence of common childhood diseases.

During July 1963-June 1964, acute illnesses or injury caused an average of 8.1 days of restricted activity per person in the population, including 3.5 days spent in bed. The rate of disability was higher among females than among males, with the average female having 1.5 more days of reduced activity and 0.8 of a day more of bed disability per person. School children between the ages 6-16 years experienced 8.6 days of restricted activity, including 3.7 days in bed. Absence from school amounted to 4.4 days per child as a result of acute illness or injury.

Currently employed persons aged 17 years and older experienced 1.6 acute conditions per person and 3.3 days of work loss associated with these acute conditions. A currently employed person is defined as a person who had a job or business or worked at any time during the 2-week period prior to the week of interview.

Persons residing in 212 standard metropolitan statistical areas in the United States (as defined for the 1960 Decennial Census) reported a slightly higher rate of incidence of acute conditions than did persons in the nonfarm population residing outside of the SMSA's. Persons residing on farms outside of these 212 areas had the lowest incidence rate. Nonfarm persons living outside of metropolitan areas had the

highest rates of restricted activity and bed disability; the farm population again had the lowest rates.

Residents of the West Region reported more acute conditions (252 per 100 persons) than did persons residing in the other regions. Persons residing in the Northeast Region had the lowest rate, 192 cases per 100 persons per year. The excess in rate for the West Region compared with the Northeast Region was greatest for persons aged 15-24 years. The rates of restricted-activity days and bed-days were also highest in the West Region and lowest in the Northeast Region.

SOURCE OF DATA

The information contained in this publication is derived from household interviews conducted by the Health Interview Survey in cooperation the U.S. Bureau of the Census in a probabilist sample of the civilian, noninstitutional popular of the United States. The sample is designenthat interviews are conducted during every of the year. During the 52-week period from 1963-June 1964, the sample was compose approximately 42,000 households containing a 134,000 persons living at the time of the inview.




Figure 1. Incidence of acute conditions per 100 persons per quarter, by condition group.

A description of the design of the Survey, of the methods used in estimation, and of general qualifications of the data obtained from surveys is presented in Appendix I. Since estimates shown in this report are based on a sample of the population rather than on the entire population, they are subject to sampling error. Therefore, particular attention should be paid to the section entitled "Reliability of Estimates." Sampling errors for most of the estimates are of relatively low magnitude. However, where an estimated number or the numerator or the denominator of a rate or percentage is small, the sampling error may be high.

Certain terms used in this report are defined in Appendix II. Some of the terms have specialized meanings for the purpose of the Survey. For example, estimates of the incidence of acute conditions include, with certain exceptions, those conditions which had started within 2 weeks prior to the week of interview and which had involved either medical attention or restricted activity. The exceptions, listed in Appendix II, are certain conditions such as heart trouble and diabetes which are always considered chronic, regardless of duration or onset.

Annual estimates of the number of disability days associated with each acute condition are derived from the number of such days experienced during the 2-week period prior to the week of interview and include all those reported, even if the illness causing the disability had its onset prior to the 2-week period. Since any particular day of disability may be associated with more than one condition, the sum of days for all conditions adds to more than the total number of person-days of disability.

The questionnaire used by the Health Interview Survey during July 1963-June 1964 is illustrated in the publication "Current Estimates From the Health Interview Survey, United States, July 1963-June 1964" (Vital and Health Statistics, Series 10, Number 13). Each condition reported in response to the "illness-recall" questions (8-14) was entered on a line of table I. Further questions were asked to obtain a better description of the specific conditions and to obtain information about the actions taken by the person, such as seeking medical attention or restricting usual activities.

The incidence rates of acute conditions per quarter for the first 16 quarters from July 1957-June 1961, shown in figure 1, are presented in *Health Statistics*, Series B, Number 33. The rates for the period July 1961-June 1963 are presented in *Vital and Health Statistics*, Series 10, Numbers 1 and 10.

AND ASSOCIATED DAYS OF DISABILITY

Condition Group, Sex, and Age

During the 12 months ending June 1964 the civilian population, excluding the residents of institutions, experienced an estimated 387.4 million acute illnesses or injuries requiring either medical attention or reduction of daily activity (table 1). The incidence rate per 100 persons was 208.5 during the year. This rate is about 5 percent lower than the rate of 218.8 reported for the previous year in Vital and Health Statistics, Series 10, Number 10. The principal cause of the reduction in rate was the lower incidence of respiratory illness with a rate of 110.0 per 100 persons per year for the current period compared with 127.2 for the previous year. It can be seen in figure 1 that the pattern of low incidence of respiratory illness in alternate years. established early in the Survey, has been reinstituted. This pattern was interrupted in the period July 1961-June 1963 during which high incidence rates for respiratory illness were noted for 2 consecutive years. It is of interest that the incidence rates per quarter during the current year were remarkably similar to those for the July 1960-June 1961 period.

The overall incidence rate would have been lower in the current year if an increase in the incidence of infective and parasitic diseases and injuries had not occurred. The rise in rate of infective and parasitic diseases was caused primarily by increased incidence of common child-hood diseases. The annual incidence rate for measles was twice that for the preceding year; the rate for rubella (German measles) was 3½

times as great. Epidemics of these conditions began early in 1964 as evidenced by the distribution of cases reported in each of the following calendar quarters:

	Measles	<u>Rubella</u>
		of conditions thousands
Total	10,020	4,705
July-September 1963 October-December 1963 January-March 1964 April-June 1964	730 480 2,599 6,211	391 106 846 3,362

The estimated 9.6 million cases of measles and rubella occurring during the period April-June 1964 is a threefold increase over the number of cases, 3.2 million, reported for the corresponding quarter in 1963. The epidemic proportions of these diseases not only add to the total incidence of acute conditions but may also contribute to the increased occurrence of serious complications associated with these diseases, such as severe infections arising from measles and birth defects resulting from rubella infection during pregnancy.

The number of injuries requiring medical attention or restriction of activities during July 1963-June 1964 was 4.5 million greater than the incidence during the preceding year. The incidence rate for the current year was 29.7 per 100 persons per year, a gain of 2.0 percentage points. The rise in rate was restricted almost exclusively to males.

The comparatively low incidence of respiratory illness was due primarily to the low incidence rates of influenza and the common cold. The rate for influenza during the current year was about 73 percent of that for the preceding year; similarly, the rate for the common cold declined by about 10 percent.

Females experienced a somewhat higher rate of acute conditions than did males, 216.4 cases per 100 females compared with 200.0 per 100 males. The higher rate among females was largely attributable to respiratory illnesses and to the group "All other acute conditions" which includes genitourinary disorders (with a sex ratio of 6 to 1) and the sex-specific group "Deliveries and disorders of pregnancy and the puerperium." The

sex ratio was reversed for the injuries gro with the rate for males exceeding that females by 13.2 percentage points.

Acute illness and injury was responsible an annual rate of restricted activity of 81 days per 100 persons, including 345.9 days sp in bed (tables 2 and 3). Respiratory illnesses injuries were the two condition groups contribut the majority of the disability days, with 6 percent of all days reported assignable to eitl of these condition groups. An estimated 7 percent of the acute conditions required : striction of activities or a combination of m ical attention and activity restriction (table Thus, the duration of the average condit requiring reduction of activity was about 5.4 da (1,507,691,000 days of restricted activity divide by 277,446,000 cases requiring restriction activity). If acute conditions requiring only med attention were also included in the denominat the average duration of restricted activity | condition would be 3.9 days.

Distributions of condition groups and asseciated disability days by age and sex are present in tables 5-7. For most of these groups both incidence and disability were highest in the agroup under 6 years of age. In general, as a advanced, the rates of incidence and disabil declined; however, in many instances the rate of disability rose again in the oldest age group (fig. 2).

Children in the school-age population (6-years of age) experienced an annual incider rate of acute conditions of 253.4 per 100 childres About 72.3 percent of the conditions were eith respiratory illnesses or infective and parasidiseases. As a result of acute conditions, average child had 8.6 days of restricted activincluding 4.4 days absent from school (table Of the 8.6 days, 3.7 were days spent in b

Currently employed persons in the populat 17 years of age and older experienced an ann incidence rate of 155.6 acute conditions per persons (table 9). A currently employed person who had a job or business or worked at time during the 2-week period prior to the woof interview. The rate of acute illness was high for female workers than for male workers and was also higher for persons in the age ground.




Figure 2. Incidence of acute conditions per 100 persons per year, by sex and age.

17-44 years than for those in the 45 years and older group. The average employee lost 3.3 days from work on account of acute illness or injury (table 10). Respiratory illness contributed 1.2 days to this total, and injuries were responsible for 1.1 days.

Geographic Distribution

For the year July 1963-June 1964, there was a change in the method of classification of the population by place of residence. The residence categories—urban, rural nonfarm, and rural farm—are no longer used to describe the distribution of the population during intercensal years. Instead, the population by place of residence is classified into the following groups: those living in the 212 standard metropolitan statistical areas (SMSA) as defined for the 1960 Decennial Census and those living outside the standard metropolitan statistical areas in places classified as farm or nonfarm.

One effect of the change in classification by place of residence is a mixing of some of the urban and rural elements of the population. Since county boundaries are used in defining SMSA's (except in New England), rural elements within the county bounds are included in the SMSA. On the other hand, urban population not included in the 212 defined SMSA's is classified as non-farm. As a means of assessing the effect of this change in residence classification on the incidence rates of acute conditions, data on acute conditions

collected during the period July 1962-June 1963 are shown in the following table according to both methods of classification:

	Number of acute conditions per 100 persons per year
All areas	218.8
UrbanRural nonfarm	223.9 217.0 176.5
	Number of acute conditions per 100 persons per year
All areas	218.8
All SMSA's Non-SMSA-nonfarm	221.6 222.2

This comparison indicates that the change in classification has not affected the distribution of acute condition data by place of residence to any great extent.

Non-SMSA-farm-----

During July 1963-June 1964, residents of all standard metropolitan statistical areas established for the 1960 Decennial Census had an incidence of acute illnesses or injuries of 212.3 conditions per 100 persons per year. Persons residing in nonfarm areas outside of SMSA's had an incidence rate similar to that for the SMSA's, 208.5 per 100 persons. Residents of non-SMSA-farm areas had a significantly lower rate (169.7) than either of the other areas (table 11).

Figure 3 shows the incidence rates by age and place of residence. It is evident that the age distribution of rates in the farm areas was quite different from the pattern in the other areas. Table A shows the effect of age adjustment of the incidence rates by place of residence, indicating that the age distribution of residents of the three areas is not sufficiently different to explain the lower rates among the farm population.

176.9




Figure 3. Incidence of acute conditions per 100 persons per year, by residence and age.

The rates of restricted-activity and beddisability days associated with acute conditions were lowest in the farm areas and highest in the other nonmetropolitan areas (tables 12 and

Table A. Comparison between unadjusted and age-adjusted incidence of acute conditions per 100 persons per year, by residence and region: United States, July 1963-June 1964

Residence and region	Un- adjusted	Age- adjusted			
	Number of acute co ditions per 100 persons per year				
All areas	208.5	•••			
Residence					
All SMSA's Non-SMSA-nonfarm Non-SMSA-farm	212.3 208.5 169.7	212.6 208.0 169.4			
Region					
Northeast North Central South West	192.2 208.9 198.7 252.1	196.6 208.9 197.0 249.7			

¹Adjusted to the age distribution of the total civilian, noninstitutional population of the United States, July 1963-June 1964.

13). The disability rates for the SMSA's v similar to those for the nonfarm areas.

Tables 14-16 indicate substantially lc incidence rates in the farm population for of the acute condition groups except for the a groups of influenza and contusions and sufficial injuries. The distribution of disability a indicates that the infective and parasitic, diges system, and injuries groups were responsible the lower rates of disability for the farm reside

During July 1963-June 1964 the resident the West Region experienced the highest rat incidence of acute conditions requiring med attention and/or activity restriction (table Persons residing in the Northeast Region the lowest rate. Figure 4 compares the incide rates for the regions during the 7 years of collection in the Health Interview Survey.




Figure 4. Incidence of acute conditions per persons per year, by region.

the exception of the period July 1957-June 1958, a year in which Asian influenza reached epidemic proportions, the West Region has displayed a substantially higher incidence rate than any of the other regions. This same pattern of high rates in the West Region is apparent when incidence is limited to activity-restricting conditions (fig. 5).

Table A shows that after age adjustment of the incidence rates for each region the rate for the West Region remained substantially greater. This indicates that differences in age distribution among the regions are not the cause. Since the reporting of acute conditions is restricted to only those conditions with medical attention and/or reduced activity, it is of interest to examine the pattern of medical care among the regions. During July-December 1962, information was obtained for the personal health expenditures of the population. Expenditures for personal health services include such items as hospital care, medical and dental services, medicines, and other health related services and products. Per capita




Figure 5. Incidence of activity-restricting acute conditions per 100 persons per year, by region.

expenditure among the four regions was as follows:

	Medical expenses per person per year
Northeast	\$137
North Central	\$121
South	\$119
West	\$150

Another measure related to the distribution of acute conditions in the major regions of the country is the rate of physician visits per person per year. Information on the volume of physician visits was obtained during July 1957-June 1959 and also during July 1963-June 1964. The rates for these two periods, by geographic region, are shown below:

	Pl	nysician	visits	per	person	per	year	
	July 1	957-June	1959		July 1	1963-	June	1964
Northeast North Central-		5.4 4.7				4.		
South West		4.7 5.7				4. 5.		

Even though the rates for the earlier period reflect increased medical attention during the Asian influenza epidemic, the patterns of medical care are similar to those for acute conditions and health expenses, with the West Region reporting the highest rates. It is quite possible that the greater use of medical facilities by residents of the West Region is sufficient to explain the difference in the incidence rates of acute conditions. On the other hand, it is possible that the converse of this statement is correct. The greater incidence of acute conditions in the West Region required greater use of medical facilities.

Figure 6 shows that the incidence rates by age were highest in the West for most ages. It is of interest to note that for children under 5 years of age and for persons 65 years of age and older the rates were substantially the same for most of the regions. The sex difference in rates by age shown in figure 2 was reflected to a large extent in each of the regions, although the Northeast and West Regions displayed a some-




Figure 6. Incidence of acute conditions per 100 persons per year, by region and age.

what different pattern from that of the population as a whole for children under 15 years and persons 65 years and older.

The rates of disability days associated with acute conditions were highest in the West and lowest among residents of the Northeast Region (tables 18, 19, 21, 22).

The condition groups contributing substantially to the high rate for the West Region were

the respiratory conditions group and injut (table 20). It should be noted that the rate infective and parasitic diseases was low in the West and North Central Regions, indicate the possibility that the measles and rubella ϵ demics were less severe in these areas that the other regions.

Seasonal Variation: Calendar Quarter

Tables 23-30 show the distribution of ra of acute conditions by calendar quarter dur July 1963-June 1964. Figure I shows the dist bution of rates by quarter for each of th years of data collection for each of the ma condition groups. It is evident that the seasc distribution of respiratory conditions establis the pattern for the rates per quarter for acute conditions. It should be noted that the cufor injuries differs markedly from that for spiratory conditions, low when the respirat rate is high and vice versa.

The increased incidence of infective and pa sitic diseases during the 2 quarters of 1 reflects the epidemics of measles and rube mentioned earlier in the report.

DETAILED TABLES

		EXTENDED LIST OF CONDITION GROUPS, BY SEX	
Table	1.	Incidence of acute conditions, percent distribution, and number of acute conditions per 100 persons per year, by sex and condition group: United States, July 1963-June 1964	12
	2.	Days of restricted activity associated with acute conditions and days of restricted activity per 100 persons per year, by sex and condition group: United States, July 1963-June 1964	13
	3.	Days of bed disability associated with acute conditions and days of bed disability per 100 persons per year, by sex and condition group: United States, July 1963-June 1964	14
		MEASURES OF THE IMPACT OF ILLNESS	
	4.	Incidence of acute conditions and number of acute conditions per 100 persons per year, by measures of the impact of illness, sex, and condition group: United States, July 1963-June 1964	15
		SEX AND AGE	
	5.	Incidence of acute conditions and number of acute conditions per 100 persons per year, by age, sex, and condition group: United States, July 1963-June 1964	16
	6.	Days of restricted activity associated with acute conditions and days of restricted activity per 100 persons per year, by age, sex, and condition group: United States, July 1963-June 1964	17
	7.	Days of bed disability associated with acute conditions and days of bed disability per 100 persons per year, by age, sex, and condition group: United States, July 1963-June 1964	18
		TIME LOST FROM SCHOOL AND WORK	
	8.	Days lost from school associated with acute conditions and days lost from school per 100 children (6-16) per year, by sex and condition group: United States, July 1963-June 1964	19
	9.	Incidence of acute conditions among currently employed persons and number of acute conditions per 100 currently employed persons per year, by age, sex, and condition group: United States, July 1963-June 1964	20
	10.	Days lost from work associated with acute conditions and days lost from work per 100 currently employed persons per year, by age; sex, and condition group: United States, July 1963-June 1964	21

Page

DETAILED TABLES-Continued

RESIDENCE

- Table 11. Incidence of acute conditions and number of acute conditions per 100 persons per year, by residence, sex, and age: United States, July 1963-June 1964------

 - 13. Days of bed disability associated with acute conditions and days of bed disability per 100 persons per year, by residence, sex, and age: United States, July 1963-June 1964------
 - 14. Incidence of acute conditions and number of acute conditions per 100 persons per year, by residence, sex, and condition group: United States, July 1963-June 1964-
 - 15. Days of restricted activity associated with acute conditions and days of restricted activity per 100 persons per year, by residence, sex, and condition group: United States, July 1963-June 1964-------

REGION

- 17. Incidence of acute conditions and number of acute conditions per 100 persons per year, by region, sex, and age: United States, July 1963-June 1964-----

- 20. Incidence of acute conditions and number of acute conditions per 100 persons per year, by region, sex, and condition group: United States, July 1963-June 1964----
- 21. Days of restricted activity associated with acute conditions and days of restricted activity per 100 persons per year, by region, sex, and condition group: United States, July 1963-June 1964-----

CALENDAR QUARTER

- 23. Incidence of acute conditions per quarter and number of acute conditions per 100 persons per quarter, by sex and age: United States, July 1963-June 1964------

DETAILED TABLES—Continued

			Page
		CALENDAR QUARTER—Continued	
Table 2	8.	Days of bed disability associated with acute conditions per quarter and days of bed disability per 100 persons per quarter, by sex and condition group: United States, July 1963-June 1964	39
29	9.	Incidence of acute respiratory conditions and associated disability days per quarter and number of acute respiratory conditions per 100 persons per quarter, by age: United States, July 1963-June 1964	40
30	0.	Incidence of injuries and associated disability days per quarter and number of injuries per 100 persons, by age: United States, July 1963-June 1964	41
		POPULATION	
3:	1.	Population used in obtaining rates shown in this publication, by residence, region, sex, and age: United States, July 1963-June 1964	42
3	2.	Population of currently employed persons, by sex and age: United States, July 1963-June 1964	43
3:	3.	Population used in obtaining rates shown in this publication, by quarter, sex, and age: United States, July 1963-June 1964	44

TABLE 1. INCIDENCE OF ACUTE CONDITIONS, PERCENT DISTRIBUTION, AND NUMBER OF ACUTE CONDITIONS PER 100 PERSC PER YEAR, BY SEX AND CONDITION GROUP: UNITED STATES, JULY 1963-JUNE 1964

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliable

Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliable of the estimates are given in Appendix I. Definitions of terms are given in Appendix II

THE VIRIESTIVE AND PARASITIC DISEASES—————————————————————————————————	of the estimates are given in Appendix 1. Definitions of terms are given in Appendix in										
ALL ACUTE CONDITIONS	CONDITION GROUP		MALE	FEMALE		MAŁE	FEMALE		MALE	FEM	
INFECTIVE AND PARASITIC DISEASES 55,283 26,002 29,281 14,3 14,4 14,1 29,8 28,9 3								DITIONS PER 100			
COMMON CHILDHOOD DISEASES—— 24,666 12,987 11,680 6.4 7.2 5.6 13.3 14.4 1 THE VIRUS, N.O.S. 24,105 10,248 13,857 6.2 5.7 6.7 13.0 11.4 1 OTHER INVECTIVE AND PARASITIC 0152ASE—— 6,511 2,767 3,744 1.7 1.5 1.8 5.5 3.1 RESPERATORY CONDITIONS—— 204,301 93,808 110,493 52.7 52.1 53.3 110.0 104.1 11 UPPER RESPIRATORY CONDITIONS—— 133,797 62,119 71,678 34.5 34.5 34.6 72.0 69.0 7 COMMON COLD—— 103,393 47,990 55,403 26.7 26.6 26.7 55.6 53.3 5 OTHER ACUTE UPPER RESPIRATORY CONDITIONS—— 61,980 27,830 33,150 16.0 15.4 16.5 33.4 30.9 2 INFLUENZA MITH DIGESTIVE 61,990 27,830 33,150 16.0 15.4 16.5 33.4 30.9 2 INFLUENZA MITH DIGESTIVE 7,954 34.5 34.6 72.0 16.4 15.7 1 RAMIFESTATIONS—— 55,408 3.695 34.66 2.2 2.1 1.7 1.7 3.5 3.4 OTHER RESPIRATORY CONDITIONS—— 3,026 1.394 1.631 0.8 0.8 1.6 1.5 15.7 1.8 DIGESTIVE SYSTEM CONDITIONS—— 2,340 1,070 1,270 0.6 0.6 0.6 0.6 1.3 1.2 DIGESTIVE SYSTEM CONDITIONS—— 20,608 9,961 10,647 5.3 5.5 5.1 11.1 11.1 1 DENTAL CONDITIONS—— 55,226 32,895 22,331 14.3 18.3 10.8 29.7 36.5 2 TRACTURES, DISLOCATIONS, SPRAINS, AND STRAINS— AND STRAINS— 16,366 9,899 4.466 4.1 5.6 5.6 2.8 8.5 11.2 CONDITIONS—— 17,168 3,592 3,576 1.9 2.0 1.7 5.9 4.0 DISLOCATIONS—— 55,226 32,895 22,331 14.3 18.3 10.8 29.7 36.5 2 DISLOCATIONS—— 55,226 32,895 22,331 14.3 18.3 10.8 29.7 36.5 2 DISLOCATIONS—— 51,931 17,517 34,423 13.4 9.7 16.6 28.0 19.4 3 DISLOCATIONS—— 11,935 10,090 5,746 4.1 5.6 2.8 8.5 11.2 CONDITIONS—— 12,603 4.907 5.67 3.3 3.8 2.7 6.8 7.7 1.2 1.2 1.2 1.2 1.2 1.2 1.2 1.2 1.2 1.2	ALL ACUTE CONDITIONS	387,358	180,182	207,175	100.0	100-0	100.0	208.5	200.0	21	
THE VIRIESTIVE AND PARASITIC DISEASES—————————————————————————————————	INFECTIVE AND PARASITIC DISEASES	55,283	26,002	29,281	14.3	14.4	14.1	29.8	28.9	3	
RESPIRATORY CONDITIONS————————————————————————————————————	THE VIRUS, N.O.SOTHER INFECTIVE AND PARASITIC	24,105	10,248	13,857	6.2	5.7	6.7	13.0	11.4	1 1	
UPPER RESPIRATORY CONDITIONS— 133,797 62,119 71,678 34.5 34.5 34.6 72.0 69.0 7 CONMON COLO— 103,393 47,990 55,403 26.7 26.6 26.7 55.6 53.3 5 0 THE ACUTE UPPER RESPIRATORY CONDITIONS— 10,980 34,150 16.0 15.4 16.5 33.4 30.9 2 1 NFLUENZA— 11,085 17 1 1,980 17,187 18.7 19.5 18.7 19.5 19.5 19.5 19.5 19.5 19.5 19.5 19.5										11	
COMMON COLD————————————————————————————————————	KESPIKATUKY CUMUTTUMS	204,301	93,808	110443	52.1	52.1	23.3	110.0	104.1		
INFLUENZA HITH DIGESTIVE ANIFESTATIONS	COMMON COLDOTHER ACUTE UPPER RESPIRATORY	103,393	47,990	55,403	26.7	26.6	26.7	55.6	53.3	5	
OTHER INFLUENZA	INFLUENZA							33.4		1 2	
OTHER ACUTE RESPIRATORY CONDITIONS————————————————————————————————————	OTHER INFLUENZAOTHER RESPIRATORY CONDITIONS	55,408 8,524 3,026	24,727 3,859 1,394	30,682 4,666 1,631	14.3 2.2 0.8	13.7 2.1 0.8	14.8 2.3 0.8	29.8 4.6 1.6	27.5 4.3 1.5	2	
DENTAL CONDITIONS————————————————————————————————————	OTHER ACUTE RESPIRATORY										
FUNCTIONAL AND SYMPTOMATIC UPPER GASTROINTESTINAL DISORDERS, N.E.C.——————————————————————————————————	DIGESTIVE SYSTEM CONDITIONS	20,608	9,961	10,647	5.3	5.5	5.1	11.1	11.1	1	
N.E.C.	FUNCTIONAL AND SYMPTOMATIC UPPER	8,711	4,363	4,349	2.2	2.4	2•1	4.7	4.8		
INJURIES	N.E.C OTHER DIGESTIVE SYSTEM										
FRACTURES, DISLOCATIONS, SPRAINS, AND STRAINS————————————————————————————————————	CONDITIONS	7,168	3,592	3,576	1.9	2.0	1.7	3.9	4.0		
AND STRAINS————————————————————————————————————	INJURIES	55,226	32 + 895	22,331	14.3	18.3	10.8	29.7	36.5	2	
INJURIES————————————————————————————————————	AND STRAINS————————————————————————————————————	5,353 11,013	3,523 6,377	1,830 4,636	1.4 2.8	2.0 3.5	0.9 2.2	2.9 5.9	3.9 7.1		
DISEASES OF THE EAR	INJURIES										
HEADACHES————————————————————————————————————	ALL OTHER ACUTE CONDITIONS	51,941	17,517	34,423	13.4	9.7	16.6	28.0	19.4	3	
PREGNANCY AND THE PUERPERIUM 5,718 5.718 1.5 2.8 3.1 DISEASES OF THE SKIN 6,845 4,025 2,820 1.8 2.2 1.4 3.7 4.5 DISEASES OF THE MUSCULOSKELETAL SYSTEM 4,506 1,962 2,544 1.2 1.1 1.2 2.4 2.2	HEADACHESGENITOURINARY DISORDERS	4,151	1,375	2,776	1.1	0.8	1.3	2.2	1.5		
	PREGNANCY AND THE PUERPERIUM DISEASES OF THE SKIN DISEASES OF THE MUSCULOSKELETAL	6,845	4,025	2,820	1.8	2.2	1.4	3.7	4.5		
										1	

NOTES: Excluded from these statistics are all conditions involving neither restricted activity nor mediattention.

N.o.s.—not otherwise specified; n.e.c.—not elsewhere classified.

TABLE 2. DAYS OF RESTRICTED ACTIVITY ASSOCIATED WITH ACUTE CONDITIONS AND DAYS OF RESTRICTED ACTIVITY PER 100
PERSONS PER YEAR, BY SEX AND CONDITION GROUP: UNITED STATES, JULY 1963-JUNE 1964

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

and established are given in Appendix 1. Definitions of terms are given in Appendix ii										
CONDITION GROUP	BOTH SEXES	MALE	FEMALE	BOTH Sexes	MALE	FEMALE				
		RESTRICTED A N THOUSANDS	CTIVITY		DAYS OF RESTRICTED ACTIVITY PER 100 PERSONS PER YEAR					
ALL ACUTE CONDITIONS	1,507,691	659,107	848,584	811.5	731.7	886.5				
INFECTIVE AND PARASITIC DISEASES	239,096	113,327	125,770	128.7	125.8	131.4				
COMMON CHILDHOOD DISEASES	132,196 67,203	66,786 28,791	65,410 38,412	71.2 36.2	74.1 32.0	68•3 40•1				
OTHER INFECTIVE AND PARASITIC DISEASES	39,698	17,750	21,948	21.4	19.7	22.9				
RESPIRATORY CONDITIONS	610,380	269,246	341,133	328.5	298.9	356.4				
UPPER RESPIRATORY CONDITIONS	330,654	144,541	186,112	178.0	160.5	194.4				
OTHER ACUTE UPPER RESPIRATORY	241,621	103,903	137,718	130.0	115.3	143.9				
INFLUENZAINFLUENZAINFLUENZA WITH DIGESTIVE	89,032 211,509	40,638 93,231	48,394 118,278	47.9 113.8	45.1 103.5	50.6 123.6				
MANIFESTATIONSOTHER INFLUENZA	16,976 194,533	7,430 85,801	9,546 108,732	9.1 104.7	8.2 95.3	10.0 113.6				
OTHER RESPIRATORY CONDITIONS PNEUMONIA BRONCHITIS	68,218 37,529 17,910	31,474 16,185 9,155	36,743 21,344 8,754	36.7 20.2 9.6	34.9 18.0 10.2	38.4 22.3 9.1				
OTHER ACUTE RESPIRATORY CONDITIONS	12,779	6,134	6,645	6.9	6.8	6.9				
DIGESTIVE SYSTEM CONDITIONS	69,488	31,741	37,746	37.4	35.2	39.4				
DENTAL CONDITIONS	17,786	9,460	8,325	9•6	10.5	8.7				
GASTROINTESTINAL DISORDERS, N.E.C	11,972	4,959	7,013	6.4	5.5	7.3				
CONDITIONS	39,730	17,322	22,407	. 21.4	19.2	23.4				
INJURIES	330,590	177,102	153,488	177.9	196.6	160.4				
FRACTURES, DISLOCATIONS, SPRAINS,	173,279	91,714	81,565	93.3	101.8	85•2				
FRACTURES AND DISLOCATIONS SPRAINS AND STRAINS	105,477 67,802	55,374 36,340	50,103 31,462	56.8 36.5	61.5 40.3	52 .3 32 . 9				
OPEN WOUNDS AND LACERATIONS CONTUSIONS AND SUPERFICIAL	44,827	25,957	18,870	24-1	28.8	19.7				
OTHER CURRENT INJURIES	49,825 62,660	24,146 35,286	25,679 27,374	26.8 33.7	26.8 39.2	26.8 28.6				
ALL OTHER ACUTE CONDITIONS	258,137	67,691	190,446	138.9	75.1	199•0				
DISEASES OF THE EAR	30,012 3,688	11,623	18,389 2,595	16.2 2.0	12.9	19•2 2•7				
GENITOURINARY DISORDERSDELIVERIES AND DISORDERS OF	36,922	6,112	30,810	19.9	6.8	32.2				
PREGNANCY AND THE PUERPERIUM DISEASES OF THE SKIN DISEASES OF THE MUSCULOSKELETAL	56, 156 21,422	10,734	56,156 10,688	30.2 11.5	11.9	58.7 11.2				
SYSTEMALL OTHER ACUTE CONDITIONS	28,489 81,447	11,620 26,508	16,869 54,939	15.3 43.8	12.9 29.4	17.6 57.4				

NOTE: N.o.s.—not otherwise specified; n.e.c.—not elsewhere classified.

TABLE 3. DAYS OF BED DISABILITY ASSOCIATED WITH ACUTE CONDITIONS AND DAYS OF BED DISABILITY PER 100 PERSON PER YEAR, BY SEX AND CONDITION GROUP: UNITED STATES, JULY 1963-JUNE 1964

Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliabil of the estimates are given in Appendix I. Definitions of terms are given in Appendix II

of the estimates	are given in Appendi	x I. Definitions	or terms are given	III Appendix II			
CONDITION GROUP	BOTH SEXES	MALE	FEMALE	BOTH Sexes	MALE	FEMALE	
		F BED DISABI N THOUSANDS	LITY	DAYS OF BED DISABILITY PER 100 PERSONS PER YEAR			
ALL ACUTE CONDITIONS	642,741	272,897	369,844	345.9	303.0	38	
INFECTIVE AND PARASITIC DISEASES	111,138	51,041	60,098	59.8	56.7	6	
COMMON CHILDHOOD DISEASES	54,517	27,978	26,539	29.3	31.1	2	
OTHER INFECTIVE AND PARASITIC DISEASES	37,799 18,822	16,000 7,063	21,799	20.3	7.8	1	
RESPIRATORY CONDITIONS	291,924	131,944	159,980	157.1	146.5	16	
RESPERATORY CONDITIONS	271,724	2324311	137,750				
UPPER RESPIRATORY CONDITIONS COMMON COLD OTHER ACUTE UPPER RESPIRATORY	131,103 89,779	60,760 42,297	70,343 47,481	70.6 48.3	67.5 47.0	7 4	
CONDITIONSINFLUENZA	41,324 117,642	18,463 50,896	22,862 66,745	22.2 63.3	20.5 56.5	2 6	
INFLUENZA WITH DIGESTIVE MANIFESTATIONS OTHER INFLUENZA	9,803 107,839	4,686 46,211	5,117 61,628	5.3 58.0	5.2 51.3	6	
OTHER RESPIRATORY CONDITIONS	43,179	20,288	22,891	23.2	22.5	2	
PNEUMONIABRONCHITIS	28,060 9,607	12,830	15,230 4,619	15.1 5.2	14.2 5.5	'	
OTHER ACUTE RESPIRATORY CONDITIONS	5,513	2,470	3,043	3.0	2.7		
DIGESTIVE SYSTEM CONDITIONS	36,387	15,007	21,381	19•6	16.7	2	
DENTAL CONDITIONS	7,096	3,775	3,322	3.8	4.2		
GASTROINTESTINAL DISORDERS, N.E.C	6,259	2,134	4,125	3.4	2.4		
OTHER DIGESTIVE SYSTEM CONDITIONS	23,032	9,098	13,934	12.4	10.1	1	
INJURIES	96,493	50,812	45,681	51.9	56.4	4	
FRACTURES, DISLOCATIONS, SPRAINS,				25.0	28.2		
FRACTURES AND DISLOCATIONS	48,138 31,296	25,442 15,615	22,697 15,682	25.9 16.8	17.3	1 2	
SPRAINS AND STRAINS OPEN WOUNDS AND LACERATIONS CONTUSIONS AND SUPERFICIAL	16,842 12,653	9,827 6,669	7,015 5,983	9.1 6.8	10.9 7.4		
INJURIESOTHER CURRENT INJURIES	13,375 22,327	6.893 11,807	6,482 10,519	7.2 12.0	7.7 13.1	1	
ALL OTHER ACUTE CONDITIONS	106,798	24,094	82,704	57.5	26.7		
DISEASES OF THE EAR	9,457	3,822	5,635	5.1	4.2		
HEADACHESGENITOURINARY DISORDERS	1,794 18,143	2,462	15,681	1.0 9.8	2.7	1	
PREGNANCY AND THE PUERPERIUM DISEASES OF THE SKIN	30,447 6,643	3,318	30,447 3,325	16•4 3•6	3.7	3	
DISEASES OF THE MUSCULOSKELETAL SYSTEM	9,714 30,600	4,526 9,302	5,187 21,298	5.2 16.5	5.0 10.3	2	

NOTE: N.o.s.—not otherwise specified; n.e.c.—not elsewhere classified.

TABLE 4. INCIDENCE OF ACUTE CONDITIONS AND NUMBER OF ACUTE CONDITIONS PER 100 PERSONS PER YEAR, BY MEASURES OF THE IMPACT OF ILLNESS, SEX, AND CONDITION GROUP: UNITED STATES, JULY 1963-JUNE 1964

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

	ule estima	wes are given in 7	Appendix I. Defin	tuons of terms a	re given in App	enaix IIJ			
	MEASURES OF THE IMPACT OF ILLNESS								
SEX AND CONDITION GROUP	TOTAL	MEDICALLY ATTENDED ONLY	MEDICALLY ATTENDED AND ACTIV- ITY RE- STRICTING	ACTIVITY RESTRICT- ING ONLY (NOT MED- ICALLY ATTENDED)	TOTAL	MEDICALLY ATTENDED ONLY	MEDICALLY ATTENDED AND ACTIV- ITY RE- STRICTING	ACTIVITY RESTRICT- ING ONLY (NOT MED- ICALLY ATTENDED)	
BOTH SEXES	IN		CUTE CONDITI	ONS			UTE CONDITI		
ALL ACUTE CONDITIONS-	387,358	109,912	140,293	137,153	208.5	59•2	75.5	73.8	
INFECTIVE AND PARASITIC DISEASES RESPIRATORY CONDITIONS UPPER RESPIRATORY	55,283 204,301	6,591 48,385	25,367 64,198	23,325 91,718	29.8 110.0	3.5 26.0	13.7 34.6	12.6 49.4	
CONDITIONSINFLUENZAOTHER RESPIRATORY	133,797 61,980	41,456 4,737	37,724 20,652	54,617 36,591	72.0 33.4	22.3 2.5	20.3 11.1	29.4 19.7	
CONDITIONS	8,524	2,193	5,822	×	4-6	1.2	3.1	×	
CONDITIONSINJURIES	20,608 55,226	8,209 25,014	6,999 22,362	5,399 7,850	11.1 29.7	4.4 13.5	3.8 12.0	2.9 4.2	
ALL OTHER ACUTE CONDITIONS	51,941	21,712	21,368	8,861	28.0	11.7	11.5	4.8	
MALE									
ALL ACUTE CONDITIONS-	180,182	53,092	64,884	62,206	200.0	58.9	72.0	69-1	
INFECTIVE AND PARASITIC DISEASES RESPIRATORY CONDITIONS UPPER RESPIRATORY	26,002 93,808	3,423 21,680	11,452 30,424	11,127 41,704	28.9 104.1	3.8 24.1	12.7 33.8	12.4 46.3	
CONDITIONS	62,119 27,830	18,715 1,922	18,512 9,283	24,892 16,625	69.0 30.9	20.8 2.1	20.6 10.3	27.6 18.5	
OTHER RESPIRATORY CONDITIONS DIGESTIVE SYSTEM	3,859	1,043	2,629	3 £	4.3	1.2	2.9	×	
INJURIES	9,961 32,895	4,151 15,358	3,278 13,669	2,532 3,867	11.1 36.5	4.6 17.0	3.6 15.2	2 • 8 4 • 3	
ALL OTHER ACUTE CONDITIONS	17,517	8,481	6,061	2,975	19•4	9.4	6•7	3.3	
FEMALE									
ALL ACUTE CONDITIONS-	207,175	56,820	75,409	74,946	216.4	59.4	78.8	78.3	
INFECTIVE AND PARASITIC DISEASES RESPIRATORY CONDITIONS UPPER RESPIRATORY	29,281 110,493	3,168 26,706	13,915 33,774	12,198 50,013	30.6 115.4	3.3 27.9	14.5 35.3	12•7 52•2	
CONDITIONS	71,678 34,150	22,741 2,815	19,212 11,369	29,725 19,966	74.9 35.7	23.8 2.9	20.1 11.9	31.1 20.9	
OTHER RESPIRATORY CONDITIONS—————	4,666	1,150	3,193	×	4.9	1.2	3.3	×	
DIGESTIVE SYSTEM CONDITIONSINJURIES	10,647 22,331	4,059 9,656	3,721 8,692	2,867 3,983	11.1 23.3	4.2 10.1	3.9 9.1	3.0 4.2	
CONDITIONS	34,423	13,230	15,308	5,885	36.0	13.8	16.0	6.1	

NOTE: Excluded from these statistics are all conditions involving $\,$ neither restricted activity nor medical attention.

TABLE 5. INCIDENCE OF ACUTE CONDITIONS AND NUMBER OF ACUTE CONDITIONS PER 100 PERSON'S PER YEAR, BY I SEX, AND CONDITION GROUP: UNITED STATES, JULY 1963-JUNE 1964

Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the bility of the estimates are given in Appendix I. Definitions of terms are given in Appendix II

bility o	f the estimates	are given in	Appendix I. I	Definitions of	terms are giv	en in Appe	oudix III			
SEX AND CONDITION GROUP	ALL AGES	UNDER 6	6-16	17-44	45 & OVER	ALL AGES	UNDER 6	6-16	17-44	1
BOTH SEXES	I		OF ACUTE THOUSAND	CONDITION S	s		R OF AC 100 PER			ΡI
ALL ACUTE CONDITIONS-	387,358	90,037	103,775	121,462	72,084	208.5	360.5	253.4	186.2	1:
INFECTIVE AND PARASITIC DISEASES—————————————————————————————————	55,283 204,301 133,797	16,318 50,358 37,958	22,658 52,366 34,044	10,728 61,229 36,289	5,578 40,348 25,507	29.8 110.0	65.3 201.6	55.3 1.27.9 83.1	16.4 93.8 55.6	:
INFLUENZAOTHER RESPIRATORY	61,980	9,570	16,892	22,803	12,715	33.4	38.3	41.2	35.0	;
CONDITIONS DIGESTIVE SYSTEM CONDITIONS	8,524 20,608	2,831 4,165	1,430 4,431	2,137 7,724	2,126 4,289	11.1	11.3	3.5 10.8	3.3	
INJURIESALL OTHER ACUTE CONDITIONS	55,226 51,941	9,380	13,394	20,361	12,091 9,778	29.7	37.6	32.7 26.7	31.2	
	22,012									
MALE										
ALL ACUTE CONDITIONS-	180,182	46,196	53,013	49,987	30,986	200.0	364.4	254.5	161.9	1
INFECTIVE AND PARASITIC DISEASES RESPIRATORY CONDITIONS UPPER RESPIRATORY	26,002 93,808	8,987 25,420	11,331 25,650	3,566 24,873	2,116 17,865	28.9 104.1	70.9 200.5	54.4 123.1	11.6 80.6	
CONDITIONS	62,119 27,830	19,414 4,631	16,454 8,646	15,284 8,636	10,968 5,916	69.0 30.9	153.1 36.5	79.0 41.5	49.5 28.0	
CONDITIONSDIGESTIVE SYSTEM	3,859	1,375	*	953	980	4.3	10.8	*	3.1	
INJURIESALL OTHER ACUTE	9,961 32,895	2,561 5,076	2,096 8,770	3,292 12,903	2,012 6,146	11.1 36.5	20.2 40.0	10.1	10.7	
CCNDITIONS	17,517	4,151	5,165	5,353	2,848	19.4	32.7	24.8	17.3	
FEMALE										
ALL ACUTE CONDITIONS-	207,175	43,841	50,761	71,475	41,098	216.4	356.6	252.2	207.9	1
INFECTIVE AND PARASITIC DISEASES	29,281 110,493	7,331 24,938	11,326 26,715	7,162 36,356	3,462 22,484	30.6 115.4	59.6 202.8	56.3 132.7	20.8 105.8	
CONDITIONS	71,678 34,150	18,544 4,938	17,590 8,246	21,005 14,167	14,539 6,799	74.9 35.7	150.8 40.2	87.4 41.0	61.1 41.2	
OTHER RESPIRATORY CONDITIONS DIGESTIVE SYSTEM	4,666	1,456	880	1,184	1,146	4.9	11.8	4.4	3.4	
CONDITIONSINJURIES	10,647 22,331	1,604 4,304	2,334 4,624	4,432 7,458	2,277 5,945	11.1 23.3	13.0 35.0	11.6 23.0	12.9 21.7	
ALL OTHER ACUTE CONDITIONS	34,423	5,664	5,762	16,067	6,930	36.0	46.1	28.6	46.7	<u> </u>

NOTE: Excluded from these statistics are all conditions involving neither restricted activity medical attention.

TABLE 6. DAYS OF RESTRICTED ACTIVITY ASSOCIATED WITH ACUTE CONDITIONS AND DAYS OF RESTRICTED ACTIVITY PER 100 PERSONS PER YEAR, BY AGE, SEX, AND CONDITION GROUP: UNITED STATES, JULY 1963-JUNE 1964

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II.

		6-7	-PP-Halle II				٠. نـــــ			
SEX AND CONDITION GROUP	ALL AGES	UNDER 6	6-16	17-44	45 & OVER	ALL AGES	UNDER 6	6-16	17-44	45 & OVER
BOTH SEXES	DAYS OF	RESTRICTE	D ACTIVII	Y IN THOU	ISANDS		YS OF RES ER 100 PE			TY
ALL ACUTE CONDITIONS-	1,507,691	240,516	351,825	465,181	450,170	811.5	963•1	859.0	713.0	824.1
INFECTIVE AND PARASITIC DISEASES RESPIRATORY CONDITIONS UPPER RESPIRATORY	-	64,825 123,960	108,691 146,487	37,035 163,876	28,545 176,057	128.7 328.5	259.6 496.4	265.4 357.7	56.8 251.2	52.3 322.3
CONDITIONS INFLUENZA OTHER RESPIRATORY	330,654 211,509	81,485 24,390	86,199 50,687	82,103 69,357	80,866 67,075	178.0 113.8	326.3 97.7	210.5 123.8	125.8 106.3	148.0 122.8
CONDITIONS DIGESTIVE SYSTEM	68,218	18,085	9,601	12,416	28,116	36.7	72.4	23.4	19.0	51.5
CONDITIONSINJURIES	69,488 330,590	10.593 14.654	10,849 53,194	21,994 119,291	26,052 143,451	37.4 177.9	42.4 58.7	26.5 129.9	33.7 182.8	47•7 262•6
ALL OTHER ACUTE CONDITIONS	258,137	26,484	32,603	122,984	76,065	138.9	106.1	79.6	188.5	139.3
MALE										
ALL ACUTE CONDITIONS-	659,107	123,173	175,157	168,979	191,798	731.7	971.5	840.9	547.4	746.4
INFECTIVE AND PARASITIC DISEASES RESPIRATORY CONDITIONS UPPER RESPIRATORY CONDITIONS INFLUENZA	113,327 269,246 144,541 93,231	35,290 61,930 42,594 11,304	53,173 69,409 41,064 24,407	13,145 61,367 30,323 26,389	11,718 76,540 30,560 31,130	125.8 298.9 160.5 103.5	278.3 488.4 335.9 89.2	255.3 333.2 197.1 117.2	42.6 198.8 98.2 85.5	45.6 297.9 118.9 121.1
OTHER RESPIRATORY CONDITIONS DIGESTIVE SYSTEM	31,474	8,031	3,938	4,655	14,850	34.9	63.3	18.9	15.1	57.8
CONDITIONSINJURIESALL OTHER ACUTE	31,741 177,102	6,631 8,526	4,966 32,763	7,166 70,441	12,978 65,372	35.2 196.6	52•3 67•2	23.8 157.3	23.2 228.2	50.5 254.4
CONDITIONS	67,691	10,797	14,846	16,859	25,190	75.1	85.2	71.3	54•6	98.0
FEMALE		: -								
ALL ACUTE CONDITIONS-	848,584	117,343	176,668	296,202	258,372	886.5	954.5	877.8	861.7	893.2
INFECTIVE AND PARASITIC DISEASES RESPIRATORY CONDITIONS UPPER RESPIRATORY	125,770 341,133	29,535 62,031	55,518 77,078	23,890 102,509	16,827 99,516	131.4 356.4	240.2 504.6	275.9 383,0	69.5 298.2	58.2 344.0
INFLUENZA OTHER RESPIRATORY	186,112 118,278	38,851 13,086	45,136 26,279	51,780 42,968	50,306 35,944	194.4 123.6	316.3 106.4	224.3 130.6	150.6 125.0	124.3
CONDITIONS DIGESTIVE SYSTEM	36,743	10,054	5,663	7,761	13,266	38.4	81.8	28-1	22.6	45.9
CONDITIONS INJURIES ALL OTHER ACUTE	37,746 153,488	3,962 6,128	5,883 20,431	14,828 48,850	13,073 78,080	39.4 160.4	32.2 49.8	29.2 101.5	43.1 142.1	45.2 269.9
CONDITIONS	190,446	15,687	17,758	106,126	50,875	199.0	127.6	88.2	308.7	175.9

TABLE 7. DAYS OF BED DISABILITY ASSOCIATED WITH ACUTE CONDITIONS AND DAYS OF BED DISABILITY FER 100 PERSONS P YEAR, BY AGE, SEX, AND CONDITION GROUP: UNITED STATES, JULY 1963-JUNE 1964

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of estimates are given in Appendix I. Definitions of terms are given in Appendix II]

estimates are given in Appendix 1. Definitions of terms are given in Appendix 11											
SEX AND CONDITION GROUP	ALL AGES	UNDER 6	6-16	17-44	45 & OVER	ALL AGES	UNDER 6	6-16	17-44	45 0V	
BOTH SEXES	DAYS OF	BED DISA	BILITY IN	THOUSAND	s	D	AYS OF BE 100 PER	D DISAB SONS PE		R	
ALL ACUTE CONDITIONS-	642,741	110,569	153,273	211,479	167,420	345.9	442.8	374.2	324.1	30	
INFECTIVE AND PARASITIC DISEASES RESPIRATORY CONDITIONS UPPER RESPIRATORY	111,138 291,924	28,880 57,798	47,776 77,676	21,448 84,089	13,034 72,361	59.8 157.1	115.6 231.4	116.7 189.7	32.9 128.9	2 13	
INFLUENZA	131,103 117,642	29,575 14,513	40,806 30,755	36,489 38,894	24,234 33,479	70.6 63.3	118.4 58.1	99.6 75.1	55.9 59.6	6	
OTHER RESPIRATORY CONDITIONS DIGESTIVE SYSTEM	43,179	13,710	6,115	8,707	14,648	23.2	54.9	14.9	13.3	2	
CONDITIONSINJURIESALL OTHER ACUTE	36,387 96,493	5,293 4,337	4,949 11,324	11,731 35,921	14,415 44,911	19.6 51.9	21.2 17.4	12.1 27.6	18.0 55.1	2 8	
CGNDITIONS	106,798	14,260	11,548	58,290	22,700	57.5	57.1	28.2	89.3	4	
MALE											
ALL ACUTE CONDITIONS-	272,897	56,358	70,618	73,704	72,216	303.0	444.5	339.0	238.7	28	
INFECTIVE AND PARASITIC DISEASES RESPIRATORY CONDITIONS UPPER RESPIRATORY CONDITIONS INFLUENZA	51,041 131,944 60,760 50,896	14,824 29,879 16,806 6,950	23,777 34,823 18,025 14,440	6,952 33,494 14,840 15,054	5,488 33,748 11,088 14,453	56.7 146.5 67.5 56.5	116.9 235.7 132.5 54.8	114.1 167.2 86.5 69.3	22.5 108.5 48.1 48.8	2 13 4 5	
OTHER RESPIRATORY CONDITIONS	20,288	6,123	2,358	3,600	8,207	22.5	48.3	11.3	11.7	3	
DIGESTIVE SYSTEM CONDITIONS INJURIES	15,007 50,812	3,338 2,521	1,981 5,396	3,417 24,016	6,271 18,880	16.7 56.4	26.3 19.9	9.5 25.9	11.1 77.8	2 7	
ALL OTHER ACUTE CONDITIONS	24,094	5,796	4,642	5,826	7,830	26.7	45.7	22.3	18.9	3	
FEMALE											
ALL ACUTE CONDITIONS-	369,844	54,210	82,655	137,775	95,204	386.4	440.9	410.7	400.8	32	
INFECTIVE AND PARASITIC DISEASES RESPIRATORY CONDITIONS UPPER RESPIRATORY	60,098 159,980	14,055 27,919	24,000 42,853	14,496 50,595	7,546 38,613	62.8 167.1	114.3 227.1	212.9	42.2 147.2	2 13	
CONDITIONS	70,343 66,745	12,769 7,564	22,781 16,315	21,648 23,840	13,145 19,026	73.5 69.7	103.9 61.5	81.1	63.0	4	
CONDITIONS————— DIGESTIVE SYSTEM	22,891	7,587	3,757	5,106	6,441	23.9	61.7	18.7	14.9	2	
CONDITIONSINJURIESALL OTHER ACUTE	21,381 45,681	1,955 1,816	2,967 5,928	8,314 11,906	8,143 26,031	22.3 47.7	15.9 14.8	14.7 29.5	24.2 34.6	9	
CONDITIONS	82,704	8,464	6,906	52,463	14,870	86.4	68•8	34.3	152.6	5	

TABLE 8. DAYS LOST FROM SCHOOL ASSOCIATED WITH ACUTE CONDITIONS AND DAYS LOST FROM SCHOOL PER 100 CHILDREN (6-16) PER YEAR, BY SEX AND CONDITION GROUP: UNITED STATES, JULY 1963-JUNE 1964

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

						
CONDITION GROUP	BOTH SEXES	MALE	FEMALE	BOTH SEXES	MALE	\$ FEMALE
	DAYS	LOST FROM IN THOUSAN			ST FROM SO HILDREN PE	
ALL ACUTE CONDITIONS	181,037	88,237	92,801	442.0	423.6	461.1
×						
INFECTIVE AND PARASITIC DISEASES	65,831	31,607	34,223	160.7	151.7	170.0
RESPIRATORY CONDITIONS	87,399	41,856	45,543	213.4	200.9	226.3
UPPER RESPIRATORY CONDITIONS	51,143	24,239		124.9		133.7
INFLUENZA	31,625	15,602	16,023	77.2		79.6
OTHER RESPIRATORY CONDITIONS	4,630	2,014	2,616	11.3	9.7	13.0
DIGESTIVE SYSTEM CONDITIONS	4,990	2,027	2,963	12.2	9.7	14.7
INJURIES	9,364	6,595	2,769	22.9	31.7	13.8
ALL OTHER ACUTE CONDITIONS	13,454	6,151	7,302	32.8	29.5	36.3
						•

TABLE 9. INCIDENCE OF ACUTE CONDITIONS AMONG CURRENTLY EMPLOYED PERSONS AND NUMBER OF ACUTE CONDITIONS PER 100 CURRENTLY EMPLOYED PERSONS PER YEAR, BY AGE, SEX, AND CONDITION GROUP: UNITE STATES, JULY 1963-JUNE 1964

[Data are based on household interviews of the civilian, noninstitutional population: The survey design, general qualifications, and informatic on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II.

on the feriality of the estimates are given in Appendix 1. Definitions of terms are given in Appendix I.										
⇒ SEX AND CONDITION GROUP .	ALL AGES- 17 & OVER	17-44	45 & OVER	ALL AGES- 17 & OVER	17-44	45 ε OVER				
BOTH SEXES		E OF ACUI IN THOUS		NUMBER OF PER 100 CU PERSON		EMPLOYE				
ALL ACUTE CONDITIONS	109,082	71,433	37,648	155.6	171.7	132.				
INFECTIVE AND PARASITIC DISEASES RESPIRATORY CONDITIONS UPPER RESPIRATORY CONDITIONS INFLUENZA OTHER RESPIRATORY CONDITIONS DIGESTIVE SYSTEM CONDITIONS LNJURIES	8,660 56,796 34,551 20,229 2,015 6,856 22,363 14,408	5,786 36,130 21,288 13,609 1,232 4,586 14,863 10,068	2,874 20,666 13,263 6,620 783 2,270 7,500 4,339	49.3 28.8 2.9 9.8	13.9 86.8 51.2 32.7 3.0 11.0 35.7 24.2	10. 72. 46. 23. 2. 8. 26.				
MALE										
ALL ACUTE CONDITIONS	67,364	43,877	23,487	147.3	161.3	126.				
INFECTIVE AND PARASITIC DISEASES RESPIRATORY CONDITIONS UPPER RESPIRATORY CONDITIONS OTHER RESPIRATORY CONDITIONS DIGESTIVE SYSTEM CONDITIONS INJURIES	4,646 34,995 21,225 12,388 1,382 4,284 16,591 6,848	3,165 21,654 12,926 7,923 805 2,712 11,591 4,754	1,480 13,341 8,299 4,465 1,572 5,000 2,094	10.2 76.5 46.4 27.1 3.0 9.4 36.3 15.0	11.6 79.6 47.5 29.1 3.0 10.0 42.6 17.5	8. 72. 44. 24. 8. 27. 11.				
FEMALE										
ALL ACUTE CONDITIONS	41,718	27,557	14,161	171.0	191.2	141.				
LNFECTIVE AND PARASITIC DISEASES RESPIRATORY CONDITIONS UPPER RESPIRATORY CONDITIONS INFLUENZA OTHER RESPIRATORY CONDITIONS	4.014 21,800 13,326 7,842	2,621 14,476 8,362 5,687	1,393 7,324 4,964 2,155	16.5 89.4 54.6 32.1	18.2 100.4 58.0 39.5	14. 73. 49. 21.				
DIGESTIVE SYSTEM CONDITIONS	2,572 5,772 7,560	1,875 3,272 5,314	2,500 2,246	10.5 23.7 31.0	13.0 22.7 36.9	25. 22.				

NOTE: Excluded from these statistics are all conditions involving neither restricted activaty nor medical attention.

TABLE 10. DAYS LOST FROM WORK ASSOCIATED WITH ACUTE CONDITIONS AND DAYS LOST FROM WORK PER 100 CURRENTLY EMPLOYED PERSONS PER YEAR, BY AGE, SEX, AND CONDITION GROUP: UNITED STATES, JULY 1963-JUNE 1964

Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II

on the reliability of the estimates are given	en in Appendix.	. Definitions	or terms are g	given in Appendi	(11)		
SEX AND CONDITION GROUP	ALL AGES- 17 & OVER	17-44	45 & OVER	ALL AGES- 17 & OVER	17-44	45 & OVER	
BOTH SEXES		.OST FROM THOUSANDS		100 CURRE	DAYS LOST FROM WORK F 100 CURRENTLY EMPLOYE PERSONS PER YEAR		
ALL ACUTE CONDITIONS	228,179	125,333	102,846	325.4	301.2	360.8	
INFECTIVE AND PARASITIC DISEASES RESPIRATORY CONDITIONS UPPER RESPIRATORY CONDITIONS INFLUENZA	20,735 87,352 39,958 37,948 9,446 12,261 75,220 32,611	11,687 48,777 23,351 21,477 3,949 5,705 38,221 20,944	9,048 38,575 16,607 16,472 5,496 6,556 36,999 11,668	29.6 124.6 57.0 54.1 13.5 17.5 107.3 46.5	28.1 117.2 56.1 51.6 9.5 13.7 91.8 50.3	31.7 135.3 58.3 57.8 19.3 23.0 129.8 40.9	
MALE	į						
ALL ACUTE CONDITIONS	147,743	75,484	72,259	323.1	277.5	390.0	
INFECTIVE AND PARASITIC DISEASES RESPIRATORY CONDITIONS UPPER RESPIRATORY CONDITIONS INFLUENZA	10,919 56,322 24,179 25,525 6,618 9,243 56,245 15,013	5,654 28,459 13,587 13,277 1,595 3,404 30,633 7,334	5,265 27,863 10,592 12,248 5,023 5,839 25,612 7,679	23.9 123.2 52.9 55.8 14.5 20.2 123.0 32.8	20-8 104-6 50-0 48-8 5-9 12-5 112-6 27-0	28.4 150.4 57.2 66.1 27.1 31.5 138.2 41.4	
FEMALE							
ALL ACUTE CONDITIONS	80,436	49,849	30,587	329.8	345.9	306•5	
INFECTIVE AND PARASITIC DISEASES—————————————————————————————————	9,816 31,030 15,779 12,423 2,828 3,018 18,974 17,598	6,033 20,318 9,764 8,199 2,354 2,301 7,587 13,610	3,782 10,712 6,014 4,224 :: 11,387 3,989	40.2 127.2 64.7 50.9 11.6 12.4 77.8 72.1	41.9 141.0 67.7 56.9 16.3 16.0 52.6 94.4	37.9 107.3 60.3 42.3 :: :114.1 40.0	

TABLE 11. INCIDENCE OF ACUTE CONDITIONS AND NUMBER OF ACUTE CONDITIONS PER 100 PERSONS PER YEAR, RESIDENCE, SEX, AND AGE: UNITED STATES, JULY 1963-JUNE 1964

Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II

				RESIDENCE	50 (Mag) (Ag)(1)	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1		
SEX AND AGE	ALL AREAS	ALL SMSA	NON-SMSA NONFARM	NON-SMSA Farm	ALL ARE A S	ALL SMSA	NON-SMSA NONFARM	NON-SI FARI
BOTH SEXES	INCI	DENCE OF AC		ONS			E CONDITIONS PER YEA	
ALL AGES	387,358	252,055	115,412	19,890	208.5	212.3	208.5	16
UNDER 5 YEARS	76,083	50,664	22,407	3,012	367.2	384.0	352.6	25(
5-14 YEARS	103,653	68,700	29,827	5,126	271.6	287.7	258.0	188
15-24 YEARS	55,836	34,966	18,250	2,620	207.1	206.7	217.8	15
25-44 YEARS	79,701	53,187	23,184	3,331	175.8	175.1	184.1	140
45-64 YEARS	52,539	33,276	14,912	4,351	139.7	136.4	141.2	164
65 & OVER YEARS	19,545	11,264	6,831	1,450	114.8	113.0	115.9	12!
MALE			į					
ALL AGES	180,182	116,350	54,021	9,812	200.0	203.2	202-0	16:
UNDER 5 YEARS	38,847	25,382	11,764	1,701	367.9	376.0	367.9	279
5-14 YEARS	53,009	35,706	14,880	2,424	273.5	295.8	252.7	171
15-24 YEAR\$	24,433	15,830	7,482	1,121	190.7	196.9	192.8	12!
25-44 YEARS	32,906	20,641	10,754	1,511	152.2	142.8	179.2	129
45-64 YEARS	23,225	14,718	6,293	2,215	127.9	126.1	123.1	162
65 & OVER YEARS	7,761	4,074	2•847	840	102.9	95.3	107.3	130
FEMALE								
ALL AGES	207,175	135,705	61,391	10,079	216.4	220.8	214.6	178
UNDER 5 YEARS	37,236	25,282	10,643	1,311	366.4	392.5	337.2	23:
5-14 YEARS	50,644	32,994	14,947	2,703	269.7	279.5	263.5	201
15-24 YEARS	31,403	19,136	10,768	1+499	222.0	215.4	239.3	191
25-44 YEARS	46,795	32,546	12,430	1,820	197.4	204.5	188.5	15;
45-64 YEARS	29,314	18,558	8,620	2,136	150.7	145.9	158.2	16
65 & OVER YBARS	11,783	7,190	3,984	×	124.3	126.2	123.0	

NOTE: Excluded from these statistics are all conditions involving neither restricted activity \boldsymbol{n} medical attention.

TABLE 12. DAYS OF RESTRICTED ACTIVITY ASSOCIATED WITH ACUTE CONDITIONS AND DAYS OF RESTRICTED ACTIVITY PER 100 PERSONS PER YEAR, BY RESIDENCE, SEX, AND AGE: UNITED STATES, JULY 1963-JUNE 1964

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II.

				RESIDENCE				
SEX AND AGE	ALL AREAS	ALL SMSA	NON-SMSA Nonfarm	NON-SMSA Farm	ALL AREAS	ALL SMSA	NON-SMSA NONFARM	NON-SMSA FARM
BOTH SEXES	DA		ICTED ACTIV	TTY		F RESTRIC		
ALL AGES	1,507,691	961,182	458,073	88,436	811.5	809.5	827.7	754.6
UNDER 5 YEARS	194,356	136,190	51,761	6,406	938.0	1,032.3	814.6	545.7
5-14 YEARS	344,445	229,622	96,912	17,911	902.6	961.7	838.2	658.5
15-24 YEARS	203,631	125,437	66,800	11,394	755.3	741.4	797.2	686.4
25-44 YEARS	315,090	213,881	88,751	12,458	695.1	704.2	704.6	526.8
45-64 YEARS	280,772	166,078	85,640	29,054	746.7	680.7	810.8	1,100.1
65 & OVER YEARS	169,397	89,974	68,210	11,213	995.2	902.4	1,157.5	967.5
MALE								
ALL AGES	659,107	417,121	196+958	45,028	731.7	728.4	736.6	741.2
UNDER 5 YEARS	99,158	68,276	26,975	3,907	939.2	1,011.3	843.5	641.5
5-14 YEARS	173,592	117,304	46,796	9,491	895.6	971.8	794.6	667•4
15-24 YEARS	79,816	48,793	25•900	5,123	622.8	607.0	667.4	571.8
25-44 YEARS	114,743	74,924	34,902	4,917	530.6	518.3	581.6	419.9
45-64 YEARS	133,331	78,327	39,038	15,966	734.5	670.8	763.4	1,172.2
65 & OVER YEARS	58,467	29,497	23,347	5,623	775.0	690.3	879.7	912.8
FEMALE								
ALL AGES	848,584	544,061	261,115	43,408	886.5	885.1	912.7	769.1
UNDER 5 YEARS	95,199	67,914	24,786	2,499	936.7	1,054.2	785.4	441.5
5-14 YEARS	170,853	112,318	50,116	8,419	909.9	951.4	883.4	648.1
15-24 YEARS	123,815	76,644	40,900	6,271	875.3	862.9	909.1	820.8
25-44 YEARS	200,346	138,957	53,849	7,541	845.1	873.0	816.5	631.6
45-64 YEARS	147,441	87,751	46,602	13,088	758.1	689.8	855.4	1,023.3
65 & OVER YEARS	110,930	60,478	44,862	5,590	1,170.3	1,061.8	1,385.1	1,029.5

TABLE 13. DAYS OF BED DISABILITY ASSOCIATED WITH ACUTE CONDITIONS AND DAYS OF BED DISABILITY PER DECRETAGE OF PERSONS PER YEAR, BY RESIDENCE, SEX, AND AGE: UNITED STATES, JULY 1963-JUNE 1964,

Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II

	Marin Control			RESIDENCE				
SEX AND AGE	ALL AREAS	ÄLL SMSA	NON-SMSA NONFARM	NON-SMSA FARM	ALL AREAS	ALL SMSA	NON-SMSA NONFARM	NON-SI FARI
BOTH SEXES	1	DAYS OF BED IN THO	DISABILITY USANDS		DAYS O		ABILITY F PER YEAR	PER 100
ALL AGES	642,741	409,302	198,802	34,637	345.9	344.7	359.2	29!
UNDER 5 YEARS	91,515	63,790	25,442	2,283	441.7	483.5	400.4	194
5-14 YEARS	147,137	94,267	43,603	9,266	385.6	394.8	377.1	341
15-24 YEARS	99,596	60,814	33,210	5,571	369.4	359.4	396.3	33!
25-44 YEARS	137,074	90,966	41,030	5,077	302.4	299.5	325.7	214
45-64 YEARS	108,400	65,382	33,923	9,096	288.3	268.0	321.2	344
65 & OVER YEARS	59,020	34,082	21,594	3,344	346.7	341.8	366.4	288
MALE								
ALL AGES	272,897	172,063	84,403	16,431	303.0	300.5	315.7	271
UNDER 5 YEARS	47,043	31,522	13,991	1,530	445.6	466.9	437.5	25:
5-14 YEARS	69,288	44,410	19,752	5,126	357.5	367.9	335.4	361
15-24 YEARS	34,102	19,854	11,758	2,490	266.1	2,47.0	303.0	27
25-44 YEARS	50,248	32,229	16,868	×	232.3	223.0	281.1	
45-64 YEARS	49,228	30,981	14,132	4,115	271.2	265.3	276.3	30:
65 & OVER YEARS	22,988	13,067	7,902	2,019	304.7	305.8	297.7	32
<u>FEMALE</u>								
ALL AGES	369,844	237,238	114,399	18,206	386.4	386.0	399.9	32:
UNDER 5 YEARS	44,472	32,268	11,451	×	437.6	500.9	362.8	
5-14 YEARS	77,849	49,857	23,852	4,141	414.6	422.3	420.4	31
15-24 YEARS	65,493	40,961	21,452	3,081	463.0	461.2	476.8	40:
25-44 YEARS	86,826	58,737	24,162	3,926	366.3	369.0	366.4	32
45-64 YEARS	59,172	34,401	19,790	4,981	304.2	270.4	363.3	38
65 & OVER YEARS	36,032	21,015	13,692	ж	380.1	368.9	422.7	

TABLE 14. INCIDENCE OF ACUTE CONDITIONS AND NUMBER OF ACUTE CONDITIONS PER 100 PERSONS PER YEAR, BY RESIDENCE, SEX, AND CONDITION GROUP: UNITED STATES, JULY 1963-JUNE 1964

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix II. Definitions of terms are given in Appendix II.

estimate	s are given in A	ppendix I. Defin	itions of terms a	re given in App	endix II				
				RESIDENCE					
SEX AND CONDITION GROUP	ALL AREAS	ALL SMSA	NON-SMSA NONFARM	NON-SMSA Farm	ALL AREAS	ALL SMSA	NON-SMSA NONFARM	NON-SMSA FARM	
BOTH SEXES	INC	INCIDENCE OF ACUTE CONDITIONS NUMBER OF ACUTE CONDITIONS PERSONS PER Y							
ALL ACUTE CONDITIONS	387,358	252,055	115,412	19,890	208.5	212.3	208.5	169.7	
INFECTIVE AND PARASITIC DISEASES— RESPIRATORY CONDITIONS————————————————————————————————————	55,283 204,301 133,797 61,980 8,524 20,608 55,226 16,366 15,835 10,421 12,603 51,941	35,748 133,437 90,128 37,951 5,359 12,209 37,292 10,946 10,880 7,069 8,398 33,369	16,767 59,839 36,685 20,338 2,815 7,438 15,246 4,753 4,189 2,637 3,667 16,121	2,767 11,024 6,984 3,691 961 2,688 % 767 716 2,450	29.8 110.0 72.0 33.4 4.6 11.1 29.7 8.8 8.5	30.1 112.4 75.9 32.0 4.5 10.3 31.4 9.2 9.2 6.0 7.1 28.1	30.3 108.1 66.3 36.7 5.1 13.4 27.5 8.6 7.6	23.6 94.1 59.6 31.5 8.2 22.9 4 6.5 6.1	
MALE	_	_	•	·					
ALL ACUTE CONDITIONS	180,182	116,350	54,021	9,812	200.0	203.2	202.0	161.5	
INFECTIVE AND PARASITIC DISEASES— RESPIRATORY CONDITIONS————————————————————————————————————	26,002 93,808 62,119 27,830 3,859 9,961 32,895 9,899 10,090 5,999 6,907 17,517	16,248 60,500 41,664 16,360 2,477 6,108 21,865 6,357 6,914 4,086 4,509 11,629	8,171 28,261 17,303 9,655 1,303 3,320 9,377 3,072 2,881 1,360 2,064 4,891	1,582 5,046 3,152 1,815 2 1,653 **	28.9 104.1 69.0 30.9 4.3 11.1 36.5 11.0 11.2 6.7 7.7 19.4	28.4 105.6 72.8 28.6 4.3 10.7 38.2 11.1 12.1 7.1 7.9 20.3	30.6 105.7 64.7 36.1 4.9 12.4 35.1 11.5 10.8 5.1 7.7	26.0 83-1 51.9 29.9 27.2 ** 27.2	
FEMALE									
ALL ACUTE CONDITIONS	207,175	135,705	61,391	10,079	216.4	220.8	214.6	178.6	
INFECTIVE AND PARASITIC DISEASES— RESPIRATORY CONDITIONS————————————————————————————————————	29,281 110,493 71,678 34,150 4,666 10,647 22,331	21,590 2,883 6,101 15,427 4,589	8,596 31,577 19,382 10,683 1,512 4,118 5,869	1.185 5.978 3.832 1,876 2 1,035	30.6 115.4 74.9 35.7 4.9 11.1 23.3	31.7 118.7 78.8 35.1 4.7 9.9 25.1	30.0 110.4 67.7 37.3 5.3. 14.4 20.5	21.0 105.9 67.9 33.2 # 18.3	
OPEN HOUNDS AND LACERATIONS CONTUSIONS AND SUPERFICIAL INJURIES OTHER CURRENT INJURIES ALL OTHER ACUTE CONDITIONS	5,746 4,422 5,697 34,423	3,966 2,983 3,889 21,740	1,308 1,277 1,603 11,231	# # # 1,452	6.0 4.6 6.0 36.0	6.5 4.9 6.3 35.4	4.6 4.5 5.6 39.3	# # 25•7	

NOTE: Excluded from these statistics are all conditions involving neither restricted activity nor medical attention.

TABLE 15. DAYS OF RESTRICTED ACTIVITY ASSOCIATED WITH ACUTE CONDITIONS AND DAYS OF RESTRICTED ACTIVITY PER 1
PERSONS PER YEAR, BY RESIDENCE, SEX, AND CONDITION GROUP: UNITED STATES, JULY 1963-JUNI 1964

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of
estimates are given in Appendix I. Definition of terms are given in Appendix II.

esumates	are given in App	endix I. Dellilli	ion of terms are	given in Append			·	
				RESIDENCE				
SEX AND CONDITION GROUP	ALL AREAS	ALL SMSA	NON-SMSA NONFARM	NON-SMSA FARM	ALL AREAS	ALL SMSA	NON-SMSA NONFARM	NON-SM FARM
BOTH SEXES	DA	YS OF RESTR IN THO		ITY			ED ACTIV	
ALL ACUTE CONDITIONS	1,507,691	961,182	458,073	88,436	811.5	809.5	827.7	754
INFECTIVE AND PARASITIC DISEASES— RESPIRATORY CONDITIONS————————————————————————————————————	239,096 610,380 330,654 211,509 68,218 69,488 330,590 173,279 44,827 49,825 62,660 258,137	157,972 382,613 211,992 125,043 45,577 41,454 218,938 114,139 28,006 30,944 45,848 160,206	68,784 187,818 97,664 72,809 17,345 24,869 96,351 53,079 12,948 16,238 14,088 80,250	12,340 39,949 20,997 13,656 5,296 3,165 15,301 6,061 3,873 2,643 2,724 17,681	128.7 328.5 178.0 113.8 36.7 37.4 177.9 93.3 24.1 26.8 33.7 138.9	133.1 322.3 178.5 105.3 38.4 34.9 184.4 96.1 23.6 26.1 38.6	31.3 44.9 174.1 95.9 23.4 29.3	340 179 116 45 27 130 51 33
MALE	230,131	100,200	80,230	17,001	130.9	154.9	145.0	
ALL ACUTE CONDITIONS	659,107	417,121	196,958	45,028	731.7	728.4	736.6	741
INFECTIVE AND PARASITIC DISEASES— RESPIRATORY CONDITIONS————————————————————————————————————	113,327 269,246 144,541 93,231 31,474 31,741 177,102	71,875 171,802 95,137 53,597 23,068 18,992 112,011	34,610 78,041 39,701 31,419 6,921 11,219 54,244	6,842 19,403 9,703 8,214 1,531 10,846	125.8 298.9 160.5 103.5 34.9 35.2 196.6	125.5 300.0 166.1 93.6 40.3 33.2 195.6	291.9 148.5 117.5 25.9 42.0	319 159 135
FRACTURES, DISLOCATIONS, SPRAINS, AND STRAINS————— OPEN WOUNDS AND LACERATIONS——— CONTUSIONS AND SUPERFICIAL	91,714 25,957	57,795 15,499	29,213 7,902	4,706 2,556	101.8 28.8	100.9 27.1		
INJURIES OTHER CURRENT INJURIES ALL OTHER ACUTE CONDITIONS	24,146 35,286 67,691	13,765 24,953 42,441	8,473 8,656 18,843	1,907 1,677 6,406	26.8 39.2 75.1	24.0 43.6 74.1	32.4	27
FEMALE								
ALL ACUTE CONDITIONS	848,584	544,061	261,115	43,408	886.5	885.1	912.7	769
INFECTIVE AND PARASITIC DISEASES— RESPIRATORY CONDITIONS————————————————————————————————————	125,770 341,133 186,112 118,278 36,743 37,746 153,488		34,174 109,777 57,964 41,390 10,424 13,650 42,107	5,498 20,546 11,294 5,442 3,810 1,634 4,455	131.4 356.4 194.4 123.6 38.4 39.4 160.4	140.1 343.0 190.1 116.2 36.6 36.5 174.0	383.7 202.6 144.7 36.4 47.7 147.2	364 200 96 67 29 78
OPEN HOUNDS AND LACERATIONS CONTUSIONS AND SUPERFICIAL INJURIES OTHER CURRENT INJURIES ALL OTHER ACUTE CONDITIONS	18,870 25,679 27,374 190,446	12,507 17,179 20,896 117,764	7,765 5,431 61,407	# # 11,275	19.7 26.8 28.6 199.0	27.9 27.9 34.0 191.6	17.6 27.1 19.0	

TABLE 16. DAYS OF BED DISABILITY ASSOCIATED WITH ACUTE CONDITIONS AND DAYS OF BED DISABILITY PER 100 PERSONS PER YEAR, BY RESIDENCE, SEX, AND CONDITION GROUP: UNITED STATES, JULY 1963-JUNE 1964

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

		,		RESIDENCE		·					
SEX AND CONDITION GROUP	ALL AREAS	ALL SMSA	NON-SMSA NONFARM	NON-SMSA FARM	ALL AREAS	ALL SMSA	NON-SMSA NONFARM	NON-SMSA FARM			
BOTH SEXES			DISABILITY	,	DAYS O		SABILITY P PER YEAR	ER 100			
BOTH SEACS		211 1110	OSANDO								
ALL ACUTE CONDITIONS	642,741	409,302	198,802	34,637	345.9	344.7	359.2	295.5			
INFECTIVE AND PARASITIC DISEASES RESPIRATORY CONDITIONS	111,138 291,924	69,845 190,204	35,154 84,099	6,140 17,621	59.8 157.1	58.8 160.2	63.5 152.0	52.4 150.3			
UPPER RESPIRATORY CONDITIONS	131,103	89,586	34,416	7,102	70.6	75.5	62.2	60.6 68.1			
OTHER RESPIRATORY CONDITIONS	117,642 43,179	71,246	38,418 11,265	7,978 2,542	63.3 23.2	60.0 24.7	69.4 20.4	21.7			
DIGESTIVE SYSTEM CONDITIONS	36,387	22,626	12,072	1,690	19.6	19.1	21.8	14.4			
FRACTURES, DISLOCATIONS,	96,493	60,577	33,198	2,718	51.9	51.0	60.0	23.2			
SPRAINS, AND STRAINS	48,138	29,733	17,277	*	25.9	25.0	31.2	×			
OREN WOUNDS AND LACERATIONS	12,653	6,533	5,572	×	6.8	5.5	10.1				
INJURIES	13,375	8,080	4,674	×	7.2	6.8	8.4	* *			
OTHER CURRENT INJURIES	22,327 106,798	16,231 66,050	5,676 34,279	6,469	12.0 57.5	13.7 55.6	10.3	55.2			
ALL OTHER ACOTE COMOTITIONS	100,130	00,000	34,217	0,,05	,,,,,	33.0		,,,,,			
MALE											
ALL ACUTE CONDITIONS	272 007	172 043	84 403	14 621	303.0	300.5	315.7	270.5			
ALL ACUTE CONDITIONS	272,897	172,063	84,403	16,431	303.0	300.5	313.1	210.5			
INFECTIVE AND PARASITIC DISEASES	51,041	30,692	16,648	3,701	56.7	53.6	62.3	60.9			
RESPIRATORY CONDITIONS	131,944	86,767	37,348	7,830	146.5	151.5	139.7	128.9			
URPER RESPIRATORY CONDITIONS	60,760	41,271	16,972	2,517	67.5 56.5	72.1 53.4	63.5 59.4	41.4 73.2			
OTHER RESPIRATORY CONDITIONS	50,896 20,288	30,580 14,915	15,872 4,505	4,445 ×	22.5	26.0	16.8	×			
DIGESTIVE SYSTEM CONDITIONS	15,007	9,840	4,352	2 014	16-7	17-2	16.3 70.1	33.2			
FRACTURES, DISLOCATIONS,	50,812	30,063	18,734	2,014	56•4	52.5	70.1				
SPRAINS, AND STRAINS	25,442	14,840	9,632	# #	28.2	25.9	36.0	* *			
OPEN WOUNDS AND LACERATIONS CONTUSIONS AND SUPERFICIAL	6,669	3,310	3,230	•	7.4	5.8	12.1				
INJURIES	6,893	4,164	2,108	x ×	7.7	7.3	7.9	×			
OTHER CURRENT INJURIESALL OTHER ACUTE CONDITIONS	11,807 24,094	7,749 14,701	3,763 7,322	2,071	13.1 26.7	13.5 25.7	14-1 27-4	34.1			
<u>FEMALE</u>		-									
ALL ACUTE CONDITIONS	369,844	237,238	114,399	18,206	386.4	386.0	399.9	322.6			
INFECTIVE AND PARASITIC DISEASES	60,098	39,153	18,506	2,439	62.8	63.7	64.7	43.2			
RESPIRATORY CONDITIONS	159,980	103,437	46,751	9,791	167.1	168.3	163.4	173.5 81.2			
UPPER RESPIRATORY CONDITIONS	70,343 66,745		17,444 22,546	4,585 3,533	73.5 69.7		78.8	62.6			
OTHER RESPIRATORY CONDITIONS	22,891	14,457	6,761	1,674	23.9	23.5	23.6	29.7			
DIGESTIVE SYSTEM CONDITIONS	21,381 45,681	12,785 30,514	7,720 14,464	×	22.3 47.7	20.8 49.6	27.0 50.6	×			
FRACTURES, DISLOCATIONS,								×			
SPRAINS, AND STRAINS OREN WOUNDS AND LACERATIONS	22,697 5,983	14,894	7,644 2,341	×	23.7 6.3	24.2 5.2	26.7 8.2	×			
CONTUSIONS AND SUPERFICIAL				×				×			
INJURIESOTHER CURRENT INJURIES	6,482 10,519	3,916 8,482	2,566 1,913	×	6.8 11.0	6.4 13.8	9.0 6.7	×			
ALL OTHER ACUTE CONDITIONS	82,704	51,348	26,958	4,398	86.4	83.5	94.2	77.9			
					L	<u> </u>	L				

TABLE 17. INCIDENCE OF ACUTE CONDITIONS AND NUMBER OF ACUTE CONDITIONS PER 100 PERSONS PER YEAR, BY REGION, SE AND AGE: UNITED STATES, JULY 1963-JUNE 1964

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

	REGION										
SEX AND AGE				1	REGI	ON I	1	r	,	ı	
	ALL REGIONS	NORTH- EAST	NORTH CENTRAL	SOUTH	WEST	ALL REGIONS	NORTH- EAST	NORTH CENTRAL	SOUTH	WES.	
BOTH SEXES	INCIDENCE OF ACUTE CONDITIONS IN THOUSANDS					NUMBER OF ACUTE CONDITIONS PER 100 PERSONS PER YEAR					
ALL AGES	387,358	89,328	110,505	112,842	74,682	208.5	192.2	208.9	198.7	252	
UNDER 5 YEARS	76,083	18,804	21,160	23,335	12,784	367.2	386.2	354.8	360.8	373	
5-14 YEARS	103,653	26,089	29,184	27,042	21,338	271.6	293.6	263.3	227.1	339	
15-24 YEARS	55,836	10,463	16,136	18,312	10,924	207.1	161.8	219.9	203.1	263	
25-44 YEARS	79,701	17,447	22,630	22,177	17,447	175.8	150.1	178.3	164.1	232	
45-64 YEARS	52,539	12,543	14,948	16,018	9,031	139.7	124.0	140.1	144.5	157	
65 & OVER YEARS	19,545	3,983	6,448	5,957	3,157	114.8	88-2	125.2	123.8	124	
MALE											
ALL AGES	180,182	40,682	51,967	52,508	35,025	200.0	182.4	199.7	192.4	242	
UNDER 5 YEARS	38,847	9,172	10,298	12,467	6,911	367.9	379.5	330.4	379.6	397.	
5-14 YEARS	53,009	13,118	15,384	13,490	11,018	273.5	295.6	272.9	222.6	339,	
15-24 YEARS	24,433	4,761	7,169	8,160	4,343	190.7	155.2	200.7	192.4	224.	
25-44 YEARS	32,906	6,639	10,051	8,773	7,443	152.2	119.0	162.0	139.2	210.	
45-64 YEARS	23,225	5,334	6,645	7•291	3,954	127.9	109.3	128.0	139.5	138,	
65 & OVER YEARS	7,761	1,657	2,420	2,327	1,357	102.9	86.4	104.9	107.3	117.	
FEMALE											
ALL AGES	207,175	48,646	58,538	60,335	39,656	216-4	201-2	217.9	204.4	261.	
UNDER 5 YEARS	37,236	9,633	10,862	10,869	5,873	366.4	393.0	381.5	341.5	349.	
5-14 YEARS	50,644	12,970	13,800	13,553	10,321	269.7	291.7	253.3	231.8	340.	
15-24 YEARS	31,403	5,702	8,967	10,152	6,582	222.0	167.9	238.1	212.6	298.	
25-44 YEARS	46,795	10,807	12,579	13,404	10,005	197.4	178.8	194.0	185.7	252.	
45-64 YEARS	29,314	7,209	8,302	8,726	5,077	150.7	137.7	151.6	149.0	176.	
65 & OVER YEARS	11,783	2,326	4,027	3,631	1,800	124.3	89.6	141.4	137.4	129.	

NOTE: Excluded from these statistics are all conditions involving neither restricted activity nor medical at tention.

TABLE 18. DAYS OF RESTRICTED ACTIVITY ASSOCIATED WITH ACUTE CONDITIONS AND DAYS OF RESTRICTED PERSONS PER YEAR, BY REGION, SEX, AND AGE: UNITED STATES, JULY 1963-JUNE 1964

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II

estimates are given in Appendix I. Definitions of terms are given in Appendix II													
	. REGION												
SEX AND AGE	ALL REGIONS	NORTH- EAST	NORTH CENTRAL	SOUTH	WEST	ALL REGIONS	NORTH- EAST	NORTH CENTRAL	SOUTH	WEST			
BOTH SEXES	DAYS OF RESTRICTED ACTIVITY IN THOUSANDS						DAYS OF RESTRICTED ACTIVITY PER 100 PERSONS PER YEAR						
ALL AGES	1,507,691	349,156	413,272	455,018	290,246	811.5	751.3	781.3	801.0	979.9			
UNDER 5 YEARS	194,356	56,687	49,751	55,936	31,982	938.0	1,164.2	834.2	864.9	934.9			
5-14 YEARS	344,445	88,242	98,475	85,634	72,093	902.6	993.2	888.4	719.1	1,147.8			
15-24 YEARS	203,631	39,010	52,585	73,339	38,696	755.3	603.4	716.7	813.4	934.5			
25-44 YEARS	315,090	72,913	88,519	91,707	61,951	695.1	627.2	697.5	678.4	826.0			
45-64 YEARS	280,772	59,906	71,146	96,747	52,973	746.7	592.1	666.8	872.9	924.5			
65 & OVER YEARS	169,397	32,397	52,795	51,654	32,551	995.2	717.7	1,024.7	1,073.7	1,279.0			
MALE	:							!					
ALL AGES	659,107	150,921	173,881	205,298	129,007	731.7	676.7	668.0	752.4	892.1			
UNDER 5 YEARS	99,158	28,388	24,721	29,222	16,827	939•2	1,174.5	793.1	889.8	967.6			
5-14 YEARS	173,592	46,733	47,369	43,245	36,245	895.6	1,053.0	840.3	713.5	1,116.9			
15-24 YEARS	79,816	14,045	21,584	27,461	16,725	622.8	457.8	604.3	647.5	864.8			
25-44 YEARS	114,743	22,669	31,983	33,458	26,634	530.6	406.2	515.4	531.0	752.4			
45-64 YEARS	133,331	30,220	33,049	46•487	23,576	734.5	619.0	636.5	889.2	826.9			
65 & OVER YEARS	58,467	8,866	15,175	25,426	9,000	775.0	462.5	658.1	1,172.2	781.3			
FEMALE													
ALL AGES	848,584	198,235	239,391	249,720	161,240	886.5	820-1	891.0	845.9	1,063.7			
UNDER 5 YEARS	95,199	28,299	25,030	26,715	15,155	936.7	1,154.6	879.2	839.3	901.0			
5-14 YEARS	170,853	41,509	51,106	42,390	35,848	909.9	933.6	938.1	724.9	1,180.8			
15-24 YEARS	123,815	24,965	31,001	45,878	21,971	875.3	734.9	823.2	960.8	995.5			
25-44 YEARS	200,346	50,244	56,536	58,249	35,317	845.1	831.2	871.9	807.1	891.6			
45-64 YEARS	147,441	29,686	38,097	50,260	29,397	758.1	567.0	695 . 5	858.4	1,021.1			
65 & OVER YEARS	110,930	23,531	37,620	26,228	23,551	1,170.3	906.1	1,321.4	992.7	1,690.7			
		L	1			<u> </u>							

TABLE 19. DAYS OF BED DISABILITY ASSOCIATED WITH ACUTE CONDITIONS AND DAYS OF BED DISABILITY PER 100 PERSONS PER YEAR, BY REGION, SEX, AND AGE: UNITED STATES, JULY 1963-JUNE 1964

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

	REGION										
SEX AND AGE	ALL REGIONS	NORTH- EAST	NORTH CENTRAL	SOUTH	WEST	ALL REGIONS	NORTH- EAST	NORTH CENTRAL	SOUTH	HES.	
BOTH SEXES	DAYS OF BED DISABILITY IN THOUSANDS					DAYS OF BED DISABILETY PER 100 PERSONS PER VEAR					
ALL AGES	642,741	136,430	176,489	200,096	129,726	345.9	293.5	333.6	352.3	438	
UNDER 5 YEARS	91,515	26,499	23+239	27,513	14,264	441.7	544.2	389.7	425.4	417.	
5-14 YEARS	147,137	34,280	43,955	36,305	32,596	385.6	385•8	396.5	304-9	519.	
15-24 YEARS	99,596	17,382	23,609	40,184	18,421	369.4	268-9	321.8	445.7	444	
25-44 YEARS	137,074	27,095	40,319	41,657	28,003	302.4	233-1	317.7	308.2	373	
45-64 YEARS	108,400	21,739	28,843	36,687	21,132	288.3	214.9	270.3	331-0	368	
65 & OVER YEARS	59,020	9,435	16,523	17,751	15,311	346.7	209.0	320.7	369.0	601	
MALE											
ALL AGES	272,897	56,178	69,508	89,780	57,431	303-0	251.9	267.0	329.1	397	
UNDER 5 YEARS	47,043	12,342	11,408	15,256	8,037	445•6	510-6	366.0	464-6	462	
5-14 YEARS	69,288	17,309	19,500	17,578	14,900	357.5	390.0	345.9	290-0	459.	
15-24 YEARS	34,102	5,148	7,968	13,541	7,445	266.1	167.8	223.1	319.3	385.	
25-44 YEARS	50,248	6,858	13,399	17,321	12,670	232.3	122.9	215.9	274.9	357.	
45-64 YEARS	49,228	11,381	11,670	16,943	9,234	271.2	233.1	224.8	324.1	323.	
65 & OVER YEARS	22,988	3,139	5,563	9,142	5,145	304.7	163.7	241.2	421.5	446	
FEMALE											
ALL AGES	369,844	80,252	106,980	110,316	72,295	386.4	332.0	398.2	373.7	476	
UNDER 5 YEARS	44,472	14,157	11,831	12,257	6,227	437.6	577.6	415.6	385.1	370	
5-14 YEARS	77,849	16,971	24,455	18,727	17,696	414.6	381.7	448.9	320.2	582	
15-24 YEARS	65,493	12,234	15,641	26,642	10,976	463.0	360.1	415.3	557.9	497	
25-44 YEARS	86,826	20,236	26,919	24,337	15,333	366.3	334.8	415.2	337.2	387	
45-64 YEARS	59,172	10,357	17,173	19,744	11,897	304.2	197.8	313.5	337-2	413	
65 & OVER YEARS	36,032	6,296	10,961	8,609	10,166	380.1	242.4	385.0	325.9	729	

TABLE 20. INCIDENCE OF ACUTE CONDITIONS AND NUMBER OF ACUTE CONDITIONS PER 100 PERSONS PER YEAR, BY REGION, SEX, AND CONDITION GROUP: UNITED STATES, JULY 1963-JUNE 1964

Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II.

- uni	given in	Appendix I.	Definitions o	f terms are gi	ven in Append	lix I <u>I</u> J				
			·		REGI	ои	•			
SEX AND CONDITION GROUP	ALL REGIONS	NORTH- EAST	NORTH CENTRAL	SOUTH	WEST	ALL REGIONS	NORTH- EAST	NORTH CENTRAL	SOUTH	WEST
BOTH SEXES	INCIDENC	E OF ACUI	E CONDIT	IONS IN TH	HOUSANDS	NUMB		JTE CONDIT	TIONS PER YEAR	100
ALL ACUTE CONDITIONS	387,358	89,328	110,505	112,842	74,682	208.5	192.2	208.9	198.7	252.1
INFECTIVE AND PARASITIC DISEASES	55,283	18,014	11,566	19,977	5,726	29-8	38.8	21.9	35,2	19.3
RESPIRATORY CONDITIONS UPPER RESPIRATORY CONDITIONS INFLUENZA OTHER RESPIRATORY CONDITIONS	204,301 133,797 61,980 8,524	41,868 34,612 5,141 2,116	63,210 36,711 24,549 1,949	54,109 39,984 11,313 2,812	45,114 22,491 20,976 1,647	110.0 72.0 33.4 4.6	90-1 74-5 11-1 4-6	119.5 69.4 46.4 3.7	95.3 70.4 19.9 5.0	152.3 75.9 70.8 5.6
DIGESTIVE SYSTEM CONDITIONS	20,608	4,851	5,461	6,701	3,595	11.1	10.4	10.3	11.8	12.1
INJURIES————————————————————————————————————	55,226 16,366 15,835	13:066 3:634 4:248	15,460 4,688 4,384	16,094 4,436 4,832	3,609 2,371	29.7 8.8 8.5	28-1 7-8 9-1	29.2 8.9 8.3	28.3 7.8 8.5	35.8 12.2 8.0
OTHER CURRENT INJURIES	10,421 12,603	1,680 3,504	3,102 3,286	3,381 3,445	2,258 2,369	5.6 6.8	3.6 7.5	5.9 6.2	6.0 6.1	7.6 8.0
ALL OTHER ACUTE CONDITIONS	51,941	11,530	14,809	15,962	9,639	28.0	24.8	28.0	28.1	32.5
MALE										
ALL ACUTE CONDITIONS	180,182	40,682	51.,967	52,508	35,025	200.0	182.4	199.7	192.4	242.2
INFECTIVE AND PARASITIC DISBASES	26,002	8,335	5,267	9,491	2,908	28.9	37.4	20.2	34.8	20.1
RESPIRATORY CONDITIONS————————————————————————————————————	93,808 62,119 27,830 3,859	18,638 15,490 2,314 834	29,622 17,466 11,346 810	24.709 18.587 4.828 1.294	20,839 10,575 9,343 921	104.1 69.0 30.9 4.3	83.6 69.5 10.4 3.7	113.8 67.1 43.6 3.1	90.6 68.1 17.7 4.7	144.1 73.1 64.6 6.4
DIGESTIVE SYSTEM CONDITIONS	9,961	2,243	3,017	3,161	1,539	11.1	10.1	11.6	11.6	10.6
INJURIESFRACTURES, DISLOCATIONS,	32,895	7,246	8,843	10,281	6,525	36.5	32.5	34.0	37.7	45-1
SPRAINS, AND STRAINS OPEN HOUNDS AND LACERATIONS CONTUSIONS AND SUPERFICIAL	9,899 10,090	2,046 2,419	2,675 2,834	2,937 3,211	2,241	11.0	9.2 10.8	10.3 10.9	10.8 11.8	15.5 11.2
OTHER CURRENT INJURIES	5,999 6,907	973 1,808	1,594 1,740	2,108 2,024	1,324 1,334	6.7 7.7	4.4 8.1	6.1 6.7	7.7 7.4	9•2 9•2
ALL OTHER ACUTE CONDITIONS	17,517	4,220	5,217	4,866	3,215	19.4	18.9	20.0	17.8	22.2
FEMALE										
ALL ACUTE CONDITIONS	207,175	48,646	58,538	60,335	39,656	216-4	201.2	217.9	204.4	261.6
INFECTIVE AND PARASITIC DISBASES	29,281	9,679	6,298	10,486	2,818	30.6	40 . 0.	23.4	35.5	18.6
RESPIRATORY CONDITIONS—— UPPER RESPIRATORY CONDITIONS—— INFLUENZA—— OTHER RESPIRATORY CONDITIONS——	110,493 71,678 34,150 4,666	23,230 19,122 2,827 1,281	33,587 19,244 13,203 1,140	29,400 21,396 6,486 1,518	24,275 11,915 11,634 726	115.4 74.9 35.7 4.9	96.1 79.1 11.7 5.3	125.0 71.6 49.1 4.2	99.6 72.5 22.0 5.1	160.1 78.6 76.8 4.8
DIGESTIVE SYSTEM CONDITIONS	10,647	2,607	2:444	3,539	2,057	11.1	10.8	9.1	12.0	13.6
INJURIESFRACTURES, DISLOCATIONS, SPRAINS, AND STRAINS	22,331 6,467	5,820 1,588	6,617 2,012	5,814 1,498	4,082 1,368	23.3 6.8	24.1	24.6 7.5	19.7 5.1	26.9 9.0
OPEN WOUNDS AND LACERATIONS————————————————————————————————————	5,746 4,422	1,829 707	1,550	1,621	745 934	. 6.0	7.6 2.9	5.8 5.6	5.5	4.9
OTHER CURRENT INJURIES	5,697	1,696	1,546	1,273	1,034	4.6 6.0	7.0	5.8	4.3 4.8	6.2 6.8
ALL OTHER ACUTE CONDITIONS	34,423	7,310	9,592	11,096	6,425	36.0	30.2	35.7	37.6	42.4

NOTE: Excluded from these statistics are all conditions involving neither restricted activity nor medical attention.

TABLE 21. DAYS OF RESTRICTED ACTIVITY ASSOCIATED WITH ACUTE CONDITIONS AND DAYS OF RESTRICTED ACTIVITY PER 100 PERSONS PER YEAR, BY REGION, SEX, AND CONDITION GROUP: UNITED STATES, JULY 1963-JUNE 1964

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

		- ippozuik ii			тел та търъви					
					REGIO	ON				
SEX AND CONDITION GROUP	ALL REGIONS	NORTH- EAST	NORTH CENTRAL	SOUTH	WEST	ALL REGIONS	NORTH- EAST	NORTH CENTRAL	SOUTH	WEST
BOTH SEXES	DAYS OF	RESTRICTE	ED ACTIVIT	ry IN THOU	ISANDS	DAYS		ICTED ACT		100
ALL ACUTE CONDITIONS	1,507,691	349,156	413,272	455,018	290,246	811.5	751.3	-781.3	801.0	979.5
INFECTIVE AND PARASITIC DISEASES	239,096	73,236	61,987	71,271	32,602	128.7	157.6	117.2	125.5	110-1
RESPIRATORY CONDITIONS——— UPPER RESPIRATORY CONDITIONS——— INFLUENZA————————————————————————————————————	610,380 330,654 211,509 68,218	122,085 83,912 16,810 21,363	177,072 90,649 70,592 15,830	171,012 101,992 50,002 19,018	140,210 54,101 74,105 12,005	328.5 178.0 113.8 36.7	262.7 180.5 36.2 46.0	334.7 171.4 133.4 29.9	301.1 179.6 88.0 33.5	473.4 182.7 250.2 40.5
DIGESTIVE SYSTEM CONDITIONS	69,488	16,755	17,387	24,673	10,674	37.4	36-1	32.9	43.4	36.(
INJURIESFRACTURES, DISLOCATIONS, SRRAINS, AND STRAINS OPEN HOUNDS AND LACERATIONS CONTUSIONS AND SUPERFICIAL	330,590 173,279 44,827	81,605 36,158 10,484	85,997 48,417 13,311	95,810 48,527 14,995	67,178 40,177 6,036	177.9 93.3 24.1	175.6 77.8 22.6	162.6 91.5 25.2	168.7 85.4 26.4	226.8 135.6 20.4
INJURIES	49,825 62,660	13,468 21,496	11,184 13,085	17,639 14,649	7,534 13,431	26.8 33.7	29-0 46-3	21.l 24.7	31.1 25.8	25.4 45.3
ALL OTHER ACUTE CONDITIONS	258,137	55,474	70,829	92,251	39,582	138.9	119-4	133.9	162.4	133.6
MALE						:				!
ALL ACUTE CONDITIONS	659,107	150,921	173,881	205,298	129,007	731.7	676.7	668.0	752.4	892-1
INFECTIVE AND PARASITIC DISEASES	113,327	32,897	25,626	36,786	18,017	125-8	147.5	98.5	134.8	124-6
RESPIRATORY CONDITIONS————————————————————————————————————	269,246 144,541 93,231 31,474	52,705 36,883 7,346 8,476	77,255 40,693 30,557 6,005	74.565 41.739 21.673 11.153	64,722 25,227 33,655 5,841	298.9 160.5 103.5 34.9	236.3 165.4 32.9 38.0	296.3 156.3 117.4 23.1	273.3 153.0 79.4 40.9	447.6 174.4 232.7 40.4
DIGESTIVE SYSTEM CONDITIONS	31,741	6+015	7,235	12,952	5,540	35-2	27.0	27.3	47.5	38-3
INJURIES	177,102 91,714 25,957	43,506 18,076 4,809	44,721 24,243 9,646	55,768 27,784 9,008	33,107 21,611 2,494	196.6 101.8 28.8	195.1 81.0 21.6	171.8 93.1 37.1	204.4 101.8 33.0	228.5 149.4 17.2
OTHER CURRENT INJURIES	24,146 35,286	7,657 12,963	4,207 6,626	9,482 9,493	2,799 6,203	26-8 39-2	34-3 58-1	16.2 25.5	34.8 34.8	19.4 42.5
ALL OTHER ACUTE CONDITIONS	67,691	15,799	19,044	25,227	7,621	75.1	70.8	73.2	92•5	52.1
<u>FEMALE</u>										
ALL ACUTE CONDITIONS	848,584	198,235	239,391	249,720	161,240	886.5	820 - 1	891.0	845.9	1,063.7
INFECTIVE AND PARASITIC DISEASES	125,770	40,339	36,360	34,485	14,585	131.4	166.9	135.3	116.8	96-2
RESPIRATORY CONDITIONS—— UPPER RESPIRATORY CONDITIONS—— INFLUENZA—— OTHER RESPIRATORY CONDITIONS——	341,133 186,112 118,278 36,743	69,381 47,029 9,464 12,888	99,817 49,957 40,035 9,825	96,447 60,253 28,329 7,866	75,488 28,874 40,450 6,165	356.4 194.4 123.6 38.4	287.0 194.6 39.2 53.3	371.5 185.9 149.0 36.6	326.7 204.1 96.0 26.6	498.0 190.5 266.9 40.7
DIGESTIVE SYSTEM CONDITIONS	37,746	10,740	10,152	11,721	5,134	39.4	44.4	37.8	397	33.9
INJURIES	153,488 81,565 18,870	38,099 18,082 5,675	41,276 24,174 3,665	40,042 20,743 5,987	34,072 18,567 3,542	160.4 85.2 19.7	157.6 74.8 23.5	90.0 13.6	135.6 70.3 20.3	224.8 122.5 23.4
INJURIESOTHER CURRENT INJURIES	25,679 27,374	5,810 8,533	6,978 6,459	8,156 5,155	4,735 7,227	26.8 28.6	24.0 35.3	26.0 24.0	27.6 17.5	31.2 47.7
ALL OTHER ACUTE CONDITIONS	190,446	39,675	51,785	67,025	31,962	199.0	164-1	192.7	227.0	210.9

TABLE 22. DAYS OF BED DISABILITY ASSOCIATED WITH ACUTE CONDITIONS AND DAYS OF BED DISABILITY PER 100 FERSONS PER YEAR, BY REGION, SEX, AND CONDITION GROUP: UNITED STATES, JULY 1963-JUNE 1964

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II.

	given in	Appendix I.	Definitions of	of terms are g	iven in Appen	dix II				
					REGIO	ON				
SEX AND CONDITION GROUP	ALL REGIONS	NORTH- EAST	NORTH CENTRAL	SOUTH	WEST	ALL REGIONS	NORTH- EAST	NORTH CENTRAL	SOUTH	WEST
<u>BOTH SEXES</u>	DAYS	OF BED D	ISABILITY	IN THOUS	ANDS	DAYS 0	F BED DI:	SABILITY I PER YEAR	PER 100 P	ERSONS
ALL ACUTE CONDITIONS	642,741	136,430	176,489	200,096	129,726	345.9	293.5	333.6	352.3	438.0
INFECTIVE AND PARASITIC DISEASES	111,138	31,318	29,178	35,374	15,268	59.8	67.4	55.2	62.3	51.5
RESPIRATORY CONDITIONS	291,924 131,103 117,642 43,179	54,967 33,287 8,421 13,260	35,240 38,678	79,305 39,090 27,178 13,036	74,327 23,486 43,365 7,476	157.1 70.6 63.3 23.2	118.3 71.6 18.1 28.5	157.5 66.6 73.1 17.8	139.6 68.8 47.8 22.9	250.9 79.3 146.4 25.2
DIGESTIVE SYSTEM CONDITIONS	36,387	9,624	9,180	12,456	5,127	19.6	20.7	17.4	21.9	17.3
INJURIES FRACTURES, DISLOCATIONS, SPRAINS, AND STRAINS OPEN WOUNDS AND LACERATIONS CONTUSIONS AND SUPERFICIAL	96,493 48,138 12,653	15,415 5,659 2,471	27,638 17,473 2,647	34,297 13,437 5,950	19,142 11,569 1,584	51.9 25.9 6.8	33.2 12.2 5.3	52.2 33.0 5.0	60.4 23.7 10.5	64.6 39.1 5.3
INJURIES	13,375 22,327	2,307 4,978	2,327 5,191	6,942 7,968	1,799 4,190	7.2 12.0	5.0 10.7	4.4 9.8	12.2 14.0	6.1 14.1
ALL OTHER ACUTE CONDITIONS	106,798	25,106	27,166	38,665	1,5,861	57-5	54.0	51.4	68.1	53.6
MALE									i	
ALL ACUTE CONDITIONS	272,897	56,178	69,508	89,780	57,431	303.0	251.9	267.0	329-1	397.1
INFECTIVE AND PARASITIC DISEASES	51,041	14,479	11,945	17,486	7,130	56.7	64.9	45.9	64.1	49.3
RESPIRATORY CONDITIONS————————————————————————————————————	131,944 60,760 50,896 20,288	24,261 14,931 4,310 5,021	36,746 16,834 15,961 3,951	36,784 16,558 12,194 8,031	34,153 12,437 18,431 3,285	146.5 67.5 56.5 22.5	108.8 66.9 19.3 22.5	141.2 64.7 61.3 15.2	134.8 60.7 44.7 29.4	236.2 86.0 127.5 22.7
DIGESTIVE SYSTEM CONDITIONS	15,007	3,716	3,201	5,746	2,344	16.7	16.7	12.3	21.1	16.2
INJURIESFRACTURES, DISLOCATIONS,	50,812	8,370	11,789	19,657	10,995	56.4	37.5	45.3	72.0	76.0
SPRAINS, AND STRAINS————————————————————————————————————	25,442 6,669	2,797	7,912 #	7,813 3,152	6,920	28.2 7.4	12.5	30.4 *	28.6 11.6	47.9
OTHER CURRENT INJURIES	6,893 11,807	1,778 2,416	2,051	3,656 5,037	2,304	7.7 13.1	8.0 10.8	7.9	13.4 18.5	15.9
ALL OTHER ACUTE CONDITIONS	24,094	5,351	5,827	10,107	2,809	26.7	24.0	22.4	37.0	19.4
<u>FEMALE</u>										
ALL ACUTE CONDITIONS	369,844	80,252	106,980	110,316	72,295	386.4	332.0	398.2	373.7	476.9
INFECTIVE AND PARASITIC DISEASES	60,098	16,839	17,233	17,888	8,138	62.8	69.7	64.1	60.6	53.7
RESPIRATORY CONDITIONS———— UPPER RESPIRATORY CONDITIONS——— INFLUENZA————————————————————————————————————	159,980 70,343 66,745	30,706 18,356 4,111	46,579 18,406 22,717	42,521 22,532 14,984	40,174 11,049 24,934	167.1 73.5 69.7	127.0 75.9 17.0	173.4 68.5 84.5	144.0 76.3 50.8	265.0 72.9 164.5
OTHER RESPIRATORY CONDITIONS DIGESTIVE SYSTEM CONDITIONS	22,891	8,239 5,908	5,457	5,005 6,710	4,191 2,784	23.9	34.1 24.4	20.3	17.0 22.7	27.6 18.4
INJURIES	45,681	7,045	15,849	14,639	8,148	47.7	29.1	59.0	49.6	53.8
FRACTURES, DISLOCATIONS, SPRAINS, AND STRAINS OPEN WOUNDS AND LACERATIONS CCNTUSIONS AND SUPERFICIAL	22,697 5,983	2,862	9,562 #	5,624 2,798	4,649	23.7	11.8	35.6	19.1 9.5	30.7
OTHER CURRENT INJURIES	6,482 10,519	2,562	1,897 3,140	3,286 2,931	1,887	6-8	10.6	7.1 11.7	11.1 9.9	* 12.4
ALL OTHER ACUTE CONDITIONS	82,704	19,754	21,339	28,558	13,052	86.4	81.7	79.4	96.7	86.1

TABLE 23. INCIDENCE OF ACUTE CONDITIONS PER QUARTER AND NUMBER OF ACUTE CONDITIONS PER 100 PER SONS PER QUARTER, BY SEX AND AGE: UNITED STATES, JULY 1963-JUNE 1964

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and informatic

on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix I.

On the re	lability of the c	Sumates are g	I Ton In Appendi	x 1. Delililat	ons of terms are	given in Appen	——————————————————————————————————————	
	196	3	196	4	19	163	1964	·
SEX AND AGE	JULY - September	OCTOBER- DECEMBER	JANUARY- March	APRIL- JUNE	JULY- SEPT E MBER	OCTOBER- DECEMBER	JANUARY- MARCH	APRIL- JUNE
BOTH SEXES	INCIDE	NCE OF ACU IN THOUS	ITE CONDITI SANDS	ONS		OF ACUTE CO PERSONS PER	ONDITIONS P R QUARTER	ER 100
ALL AGES	68,161	104,972	118,628	95,597	36.9	56.6	63.7	51-
UNDER 5 YEARS	14,523	22,542	21,871	17,147	70-1	108.5	105.5	83.
5-14 YEARS	16,472	27,203	31,216	28,762	43.5	71.4	81.6	74.
15-24 YEARS	9,268	15,293	17,820	13,455	35.0	57.0	65.7	49.
25-44 YEARS	15,709	21,930	23,444	18,619	34.6	48.4	51.7	41.
45-64 YEARS	8,852	13,393	17,909	12,385	23.7	35.7	47.5	32.
65 & OVER YEARS	3,336	4,611	6,369	5,229	19.7	27.1	37.4	30.
MALE								
ALL AGES	31,896	49,014	54,512	44,761	35.6	54.5	60.4	49.
UNDER 5 YEARS	7,602	11,192	10,753	9,300	72.0	105.8	101.8	88.
5-14 YEARS	8,677	14,498	15,252	14,582	45.1	75.0	78.5	74.
15-24 YEARS	3,955	7,039	7,325	6,114	31.5	55.2	56.8	46.
25-44 YEARS	6,512	8,590	9,819	7,986	30.1	39.7	45.4	37.
45-64 YEARS	3,859	5,771	8,841	4,755	21.4	31.8	48.6	26.
65 & OVER YEARS	æ	×	2,523	×	ж	×	33.4	
FEMALE								
ALL AGES	36,265	55,959	64,116	50,836	38.1	58.6	66.9	52.
UNDER 5 YEARS	6•921	11,351	11,117	7,847	68.1	111-4	109.4	77.
5-14 YEARS	7,795	12,706	15,964	14,179	41.8	67.8	84.8	75.
15-24 YEARS	5,313	8,253	10,495	7,341	38.2	58.7	73-8	51.
25-44 YEARS	9,197	13,340	13,625	10,633	38.8	56.3	57.5	44.
45-64 YEARS	4,994	7,622	9,068	7,630	25.9	39.3	46.5	39.
65 & OVER YEARS	ж	-2,687	3,846	3,205	×	28.4	40.5	33.

NOTE: Excluded from these statistics are all conditions involving neither restricted activity nor medical attention.

TABLE 24. INCIDENCE OF ACUTE CONDITIONS PER QUARTER AND NUMBER OF ACUTE CONDITIONS PER 100 PERSONS PER QUARTER, BY SEX AND CONDITION GROUP: UNITED STATES, JULY 1963-JUNE 1964

bility o	f the estimates a	re given in App	pendix I.Defini	tions of terms	are given in App	oendix II		
	19	63	19	64	19	63	19	64
SEX AND CONDITION GROUP	JULY- SEPTEMBER	OCTOBER- DECEMBER	JANUARY- MARCH	APRIL- JUNE	JULY- SEPTEMBER	OCTOBER- DECEMBER	JANUARY- MARCH	APRIL- JUNE
BOTH SEXES	INCIDE	NCE OF ACL IN THOUS		ONS		OF ACUTE CO PERSONS PER		ER 100
ALL ACUTE CONDITIONS-	68,161	104,972	118,628	95,597	36.9	56.6	63.7	51.2
INFECTIVE AND PARASITIC DISEASES	7,598 24,907	10,090 65,856	16,507 73,027	21,088 40,511	4.1 13.5	5.4 35.5	8.9 39.2	11.3 21.7
CONDITIONSINFLUENZAOTHER RESPIRATORY	17,123 6,685	42,789 20,688	47,596 22,471	26,288 12,135	9.3 3.6	23.1 11.2	25.6 12.1	14.1 6.5
CONDITIONS DIGESTIVE SYSTEM CCNDITIONS	5,759	4,475	2,960 5,127	5,248	3.1	2.4	1.6 2.8	2.8
INJURIES ALL OTHER ACUTE CONDITIONS	16,195 13,702	12,978 11,574	11,156	14,897 13,853	8.8 7.4	7•0 6•2	6.0 6.9	8.0 7.4
MALE								
ALL ACUTE CONDITIONS-	31,896	49,014	54,512	44,761	35.6	54.5	60.4	49.5
INFECTIVE AND PARASITIC DISEASESRESPIRATORY CONDITIONS UPPER RESPIRATORY	3,557 11,059	4,858 30,477	7,811 33,835	9,776 18,438	4.0 12.3	5.4 33.9	8.7 37.5	10.8 20.4
CONDITIONS	7,501 2,994	20,177 9,220	22,551 10,208	11,889 5,408	8.4 3.3	22.4 10.3	25.0 11.3	13-1 6-0
CONDITIONS DIGESTIVE SYSTEM CONDITIONS INJURIES ALL OTHER ACUTE CONDITIONS	2,965 9,542 4,772	* 7,438 3,890	# 6,308 4,313	* 9,607 4,541	3.3 10.7 5.3	# 8.3 4.3	# 7.0 4.8	# 10•6 5•0
FEMALE								
ALL ACUTE CONDITIONS-	36,265	55,959	64,116	50,836	38.1	58.6	66.9	52.8
INFECTIVE AND PARASITIC DISEASES	4,041 13,848	5,232 35,379	8,697 39,192	11,312 22,073	, 4•2 14•6	5•5 3 7• 0	9•1 40•9	11.8 22.9
CONDITIONSINFLUENZAOTHER RESPIRATORY	9,622 3,691	22,613 11,469	25,044 12,263	14,399 6,727	10.1 3.9	23.7 12.0	26.1 12.8	15.0 7.0
CONDITIONS————— DIGESTIVE SYSTEM CONDITIONS—————	2,793	x	¥ 2,881	2,849	* 2.9	×	# 3 . 0	# 3.0
INJURIESALL OTHER ACUTE CONDITEINS	6,653 8,930	5,540 7,684	4,848 8,498	5,291 9,311	7.0 9.4	5.8 8.0	5.1 8.9	5.5 9.7
				-,		5.5		

NOTE: Excluded from these statistics are all conditions involving neither restricted activity nor medical attention.

TABLE 25. DAYS OF RESTRICTED ACITIVITY ASSOCIATED WITH ACUTE CONDITIONS PER QUARTER AND DA OF RESTRICTED ACTIVITY PER 100 PERSONS PER QUARTER, BY SEX AND AGE: UNITED STATES, JULY 196 JUNE 1964

on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II									
65V 4VD 445	19	63	190	54	19	63	19	64	
SEX AND AGE	JULY- SEPTEMBER	OCTOBER- DECEMBER	JANUARY- March	APRIL-	JULY- SEPTEMBER	OCTOBER- DECEMBER	JANUARY- March	APR I JUN	
BOTH SEXES	DAYS	OF RESTRI IN THOU		ITY		ESTRICTED ERSONS PER		ER 10	
ALL AGES	262,471	370,827	469,437	404,957	142.1	199.9	252.2	216	
UNDER 5 YEARS	31,720	47,062	62,305	53,269	153.1	226.6	300.7	257	
5-14 YEARS	46,338	81,403	112,323	104,381	122.3	213.8	293.7	271	
15-24 YEARS	31,150	54,959	62,154	55,369	117.7	205.0	229.1	201	
25-44 YEARS	69,425	85,333	87,439	72,894	153.0	188.3	192.9	160	
45-64 YEARS	49,796	66,840	92,804	71,332	133.3	178.1	246.3	188	
65 & OVER YEARS	34,042	35,230	52,413	47,712	200.9	207.4	307.5	278	
MALE									
ALL AGES	107,342	156,310	215,090	180,366	119.8	173.8	238.4	199	
UNDER 5 YEARS	15,861	22,986	30,340	29,970	150.3	217.2	287.3	284	
5-14 YEARS	21,950	39,890	54,908	56,844	114.0	206.2	282.7	291	
15-24 YEARS	12,377	22,920	23,225	21,295	98.6	179.9	180.0	163	
25-44 YEARS	21,606	31,109	35,603	26,425	99.8	143.9	164.7	122	
45-64 YEARS	24,290	30,527	48,018	30,496	134.5	168-4	264.1	167	
65 & OVER YEARS	11,259	8,878	22,995	15,336	149.7	117.8	304.7	202	
FEMALE									
ALL AGES	155,129	214,517	254,347	224,591	163.0	224.5	265.2	233	
UNDER 5 YEARS	15,860	24,076	31,964	23,299	156.0	236.3	314.5	230	
5-14 YEARS	24,388	41,513	57,415	47,537	130.8	221.6	305.1	251	
15-24 YEARS	18,773	32,039	38,929	34,074	135.0	227.8	273.7	237	
25-44 YEARS	47,819	54,223	51,836	46,468	201.7	228.7	218.7	196	
45-64 YEARS	25,506	36,314	44,786	40,836	132.1	187.1	229.8	208	
65 & OVER YEARS	22,784	26,352	29,418	32,376	241.9	278.8	309.8	339	

TABLE 26. DAYS OF RESTRICTED ACTIVITY ASSOCIATED WITH ACUTE CONDITIONS PER QUARTER AND DAYS OF RESTRICTED ACTIVITY PER 100 PERSONS PER QUARTER, BY SEX AND CONDITION GROUP: UNITED STATES, JULY 1963-JUNE 1964

bility of	f the estimates a	re given in App	endix I. Defin	itions of term	s are given in Ap	ppendix II		
CEN AND CONDITION COOLS	19	63	19	164	19	63	19	64
SEX AND CONDITION GROUP	JULY- SEPTEMBER	JOCTOBER- DECEMBER	JANUARY- MARCH	APRIL- JUNE	JULY- SEPTEMBER	OCTOBER- DECEMBER	JANUARY- March	APRIL- JUNE
BOTH SEXES	DAYS	OF RESTRI IN THOU		/ITY		ESTRICTED PERSONS PER	ACTIVITY P QUARTER	ER 100
ALL ACUTE CONDITIONS-	262,471	370,827	469,437	404,957	142.1	199.9	252.2	216.9
INFECTIVE AND PARASITIC DISEASES RESPIRATORY CONDITIONS UPPER RESPIRATORY CONDITIONS	28,606 63,227 35,780	33,807 187,932 103,365	79,757 229,822	96,927 129,399	15.5 34.2	18.2 101.3	42.8 123.5.	51.9 69.3
INFLUENZAOTHER RESPIRATORY	16,078	65,279	126,431 85,177	65,079 44,976	19.4 8.7	55.7 35.2	67 . 9 45 . 8	34.9 24.1
CONDITIONS DIGESTIVE SYSTEM CONDITIONS	11,370	19,289	18,215	19,345 17,489	6.2	10.4	9-8	10.4
INJURIESALL OTHER ACUTE	82,019	75,910	80,404	92,257	9.3 44.4	8.7 40.9	10.1 43.2	9•4 49•4
CONDITIONS	71,509	57,050	60,693	68,885	38.7	30.8	32.6	36.9
MALE			:				-	
ALL ACUTE CONDITIONS-	107,342	156,310	215,090	180,366	119.8	173.8	238.4	199-3
INFECTIVE AND PARASITIC DISEASES RESPIRATORY CONDITIONS UPPER RESPIRATORY	13,355 28,307	15,828 78,494	37,838 105,792	46,306 56,653	14.9 31.6	17.6 87.3	41.9 117.2	51.2 62.6
CONDITIONS INFLUENZA OTHER RESPIRATORY	15,256 7,729	43,798 27,744	58,573 38,911	26,914 18,847	17.0 8.6	48.7 30.8	64.9 43.1	29.7 20.8
CONDITIONS DIGESTIVE SYSTEM	×	6,952	8,308	10,893	36	7.7	9•2	12.0
INJURIESALL OTHER ACUTE	7,674 43,680	6,293 40,435	7,803 44,142	9,970 48,844	8.6 48.8	7.0 45.0	8•6 48•9	11.0 54.0
CONDITIONS	14,325	15,258	19,514	18,593	16.0	17.0	21.6	20.5
FEMALE	:							
ALL ACUTE CONDITIONS-	155,129	214,517	254,347	224,591	163.0	224.5	265.2	233.4
INFECTIVE AND PARASITIC DISEASES RESPIRATORY CONDITIONS UPPER RESPIRATORY	15,251 34,920	17,978 109,438	41,919 124,030	50,621 72,746	16.0 36.7	18.8 114.5	43.7 129.3	52.6 75.6
CONDITIONSINFLUENZAOTHER RESPIRATORY	20,523 8,349	59,567 37,535	67,857 46,266	38,165 26,129	21.6 8.8	62•3 39•3	70.8 48.2	39.7 27.2
CONDITIONSDIGESTIVE SYSTEM	6,048	12,336	9,907	8,452	6.4	12.9	10.3	8.8
INJURIESALL OTHER ACUTE	9,435 38,339	9,835 35,475	10,958 36,261	7,519 43,413	9.9 40.3	10.3 37.1	11.4 37.8	7.8 45.1
CONDITIONS	57,183	41,791	41,179	50,292	60.1	43.7	42.9	52.3

TABLE 27. DAYS OF BED DISABILITY ASSOCIATED WITH ACUTE CONDITIONS PER QUARTER AND DAYS OF B DISABILITY PER 100 PERSONS PER QUARTER, BY SEX AND AGE: UNITED STATES, JULY 1953-JUNE 1964

[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifications, and informal on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II									
	19	63	19	64	19	63	19	64	
SEX AND AGE	JULY- SEPTEMBER	OCTOBER- DECEMBER	JANUARY- March	APRIL- JUNE	JULY- SEPTEMBER	OCTOBER- DECEMBER	JANUARY- MARCH	APRI JUN	
BOTH SEXES	D	AYS OF BED IN THO	DISABILIT USANDS	Y		BED DISAE	BILITY PER QUARTER	100	
ALL AGES	109,812	152,081	207,462	173,385	59.4	82•0	111.5	92	
UNDER 5 YEARS	16,524	22,037	25,530	27,424	79.7	106.1	123.2	132	
5-14 YEARS	17,821	37,577	48,745	42,993	47.0	98.7	127.5	112	
15-24 YEARS	13,785	24,366	35,046	26,398	52.1	90.9	129.2	96	
25-44 YEARS	28,338	34,159	42,143	32,434	62.5	75.4	93.0	71	
45-64 YEARS	19,013	22,171	38,941	28,275	50.9	59.1	103.4	74	
65 & OVER YEARS	14,331	11,771	17,056	15,861	84.6	69.3	100.1	92	
MALE	44 052	43.200	90 045	72 770	£2 2	70.2	98.7	81	
ALL AGES	46,853	63,209	89,065	73,770	52.3	70.3	70 • 1	91	
UNDER 5 YEARS	9,195	10,996	12,003	14,849	87.1	103.9	113.7	141	
5-14 YEARS	6,932	17,692	23,306	21,358	36.0	91.5	120.0	109	
15-24 YEARS	×	9,542	10,758	9,367	×	74.9	83.4	71	
25-44 YEARS	9,023	12,533	16,557	12,135	41.7	58.0	76.6	56	
45-64 YEARS	11,190	8,754	19,051	10,233	62.0	48.3	104.8	56	
65 & OVER YEARS	6,077	×	7,390	×	80.8	×	97.9		
FEMALE							·		
ALL AGES	62,959	88,872	118,397	99,615	66.2	93.0	123.5	103	
UNDER 5 YEARS	7,329	11,041	13,527	12,575	72.1	108.4	133.1	124	
5-14 YEARS	10,889	19,885	25,439	21,636	58.4	106.1	135.2	114	
15-24 YEARS	9,350	14,824	24,289	17,031	67.2	105.4	170.8	118	
25-44 YEARS	19,315	21,626	25,586	20,299	81.5	91.2	107.9	85	
45-64 YEARS	7,823	13,417	19,890	18,042	40.5	69.1	102.0	92	
65 & OVER YEARS	8,255	8,079	9,666	10,032	87.7	85.5	101.8	105	

TABLE 28. DAYS OF BED DISABILITY ASSOCIATED WITH ACUTE CONDITIONS PER QUARTER AND DAYS OF BED DISABILITY PER 100 PERSONS PER QUARTER, BY SEX AND CONDITION GROUP: UNITED STATES, JULY 1963-JUNE 1964

SEX AND CONDITION GROUP JULY	bility	of the estimates	are given in A	ppendix I.Defin	itions of term	s are given in A	ppendix II		
DAYS OF SEPTEMBER DAYS OF SED DISABILITY DAYS OF SED DISABILITY PER 100		19	63	19	64	19	63	19	64
### ALL ACUTE CONDITIONS— ALL ACUTE CONDITIONS— 109,812 152,081 207,462 173,385 59.4 82.0 111.5 92.9 INFECTIVE AND PARASITIC DISEASES— ESPIRATORY CONDITIONS— 14,258 39,770 49,875 27,200 7.7 21.4 26.8 14.6 17.0 59.1 34.8 UPPER RESPIRATORY CONDITIONS— 14,258 39,770 49,875 27,200 7.7 21.4 26.8 14.6 15.5 26.1 13.8 UPPER RESPIRATORY CONDITIONS— 14,258 39,770 49,875 27,200 7.7 21.4 26.8 14.6 13.5 26.1 13.8 UPPER RESPIRATORY CONDITIONS— 17,830 8,941 10,842 4,771 11,958 3.6 7.0 6.2 6.4 UNDITIONS— 7,830 8,941 10,842 4,771 4,2 4.8 5.8 4.7 INJURIES— 25,451 16,307 27,065 27,670 13.8 8.8 14.5 14.8 ALL ACUTE CONDITIONS— 30,228 21,785 27,538 27,247 16.4 11.7 14.8 14.6 **MALE** ALL ACUTE CONDITIONS— 13,561 37,801 49,803 30,783 15.1 42.0 55.2 34.0 UPPER RESPIRATORY CONDITIONS— 46,853 63,209 89,065 73,770 52.3 70.3 98.7 81.5 INFECTIVE AND PARASITIC DISEASES— RESPIRATORY CONDITIONS— 13,561 37,801 49,803 30,783 15.1 42.0 55.2 34.0 UPPER RESPIRATORY CONDITIONS— 6,210 18,802 22,869 12,879 6.9 20.9 25.3 14.2 UPPER RESPIRATORY CONDITIONS— 14,544 21,536 10,750 % 16.2 23.9 11.9 UPPER RESPIRATORY CONDITIONS— CONDITIONS— 2 % % % % % % % % % % % % % % % % % %	SEX AND CONDITION GROUP								
INFECTIVE AND PARASITIC 16,592 17,915 31,920 44,711 9.0 9.7 17.1 23.9 34.8 10.008 16.1 47.0 59.1 34.8 10.008 16.1 47.0 6.2 6.4	BOTH SEXES	ι			Υ				100
DISEASES 16,992 17,915 31,920 44,711 9.0 9.7 17.1 23.9	ALL ACUTE CONDITIONS-	109,812	152,081	207,462	173,385	59.4	82.0	111.5	92.9
THELDENZA— OTHER RESPIRATORY CONDITIONS— C	DISEASES	29,711	87,133	110,098	64,983	16-1	47.0	59.1	34.8
DIGESTIVE SYSTEM CONDITIONS 25,451 16,307 27,065 27,670 13.8 8.8 14.5 14.8 14.5 14.8 14.6 14.6 11.7 14.8 14.6 14.	OTHER RESPIRATORY			48,649	25,824	4.8	18.5	26.1	13.8
MALE	CONDITIONS						4-8	5.8	4.7
ALL ACUTE CONDITIONS— 46,853 63,209 89,065 73,770 52.3 70.3 98.7 81.5 INFECTIVE AND PARASITIC DISEASES— CONDITIONS— 7,884 9,183 14,725 19,249 8.8 10.2 16.3 21.3 RESPIRATORY CONDITIONS— 6,210 18,802 22,869 12,879 6.9 20.9 25.3 14.2 UNPER RESPIRATORY CONDITIONS— 2 2 2,869 12,879 6.9 20.9 25.3 14.2 UNFLUENZA————————————————————————————————————		30,228	21,785	27,538	27,247	16.4	11.7	14.8	14.6
INFECTIVE AND PARASITIC DISEASES— OTHER RESPIRATORY CONDITIONS— CO	MALE								
DISEASES	ALL ACUTE CONDITIONS-	46,853	63,209	89,065	73,770	52.3	70.3	98.7	81.5
INFLUENZA————————————————————————————————————	DISEASES	13,561	37,801	49,800	30,783	15.1	42.0	55.2	34.0
DIGESTIVE SYSTEM	INFLUENZAOTHER RESPIRATORY	×	14,544	21,536		×	16.2		
ALL OTHER ACUTE CONDITIONS————————————————————————————————————	DIGESTIVE SYSTEM CONDITIONS	ĸ	×	×	×				
ALL ACUTE CONDITIONS— 62,959 88,872 118,397 99,615 66.2 93.0 123.5 103.5 INFECTIVE AND PARASITIC OISEASES—————————————————————————————————	ALL OTHER ACUTE			•					
INFECTIVE AND PARASITIC OISEASES—————————————————————————————————	FEMALE								
DISEASES	ALL ACUTE CONDITIONS-	62,959	88,872	118,397	99,615	66.2	93.0	123.5	103.5
INFLUENZA	DISEASES RESPIRATORY CONDITIONS UPPER RESPIRATORY	16,150	49,332	60,298	34,200	17.0	51.6	62.9	35.5
DIGESTIVE SYSTEM CONDITIONS	INFLUENZA OTHER RESPIRATORY	x	19,825	27,113	15,074	×	20.7	28.3	15.7
ALL OTHER ACUTE	DIGESTIVE SYSTEM CONDITIONS	×	×	7,156	×	×	×	7.5	¥
	ALL OTHER ACUTE				i				

TABLE 29. INCIDENCE OF ACUTE RESPIRATORY CONDITIONS AND ASSOCIATED DISABILITY DAYS PER QUARTER AND NUMBER OF ACUTE RESPIRATORY CONDITIONS PER 100 PERSONS PER QUARTER BY AGE: UNITED STATES JULY 1963-JUNE 1964

	19	63	196	54	19	63	19	64		
AGE	JULY- SEPTEMBER	OCTOBER- DECEMBER	JANUARY- MARCH	APRIL- JUNE	JULY- SEPTEMBER	OCTOBER- DECEMBER	JANUARY- March	APRIL- JUNE		
		NCE OF ACU NDITIONS I			NUMBER OF PER 1		PIRATORY CO PER QUART			
ALL AGES	24,907	65,856	73,027	40,511	13.5	35.5	39.2	21.		
UNDER 5 YEARS	6,139	15,642	13,597	7,887	29.6	75.3	65.6	38.		
5-14 YEARS	6,228	17,625	17,586	10,253	16.4	46.3	46.0	26.		
15-24 YEARS	2,936	8,532	10,955	5,473	11.1	31.8	40.4	19.		
25-44 YEARS	5,330	13,198	14,098	8,473	11-8	29.1	31.1	18.		
45-64 YEARS	3,168	7,881	12,341	5,732	8.5	21.0	32.8	15.		
65 & OVER YEARS	×	2,977	4,449	2,694	ж	17.5	26.1	15.		
	DAYS	OF RESTRI		'ITY	DAYS OF RESTRICTED ACTIVITY PER 100 PERSONS PER QUARTER					
ALL AGES	63,227	187,932	229,822	129,399	34.2	101.3	123.5	69.		
UNDER 5 YEARS	11,248	32,081	35,395	23,306	54.3	154.5	170.8	112.		
5-14 YEARS	16,068	50,244	50,501	28,793	42.4	131.9	132.1	75.		
15-24 YEARS	7,229	20,347	29,200	13,903	27.3	75.9	107.6	50•		
25-44 YEARS	12,411	38,120	42,277	23,199	27.4	84.1	93.3	51.		
45-64 YEARS	9,811	30,978	47,706	20,407	26.3	82.5	126.6	54.		
65 & OVER YEARS	6,459	16,163	24,742	19,790	38.1	95.1	145.1	115.		
	DA	YS OF BED IN THOU		,		BED DISAE	BILITY PER QUARTER	100		
ALL AGES	29,711	87,133	110,098	64,983	16.1	47.0	59.1	34.		
UNDER 5 YEARS	×	15,327	14,057	12,642	36	73.8	67.8	61.		
5-14 YEARS	7,970	25,819	25,454	15,665	21.0	67.8	66.6	40.		
15-24 YEARS	×	11,054	17,270	8,264	×	41.2	63.7	30.		
25-44 YEARS	6,003	18,041	23,078	10,535	13.2	39.8	50.9	23.		
45-64 YEARS	×	11,970	20,782	11,183	×	31.9	55.2	29.		
65 & OVER YEARS	×	×	9,456	6,694	×	×	55.5	39.		

NOTE: Excluded from these statistics are all conditions involving neither restricted activit nor medical attention.

TABLE 30. INCIDENCE OF INJURIES AND ASSOCIATED DISABILITY DAYS PER QUARTER AND NUMBER OF INJURIES PER 100 PERSONS, BY AGE: UNITED STATES, JULY 1963-JUNE 1964

on the reliability of the estimates are given in Appendix 1. Definitions of terms are given in Appendix II										
	19	63	19	64	19	63	19	64		
AGE	JULY- SEPTEMBER	OCTOBER- DECEMBER	JANUARY- March	APRIL- JUNE	JULY- SEPTEMBER	OCTOBER- DECEMBER	JANUARY- March	APRIL- JUNE		
	I	NCIDENCE O				R OF INJUR ERSONS PER		0		
ALL AGES	16,195	12,978	11,156	14,897	8.8	7.0	6.0	8.0		
UNDER 15 YEARS	6,672	4,372	3,374	5,924	11.4	7.4	5.7	10.0		
15-44 YEARS	6,247	5,738	5,102	5,706	8.7	8.0	7.0	7.8		
45 & OVER YEARS	3,277	2,867	2,680	3,268	6.0	5.3	4.9	5.9		
	DAYS	OF RESTRI		ITY	DAYS OF RESTRICTED ACTIVITY PER 100 PERSONS PER QUARTER					
ALL AGES	82,019	75,910	80,404	92,257	44.4	40•9	43.2	49.4		
UNDER 15 YEARS	16,754	11,541	8,723	15,862	28.6	19.6	14.8	26.9		
15-44 YEARS	28,726	35,088	33,849	36,597	40•0	48.6	46.7	50.3		
45 & OVER YEARS	36,540	29,281	37,832	39,799	67.3	53.7	69.1	72.5		
	DA	YS OF BED IN THOU				BED DISAB RSONS PER		100		
ALL AGES	25,451	16,307	27,065	27,670	13.8	8.8	14.5	14.8		
UNDER 15 YEARS	3 2	ж	ж	×	×	×	×	n		
15-44 YEARS	7,885	8,295	13,362	11,025	11.0	11.5	18.4	15.2		
45 & OVER YEARS	13,478	6,709	12,095	12,629	24.8	12.3	22.1	. 23.0		

 ${\tt NOTE:}$ Excluded from these statistics are all conditions involving neither restricted activity nor medical attention.

TABLE 31. POPULATION USED IN OBTAINING RATES SHOWN IN THIS PUBLICATION, BY RESIDENCE, REGION, SEX AND AGE: UNITED STATES, JULY 1963-JUNE 1964

on the reliability	of the estimat	es are given	in Appendix I	. Definitions	of terms are gi	ven in Append	ix IIJ	
		RESID	ENCE			REGI	ON	
SEX AND AGE	TOTAL	ALL SMSA	NON-SMSA NONFARM	NON-SMSA FARM	NORTH- East	NORTH CENTRAL	SOUTH	WES.
BOTH SEXES			PC	PULATION	IN THOUSAN	IDS		
ALL AGES	185,797	118,731	55,346	11,720	46,476	52,898	56,804	29,6
UNDER 5 YEARS	20,721 38,160 26,960	13,193 23,877 16,920	6,354 11,562 8,379 12,596	1,174 2,720 1,660	4,869 8,885 6,465 11,626	5,964 11,085 7,337 12,690	6,467 11,909 9,016 13,518	3,4, 6,2 4,1 7,5
25-44 YEARS	45,333 54,624 37,602 17,022	30,372 34,368 24,399 9,970	16,455 10,562 5,893	2,365 3,800 2,641 1,159	14,632 10,118 4,514	15,823 10,670 5,152	15,894 11,083 4,811	8,2 5,7 2,5
UNDER 6 YEARS	24,973 40,956 65,244	15,901 25,496 42,966	7,637 12,410 18,844	1,435 3,051 3,434	5,790 9,681 16,373	7,221 11,786 18,069	7,807 12,857 20,246	4,1 6,6 10,5
MALE								
ALL AGES	90,078	57,266	26,737	6,075	22,303	26,029	27,284	14,4
UNDER 5 YEARS	10,558 19,382 12,815 21,627 25,696 18,153 7,544	6,751 12,071 8,039 14,455 15,950 11,676 4,273	3,198 5,889 3,881 6,001 7,768 5,114 2,654	609 1,422 896 1,171 1,978 1,362 616	2,417 4,438 3,068 5,581 6,799 4,882	3,117 5,637 3,572 6,206 7,498 5,192 2,306	3,284 6,061 4,241 6,301 7,397 5,228 2,169	1,7 3,2 1,9 3,5 4,0 2,8 1,1
UNDER 6 YEARS	12,679 20,830 30,872	8,097 12,915 20,304	3,854 6,313 8,802	729 1,602 1,766	2,866 4,882 7,757	3,758 6,002 8,772	3,960 6,524 9,403	2,0° 3,4; 4,9;
FEMALE								
ALL AGES	95,720	61,466	28,610	5,644	24,173	26,869	29,520	15,1
UNDER 5 YEARS	10,163 18,778 14,145 23,706 28,928 19,449 9,479	.6,442 11,806 8,882 15,918 18,419 12,722 5,696	3,156 5,673 4,499 6,595 8,687 5,448 3,239	566 1,299 764 1,194 1,822 1,279 543	2,451 4,446 3,397 6,045 7,833 5,236 2,597	2,847 5,448 3,766 6,484 8,325 5,478 2,847	3,183 5,848 4,775 7,217 8,497 5,855 2,642	1,6 3,0 2,2 3,9 4,2 2,8 1,3
UNDER 6 YEARS	12,294 20,126 34,373	7,804 12,581 22,662	3,783 6,097 10,042	706 1,449 1,668	2,924 4,799 8,616	3,464 5,783 9,297	3,848 6,333 10,842	2,0 3,2 5,6

NOTE: For official population estimates for more general use, see Bureau of the Censt reports on the civilian population of the United States, in Current Population Reports, Serie P-20, P-25, and P-60.

TABLE 32. POPULATION OF CURRENTLY EMPLOYED PERSONS, BY SEX AND AGE: UNITED STATES, JULY 1963-JUNE 1964

AGE	BOTH SEXES	MALE	FEMALE	
	POPULATION IN THOUSANDS			
ALL AGES-17 YEARS & OVER	70,122	45,730	24,392	
17-44 YEARS	41,614 28,508	27,201 18,529	14,413 9,979	

NOTE: For official population estimates for more general use, see Bureau of Census reports on the civilian population of the United States, in <u>Current Population Reports</u>, Series P-20, P-25, and P-60; and Bureau of Labor Statistics monthly report, <u>Employment and Earnings</u>.

TABLE 33. POPULATION USED IN OBTAINING RATES SHOWN IN THIS PUBLICATION, BY QUARTER, SEX, AND ACCURATE UNITED STATES, JULY 1963-JUNE 1964

on the reliability of the estimates are given in A	Appendix 1. Definition	ons of terms are grad	an in Appendix ii	
	1963		1964	
SEX AND AGE	JULY- SEPTEMBER	OCTOBER- DECEMBER	JANUARY- MARCH	APRIL- JUNE
BOTH SEXES	POPULATION IN THOUSANDS			
ALL AGES	184,754	185,502	186,135	186,7
UNDER 5 YEARS	20,720	20,769	20,722	20,0
5-14 YEARS	37,902	38,079	38,241	38,4
15-24 YEARS	26,461	26,807	27,128	27,4
25-44 YEARS	45,361	45,325	45,324	45•:
45-64 YEARS	37,369	37,532	37,673	37.1
65 & OVER YEARS	16,941	16,990	17,046	17,
MALE				
ALL AGES	89,590	89,944	90,239	90•
UNDER 5 YEARS	10,553	10,581	10,560	10•!
5-14 YEARS	19,253	19,342	19,423	19,
15-24 YEARS	12,551	12,741	12,906	13,(
25-44 YEARS	21,656	21,618	21,619	21,0
45-64 YEARS	18,054	18,124	18,182	18.
65 & OVER YEARS	7,523	7,539	7,548	7•!
FEMALE				
ALL AGES	95,163	95,558	95,896	96,
UNDER 5 YEARS	10,167	10,188	10,162	10,
5-14 YEARS	18,649	18,737	18,818	18.
15-24 YEARS	13,909	14,066	14,222	14.
25-44 YEARS	23,705	23,707	23,705	23,
45-64 YEARS	19,315	19,408	19,491	19,
65 & OVER YEARS	9,418	9,451	9,497	9,:
	i		.	<u> </u>

NOTE: For official population estimates for more general use, see Bureau of the Census r ports on the civilian population of the United States, in <u>Current Population Reports</u>, Series P-P-25, and P-60.

APPENDIX I

TECHNICAL NOTES ON METHODS

Background of This Report

This report is one of a series of statistical reports prepared by the National Health Survey. It is based on information collected in a continuing nationwide sample of households in the Health Interview Survey, a major part of the program.

The Health Interview Survey utilizes a questionnaire which, in addition to personal and demographic characteristics, obtains information on illnesses, injuries, chronic conditions and impairments, and other health topics. As data relating to each of these various broad topics are tabulated and analyzed, separate reports are issued which cover one or more of the specific topics. The present report is based on the consolidated sample for 52 weeks of interviewing ending June 1964.

The population covered by the sample for the Health Interview Survey is the civilian, noninstitutional population of the United States living at the time of the interview. The sample does not include members of the Armed Forces, U.S. nationals living in foreign countries, or crews of vessels. It should also be noted that the estimates shown do not represent a complete inventory of injuries for the specified calendar period since no adjustment has been made for persons who incurred injuries during the 2-week-recall period but who died prior to the interview.

Statistical Design of the Health Interview Survey

General plan.—The sampling plan of the survey follows a multistage probability design which permits a continuous sampling of the civilian population of the United States. The first stage of this design consists of drawing a sample of 357 from about 1,900 geographically defined primary sampling units (PSU's) into which the United States has been divided. A PSU is a county, a group of contiguous counties, or a standard metropolitan statistical area.

With no loss in general understanding, the remaining stages can be combined and treated in this discussion as an ultimate stage. Within PSU's then, ultimate stage units called segments are defined in such a man-

ner that each segment contains an expected nine households. A segment consists of a cluster of neighboring households or addresses. Two general types of segments are used: (1) area segments which are defined geographically, and (2) B segments which are defined from a list of addresses from the Decennial Census and Survey of Construction. Each week a random sample of about 90 segments is drawn. In the approximate 800 households in those segments, household members are interviewed concerning factors related to health.

Since the household members interviewed each week are a representative sample of the population, samples for successive weeks can be combined into larger samples. Thus the design permits both continuous measurement of characteristics of high incidence or prevalence in the population and, through the larger consolidated samples, more detailed analysis of less common characteristics and smaller categories. The continuous collection has administrative and operational advantages as well as technical assets, since it permits field work to be handled with an experienced, stable staff.

Sample size and geographic detail.—The national sample plan for the 12-month period ending June 1964 included about 134,000 persons from 42,000 households in about 4,700 segments.

The overall sample was designed in such a fashion that tabulations can be provided for each of the major geographic regions and for urban and rural sectors of the United States.

Collection of data.—Field operations for the household survey are performed by the Bureau of the Census under specifications established by the National Center for Health Statistics. In accordance with these specifications the Bureau of the Census selects the sample; conducts the field interviewing as an agent of the Center; and performs a manual edit and coding of the questionnaires. The Health Survey, using Center electronic computers, carries out further editing and tabulates the edited data.

Estimating methods.—Each statistic produced by the survey—for example, the number of work-loss days occurring in a specified period—is the result of two stages of ratio estimation. In the first of these, the control factor is the ratio of the 1960 decennial population count to the 1960 estimated population in the Na-

tional Health Survey's first-stage sample of PSU's. These factors are applied for some 25 color-residence classes.

Later, ratios of sample-produced estimates of the population to official Bureau of the Census figures for current population in about 60 age-sex-color classes are computed and serve as second-stage factors for ratio estimating.

The effect of the ratio-estimating process is to make the sample more closely representative of the population by age, sex, color,\and residence, thus reducing sampling variance.

As noted, each week's sample represents the population living during that week and characteristics of that population. Consolidation of samples over a time period, say a calendar quarter, produces estimates of average characteristics of the U.S. population for that calendar quarter. Similarly, population data for a year are averages of the four quarterly figures.

For statistics measuring the number of occurrences during a specified time period, such as the number of cases of influenza or number of bed-disability days. a similar computational procedure is used, but the statistics are interpreted differently. For these items, the questionnaire asks for the respondent's experience during the 2 calendar weeks prior to the week of interview. In such instances the estimated quarterly total for the statistic is simply 6.5 times the average 2-week estimate produced by the 13 successive samples taken during the period. The annual total is the sum of the four quarters. Thus, the experience of persons interviewed during a year-experience which actually occurred for each person in a 2-calendar-week interval prior to week of interview-is treated as though it measured the total of such experience during the year. Such interpretation leads to no significant bias.

General Qualifications

Nonresponse.—Data were adjusted for nonresponse by a procedure which imputes to persons in a household which was not interviewed the characteristics of persons in households in the same segment which were interviewed. The total noninterview rate was 5 percent; 1 percent was refusal; and the remainder was primarily due to the failure to find any eligible household respondent after repeated trials.

The interview process.—The statistics presented in this report are based on replies secured in interviews of persons in the sampled households. Each person 19 years of age and over, available at the time of interview, was interviewed individually. Proxy respondents within the household were employed for children and for adults not available at the time of the interview, provided the respondent was closely related to the person about whom information was being obtained.

There are limitations to the accuracy of diagnost and other information collected in household interviews. For diagnostic information, the household respondent can, at best, pass on to the interviewer on the information the physician has given the family. For conditions not medically attended, diagnostic information is often no more than a description of symptom However, other facts, such as the number of disability days caused by the condition, can be obtained more accurately from household members than from any other source, since only the persons concerned are in a postion to report this information.

Rounding of numbers.—The original tabulations which the data in this report are based show all est mates to the nearest whole unit. All consolidation were made from the original tabulations using the entimates to the nearest unit. In the final published table the figures are rounded to the nearest thousand, althoug these are not necessarily accurate to that detail. Do vised statistics such as rates and percent distributions are computed after the estimates on which the are based have been rounded to the nearest thousan

Population figures.—Some of the published table include population figures for specified categories. E: cept for certain overall totals by age and sex, which are adjusted to independent estimates, these figure are based on the sample of households in the Nation Health Survey. These are given primarily to providenominators for rate computation, and for this put pose are more appropriate for use with the accon panying measures of health characteristics than other population data that may be available. In some instance these will permit users to recombine published da into classes more suitable to their specific needs. Wi the exception of the overall totals by age and sex, men tioned above, the population figures differ from corre sponding figures (which are derived from | different sources) published in reports of the Bureau of the Cei sus. For population data for general use, see the off cial estimates presented in Bureau of the Census re ports in the P-20, P-25, and P-60 series.

Reliability of Estimates


Since the estimates are based on a sample, the will differ somewhat from the figures that would have been obtained if a complete census had been taken using the same schedules, instructions, and interviewing personnel and procedures. As in any survey, the result are also subject to measurement error.

The standard error is primarily a measure sampling variability, that is, the variations that mig occur by chance because only a sample of the population is surveyed. As calculated for this report, t standard error also reflects part of the variation whice arises in the measurement process. It does not include estimates of any biases which might lie in the dat

The chances are about 68 out of 100 that an estimate from the sample would differ from a complete census by less than the standard error. The chances are about 95 out of 100 that the difference would be less than twice the standard error and about 99 out of 100 that it would be less than 2½ times as large.

The relative standard error of an estimate is obtained by dividing the standard error of the estimate by the estimate itself and is expressed as a percentage of the estimate. Charts from which the relative standard errors can be determined for estimates shown in the

report are presented in *Vital and Health Statistics*, Series 10, No. 10, pp. 51-56. In order to derive relative errors which would be applicable to a wide variety of health statistics and which could be prepared at a moderate cost, a number of approximations were required. As a result, the charts provide an estimate of the approximate relative standard error rather than the precise error for any specific aggregate or percentage. Additional information and a guide to the use of the standard error charts are included in the publication referred to above.



APPENDIX II

DEFINITIONS OF CERTAIN TERMS USED IN THIS REPORT

Terms Relating to Conditions

Condition.—A morbidity condition, or simply a condition, is any entry on the questionnaire which describes a departure from a state of physical or mental wellbeing. It results from a positive response to one of a series of "illness-recall" questions. In the coding and tabulating process, conditions are selected or classified according to a number of different criteria, such as, whether they were medically attended; whether they resulted in disability; whether they were acute or chronic; or according to the type of disease, injury, impairment, or symptom reported. For the purposes of each published report or set of tables, only those conditions recorded on the questionnaire which satisfy certain stated criteria are included.

Acute conditions are classified by type according to the International Classification of Diseases, 1955 Revision, with certain modifications adopted to make the code more suitable for a household-interview-type survey.

Acute condition.—An acute condition is defined as a condition which has lasted less than 3 months and which has involved either medical attention or restricted activity. Because of the procedures used to estimate incidence, the acute conditions included in this report are the conditions which had their onset during the 2 weeks prior to the interview week and which involved either medical attention or restricted activity during that 2-week period. However, it excludes certain conditions which are always classified as chronic (listed below) even though the onset occurred within 3 months.

Conditions always classified as chronic:

Stomach ulcer Asthma Any other chronic stomach Hav fever trouble Tuberculosis Kidney stones or chronic Chronic bronchitis kidney trouble Repeated attacks of sinus Arthritis or rheumatism trouble Mental illness Rheumatic fever Hardening of the arteries Diabetes High blood pressure Thyroid trouble or goiter Heart trouble Any allergy Epilepsy Stroke Trouble with varicose Chronic nervous trouble Cancer veins

Hemorrhoids or piles
Tumor, cyst, or growth
Chronic gallbladder or
liver trouble
Deafness or serious
trouble with hearing
Serious trouble with seeing, even when wearing
glasses
Cleft palate
Any speech defect
Missing fingers, hand, or
arm—toes, foot, or leg
Palsy

Chronic skin trouble
Hernia or rupture
Prostate trouble
Paralysis of any kind
Repeated trouble with
back or spine
Club foot
Permanent stiffness or
any deformity of the foc
leg, fingers, arm, or
back
Condition present since
birth

Condition groups.—Conditions are classified a cording to the International Classification of Disease 1955 Revision, with certain modifications adopted make the code more suitable for a household-intervis survey. In this report, all tables which have data classified by type of condition employ a five-category regrouping plus several selected subgroups. The International Classification code numbers included in eacategory are shown below:

International

Condition Groups		Classification		
		Code Numbers		
Ι	Infective and parasitic			
	diseases	020-138		
II	Respiratory conditions	470-501, 511, 517-525, 527, 783		
	Upper respiratory	470-475, 511, 517		
	Influenza	480-483		
	Other respiratory	490-501, 518-525. 527, 783		
II	Digestive system			
	conditions	530-539, 543-553, 570, 571, 573-587, 784, 785		
IV	Injuries	N800-1885, N890-N895		
		N900-N994, N996-N999		
v	Other conditions	All other acute code numbers		

.

Onset of condition.—A condition is considered to have had its onset when it was first noticed. This could be the time the person first felt sick or became injured, or it could be the time when the person or his family was first told by a physician that he had a condition of which he was previously unaware.

Incidence of conditions.—The incidence of conditions is the estimated number of conditions having their onset in a specified time period. As previously mentioned, minor acute conditions, involving neither restricted activity nor medical attention are excluded from the statistics. The incidence data shown in some reports are further limited to various subclasses of conditions, such as "incidence of conditions involving bed disability."

Activity-restricting condition.—An activity restricting condition is a condition which has caused at least 1 day of restricted activity during the 2 calendar weeks before the interview week. (See definition of "Restricted-activity day.") The incidence of acute activity-restricting conditions is estimated from the number of such conditions reported as having started in the 2-week period, but a condition which did not result in restricted activity until after the end of the 2-week period in which it had its onset is not included.

Bed-disabling condition.—A condition involving at least 1 day of bed disability is called a bed-disabling condition. (See definition of "Bed-disability day.") The incidence of acute bed-disabling conditions is defined in a manner analogous to the incidence of acute activity-restricting conditions.

Medically attended condition.—A condition is considered to be medically attended if a physician has been consulted about it either at its onset or at any time thereafter. Medical attention includes consultation either in person or by telephone for treatment or advice. Advice from the physician transmitted to the patient through the nurse is counted as well as visits to physicians in clinics or hospitals. If during the course of a single visit the physician is consulted about more than one condition for each of several patients, each condition of each patient is counted as medically attended.

Discussions of a child's condition by the physician and a responsible member of the household are considered as medical attention even if the child was not seen at that time.

For the purpose of this definition, the term "physician" includes doctors of medicine and osteopathic physicians.

Terms Relating to Disability

<u>Disability</u>.—Disability is the general term used to describe any temporary or long-term reduction of a person's activity as a result of an acute or chronic condition.

• Disability days are classified according to whether they are days of restricted activity, bed-days, workloss days, or school-loss days. All days of bed disability are, by definition, days of restricted activity. The converse form of this statement is, of course, not true. Days lost from work and days lost from school are also days of restricted activity for the working and schoolage populations. Hence, restricted activity is the most inclusive term used in describing disability days.

Condition-days of restricted activity, bed disability, etc.—Condition-days of restricted activity, bed disability, and so forth are days of the various forms of disability associated with any one condition. Since any particular day of disability may be associated with more than one condition, the sum of days for all conditions adds to more than the total number of person-days of disability.

Restricted-activity day.—A day of restricted activity is one on which a person substantially reduces the amount of activity normal for that day because of a specific illness or injury. The type of reduction varies with the age and occupation of the individual as well as with the day of the week or season of the year. Restricted activity covers the range from substantial reduction to complete inactivity for the entire day.

Bed-disability day.—A day of bed disability is one on which a person stays in bed for all or most of the day because of a specific illness or injury. It is considered to be a day only if the period of bed disability includes more than half of the daylight hours. All hospital days for inpatients are considered to be days of bed disability even if the patient was not actually in bed at the hospital.

Work-loss day.—A day lost from work is a normal working day on which a person did not work at his job or business because of a specific illness or injury. If the person's regular work day is less than a whole day and the entire work day was lost, it would be counted as a whole work day lost. The number of days lost from work is determined only for persons 17 years of age or over who reported that at any time during the 2-week period covered by the interview they either worked at, or had a job or business. (See definition of "Currently employed persons.")

School-loss day.—A day lost from school is a normal school day on which a child did not attend school because of a specific illness or injury. The number of days lost from school is determined only for children 6-16 years of age.

Demographic Terms

Age.—The age recorded for each person is the age at last birthday. Age is recorded in single years and grouped in a variety of distributions depending upon the purpose of the table.

Quarter.—The quarters used by the National Health Survey are actually 13-week periods rather than 3 calendar months. Since each 13-week period begins on a Monday and ends on a Sunday, the actual dates of the beginning and end of each 13-week period may overlap into another calendar quarter. Therefore, the time periods in the table headings are the approximate rather than the precise periods during which the interviewing was conducted.

Currently employed persons.—Currently employed persons are all persons 17 years of age or over who reported that at any time during the 2-week period covered by the interview they either worked at, or had a job or business. Current employment includes paid work as an employee of someone else, self-employment in business, farming, or professional practice, and unpaid work in a family business or farm. Persons who were temporarily absent from their job or business because of a temporary illness, vacation, strike, or bad weather are considered as currently employed if they expected to work as soon as the particular event causing their absence no longer existed.

Free-lance workers are considered as having a job if they had a definite arrangement with one or more employees to work for pay according to a weekly or monthly schedule, either full-time or part-time. Excluded from the currently employed population are such persons who have no definite employment schedule but who work only when their services are needed.

Also excluded from the currently employed population are (1) persons who were not working, even though having a job or business, but were on layoff or looking for work, (2) persons receiving revenue from an enterprise in whose operation they did not participate, (3) persons doing housework or charity work for which they received no pay, and (4) seasonal workers during the unemployment season.

The number of currently employed persons estimated by the National Health Survey (NHS) will differ from the estimates prepared by the Current Population Survey (CPS), Bureau of the Census, for several reasons. In addition to sampling variability they include three primary conceptual differences, namely: (1) NHS estimates are for persons 17 years of age or over; CPS estimates are for persons 14 years of age or over. (2) NHS uses a 2-week-reference period while CPS uses a 1-week-reference period. (3) NHS is a continuing

survey with separate samples taken weekly, while C is a monthly sample taken for the survey week which i cludes the 12th of the month.

Residence.—The place of residence of a member of the civilian, noninstitutional population is class fied as inside a standard metropolitan statistic area (SMSA) or outside an SMSA, according to far or nonfarm residence.

Standard metropolitan statistical areas.—The definitions and titles of SMSA are established by t U.S. Bureau of the Budget with the advice of t Federal Committee on Standard Metropolitan Statitical Areas. There were 212 SMSA's, as defin for the 1960 Decennial Census, for which data m be provided for places of residence in the Hea Interview Survey.

The definition of an individual SMSA involutive considerations: first, a city or cities of specific population which constitute the central city and identify the county in which it is located as the cutral county; and, second, economic and social alationships with contiguous counties except in N England which are metropolitan in character, that the periphery of the specific metropolitan as may be determined. SMSA's are not limited by St boundaries.

Farm and nonfarm residence.—The populat residing outside SMSA is subdivided into the fa population, which comprises all non-SMSA reside living on farms, and the nonfarm population, wh comprises the remaining non-SMSA population. I farm population includes persons living on plac of 10 or more acres from which sales of farm products amounted to \$50 or more during the previous 12 months or on places of less than 10 acres from which sales of farm products amounted to \$250 more during the preceding 12 months. Other pasons living in non-SMSA territory were classiff as nonfarm if their household paid rent for the household paid rent for the household paid rent for the household paid used for far ing.

Sales of farm products refer to the gross: ceipts from the sale of field crops, vegetables, fru: nuts, livestock and livestock products (milk, we etc.), poultry and poultry products, and nursery forest products produced on the place and sold any time during the preceding 12 months.

Region.—For the purpose of classifying the population by geographic area, the States are grouped into four regions. These regions, which correspond to those used by the Bureau of the Census, are as follows:

Region

States Included

Northeast----- Maine, New Hampshire, Vermont, Massachusetts, Rhode Island, Connecticut, New York, New Jersey, Pennsylvania

North Central --- Michigan, Ohio, Indiana, Illinois, Wisconsin, Minnesota, Iowa, Missouri, North Dakota, South Dakota, Nebraska, Kansas South----- Delaware, Maryland, District of Columbia, Virginia, West Virginia, North Carolina, South Carolina, Georgia, Florida, Kentucky, Tennessee, Alabama, Mississippi, Arkansas, Louisiana, Oklahoma, Texas

West----- Montana, Idaho, Wyoming, Colorado, New Mexico, Arizona, Utah, Nevada, Alaska, Washington, Oregon, California, Hawaii

_____000____

REPORTS FROM THE NATIONAL CENTER FOR HEALTH STATISTICS Public Health Service Publication No. 1000

Series 1. Programs and collection procedures

- No. 1. Origin, Program, and Operation of the U.S. National Health Survey. 35 cents.
- No. 2. Health Survey Procedure: Concepts, Questionnaire Development, and Definitions in the Health Interview Survey. 45 cents.
- No. 3. Development and Maintenance of a National Inventory of Hospitals and Institutions. 25 cents.

Series 2. Data evaluation and methods research

- No. 1. Comparison of Two-Vision Testing Devices. 30 cents.
- No. 2. Measurement of Personal Health Expenditures. 45 cents.
- No. 3. The One-Hour Glucose Tolerance Test. 30 cents.
- No. 4. Comparison of Two Methods of Constructing Abridged Life Tables. 15 cents.

Series 3. Analytical studies

- No. 1. The Change in Mortality Trend in the United States. 35 cents.
- No. 2. Recent Mortality Trends in Chile. 30 cents.

Series 4. Documents and committee reports

No reports to date.

Series 10. Data From the Health Interview Survey

- No. 1. Acute Conditions, Incidence and Associated Disability, United States, July 1961-June 1962. 40 cents.
- No. 2. Family Income in Relation to Selected Health Characteristics, United States. 40 cents.
- No. 3. Length of Convalescence After Surgery, United States, July 1960-June 1961. 35 cents.
- No. 4. Disability Days, United States, July 1961-June 1962. 40 cents.
- No. 5. Current Estimates From the Health Interview Survey, United States, July 1962-June 1963. 35 cents.
- No. 6. Impairments Due to Injury, by Class and Type of Accident, United States, July 1959-June 1961. 25 cents.
- No. 7. Disability Among Persons in the Labor Force, by Employment Status, United States, July 1961-June 1962. 40 cents.
- No. 8. Types of Injuries, Incidence and Associated Disability, United States, July 1957-June 1961. 35 cents.
- No. 9. Medical Care, Health Status, and Family Income, United States. 55 cents.
- No. 10. Acute Conditions, Incidence and Associated Disability, United States, July 1962-June 1963. 45 cents.
- No. 11. Health Insurance Coverage, United States, July 1962-June 1963. 35 cents.
- No. 12. Bed Disability Among the Chronically Limited, United States, July 1957-June 1961. 45 cents.
- No. 13. Current Estimates From the Health Interview Survey, United States, July 1963-June 1964, 40 cents.
- No. 14. Illness, Disability, and Hospitalization Among Veterans, United States, July 1957-June 1961.
- No. 15. Acute Conditions, Incidence and Associated Disability, United States, July 1963-June 1964.
- No. 16. Health Insurance, Type of Insuring Organization and Multiple Coverage, United States, July 1962-June 1963.

Series 11. Data From the Health Examination Survey

- No. 1. Cycle I of the Health Examination Survey: Sample and Response, United States, 1960-1962. 30 cents.
- No. 2. Glucose Tolerance of Adults, United States, 1960-1962. 25 cents.
- No. 3. Binocular Visual Acuity of Adults, United States, 1960-1962. 25 cents.
- No. 4. Blood Pressure of Adults, by Age and Sex, United States, 1960-1962. 35 cents.
- No. 5. Blood Pressure of Adults, by Race and Region, United States, 1960-1962. 25 cents.
- No. 6. Heart Disease in Adults, United States, 1960-1962. 35 cents.
- No. 7. Selected Dental Findings in Adults, United States, 1960-1962. 30 c ats.

Series 12. Data From the Health Records Survey

No reports to date.

Series 20. Data on mortality

No reports to date.

Series 21. Data on natality

No. 1. Natality Statistics Analysis, United States, 1962. 45 cents.

Series 22. Data on marriage and divorce

No reports to date.

Series 23. Data from the program of sample surveys related to vital records

No reports to date.