Utilization of Short-Stay Hospitals Summary of Nonmedical Statistics United States - 1971

Statistics are presented on the utilization of short-stay hospitals based on data collected in the Hospital Discharge Survey from a national sample of hospital records of discharged patients. Discharges, days of care, and average length of stay are distributed by each of the variables age, sex, and color of patient and by geographic region, bed size, and type of ownership (control) of hospital.

DHEW Publication No. (HRA) 75-1768

U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE
Public Health Service

Health Resources Administration
National Center for Health Statistics
Rockville, Md. August 1974



Library of Congress Cataloging in Publication Data

Lewis, W. Frank

Utilization of short-stay hospitals.

(Vital and health statistics. Series 13: Data from the National Health survey, no. 17) (DHEW publication no. (HRA) 75-1768)

Supt. of Docs. no.: HE 20.2210: 13/17.

Bibliography: p.

1. Hospital utilization—United States—Statistics. I. United States. National Center for Health Statistics. II. Title. III. Series: United States. National Center for Health Statistics. Vital and health statistics. Series 13: Data from the National Health Survey. Data from the hospital discharge survey, no. 17. IV. Series: United States. Dept. of Health, Education, and Welfare. Publication no. (HRA) 75-1768. [DNLM: 1. Hospitalization—Statistics. 2. Hospitals, Special—Utilization—U.S. W2A N148vm no. 2 etc.]

RA407.3.A349 no. 17 [RA981.A2] 362.1'1'0973s

ISBN 0-8406-0003-8

[362.1'1'0973]

74-2233

NATIONAL CENTER FOR HEALTH STATISTICS

EDWARD B. PERRIN, Ph.D., Director

PHILIP S. LAWRENCE, Sc.D., Deputy Director
JACOB J. FELDMAN, Ph.D., Acting Associate Director for Analysis
GAIL F. FISHER, Associate Director for the Cooperative Health Statistics System
ELIJAH L. WHITE, Associate Director for Data Systems
IWAO M. MORIYAMA, Ph.D., Associate Director for International Statistics
EDWARD E. MINTY, Associate Director for Management
ROBERT A. ISRAEL, Associate Director for Operations
QUENTIN R. REMEIN, Associate Director for Program Development
PHILIP S. LAWRENCE, Sc.D., Acting Associate Director for Research
ALICE HAYWOOD, Information Officer

DIVISION OF HEALTH RESOURCES UTILIZATION STATISTICS

SIEGFRIED A. HOERMANN, Director WILLIAM F. STEWART, Acting Deputy Director TOY C. TAIRA, Acting Chief, Hospital Care Statistics Branch

COOPERATION OF THE BUREAU OF THE CENSUS

Under the legislation establishing the National Health Survey, the Public Health Service is authorized to use, insofar as possible, the services or facilities of other Federal, State, or private agencies. In accordance with specifications established by the National Center for Health Statistics, the Bureau of the Census, under a contractual arrangement, participated in planning the survey and collecting the data.

Vital and Health Statistics-Series 13-No. 17

DHEW Publication No. (HRA) 75-1768

Library of Congress Catalog Card Number 74-2233

CONTENTS

	Page
Introduction	1
Selected Findings	1
Discharge and Discharge Rates	2
Age and Sex	
Color	2
Bed Size of Hospital	4
Type of Ownership of Hospital	4
Days of Care and Length of Stay	5
Age and Sex	5
Color	5
Bed Size of Hospital	5
Type of Ownership of Hospital	6
Geographic Region	6
Age and Sex	6
Color	7
Conclusion	9
References	12
List of Detailed Tables	13
Appendix I. Technical Notes on Methods	33 33 36 36 37
Appendix II. Definitions of Certain Terms Used in This Report Terms Relating to Hospitalization	40 40 40 40

UTILIZATION OF SHORT-STAY HOSPITALS, SUMMARY OF NONMEDICAL STATISTICS

W. Frank Lewis, Division of Health Resources Utilization Statistics

INTRODUCTION

This report presents estimates on the utilization of non-Federal short-stay hospitals in the United States based on information collected in the Hospital Discharge Survey, a continuous nation-wide survey conducted by the National Center for Health Statistics. Data were abstracted from about 200,000 hospital records of inpatients discharged from 379 hospitals that participated in the survey.

Results of the survey permit generation of four basic types of reports: nonmedical, diagnostic, surgical, and patient charges. Reports are published in Series 13 of the *Vital and Health Statistics* reports and as selected supplements of *Monthly Vital Statistics Reports.*¹⁻¹⁶ Estimates shown in this report are for patients in non-Federal short-stay hospitals, excluding newborn infants, discharged during 1971. Nonmedical data are presented on the number and rate of discharges and of days of care, and average length of stay for patients discharged, by age, sex, and color and by geographic region, bed size, and type of ownership (control) of hospitals.

This nonmedical report will be divided primarily into three areas: an analysis of hospital discharges, an analysis of days of care and length of stay, and some comparisons between the four regions on selected variables. Since the estimates are based on a sample of discharges from participating hospitals rather than on all discharges from all short-stay hospitals, they are subject to sampling error. Tables and graphs of approximate sampling errors and instructions for their use are given in the section "Reliability of Estimates" in appendix I.

Appendix II contains definitions of terms relating to hospitalization and the characteristics of patients and hospitals surveyed. Since several of these terms have specialized meaning in the Hospital Discharge Survey, familiarity with the definitions will aid in interpreting the data.

SELECTED FINDINGS

An estimated 29.5 million inpatients were discharged from non-Federal short-stay hospitals in 1971. These patients received an estimated 231.0 million days of care, with an average length of stay of 7.8 days per hospital episode. In terms of annual rates, the rate of days of care per 1,000 persons in the civilian noninstitutionalized population was 1,143.1, and there was a discharge rate of 145.8 per 1,000 persons. Approximately three-fourths (73.3 percent) of the discharges in 1971 were from voluntary nonprofit hospitals. State and local government hospitals accounted for 22.5 percent of the discharges and proprietary hospitals for only 4.2 percent.

Patients under 15 years of age accounted for 13.7 percent of all discharges and had a rate of discharge per 1,000 population of 70.2. This contrasted with the population 65 years of age and over, whose rate of discharge was 305.7 per 1,000 population. Differences in hospital utilization by sex were also noted. Rates of discharge and of days of care were higher for females than for males. The discharge rate for females of 169.6 per 1,000 population was 42 percent higher than that for males, 119.6 per 1,000 population, but with hospitalization for deliveries excluded the discharge rate for females was only 15 per-

cent higher than the rate for males. For the group 1-14 years of age, however, the discharge rate for males was higher than for females, 76.9 compared to 63.0 per 1,000 population.

Hospitalization utilization figures by color are grouped in the categories "white," "all other," and "color not stated." Since the number of discharged patients for whom color was not stated is slightly larger than the all other group, data analysis by color must be interpreted with caution. Based on the estimates of patients discharged for whom color was stated, those identified as white outnumbered the all other group by about 7 to 1. As a group white patients were older than all other patients but each age-sex group had shorter average lengths of stay than did all other patients.

The age distribution within hospitals varied by the size of the hospital. The smallest hospitals had proportionately fewer patients 15-64 years of age than did the largest hospitals, in which only 15.9 percent of the patients were age 65 years and over. Average length of stay increased with hospital size from 6.5 days in the smallest hospitals to 9.1 days in hospitals with 500 beds or more.

Regional differences were apparent in number of discharges, ranging from 4.2 million in the West Region to 9.2 million in the North Central Region. Average length of stay was longest in the Northeast Region, 9.0 days, and lowest in the West, where length of stay averaged only 6.5 days.

DISCHARGES AND DISCHARGE RATES

Age and Sex

Patients under 15 years of age accounted for an estimated 4.0 million or 13.7 percent of all discharged patients from short-stay hospitals in 1971. Of these, 2.4 percent were less than 1 year old, 3.8 percent were 1-4 years old, and 7.5 percent were from 5-14 years of age (figure 1). The discharge rate for the group under 15 years of age was the lowest for any age group with a rate of 70.2 per 1,000 population (table A). In contrast, the discharge rate for persons 65 years and older was 305.7 discharges per 1,000 population.

Males 65 years and over were discharged at a rate of 328.9 per 1,000 compared to the lower




Figure 1. Percent distribution of patients discharged from shortstay hospitals by age, according to sex: United States, 1971.

rate for females of 288.2 per 1,000. For all age groups, however, discharge rates for females excluding deliveries was higher than for males, 137.7 versus 119.6 per 1,000. There were more male than female discharges in each age bracket under 15 years of age. At under 1 year of age males outnumbered females 409,000 to 294,000; ages 1-4 years, 644,000 to 488,000; and ages 5-14 years, 1,196,000 to 992,000 (table B). Within these young age groups there was a higher percent of the total male population than of the female population (figure 1) with 19.3 percent of the male discharges in the age group under 15 years compared to 10.0 percent for females.

As shown in table A, the effect of deliveries can be seen in the trend of discharge rates by age. The rates for males and for females excluding deliveries increased consistently with increasing age.

Color

Data for patients discharged by color are shown in table 1 according to the three categories "white," "all other," and "color not stated." An estimated 22.5 million white patients and 3.3 million all other patients for whom color was reported were discharged from short-stay hos-

Table A. Number and rate of discharges and of days of care and average length of stay for patients discharged from short-stay hospitals, by age and sex: United States, 1971

			Female			
Age	Both sexes ¹	Male	Including deliveries	Excluding deliveries		
	Number	of discha	rges in thousands			
All ages	29,459	11,644	17,767	14,431		
Under 15 years	4,029 12,605 6,840 5,986	3,467 3,232	9,118 3,596	3,590		
	Rate of di	scharges	per 1,000 po	pulation		
All ages	145.8	119.6	169.6	137.7		
Under 15 years	70.2 151.4 163.3 305.7	86.7 162.3		62.5 134.1 163.4 288.2		
	Number o	f days of	care in tho	usands		
All ages	231,017	97,723	132,906	119,374		
Under 15 years	18,773 72,359 64,304 75,581	23,866 30,595	33,606	8,130 34,887 33,580 42,778		
	Rate of day	s of care	per 1,000 p	opulation		
All ages	1,143.1	1,004.0	1,268.7	1,139.5		
Under 15 years	1,535.0	596.8	1,529.0	288.9 806.5 1,527.8 3,759.0		
	Averag	e length	of stay in d	ays		
All ages	7.8	8.4	7.5	8.3		
Under 15 years	4.7 5.7 9.4 12.6	4.7 6.9 9.5 12.1	4.6 5.3 9.3 13.0	4.6 6.0 9.4 13.0		

 $^{^{1} \, \}mathrm{Includes}$ discharge data for which sex was not stated.

Table B. Number of patients under 15 years of age discharged from short-stay hospitals, by age and sex: United States, 1971

Age	Both sexes ¹	Male	Female includ- ing de- liveries		
Under 15 years	Number in thousands				
Under 1 year 1-4 years 5-14 years	705 1,133 2,192	409 644 1,196	294 488 992		

¹Includes discharge data for which sex was not stated.

pitals in 1971, white patients outnumbering all other patients by about 7 to 1. Color was not stated in the medical record summary sheets for about 3.7 million patients, a number greater than that for patients identified as all other. The distribution for those for whom color was not stated suggests that they were proportional by color to those for whom it was stated.

There were more white patients than all other patients 45 years and over, 46.0 percent and 29.2 percent, respectively. Approximately 1 in 5 of the estimated white patients were age 65 and over as compared to 1 in 8 among patients of races other than white (figure 2).

White patients included 40 percent males and 60 percent females compared with 36 percent males and 64 percent females in the all other category. A smaller percentage of white females hospitalized for deliveries than all other patients accounted for most of this difference.

Bed Size of Hospital

For all hospital sizes the percent of patients discharged during 1971 under 15 years of age was approximately the same, between 13 and 15 percent; however, they differed appreciably in the percent distributions for the three age groups 15 years and older (table 3). The smallest hospitals




Figure 2. Percent of discharges from short-stay hospitals, by color and age: United States, 1971.

had proportionately fewer discharged patients 15-64 years old than did the largest hospitals. On the other hand, for the age group 65 years and over, the smallest hospitals had more patients discharged (26.6 percent) than the largest hospitals (15.9 percent). In other words, the smaller the hospital the more likely it was that a larger percent of its discharged patients would be found in the oldest age group, and the larger the hospital the larger the relative percent of patients to be found in the age group 15-64 years.

This age and hospital size trend was true for males and, with the exception of the age group 45-64 years, for females. The percents of female discharges 45-64 years old were essentially the same (19.2, 19.7, 19.9, 21.6, 20.7) for each size hospital. When deliveries were excluded, the trend of increased percent of patients with increased size of hospital was still not very apparent for this age group (23.0, 24.2, 24.8, 26.4, 26.0).

Type of Ownership of Hospital

Voluntary nonprofit hospitals (operated by church or other nonprofit organizations) cared for 21.6 million patients in 1971, or 73 percent

of all patients discharged from short-stay hospitals (table 4). Government hospitals (controlled by State or local government) accounted for 6.6 million discharges, or 23 percent of all patients, and proprietary hospitals discharged 1.2 million, or 4 percent. There were relatively few differences in the percent distributions by age and sex. However, the relative proportion of male to female discharges varied with the type of hospital, government hospitals recording female discharges 49 percent greater than for males (3,961,000 and 2,660,000, respectively), and voluntary and proprietary hospitals showing approximately a 58percent difference. When deliveries are excluded, the number of discharges for females is around 24 percent greater than for males for all types of hospitals. There was a noticeable difference between voluntary and government hospitals in the distribution of their patients aged 15-44 years, with government hospitals having 45.6 percent and voluntary hospitals only 41.9 percent of their patients in this age group.

DAYS OF CARE AND LENGTH OF STAY

Age and Sex

The rates of days of care by age groups ranged from 234.0 days per 1,000 population aged 5-14 years to 5,119.8 days for persons 75 years and over (table 6). Starting with age group 5-14 years, the days of care rate increased with each advance in age. Age groups under 15 years represented 8.2 percent of all days of care; ages 15-44, 31.4 percent; ages 45-64, 27.9 percent; and ages 65 and over, 32.7 percent of all days of care.

The average length of stay for patients discharged during 1971 was 7.8 days. Average length of stay increased with each successive age group from 4.7 days for patients under age 15 to 12.6 days for patients aged 65 years and over. For all age groups, about two-thirds were discharged within a week (table 7).

The days of care rates per 1,000 population were lower for females than for males under 15 and over 54 years of age (table 6). Deliveries exerted less influence on the days of care rate

than on the rate of discharges for females because of the relatively short average length of stay. The average length of stay for females aged 15-44 years is lowered by approximately two-thirds of a day when deliveries are included. Generally speaking, females under 65 years of age had shorter lengths of stay than males, and females over 64 years had longer lengths of stay (table 9).

Color

Differences between the age and sex distributions of days of care utilized by white and all other patients for whom color was stated are found in table 8. Patients under age 15 years in the all other group used a larger proportion (13.1 percent) of days of care than did white patients in this age group, who used 7.4 percent. Among white patients, approximately 36 percent of the days of care were provided for patients under 45 years of age; among all others about 55 percent were provided for this age group. For ages 65 years and over white patients—both men and women—used a considerably larger percent of days of care than did all other patients, 34.9 percent and 21.1 percent, respectively.

There was little difference in the average length of stay for all discharges by color, with white patients averaging 7.9 days and all other patients 8.1 days per stay (table 9). Average length of stay for the two groups was about the same because the white discharged patients included a larger percent of older patients with longer hospital stays than did the all other group. For every age and sex group, however, the average stay was significantly shorter for white patients than for all others. Regardless of color status, males had a longer length of stay than did females including deliveries. When deliveries are excluded, all other males average 1 day longer than all other females, primarily because of the 8.4 average length of stay for age group 15-44 years. White males and those with color not stated had approximately the same lengths of stay as did females excluding deliveries.

Bed Size of Hospital

Days of care for hospital size also varied by age. For patients aged 65 years and over, reported days of care ranged from 25.0 percent

in the largest hospitals to 43.8 percent in the smallest hospitals (table 12). Days of care for patients of ages 15-44 was 25.7 percent in hospitals with fewer than 100 beds and 36.3 percent in those with 500 beds or more.

Average length of stay increased as age and hospital size increased, ranging from 6.5 days in the smallest hospitals to 9.1 days in the largest hospitals and from 4.7 days for the youngest group to 12.6 days for the oldest group. This was true for both sexes (table 13). The shortest length of stay, 3.8 days, was for the age group under 15 discharged from hospitals with 6-99 beds. The longest length of stay was 14.3 days for the group 65 years and over discharged from hospitals with 500 beds or more.

The pattern of length of stay increasing by size of hospital was true for each of the four regions. The trend was most evident in the Northeast and West Regions, where average length of stay in the largest hospitals exceeded that in the smallest hospitals by 3.5 days (table 16). This trend was more pronounced for males than for females in each of the four regions, regardless of delivery status. For the male episodes in the 15-44 age group, the average length of stay in the largest hospitals was 74 percent longer or more than that in the smallest hospitals in each of the regions.

Type of Ownership of Hospital

The 231.0 million days of care utilized in 1971 were distributed by ownership of hospital as follows: voluntary nonprofit hospitals provided 173.5 million days, or 75.1 percent; government hospitals provided 49.0 million days, or 21.2 percent; and proprietary hospitals provided 8.5 million days, or 3.7 percent (table 14).

The relative proportion of males to females varied considerably among the three types of hospitals for days of care provided. In government hospitals days of care provided for females including deliveries was 27 percent greater than the days for males, for voluntary nonprofit hospitals days of care provided for females was 38 percent greater than for males, and for proprietary hospitals days of care for females was 50 percent greater than for males.

Average length of stay was consistently shorter in proprietary hospitals than involuntary nonprofit hospitals for both sexes and all age groups. Average length of stay in government hospitals was also shorter than in voluntary nonprofit hospitals for both sexes and all age groups except under 15 years, where average length of stay was longer in the government hospitals (table 15). For all age groups under 65 years, the average length of stay in proprietary hospitals was shorter than in the other hospital ownership groups for both sex groups and all age groups. The difference between length of stay for male and female including deliveries was also smallest for proprietary hospitals, about half a day compared to approximately 1 day for voluntary nonprofit and government hospitals. The average length of stay is about the same for both sexes when deliveries are excluded. For the group 15-44 years old, the average length of stay for males varied between half a day and 2 days longer than that for females, regardless of delivery status, for each type of hospital.

GEOGRAPHIC REGION

Age and Sex

The number of discharges in 1971 by geographic region ranged from 4.2 million in the West Region to 9.2 million in the North Central Region (table 5). The number of discharges per 1,000 population ranged from an estimated 122.6 in the West Region to 162.9 in the North Central Region; among the age groups the greatest relative difference is found in the group less than 15 years, 85.1 in the North Central and 54.4 in the West per 1,000 population (table C).

The number of days of care per 1,000 population followed a similar pattern, again being lowest in the West Region and highest in the North Central Region. The rates were 790.9 days and 1,304.4 days, respectively, a difference of 65.0 percent. For patients under age 15 years, these two regions differed even more significantly, with the days of care per 1,000 in the North Central Region being 99 percent higher than those in the West Region (393.2 and 197.4 days of care).

Average length of stay in days was highest in the Northeast Region and again lowest in the West

Table C. Rate of discharges and of days of care and average length of stay for patients discharged from short-stay hospitals, by age and geographic region: United States, 1971

Age	All regions	Northeast	North Central	South	West
	Rate	of discharg	ge per 1,0	000 popula	ation
All ages	145.8	141.6	162.9	146.5	122.6
Under 15 years	70.2 151.4 163.3 305.7	152.5 153.3	166.3 186.4	150.7 162.3	127.7 142.6
	Rate of	days of ca	re per 1,	000 popul	ation
All ages	1,143.1	1,275.8	1,304.4	1,089.0	790.9
Under 15 years	327.2 869.2 1,535.0 3,860.3	934.5 1,681.3	393.2 993.7 1,789.9 4,278.3	857.3 1,417.1	606.4 1,102.7
	A	werage leng	th of sta	y in days	3
All ages	7.8	9.0	8.0	7.4	6.5
Under 15 years	4.7 5.7 9.4 12.6	5.2 6.1 11.0 15.2		5.7	4.7

Region, being 9.0 days and 6.5 days, respectively. This pattern was consistent for all age groups, with the difference between the Northeast and the West Regions being again greatest in the group under 15 years, 5.2 and 3.6 days, respectively, a difference of 44.4 percent.

The average length of stay showed the same age and regional trends, with length of stay for both sexes being longest in the Northeast Region for the age group over 65 years, and shortest in the West Region for the group less than 15 years. Average length of stay for males was slightly longer than for females in all regions. Females 65 years and over in each of the four regions had longer stays than did males in this age group (table 11).

Color

When color is considered, differences were found among the regions in the rate of discharges.

The Northeast Region had the highest proportion of white discharges (82.7 percent) and the North Central Region the lowest (71.0 percent) (figure 3). The South had the smallest proportion of its patients in the not stated category, 6.4 percent, in contrast to the North Central Region, which had 21.2 percent listed as not stated.

Deliveries represented a smaller proportion of white patients than of all other patients hospitalized, 10.5 percent and 17.4 percent, respectively (figure 4). With the exception of the West Region, deliveries represented a smaller proportion of the total discharges for white patients than for all others.

Average length of stay by color was also not consistent among regions. In the Northeast, South, and West Regions white patients had shorter stays than all others (figure 5), whereas the North Central Region showed no apparent color variation. Excluding deliveries, about half the white




Figure 3. Percent of patients discharged from short-stay hospitals, by geographic region and color: United States, 1971.

patients hospitalized were 45 years of age and over in every region (figure 6), whereas only 35 percent of all other patients were 45 years or older.

For the population 15 years and over, regional differences were found in the all other groups. Over 70 percent of the Northeast and North Central discharges were in the 15-44 age group contrasted




Figure 4. Percent of deliveries of total discharges from shortstay hospitals, by geographic region and color: United States, 1971.




Figure 5. Average length of stay for patients discharged from short-stay hospitals, by geographic region and color: United States, 1971.

to the South and West, which had only 62.2 and 56.5 percent, respectively, of their discharges in this age group (table D).

The average length of stay for all others varied considerably for the group 65 years and over among regions, with 11.6 days of care in the West and 21.7 in the Northeast (table E).




Figure 6. Percent of patients aged 45 years and over of total inpatients discharged from short-stay hospitals, excluding deliveries, by geographic region and color: United States, 1971.

Table D. Number and percent distribution of patients 15 years of age and over discharged from short-stay hospitals by geographic region and age, according to color: United States, 1971

Geographic region and age	Total	White	All other	Color not stated	Total	White	All other	Color not stated
United States	Nu	mber in t	housand	ls	Per	cent dis	tributi	.on
15 years and 'over-	25,431	19,477	2,787	3,166	100.0	100.0	100.0	100.0
15-44 years	12,605 6,840 5,986	9,159 5,430 4,888	1,812 570 405	1,633 840 693	50.0 26.9 23.5	47.0 27.9 25.1	65.0 20.5 14.5	51.6 26.5 21.9
Northeast 15 years and over	6,061	5,020	562	479	100.0	100.0	100.0	100.0
15-44 years 45-64 years65 years and over	3,006 1,665 1,390	2,362 1,429 1,229	397 108 57	247 128 104	49.6 27.5 22.9	47.1 28.5 24.5	70.7 19.2 10.1	51.5 26.7 21.7
North Central								
15 years and over	7,785	5,556	583	1,648	100.0	100.0	100.0	100.0
15-44 years	3,838 2,132 1,815	2,593 1,583 1,380	415 110 58	830 440 378	49.3 27.4 23.3	46.7 28.5 24.8	71.2 18.9 9.9	50.4 26.7 22.9
South								
15 years and over	7,882	6,098	1,262	521	100.0	100.0	100.0	100.0
15-44 years	3,887 2,046 1,949	2,829 1,673 1,596	785 251 226	273 122 126	49.3 26.0 24.7	46.4 27.4 26.2	62.2 19.9 17.9	52.4 23.3 24.2
West								
15 years and over	3,702	2,802	381	520	100.0	100.0	100.0	100.0
15-44 years	1,873 996 833	1,375 744 683	215 101 65	284 151 85	50.6 26.9 22.5	49.1 26.6 24.4	56.5 26.5 17.0	54.6 29.0 16.4

CONCLUSION

Analysis of estimates on the utilization of short-stay hospitals in the United States for 1971 in terms of age, sex, race, hospital size and ownership, and regions based on the Hospital Discharge Survey can be summarized as follows: 1. Patients under 15 years of age had the lowest rates of discharges and days of care per 1,000 population and the shortest average length of stay of any age group. Rates of discharges and days of care and average length of stay increased with age.

Table E. Average length of stay for patients 15 years of age and over discharged from short-stay hospitals, by geographic region, age, and color: United States, 1971

Geographic region and age	Total	White	All other	Color not stated
United States	Aver	age length	of stay in	days
15 years and over	8.3	8.4	8.5	8.0
15-44 years	5.7 9.4 12.6	5.6 9.3 12.6	6.3 11.5 14.1	5.9 8.9 12.1
Northeast	:			
15 years and over	9.5	9.6	10.2	8.6
15-44 years	6.1 11.0 15.2	6.0 10.8 15.0	7.3 14.7 21.7	5.8 9.5 14.4
North Central				
15 years and over	8.6	8.7	8.5	8.3
15-44 years	6.0 9.6 13.0	5.9 9.5 13.1	6.5 12.6 15.0	5.9 9.2 12.6
South			:	<u> </u>
15 years and over	7.9	7.8	8.1	8.0
15-44 years	5.7 8.7 11.3	5.5 8.5 11.2	5.9 10.7 12.8	6.8 8.3 10.1
West				
15 years and over	6.9	6.8	7.3	6.6
15-44 years	4.7 7.7 10.6	4.6 7.6 10.6	5.4 8.6 11.6	5.0 7.7 9.9

^{2.} Female patients overall had higher discharge and days of care rates than did male, but male rates were higher than those of females for age groups less than 15 and over 64 years old. Average length of stay was longer for female patients aged 65 years and over.

^{3.} White patients were generally older than patients in the all other category, had a greater proportion of female patients (but with proportionally fewer deliveries), and had a shorter average length of stay for each age and sex group.

- 4. Small hospitals tended to have a greater proportion of their patients in the oldest age group, and large hospitals had proportionately more of their patients in the 15-64 age group. Average length of stay was shortest in the smallest hospitals and increased steadily with hospital size.
- 5. Voluntary hospitals cared for almost three-quarters of all patients discharged and reported female discharges 53 percent greater than for male. Voluntary nonprofit hospitals had the longest average length of stay and proprietary hospitals the shortest.
- 6. The North Central Region had the highest rate of discharges and days of care per

- 1,000 population, and the West Region had the lowest rates. Average length of stay was also lowest in the West Region.
- 7. In general, patients under 15 years of age admitted to hospitals with less than 100 beds in the West Region had the shortest hospital episodes, whereas female patients 65 years old and over admitted to hospitals with 500 beds or more located in the Northeast Region had the longest average length of stay.

A more complete analysis of the interrelationships among these variables and their effects on reported estimates is not possible due to the sampling errors inherent in the statistical design.

REFERENCES

¹National Center for Health Statistics: Utilization of short-stay hospitals, summary of nonmedical statistics, United States, 1965. *Vital and Health Statistics*. PHS Pub. No. 1000-Series 13-No. 2. Public Health Service. Washington. U.S. Government Printing Office, Aug. 1967.

²National Center for Health Statistics: Utilization of short-stay hospitals by characteristics of discharged patients, United States, 1965. *Vital and Health Statistics*. PHS Pub. No. 1000-Series 13-No. 3. Public Health Service. Washington. U.S. Government Printing Office, Dec. 1967.

³National Center for Health Statistics: Patients discharged from short-stay hospitals by size and type of ownership, United States, 1965. *Vital and Health Statistics*. PHS Pub. No. 1000-Series 13-No. 4. Public Health Service. Washington. U.S. Government Printing Office, Dec. 1968.

⁴National Center for Health Statistics: Regional utilization of short-stay hospitals, United States, 1965. *Vital and Health Statistics*. PHS Pub. No. 1000-Series 13-No. 5. Public Health Service. Washington. U.S. Government Printing Office, June 1969.

⁵National Center for Health Statistics: Inpatient utilization of short-stay hospitals by diagnosis, United States, 1965. Vital and Health Statistics. PHS Pub. No. 1000-Series 13-No. 6. Public Health Service. Washington. U.S. Government Printing Office, May 1970.

⁶National Center for Health Statistics: Surgical operations in short-stay hospitals for discharged patients, United States, 1965. *Vital and Health Statistics*. PHS Pub. No. 1000-Series 13-No. 7. Public Health Service. Washington. U.S. Government Printing Office, Apr. 1971.

⁷National Center for Health Statistics: Utilization of short-stay hospitals, summary of nonmedical statistics, United States, 1966. *Vital and Health Statistics*. Series 13-No. 8. DHEW Pub. No. (HSM) 72-1006. Health Services and Mental Health Administration. Washington. U.S. Government Printing Office, Sept. 1971.

⁸ National Center for Health Statistics: Utilization of short-stay hospitals, summary of nonmedical statistics, United States, 1967. *Vital and Health Statistics*. Series 13-No. 9. DHEW Pub. No. (HSM) 72-1058. Health Services and Mental Health Administration. Washington. U.S. Government Printing Office, May 1972.

⁹National Center for Health Statistics: Inpatient utilization of short-stay hospitals in each geographic division, United States, *Vital and Health Statistics*. Series 13-No. 10. DHEW Pub. No. (HSM) 73-1761. Health Services and Mental Health Administration. Washington. U.S. Government Printing Office, Nov. 1972.

¹⁰National Center for Health Statistics: Surgical operations in short-stay hospitals, United States, 1968. *Vital and Health Statistics*. Series 13-No. 11. DHEW Pub. No. (HSM) 73-1762. Health Services and Mental Health Administration. Washington. U.S. Government Printing Office, Jan. 1973.

¹¹National Center for Health Statistics: Inpatient utilization of short-stay hospitals by diagnosis, United States, 1968. Vital and Health Statistics. Series 13-No. 12. DHEW Pub. No. (HSM) 73-1763. Health Services and Mental Health Administration. Washington. U.S. Government Printing Office, Mar. 1973.

¹²National Center for Health Statistics: Average length of stay in short-stay hospitals: demographic factors, United States, 1968. Vitàl and Health Statistics. Series 13-No. 13. DHEW Pub. No. (HSM) 73-1764. Health Services and Mental Health Administration. Washington. U.S. Government Printing Office, Apr. 1973.

¹³National Center for Health Statistics: Utilization of short-stay hospitals, summary of nonmedical statistics: United States, 1969, *Monthly Vital Statistics Reports.* Vol. 21, No. 6, Supplement (HSM) 73-1126. Rockville, Md. Health Services and Mental Health Administration, Sept. 19, 1972.

¹⁴National Center for Health Statistics: Utilization of short-stay hospitals, summary of nonmedical statistics, United States, 1973. *Vital and Health Statistics*. Series 13, No. 14. DHEW Pub. No. (HRA) 74-1765. Rockville, Md. Health Resources Administration, Aug. 1973.

¹⁵National Center for Health Statistics: Utilization of short-stay hospitals, summary of nonmedical statistics: United States, 1971, Monthly Vital Statistics Reports. Vol. 22, No. 4, Supplement (HSM) 73-1134. Rockville, Md. Health Services and Mental Health Administration, July 5, 1973.

¹⁶National Center for Health Statistics: Development and maintenance of a national inventory of hospitals and institutions. *Vital and Health Statistics*. PHS Pub. No. 1000-Series 1-No. 3. Public Health Services. Washington. U.S. Government Printing Office, Feb. 1965.

LIST OF DETAILED TABLES

			Page
Tab1e	1.	Number and percent distribution of patients discharged from short-stay hospitals by color and age of patient, according to sex: United States, 1971	14
	2.	Number and percent distribution of patients discharged from short-stay hospitals by geographic region and age, according to sex: United States, 1971	15
	3.	Number and percent distribution of patients discharged from short-stay hospitals by bed size of hospital and age of patient, according to sex: United States, 1971-	16
	4.	Number and percent distribution of patients discharged from short-stay hospitals by type of ownership of hospital and age of patient, according to sex: United States, 1971	17
	5.	Number of patients discharged from short-stay hospitals and days of care, by sex, age, geographic region, and bed size of hospital: United States, 1971	18
	6.	Number, percent distribution, and rate of days of care, average number of hospital beds occupied daily, and average length of stay for patients discharged from short-stay hospitals, by sex and age: United States, 1971	21
	7.	Number and percent distribution of patients discharged from short-stay hospitals by age and length of stay, according to sex: United States, 1971	22
	8.	Number and percent distribution of days of care for patients discharged from short-stay hospitals by color and age of patient, according to sex: United States, 1971	24
	9.	Average length of stay for patients discharged from short-stay hospitals by color, age, and sex: United States, 1971	25
	10.	Number and percent distribution of days of care for patients discharged from short-stay hospitals by geographic region and age, according to sex: United States, 1971	26
	11.	Average length of stay for patients discharged from short-stay hospitals by geographic region, age, and sex: United States, 1971	27
	12.	Number and percent distribution of days of care for patients discharged from short-stay hospitals by bed size of hospital and age of patient, according to sex: United States, 1971	28
	13.	Average length of stay for patients discharged from short-stay hospitals by bed size of hospital, age of patient, and sex: United States, 1971	29
	14.	Number and percent distribution of days of care for patients discharged from short-stay hospitals by type of ownership of hospital and age of patient, according to sex: United States, 1971	30
	15.	Average length of stay for patients discharged from short-stay hospitals by type of ownership of hospital, age of patient, and sex: United States, 1971	31
	16.	Average length of stay for patients discharged from short-stay hospitals by sex,	32

TABLE 1. NUMBER AND PERCENT DISTRIBUTION OF PATIENTS DISCHARGED FROM SHORT-STAY HOSPITALS BY COLOR AND AGE OF PATIENT. ACCORDING TO SEX: UNITED STATES, 1971

COLOR·AND AGE	1/ BOTH SEXES	MALE	FEMALE INCLUD- ING DELIV- ERIES	FEMALE EXCLUD- ING DELIV- ERIES	1/ BOTH SEXES	MALE	FEMALE INCLUD- ING DELIV- ERIES	FEMALE EXCLUD- ING DELIV- ERIES
TOTAL	вмии		ENTS DISCH USANDS	IARGED	!	PERCENT DI	STPIBUTION	
ALL AGES	25,459	11,644	17.767	14,431	100.0	100.0	100.3	100.0
UNDER 15 YEARS 15-44 YEARS 45-64 YEARS 65 YEARS AND OVER	4,029 12,605 6,840 5,986	2,249 3,467 3,232 2,696	1,773 9,118 3,596 3,280	1,759 5,801 3,590 3,280	13.7 42.8 23.2 20.3	19.3 29.8 27.8 23.2	10.0 51.3 20.2 18.5	12.2 40.2 24.9 22.7
ALL AGES	22,451	9,011	13,435	11,086	100.0	100.0	100.0	100.0
UNDER 15 YEARS 15-44 YEARS 45-64 YEARS 65 YEARS AND OVER	2,974 9,159 5,430 4,888	1,652 2,585 2,568 2,206	1,322 6,571 2,861 2,680	1,317 4,231 2,858 2,680	13.2 40.8 24.2 21.8	18.3 28.7 28.5 24.5	9.8 48.9 21.3 20.0	11.9 38.2 25.8 24.2
ALL OTHER								
ALL AGES	3,338	1,198	2,138	1,558	100.0	100.0	100.0	100.0
UNDER 15 YEARS 15-44 YEARS 45-64 YEARS 65 YEARS AND OVER	551 1,812 570 405	321 422 266 189	230 1•389 303 215	222 819 302 215	16.5 54.3 17.1 12.1	26.8 35.3 22.2 15.8	10.8 65.0 14.2 10.1	14.2 52.6 19.4 13.8
COLOR NOT STATED								
ALL AGES	3,670	1,435	2,195	1,786	100.0	100.0	100.0	100.0
UNDER 15 YEARS 15-44 YEARS 45-64 YEARS 65 YEARS AND OVER	504 1,633 840 693	276 460 397 301	221 1,157 432 384	221 751 431 384	13.7 44.5 22.9 18.9	19.2 32.1 27.7 21.0	10.1 52.7 19.7 17.5	12.4 42.0 24.1 21.5

^{1/} INCLUDES DISCHARGE DATA FOR WHICH SEX WAS NOT STATED.

TABLE 2. NUMBER AND PERCENT DISTRIBUTION OF PATIENTS DISCHARGED FROM SHORT-STAY HOSPITALS BY GEOGRAPHIC REGION AND AGE, ACCORDING TO SEX: UNITED STATES, 1971

REGION AND AGE	1/ BOTH SEXES	MALE	FEMALE INCLUD- ING DELIV- ERIES	FEMALE EXCLUD- ING DELIV- ERIES	1/ BOTH SEXES	MALE	FEMALE INCLUD- ING DELIV- ERIES	FEMALE EXCLUD- ING DELIV- ERIES
UNITED STATES	NUMB		ENTS DISCH	ARGED	PERCENT DISTRIBUTION			
ALL AGES	29,459	11,644	17,767	14,431	100.0	100.0	100.0	100.0
UNDER 15 YEARS 15-44 YEARS 45-64 YEARS 65 YEARS AND OVER	4,029 12,605 6,840 5,986	2,249 3,467 3,232 2,696	1,773 9,118 3,596 3,280	1,759 5,801 3,590 3,280	13.7 42.8 23.2 20.3	19.3 29.8 27.8 23.2	10.0 51.3 20.2 18.5	12.2 40.2 24.9 22.7
NORTHEAST ALL AGES	6,912	2,716	4,180	3,378	100.0	100.0	100.0	100.0
UNDER 15 YEARS 15-44 YEARS 45-64 YEARS 65 YEARS AND OVER	851 3,006 1,665 1,390	483 788 810 635	365 2•212 852 752	363 1,413 851 752	12.3 43.5 24.1 20.1	17.8 29.0 29.8 23.4	8.7 52.9 20.4 18.0	10.7 41.8 25.2 22.3
NORTH CENTRAL					:			
ALL AGES	9,171	3,611	5,545	4,508	100.0	100.0	100.0	100.0
UNDER 15 YEARS 15-44 YEARS 45-64 YEARS 65 YEARS AND OVER	1,385 3,838 2,133 1,815	772 1,050 986 803	609 2,783 1,143 1,009	605 1,752 1,141 1,009	15.1 41.9 23.3 19.8	21.4 29.1 27.3 22.2	11.0 50.2 20.6 18.2	13.4 38.9 25.3 22.4
SOUTH								
ALL AGES	9,136	3,629	5•495	4,501	100.0	100.0	100.0	100.0
UNDER 15 YEARS 15-44 YEARS 45-64 YEARS 65 YEARS AND OVER	1,255 3,887 2,046 1,949	698 1,096 962 873	556 2,786 1,081 1,072	550 1,800 1,079 1,072	13.7 42.5 22.4 21.3	19.2 30.2 26.5 24.1	10.1 50.7 19.7 19.5	12.2 40.0 24.0 23.8
WEST				ŀ				
ALL AGES	4,241	1,689	2,547	2,043	100.0	100.0	100.0	100.0
UNDER 15 YEARS 15-44 YEARS 45-64 YEARS 65 YEARS AND OVER	538 1,873 996 833	295 533 474 385	243 1,338 519 447	242 835 519 447	12.7 44.2 23.5 19.6	17.5 31.6 28.1 22.8	9.5 52.5 20.4 17.6	11.8 40.9 25.4 21.9

^{1/} INCLUDES DISCHARGE DATA FOR WHICH SEX WAS NOT STATED.

TABLE 3. NUMBER AND PERCENT DISTRIBUTION OF PATIENTS DISCHARGED FROM SHORT-STAY HOSPITALS BY BED SIZE OF HOSPITAL AND AGE OF PATIENT, ACCORDING TO SEX: UNITED STATES, 1971

BED SIZE OF HOSPITAL AND AGE	1/ BOTH SEXES	MALE	FEMALE INCLUD- ING DELIV- ERIES	FEMALE EXCLUD- ING DELIV- ERIES	1/ BOTH SEXES	MALE	FEMALE INCLUD- ING DELIV- ERIES	FEMALE EXCLUD- ING DELIV- ERIES
ALL SIZES	NUMB		ENTS DISCH	IARGED	PERCENT DISTRIBUTION			
ALL AGES	29,459	11,644	17,767	14,431	100.0	100.0	100.0	100.0
UNDER 15 YEARS 15-44 YEARS 45-64 YEARS 65 YEARS AND OVER	4,029 12,605 6,840 5,986	2,249 3,467 3,232 2,696	1,773 9,118 3,596 3,280	1,759 5,801 3,590 3,280	13.7 42.8 23.2 20.3	19.3 29.8 27.8 23.2	10.0 51.3 20.2 18.5	12.2 40.2 24.9 22.7
6-99 BEDS								
ALL AGES	5,829	2,310	3,510	2,921	100.0	100.0	100.0	100.0
UNDER 15 YEARS 15-44 YEARS 45-64 YEARS 65 YEARS AND OVER	781 2•253 1•244 1•552	438 623 569 680	341 1,627 672 870	339 1,041 672 870	13.4 38.6 21.3 26.6	19.0 27.0 24.6 29.4	9.7 46.4 19.2 24.8	11.6 35.6 23.0 29.8
100-199 BEDS								
ALL AGES	6,054	2,354	3,692	3,004	100.0	100.0	100.0	100.0
UNDER 15 YEARS 15-44 YEARS 45-64 YEARS 65 YEARS AND OVER	852 2,519 1,371 1,312	462 657 641 593	389 1•859 727 717	386 1,175 726 717	14.1 41.6 22.6 21.7	19.6 27.9 27.2 25.2	10.5 50.3 19.7 19.4	12.9 39.1 24.2 23.9
200-299 BEDS								
ALL AGES	5,186	2,039	3,137	2,517	100.0	100.0	100.0	100.0
UNDER 15 YEARS 15-44 YEARS 45-64 YEARS 65 YEARS AND OVER	772 2,244 1,197 973	425 603 570 441	346 1,637 624 530	344 1,019 624 530	14.9 43.3 23.1 18.8	20.9 29.6 28.0 21.6	11.0 52.2 19.9 16.9	13.7 40.5 24.8 21.1
300-499 BEDS							"	
ALL AGES	6,877	2,762	4,100	3,347	100.0	100.0	100.0	100.0
UNDER 15 YEARS 15-44 YEARS 45-64 YEARS 65 YEARS AND OVER	918 2,985 1,701 1,272	522 857 814 570	394 2•121 884 700	391 1,373 882 700	13.4 43.4 24.7 18.5	18.9 31.0 29.5 20.6	9.6 51.7 21.6 17.1	11.7 41.0 26.4 20.9
500 BEDS OR MORE								
ALL AGES	5,513	2,178	3,328	2,642	100.0	100.0	100.0	100.0
UNDER 15 YEARS 15-44 YEARS 45-64 YEARS 65 YEARS AND OVER	705 2,604 1,326 877	400 728 637 413	304 1,873 688 463	299 1,194 686 463	12.8 47.2 24.1 15.9	18.4 33.4 29.3 18.9	9.1 56.3 20.7 13.9	11.3 45.2 26.0 17.5

^{1/} INCLUDES DISCHARGE DATA FOR WHICH SEX WAS NOT STATED.

TABLE 4. NUMBER AND PERCENT DISTRIBUTION OF PATIENTS DISCHARGED FROM SHORT-STAY HOSPITALS BY TYPE OF OWNERSHIP OF HOSPITAL AND AGE OF PATIENT, ACCORDING TO SEX: UNITED STATES, 1971

TYPE OF GWNERSHIP AND AGE	1/ BOTH SEXES	MALE	FEMALE INCLUD- ING DELIV- ERIES	FEMALE EXCLUD- ING DELIV- ERIES	1/ BOTH SEXES	MALE	FEMALE INCLUD- ING DELIV- EPIES	FEMALE EXCLUD- ING DELIV- ERIES
ALL TYPES	NUMB		CHARGED PAI DUSANDS	IENTS		PERCENT DI	STPIBUTIO	N
ALL AGES	29,459	11,644	17,767	14,431	100.0	100.0	100.0	100.0
UNDER 15 YEARS 15-44 YEARS 45-64 YEARS 65 YEARS AND OVER	4,029 12,605 6,840 5,986	2,249 3,467 3,232 2,696	1,773 9,118 3,596 3,280	1,759 5,801 3,590 3,280	13.7 42.8 23.2 20.3	19.3 29.8 27.8 23.2	10.0 51.3 20.2 18.5	12.2 40.2 24.9 22.7
ALL AGES	21,589	8,502	13,048	10,667	100.0	100.0	100.0	100.0
UNDFR 15 YEARS 15-44 YEARS 45-64 YEARS 65 YEARS AND OVER	2,959 9,049 5,182 4,399	1.634 2.456 2.445 1.968	1,320 6,577 2,728 2,423	1,312 4,207 2,725 2,423	13.7 41.9 24.0 20.4	19.2 28.9 28.8 23.1	10.1 50.4 20.9 18.6	12.3 39.4 25.5 22.7
GOVERNMENT								
ALL AGES	6,629	2,660	3,961	3,120	100.0	100.0	100.0	100.0
UNDER 15 YEARS 15-44 YEARS 45-64 YEARS 65 YEARS AND OVER	909 3,023 1,390 1,306	528 867 662 603	381 2•153 727 701	376 1,318 724 701	13.7 45.6 21.0 19.7	19.8 32.6 24.9 22.7	9.6 54.3 18.3 17.7	12.0 42.3 23.2 22.5
PROPRIETARY								
ALL AGES	1,241	481	758	644	130.0	100.0	100.0	100.0
UNDER 15 YEARS 15-44 YEARS 45-64 YEARS 65 YEARS AND OVER	160 533 267 281	87 144 125 125	72 389 141 156	72 275 141 156	12.9 43.0 21.5 22.6	18.2 29.9 26.0 26.0	9.5 51.3 18.6 20.5	11.2 42.7 21.9 24.2

^{1/} INCLUDES DISCHARGE DATA FOR WHICH SEX WAS NOT STATED.

TABLE 5. NUMBER OF PATIENTS DISCHARGED FROM SHORT-STAY HOSPITALS AND DAYS OF CAPE, BY SEX, AGE, CEOGRAPHIC REGION, AND BED SIZE OF HOSPITAL: UNITED STATES, 1971

(DISCHARGES FROM NONFEDERAL SHORT-STAY HOSPITALS. EXCLUDES NEWBORN INFANTS)

	BED SIZE OF HOSPITAL							
-				0 3126 0	11035114	- I		
SEX. AGE. AND REGION	ALL SIZES	6-99 BEDS	100- 499 BEDS	500 BEDS OR MORE	ALL SIZES	6-99 BEDS	100- 499 BEDS	500 BEDS OR MORE
1/ BOTH SEXES	NUMBER	OF PATIE		HARGED	NUMBER OF DAYS OF CARE IN THOUSANDS			
UNITED STATES	29,459	5,829	18,118	5,513	231,017	37,792	143,213	50,012
UNDER 15 YEARS	4,029 12,605 6,840	781 2,253 1,244	2,543 7,748 4,270 3,558	705 2,604 1,326 877	18,773 72,359 64,304 75,581	2,958 9,722 8,542 16,570	11,265 44,459 40,976 46,513	4,55C 18,179 14,785 12,498
65 YEARS AND OVER	5,986	1,552			62,283	4,033	44,634	13,615
NORTHEAST	6,912	568	5,060	1,283		ļ	<u> </u>	
UNDER 15 YEARS	851 3,006 1,665 1,390	67 220 139 143	2,160 1,225 1,031	141 626 301 215	4,450 18,417 18,264 21,152	258 1,007 1,031 1,737	3,116 12,831 13,205 15,482	1,075 4,579 4,028 3,933
NORTH CENTRAL	9,171	1,500	5,885	1,785	73,449	10,597	46,828	16,025
UNDER 15 YEARS	1,385 3,838 2,133 1,815	207 556 331 407	907 2,511 1,340 1,126	271 771 462 282	6,396 22,930 20,477 23,646	788 2,619 2,393 4,798		1,626 5,502 5,046 3,851
SOUTH	9,136	2,707	4,431	1,998	67 , 930	17,875	33,465	16,590
UNDER 15 YEARS	1,255 3,887 2,046 1,949	351 998 568 790	647 1,916 1,025 844	257 974 452 315	5,972 22,115 17,864 21,979	1,460 4,464 3,936 8,014	11,242 9,308	1,646 6,409 4,619 3,916
WEST	4,241	1,053	2,741	446	27,355	5,287	18,286	3,781
UNDER 15 YEARS	538 1,873 996 833	157 480 206 212	345 1,161 679 556	37 233 111 65	1,954 8,897 7,699 8,804	452 1,632 1,182 2,020	5,576 5,425	202 1,689 1,092 798
MALE								
UNITED STATES	11,644	2,310	7,156	2,178	97,723	15,174	60,100	22,449
UNDER 15 YEARS	2,249 3,467 3,232 2,696	438 623 569 680	1,410 2,117 2,025 1,604	400 728 637 413	30,595 32,696	3,742 6,890	14,336 19,458 20,047	2,591 6,704 7,395 5,760
NORTHEAST	2,716	243	1,979	493	26,546	1,641	18,689	6,216
UNDER 15 YEARS	483 788 810 635	38 72 75 59	589			498	4,006 6,344	581 1,669 2,086 1,880

^{1/} INCLUDES DISCHARGE DATA FOR WHICH SEX WAS NOT STATED.

TABLE 5. NUMBER OF PATIENTS DISCHARGED FROM SHURT-STAY HOSPITALS AND DAYS OF CARE, BY SEX, AGE, GEOGRAPHIC REGION, AND BED SIZE OF HOSPITAL: UNITED STATES, 1971--CON.

				-D 6145 6	NE HOCDITA			
			B1	ED 217E C	F HOSPITA	L }	r	
SEX, AGE, AND REGION	ALL SIZES	6-99 BEDS	100- 499 BEDS	500 BEDS OR MORE	ALL SIZES	6-99 BEDS	100- 499 BEDS	500 BEDS OR MORE
MALECON.	NUMBER	OF PATION		CHARGED	NUM	BER OF DA	AYS OF CA USANDS	RE
NURTH CENTRAL	3,611	584	2,289	738	30,370	4,073	19,033	7,264
UNDER 15 YEARS	772 1,050 986 803	119 146 141 178	501 677 617 494	152 227 228 131	3,644 7,315 9,498 9,913	442 696 1,007 1,928	2,268 4,571 5,964 6,230	933 2,048 2,527 1,756
SOUTH	3,629	1,091	1,770	767	28,674	7,268	14,246	7,160
UNDER 15 YEARS	698 1,096 962 873	197 286 258 351	351 542 492 385	150 268 212 138	3,358 7,291 8,381 9,645	834 1,348 1,649 3,436	1,565 3,750 4,447 4,485	958 2,194 2,284 1,724
WEST	1,689	391	1,118	179	12,132	2,192	8,131	1,809
UNDER 15 YEARS	295 533 474 385	85 119 95 92	190 339 328 261	20 76 51 32	1,072 3,248 3,788 4,025	281 445 587 878	672 2,010 2,703 2,747	119 793 497 400
FEMALE INCLUDING DELIVERIES								
UNITED STATES	17,767	3,510	10,929	3,328	132,906	22,564	82,823	27,519
UNDER 15 YEARS	1,773 9,118 3,596 3,280	341 1,627 672 870	1,129 5,617 2,235 1,947	304 1,873 688 463	8,183 48,339 33,606 42,778	1,236 6,882 4,786 9,660	4,994 30,002 21,436 26,390	1,953 11,455 7,384 6,727
NORTHEAST	4,180	323	3,070	788	35,568	2,376	25,817	7,375
UNDER 15 YEARS	365 2,212 852 752	28 147 64 84	274 1,597 634 565	62 468 155 103	1,949 12,332 9,291 11,996	99 664 529 1,084	1,359 8,771 6,821 8,866	492 2,897 1,940 2,046
NORTH CENTRAL	5,545	913	3,586	1,045	42,973	6,507	27,713	8,753
UNDER 15 YEARS	609 2,783 1,143 1,009	87 409 189 228	404 1,830 722 631	118 544 233 150	2,743 15,572 10,951 13,706	343 1,921 1,380 2,862	1,7C7 10,201 7,053 8,751	692 3,451 2,518 2,092
SOUTH	5,495	1,612	2,654	1.228	39,172	10,588	19,163	9,420
UNDER 15 YEARS	556 2,786 1,081 1,072	153 711 310 438	296 1,371 531 456	107 704 240 177	2,608 14,801 9,465 12,297	623 3,110 2,284 4,571	1,300 7,479 4,849 5,535	685 4,212 2,333 2,190

TABLE 5. NUMBER OF PATIENTS DISCHARGED FROM SHORT-STAY HOSPITALS AND DAYS OF CARE, BY SEX, AGE, GEOGRAPHIC REGION, AND BED SIZE OF HOSPITAL: UNITED STATES, 1971--CON.

			ВЕ	D SIZE O	F HOSPITA	L			
SEX. AGE. AND REGION	ALL SIZES	6-99 BEDS	100- 499 BEDS	500 BEDS OR MORE	ALL SIZES	6-99 BEDS	100- 499 BEDS	500 BEDS OR MORE	
FEMALE INCLUDING DELIVERIES—CCN.	NUMBER	OF PATIE		HARGED	NUMBER OF DAYS OF CARE IN THOUSANDS				
WEST	2,547	562	1,619	267	15,193	3,093	10,130	1,970	
UNDER 15 YEARS	245 1+338 519 447	72 360 110 120	155 820 349 295	16 157 61 32	882 5,633 3,899 4,779	171 1,187 593 1,142	628 3,552 2,712 3,238	83 895 594 399	
FEMALE FXCLUDING DELIVERIES									
UNITED STATES	14,431	2,921	8,867	2,642	119,374	20,486	74,332	24,556	
UNDER 15 YEARS	1,759 5,801 3,590 3,280	339 1,041 672 870	1,121 3,567 2,232 1,947	299 1,194 686 463	8,130 34,887 33,580 42,778	1,231 4,812 4,783 9,660	4,964 21,556 21,421 26,390	1,935 8,518 7,376 6,727	
NORTHEAST	3,378	268	2,481	630	31,882	2,138	23,160	6,585	
UNDER 15 YEARS	363 1,413 851 752	28 92 64 84	274 1,009 633 565	61 312 154 103	1,938 8,664 9,285 11,996	98 427 529 1,084	1,353 6,124 6,817 8,866	487 2,113 1,939 2,046	
NORTH CENTRAL	4,508	768	2,889	851	38,401	5,928	24,589	7,884	
UNDER 15 YEARS	605 1,752 1,141 1,009	87 265 188 228	401 1,137 720 631	117 350 233 150	2,727 11,025 10,943 13,706	342 1,346 1,378 2,862	1,695 7,093 7,049 8,751	690 2,586 2,515 2,092	
SOUTH	4,501	1,376	2,177	948	35,546	9,824	17,412	8,31C	
UNDER 15 YEARS	550 1,800 1,079 1,072	152 475 310 438	293 898 530 456	105 427 239 177	2,586 11,208 9,454 12,297	621 2,348 2,284 4,571	1,288 5,745 4,844 5,535	676 3,116 2,327 2,190	
WEST	2,043	510	1,321	213	13,545	2,596	9,171	1,778	
UNDER 15 YEARS	242 835 519 447	71 209 110 120	154 522 349 295	16 104 61 32	879 3,989 3,898 4,779	169 692 592 1,142	627 2,594 2,712 3,238	82 704 594 399	

TABLE 6. NUMBER, PERCENT DISTRIBUTION, AND RATE OF DAYS OF CARE, AVERAGE NUMBER OF HOSPITAL BEDS OCCUPIED DAILY, AND AVERAGE LENGTH OF STAY FOR PATIENTS DISCHARGED FROM SHORT-STAY HOSPITALS, BY SEX AND AGE: UNITED STATES, 1971

			· ·		
		DAYS OF CARE			
SEX AND AGE	NUMBER IN THOUSANDS	PERCENT DISTRIBUTION	RATE PER 1,000 POPULATION	1/ NUMBER OF HOSPITAL BEDS OCCUPIED DAILY	AVERAGE LENGTH OF STAY IN DAYS
2/ BOTH SEXES					
ALL AGES	231,017	100.0	1,143.1	313.2	7.8
UNDER 1 YEAR	4,610 4,781 9,381 25,096 23,749 23,515 30,380 33,924 38,518	2.0 2.1 4.1 10.9 10.3 10.2 13.2 14.7	1,266.8 350.6 234.0 700.4 950.3 1,048.4 1,309.0 1,815.8 3,121.7	347.1 96.0 64.1 191.9 260.3 287.2 358.6 497.5 855.3	6.5 4.2 4.3 4.8 5.6 7.4 8.7 10.2
75 YEARS AND OVER	37,063	16.0	5,119.8	1,402.7	13.3
MALE					
ALL AGES	97,723	100.0	1,004.0	275.1	8.4
UNDER 1 YEAR	2,620 2,635 5,311 7,797 7,048 9,021 13,625 16,970 17,870 14,826	2.7 2.7 5.4 8.0 7.2 9.2 13.9 17.4 18.3	1,406.9 378.9 260.2 452.5 586.9 838.9 1,225.0 1,930.6 3,328.4 5,238.8	385.5 103.8 71.3 124.0 160.8 229.8 335.6 528.9 911.9 1,435.3	6.4 4.1 4.4 6.1 6.9 7.7 8.8 10.1 11.7
FEMALE	ļ				
ALL AGES	132,906	100.0	1,268.7	347.6	7.5
UNDER 1 YEAR	1,980 2,144 4,058 17,285 16,662 14,392 16,701 16,905 20,581	1.5 1.6 3.1 13.0 12.5 10.8 12.6 12.7	1,114.5 320.7 206.2 929.3 1,283.5 1,282.5 1,382.0 1,708.6 2,952.4 5,034.3	305.3 87.9 56.5 254.6 351.6 337.7 378.6 468.1 808.9	6.7 4.4 4.1 5.2 7.3 8.6 10.3 12.3

^{1/} EXPRESSED AS DAILY NUMBER OF BEDS OCCUPIED PER 100,000 CIVILIAN, NONINSTITUTIONALIZED POPULATION.

^{2/} INCLUDES DISCHARGE DATA FOR WHICH SEX WAS NOT STATED.

TABLE 7. NUMBER AND PERCENT DISTRIBUTION OF PATIENTS DISCHARGED FROM SHORT-STAY HOSPITALS BY AGE AND LENGTH OF STAY, ACCORDING TO SEX: UNITED STATES, 1971--CON.

(DIOCHANGE)		COUNTRY ONC	KI-SIAI F	100. ITALU	EVCTORES	1121103111	2111 /211107			
AGE AND LENGTH OF STAY	1/ BOTH SEXES	MALE	FEMALE INCLUD- ING DELIV- ERIES	FEMALE EXCLUD- ING DELIV- ERIES	1/ BOTH SEXES	MALE	FEMALE INCLUD- ING DELIV- ERIES	FEMALE EXCLUD- ING DELIV- EPIES		
<u>ALL AGES</u>	NUM	OF DISCH IN THOU	IARGED PAT ISANDS	IENTS	1	PERCENT DISTRIBUTION				
ALL STAYS	29,459	11,644	17,767	14,431	130.0	100.0	100.0	100.0		
LESS THAN 1 DAY 1 DAY 2 DAYS 3 DAYS 4 DAYS 5-6 DAYS 7-8 DAYS 9-10 DAYS 11-20 DAYS 21-30 DAYS 31 DAYS OR MORE.	720 2,205 4,221 3,717 3,221 4,325 2,959 1,978 4,063 1,199 852	297 957 1,622 1,209 1,049 1,726 1,204 825 1,819 554 383	421 1,245 2,593 2,504 2,168 2,588 1,752 1,147 2,238 642 468	412 1,134 2,165 1,504 1,254 1,968 1,581 1,105 2,209 637 462	2.4 7.5 14.3 12.6 10.9 14.7 10.0 6.7 13.8 4.1 2.9	2.6 8.2 13.9 10.4 9.0 14.8 10.3 7.1 15.6 4.8	2.4 7.0 14.6 14.1 12.2 14.6 9.9 6.5 12.6	2.9 7.9 15.0 10.4 8.7 13.6 11.0 7.7 15.3 4.4		
UNDER 15 YEARS ALL STAYS	4,029	2,249	1,773	1,759	100.0	100.0	100.C	100.0		
LESS THAN 1 DAY	149 675 1,082 524 392 500 263 143 211 46 44	80 381 589 294 220 288 148 83 120 23 25	69 294 491 229 170 212 115 60 90 23 19	69 293 489 224 168 210 114 60 90 23	3.7 16.8 26.9 13.0 9.7 12.4 6.5 3.5 5.2 1.1	3.5 16.9 26.2 13.1 9.8 12.8 6.6 3.7 5.3 1.0	3.9 16.6 27.7 12.9 9.6 12.0 6.5 3.4 1.3	3.9 16.7 27.8 12.7 9.6 11.9 6.5 3.4 5.1 1.3		
15-44 YEARS ALL STAYS	12,605	3,467	9,118	5,801	100.0	100.0	100.0	100.0		
LESS THAN 1 DAY	382 1,032 2,082 2,175 1,840 2,044 1,120 611 947 213	111 320 569 438 365 577 338 201 378	270 710 1.511 1.735 1.473 1.461 781 409 568 117	261 600 1,085 742 563 843 611 367 539 112	3.0 8.2 16.5 17.3 14.6 16.2 8.9 4.9 7.5	3.2 9.2 16.4 12.6 10.5 16.6 9.7 5.8	3.0 7.8 16.6 19.0 16.2 16.0 8.6 4.5 6.2	4.5 10.4 18.7 12.8 9.7 14.5 10.5 6.3 9.3		
31 DAYS OR MORE	160	76	83	77	1.3	2.2		1.3		

^{1/} INCLUDES DISCHARGE DATA FOR WHICH SEX WAS NOT STATED.

TABLE 7. NUMBER AND PERCENT DISTRIBUTION OF PATIENTS DISCHARGED FROM SHORT-STAY HOSPITALS BY AGE AND LENGTH OF STAY, ACCORDING TO SEX: UNITED STATES, 1971--CON.

101001741000	TRON NOW	EDC AL SITE	,,,, ,,,,, ,	100. I . ALU	ENGLODES	MEMBORN 1	IIII AIII 37	
AGE AND LENGTH OF STAY	1/ BOTH SEXES	MALE	FEMALE INCLUD- ING DELIV- ERIES	FEMALE EXCLUD- ING DELIV- ERIES	1/ BOTH SEXES	MALE	FEMALE INCLUD- ING DELIV- ERIES	FEMALE EXCLUD- ING DELIV- ERIES
45-64 YEARS	NUMBER OF DISCHARGED PATIENTS IN THQUSANDS				PERCENT DISTRIBUTION			
ALL STAYS	6,840	3,232	3,596	3,590	100.0	100.0	100.C	100.0
LESS THAN 1 DAY	103 309 733 642 567 1.014 865 663 1.332 381 232	60 158 315 299 272 516 395 291 623 192 111	43 150 417 341 294 496 469 369 707 189 121	43 150 416 340 293 495 469 369 707 188 121	1.5 4.5 10.7 9.4 8.3 14.8 12.6 9.7 19.5 5.6	1.9 4.9 9.8 9.3 8.4 16.0 12.2 9.0 19.3 5.9 3.4	1.2 4.2 11.6 9.5 8.2 13.8 13.0 10.3 19.7 5.2	1.2 4.2 11.6 9.5 8.2 13.8 13.1 10.3 19.7 5.2 3.4
ALL STAYS	5,986	2,696	3,280	3,280	100.0	100.0	100.0	100.0
LESS THAN 1 DAY 1 DAY 2 DAYS 3 DAYS 4 DAYS 5-6 DAYS 7-8 DAYS 9-10 DAYS 11-20 DAYS 21-30 DAYS 31 DAYS OR MORE	86 188 325 377 422 767 711 560 1,573 559 416	46 98 149 177 191 346 323 250 699 245 171	40 91 174 199 230 420 387 308 872 314 245	40 91 174 199 230 420 387 308 872 314 245	1.4 3.1 5.4 6.3 7.1 12.8 11.9 9.4 26.3 9.3 6.9	1.7 3.6 5.5 6.6 7.1 12.8 12.0 9.3 25.9 9.1 6.3	1.2 2.8 5.3 6.1 7.0 12.8 11.8 9.4 26.6 9.6 7.5	1.2 2.8 5.3 6.1 7.0 12.8 11.8 9.4 26.6 9.6 7.5

^{1/} INCLUDES DISCHARGE DATA FOR WHICH SEX WAS NOT STATED.

TABLE 8. NUMBER AND PERCENT DISTRIBUTION OF DAYS OF CARE FOR PATIENTS DISCHARGED FROM SHORT-STAY HOSPITALS BY COLOR AND AGE OF PATIENT, ACCORDING TO SEX: UNITED STATES, 1971

									
COLOR AND AGE	1/ BOTH SEXES	MALE	FEMALE INCLUD- ING DELIV- ERIES	FEMALE EXCLUD- ING DELIV- ERIES	1/ BOTH SEXES	MALE	FEMALE INCLUD- ING DELIV- ER IES	FEMALE EXCLUD- ING DELIV- ERIES	
TOTAL	N	UMBER OF E	DAYS OF CA	ARE	PERCENT DISTRIBUTION				
ALL AGES	231,017	97,723	132,906	119,374	100.0	100.0	100.0	100.0	
UNDER 15 YEARS	18,773 72,359 64,304 75,581	10,565 23,866 30,595 32,696	8,183 48,339 33,606 42,778	8,130 34,887 33,580 42,778	8.1 31.3 27.8 32.7	10.8 24.4 31.3 33.5	6.2 36.4 25.3 32.2	6.8 29.2 28.1 35.8	
<u>WHITE</u> ALL AGES	176,252	75,036	101,183	91•740	100.0	100.0	100.0	100.0	
UNDER 15 YEARS	13,083 51,375 50,339 61,454	7,285 17,118 24,005 26,628	5,796 34,240 26,331 34,816	5,776 24,833 26,315 34,816	7.4 29.1 28.6 34.9	9.7 22.8 32.0 35.5	5.7 33.8 26.0 34.4	6.3 27.1 28.7 38.0	
ALL OTHER ALL AGES	27,184	11,451	15,720	13,378	100.0	100.0	100.0	100.0	
UNDER 15 YEARS 15-44 YEARS 45-64 YEARS 65 YEARS AND OVER	3,556 11,377 6,523 5,728	2,065 3,567 3,140 2,680	1,488 7,807 3,381 3,044	1,458 5,502 3,374 3,044	13.1 41.9 24.0 21.1	18.0 31.1 27.4 23.4	9.5 49.7 21.5 19.4	10.9 41.1 25.2 22.8	
COLOR NOT STATED									
ALL AGES	27,581	11,235	16,002	14,257	100.0	100.0	100.0	100.0	
UNDER 15 YEARS 15-44 YEARS 45-64 YEARS 65 YEARS AND OVER	2,134 9,607 7,442 8,398	1,215 3,182 3,450 3,388	899 6•292 3•894 4•918	896 4,552 3,891 4,918	7.7 34.8 27.0 30.4	10.8 28.3 30.7 30.2	5.6 39.3 24.3 30.7	6.3 31.9 27.3 34.5	

^{1/} INCLUDES DISCHARGE DATA FOR WHICH SEX WAS NOT STATED.

TABLE 9. AVERAGE LENGTH OF STAY FOR PATIENTS DISCHARGED FROM SHORT-STAY HOSPITALS BY COLOR, AGE, AND SEX: UNITED STATES, 1971

the same of the sa								
COLOR AND AGE	1/ BOTH SEXES	MALE	FEMALE INCLUDING DELIVERIES	FEMALE EXCLUDING DELIVERIES				
<u>TCTAL</u>	AVERAGE LENGTH OF STAY IN DAYS							
ALL AGES	7.7	4	7.5	R.3				
UNDER 15 YEARS	4.7 5.7 9.4 12.6	4.7 6.9 9.5 12.1	4.6 5.3 9.3 13.0	4.6 6.0 9.4 13.0				
<u>WHITE</u>								
ALL AGES	7.9	8.3	7.5	8.3				
UNDER 15 YEARS	4.4 5.6 9.3 12.6	4.4 6.6 9.3 12.1	4.4 5.2 9.2 13.0	4.4 5.9 9.2 13.0				
ALL CTHER								
ALL AGES	8.1	9.6	7.4	8.6				
UNDER 15 YEARS	6.5 6.3 11.5 14.1	6.4 8.4 11.8 14.2	6.5 5.6 11.1 14.1	6.6 6.7 11.2 14.1				
COLOR NOT STATED								
ALL AGES	7.5	7.8	7.3	8.0				
UNDER 15 YEARS	4.2 [.] 5.9 8.9 12.1	4.4 6.9 8.7 11.2	4.1 5.4 9.0 12.8	4.1 6.1 9.0 12.8				

^{1/} INCLUDES DISCHARGE DATA FOR WHICH SEX WAS NOT STATED.

TABLE 10. NUMBER AND PERCENT DISTRIBUTION OF DAYS OF CARE FOR PATIENTS DISCHARGED FROM SHOPT-STAY HOSPITALS BY GEOGRAPHIC REGION AND AGE, ACCORDING TO SEX: UNITED STATES, 1971

REGION AND AGE	1/ BOTH SEXES	MALE	FEMALE INCLUD- ING DELIV- ERIES	FEMALE EXCLUD- ING DELIV- ERIES	1/ BOTH SEXES	MALE	FEMALE INCLUD- ING DELIV- ERIES	FEMALE EXCLUD- ING DELIV- ERIES	
UNITED STATES	NUMBER OF DAYS OF CARE IN THOUSANDS .				PERCENT DISTRIBUTION				
ALL AGES	231,017	97,723	132,906	119,374	100.0	100.0	100.0	100.0	
UNDER 15 YEARS	18,773 72,359 64,304 75,581	10,565 23,866 30,595 32,696	8,183 48,339 33,606 42,778	8,130 34,887 33,580 42,778	8.1 31.3 27.8 32.7	10.8 24.4 31.3 33.5	6.2 35.4 25.3 32.2	6.8 29.2 28.1 35.8	
NORTHEAST									
ALL AGES	62,283	26,546	35,568	31,882	100.0	100.0	100.0	100.0	
UNDER 15 YEARS	4,450 18,417 18,264 21,152	2,493 6,012 8,928 9,113	1,949 12,332 9,291 11,996	1,938 8,664 9,285 11,996	7•1 29•6 29•3 34•0	9.4 22.6 33.6 34.3	5.5 34.7 26.1 33.7	6.1 27.2 29.1 37.6	
NORTH CENTRAL									
ALL AGES	73,449	30,370	42,973	38,401	100.0	100.0	100.0	100.0	
UNDER 15 YEARS	6,396 22,930 20,477 23,646	3,644 7,315 9,498 9,913	2,743 15,572 10,951 13,706	2,727 11,025 10,943 13,706	8.7 31.2 27.9 32.2	12.0 24.1 31.3 32.6	6.4 36.2 25.5 31.9	7.1 28.7 28.5 35.7	
SOUTH									
ALL AGES	67,930	28,674	39,172	35,546	100.0	100.0	100.0	100.0	
UNDER 15 YEARS	5,972 22,115 17,864 21,979	3,358 7,291 8,381 9,645	2,608 14,801 9,465 12,297	2,586 11,208 9,454 12,297	8.8 32.6 26.3 32.4	11.7 25.4 29.2 33.6	6.7 37.8 24.2 31.4	7.3 31.5 26.6 34.6	
WEST									
ALL AGES	27,355	12,132	15,193	13,545	100.0	100.0	100.0	100.0	
UNDER 15 YEARS	1.954 8.897 7.699 8.804	1,072 3,248 3,788 4,025	882 5,633 3,899 4,779	879 3,989 3,898 4,779	7.1 32.5 28.1 32.2	8.8 26.8 31.2 33.2	5.8 37.1 25.7 31.5	6.5 29.5 28.8 35.3	

^{1/} INCLUDES DISCHARGE DATA FOR WHICH SEX WAS NOT STATED.

TABLE 11. AVERAGE LENGTH OF STAY FOR PATIENTS DISCHARGED FROM SHORT-STAY HOSPITALS BY GEOGRAPHIC REGION, AGE, AND SEX: UNITED STATES, 1971

REGION AND AGE	1/ BOTH SEXES	MALE	FEMALE INCLUDING DELIVERIES	FEMALE EXCLUDING DELIVERIES
<u>UNITED STATES</u>	AV	ERAGE LENGTH	OF STAY IN D	AYS
ALL AGES	7.8	8.4	7.5	8.3
UNDER 15 YEARS	4.7 5.7 9.4 12.6	4.7 6.9 9.5 12.1	4.6 5.3 9.3 13.0	4.6 6.0 9.4 13.0
NORTHEAST ALL AGES	9.0	9.8	8.5	9.4
UNDER 15 YEARS	5.2 6.1 11.0 15.2	5.2 7.6 11.0 14.4	5.3 5.6 10.9 16.0	5.3 6.1 10.9 16.0
NORTH CENTRAL				
ALL AGES	8.0	8.4	7.8	8.5
UNDER 15 YEARS	4.6 6.0 9.6 13.0	4.7 7.0 9.6 12.4	4.5 5.6 9.6 13.6	4.5 6.3 9.6 13.6
<u>SOUTH</u>				
ALL AGES	7.4	7.9	7.1	7.9
UNDER 15 YEARS	4.8 5.7 8.7 11.3	4.8 6.7 8.7 11.0	4.7 5.3 8.8 11.5	4.7 6.2 8.8 11.5
WEST				
ALL AGES	6.5	7.2	6.0	6.6
UNDER 15 YEARS	3.6 4.7 7.7 10.6	3.6 6.1 8.0 10.4	3.6 4.2 7.5 10.7	3.6 4.8 7.5 10.7

^{1/} INCLUDES DISCHARGE DATA FOR WHICH SEX WAS NOT STATED.

TABLE 12. NUMBER AND PERCENT DISTRIBUTION OF DAYS OF CARE FOR PATIENTS DISCHAFGED FROM SHORT-STAY HOSPITALS BY BED SIZE OF HOSPITAL AND AGE OF PATIENT, ACCORDING TO SEX: UNITED STATES, 1971

(DISCHARGES FROM NONFEDERAL SHORT-STAY HOSPITALS. EXCLUDES NEWBORN INFANTS)

BED SIZE OF HOSPITAL AND AGE	1/ BOTH SEXES	MALE	FEMALE INCLUD- ING DELIV- ERIES	FEMALE EXCLUD- ING DELIV- ERIES	1/ BOTH SEXES	MALE	FEMALE INCLUD- ING DELIV- EFIES	FEMALE EXCLUD- ING DELIV- EPIES	
ALL SIZES	N	UMBER OF L	IAYS OF CA	RE	∼ERCENT DISTRIBUTION				
ALL AGES	231,017	97,723	132,906	119,374	100.0	100.0	100.0	100.0	
UNDER 15 YEARS	18,773 72,359 64,304 75,581	10,565 23,866 30,595 32,696	8,183 48,339 33,606 42,778	8,130 34,887 33,580 42,778	8.1 31.3 27.8 32.7	10.8 24.4 31.3 33.5	6.2 36.4 25.3 32.2	6.8 29.2 28.1 35.8	
6-99 BEDS									
ALL AGES	37,792	15,174	22,564	20,486	100.0	100.0	100.0	100.0	
UNDER 15 YEARS 15-44 YEARS 45-64 YEARS 65 YEARS AND OVER	2,958 9,722 8,542 16,570	1,716 2,826 3,742 6,890	1,236 6,882 4,786 9,660	1,231 4,812 4,783 9,660	7.8 25.7 22.6 43.8	11.3 18.6 24.7 45.4	5.5 30.5 21.2 42.8	6.0 23.5 23.3 47.2	
100-199 BEDS							:		
ALL AGES	44,274	18,142	26,062	23,348	100.0	100.0	100.0	100.0	
UNDER 15 YEARS 15-44 YEARS 45-64 YEARS 65 YEARS AND OVER	3,656 12,777 11,642 16,198	1,958 3,729 5,523 6,932	1,695 9,029 6,095 9,242	1,682 6,333 6,089 9,242	8.3 28.9 26.3 36.6	10.8 20.6 30.4 38.2	6.5 34.6 23.4 35.5	7.2 27.1 26.1 39.6	
200-299 BEDS									
ALL AGES	40,775	16,822	23,872	21,346	100.0	100.0	100.0	100.0	
UNDER 15 YEARS 15-44 YEARS 45-64 YEARS 65 YEARS AND OVER	3,351 12,554 12,050 12,820	1,860 3,917 5,618 5,427	1,489 8,605 6,406 7,373	1,483 6,085 6,404 7,373	8.2 30.8 29.6 31.4	11.1 23.3 33.4 32.3	6.2 36.0 26.8 30.9	6.9 28.5 30.0 34.5	
300-499 BEDS						:			
ALL AGES	58,165	25,136	32,889	29,638	100.0	100.0	100.0	100.0	
UNDER 15 YEARS 15-44 YEARS 45-64 YEARS 65 YEARS AND OVER	4,259 19,128 17,284 17,494	2,441 6,691 8,317 7,688	1,810 12,368 8,935 9,775	1,798 9,138 8,928 9,775	7.3 32.9 29.7 30.1	9.7 26.6 33.1 30.6	5.5 37.6 27.2 29.7	6.1 30.8 30.1 33.0	
500 BEDS OR MORE									
ALL AGES	50,012	22,449	27,519	24,556	100.0	100.0	100.0	100.0	
UNDER 15 YEARS	4,550 18,179 14,785 12,498	2,591 6,704 7,395 5,760	1,953 11,455 7,384 6,727	1,935 8,518 7,376 6,727	9.1 36.3 29.6 25.0	11.5 29.9 32.9 25.7	7.1 41.6 26.8 24.4	7.9 34.7 30.0 27.4	

^{1/} INCLUDES DISCHARGE DATA FOR WHICH SEX WAS NOT STATED.

TABLE 13. AVERAGE LENGTH OF STAY FOR PATIENTS DISCHARGED FROM SHOPT-STAY HOSPITALS BY BED SIZE CF HOSPITAL, AGE UF PATIENT, AND SEX: UNITED STATES, 1971

BED SIZE OF HOSPITAL AND AGE	1/ BOTH SEXES	MALE	FEMALE INCLUDING DELIVERIES	FEMALE EXCLUDING DELIVERIES
ALL SIZES		AVERAGE LENGTH	OF STAY IN DAY	'S
ALL AGES	7.8	8.4	7.5	8.3
UNDER 15 YEARS	4.7 5.7 9.4 12.6	4.7 6.9 9.5 12.1	4.6 5.3 9.3 13.0	4.6 6.0 9.4 13.0
<u>6-99 BFNS</u>				
ALL AGES	6.5	6.6	6.4	7.0
UNDER 15 YEARS	3.8 4.3 6.9 10.7	3.9 4.5 6.6 10.1	3.6 4.2 7.1 11.1	3.6 4.6 7.1 11.1
100-199 BEDS ALL AGES	7.3	7.7	7.1	7.0
UNDER 15 YEARS	4.3 5.1 8.5 12.3	4.2 5.7 8.6 11.7	4.4 4.9 8.4 12.9	7.8 4.4 5.4 8.4 12.9
200-299 BEDS				
ALL AGES	7.9	8.2	7.6	8.5
UNDER 15 YEARS	4.3 5.6 10.1 13.2	4.4 6.5 9.9 12.3	4.3 5.3 10.3 13.9	4.3 6.0 10.3 13.9
ALL AGES	8.5	9.1	8.0	8.9
UNDER 15 YEARS	4.6 6.4 10.2 13.7	4.7 7.8 10.2 13.5	4.6 5.8 10.1 14.0	4.6 6.7 10.1 14.0
500 BEDS CR MORE				
ALL AGES	9.1	10.3	8.3	9.3
UNDER 15 YEARS	6.4 7.0 11.1 14.3	6.5 9.2 11.6 14.0	6.4 6.1 10.7 14.5	6.5 7.1 10.7 14.5

^{1/} INCLUDES DISCHARGE DATA FOR WHICH SEX WAS NOT STATED.

TABLE 14. NUMBER AND PERCENT DISTRIBUTION OF DAYS OF CARE FOR PATIENTS DISCHAPGED FROM SHOPT-STAY HOSPITALS BY TYPE OF OWNERSHIP CF HUSPITAL AND AGE OF PATIENT, ACCORDING TO SEX: UNITED STATES, 1971

(DISCHARGES FROM NONFEDERAL SHORT-STAY HOSPITALS. EXCLUDES NEWBORN INFANTS)

TYPE OF OWNERSHIP AND AGE	1/ BOTH SEXES	MALE	FEMALE INCLUD- ING DELIV- ERIES	FEMALE EXCLUD- ING DELIV- ERIES	1/ BOTH SEXES	MALE	FEMALE INCLUD- ING DELIV- EPIES	FEMALE EXCLUD- ING DELIV- ERIES
ALL TYPES	N	UMBER OF D IN TH	AYS OF CA	APE .	PERCENT DISTRIBUTION			
ALL 'AGES	231,017	97,723	132,906	119,374	100.0	100.0	100.0	100.0
UNDER 15 YEARS	18,773 72,359 64,304 75,581	10,565 23,866 30,595 32,696	8,183 48,339 33,606 42,778	8,130 34,887 33,580 42,778	8.1 31.3 27.8 32.7	10.8 24.4 31.3 33.5	6.2 36.4 25.3 32.2	6.8 29.2 28.1 35.8
VOLUNTARY NONPROFIT ALL AGES	173,457	72,715	100,430	90,441	100.0	100.0	100.0	100.0
UNDER 15 YEARS	13,571 53,162 49,711 57,013	7,514 17,096 23,606 24,499	6,039 35,933 26,024 32,435	6,004 25,997 26,006 32,435	7.8 30.6 28.7 32.9	10.3 23.5 32.5 33.7	6.0 35.8 25.9 32.3	6.6 28.7 28.8 35.9
GOVERNMENT ALL AGES	49,024	21,597	27,366	24,262	100.0	100.0	100.0	100.0
UNDER 15 YEARS	4,629 16,638 12,475 15,283	2,737 5,967 6,034 6,860	1,887 10,651 6,427 8,400	1,870 7,573 6,419 8,400	9.4 33.9 25.4 31.2	12.7 27.6 27.9 31.8	6.9 38.9 23.5 30.7	7 • 7 31 • 2 26 • 5 34 • 6
PROPRIETARY								
ALL AGES	8,535	3,411	5,109	4,671	100.0	100.0	100.0	100.0
UNDER 15 YEARS	572 2,559 2,118 3,285	315 804 954 1,338	256 1,755 1,155 1,943	256 1,317 1,155 1,943	6.7 30.0 24.8 38.5	9.2 23.6 28.0 39.2	5.0 34.3 22.6 38.0	5.5 28.2 24.7 41.6

^{1/} INCLUDES DISCHARGE DATA FOR WHICH SEX WAS NOT STATED.

TABLE 15. AVERAGE LENGTH OF STAY FOR PATIENTS DISCHARGED FROM SHORT-STAY HOSPITALS BY TYPE OF OWNERSHIP OF HOSPITAL, AGE OF PATIENT, AND SEX: UNITED STATES, 1971

(DISCHARGES FROM NONFEDERAL SHORT-STAY HOSPITALS. EXCLUDES NEWBORN INFANTS)

TYPE OF OWNERSHIP AND AGE	1/ BOTH SEXES	MALE	FEMALE INCLUDING DEL IVEP IES	FEMALE EXCLUDING DELIVERIES
ALL TYPES	AVERAGE LENGTH OF STAY IN DAYS			
ALL AGES	7.8	8.4	7.5	8.3
UNDER 15 YEARS	4.7 5.7 9.4 12.6 8.0 4.6 5.9 9.6	4.7 6.9 9.5 12.1 8.6 4.6 7.0 9.7	4.6 5.3 9.3 13.0 7.7 4.6 5.5	4.6 6.0 9.4 13.0 8.5 4.6 6.2 9.5
65 YEARS AND OVER	13.0	12.4	13.4	13.4
GOVERNMENT ALL AGES UNDER 15 YEARS 15-44 YEARS 45-64 YEARS 65 YEARS AND OVER	7.4 5.1 5.5 9.0 11.7	8.1 5.2 6.9 9.1 11.4	5.0 4.9 8.8 12.0	7.8 5.0 5.7 8.9 12.0
PROPRIETARY				
ALL AGES	6.9	7.1	6.7	7.3
UNDER 15 YEARS	3.6 4.8 7.9 11.7	3.6 5.6 7.6 10.7	3.6 4.5 8.2 12.5	3.6 4.8 8.2 12.5

^{1/} INCLUDES DISCHARGE DATA FOR WHICH SEX WAS NOT STATED.

TABLE 16. AVERAGE LENGTH OF STAY FOR PATIENTS DISCHARGED FROM SHORT-STAY HOSPITALS BY SEX, AGE, GEOGRAPHIC REGION, AND BED SIZE OF HOSPITAL: UNITED STATES, 1971

(DISCHARGES FROM NONFEDERAL SHORT-STAY HOSPITALS. EXCLUDES NEWBORN INFANTS)

		NO	ORTHEA	ST	NORTH CENTRAL		SOUTH		WEST				
SEX AND AGE	TOTAL	6-99 BEDS	100- 499 BEDS	500 BEDS OR MORE	6-99 BEDS	100- 499 PENS	500 BEDS OR MORE	6-99 BEDS	100- 499 BEDS	500 BEDS OR MORE	6-99 3EDS	100 499 BEDS	500 BEDS OR MORE
1/ BOTH SEXES		AVERAGE LENGTH OF STAY IN DAYS											
ALL AGES	7.8	7.1	8.8	10.6	7.1	8.0	9.0	6.6	7.6	8.3	5.0	6.7	8.5
UNDER 15 YEARS 15-44 YEARS 45-64 YEARS 65+ YEARS	4.7 5.7 9.4 12.6	3.9 4.6 7.4 12.1	4.8 5.9 10.8 15.0	7.6 7.3 13.4 18.3	3.8 4.7 7.2 11.8	4.4 5.9 9.7 13.3	6.0 7.1 10.9 13.7	4.2 4.5 6.9 10.1	4.4 5.9 9.1 11.9	6.4 6.6 10.2 12.4	2.9 3.4 5.8 9.5	3.8 4.8 8.0 10.8	5.5 7.2 9.8 12.3
ALL AGES	8.4	6.7	9.4	12.6	7.0	8.3	9.8	6.7	8.0	9.3	5.6	7.3	10.1
UNDER 15 YEARS 15-44 YEARS 45-64 YEARS 65+ YEARS	4.7 6.9 9.5 12.1	4.2 4.7 6.7 11.0	4.8 7.2 10.8 14.2	7.5 10.6 14.3 16.8	3.7 4.8 7.1 10.8	4.5 6.7 9.7 12.6	6.1 9.0 11.1 13.4	4.2 4.7 6.4 9.8	4.5 6.9 9.0 11.6	6.4 8.2 10.8 12.5	3.3 3.7 6.2 9.5	3.5 5.9 8.2 10.5	5.8 10.5 9.8 12.3
FEMALE INCLUD- ING DELIVERIES													
ALL AGES	7.5	7.4	8.4	9.4	7.1	7.7	8.4	6.6	7.2	7.7	4.7	6.3	7.4
UNDER 15 YEARS 15-44 YEARS 45-64 YEARS 65+ YEARS	4.6 5.3 9.3 13.0	3.5 4.5 8.3 13.0	5.0 5.5 10.8 15.7	7.9 6.2 12.5 19.8	3.9 4.7 7.3 12.5	4.2 5.6 9.8 13.9	5.9 6.3 10.8 13.9	4.1 4.4 7.4 10.4	4.4 5.5 9.1 12.1	6.4 6.0 9.7 12.4	2.4 3.3 5.4 9.6	4.1 4.3 7.8 11.0	5.1 5.7 9.8 12.3
FEMALE EXCLUD- ING DELIVERIES													
ALL AGES	8.3	8.0	9.3	10.5	7.7	8.5	9.3	7.1	8.0	8.8	5.1	6.9	8.3
UNDER 15 YEARS 15-44 YEARS 45-64 YEARS 65+ YEARS	4.6 6.0 9.4 13.0	3.5 4.6 8.3 13.0	4.9 6.1 10.8 15.7	8.0 6.8 12.6 19.8	3.9 5.1 7.3 12.5	4.2 6.2 9.8 13.9	5.9 7.4 10.8 13.9	4.1 4.9 7.4 10.4	4.4 6.4 9.1 12.1	6.5 7.3 9.7 12.4	2.4 3.3 5.4 9.6	4.1 5.0 7.8 11.0	5.2 6.7 9.8 12.3

^{1/} INCLUDES DISCHARGE DATA FOR WHICH SEX WAS NOT STATED.

APPENDIX I

TECHNICAL NOTES ON METHODS

Statistical Design of the Hospital Discharge Survey

Scope of the survey.—The scope of the Hospital Discharge Survey (HDS) encompasses patients discharged from noninstitutional hospitals which have six beds or more for inpatient use, are located in the 50 States and the District of Columbia, and have an average length of stay of less than 30 days. Although all discharges of inpatients from these hospitals are within the scope of the survey, all newborn infants are excluded from this report.

Sampling frame and size of hospital.—The universe (sampling frame) for the Hospital Discharge Survey consists of the short-stay hospitals excluding military and Veterans Administration hospitals, which are included in the Master Facility Inventory of Hospitals and Institutions (MFI). A detailed description of how the MFI was developed, its content, plans for maintaining it, and procedures for assessing the completeness of its coverage is published in an earlier report. ¹⁶

There were 6,965 hospitals in the universe. The distribution of short-stay hospitals by bed size and region in the universe and in the HDS sample is shown in table I. The sample for 1971 consisted of 465 hospitals, of which 26 were ruled out of scope of the survey because they failed to meet the definition of a short-stay hospital and 60 refused to participate. Estimates are based on 198,223 abstracts received from the remaining 379 hospitals that participated during 1971.

Sample design.—All hospitals with 1,000 beds or more in the universe of short-stay hospitals were selected with certainty in the sample. All hospitals with fewer than 1,000 beds were stratified, the primary strata being the 24 size-by-region classes shown in table I. Within each of these 24 primary strata, the allocation of the hospitals was made through a controlled selection technique so that hospitals in the sample would be properly distributed with regard to ownership and geographic division. Sample hospitals were drawn with probabilities ranging from certainty for the largest hospitals to 1 in 40 for the smallest hospitals.

The within-hospital sampling ratio for selecting discharges varied inversely with the probability of se-

NOTE: The list of references follows the text.

lection of the hospital. The smallest sampling fraction of discharged patients was taken in the largest hospitals, and the largest fraction was taken in the smallest hospitals. This was done to compensate for the fact that hospitals were selected with probabilities proportionate to their size class and to assure that the overall probability of selecting a discharge would be approximately the same in all hospitals.

In all hospitals the daily listing sheet of discharges was the frame from which the subsamples of discharges were selected within the sample hospitals. The sample discharges were selected by a random technique, usually on the basis of the terminal digit(s) of the patient's medical record number—a number assigned when the patient was admitted to the hospital. If the hospital's daily discharge listing did not show the medical record numbers, the sample was selected by starting with a randomly selected discharge and taking every Kth discharge thereafter.

Estimation.—Statistics produced by the HDS are derived by a complex procedure. The basic unit of estimation is the sample patient abstract. The estimating procedure used to produce essentially unbiased national estimates has three principal components: (1) inflation of reciprocals of the probabilities of sample selection, (2) adjustment for nonresponse, and (3) ratio adjustments to fixed totals. These components are described in appendix I of two earlier publications. ^{1,2}

Data collection.—Depending on the study procedure agreed on with the hospital administrator, the sample selection and the transcription of information from the hospital records to the abstract forms were performed either by the hospital staff or by representatives of the National Center for Health Statistics (NCHS) or by both. In more than three-fourths of the hospitals that participated in the HDS during 1971, this work was performed by the medical records department of the hospital. In the remaining hospitals, nearly all the work was performed by personnel of the U.S. Bureau of the Census acting for NCHS.

Nearly all survey hospitals transcribed data from hospital records to the form shown in figure I.

Data processing and editing of data.—Shipments of completed abstract forms for each sample hospital were transmitted along with sample selection control sheets to NCHS for processing. Every shipment of

Table I. Distribution of short-stay hospitals in the universe (MFI), in the Hospital Discharge Survey sample, and participating in the survey, by bed size of hospital and geographic region: United States, 1971

Bed size of hospital	All regions	Northeast	North Central	South	West		
All sizes	Number of hospitals						
Universe Total sample Survey participants	6,965 465 379	1,107 123 108	1,979 139 121	2,620 135 102	1,259 68 48		
6-49 beds					į		
Universe Total sample Survey participants	3,113 59 35	199 7 5	830 17 13	1,438 23 11	646 12 6		
50-99 beds							
Universe Total sample Survey participants	1,623 66 56	288 12 10	442 18 17	587 24 20	306 12 9		
100-199 beds							
Universe Total sample Survey participants	1,144 95 80	277 24 23	378 30 26	332 29 24	157 12 7		
200-299 beds							
Universe Total sample Survey participants	552 83 68	182 29 27	151 24 21	134 18 12	85 12 8		
300-499 beds							
Universe Total sample Survey participants	386 89 - 75	110 24 19	129 29 25	96 24 21	51 12 10		
500-999 beds							
Universe Total sample Survey participants	129 55 47	42 18 15	46 18 16	28 12 9	13 7 7		
1,000 beds or more							
Universe Total sample Survey participants	18 18 18	9 9 9	3 3 3	5 5 5	1 1 1		

abstracts was reviewed; each abstract form was checked for completeness; and when necessary, problems were referred to the hospitals for clarification and correction.

Final editing was done by computer inspection of the demographic data compared with the category code assigned each abstract. If the patient's sex was left blank, it was coded and tabulated as "not stated," except in those cases known to be deliveries.

Very few rejects were encountered; those found

were corrected by inspection of data on the computer tape. If age was left blank, it was imputed by assigning the patient an age consistent with the ages of other patients with the same category code. If the dates of admission or discharge were not given and if they could not be obtained from the monthly sample listing sheet transmitted by the sample hospital, a length of stay was imputed by assigning the patient a stay consistent with the stays of other patients of the same age. Other

CONFIDENTIAL - All information which would permit identification of an individual or of an establishment will be held confidential, will be used only by persons engaged in and for the purposes of the survey and will not be disclosed or released to other persons or used for any other purpose.

DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE Public Health Service Health Services and Mental Health Administration

National Center for Health Statistics

MEDICAL ABSTRACT - HOSPITAL DISCHARGE SURVEY

. Patie	nt Identification				
1.	Hospital number	4. Date of admission			
2.	HDS number	5. Date of discharge	Month	Day	Year
3.	Medical record number	o. Date of discharge	Month	Day	Year
l. Patie	nt Characteristics				
1. Date of birth:		2. Age (complete ONL)	1	(1	years
	Month Day Ye	ar if date of birth not		•	months
				(3	days
3.	Sex: 1 Male 2 Female				
4.	Race or color: 1 White 2 Negro	3 Other nonwhite 4 0	'Nonwhite'	5 🗌 Not	stated
5.	Marital status: 1 Married 2 Single	e 3 Widowed 4 Divorced	5 Separa	ited 6 🗆 N	ot stated
6.	Discharge status: 1 Alive 2	Dead			
2.	Operations:			∏ see	reverse sid
complete	ed by	Date		see	reverse sid
OR NCF	IS USE ONLY				*
iagnose	es				
			_ 		
peration	18				
				GPO	· 1971 () - 450-5

Figure I. Medical abstract for the Hospital Discharge Survey.

Table II. Civilian noninstitutional population by age and sex: United States, July 1, 1971

[Consistent with Current Poulation Reports, Series P-25, No.-490]

		· · · · · · · · · · · · · · · · · · ·	
Age group	Both sexes	Male	Female
Total	202,090	97,330	104,760
0-14 years Under 1 year 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-64 years 55-64 years	57,368 3,639 13,639 40,089 83,252 35,830 24,992 22,430 41,891 23,208 18,683	29,227 1,862 6,953 20,412 39,992 17,229 12,010 10,753 19,912 11,123 8,790	28,141 1,777 6,686 19,678 43,259 18,600 12,982 11,677 21,979 12,085 9,894
65 years and over	19,579 12,339 7,239	8,199 5,369 2,830	11,380 6,971 4,409

missing demographic items were coded and tabulated as not stated.

Population Estimates

The base populations used in computing rates are unpublished estimates for the U.S. civilian, noninstitutionalized population as of July 1, 1971, provided by the U.S. Bureau of the Census.

The population estimates for the United States by age and sex (table II) and by age and geographic region (table III) are consistent with the estimates of the civilian population published by the U.S. Bureau of the Census in *Current Population Reports*, Series P-25. However, they are not official population estimates of the U.S. Bureau of the Census. Estimates of the regional populations by age and sex were provided by the U.S. Bureau of the Census specifically for use in the HDS for computing rates.

General Qualifications

Rounding of numbers.—Estimates of the number of discharges and number of days of care were rounded to the nearest thousand for tabular presentation. Percents

Table III. Civilian noninstitutional population by geographic region, sex, and age: United States, July 1, 1971

Sex and age	All regions	Northeast	North Central	South	West			
<u>Total</u>		In thousands						
All ages	202,090	48,818	56,310	62,376	34,586			
Under 65 years Under 15 years 15-44 years 45-64 years 65 years and over	182,511 57,368 83,252 41,891 19,579	43,787 13,216 19,708 10,863 5,031	50,783 16,267 23,076 11,440 5,527	56,385 17,984 25,795 12,606 5,991	31,555 9,900 14,673 6,982 3,030			
Male	07 220	02.400	07.050	00.070				
All ages	97,330	23,400	27,358	29,818	16,754			
Under 65 years	89,131 29,227 39,992 19,912 8,199	21,341 6,744 9,486 5,111 2,059	25,024 8,296 11,237 5,491 2,334	27,302 9,145 12,228 5,929 2,516	15,464 5,042 7,041 3,381 1,290			
Female								
All ages	104,760	25,418	28,952	32,558	17,832			
Under 65 years	93,379 28,141 43,259 21,979 11,380	22,446 6,472 10,222 5,752 2,972	25,759 7,971 11,839 5,949 3,193	29,083 8,839 13,567 6,677 3,475	16,091 4,858 7,632 3,601 1,741			

and rates were calculated on the basis of unrounded estimates. Due to rounding, detailed figures within tables do not always add to totals.

Patient characteristics not stated.—Age and/or sex was not stated for less than 1 percent of all discharges. However, color was not stated for approximately 12 percent of the patients discharged. The proportion of sample hospital records with color not stated varied considerably among the sample hospitals.

Reliability of Estimates

Estimates from sample surveys such as the Hospital Discharge Survey are subject to two types of errors—measurement or nonsampling errors and sampling errors. Measurement errors can occur in a complete count or census as well as in a sample survey.

Sampling errors, on the other hand, occur because a sample instead of a complete count is taken.

Measurement errors.—Measurement errors include those due to hospital nonresponse, missing abstracts, information incompletely or inaccurately recorded on abstract forms, and processing errors. Some of these have been discussed in earlier sections.

Sampling errors.—The standard error in this survey is primarily a measure of the sampling variability that occurs by chance because the estimates are based

Table IV. Approximate standard errors of percentages shown in this report for discharges: patient characteristics classified by geographic region and for all hospitals

[Standard errors for patient characteristics classified by size of hospital are 1½ times and by type of ownership are 3½ times the standard errors shown in this table]

Number of	Estimate percent								
discharges (base of percent)	2 or 98	4 or 96	10 or 90	20 or 80	30 or 70	50			
	Standard error expressed in percentage points								
100,000 200,000 600,000 1,000,000 2,000,000 6,000,000 10,000,000 20,000,000 30,000,000	1.4 1.0 0.6 0.5 0.3 0.2 0.1 0.1	2.0 1.4 0.8 0.6 0.5 0.3 0.2 0.1	3.1 2.2 1.3 1.0 0.7 0.4 0.3 0.2 0.2	4.2 3.0 1.7 1.3 0.9 0.5 0.4 0.3 0.2	4.8 3.4 2.0 1.5 1.1 0.6 0.5 0.3	5.2 3.7 2.1 1.7 1.2 0.7 0.5 0.4 0.3			

Illustration of use of table IV. Table 1 shows that 28.5 percent of the 9,011,000 white male patients discharged during 1971 from all hospitals were aged 45-64 years. Linear interpolation between the values shown in table IV will yield an approximate standard error of 0.5 percent for an estimate of 28.5 percent with a base of 9,011,000.

on subsamples of discharges within a sample of shortstay hospitals rather than on all discharges from all short-stay hospitals. The relative standard error of an estimate is obtained by dividing the standard error of the estimate by the estimate itself and is expressed as a percentage of the estimate.

The chances are about 68 out of 100 that the value obtained in a complete enumeration is contained in the interval represented by the estimate plus and minus one standard error of the estimate; 95 out of 100 for two standard errors; and 99 out of 100 for 2% standard errors. Applying the illustration at the bottom of figure II, the chances are about 68 out of 100 that the value that would be obtained in a complete enumeration is contained in the interval 3,887,000 + 5.1 percent of 3,887,000 (between 3,688,163 and 4,085,237); 99 out of 100 for the interval 3,887,000 + 5.1 percent of 3,887,000 multiplied by 2.5.

The standard error of one statistic is generally different from that of another even when the two come from the same survey. In order to derive standard errors that would be applicable to a wide variety of statistics and that could be prepared at a moderate cost, a number of approximations were required. As a result, figure II and tables IV and V provide general

Table V. Approximate standard errors of percentages shown in this report for days of care: patient characteristics classified by geographic region and for all hospitals

[Standard errors for patient characteristics classified by size of hospital are 1½ times and by type of ownership are 2½ times the standard errors shown in this table]

Number of days of care (base of percent)	Estimated percent								
	2 or 98	4 or 96	10 or 90	20 or 80	30 or 70	50			
	Standard error expressed in percentage points								
1,000,000 2,000,000 6,000,000 10,000,000 20,000,000 100,000,000 200,000,000 300,000,000	1.8 1.3 0.7 0.6 0.4 0.2 0.2 0.1	1.8 1.0 0.8	4.0 2.8 1.6 1.2 0.9 0.5 0.4 0.3 0.2	5.2 3.7 2.1 1.6 1.2 0.7 0.5 0.4 0.3	6.0 4.2 2.4 1.9 1.3 0.8 0.6 0.4 0.3	6.5 4.6 2.7 2.1 1.5 0.8 0.7 0.5 0.4			

Illustration of use of table V: Table 12 shows that of the 22,449,000 days of care provided for males discharged during 1971 from hospitals with 500 beds or more 25.7 percent of the days were utilized by patients 65 years and over. Linear interpolation between the values shown in table V will yield an approximate standard error of 1.9 percent for an estimate of 25.7 percent with a base of 22,449,000.

Figure II. Approximate relative standard errors of estimated number of patients discharged for patient characteristics, by geographic region, and/or size of hospital, and type of ownership and for all hospitals.




Illustration of use of figure II: As shown in table 2, an estimated 3,887,000 patients aged 15-44 years were discharged during 1971 within the South Region. The relative standard error of this estimate as read from the line "Region and/or size groups" is approximately 5.1 percent: the standard error of 3,887,000 is 198,237 (5.1 percent of 3,887,000).

standard errors for a wide variety of estimates rather than the specific error for any statistic.

The relative standard errors and approximate standard errors of percentages that have been prepared for this report are applicable to estimates of discharges for patient characteristics (age, sex, color, and discharge status, and cross-classifications, e.g., age by sex) cross-classified by one of three hospital groupings as follows: (1) by region (e.g., Northeast) and/or size (e.g., 6-99 beds), (2) by type of ownership (e.g., government), or (3) by hospitals summed over all regions, size and ownership groups (all hospitals). The particular figure or table to which one refers to obtain a sampling error is contingent upon both the type of estimate (e.g., discharges) and the hospital grouping with which the patient characteristic(s) is cross-classified. The procedures that apply are as follows:

- Approximate relative standard errors of estimated number of discharges are obtained from the curves shown on figure II.
- Approximate relative standard errors of number of days of care are obtained from the curves shown in figure III.
- Approximate standard errors of estimated percentages of discharges when the characteristic(s) used to form the numerator of the percentage is a subclass of the denominator are shown in table IV.
- 4. Approximate standard errors of estimated percentages of days of care when the characteristic(s) used to form the numerator is a subclass of the denominator are shown in table V.

Figure III. Approximate relative standard errors of estimated number of days of care for patient characteristics, by geographic region and/or size of hospital, and type of ownership and for all hospitals.




Illustration of use of figure III: As shown in table 14, an estimated 2,559,000 days of care during 1971 were provided to patients aged 15-44 years in proprietary hospitals. The relative standard error of this estimate as read from the line "Ownership groups" is approximately 22.0 percent: the standard error is 562,980 (22.0 percent of 2,559,000).

Approximate standard errors of average lengths of stay can be calculated as in the following example:

Suppose the standard error $(\sigma_{R'})$ of the average length of stay during 1971 for males aged 15-44 years for all hospitals is desired. The estimated number of discharges for this statistic is 3,467,000 (table 1) and the estimated number of days of care is 23,866,000 (table 10).

Let
$$R' = \frac{Number\ of\ days\ of\ care}{Number\ of\ discharge}$$
$$= \frac{X'}{Y'} = \frac{23,866,000}{3.467,000} = 6.9\ days$$

The relative standard error $(V_{\rm X})$ of 23,866,000 (from all hospitals curve in figure III) is 2.9 percent, or .029 $V_{\rm X'}^2(.029)^2$. The relative standard error $(V_{\rm Y'})$ of 3.467,000 (from all hospitals curve in figure II) is 3.2 percent, or .032; $V_{\rm Y'}=(.032)^2$. The sample correlation coefficient (r) which measures the closeness of the relation between the estimated number of days of care and the estimated number of discharges has been computed to be 0.75.

$$\begin{split} {V_{\text{R}}}^2 &= {V_{\text{X}}}^2 + {V_{\text{Y}}}^2 - 2r V_{\text{X}}, \ V_{\text{Y}}, \\ &= \left(.029 \right)^2 + \left(.032 \right)^2 - 1.5 \left(.029 \times .032 \right) \\ &= .00084 + .00102 - .00139 = .00047 \\ V_{\text{R}}^2 &= \sqrt{.00047} = .022 \\ \sigma_{\text{B}}^2 &= R^2 \times V_{\text{B}}^2 = 6.9 \times .022 = 0.15 \ days \end{split}$$

APPENDIX II

DEFINITIONS OF CERTAIN TERMS USED IN THIS REPORT

Terms Relating to Hospitalization

Patient.—A person who is formally admitted to the inpatient service of a short-stay hospital for observation, care, diagnosis, or treatment. In this report the number of patients refers to the number of discharges during 1971 including any multiple discharges of the same individual from one short-stay hospital or more. All newborn infants, defined as those admitted by birth to the hospital, are excluded from this report. "Inpatient" and "patient" are used synonymously.

Patient under 1 year of age.—Includes infants admitted on the day of birth, directly or by transfer from another medical facility, with or without mention of a disease, disorder, or immaturity.

Discharge.—The formal release of an inpatient by a hospital, that is, the termination of a period of hospitalization by death or by disposition to place of residence, nursing home, or another hospital. In this report, "discharges" and "patients (or inpatients) discharged" are used synonymously.

Discharge rate. -- The ratio of the number of hospital discharges during a year to the number of persons in the civilian, noninstitutionalized population July 1 of that year.

Days of care.—The total number of inpatient days accumulated at time of discharge by patients discharged from short-stay hospitals during a year. A stay of less than 1 day (inpatient admission and discharge on the same day) is counted as 1 day in the summation of total days of care. For patients admitted and discharged on different days, the number of days of care is computed by counting all days from (and including) the date of admission to (but not including) the date of discharge.

Rate of days of care.—The ratio of the number of inpatient days accumulated at time of discharge by patients discharged from short-stay hospitals during a year to the number of persons in the civilian, non-institutionalized population July 1 of that year.

Average length of stay.—The total number of inpatient days accumulated at time of discharge by patients discharged during 1971 divided by the number of patients discharged. "Average stay," "duration of stay," and "length of stay" are used interchangeably.

Hospitals and Hospital Characteristics

Short-stay hospitals.—General and short-term special hospitals having six beds or more for inpatient use and an average (mean) length of stay of less than 30 days. Federal hospitals and hospital units of institutions are not included. "Hospitals" and "short-stay hospitals" are used synonymously.

Size of hospital.—Measured by the number of beds, cribs, and pediatric bassinets regularly maintained (set up and staffed for use) for inpatients; bassinets for newborn infants are not included. In this report the classification of hospitals by bed size is based on the number of beds at or near midyear reported by the hospitals.

Location of hospitals.—See "Geographic region."

Type of ownership of hospital.—Refers to the type of organization that controls and operates the hospital. The hospitals are grouped as follows:

Voluntary hospitals.—Hospitals operated by a church or another nonprofit organization.

Government hospitals. — Hospitals operated by State or local governments.

Proprietary hospitals.—Hospitals controlled by individuals, partnerships, or corporations for profit.

Demographic Terms

Age.-Refers to age at birthday prior to admission to the hospital inpatient service.

Color.—In this report patients are classified into two groups, "white" and "all other." The all other classification includes all categories other than white; some groups are too small to be presented separately for statistical purposes. White includes Mexican and Puerto Rican unless specifically identified as all other.

Geographic region.—In this report hospitals are classified by location according to the four geographic regions of the United States which correspond to those used by the U.S. Bureau of the Census.

Region States Included	South Delaware, Maryland, District of Columbia, Virginia, West Virginia, North Carolina,	
Northeast Maine, New Hampshire, Vermont Massachusetts, Rhode Island, Connecticut, New York, New Jersey and Pennsylvania North CentralMichigan, Ohio, Illinois, Indiana Wisconsin, Minnesota, Iowa, Missouri, North Dakota, South Dakota, Nebraska, and Kansas	South Carolina, Georgia, Florid Kentucky, Tennessee, Alabama, Mississippi, Arkansas, Louisiana, Oklahoma, and Texas	

VITAL AND HEALTH STATISTICS PUBLICATION SERIES

Originally Public Health Service Publication No. 1000

- Series 1. Programs and collection procedures.—Reports which describe the general programs of the National Center for Health Statistics and its offices and divisions, data collection methods used, definitions, and other material necessary for understanding the data.
- Series 2. Data evaluation and methods research.—Studies of new statistical methodology including: experimental tests of new survey methods, studies of vital statistics collection methods, new analytical techniques, objective evaluations of reliability of collected data, contributions to statistical theory.
- Series 3. Analytical studies.—Reports presenting analytical or interpretive studies based on vital and health statistics, carrying the analysis further than the expository types of reports in the other series.
- Series 4. Documents and committee reports.—Final reports of major committees concerned with vital and health statistics, and documents such as recommended model vital registration laws and revised birth and death certificates.
- Series 10. Data from the Health Interview Survey.—Statistics on illness, accidental injuries, disability, use of hospital, medical, dental, and other services, and other health-related topics, based on data collected in a continuing national household interview survey.
- Series 11. Data from the Health Examination Survey.—Data from direct examination, testing, and measurement of national samples of the civilian, noninstitutional population provide the basis for two types of reports: (1) estimates of the medically defined prevalence of specific diseases in the United States and the distributions of the population with respect to physical, physiological, and psychological characteristics; and (2) analysis of relationships among the various measurements without reference to an explicit finite universe of persons.
- Series 12. Data from the Institutional Population Surveys —Statistics relating to the health characteristics of persons in institutions, and their medical, nursing, and personal care received, based on national samples of establishments providing these services and samples of the residents or patients.
- Series 13. Palo from the Hospital Discharge Survey.—Statistics relating to discharged patients in short-stay hospitals, based on a sample of patient records in a national sample of hospitals.
- Series 14. Dala on health resources: manpower and facilities.—Statistics on the numbers, geographic distribution, and characteristics of health resources including physicians, dentists, nurses, other health occupations, hospitals, nursing homes, and outpatient facilities.
- Series 20. Data on mortality.—Various statistics on mortality other than as included in regular annual or monthly reports—special analyses by cause of death, age, and other demographic variables, also geographic and time series analyses.
- Scries 21. Data on natality, marriage, and divorce.—Various statistics on natality, marriage, and divorce other than as included in regular annual or monthly reports—special analyses by demographic variables, also geographic and time series analyses, studies of fertility.
- Series 22. Data from the National Natality and Mortality Surveys.—Statistics on characteristics of births and deaths not available from the vital records, based on sample surveys stemming from these records, including such topics as mortality by socioeconomic class, hospital experience in the last year of life, medical care during pregnancy, health insurance coverage, etc.

For a list of titles of reports published in these series, write to:

Office of Information
National Center for Health Statistics
Public Health Service, HRA
Rockville, Md. 20852

DHEW Publication No. (HRA) 75-1768 Series 13 - No. 17

U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE

Public Health Service

HEALTH RESOURCES ADMINISTRATION 5600 Fishers Lane Rockville, Md. 20852

OFFICIAL BUSINESS
Penalty for Private Use, \$300

POSTAGE AND FEES PAID U.S. DEPARTMENT OF H.E.W.

HEW 390

THIRD CLASS BLK, RATE

