

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
All causes											
All origins 3/	848.5	683.4	33.3	15.3	19.2	66.9	95.0	96.2	113.5	165.9	240.5
Male	846.4	749.8	37.0	16.7	22.8	93.7	141.1	135.6	151.2	213.7	304.8
Female	850.4	613.9	29.5	13.9	15.3	38.5	46.8	55.8	75.2	118.2	177.2
Hispanic	306.8	572.7	30.6	13.0	16.6	63.1	86.9	80.6	88.7	129.5	185.3
Male	332.9	624.4	33.8	14.9	18.6	92.0	128.2	115.2	121.1	175.5	249.0
Female	279.0	518.9	27.2	11.0	14.5	31.1	36.1	39.4	51.3	79.0	118.3
Mexican	224.4	521.5	30.7	14.0	16.1	34.1	86.8	46.8	80.6	119.0	172.3
Male	282.8	550.0	33.7	15.7	17.3	99.3	128.5	112.7	109.9	161.4	223.5
Female	230.8	490.5	27.6	12.1	14.9	30.4	34.0	34.7	43.9	69.8	112.7
Puerto Rican	398.1	*	27.8	11.9	21.1	33.0	104.7	52.0	151.4	224.1	318.3
Male	537.6	*	*	*	*	93.9	156.4	165.9	199.7	291.6	476.6
Female	390.5	*	*	*	*	*	*	57.9	105.7	157.6	183.7
Cuban	760.1	*	*	*	*	*	*	26.8	*	110.4	*
Male	861.9	*	*	*	*	*	*	*	*	*	*
Female	859.1	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	255.6	716.5	31.7	10.3	16.4	27.2	85.0	42.4	91.0	123.9	163.3
Male	325.4	*	35.3	12.1	20.1	71.4	123.6	115.1	131.9	176.0	231.8
Female	274.7	*	27.6	8.6	12.9	31.7	40.0	47.2	52.2	73.8	101.4

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
All causes											
Non-Hispanic 5/	926.2	703.5	33.8	15.8	19.6	67.3	96.2	99.2	117.5	170.6	245.8
Male	923.6	773.2	37.6	17.0	23.6	93.6	143.3	139.8	156.2	218.5	309.6
Female	928.6	630.4	29.9	14.4	15.4	39.7	48.7	58.8	79.1	123.4	183.5
Non-Hispanic white	991.1	555.4	30.1	14.5	18.0	63.4	85.4	87.2	104.4	153.1	220.9
Male	975.6	611.6	33.7	15.2	21.7	86.1	125.6	122.4	139.3	197.9	282.4
Female	1,006.1	496.4	26.3	13.7	14.1	39.6	44.3	51.4	69.0	108.2	159.7
Non-Hispanic black	798.1	1,354.4	49.5	21.4	26.6	88.7	156.7	174.9	207.1	294.6	438.1
Male	849.7	1,484.2	55.1	24.6	32.5	133.9	244.7	255.7	279.6	375.1	538.9
Female	751.2	1,217.8	43.6	18.1	20.5	42.0	72.2	101.4	142.2	223.2	349.0

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Septicemia (A40-A41)											
Non-Hispanic 5/	12.4	8.1	0.7	0.1	0.2	0.3	0.4	0.6	1.0	1.5	2.3
Male	11.2	8.8	0.9	*	*	0.3	0.5	0.6	0.9	1.6	2.6
Female	13.5	7.4	0.6	*	*	0.3	0.4	0.5	1.1	1.3	2.1
Non-Hispanic white	12.3	5.1	0.7	*	*	0.2	0.3	0.4	0.7	1.1	1.8
Male	11.0	5.8	0.9	*	*	*	*	0.5	0.6	1.2	2.1
Female	13.6	4.4	0.5	*	*	*	*	0.4	0.8	0.9	1.5
Non-Hispanic black	16.4	20.3	0.9	*	*	*	0.8	1.1	2.7	3.9	5.7
Male	15.4	21.4	*	*	*	*	*	*	2.6	3.9	5.9
Female	17.2	19.1	*	*	*	*	*	*	2.7	3.9	5.6

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Human immunodeficiency virus (HIV) disease (B20-B24)											
Non-Hispanic 5/	4.9	*	*	*	*	0.2	1.0	3.2	7.4	12.3	13.2
Male	7.5	*	*	*	*	*	0.8	3.8	10.3	18.4	19.6
Female	2.5	*	*	*	*	*	1.1	2.6	4.4	6.3	6.9
Non-Hispanic white	2.2	*	*	*	*	*	0.2	0.9	3.0	6.0	6.1
Male	3.8	*	*	*	*	*	*	1.3	4.9	10.2	10.5
Female	0.6	*	*	*	*	*	*	0.5	1.2	1.7	1.7
Non-Hispanic black	21.6	*	*	*	*	1.1	4.6	15.2	32.3	50.4	59.7
Male	30.8	*	*	*	*	*	3.7	17.6	43.5	71.6	84.0
Female	13.3	*	*	*	*	*	5.4	13.0	22.3	31.5	38.2

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malignant neoplasms (C00-C97)											
All origins 3/	194.4	1.6	2.7	2.4	2.5	3.6	4.9	7.3	12.6	24.6	48.7
Male	205.3	1.5	2.9	2.4	2.6	4.3	5.7	7.2	11.2	21.0	44.0
Female	183.9	1.8	2.5	2.4	2.3	2.9	4.1	7.4	14.0	28.1	53.5
Hispanic	60.5	*	2.8	2.4	2.7	4.5	4.1	5.7	8.9	18.3	34.4
Male	62.2	*	2.9	2.6	2.4	5.7	5.4	5.2	7.5	15.1	29.9
Female	58.7	*	2.7	2.1	3.0	3.2	2.7	6.4	10.4	21.8	39.2
Mexican	42.8	*	2.7	2.7	2.8	2.2	4.4	3.4	8.2	17.5	33.0
Male	49.7	*	2.5	2.9	2.5	5.8	5.8	4.5	7.0	13.9	27.0
Female	48.6	*	2.9	2.5	3.2	2.8	2.7	6.9	9.7	21.8	39.9
Puerto Rican	74.7	*	*	*	*	*	*	*	11.7	23.0	47.5
Male	99.7	*	*	*	*	*	*	*	*	*	*
Female	74.4	*	*	*	*	*	*	*	*	*	*
Cuban	163.4	*	*	*	*	*	*	*	*	*	*
Male	208.2	*	*	*	*	*	*	*	*	*	*
Female	161.4	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	54.4	*	*	*	*	2.7	3.1	3.1	9.4	18.2	30.6
Male	63.2	*	*	*	*	*	*	7.1	8.3	15.6	27.4
Female	64.5	*	*	*	*	*	*	*	10.3	20.7	33.5

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malignant neoplasms (C00-C97)											
Non-Hispanic 5/	213.9	1.6	2.7	2.4	2.4	3.4	5.1	7.7	13.3	25.4	50.3
Male	227.2	1.6	2.9	2.4	2.7	4.0	5.8	7.7	11.9	21.8	45.5
Female	201.2	1.6	2.5	2.5	2.2	2.8	4.4	7.6	14.6	29.0	55.0
Non-Hispanic white	230.6	1.8	2.9	2.5	2.3	3.4	5.1	7.4	12.8	24.5	47.7
Male	243.4	1.7	3.1	2.3	2.4	4.0	5.8	7.7	12.0	21.2	43.2
Female	218.4	1.8	2.8	2.6	2.2	2.8	4.4	7.2	13.7	27.7	52.2
Non-Hispanic black	172.8	*	2.0	2.4	2.9	3.7	5.3	9.6	17.0	33.7	72.2
Male	190.8	*	2.4	2.6	3.4	4.3	6.1	8.9	12.9	27.6	64.3
Female	156.5	*	*	2.2	2.3	3.0	4.5	10.3	20.8	39.2	79.2

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malignant neoplasm of stomach (C16)											
Non-Hispanic 5/	4.5	*	*	*	*	*	*	*	0.4	0.7	1.1
Male	5.4	*	*	*	*	*	*	*	0.5	0.8	1.2
Female	3.6	*	*	*	*	*	*	*	0.3	0.6	1.0
Non-Hispanic white	4.2	*	*	*	*	*	*	*	0.3	0.5	0.8
Male	5.0	*	*	*	*	*	*	*	0.3	0.6	1.0
Female	3.4	*	*	*	*	*	*	*	*	0.4	0.7
Non-Hispanic black	5.8	*	*	*	*	*	*	*	0.8	1.7	2.6
Male	6.9	*	*	*	*	*	*	*	*	1.9	2.9
Female	4.7	*	*	*	*	*	*	*	*	1.6	2.3

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malignant neoplasms of colon, rectum and anus (C18-C21)											
Non-Hispanic 5/	22.0	*	*	*	*	*	0.2	0.5	1.2	2.4	4.2
Male	22.3	*	*	*	*	*	*	0.4	1.3	2.7	4.4
Female	21.7	*	*	*	*	*	*	0.5	1.0	2.0	3.9
Non-Hispanic white	23.5	*	*	*	*	*	0.2	0.5	1.1	2.2	3.8
Male	23.9	*	*	*	*	*	*	0.4	1.3	2.6	4.0
Female	23.1	*	*	*	*	*	*	0.5	0.9	1.9	3.7
Non-Hispanic black	19.2	*	*	*	*	*	*	*	1.6	3.4	6.6
Male	19.0	*	*	*	*	*	*	*	1.9	4.0	7.6
Female	19.4	*	*	*	*	*	*	*	*	3.0	5.7

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malignant neoplasms of trachea, bronchus and lung (C33-C34)											
Non-Hispanic 5/	61.3	*	*	*	*	*	*	0.2	0.6	3.2	10.2
Male	72.6	*	*	*	*	*	*	*	0.6	3.1	11.3
Female	50.5	*	*	*	*	*	*	*	0.7	3.3	9.0
Non-Hispanic white	67.3	*	*	*	*	*	*	0.2	0.7	3.2	9.8
Male	78.4	*	*	*	*	*	*	*	0.6	3.0	10.5
Female	56.6	*	*	*	*	*	*	*	0.7	3.4	9.1
Non-Hispanic black	44.6	*	*	*	*	*	*	*	*	3.6	14.9
Male	59.2	*	*	*	*	*	*	*	*	4.0	18.8
Female	31.3	*	*	*	*	*	*	*	*	3.3	11.4

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malignant neoplasm of breast (C50)											
All origins 3/	14.7	*	*	*	*	*	*	0.5	1.8	4.3	8.2
Male	0.3	*	*	*	*	*	*	*	*	*	*
Female	28.6	*	*	*	*	*	*	0.9	3.6	8.6	16.2
Hispanic	4.8	*	*	*	*	*	*	*	1.1	3.5	5.7
Male	*	*	*	*	*	*	*	*	*	*	*
Female	9.7	*	*	*	*	*	*	*	2.5	7.3	11.5
Mexican	3.3	*	*	*	*	*	*	*	1.2	3.4	5.5
Male	*	*	*	*	*	*	*	*	*	*	*
Female	7.8	*	*	*	*	*	*	*	2.6	7.1	11.8
Puerto Rican	6.7	*	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*	*	*
Female	14.6	*	*	*	*	*	*	*	*	*	*
Cuban	10.9	*	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*	*	*
Female	24.8	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	4.6	*	*	*	*	*	*	*	*	3.7	4.3
Male	*	*	*	*	*	*	*	*	*	*	*
Female	10.7	*	*	*	*	*	*	*	*	7.3	8.2

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES,

1989), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malignant neoplasm of breast (C50)											
Non-Hispanic 5/	16.1	*	*	*	*	*	*	0.5	1.9	4.4	8.5
Male	0.3	*	*	*	*	*	*	*	*	*	*
Female	31.1	*	*	*	*	*	*	0.9	3.8	8.8	16.7
Non-Hispanic white	17.0	*	*	*	*	*	*	0.4	1.6	3.9	7.5
Male	0.3	*	*	*	*	*	*	*	*	*	*
Female	32.9	*	*	*	*	*	*	0.7	3.2	7.8	15.0
Non-Hispanic black	15.5	*	*	*	*	*	*	1.2	4.2	7.8	15.2
Male	0.4	*	*	*	*	*	*	*	*	*	*
Female	29.2	*	*	*	*	*	*	2.3	7.8	14.7	28.3

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malignant neoplasm of cervix uteri (C53)											
Non-Hispanic 5/	1.5	*	*	*	*	*	*	0.3	0.8	1.3	1.8
Female	2.9	*	*	*	*	*	*	0.5	1.5	2.5	3.5
Non-Hispanic white	1.4	*	*	*	*	*	*	0.2	0.7	1.2	1.6
Female	2.7	*	*	*	*	*	*	0.5	1.4	2.4	3.2
Non-Hispanic black	2.2	*	*	*	*	*	*	*	1.2	1.9	3.0
Female	4.2	*	*	*	*	*	*	*	2.2	3.6	5.6

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malignant neoplasm of ovary (C56)											
Non-Hispanic 5/	5.6	*	*	*	*	*	*	0.2	0.4	0.7	1.4
Female	10.9	*	*	*	*	*	*	0.4	0.7	1.4	2.8
Non-Hispanic white	6.2	*	*	*	*	*	*	0.2	0.3	0.7	1.5
Female	12.2	*	*	*	*	*	*	0.4	0.7	1.4	2.9
Non-Hispanic black	3.2	*	*	*	*	*	*	*	*	0.9	1.4
Female	6.1	*	*	*	*	*	*	*	*	1.7	2.7

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)											
Non-Hispanic 5/	21.6	0.6	0.9	0.7	0.9	1.4	2.0	2.3	2.7	3.5	4.8
Male	23.5	*	1.0	0.8	1.1	1.6	2.3	2.6	3.2	4.2	5.8
Female	19.7	*	0.9	0.6	0.7	1.1	1.7	2.0	2.3	2.8	3.7
Non-Hispanic white	23.8	*	1.1	0.7	0.9	1.4	2.0	2.1	2.6	3.4	4.5
Male	25.9	*	1.0	0.7	1.0	1.6	2.3	2.3	3.2	3.9	5.5
Female	21.8	*	1.1	0.6	0.8	1.1	1.7	1.8	2.1	2.9	3.5
Non-Hispanic black	14.5	*	*	0.8	1.1	1.5	2.2	3.3	3.7	4.7	6.8
Male	15.6	*	*	*	1.7	1.8	2.2	3.9	4.1	6.8	8.4
Female	13.5	*	*	*	*	*	2.2	2.8	3.3	2.9	5.4

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Leukemia (C91-C95)											
Non-Hispanic 5/	8.2	*	0.9	0.7	0.8	1.0	1.2	1.2	1.3	1.5	1.8
Male	9.3	*	0.9	0.7	0.9	1.1	1.4	1.4	1.5	1.7	2.0
Female	7.2	*	0.9	0.6	0.6	0.9	1.0	1.0	1.1	1.4	1.7
Non-Hispanic white	9.1	*	1.0	0.6	0.7	1.0	1.2	1.0	1.3	1.5	1.8
Male	10.3	*	1.0	0.6	0.8	1.1	1.4	1.2	1.5	1.6	1.9
Female	8.0	*	1.1	0.6	0.7	0.8	1.0	0.9	1.1	1.4	1.7
Non-Hispanic black	4.9	*	*	0.8	0.9	0.9	1.2	1.5	1.4	1.9	2.3
Male	5.4	*	*	*	1.5	*	*	1.8	1.8	2.4	2.7
Female	4.4	*	*	*	*	*	*	*	*	1.4	2.0

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)											
Non-Hispanic 5/	24.1	*	0.8	0.6	0.8	1.2	1.4	1.8	2.1	3.3	6.0
Male	24.9	*	1.0	0.7	0.8	1.4	1.9	2.2	2.5	3.6	6.6
Female	23.3	*	0.6	0.6	0.7	0.9	0.9	1.4	1.7	3.0	5.4
Non-Hispanic white	26.0	*	0.9	0.6	0.7	1.2	1.5	1.9	2.0	3.2	5.8
Male	26.8	*	1.1	0.7	0.8	1.4	2.0	2.3	2.4	3.5	6.4
Female	25.3	*	0.7	0.6	0.7	0.9	1.0	1.5	1.6	2.8	5.1
Non-Hispanic black	18.9	*	*	*	0.8	1.2	1.4	2.1	2.7	4.3	7.8
Male	20.0	*	*	*	*	1.5	2.1	2.6	3.0	4.4	8.3
Female	17.8	*	*	*	*	*	*	1.6	2.4	4.3	7.4

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Anemias (D50-D64)											
Non-Hispanic 5/	1.8	*	*	0.1	0.2	0.2	0.3	0.4	0.5	0.4	0.6
Male	1.4	*	*	*	*	0.3	0.4	0.5	0.6	0.4	0.5
Female	2.2	*	*	*	*	*	0.3	0.3	0.4	0.4	0.8
Non-Hispanic white	1.7	*	*	*	*	*	*	*	*	*	0.2
Male	1.3	*	*	*	*	*	*	*	*	*	*
Female	2.1	*	*	*	*	*	*	*	*	*	0.3
Non-Hispanic black	2.8	*	*	*	*	0.9	1.6	2.0	2.5	2.2	3.3
Male	2.6	*	*	*	*	*	1.7	2.6	3.4	2.3	2.8
Female	2.9	*	*	*	*	*	1.5	*	1.8	2.1	3.8

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Diabetes mellitus (E10-E14)											
Non-Hispanic 5/	26.4	*	*	*	0.2	0.2	0.6	1.3	2.0	3.5	5.5
Male	25.0	*	*	*	*	0.3	0.6	1.4	2.4	4.1	6.7
Female	27.8	*	*	*	*	*	0.5	1.1	1.6	2.8	4.4
Non-Hispanic white	26.0	*	*	*	*	0.2	0.4	0.9	1.7	2.9	4.8
Male	25.1	*	*	*	*	*	0.5	0.9	1.9	3.4	6.2
Female	26.9	*	*	*	*	*	*	0.9	1.5	2.4	3.5
Non-Hispanic black	34.1	*	*	*	*	*	1.4	3.3	4.4	7.2	11.0
Male	29.4	*	*	*	*	*	1.5	4.1	5.8	9.0	12.0
Female	38.4	*	*	*	*	*	*	2.6	3.0	5.6	10.1

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Major cardiovascular diseases (I00-I78)											
All origins 3/	323.9	14.9	1.9	0.7	1.1	2.2	4.3	7.1	13.0	25.0	48.8
Male	306.2	15.3	1.9	0.6	1.1	2.7	5.3	8.8	16.5	33.1	66.5
Female	340.9	14.5	1.8	0.8	1.1	1.7	3.2	5.5	9.5	17.0	31.5
Hispanic	96.2	12.0	1.5	0.7	0.9	2.1	2.8	4.4	8.6	14.1	26.7
Male	95.4	12.7	1.4	*	*	2.4	3.5	5.6	11.2	19.0	36.5
Female	97.1	11.2	1.6	*	*	1.7	1.8	3.0	5.6	8.7	16.5
Mexican	66.0	10.4	1.4	0.8	*	0.9	2.9	2.5	8.2	13.6	26.9
Male	76.6	11.2	*	*	*	2.1	3.7	5.2	10.8	17.7	35.2
Female	75.2	9.5	*	*	*	*	1.8	2.9	4.9	8.8	17.2
Puerto Rican	127.6	*	*	*	*	*	*	4.5	13.7	21.5	38.5
Male	157.2	*	*	*	*	*	*	*	*	*	*
Female	138.8	*	*	*	*	*	*	*	*	*	*
Cuban	331.1	*	*	*	*	*	*	*	*	*	*
Male	352.7	*	*	*	*	*	*	*	*	*	*
Female	397.4	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	76.9	*	*	*	*	*	*	1.8	8.1	12.0	21.2
Male	87.9	*	*	*	*	*	*	5.4	11.7	17.3	30.8
Female	92.6	*	*	*	*	*	*	*	*	7.0	12.5

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Major cardiovascular diseases (I00-I78)

Non-Hispanic 5/	356.7	15.5	1.9	0.7	1.1	2.2	4.5	7.7	13.8	26.6	51.3
Male	338.1	15.8	2.0	0.6	1.2	2.7	5.6	9.5	17.4	35.2	69.8
Female	374.3	15.2	1.8	0.8	1.1	1.7	3.4	6.0	10.1	18.2	33.1
Non-Hispanic white	385.8	12.1	1.5	0.6	1.0	1.9	3.5	6.2	11.3	22.5	43.3
Male	363.0	13.0	1.6	0.6	1.1	2.2	4.5	7.8	14.6	30.4	61.8
Female	407.6	11.0	1.5	0.7	0.9	1.5	2.5	4.6	8.0	14.6	24.8
Non-Hispanic black	289.7	30.1	3.7	1.0	1.6	4.1	9.8	17.1	29.5	53.3	108.4
Male	281.2	29.7	3.8	*	1.6	5.2	11.7	21.0	36.4	67.9	132.7
Female	297.4	30.6	3.6	1.3	1.5	2.9	7.9	13.6	23.3	40.4	86.9

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Diseases of heart (I00-I09,I11,I13,I20-I51)											
Non-Hispanic 5/	270.7	12.3	1.5	0.5	0.9	1.7	3.6	6.1	10.9	21.3	41.4
Male	268.0	12.0	1.7	0.4	0.9	2.2	4.6	7.6	14.4	29.3	59.2
Female	273.3	12.7	1.4	0.6	0.8	1.3	2.6	4.5	7.5	13.4	24.2
Non-Hispanic white	294.1	9.8	1.2	0.4	0.8	1.4	2.7	4.8	9.0	18.4	36.2
Male	289.5	10.3	1.3	0.4	0.8	1.7	3.5	6.3	12.1	26.0	53.9
Female	298.4	9.2	1.1	0.5	0.7	1.1	1.9	3.4	5.9	10.8	18.5
Non-Hispanic black	215.5	22.9	3.1	0.8	1.3	3.5	7.8	13.5	23.5	41.3	81.6
Male	215.7	20.7	3.4	*	1.4	4.5	10.0	16.7	30.3	54.1	104.2
Female	215.4	25.2	2.9	*	*	2.4	5.8	10.7	17.5	30.0	61.6

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Hypertensive heart disease (I11)											
Non-Hispanic 5/	9.4	*	*	*	*	*	*	0.4	1.0	1.6	3.4
Male	8.1	*	*	*	*	*	*	0.4	1.4	2.1	4.6
Female	10.7	*	*	*	*	*	*	0.3	0.6	1.1	2.1
Non-Hispanic white	8.6	*	*	*	*	*	*	0.2	0.6	1.0	2.1
Male	6.9	*	*	*	*	*	*	*	0.9	1.5	3.1
Female	10.2	*	*	*	*	*	*	*	*	0.6	1.0
Non-Hispanic black	16.5	*	*	*	*	*	*	1.1	3.3	5.2	11.8
Male	16.3	*	*	*	*	*	*	*	4.2	6.0	15.1
Female	16.7	*	*	*	*	*	*	*	2.4	4.6	8.9

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Ischemic heart diseases (I20-I25)											
Non-Hispanic 5/	193.6	0.7	*	*	*	0.2	0.7	1.6	4.4	10.8	24.9
Male	200.4	*	*	*	*	0.2	1.0	2.4	6.4	16.3	38.2
Female	187.2	*	*	*	*	*	0.4	0.8	2.3	5.3	11.9
Non-Hispanic white	212.6	*	*	*	*	*	0.6	1.4	4.1	10.4	23.6
Male	219.6	*	*	*	*	*	0.9	2.2	6.0	15.8	37.2
Female	205.8	*	*	*	*	*	*	0.6	2.1	4.9	10.1
Non-Hispanic black	141.2	*	*	*	*	*	1.5	2.9	6.6	15.4	37.5
Male	143.2	*	*	*	*	*	2.0	4.0	9.5	22.5	52.0
Female	139.4	*	*	*	*	*	*	1.8	4.0	9.2	24.7

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Acute myocardial infarction (I21-I22)											
Non-Hispanic 5/	71.4	*	*	*	*	*	0.3	0.6	1.9	4.6	10.6
Male	75.9	*	*	*	*	*	0.4	0.8	2.7	6.9	16.0
Female	67.1	*	*	*	*	*	*	0.4	1.1	2.3	5.3
Non-Hispanic white	78.4	*	*	*	*	*	0.2	0.5	1.9	4.5	10.3
Male	83.7	*	*	*	*	*	0.3	0.6	2.7	6.8	16.1
Female	73.4	*	*	*	*	*	*	0.4	1.1	2.2	4.5
Non-Hispanic black	51.6	*	*	*	*	*	*	1.1	2.6	6.0	14.6
Male	50.9	*	*	*	*	*	*	*	3.5	8.3	18.8
Female	52.3	*	*	*	*	*	*	*	1.8	3.9	10.9

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Other forms of chronic ischemic heart disease (I20,I25)											
Non-Hispanic 5/	120.9	*	*	*	*	*	0.4	1.0	2.4	6.1	14.0
Male	123.1	*	*	*	*	*	0.6	1.6	3.6	9.3	21.8
Female	118.8	*	*	*	*	*	*	0.4	1.1	2.9	6.5
Non-Hispanic white	132.7	*	*	*	*	*	0.3	0.9	2.1	5.7	13.1
Male	134.4	*	*	*	*	*	0.5	1.6	3.2	8.8	20.7
Female	131.1	*	*	*	*	*	*	*	1.0	2.6	5.5
Non-Hispanic black	88.2	*	*	*	*	*	0.8	1.8	3.9	9.2	22.2
Male	90.8	*	*	*	*	*	*	2.5	6.0	13.8	32.1
Female	86.0	*	*	*	*	*	*	*	2.0	5.0	13.4

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Atherosclerotic cardiovascular disease, so described (I25.0)											
Non-Hispanic 5/	26.0	*	*	*	*	*	*	0.2	0.7	1.9	4.9
Male	27.5	*	*	*	*	*	*	0.3	1.0	3.1	7.6
Female	24.6	*	*	*	*	*	*	*	0.4	0.8	2.2
Non-Hispanic white	26.8	*	*	*	*	*	*	0.2	0.7	1.8	4.4
Male	27.8	*	*	*	*	*	*	*	1.0	2.9	7.0
Female	25.8	*	*	*	*	*	*	*	0.3	0.7	1.8
Non-Hispanic black	28.5	*	*	*	*	*	*	*	1.0	3.2	8.5
Male	32.3	*	*	*	*	*	*	*	*	4.8	12.8
Female	25.0	*	*	*	*	*	*	*	*	1.7	4.7

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

Non-Hispanic 5/	94.9	*	*	*	*	*	0.3	0.8	1.6	4.1	9.2
Male	95.6	*	*	*	*	*	0.4	1.3	2.5	6.2	14.1
Female	94.2	*	*	*	*	*	*	0.3	0.8	2.1	4.3
Non-Hispanic white	106.0	*	*	*	*	*	0.2	0.7	1.5	3.9	8.7
Male	106.6	*	*	*	*	*	0.4	1.3	2.3	5.9	13.7
Female	105.3	*	*	*	*	*	*	*	0.7	1.9	3.7
Non-Hispanic black	59.7	*	*	*	*	*	*	1.3	2.9	6.0	13.7
Male	58.4	*	*	*	*	*	*	1.9	4.6	9.0	19.4
Female	60.9	*	*	*	*	*	*	*	*	3.4	8.7

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Other heart diseases (I26-I51)											
Non-Hispanic 5/	65.2	11.6	1.5	0.5	0.8	1.5	2.7	4.0	5.4	8.5	12.5
Male	57.8	11.0	1.6	0.4	0.8	1.9	3.3	4.7	6.4	10.6	15.6
Female	72.3	12.2	1.3	0.5	0.7	1.1	2.1	3.3	4.4	6.5	9.5
Non-Hispanic white	70.6	9.1	1.1	0.4	0.7	1.3	2.1	3.1	4.3	6.8	10.1
Male	61.4	9.4	1.2	0.3	0.8	1.5	2.6	3.7	5.0	8.5	13.2
Female	79.4	8.8	1.0	0.5	0.6	1.0	1.6	2.6	3.5	5.1	7.0
Non-Hispanic black	54.7	21.3	3.0	0.8	1.2	3.0	5.8	9.1	12.8	19.1	30.0
Male	53.5	18.8	3.3	*	1.3	3.9	7.3	11.1	15.5	24.1	34.6
Female	55.8	23.8	2.7	*	*	2.1	4.3	7.2	10.4	14.7	26.0

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

Non-Hispanic 5/	42.1	10.1	1.2	0.3	0.7	1.3	2.4	3.6	4.9	7.5	11.1
Male	39.6	9.3	1.4	0.3	0.7	1.7	3.1	4.2	5.8	9.4	14.1
Female	44.5	11.0	1.0	0.4	0.6	1.0	1.8	3.0	4.0	5.7	8.1
Non-Hispanic white	44.7	8.1	0.9	0.3	0.6	1.1	1.9	2.8	3.9	6.1	9.1
Male	41.3	8.1	1.0	*	0.7	1.3	2.4	3.4	4.5	7.7	12.0
Female	47.9	8.1	0.7	*	0.5	0.8	1.4	2.2	3.2	4.4	6.1
Non-Hispanic black	39.3	18.2	2.4	*	1.1	2.7	5.1	8.3	11.7	17.0	25.8
Male	40.3	16.0	2.7	*	1.2	3.5	6.6	9.7	14.1	21.1	30.8
Female	38.5	20.5	2.1	*	*	1.8	3.7	6.9	9.6	13.3	21.4

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Essential (primary) hypertension and hypertensive renal disease (I10,I12)											
Non-Hispanic 5/	7.4	*	*	*	*	*	*	0.2	0.4	0.5	1.1
Male	5.7	*	*	*	*	*	*	*	0.4	0.6	1.3
Female	9.0	*	*	*	*	*	*	*	0.4	0.4	0.8
Non-Hispanic white	7.0	*	*	*	*	*	*	*	0.2	0.3	0.5
Male	5.1	*	*	*	*	*	*	*	*	0.3	0.7
Female	8.8	*	*	*	*	*	*	*	*	*	0.3
Non-Hispanic black	11.3	*	*	*	*	*	*	*	1.4	1.9	4.5
Male	10.0	*	*	*	*	*	*	*	*	2.3	5.5
Female	12.4	*	*	*	*	*	*	*	1.4	1.5	3.5

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Cerebrovascular diseases (I60-I69)											
Non-Hispanic 5/	63.2	2.8	0.3	0.2	0.2	0.3	0.6	1.1	1.9	3.8	7.3
Male	49.7	3.3	0.3	*	*	0.3	0.6	1.1	1.9	3.9	7.3
Female	76.1	2.4	0.4	*	*	0.3	0.6	1.0	2.0	3.7	7.2
Non-Hispanic white	67.6	1.9	0.3	0.2	0.2	0.3	0.5	0.9	1.6	3.0	5.4
Male	52.0	2.2	*	*	*	0.3	0.6	0.8	1.7	2.9	5.6
Female	82.6	*	*	*	*	0.3	0.4	0.9	1.6	3.0	5.1
Non-Hispanic black	52.9	6.9	*	*	*	*	1.4	2.3	3.6	8.3	18.9
Male	46.2	8.3	*	*	*	*	*	2.8	3.4	8.9	18.2
Female	59.0	*	*	*	*	*	1.8	1.8	3.9	7.7	19.6

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Influenza and pneumonia (J10-J18)											
Non-Hispanic 5/	23.8	7.5	0.6	0.2	0.2	0.3	0.6	0.7	1.2	1.9	2.7
Male	21.5	8.2	0.6	0.2	0.2	0.3	0.8	0.7	1.4	2.1	3.1
Female	26.1	6.7	0.6	0.3	0.2	0.3	0.5	0.7	0.9	1.7	2.3
Non-Hispanic white	26.2	4.9	0.5	0.2	0.2	0.3	0.6	0.6	1.0	1.6	2.2
Male	23.1	5.8	0.6	*	*	*	0.7	0.6	1.2	1.7	2.5
Female	29.2	3.9	0.5	*	*	0.3	0.4	0.7	0.8	1.5	2.0
Non-Hispanic black	16.0	16.9	1.0	*	*	*	1.0	1.2	2.1	3.9	6.0
Male	16.3	16.6	*	*	*	*	*	*	2.8	4.7	7.1
Female	15.6	17.1	*	*	*	*	*	*	1.5	3.2	4.9

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Pneumonia (J12-J18)											
Non-Hispanic 5/	23.8	7.2	0.6	0.2	0.2	0.3	0.6	0.7	1.1	1.9	2.7
Male	21.4	7.8	0.6	*	0.2	0.3	0.7	0.7	1.3	2.1	3.0
Female	26.0	6.6	0.6	*	*	0.3	0.5	0.7	0.9	1.7	2.3
Non-Hispanic white	26.1	4.6	0.5	0.2	0.2	0.3	0.5	0.6	1.0	1.6	2.2
Male	23.0	5.3	0.6	*	*	*	0.7	0.5	1.1	1.7	2.5
Female	29.1	3.8	0.4	*	*	0.3	0.4	0.7	0.8	1.5	1.9
Non-Hispanic black	15.9	16.9	1.0	*	*	*	1.0	1.2	2.1	3.9	5.9
Male	16.3	16.6	*	*	*	*	*	*	2.7	4.7	7.1
Female	15.6	17.1	*	*	*	*	*	*	1.5	3.2	4.8

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Chronic lower respiratory diseases (J40-J47)

Non-Hispanic 5/	48.4	1.1	0.3	0.2	0.3	0.4	0.6	0.6	1.0	1.7	2.9
Male	48.0	*	0.4	0.3	0.4	0.4	0.7	0.6	1.0	1.7	2.7
Female	48.7	*	*	*	0.3	0.3	0.5	0.6	1.0	1.7	3.0
Non-Hispanic white	55.9	*	*	*	0.2	0.3	0.3	0.4	0.7	1.2	2.6
Male	54.6	*	*	*	*	*	0.4	0.4	0.6	1.3	2.4
Female	57.2	*	*	*	*	*	*	0.3	0.8	1.2	2.8
Non-Hispanic black	21.1	*	0.9	0.7	0.9	1.0	1.7	1.9	3.1	4.7	5.1
Male	24.5	*	*	*	*	1.4	1.9	2.2	3.4	4.4	5.1
Female	18.0	*	*	*	*	*	1.5	1.6	2.7	4.9	5.2

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Asthma (J45-J46)											
Non-Hispanic 5/	1.6	*	0.2	0.2	0.3	0.3	0.5	0.5	0.8	1.0	1.1
Male	1.1	*	*	0.3	0.3	0.4	0.5	0.6	0.7	0.9	1.0
Female	2.1	*	*	*	0.3	0.3	0.4	0.5	0.8	1.1	1.3
Non-Hispanic white	1.4	*	*	*	0.2	0.2	0.3	0.3	0.5	0.6	0.8
Male	0.9	*	*	*	*	*	*	0.3	0.3	0.5	0.6
Female	1.9	*	*	*	*	*	*	*	0.6	0.6	1.0
Non-Hispanic black	3.1	*	*	0.7	0.9	0.9	1.4	1.8	2.7	3.5	3.1
Male	2.7	*	*	*	*	*	1.6	2.0	3.0	3.1	2.9
Female	3.4	*	*	*	*	*	*	1.5	2.4	3.9	3.3

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Non-Hispanic 5/	10.0	7.8	0.6	0.2	0.2	0.3	0.4	0.6	0.6	1.1	1.6
Male	9.8	10.0	0.6	*	0.2	0.3	0.5	0.5	0.6	1.0	1.8
Female	10.2	5.5	0.6	*	0.2	0.3	0.4	0.7	0.6	1.1	1.4
Non-Hispanic white	11.0	4.8	0.5	0.2	0.2	0.3	0.4	0.5	0.5	0.8	1.4
Male	10.8	6.2	0.4	*	*	0.3	0.4	0.4	0.4	0.8	1.7
Female	11.3	3.4	0.5	*	*	*	0.3	0.6	0.5	0.8	1.1
Non-Hispanic black	6.9	19.0	1.1	*	*	*	0.8	1.2	1.4	2.6	3.3
Male	6.6	24.3	*	*	*	*	*	*	1.6	2.0	3.2
Female	7.2	13.4	*	*	*	*	*	*	*	3.0	3.4

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Chronic liver disease and cirrhosis (K70,K73-K74)

Non-Hispanic 5/	9.5	*	*	*	*	*	*	0.4	1.4	4.8	9.2
Male	12.3	*	*	*	*	*	*	0.5	1.8	5.9	12.1
Female	6.9	*	*	*	*	*	*	0.3	1.0	3.7	6.5
Non-Hispanic white	10.1	*	*	*	*	*	*	0.3	1.4	4.8	9.0
Male	13.0	*	*	*	*	*	*	0.4	1.8	5.8	11.7
Female	7.4	*	*	*	*	*	*	*	1.0	3.8	6.3
Non-Hispanic black	7.7	*	*	*	*	*	*	*	1.0	4.2	10.9
Male	10.7	*	*	*	*	*	*	*	*	5.8	15.1
Female	4.9	*	*	*	*	*	*	*	*	2.7	7.3

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Alcoholic liver disease (K70)											
Non-Hispanic 5/	4.2	*	*	*	*	*	*	0.2	1.0	3.2	5.8
Male	6.2	*	*	*	*	*	*	*	1.2	4.0	7.7
Female	2.4	*	*	*	*	*	*	*	0.7	2.4	3.9
Non-Hispanic white	4.4	*	*	*	*	*	*	*	1.0	3.2	5.7
Male	6.4	*	*	*	*	*	*	*	1.2	4.0	7.6
Female	2.4	*	*	*	*	*	*	*	0.7	2.5	3.9
Non-Hispanic black	3.7	*	*	*	*	*	*	*	*	2.6	6.1
Male	5.5	*	*	*	*	*	*	*	*	3.8	8.5
Female	2.2	*	*	*	*	*	*	*	*	1.6	4.1

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

ALL AGES 1/
 UNDER 1 YEAR 2/
 1-4 YEARS
 5-9 YEARS
 10-14 YEARS
 15-19 YEARS
 20-24 YEARS
 25-29 YEARS
 30-34 YEARS
 35-39 YEARS
 40-44 YEARS

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Non-Hispanic 5/	15.1	3.5	*	*	*	*	0.4	0.5	0.7	1.4	2.1
Male	14.8	3.7	*	*	*	*	0.3	0.5	0.7	1.5	2.2
Female	15.4	3.3	*	*	*	*	0.4	0.4	0.8	1.3	1.9
Non-Hispanic white	14.9	2.4	*	*	*	*	0.3	0.3	0.5	0.9	1.3
Male	14.8	2.5	*	*	*	*	*	*	0.4	1.2	1.5
Female	15.0	2.3	*	*	*	*	0.3	*	0.6	0.7	1.2
Non-Hispanic black	20.2	8.5	*	*	*	*	0.8	1.4	2.2	4.1	6.9
Male	18.6	8.9	*	*	*	*	*	1.8	2.6	3.4	7.2
Female	21.7	8.1	*	*	*	*	*	*	1.9	4.6	6.7

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Renal failure (N17-N19)											
Non-Hispanic 5/	14.9	3.3	*	*	*	*	0.3	0.4	0.7	1.3	2.0
Male	14.6	3.5	*	*	*	*	0.3	0.5	0.7	1.4	2.2
Female	15.2	3.1	*	*	*	*	0.3	0.4	0.7	1.3	1.8
Non-Hispanic white	14.6	2.2	*	*	*	*	0.2	0.3	0.4	0.9	1.3
Male	14.5	2.3	*	*	*	*	*	*	0.4	1.1	1.5
Female	14.7	2.1	*	*	*	*	*	*	0.5	0.7	1.1
Non-Hispanic black	20.0	8.5	*	*	*	*	0.8	1.3	2.2	3.9	6.8
Male	18.4	8.9	*	*	*	*	*	1.7	2.6	3.3	7.0
Female	21.4	8.1	*	*	*	*	*	*	1.8	4.5	6.6

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Pregnancy, childbirth and the puerperium (000-099)

All origins 3/	0.1	*	0.2	0.4	0.5	0.5	0.3	0.1
Female	0.3	*	0.4	0.8	1.0	0.9	0.7	0.3
Hispanic	0.2	*	*	*	0.6	*	*	*
Female	0.5	*	*	*	1.3	*	*	*
Mexican	0.2	*	*	*	*	*	*	*
Female	0.5	*	*	*	*	*	*	*
Puerto Rican	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*
Cuban	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*
Other Hispanic 4/	0.3	*	*	*	*	*	*	*
Female	0.6	*	*	*	*	*	*	*

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Pregnancy, childbirth and the puerperium (000-099)

Non-Hispanic 5/	0.1	*	0.2	0.3	0.5	0.5	0.3	0.1
Female	0.3	*	0.3	0.7	0.9	0.9	0.6	0.3
Non-Hispanic white	0.1	*	*	0.2	0.3	0.3	0.2	*
Female	0.2	*	*	0.4	0.5	0.7	0.3	*
Non-Hispanic black	0.4	*	*	1.1	1.4	1.1	1.0	*
Female	0.8	*	*	2.2	2.7	2.2	1.9	*

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Pregnancy with abortive outcome (000-007)

All origins 3/	0.0	*	*	*	*	*	*	*
Female	0.0	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*
Mexican	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*
Puerto Rican	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*
Cuban	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*
Other Hispanic 4/	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Pregnancy with abortive outcome (000-007)

Non-Hispanic 5/	0.0	*	*	*	*	*	*	*
Female	0.0	*	*	*	*	*	*	*
Non-Hispanic white	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*
Non-Hispanic black	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

ALL AGES 1/
 UNDER 1 YEAR 2/
 1-4 YEARS
 5-9 YEARS
 10-14 YEARS
 15-19 YEARS
 20-24 YEARS
 25-29 YEARS
 30-34 YEARS
 35-39 YEARS
 40-44 YEARS

Other complications of pregnancy, childbirth and the puerperium (010-099)

All origins 3/	0.1	*	0.2	0.3	0.5	0.4	0.3	0.1
Female	0.3	*	0.3	0.7	0.9	0.9	0.6	0.3
Hispanic	0.2	*	*	*	0.6	*	*	*
Female	0.5	*	*	*	1.3	*	*	*
Mexican	0.2	*	*	*	*	*	*	*
Female	0.5	*	*	*	*	*	*	*
Puerto Rican	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*
Cuban	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*
Other Hispanic 4/	0.3	*	*	*	*	*	*	*
Female	0.6	*	*	*	*	*	*	*

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other complications of pregnancy, childbirth and the puerperium (010-099)

Non-Hispanic 5/	0.1	*	0.1	0.3	0.4	0.4	0.3	0.1
Female	0.2	*	0.3	0.6	0.9	0.8	0.6	0.2
Non-Hispanic white	0.1	*	*	0.2	0.2	0.3	0.1	*
Female	0.1	*	*	0.3	0.5	0.6	0.3	*
Non-Hispanic black	0.4	*	*	1.0	1.3	1.0	0.9	*
Female	0.7	*	*	2.0	2.5	1.9	1.6	*

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

Non-Hispanic 5/	3.5	132.8	3.7	0.9	0.9	1.3	1.3	1.3	1.2	1.2	1.5
Male	3.7	139.4	3.6	1.0	1.0	1.6	1.5	1.3	1.4	1.3	1.6
Female	3.3	125.8	3.7	0.9	0.9	0.9	1.1	1.2	1.1	1.2	1.4
Non-Hispanic white	3.4	127.4	3.1	0.9	0.9	1.2	1.2	1.3	1.3	1.2	1.5
Male	3.6	135.5	3.1	1.0	1.0	1.5	1.3	1.5	1.5	1.3	1.6
Female	3.2	118.8	3.2	0.8	0.7	0.9	1.0	1.1	1.2	1.1	1.4
Non-Hispanic black	4.7	163.5	5.4	1.0	1.2	2.0	1.9	1.4	1.2	1.5	1.7
Male	5.1	167.5	5.5	*	*	2.7	2.3	*	*	*	2.2
Female	4.4	159.2	5.4	*	1.4	1.4	1.5	1.8	*	1.7	*

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Non-Hispanic 5/	12.1	92.6	1.8	0.4	0.5	1.7	2.9	3.4	4.1	6.2	7.8
Male	11.8	106.2	2.1	0.4	0.5	2.2	3.9	4.6	5.3	7.9	9.9
Female	12.3	78.3	1.5	0.3	0.4	1.1	1.9	2.2	2.9	4.6	5.7
Non-Hispanic white	12.4	73.1	1.5	0.3	0.4	1.7	2.8	3.3	3.9	6.0	7.7
Male	11.8	85.2	1.7	*	0.5	2.1	3.8	4.6	5.1	7.5	9.9
Female	13.1	60.4	1.3	*	*	1.2	1.9	1.9	2.8	4.5	5.4
Non-Hispanic black	12.9	176.9	3.0	0.6	0.7	1.7	3.8	4.8	5.5	8.5	10.1
Male	14.2	195.9	3.7	*	*	2.4	5.3	5.6	6.6	10.9	12.0
Female	11.7	156.8	2.2	*	*	*	2.4	4.1	4.5	6.4	8.4

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

All origins 3/	65.5	28.0	4.2	1.9	2.3	3.8	5.2	6.5	8.9	14.1	21.7
Male	53.8	31.4	4.6	1.8	2.5	4.4	6.0	7.3	9.9	16.8	25.7
Female	76.7	24.5	3.8	2.1	2.0	3.1	4.4	5.6	8.0	11.5	17.7
Hispanic	21.2	22.9	3.7	1.7	1.8	3.2	4.1	4.6	5.8	9.7	15.4
Male	20.2	25.4	4.3	1.6	2.0	3.9	4.2	5.0	7.1	12.6	20.0
Female	22.3	20.2	3.1	1.8	1.5	2.5	3.9	4.0	4.3	6.5	10.6
Mexican	15.3	22.5	3.6	1.8	1.7	1.6	3.8	2.5	5.5	8.3	14.6
Male	16.8	25.1	4.2	1.8	*	3.8	4.1	4.8	6.7	10.7	18.8
Female	18.5	19.8	3.0	1.9	1.8	2.2	3.4	3.5	3.9	5.6	9.6
Puerto Rican	28.0	*	*	*	*	*	*	*	10.1	18.9	26.1
Male	35.2	*	*	*	*	*	*	*	*	*	*
Female	29.8	*	*	*	*	*	*	*	*	*	*
Cuban	47.2	*	*	*	*	*	*	*	*	*	*
Male	42.3	*	*	*	*	*	*	*	*	*	*
Female	64.9	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	19.1	*	*	*	*	*	4.3	2.6	5.3	10.5	13.7
Male	21.8	*	*	*	*	*	*	5.6	*	15.1	17.3
Female	23.1	*	*	*	*	*	*	*	*	6.2	10.5

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Non-Hispanic 5/	71.8	29.0	4.3	2.0	2.3	3.9	5.4	6.9	9.5	14.8	22.3
Male	58.8	32.7	4.6	1.8	2.6	4.5	6.4	7.8	10.4	17.4	26.2
Female	84.2	25.2	4.0	2.1	2.1	3.2	4.5	6.0	8.6	12.1	18.5
Non-Hispanic white	77.3	23.2	3.9	2.0	2.2	3.6	5.0	6.0	8.2	13.0	19.6
Male	61.9	25.3	4.4	1.9	2.4	4.4	6.2	7.0	9.3	15.8	23.5
Female	92.1	21.0	3.4	2.1	2.0	2.8	3.8	5.0	7.2	10.2	15.8
Non-Hispanic black	61.1	53.8	5.6	2.0	3.0	5.4	8.0	12.0	17.3	27.4	42.3
Male	56.4	61.3	5.2	1.6	3.4	5.6	8.1	12.9	18.2	30.5	48.2
Female	65.2	46.0	6.0	2.5	2.7	5.2	8.0	11.2	16.5	24.6	37.2

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Accidents (unintentional injuries) (V01-X59,Y85-Y86)											
All origins 3/	35.7	24.2	11.2	6.4	7.4	32.8	39.5	31.4	28.5	34.1	36.7
Male	47.2	26.2	13.3	7.6	9.5	45.3	60.7	48.0	43.1	49.5	54.1
Female	24.5	22.1	8.9	5.0	5.3	19.5	17.2	14.3	13.8	18.7	19.6
Hispanic	25.8	18.8	10.3	5.0	5.7	27.8	37.0	30.2	26.1	31.6	33.1
Male	37.6	18.0	12.7	6.0	6.4	40.2	56.9	47.0	40.2	48.9	51.7
Female	13.2	19.7	7.8	4.0	5.0	14.0	12.6	10.2	10.0	12.6	13.5
Mexican	21.4	17.3	10.5	5.4	5.8	15.1	37.7	18.0	24.6	29.5	31.1
Male	36.2	15.4	12.7	6.2	6.4	43.2	58.0	47.3	37.4	45.5	47.3
Female	11.9	19.4	8.3	4.4	5.1	14.4	12.1	8.8	8.7	11.0	12.1
Puerto Rican	24.1	*	*	*	*	11.4	33.6	15.6	35.5	44.1	45.8
Male	43.2	*	*	*	*	*	49.8	*	54.5	70.5	*
Female	14.0	*	*	*	*	*	*	*	*	*	*
Cuban	22.5	*	*	*	*	*	*	*	*	*	*
Male	34.2	*	*	*	*	*	*	*	*	*	*
Female	16.7	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	24.3	*	10.2	4.9	5.6	12.2	37.5	16.4	28.1	33.4	33.3
Male	40.9	*	12.7	*	*	32.9	58.1	47.2	45.6	51.7	54.7
Female	16.1	*	*	*	*	13.1	13.5	14.6	11.4	15.8	13.9

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Non-Hispanic 5/	36.9	25.4	11.3	6.6	7.7	33.6	39.8	31.5	28.8	34.3	36.9
Male	48.5	28.0	13.4	7.9	10.0	46.1	61.4	48.1	43.4	49.3	54.0
Female	26.0	22.6	9.2	5.3	5.3	20.5	18.1	15.1	14.4	19.6	20.2
Non-Hispanic white	38.5	20.9	10.9	5.9	7.3	36.0	42.2	32.9	29.6	34.9	36.8
Male	49.7	22.9	13.2	6.7	9.5	48.8	65.0	49.7	44.2	49.7	53.7
Female	27.8	18.7	8.5	5.0	5.1	22.5	18.9	15.8	14.7	20.0	19.9
Non-Hispanic black	34.4	44.5	14.2	9.8	9.7	25.3	34.2	31.3	30.9	36.5	45.3
Male	49.5	50.4	15.8	13.0	13.2	37.0	52.9	49.5	48.0	54.3	68.4
Female	20.6	38.3	12.4	6.4	6.0	13.3	16.3	14.6	15.5	20.7	24.9

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Transport accidents (V01-V99,Y85)											
All origins 3/	16.6	3.7	4.4	3.8	5.0	26.6	29.2	20.9	16.9	17.4	17.3
Male	23.5	3.3	5.0	4.3	6.2	35.4	43.9	31.4	25.4	24.7	25.4
Female	10.0	4.2	3.8	3.3	3.8	17.3	13.8	10.1	8.3	10.2	9.4
Hispanic	15.6	5.3	5.3	3.3	4.3	22.7	29.0	21.6	15.9	17.3	17.3
Male	22.8	*	6.0	3.8	4.9	31.7	44.1	33.2	24.1	25.8	26.0
Female	8.0	6.7	4.6	2.7	3.7	12.7	10.5	7.8	6.4	7.9	8.2
Mexican	13.7	5.0	5.7	3.4	4.3	12.5	30.4	13.4	16.4	17.1	17.5
Male	23.3	*	6.5	3.7	5.0	34.5	46.1	34.8	24.3	25.1	25.5
Female	7.5	*	4.8	3.0	3.6	13.1	10.6	6.9	6.4	7.7	8.2
Puerto Rican	9.7	*	*	*	*	8.8	21.7	7.3	14.0	13.5	*
Male	16.6	*	*	*	*	*	*	*	*	*	*
Female	6.3	*	*	*	*	*	*	*	*	*	*
Cuban	12.2	*	*	*	*	*	*	*	*	*	*
Male	19.9	*	*	*	*	*	*	*	*	*	*
Female	7.7	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	14.6	*	5.0	4.0	4.8	9.6	28.6	11.6	15.5	19.4	18.8
Male	24.1	*	*	*	*	24.9	44.6	31.9	24.8	29.6	30.2
Female	10.2	*	*	*	*	11.8	9.9	12.4	6.6	9.7	8.5

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY

BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Transport accidents (V01-V99,Y85)											
Non-Hispanic 5/	16.7	3.4	4.2	3.9	5.2	27.2	29.2	20.7	17.1	17.4	17.2
Male	23.5	3.2	4.7	4.4	6.5	36.0	43.7	30.9	25.6	24.4	25.2
Female	10.2	3.6	3.6	3.4	3.8	18.0	14.4	10.5	8.6	10.5	9.5
Non-Hispanic white	17.0	3.1	3.8	3.6	5.2	29.6	30.3	21.3	17.1	17.5	17.3
Male	23.8	2.9	4.4	4.0	6.4	38.6	45.5	31.5	25.4	24.2	25.0
Female	10.6	3.3	3.2	3.2	3.8	20.0	14.8	10.9	8.7	10.8	9.6
Non-Hispanic black	16.3	*	5.4	5.5	5.4	19.3	26.7	21.6	19.6	18.5	19.8
Male	24.5	*	5.8	6.9	7.3	27.2	40.5	34.1	31.1	27.9	31.5
Female	8.9	*	5.0	4.1	3.4	11.1	13.5	10.1	9.2	10.2	9.5

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)											
All origins 3/	15.4	3.6	4.1	3.6	4.6	25.7	28.0	19.6	15.6	15.9	15.5
Male	21.5	3.2	4.5	4.0	5.7	34.0	42.1	29.5	23.1	22.3	22.5
Female	9.5	4.0	3.6	3.0	3.5	16.9	13.4	9.5	8.0	9.6	8.7
Hispanic	14.3	4.9	4.7	3.0	4.0	22.1	27.7	20.2	14.5	15.4	15.2
Male	21.0	*	5.3	3.6	4.5	30.8	42.1	31.1	22.2	22.9	23.4
Female	7.2	6.2	4.1	2.5	3.5	12.5	10.1	7.1	5.8	7.1	6.6
Mexican	13.0	5.0	5.3	3.2	4.2	12.3	29.2	12.5	15.7	16.1	16.1
Male	21.9	*	6.1	3.6	4.8	33.8	43.9	32.5	23.3	23.5	23.4
Female	7.3	*	4.6	2.8	3.5	13.1	10.6	6.6	6.2	7.5	7.6
Puerto Rican	8.9	*	*	*	*	8.3	20.9	7.2	13.3	12.4	*
Male	15.2	*	*	*	*	*	*	*	*	*	*
Female	5.8	*	*	*	*	*	*	*	*	*	*
Cuban	11.8	*	*	*	*	*	*	*	*	*	*
Male	19.2	*	*	*	*	*	*	*	*	*	*
Female	7.4	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	11.8	*	*	3.3	4.1	9.1	26.7	10.5	11.9	14.9	14.5
Male	20.3	*	*	*	*	23.4	42.6	29.4	19.5	22.9	25.5
Female	7.3	*	*	*	*	11.1	8.1	10.5	*	7.3	*

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)											
Non-Hispanic 5/	15.5	3.2	3.9	3.7	4.7	26.3	28.0	19.4	15.7	16.0	15.5
Male	21.5	3.1	4.3	4.1	5.9	34.5	41.9	29.0	23.2	22.1	22.3
Female	9.8	3.4	3.5	3.2	3.5	17.6	14.0	9.9	8.3	9.9	8.9
Non-Hispanic white	15.8	3.0	3.5	3.4	4.7	28.5	29.0	19.9	15.7	16.0	15.5
Male	21.7	2.9	3.9	3.7	5.9	36.9	43.5	29.3	22.9	21.8	22.0
Female	10.1	3.1	3.1	3.0	3.5	19.6	14.2	10.3	8.4	10.2	8.9
Non-Hispanic black	15.2	*	5.3	5.1	4.9	18.6	26.1	20.7	18.5	17.2	18.1
Male	22.7	*	5.6	6.5	6.5	26.1	39.4	32.7	29.3	25.6	28.2
Female	8.5	*	4.9	3.7	3.2	10.8	13.4	9.7	8.8	9.7	9.1

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN,

RACE FOR NON-HISPANIC POPULATION, AND SEX	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Nontransport accidents (W00-X59,Y86)											
Non-Hispanic 5/	20.2	22.0	7.2	2.7	2.6	6.4	10.7	10.9	11.8	16.9	19.7
Male	25.0	24.9	8.7	3.5	3.6	10.1	17.7	17.2	17.8	24.8	28.8
Female	15.7	19.1	5.6	1.9	1.5	2.5	3.6	4.6	5.8	9.1	10.7
Non-Hispanic white	21.5	17.8	7.1	2.3	2.2	6.5	11.9	11.6	12.5	17.4	19.5
Male	25.9	20.0	8.8	2.8	3.0	10.2	19.6	18.2	18.9	25.6	28.7
Female	17.2	15.5	5.3	1.8	1.2	2.5	4.1	4.9	6.0	9.2	10.3
Non-Hispanic black	18.1	41.6	8.7	4.3	4.3	6.1	7.5	9.7	11.3	18.0	25.5
Male	25.0	48.2	10.0	6.2	5.9	9.8	12.4	15.4	16.9	26.4	36.9
Female	11.8	34.6	7.4	2.3	2.6	2.2	2.8	4.5	6.3	10.5	15.4

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Accidental drowning and submersion (W65-W74)											
Non-Hispanic 5/	1.1	1.6	3.1	0.8	0.9	1.6	1.3	1.1	0.7	1.0	1.1
Male	1.8	1.3	3.8	1.1	1.2	2.9	2.4	2.0	1.3	1.6	1.8
Female	0.5	1.9	2.4	0.5	0.5	0.3	0.3	0.3	*	0.3	0.4
Non-Hispanic white	1.1	1.5	3.5	0.6	0.5	1.3	1.3	1.0	0.7	0.9	1.0
Male	1.6	*	4.3	0.6	0.7	2.2	2.2	1.7	1.1	1.4	1.7
Female	0.5	1.7	2.6	0.5	*	*	0.4	*	*	0.4	0.4
Non-Hispanic black	1.6	*	2.0	1.7	2.4	3.0	1.6	1.3	1.1	1.2	1.3
Male	2.8	*	2.4	2.8	3.4	5.4	3.2	2.8	2.2	2.2	2.2
Female	0.5	*	*	*	1.4	*	*	*	*	*	*

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Accidental exposure to smoke, fire and flames (X00-X09)											
Non-Hispanic 5/	1.3	1.3	1.7	0.9	0.5	0.4	0.7	0.7	0.7	0.9	1.1
Male	1.6	1.5	2.0	1.1	0.5	0.5	1.0	0.8	0.9	1.3	1.6
Female	1.0	*	1.3	0.6	0.4	0.4	0.4	0.5	0.5	0.6	0.7
Non-Hispanic white	1.1	1.1	1.3	0.7	0.4	0.4	0.7	0.5	0.7	0.8	1.0
Male	1.4	*	1.7	0.9	0.4	0.5	1.0	0.8	0.9	1.1	1.5
Female	0.9	*	0.9	0.5	0.4	0.3	0.3	*	0.5	0.5	0.6
Non-Hispanic black	2.3	*	3.4	1.8	0.8	*	1.0	1.4	1.0	1.6	2.1
Male	3.0	*	3.7	2.3	*	*	*	*	*	2.6	3.0
Female	1.8	*	3.1	1.3	*	*	*	1.5	*	*	*

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Accidental poisoning and exposure to noxious substances (X40-X49)											
Non-Hispanic 5/	5.1	*	0.2	*	0.1	2.1	5.3	6.0	7.1	10.6	12.2
Male	7.2	*	*	*	*	3.1	8.5	9.1	10.1	14.7	17.0
Female	3.1	*	*	*	*	1.0	2.1	2.9	4.1	6.5	7.6
Non-Hispanic white	5.2	*	*	*	*	2.4	6.2	6.9	7.7	11.0	12.3
Male	7.3	*	*	*	*	3.8	9.9	10.3	11.2	15.3	17.2
Female	3.1	*	*	*	*	1.0	2.5	3.4	4.2	6.7	7.4
Non-Hispanic black	6.0	*	*	*	*	0.9	2.4	4.1	6.4	11.2	15.6
Male	8.6	*	*	*	*	*	3.9	6.8	8.4	15.5	20.9
Female	3.6	*	*	*	*	*	*	1.7	4.7	7.4	10.9

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

Non-Hispanic 5/	6.7	18.3	1.9	0.6	0.7	1.3	2.0	1.9	2.2	2.9	3.3
Male	7.7	21.1	2.1	0.8	1.1	1.9	3.4	3.2	3.8	4.8	5.3
Female	5.8	15.4	1.6	0.4	0.3	0.6	0.6	0.6	0.6	1.2	1.3
Non-Hispanic white	7.2	14.5	1.7	0.7	0.7	1.4	2.2	2.0	2.2	3.1	3.2
Male	8.1	16.8	2.0	0.9	1.1	2.1	3.7	3.3	3.8	5.1	5.2
Female	6.4	12.1	1.5	0.5	0.3	0.7	0.7	0.6	0.7	1.1	1.2
Non-Hispanic black	5.5	35.5	2.6	*	0.8	0.8	1.4	1.9	2.0	2.6	4.0
Male	7.0	41.8	2.9	*	*	*	2.1	2.9	3.7	3.8	6.7
Female	4.2	28.9	2.3	*	*	*	*	*	*	1.6	1.6

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) 6/											
Non-Hispanic 5/	11.6	*	1.3	8.4	13.0	13.8	14.1	15.3	16.1
Male	19.0	*	2.0	13.8	22.7	23.3	22.9	24.2	24.9
Female	4.5	*	0.6	2.7	3.3	4.4	5.3	6.5	7.5
Non-Hispanic white	13.0	*	1.3	9.3	13.9	15.1	15.7	17.2	18.2
Male	21.4	*	2.1	15.4	24.1	25.3	25.2	26.8	27.9
Female	5.0	*	0.6	3.0	3.6	4.8	6.1	7.6	8.5
Non-Hispanic black	5.3	*	1.2	4.5	10.5	9.7	8.7	8.1	6.9
Male	9.3	*	1.7	7.5	20.0	17.7	15.5	14.2	10.8
Female	1.7	*	*	*	*	2.4	2.5	2.6	3.4

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0) /6

Non-Hispanic 5/	5.1	*	0.9	3.9	5.8	6.4	7.1	8.3	8.5
Male	7.5	*	1.3	6.0	9.5	10.2	10.9	12.4	12.0
Female	2.9	*	0.4	1.7	2.2	2.7	3.4	4.2	5.1
Non-Hispanic white	5.6	*	0.8	4.4	6.5	7.0	7.8	9.2	9.6
Male	8.2	*	1.2	6.8	10.5	11.0	11.9	13.6	13.4
Female	3.2	*	0.4	1.7	2.4	2.9	3.8	4.8	5.8
Non-Hispanic black	2.3	*	0.8	1.5	3.1	3.8	4.5	4.3	3.6
Male	3.7	*	1.2	2.3	5.6	6.8	7.4	7.0	5.2
Female	1.1	*	*	*	*	*	1.9	1.9	2.2

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) /6											
All origins 3/	7.1	8.2	2.7	0.7	0.9	9.4	17.3	14.9	11.5	10.4	8.6
Male	11.1	9.5	3.0	0.7	1.1	15.7	28.9	24.1	17.7	15.3	12.6
Female	3.3	6.9	2.4	0.6	0.7	2.7	5.1	5.5	5.2	5.5	4.7
Hispanic	9.0	6.7	2.3	0.5	1.1	13.8	22.3	16.1	11.7	11.0	9.6
Male	14.5	7.9	2.5	*	1.5	23.9	36.1	25.3	17.6	16.5	15.2
Female	3.2	5.5	2.1	*	*	2.6	5.3	5.2	4.9	5.0	3.7
Mexican	7.5	5.2	2.3	*	1.1	8.0	21.6	9.5	10.2	9.8	9.0
Male	14.1	*	2.3	*	*	27.5	35.0	25.9	15.7	15.1	14.1
Female	2.5	*	2.2	*	*	2.6	4.7	3.4	3.2	3.6	*
Puerto Rican	11.4	*	*	*	*	8.5	34.8	11.9	21.5	17.9	11.1
Male	20.8	*	*	*	*	*	58.3	*	*	*	*
Female	6.2	*	*	*	*	*	*	*	*	*	*
Cuban	6.0	*	*	*	*	*	*	*	*	*	*
Male	9.9	*	*	*	*	*	*	*	*	*	*
Female	3.6	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	7.7	*	*	*	*	4.0	21.3	7.6	12.3	12.0	10.3
Male	14.1	*	*	*	*	12.4	34.3	21.2	18.4	17.9	17.3
Female	4.0	*	*	*	*	*	*	*	6.6	6.4	*

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) /6											
Non-Hispanic 5/	6.7	8.4	2.8	0.7	0.9	8.5	16.0	14.4	11.3	10.1	8.4
Male	10.4	9.8	3.2	0.7	1.0	14.0	26.8	23.4	17.5	14.9	12.1
Female	3.2	6.9	2.4	0.7	0.7	2.7	5.0	5.4	5.2	5.5	4.7
Non-Hispanic white	4.0	5.8	1.7	0.7	0.6	2.9	5.5	6.5	6.6	7.0	6.2
Male	5.6	6.8	2.0	0.6	0.6	3.9	7.5	9.3	9.4	10.3	8.7
Female	2.4	4.7	1.5	0.7	0.6	1.9	3.5	3.6	3.8	3.7	3.7
Non-Hispanic black	22.7	19.5	7.1	0.9	1.9	33.6	64.9	54.6	36.5	29.0	22.0
Male	39.3	21.7	8.4	*	2.8	59.7	120.3	99.7	64.5	44.2	34.7
Female	7.6	17.1	5.7	*	*	6.6	11.8	13.6	11.4	15.6	10.7

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)											
Non-Hispanic 5/	3.7	*	0.3	0.3	0.5	6.7	12.7	9.8	6.5	5.0	3.4
Male	6.3	*	0.3	0.4	0.8	11.7	22.6	17.1	10.8	7.8	5.2
Female	1.2	*	0.3	0.3	0.3	1.4	2.6	2.6	2.1	2.3	1.7
Non-Hispanic white	1.6	*	0.2	0.3	0.4	1.8	3.2	2.8	2.6	2.7	2.1
Male	2.2	*	*	0.4	0.5	2.6	4.7	4.3	3.6	3.9	2.9
Female	0.9	*	*	*	*	0.9	1.6	1.4	1.6	1.5	1.3
Non-Hispanic black	16.3	*	*	*	1.4	29.5	57.6	45.3	27.9	19.1	11.8
Male	30.6	*	*	*	2.2	54.2	110.3	86.3	53.2	32.6	20.7
Female	3.2	*	*	*	*	4.0	7.2	8.0	5.2	7.0	4.0

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02.X85-X92,X96-Y09,Y87.1) /6

Non-Hispanic 5/	3.1	8.2	2.4	0.4	0.3	1.8	3.3	4.6	4.9	5.1	4.9
Male	4.2	9.5	2.8	0.4	0.2	2.2	4.2	6.3	6.7	7.1	6.9
Female	2.0	6.8	2.0	0.4	0.4	1.3	2.4	2.8	3.1	3.2	3.0
Non-Hispanic white	2.4	5.6	1.5	0.3	0.3	1.2	2.4	3.6	4.0	4.3	4.1
Male	3.4	6.7	1.8	*	*	1.3	2.8	5.0	5.7	6.4	5.8
Female	1.5	4.5	1.2	0.4	0.4	1.0	1.9	2.2	2.3	2.2	2.4
Non-Hispanic black	6.4	19.1	6.3	*	*	4.1	7.3	9.3	8.6	10.0	10.2
Male	8.7	21.4	7.7	*	*	5.6	10.0	13.4	11.3	11.6	14.1
Female	4.4	16.8	5.0	*	*	2.6	4.6	5.6	6.2	8.6	6.7

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Non-Hispanic 5/	1.6	2.4	0.3	*	0.2	0.7	1.6	1.8	2.2	3.2	3.3
Male	2.1	2.7	*	*	0.3	1.0	2.5	2.7	2.9	4.2	4.5
Female	1.0	2.0	0.4	*	*	0.3	0.7	1.0	1.5	2.3	2.2
Non-Hispanic white	1.6	1.7	0.3	*	0.2	0.8	1.8	2.0	2.4	3.3	3.4
Male	2.1	2.0	*	*	0.3	1.2	2.6	2.8	3.1	4.1	4.4
Female	1.1	*	*	*	*	0.4	0.9	1.1	1.6	2.5	2.4
Non-Hispanic black	1.8	5.9	*	*	*	*	1.1	1.5	2.0	3.9	4.0
Male	2.8	6.4	*	*	*	*	2.0	2.5	2.8	6.0	6.6
Female	0.9	*	*	*	*	*	*	*	*	2.1	1.8

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

Non-Hispanic 5/	1.5	2.4	0.3	*	0.2	0.6	1.4	1.6	2.1	3.1	3.2
Male	2.0	2.7	*	*	0.2	0.8	2.0	2.3	2.8	4.0	4.4
Female	1.0	2.0	0.4	*	*	0.3	0.7	0.9	1.5	2.2	2.1
Non-Hispanic white	1.5	1.7	0.3	*	0.2	0.7	1.6	1.8	2.3	3.1	3.3
Male	2.0	2.0	*	*	*	1.0	2.2	2.5	3.0	3.9	4.3
Female	1.1	*	*	*	*	0.4	0.9	1.1	1.6	2.3	2.3
Non-Hispanic black	1.7	5.9	*	*	*	*	0.8	1.3	2.0	3.9	4.0
Male	2.6	6.4	*	*	*	*	1.5	2.3	2.7	6.0	6.5
Female	0.9	*	*	*	*	*	*	*	*	2.1	1.7

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85+ YEARS

All causes

All origins 3/	354.8	512.4	771.8	1,210.7	1,869.7	2,878.3	4,494.0	7,151.9	15,112.8
Male	455.4	646.8	958.4	1,496.8	2,315.6	3,587.4	5,603.2	8,781.9	16,744.8
Female	257.2	383.7	596.9	951.3	1,485.9	2,314.8	3,717.6	6,175.1	14,429.9
Hispanic	280.9	403.9	580.6	910.1	1,412.8	2,229.5	3,556.3	5,501.5	11,821.8
Male	379.7	529.6	751.2	1,158.6	1,817.5	2,791.5	4,456.8	6,783.5	12,888.3
Female	181.7	283.8	426.0	695.1	1,085.0	1,800.8	2,917.0	4,707.4	11,300.5
Mexican	259.1	370.7	548.7	931.7	1,448.8	2,394.4	3,682.8	5,811.4	*
Male	350.1	455.5	673.3	1,234.4	1,715.4	2,957.1	*	*	*
Female	166.1	278.8	423.1	691.4	1,204.0	1,961.5	3,051.4	*	*
Puerto Rican	474.3	563.4	1,024.6	1,074.2	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	291.4	341.8	*	*	*	*	*	*	*
Cuban	255.0	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	261.5	411.0	482.1	814.9	1,364.6	2,123.5	*	*	*
Male	353.1	577.7	631.2	*	*	*	*	*	*
Female	173.6	273.7	359.7	*	*	*	*	*	*

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	--------------

All causes

Non-Hispanic 5/	360.4	519.3	783.7	1,228.7	1,896.2	2,908.7	4,528.0	7,201.4	15,199.0
Male	460.5	653.3	969.6	1,514.1	2,341.5	3,621.9	5,644.0	8,842.8	16,866.3
Female	263.6	391.0	608.9	968.9	1,511.2	2,340.5	3,747.4	6,219.1	14,505.6
Non-Hispanic white	324.7	474.0	730.8	1,169.7	1,827.4	2,849.2	4,476.8	7,196.1	15,371.2
Male	417.3	596.5	901.6	1,439.8	2,256.8	3,543.1	5,580.2	8,828.0	17,055.3
Female	233.1	353.9	567.1	919.2	1,448.1	2,285.2	3,695.5	6,211.0	14,670.6
Non-Hispanic black	656.8	946.1	1,351.9	1,920.5	2,777.9	3,951.1	5,712.5	8,098.0	14,660.8
Male	846.0	1,222.7	1,746.2	2,456.8	3,547.9	5,105.6	7,313.8	10,346.7	16,576.2
Female	493.1	710.9	1,028.9	1,498.8	2,209.4	3,182.0	4,756.8	6,952.7	13,951.4

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS.]

FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	--------------

Septicemia (A40-A41)

Non-Hispanic 5/	4.1	6.3	10.1	15.6	25.6	41.3	67.8	104.7	206.8
Male	4.3	6.7	10.9	16.6	28.0	47.1	78.8	123.1	226.8
Female	3.8	6.0	9.3	14.8	23.5	36.6	60.2	93.6	198.5
Non-Hispanic white	3.1	5.0	8.2	13.2	22.3	36.9	62.4	99.1	195.6
Male	3.3	5.2	8.6	14.2	24.2	41.6	72.4	116.6	213.6
Female	2.9	4.7	7.9	12.3	20.6	33.1	55.2	88.6	188.1
Non-Hispanic black	10.9	17.1	27.6	38.9	58.2	90.9	141.2	194.2	368.6
Male	12.0	19.1	34.0	41.8	67.5	112.6	173.0	237.9	430.8
Female	10.0	15.5	22.4	36.6	51.4	76.4	122.2	172.0	345.6

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	--------------

Viral hepatitis (B15-B19)

Non-Hispanic 5/	5.0	5.0	3.5	3.4	4.0	4.4	4.7	4.4	3.6
Male	7.7	7.8	5.0	4.0	4.6	4.7	4.8	5.3	4.3
Female	2.4	2.3	2.2	2.8	3.4	4.3	4.6	3.9	3.3
Non-Hispanic white	4.8	4.4	3.0	2.9	3.4	3.9	4.4	4.2	3.3
Male	7.4	7.0	4.3	3.6	4.3	4.4	4.9	5.4	4.4
Female	2.2	1.8	1.7	2.3	2.7	3.4	4.1	3.5	2.9
Non-Hispanic black	6.8	9.6	7.3	6.2	5.1	6.2	6.2	*	*
Male	10.1	14.3	10.1	7.5	*	*	*	*	*
Female	3.9	5.6	4.9	5.1	5.4	6.7	7.4	*	*

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	--------------

Human immunodeficiency virus (HIV) disease (B20-B24)

Non-Hispanic 5/	11.5	8.1	5.3	4.1	2.4	1.3	0.8	*	*
Male	18.0	13.2	9.1	6.6	3.6	2.5	1.6	*	*
Female	5.1	3.2	1.8	1.8	1.4	*	*	*	*
Non-Hispanic white	5.0	3.4	2.6	1.5	0.8	0.6	0.3	*	*
Male	8.8	6.2	4.8	2.6	1.4	1.4	*	*	*
Female	1.2	0.6	0.6	0.4	*	*	*	*	*
Non-Hispanic black	57.6	45.0	29.3	26.3	16.2	8.5	6.4	*	*
Male	88.7	72.3	50.7	43.7	25.8	15.5	12.8	*	*
Female	30.6	21.7	11.7	12.7	9.0	*	*	*	*

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	--------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Non-Hispanic 5/	1.4	1.9	2.3	4.0	5.7	6.9	9.3	12.4	20.3
Male	2.0	2.3	2.9	4.6	6.2	7.3	11.4	16.0	24.0
Female	0.9	1.5	1.8	3.4	5.2	6.5	7.9	10.2	18.8
Non-Hispanic white	1.3	1.7	2.0	3.7	5.5	6.6	9.2	12.2	20.2
Male	1.8	2.1	2.5	4.3	5.9	7.1	11.3	16.0	24.2
Female	0.7	1.4	1.5	3.1	5.1	6.2	7.7	10.0	18.6
Non-Hispanic black	2.6	3.5	4.5	6.2	7.1	9.8	9.9	13.5	20.9
Male	3.7	4.4	5.5	7.5	9.6	10.0	10.4	16.4	*
Female	1.6	2.8	3.7	5.1	5.2	9.6	9.6	12.1	20.5

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Malignant neoplasms (C00-C97)									
All origins 3/	92.4	164.8	280.7	453.3	673.2	943.5	1,211.7	1,465.9	1,765.6
Male	91.7	174.2	311.6	526.7	814.1	1,178.0	1,568.4	1,978.7	2,613.6
Female	93.2	155.9	251.7	386.7	551.9	757.2	962.0	1,158.6	1,410.7
Hispanic	61.7	110.3	172.3	291.3	413.5	616.9	846.0	1,033.1	1,363.7
Male	60.1	109.9	189.4	340.6	519.1	802.4	1,171.7	1,423.2	1,967.4
Female	63.3	110.7	156.7	248.7	328.0	475.3	614.9	791.4	1,068.6
Mexican	57.2	99.3	152.2	280.0	408.8	633.1	873.2	1,105.8	*
Male	55.0	91.8	156.0	344.4	476.0	815.8	*	*	*
Female	59.5	107.4	148.4	228.9	347.0	492.4	627.8	*	*
Puerto Rican	85.1	139.4	276.9	322.0	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Cuban	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	61.6	119.7	164.9	294.8	431.1	650.7	*	*	*
Male	57.7	119.8	178.1	*	*	*	*	*	*
Female	65.4	119.6	154.1	*	*	*	*	*	*

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	--------------

Malignant neoplasms (C00-C97)

Non-Hispanic 5/	95.3	169.1	288.6	464.3	690.0	960.9	1,226.9	1,480.7	1,776.6
Male	94.7	179.2	320.0	538.6	832.2	1,196.9	1,584.2	1,998.0	2,634.8
Female	95.8	159.5	259.0	396.6	567.0	773.0	976.9	1,171.2	1,419.6
Non-Hispanic white	90.2	162.2	279.7	458.2	682.7	955.9	1,222.7	1,473.0	1,775.2
Male	89.1	169.4	305.9	525.6	815.0	1,178.6	1,562.9	1,968.7	2,616.8
Female	91.3	155.1	254.6	395.8	565.8	775.0	981.8	1,173.8	1,425.1
Non-Hispanic black	141.4	246.5	412.8	598.1	871.7	1,172.3	1,446.2	1,735.8	1,947.5
Male	147.2	286.7	504.5	756.4	1,154.2	1,619.6	2,100.9	2,672.1	3,238.5
Female	136.3	212.4	337.8	473.7	663.1	874.2	1,055.5	1,258.9	1,469.3

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)									
Non-Hispanic 5/	2.1	3.6	6.3	8.1	9.5	11.8	13.6	15.4	18.4
Male	3.3	6.0	10.4	13.3	14.5	18.2	19.5	22.4	25.8
Female	1.0	1.4	2.5	3.4	5.3	6.8	9.4	11.3	15.3
Non-Hispanic white	1.8	3.3	5.7	7.5	8.9	11.6	13.6	15.7	18.9
Male	2.7	5.4	9.2	11.9	13.4	17.4	19.2	22.5	26.4
Female	0.8	1.2	2.3	3.3	5.0	6.9	9.7	11.6	15.7
Non-Hispanic black	4.4	6.4	12.5	13.4	15.5	16.0	14.2	13.5	13.1
Male	7.1	11.2	22.7	25.5	25.5	29.6	24.6	21.3	*
Female	2.0	2.3	4.1	3.9	8.1	6.9	7.9	9.6	10.7

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Malignant neoplasm of esophagus (C15)									
Non-Hispanic 5/	2.5	5.1	8.2	13.3	19.3	23.5	26.7	26.5	26.0
Male	4.3	9.0	14.3	23.2	33.8	40.8	47.6	48.9	48.4
Female	0.8	1.5	2.4	4.3	6.8	9.7	12.1	13.1	16.7
Non-Hispanic white	2.4	4.8	7.4	12.6	18.5	22.7	26.8	26.8	26.5
Male	4.1	8.3	13.1	22.1	32.5	39.8	48.1	49.5	49.0
Female	0.6	1.3	1.9	3.7	6.2	8.8	11.8	13.2	17.1
Non-Hispanic black	3.9	8.8	16.3	22.2	30.1	36.1	31.2	28.1	24.0
Male	6.2	15.7	28.3	39.1	52.6	60.9	55.9	54.8	47.4
Female	1.8	2.9	6.5	8.9	13.5	19.5	16.4	14.6	15.4

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Malignant neoplasm of stomach (C16)									
Non-Hispanic 5/	2.0	3.4	5.0	8.7	13.2	18.6	26.2	33.9	44.7
Male	2.6	4.6	7.2	13.1	18.7	26.3	38.3	48.2	61.0
Female	1.3	2.2	2.9	4.8	8.4	12.5	17.8	25.3	37.9
Non-Hispanic white	1.6	2.6	4.3	7.3	10.8	16.3	23.0	29.8	40.0
Male	2.1	3.6	6.1	10.9	15.1	23.0	34.2	42.4	55.7
Female	1.0	1.7	2.4	3.9	7.1	10.9	15.0	22.2	33.5
Non-Hispanic black	3.8	7.5	9.3	18.0	27.8	35.0	51.7	70.6	89.3
Male	5.2	11.6	13.8	28.0	44.8	55.1	76.8	107.1	106.3
Female	2.6	4.1	5.5	10.2	15.2	21.6	36.8	52.1	83.0

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Malignant neoplasms of colon, rectum and anus (C18-C21)									
Non-Hispanic 5/	8.8	15.6	26.0	39.9	64.4	89.9	122.4	167.9	257.4
Male	9.9	17.6	31.5	48.4	80.5	111.7	151.0	204.3	299.6
Female	7.6	13.6	20.9	32.2	50.5	72.6	102.4	146.1	239.9
Non-Hispanic white	8.0	14.5	24.4	38.4	62.2	87.8	119.2	165.7	256.9
Male	9.1	16.8	29.9	47.0	78.0	109.8	148.2	201.9	298.2
Female	6.9	12.3	19.2	30.4	48.2	69.9	98.6	143.8	239.8
Non-Hispanic black	14.8	25.2	44.0	59.0	92.8	122.1	174.5	213.3	289.0
Male	16.8	27.0	52.3	68.3	117.0	148.7	209.5	264.9	352.3
Female	13.1	23.7	37.2	51.6	75.0	104.4	153.6	187.0	265.6

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Malignant neoplasms of liver and intrahepatic bile ducts (C22)									
Non-Hispanic 5/	3.6	5.6	7.1	11.1	15.7	20.6	25.7	31.0	33.2
Male	5.6	9.4	10.6	16.6	22.6	29.4	37.6	42.9	49.7
Female	1.7	2.0	3.7	6.1	9.7	13.7	17.4	23.9	26.4
Non-Hispanic white	2.9	4.6	5.7	9.9	14.2	19.2	24.6	29.2	32.4
Male	4.3	7.5	8.4	14.9	21.0	27.6	35.9	41.3	48.9
Female	1.4	1.7	3.2	5.2	8.2	12.3	16.6	22.0	25.5
Non-Hispanic black	7.3	10.9	15.6	14.5	20.1	24.1	26.2	41.7	35.3
Male	11.8	19.1	26.1	21.7	28.1	33.1	38.4	49.9	42.7
Female	3.3	3.9	7.0	8.9	14.2	18.1	19.0	37.5	32.5

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Malignant neoplasm of pancreas (C25)									
Non-Hispanic 5/	4.2	8.8	15.1	25.3	37.4	53.0	67.6	84.6	100.6
Male	5.1	11.1	18.9	31.1	43.9	59.3	77.7	95.5	106.4
Female	3.2	6.6	11.6	20.0	31.8	48.0	60.5	78.0	98.2
Non-Hispanic white	3.9	8.6	14.6	25.3	36.7	51.8	66.7	83.6	99.4
Male	4.9	10.8	18.6	31.0	43.1	58.3	77.6	95.1	108.0
Female	3.0	6.4	10.9	19.9	31.0	46.6	58.9	76.7	95.8
Non-Hispanic black	6.5	12.5	21.1	31.9	48.9	71.5	85.2	106.0	122.0
Male	8.0	15.6	24.6	38.9	60.2	78.8	94.8	110.4	100.5
Female	5.2	9.8	18.2	26.5	40.6	66.5	79.5	103.7	130.0

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Malignant neoplasm of larynx (C32)									
Non-Hispanic 5/	0.7	1.6	3.0	4.6	5.5	7.4	7.5	7.0	6.7
Male	1.2	2.6	5.0	7.6	9.4	13.0	14.5	14.0	18.4
Female	0.3	0.5	1.1	1.8	2.3	3.0	2.6	2.8	1.8
Non-Hispanic white	0.5	1.3	2.6	4.2	5.2	6.8	7.0	6.8	6.5
Male	0.9	2.1	4.3	7.0	8.6	11.9	13.3	13.3	18.3
Female	*	0.5	0.9	1.6	2.1	2.7	2.6	2.9	1.6
Non-Hispanic black	2.0	3.9	7.1	8.9	9.6	13.9	13.6	9.4	8.4
Male	3.7	7.1	12.9	15.9	17.7	26.2	31.3	22.9	*
Female	*	*	*	3.5	*	5.7	*	*	*

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	--------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

Non-Hispanic 5/	21.9	44.4	93.2	161.7	249.8	334.1	382.4	375.1	305.8
Male	25.9	54.6	114.3	203.5	324.7	447.4	533.1	549.4	518.4
Female	17.9	34.6	73.3	123.7	185.0	243.9	276.9	270.8	217.4
Non-Hispanic white	20.7	43.2	92.7	163.9	253.2	339.3	386.7	377.9	308.6
Male	23.8	51.4	110.5	201.8	322.2	446.0	530.5	545.9	520.0
Female	17.7	35.1	75.5	128.8	192.3	252.7	284.9	276.5	220.6
Non-Hispanic black	33.6	64.6	121.4	185.0	277.6	353.4	401.1	375.0	281.6
Male	46.2	93.9	177.0	271.9	426.0	561.5	668.4	667.2	550.9
Female	22.6	39.8	75.8	116.7	168.1	214.7	241.6	226.1	181.8

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Malignant neoplasm of breast (C50)									
All origins 3/	13.4	20.6	27.1	34.1	41.7	54.0	69.1	88.0	134.2
Male	*	0.4	0.5	0.8	1.2	1.7	2.3	2.4	3.5
Female	26.3	40.1	52.0	64.3	76.5	95.6	115.9	139.4	188.9
Hispanic	9.6	15.2	20.5	24.1	25.1	31.7	38.8	40.7	68.8
Male	*	*	*	*	*	*	*	*	*
Female	19.2	29.5	39.1	44.7	44.2	55.5	64.9	64.3	102.4
Mexican	8.3	14.1	17.6	21.0	25.5	28.3	34.0	*	*
Male	*	*	*	*	*	*	*	*	*
Female	16.8	29.4	35.4	37.7	47.9	49.4	*	*	*
Puerto Rican	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Cuban	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	10.4	17.9	20.3	29.8	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	20.3	32.1	36.9	*	*	*	*	*	*

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Malignant neoplasm of breast (C50)									
Non-Hispanic 5/	13.8	21.1	27.6	34.7	42.7	55.3	70.4	89.8	136.4
Male	*	0.4	0.5	0.8	1.2	1.8	2.3	2.4	3.7
Female	27.0	40.9	53.0	65.7	78.7	97.9	118.0	142.1	191.6
Non-Hispanic white	12.6	19.8	26.2	33.9	41.9	54.9	70.8	89.9	137.1
Male	*	0.3	0.5	0.8	1.1	1.7	2.1	2.4	3.5
Female	24.9	38.8	50.8	64.6	78.0	98.1	119.4	142.8	192.6
Non-Hispanic black	23.2	33.7	43.4	48.5	58.9	71.7	83.8	107.3	153.9
Male	*	*	*	*	*	*	*	*	*
Female	42.9	61.5	78.3	85.6	101.0	117.2	131.3	160.3	208.3

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Malignant neoplasm of cervix uteri (C53)									
Non-Hispanic 5/	2.2	2.6	2.5	3.1	3.0	3.4	3.9	5.4	7.0
Female	4.3	5.0	4.9	5.9	5.5	6.1	6.7	8.6	10.0
Non-Hispanic white	2.0	2.3	2.2	2.7	2.4	2.9	3.3	4.5	6.1
Female	4.0	4.5	4.2	5.2	4.4	5.3	5.6	7.3	8.6
Non-Hispanic black	3.3	4.5	4.7	6.1	8.0	7.8	9.7	15.5	19.0
Female	6.1	8.3	8.6	10.8	13.8	13.1	15.6	23.3	26.1

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	--------------

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

Non-Hispanic 5/	0.9	2.0	3.2	5.9	8.4	11.5	14.7	19.3	24.8
Female	1.8	3.9	6.2	11.2	15.6	20.6	25.0	30.9	35.1
Non-Hispanic white	0.9	1.9	3.1	5.2	7.3	10.5	13.9	18.7	24.5
Female	1.7	3.7	6.1	10.0	13.7	19.0	23.8	30.0	34.7
Non-Hispanic black	1.4	3.0	4.6	11.7	19.1	24.6	26.6	31.7	32.8
Female	2.7	5.6	8.3	20.9	33.3	41.1	42.4	47.9	44.9

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Malignant neoplasm of ovary (C56)									
Non-Hispanic 5/	3.0	5.7	8.5	13.2	17.8	23.0	30.9	37.8	39.8
Female	5.9	11.1	16.4	25.2	33.3	41.4	52.4	60.4	56.4
Non-Hispanic white	3.2	5.9	8.7	13.4	18.3	23.8	31.8	39.0	40.6
Female	6.4	11.8	17.1	25.9	34.4	43.2	54.3	62.5	57.5
Non-Hispanic black	2.0	4.5	8.1	12.8	16.6	20.8	26.4	28.4	36.5
Female	3.7	8.3	14.8	22.8	28.9	34.6	42.2	42.9	50.0

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Malignant neoplasm of prostate (C61)									
Non-Hispanic 5/	0.5	1.7	4.6	11.5	25.1	47.9	82.3	134.0	201.6
Male	1.0	3.5	9.5	24.1	54.1	108.1	199.9	357.9	686.3
Non-Hispanic white	0.4	1.4	3.9	9.7	22.0	42.4	74.8	123.6	189.5
Male	0.8	2.8	8.0	20.2	46.9	94.5	180.5	328.4	645.0
Non-Hispanic black	1.2	4.9	11.8	29.3	58.2	116.5	182.0	283.4	365.8
Male	2.5	10.7	26.2	66.5	137.0	291.3	486.8	839.7	1,353.6

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Malignant neoplasms of kidney and renal pelvis (C64-C65)									
Non-Hispanic 5/	2.4	4.4	7.3	12.0	14.8	21.0	24.4	29.3	32.3
Male	3.4	6.5	10.8	17.4	21.2	30.1	34.3	42.4	52.6
Female	1.4	2.4	4.0	7.1	9.2	13.7	17.4	21.5	23.8
Non-Hispanic white	2.4	4.4	7.5	12.2	15.4	21.5	24.7	29.7	32.8
Male	3.6	6.5	11.1	17.3	22.0	30.8	34.3	42.9	54.3
Female	1.3	2.4	4.0	7.5	9.6	14.0	17.9	21.7	23.8
Non-Hispanic black	2.6	5.1	7.5	12.5	12.0	19.4	24.3	28.7	30.0
Male	3.5	8.0	10.8	21.1	17.5	29.6	39.3	43.3	46.2
Female	1.9	2.7	4.9	5.8	7.9	12.6	15.3	21.2	24.0

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Malignant neoplasm of bladder (C67)									
Non-Hispanic 5/	0.8	1.8	3.6	7.3	12.4	19.5	30.0	45.5	69.6
Male	1.2	2.8	5.4	11.6	20.1	32.6	52.2	82.2	136.4
Female	0.4	0.9	1.8	3.3	5.7	9.1	14.5	23.6	41.9
Non-Hispanic white	0.8	1.9	3.7	7.6	13.0	20.4	31.0	46.4	71.5
Male	1.2	2.9	5.6	12.4	21.4	34.3	54.7	85.0	143.0
Female	0.3	0.9	1.8	3.2	5.6	9.2	14.3	23.1	41.7
Non-Hispanic black	0.8	1.8	3.5	7.1	10.3	14.3	24.8	41.7	55.6
Male	*	2.9	4.3	9.4	13.0	21.0	34.1	58.9	78.5
Female	*	*	2.8	5.3	8.3	9.9	19.2	32.9	47.1

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	--------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Non-Hispanic 5/	4.2	6.3	8.6	11.8	14.5	17.1	20.6	20.7	16.5
Male	5.1	7.9	11.0	14.3	17.7	21.4	25.8	26.7	20.5
Female	3.3	4.9	6.3	9.5	11.7	13.6	16.9	17.1	14.9
Non-Hispanic white	4.6	7.0	9.2	12.9	15.6	18.2	22.0	21.8	17.2
Male	5.6	8.6	11.8	15.4	19.1	22.8	27.2	28.0	20.9
Female	3.6	5.4	6.7	10.6	12.4	14.5	18.3	18.0	15.7
Non-Hispanic black	2.7	3.5	5.5	6.3	9.2	9.6	9.2	11.0	11.9
Male	3.1	4.6	7.1	9.0	9.4	11.7	11.9	*	*
Female	2.3	2.7	4.2	4.3	9.0	8.3	7.6	10.4	9.8

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)									
Non-Hispanic 5/	8.2	13.1	23.1	38.3	59.7	93.2	130.3	170.5	202.1
Male	10.0	16.2	28.1	48.0	76.3	118.7	166.4	223.6	278.8
Female	6.5	10.2	18.5	29.5	45.4	72.9	105.0	138.7	170.2
Non-Hispanic white	8.0	12.9	22.6	38.2	60.0	95.2	133.3	174.9	208.2
Male	9.9	16.0	27.4	47.9	76.7	121.2	170.6	229.3	288.7
Female	6.2	9.8	18.0	29.3	45.2	74.0	106.8	142.1	174.6
Non-Hispanic black	10.7	16.9	31.0	44.9	68.4	92.2	117.1	144.0	157.3
Male	12.1	21.4	38.5	57.1	89.4	115.7	142.2	194.6	206.7
Female	9.5	13.0	24.8	35.3	52.9	76.6	102.1	118.3	139.0

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Non-Hodgkin's lymphoma (C82-C85)									
Non-Hispanic 5/	3.5	5.6	9.8	15.6	23.5	36.7	51.2	70.6	80.3
Male	4.5	7.5	12.1	19.1	29.7	45.5	61.3	88.2	108.1
Female	2.5	3.8	7.6	12.4	18.1	29.8	44.1	60.1	68.7
Non-Hispanic white	3.4	5.7	9.9	16.3	24.5	38.7	54.0	74.0	84.1
Male	4.5	7.6	12.1	19.8	30.9	47.5	64.3	92.2	113.1
Female	2.4	3.9	7.8	13.1	18.8	31.6	46.7	63.0	72.0
Non-Hispanic black	4.3	6.1	9.9	12.1	19.3	24.0	25.9	38.1	40.0
Male	5.1	8.8	12.4	15.5	25.0	32.4	30.8	48.2	55.4
Female	3.7	3.7	7.9	9.5	15.0	18.4	22.9	32.9	34.2

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Leukemia (C91-C95)									
Non-Hispanic 5/	3.1	4.7	7.8	13.4	21.0	33.9	48.4	62.2	82.7
Male	3.4	5.3	9.5	17.7	28.3	46.0	66.9	88.7	119.3
Female	2.8	4.1	6.2	9.5	14.7	24.3	35.5	46.4	67.5
Non-Hispanic white	3.1	4.6	7.9	13.7	21.4	35.2	49.8	64.8	86.1
Male	3.5	5.2	9.7	18.1	28.6	47.8	69.0	92.0	125.3
Female	2.7	3.9	6.2	9.7	15.0	25.0	36.2	48.4	69.8
Non-Hispanic black	3.1	5.7	8.1	12.5	22.4	28.4	41.8	42.5	57.1
Male	2.9	6.8	10.1	17.6	32.6	36.8	52.6	62.1	75.1
Female	3.4	4.8	6.5	8.5	14.8	22.7	35.4	32.5	50.5

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85+ YEARS

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

Non-Hispanic 5/	1.2	2.3	5.2	8.5	14.1	21.2	28.5	34.8	36.5
Male	1.6	2.8	6.1	10.2	16.8	25.3	35.6	43.4	48.4
Female	0.9	1.9	4.3	6.9	11.8	17.9	23.5	29.7	31.5
Non-Hispanic white	1.0	2.0	4.4	7.4	12.9	19.9	27.2	33.1	35.1
Male	1.4	2.5	5.2	9.1	15.5	23.9	34.6	41.6	47.3
Female	0.7	1.6	3.7	5.8	10.6	16.6	22.0	28.0	30.0
Non-Hispanic black	2.7	4.8	12.1	19.3	26.2	38.8	47.7	61.8	59.3
Male	3.5	5.4	15.0	23.0	31.0	45.4	56.9	81.8	72.8
Female	2.0	4.4	9.8	16.4	22.7	34.4	42.2	51.6	54.3

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	--------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Non-Hispanic 5/	10.8	18.4	30.9	47.2	68.5	98.8	134.8	172.0	235.3
Male	12.4	22.0	35.7	56.3	81.2	121.3	163.7	212.5	297.9
Female	9.3	15.0	26.4	38.8	57.5	81.0	114.6	147.7	209.3
Non-Hispanic white	10.1	17.5	29.7	45.9	67.7	98.1	135.5	172.5	238.9
Male	11.6	20.5	33.8	54.3	79.8	120.8	164.4	213.7	302.9
Female	8.6	14.4	25.8	38.1	57.1	79.6	115.1	147.6	212.3
Non-Hispanic black	17.2	28.4	45.2	65.3	86.5	120.7	143.0	182.4	218.5
Male	19.4	36.6	58.6	82.6	111.8	150.5	181.6	218.3	287.6
Female	15.2	21.5	34.2	51.8	67.9	100.9	120.0	164.1	192.9

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	--------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Non-Hispanic 5/	1.5	2.3	4.0	6.3	10.7	18.3	30.0	48.3	83.5
Male	1.7	2.4	4.2	7.5	13.1	23.0	40.1	62.7	109.0
Female	1.3	2.1	3.8	5.3	8.6	14.6	22.9	39.7	72.9
Non-Hispanic white	1.5	2.3	4.0	6.3	11.0	18.6	30.9	49.4	84.8
Male	1.8	2.5	4.3	7.4	13.5	23.6	41.3	63.9	111.4
Female	1.3	2.1	3.8	5.3	8.7	14.5	23.5	40.6	73.7
Non-Hispanic black	1.5	2.8	4.6	7.7	11.0	18.9	25.0	42.8	72.7
Male	*	2.5	4.5	9.4	13.3	23.4	32.2	59.7	87.8
Female	*	3.0	4.6	6.4	9.2	15.8	20.7	34.1	67.2

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Anemias (D50-D64)									
Non-Hispanic 5/	0.6	0.7	0.8	1.2	1.9	3.6	7.1	13.7	42.5
Male	0.6	0.7	0.8	1.1	2.0	3.9	7.1	14.4	40.8
Female	0.5	0.7	0.9	1.3	1.9	3.3	7.0	13.3	43.1
Non-Hispanic white	0.2	0.4	0.5	1.1	1.7	3.3	6.7	13.2	42.1
Male	*	0.4	0.5	1.0	1.8	3.4	6.9	14.2	41.2
Female	*	0.4	0.5	1.1	1.7	3.1	6.5	12.7	42.4
Non-Hispanic black	3.3	2.9	3.3	2.7	4.0	7.4	12.6	21.5	49.3
Male	4.0	3.2	3.4	*	*	10.7	12.3	20.4	42.7
Female	2.7	2.7	3.3	*	4.0	5.3	12.7	22.1	51.8

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	--------------

Diabetes mellitus (E10-E14)

Non-Hispanic 5/	9.7	17.6	28.7	46.4	71.9	104.7	150.1	213.9	314.8
Male	11.7	21.0	33.6	52.4	81.0	117.1	166.4	238.1	338.9
Female	7.7	14.4	24.1	41.0	64.1	94.7	138.7	199.4	304.7
Non-Hispanic white	8.1	14.7	23.9	39.4	62.0	92.7	136.3	199.9	297.4
Male	9.9	17.9	29.0	45.9	72.8	108.6	155.6	226.9	332.1
Female	6.3	11.6	18.9	33.3	52.5	79.7	122.6	183.7	283.0
Non-Hispanic black	21.3	41.0	70.6	106.6	159.4	229.9	307.4	385.5	533.8
Male	25.8	46.4	77.8	114.5	163.9	223.4	310.5	404.7	452.7
Female	17.4	36.4	64.8	100.4	156.0	234.2	305.5	375.6	563.8

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85+ YEARS

Nutritional deficiencies (E40-E64)

Non-Hispanic 5/	0.2	0.4	0.5	0.8	1.4	2.9	5.9	12.9	43.0
Male	0.3	0.3	0.5	0.9	1.5	3.0	5.5	13.2	34.8
Female	0.2	0.4	0.4	0.8	1.3	2.7	6.2	12.7	46.4
Non-Hispanic white	0.2	0.3	0.4	0.8	1.2	2.7	5.4	12.0	42.2
Male	*	*	0.4	0.8	1.2	2.7	5.1	12.3	33.3
Female	*	0.3	0.4	0.8	1.3	2.7	5.7	11.8	45.9
Non-Hispanic black	*	1.1	*	*	3.1	4.5	11.0	24.0	55.2
Male	*	*	*	*	*	6.9	11.9	24.5	58.9
Female	*	*	*	*	*	*	10.5	23.7	53.9

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Malnutrition (E40-E46)									
Non-Hispanic 5/	0.2	0.3	0.5	0.8	1.3	2.6	5.4	12.1	40.2
Male	0.2	0.3	0.5	0.8	1.4	2.7	4.9	12.7	33.2
Female	*	0.4	0.4	0.8	1.3	2.6	5.7	11.7	43.2
Non-Hispanic white	0.2	0.2	0.4	0.8	1.2	2.5	5.0	11.2	39.4
Male	*	*	0.4	0.7	1.1	2.5	4.5	11.8	31.7
Female	*	0.3	0.4	0.8	1.2	2.5	5.3	10.9	42.7
Non-Hispanic black	*	*	*	*	2.9	4.1	10.3	22.3	52.1
Male	*	*	*	*	*	*	11.4	23.7	56.6
Female	*	*	*	*	*	*	9.6	21.7	50.5

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85+ YEARS

Parkinson's disease (G20-G21)

Non-Hispanic 5/	*	0.2	0.7	1.9	5.9	18.4	47.1	91.8	135.6
Male	*	0.3	1.1	2.8	8.3	27.1	69.9	142.9	227.8
Female	*	*	0.4	1.0	3.8	11.5	31.1	61.3	97.3
Non-Hispanic white	*	0.2	0.8	2.0	6.3	19.7	50.4	97.5	143.5
Male	*	0.3	1.1	3.1	8.9	28.7	74.7	151.0	241.7
Female	*	*	0.4	0.9	4.0	12.4	33.1	65.2	102.7
Non-Hispanic black	*	*	*	*	2.8	9.1	19.3	35.6	59.0
Male	*	*	*	*	*	15.1	26.1	58.9	99.3
Female	*	*	*	*	*	5.0	15.3	23.7	44.1

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	--------------

Alzheimer's disease (G30)

Non-Hispanic 5/	*	0.3	1.2	3.4	9.3	30.0	90.0	236.3	720.5
Male	*	0.3	1.0	3.3	9.2	28.5	86.6	220.3	565.2
Female	*	0.3	1.4	3.5	9.3	31.1	92.3	245.8	785.1
Non-Hispanic white	*	0.4	1.4	3.7	9.6	31.5	92.5	244.5	749.2
Male	*	0.3	1.0	3.5	9.4	29.7	89.2	226.7	587.2
Female	*	0.4	1.7	3.9	9.8	32.9	94.9	255.3	816.6
Non-Hispanic black	*	*	*	1.8	8.9	23.8	88.4	186.8	513.2
Male	*	*	*	*	9.9	24.8	86.3	188.1	420.4
Female	*	*	*	*	8.1	23.2	89.7	186.2	547.5

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85+ YEARS

Major cardiovascular diseases (I00-I78)

All origins 3/	85.6	144.9	234.9	385.7	619.6	1,017.5	1,759.4	3,133.9	7,673.7
Male	121.4	204.7	327.6	526.3	821.4	1,325.6	2,223.2	3,774.2	7,936.0
Female	50.7	87.6	148.1	258.3	445.9	772.7	1,434.7	2,750.2	7,563.9
Hispanic	54.0	103.2	165.7	283.6	490.6	837.6	1,459.3	2,475.8	6,036.2
Male	77.7	146.6	233.8	383.5	672.6	1,084.7	1,808.6	3,004.8	6,155.6
Female	30.2	61.7	104.1	197.2	343.2	649.1	1,211.3	2,148.2	5,977.9
Mexican	51.7	96.8	154.1	285.9	493.1	877.3	1,487.8	2,533.1	*
Male	74.9	130.1	204.4	391.9	635.5	1,126.5	*	*	*
Female	27.9	60.7	103.4	201.8	362.4	685.4	1,229.5	*	*
Puerto Rican	72.1	132.2	309.9	362.0	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Cuban	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	49.0	99.5	132.7	238.7	445.5	765.6	*	*	*
Male	69.0	153.1	201.9	*	*	*	*	*	*
Female	29.9	55.4	76.0	*	*	*	*	*	*

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	--------------

Major cardiovascular diseases (I00-I78)

Non-Hispanic 5/	88.3	147.8	239.3	391.7	626.6	1,025.2	1,768.6	3,152.1	7,715.2
Male	125.4	208.7	333.1	533.8	828.4	1,335.1	2,236.3	3,795.4	7,990.6
Female	52.6	89.5	151.1	262.3	452.1	778.4	1,441.4	2,767.2	7,600.7
Non-Hispanic white	76.0	129.1	215.0	359.1	584.0	984.9	1,727.4	3,139.5	7,811.4
Male	111.7	188.0	305.3	498.8	785.3	1,292.9	2,203.7	3,797.2	8,116.3
Female	40.6	71.4	128.4	229.5	406.1	734.6	1,390.3	2,742.5	7,684.6
Non-Hispanic black	185.9	313.3	478.6	725.1	1,073.1	1,567.7	2,444.2	3,650.2	7,366.6
Male	242.7	413.4	637.0	947.4	1,361.6	2,006.9	2,966.0	4,316.4	7,413.8
Female	136.8	228.2	348.9	550.3	860.0	1,275.0	2,132.8	3,310.8	7,349.1

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Diseases of heart (I00-I09,I11,I13,I20-I51)									
All origins 3/	69.3	119.5	194.2	314.1	494.4	787.7	1,321.2	2,309.0	5,664.2
Male	102.6	175.2	279.7	440.9	672.2	1,054.4	1,732.2	2,882.7	6,040.5
Female	37.0	66.1	114.2	199.0	341.4	575.8	1,033.5	1,965.3	5,506.8
Hispanic	38.3	77.1	129.3	225.5	391.0	646.2	1,097.2	1,866.0	4,607.5
Male	58.7	115.3	191.0	311.8	547.1	860.1	1,385.1	2,307.5	4,784.3
Female	17.8	40.7	73.5	150.9	264.6	483.0	892.8	1,592.5	4,521.1
Mexican	35.7	71.6	119.8	220.5	385.3	663.7	1,083.7	1,840.6	*
Male	56.0	103.0	165.8	308.6	503.3	862.5	*	*	*
Female	14.9	37.5	73.4	150.5	276.9	510.7	875.1	*	*
Puerto Rican	55.4	103.6	248.9	295.6	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Cuban	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	33.8	69.5	99.8	194.9	347.4	563.0	*	*	*
Male	50.1	113.4	164.9	*	*	*	*	*	*
Female	18.2	33.3	46.3	*	*	*	*	*	*

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Diseases of heart (I00-I09,I11,I13,I20-I51)									
Non-Hispanic 5/	72.1	122.6	198.4	319.3	499.9	793.7	1,327.9	2,320.3	5,688.6
Male	106.6	179.5	285.1	448.0	678.1	1,061.9	1,743.4	2,898.2	6,077.8
Female	38.7	68.0	117.0	202.2	345.9	580.0	1,037.2	1,974.4	5,526.8
Non-Hispanic white	64.0	110.9	181.6	298.4	471.8	769.0	1,302.6	2,318.8	5,776.0
Male	97.9	166.9	265.9	426.1	650.7	1,037.0	1,726.7	2,911.3	6,195.4
Female	30.5	56.0	100.8	180.0	313.7	551.2	1,002.4	1,961.1	5,601.6
Non-Hispanic black	140.3	236.6	375.4	554.8	820.9	1,173.2	1,795.1	2,634.8	5,307.7
Male	188.8	320.2	510.4	740.5	1,053.7	1,531.1	2,230.3	3,202.0	5,495.4
Female	98.4	165.5	264.8	408.8	649.0	934.8	1,535.3	2,345.9	5,238.2

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)									
Non-Hispanic 5/	0.5	0.7	1.2	1.7	3.2	5.0	8.2	11.6	18.0
Male	0.5	0.5	1.1	1.2	2.3	3.3	6.0	7.8	13.1
Female	0.6	0.9	1.3	2.1	4.0	6.3	9.8	13.9	20.0
Non-Hispanic white	0.4	0.6	1.1	1.7	3.1	5.0	8.6	12.4	19.0
Male	0.4	0.4	1.0	1.2	2.2	3.4	6.3	8.2	13.7
Female	0.4	0.7	1.1	2.1	4.0	6.3	10.2	14.9	21.2
Non-Hispanic black	1.1	1.7	2.2	2.0	3.4	5.2	5.0	*	6.9
Male	*	*	*	*	*	*	*	*	*
Female	*	2.3	2.8	*	3.8	7.3	6.5	*	*

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS.]

FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Hypertensive heart disease (I11)									
Non-Hispanic 5/	5.2	7.6	9.7	12.4	16.3	23.2	36.4	66.0	190.3
Male	7.2	10.2	12.7	16.2	20.4	26.5	38.6	63.4	133.4
Female	3.3	5.2	6.8	9.0	12.7	20.5	34.8	67.6	214.0
Non-Hispanic white	3.3	5.0	6.7	8.8	11.8	17.9	30.4	58.9	180.4
Male	4.8	7.1	9.4	12.0	15.4	21.1	32.5	55.5	122.0
Female	1.8	2.9	4.2	5.8	8.6	15.3	28.9	60.9	204.7
Non-Hispanic black	18.9	28.5	35.7	44.1	56.4	78.2	106.1	155.9	328.7
Male	25.5	36.5	45.2	56.4	71.1	91.2	116.6	173.3	288.7
Female	13.2	21.7	28.0	34.3	45.6	69.5	99.9	147.0	343.5

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Hypertensive heart and renal disease (I13)									
Non-Hispanic 5/	0.3	0.4	0.8	1.2	1.7	2.7	5.0	9.2	23.1
Male	0.5	0.6	0.9	1.1	2.0	2.8	5.3	9.0	23.8
Female	*	0.3	0.7	1.3	1.4	2.7	4.8	9.3	22.8
Non-Hispanic white	*	0.2	0.3	0.6	0.9	1.8	3.8	8.1	20.7
Male	*	*	0.4	0.5	1.1	1.9	4.2	8.3	22.2
Female	*	*	*	0.7	0.8	1.8	3.6	7.9	20.1
Non-Hispanic black	1.9	2.4	4.4	5.9	8.6	11.8	18.4	22.9	50.6
Male	3.0	3.6	5.6	6.5	10.9	13.4	18.5	18.8	45.0
Female	*	*	3.4	5.4	6.9	10.8	18.4	25.0	52.6

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Ischemic heart diseases (I20-I25)									
All origins 3/	45.6	84.8	142.8	236.6	372.4	591.8	984.3	1,685.2	3,958.2
Male	71.6	131.1	215.5	344.4	525.7	820.3	1,337.5	2,180.0	4,395.8
Female	20.3	40.3	74.8	138.8	240.5	410.2	737.0	1,388.7	3,775.1
Hispanic	25.8	55.5	98.1	172.8	306.5	518.7	881.4	1,481.0	3,620.2
Male	41.3	85.6	152.6	245.2	441.9	712.3	1,147.8	1,860.8	3,828.2
Female	10.1	26.6	48.7	110.0	196.9	371.0	692.3	1,245.8	3,518.6
Mexican	24.6	52.6	90.4	167.8	304.5	530.1	872.2	1,440.7	*
Male	39.4	78.0	131.9	244.5	405.9	709.3	*	*	*
Female	9.4	25.1	48.5	106.8	211.4	392.2	664.0	*	*
Puerto Rican	36.0	72.2	186.9	228.3	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Cuban	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	21.0	48.7	76.7	146.0	255.4	436.4	*	*	*
Male	35.6	83.6	131.4	*	*	*	*	*	*
Female	*	19.8	31.8	*	*	*	*	*	*

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Ischemic heart diseases (I20-I25)									
Non-Hispanic 5/	47.4	86.9	145.7	240.2	375.6	594.1	985.9	1,688.1	3,959.4
Male	74.5	134.5	219.2	349.7	529.1	823.6	1,342.0	2,185.7	4,408.5
Female	21.2	41.4	76.6	140.6	242.9	411.4	736.7	1,390.3	3,772.7
Non-Hispanic white	44.8	82.1	137.8	228.7	359.7	581.5	973.2	1,692.1	4,022.8
Male	72.2	129.4	210.4	338.5	514.8	812.4	1,338.8	2,207.1	4,506.9
Female	17.7	35.8	68.1	126.7	222.7	393.8	714.3	1,381.2	3,821.4
Non-Hispanic black	74.3	141.7	239.4	381.7	570.0	820.0	1,257.4	1,844.5	3,628.7
Male	104.5	203.8	337.8	522.1	752.5	1,102.1	1,591.8	2,254.3	3,791.8
Female	48.2	88.8	158.8	271.3	435.3	632.0	1,057.8	1,635.8	3,568.3

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Acute myocardial infarction (I21-I22)									
Non-Hispanic 5/	20.6	37.8	63.2	102.1	156.7	237.5	375.3	614.1	1,265.3
Male	32.1	57.7	94.0	146.7	217.3	324.5	498.6	779.7	1,463.9
Female	9.5	18.7	34.2	61.6	104.3	168.3	288.9	515.1	1,182.7
Non-Hispanic white	20.2	36.7	61.6	100.5	153.1	234.9	372.6	614.8	1,276.8
Male	32.4	57.3	93.3	147.1	216.2	324.4	500.3	786.9	1,483.2
Female	8.1	16.5	31.1	57.2	97.4	162.3	282.2	510.9	1,190.9
Non-Hispanic black	28.6	54.3	90.9	137.8	214.1	308.1	458.0	677.1	1,247.0
Male	37.5	72.9	122.5	176.6	267.8	390.8	556.0	802.9	1,389.4
Female	20.8	38.5	65.0	107.3	174.5	253.0	399.5	613.0	1,194.3

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Other acute ischemic heart diseases (I24)									
Non-Hispanic 5/	0.6	1.1	1.5	2.5	3.0	4.3	5.5	9.5	22.0
Male	1.0	1.7	2.0	3.7	4.2	5.6	6.9	11.6	20.9
Female	0.3	0.5	1.0	1.4	2.1	3.2	4.5	8.2	22.4
Non-Hispanic white	0.6	0.9	1.3	2.2	2.7	4.2	5.3	9.4	22.0
Male	0.9	1.4	1.9	3.3	3.6	5.5	6.7	11.9	20.9
Female	*	0.5	0.8	1.2	2.0	3.1	4.4	7.9	22.4
Non-Hispanic black	1.1	2.2	2.9	5.3	5.4	6.3	6.9	11.0	24.3
Male	*	3.7	3.5	8.4	9.4	7.9	11.4	*	24.3
Female	*	*	*	*	*	5.3	*	12.1	24.4

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Other forms of chronic ischemic heart disease (I20,I25)									
Non-Hispanic 5/	26.1	48.0	81.1	135.6	215.9	352.3	605.2	1,064.5	2,672.2
Male	41.4	75.1	123.2	199.4	307.7	493.5	836.5	1,394.4	2,923.7
Female	11.4	22.1	41.5	77.6	136.6	239.9	443.3	867.0	2,567.6
Non-Hispanic white	24.0	44.5	74.9	126.0	203.8	342.4	595.3	1,067.8	2,724.1
Male	39.0	70.6	115.2	188.2	295.0	482.6	831.9	1,408.3	3,002.8
Female	9.3	18.8	36.2	68.2	123.3	228.4	427.7	862.3	2,608.1
Non-Hispanic black	44.6	85.2	145.6	238.5	350.5	505.5	792.5	1,156.4	2,357.3
Male	65.5	127.3	211.7	337.1	475.2	703.4	1,024.4	1,442.4	2,378.1
Female	26.6	49.4	91.4	161.0	258.3	373.7	654.1	1,010.7	2,349.6

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Atherosclerotic cardiovascular disease, so described (I25.0)									
Non-Hispanic 5/	9.1	17.5	27.7	40.5	54.8	75.9	121.3	197.5	494.8
Male	14.9	28.2	43.2	61.4	77.6	106.0	158.8	247.2	503.0
Female	3.4	7.2	13.1	21.5	35.1	51.9	95.0	167.7	491.4
Non-Hispanic white	8.1	15.4	23.6	34.4	46.7	68.0	111.2	190.8	490.7
Male	13.6	25.3	37.7	52.9	66.9	95.2	146.4	239.6	499.4
Female	2.6	5.7	10.1	17.3	28.8	45.9	86.3	161.3	487.0
Non-Hispanic black	17.1	36.3	64.5	95.5	129.4	164.1	244.8	300.8	595.9
Male	26.3	56.7	97.3	145.3	185.0	241.7	332.8	385.9	609.8
Female	9.2	18.9	37.6	56.4	88.3	112.4	192.4	257.4	590.7

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85+ YEARS

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

Non-Hispanic 5/	17.1	30.6	53.4	95.1	161.1	276.4	483.9	867.0	2,177.4
Male	26.5	46.9	79.9	138.0	230.1	387.4	677.7	1,147.3	2,420.7
Female	8.0	14.9	28.4	56.1	101.5	188.0	348.3	699.3	2,076.2
Non-Hispanic white	16.0	29.1	51.3	91.6	157.1	274.4	484.0	877.0	2,233.4
Male	25.4	45.4	77.5	135.4	228.0	387.4	685.5	1,168.6	2,503.4
Female	6.7	13.1	26.2	51.0	94.5	182.5	341.4	701.0	2,121.0
Non-Hispanic black	27.5	48.9	81.1	143.0	221.1	341.4	547.6	855.6	1,761.4
Male	39.2	70.5	114.5	191.9	290.3	461.7	691.6	1,056.4	1,768.3
Female	17.4	30.5	53.8	104.7	170.0	261.3	461.7	753.4	1,758.9

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Other heart diseases (I26-I51)									
Non-Hispanic 5/	18.7	26.9	41.1	63.8	103.1	168.7	292.4	545.3	1,497.8
Male	24.0	33.8	51.2	79.7	124.2	205.6	351.5	632.2	1,499.0
Female	13.4	20.3	31.5	49.3	84.9	139.2	251.1	493.4	1,497.3
Non-Hispanic white	15.4	23.0	35.7	58.7	96.3	162.8	286.7	547.4	1,533.2
Male	20.3	29.7	44.8	73.9	117.3	198.2	344.9	632.2	1,530.7
Female	10.6	16.5	27.1	44.6	77.7	134.0	245.4	496.2	1,534.2
Non-Hispanic black	44.0	62.3	93.6	121.1	182.4	258.0	408.2	608.1	1,292.9
Male	54.7	75.2	120.1	154.0	216.3	322.3	501.0	753.1	1,362.9
Female	34.8	51.3	71.9	95.3	157.3	215.2	352.8	534.3	1,267.0

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Heart failure (I50)									
Non-Hispanic 5/	1.8	3.4	6.8	13.3	24.4	46.8	94.5	202.2	685.6
Male	2.3	4.2	8.4	15.3	28.5	55.8	110.7	230.4	658.0
Female	1.3	2.7	5.2	11.5	20.8	39.6	83.2	185.4	697.0
Non-Hispanic white	1.4	2.8	5.9	12.2	22.9	45.5	93.9	204.3	706.8
Male	1.9	3.5	7.5	14.4	26.9	54.4	110.5	232.1	679.1
Female	0.8	2.1	4.3	10.2	19.4	38.2	82.2	187.4	718.4
Non-Hispanic black	4.7	8.9	16.0	25.1	42.1	72.3	122.2	216.6	551.2
Male	5.2	10.7	19.3	26.8	50.5	87.1	141.7	260.8	547.5
Female	4.2	7.3	13.3	23.8	35.8	62.4	110.6	194.1	552.6

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	--------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

Non-Hispanic 5/	16.2	22.7	33.4	49.2	77.1	119.3	194.6	338.8	807.8
Male	20.8	28.6	41.7	62.8	93.9	146.8	236.4	396.2	835.3
Female	11.7	17.1	25.6	36.9	62.6	97.4	165.3	304.4	796.3
Non-Hispanic white	13.4	19.6	29.1	45.3	72.0	114.9	189.5	338.8	821.8
Male	17.6	25.4	36.4	58.0	88.9	141.1	230.1	394.4	845.6
Female	9.2	13.9	22.1	33.5	57.0	93.6	160.7	305.2	811.9
Non-Hispanic black	37.7	51.0	75.4	93.3	137.0	182.1	280.5	386.8	737.9
Male	46.8	61.6	98.2	123.3	161.0	229.6	352.7	484.9	813.1
Female	29.7	42.1	56.7	69.7	119.3	150.5	237.4	336.9	710.1

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	--------------

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

Non-Hispanic 5/	1.9	3.0	4.5	7.5	12.3	18.7	35.2	65.0	172.0
Male	2.3	3.8	5.3	8.3	13.8	19.6	35.6	60.6	151.1
Female	1.4	2.3	3.8	6.8	10.9	18.0	35.0	67.6	180.7
Non-Hispanic white	1.0	1.8	2.9	5.2	8.9	14.7	30.3	58.1	162.8
Male	1.3	2.2	3.5	5.9	10.2	15.6	30.8	53.4	142.1
Female	0.7	1.3	2.3	4.5	7.8	14.0	29.9	60.9	171.5
Non-Hispanic black	7.9	12.9	18.3	27.6	41.2	57.4	90.7	150.1	291.5
Male	10.2	16.7	21.9	30.9	49.5	63.4	95.8	161.1	277.2
Female	5.9	9.6	15.4	25.0	35.0	53.4	87.7	144.5	296.9

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Cerebrovascular diseases (I60-I69)									
All origins 3/	12.1	18.5	28.8	49.6	85.7	164.3	320.2	622.4	1,500.2
Male	13.2	20.5	32.8	57.2	98.5	187.9	349.0	654.6	1,380.2
Female	10.9	16.5	25.1	42.7	74.7	145.6	300.0	603.2	1,550.4
Hispanic	12.4	20.3	28.6	44.5	74.7	145.5	271.5	463.4	1,042.9
Male	14.6	24.1	34.2	56.2	95.3	168.4	311.0	529.8	1,005.6
Female	10.2	16.7	23.5	34.3	58.0	128.1	243.5	422.2	1,061.2
Mexican	12.7	20.0	26.7	50.3	83.5	167.6	305.2	529.7	*
Male	14.6	21.6	30.7	67.3	104.9	202.4	*	*	*
Female	10.8	18.2	22.7	36.9	63.8	140.9	273.4	*	*
Puerto Rican	*	21.8	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Cuban	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	12.1	23.0	25.4	34.8	68.4	152.0	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	19.3	*	*	*	*	*	*	*

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Cerebrovascular diseases (I60-I69)									
Non-Hispanic 5/	12.0	18.2	28.8	49.8	86.3	165.1	321.7	627.7	1,514.0
Male	13.1	20.2	32.5	56.9	98.5	188.7	350.0	658.1	1,392.9
Female	10.9	16.4	25.2	43.2	75.8	146.3	302.0	609.6	1,564.4
Non-Hispanic white	8.8	13.0	23.4	41.1	74.9	153.2	310.4	621.9	1,525.0
Male	9.6	14.4	26.0	46.9	85.6	174.7	337.7	651.5	1,399.0
Female	8.0	11.6	20.8	35.7	65.4	135.7	291.1	604.0	1,577.4
Non-Hispanic black	33.1	56.2	71.9	119.3	181.5	281.4	465.4	725.7	1,465.5
Male	37.3	65.4	89.2	145.9	222.5	340.9	530.4	776.0	1,359.4
Female	29.6	48.4	57.8	98.4	151.2	241.8	426.6	700.1	1,504.8

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Atherosclerosis (I70)									
Non-Hispanic 5/	0.3	0.6	1.4	2.6	5.1	10.0	22.0	49.6	180.5
Male	0.5	0.8	1.7	3.4	6.4	12.4	25.0	53.6	168.2
Female	0.2	0.5	1.1	1.9	4.0	8.2	19.9	47.3	185.5
Non-Hispanic white	0.3	0.6	1.4	2.4	5.1	10.0	22.4	50.8	187.4
Male	0.4	0.7	1.7	3.2	6.4	12.4	25.1	54.4	175.2
Female	*	0.5	1.1	1.6	4.0	8.1	20.4	48.6	192.6
Non-Hispanic black	*	1.0	2.0	5.2	6.3	12.1	23.2	42.5	126.1
Male	*	*	*	6.1	7.8	15.1	30.8	48.2	113.2
Female	*	*	*	4.4	5.2	10.1	18.7	39.6	130.9

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Other diseases of circulatory system (I71-I78)									
Non-Hispanic 5/	2.1	3.3	6.2	12.5	23.0	37.7	61.7	89.6	160.1
Male	2.9	4.4	8.5	17.3	31.6	52.6	82.2	124.9	200.5
Female	1.2	2.2	3.9	8.1	15.6	25.9	47.3	68.4	143.3
Non-Hispanic white	1.8	2.9	5.7	12.1	23.3	38.0	61.8	90.0	160.1
Male	2.5	3.8	8.2	16.9	32.5	53.3	83.4	126.6	204.7
Female	1.1	1.9	3.4	7.7	15.3	25.5	46.5	67.9	141.6
Non-Hispanic black	4.1	6.6	10.9	18.2	23.2	43.5	69.8	97.1	175.7
Male	5.7	9.6	13.2	24.0	28.1	56.5	78.7	129.2	168.6
Female	2.7	4.1	9.0	13.6	19.6	34.9	64.5	80.8	178.4

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Aortic aneurysm and dissection (I71)									
Non-Hispanic 5/	1.4	2.1	3.9	8.1	15.5	24.5	39.8	56.4	79.0
Male	2.2	3.2	6.0	12.4	22.8	37.0	57.2	84.2	117.0
Female	0.7	0.9	1.9	4.1	9.2	14.6	27.7	39.8	63.2
Non-Hispanic white	1.2	1.7	3.7	8.1	16.1	25.5	40.7	58.1	81.3
Male	1.9	2.8	5.9	12.3	23.9	38.5	58.7	87.4	122.9
Female	0.5	0.7	1.6	4.1	9.2	14.9	27.8	40.5	64.0
Non-Hispanic black	2.8	4.5	6.2	9.5	11.6	18.6	34.5	43.3	55.9
Male	4.1	7.0	8.5	14.8	14.9	26.5	42.7	55.6	49.7
Female	1.7	2.3	4.4	5.3	9.2	13.3	29.7	37.1	58.2

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	--------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

Non-Hispanic 5/	0.6	1.2	2.3	4.4	7.5	13.2	21.9	33.1	81.1
Male	0.7	1.2	2.5	4.9	8.8	15.6	25.0	40.7	83.5
Female	0.6	1.2	2.1	4.0	6.4	11.3	19.6	28.6	80.1
Non-Hispanic white	0.6	1.1	2.1	4.0	7.3	12.5	21.1	31.9	78.8
Male	0.6	1.0	2.3	4.5	8.6	14.8	24.6	39.2	81.8
Female	0.5	1.2	1.8	3.6	6.1	10.6	18.7	27.5	77.6
Non-Hispanic black	1.3	2.2	4.6	8.7	11.6	24.9	35.3	53.8	119.9
Male	*	2.6	4.7	9.2	13.3	30.0	36.0	73.6	119.0
Female	*	*	4.6	8.4	10.4	21.6	34.8	43.7	120.2

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	--------------

Other disorders of circulatory system (I80-I99)

Non-Hispanic 5/	1.1	1.3	2.0	2.7	3.8	5.4	7.8	11.9	25.8
Male	1.2	1.5	2.3	3.0	3.9	6.1	8.3	12.0	21.6
Female	0.9	1.0	1.8	2.4	3.8	4.9	7.4	11.8	27.6
Non-Hispanic white	0.9	1.1	1.8	2.4	3.5	5.1	7.3	11.8	25.7
Male	1.2	1.3	2.1	2.5	3.6	5.7	7.8	11.9	21.2
Female	0.7	1.0	1.6	2.3	3.5	4.7	6.9	11.8	27.6
Non-Hispanic black	2.2	2.4	4.0	6.2	7.1	8.9	14.0	14.9	29.7
Male	1.8	3.1	4.8	8.8	8.1	10.7	16.1	*	28.9
Female	2.5	*	3.3	4.1	6.3	7.8	12.7	14.6	29.9

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Influenza and pneumonia (J10-J18)									
Non-Hispanic 5/	4.0	5.4	8.2	14.0	24.4	48.8	101.0	216.7	684.8
Male	4.8	6.7	9.9	17.2	30.5	61.1	130.6	274.2	808.2
Female	3.2	4.1	6.5	11.1	19.1	39.0	80.2	182.3	633.5
Non-Hispanic white	3.3	4.8	7.5	12.9	23.2	47.4	98.2	216.2	693.9
Male	4.0	5.7	8.9	15.3	28.3	58.5	125.1	271.6	813.9
Female	2.6	3.9	6.1	10.6	18.7	38.3	79.2	182.8	644.0
Non-Hispanic black	9.2	10.8	15.4	23.7	34.9	66.2	133.9	224.9	583.1
Male	11.6	15.9	20.8	34.3	50.8	95.4	203.4	309.9	718.4
Female	7.1	6.6	11.1	15.4	23.1	46.8	92.5	181.6	533.0

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Pneumonia (J12-J18)									
Non-Hispanic 5/	4.0	5.4	8.1	13.9	24.3	48.7	100.8	216.0	682.5
Male	4.8	6.7	9.8	17.1	30.4	60.9	130.4	273.6	805.9
Female	3.1	4.1	6.5	10.9	19.0	38.9	80.0	181.6	631.3
Non-Hispanic white	3.3	4.7	7.4	12.8	23.1	47.3	98.0	215.5	691.5
Male	4.0	5.6	8.8	15.3	28.3	58.4	124.9	270.9	811.5
Female	2.6	3.9	6.0	10.5	18.6	38.2	78.9	182.1	641.7
Non-Hispanic black	9.2	10.8	15.4	23.5	34.6	66.2	133.9	223.8	581.5
Male	11.6	15.9	20.6	33.9	50.3	95.4	203.4	309.9	716.1
Female	7.1	6.6	11.1	15.4	23.1	46.8	92.5	179.9	531.7

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	--------------

Chronic lower respiratory diseases (J40-J47)

Non-Hispanic 5/	5.7	12.4	30.9	65.9	129.7	223.3	337.5	461.7	649.4
Male	6.1	12.9	32.9	69.9	142.2	261.1	416.4	594.6	924.8
Female	5.3	12.0	29.0	62.3	118.9	193.3	282.4	382.2	534.8
Non-Hispanic white	5.4	12.3	31.8	69.9	137.9	236.4	355.6	483.3	676.9
Male	5.9	12.5	33.4	72.5	147.9	270.1	429.6	611.8	951.5
Female	4.8	12.1	30.3	67.4	129.1	209.0	303.2	405.8	562.7
Non-Hispanic black	9.3	16.8	32.3	52.7	95.9	150.7	208.7	279.2	379.3
Male	9.1	19.0	37.8	66.7	129.2	224.1	333.2	475.9	682.6
Female	9.5	14.9	27.8	41.7	71.4	101.9	134.4	179.1	266.9

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Emphysema (J43)									
Non-Hispanic 5/	0.7	1.8	5.3	10.7	20.8	34.6	48.2	56.6	62.8
Male	0.9	2.1	6.0	11.4	23.4	41.8	60.6	76.0	102.7
Female	0.5	1.5	4.5	10.1	18.6	28.9	39.6	44.9	46.3
Non-Hispanic white	0.8	1.8	5.5	11.6	22.7	36.9	51.4	59.9	65.9
Male	1.0	2.0	6.1	12.0	25.1	43.3	63.4	79.2	106.6
Female	0.5	1.5	4.9	11.3	20.6	31.7	42.9	48.3	48.9
Non-Hispanic black	*	2.4	5.0	6.3	11.7	20.4	21.8	26.5	35.9
Male	*	3.0	6.9	9.2	15.9	32.4	34.6	49.9	69.3
Female	*	*	3.4	4.1	8.7	12.4	14.1	14.6	23.5

ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX: UNITED STATES, 2001

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Asthma (J45-J46)									
Non-Hispanic 5/	1.4	1.7	1.8	2.6	3.0	3.9	5.0	8.1	16.1
Male	1.0	1.3	1.1	1.9	2.3	2.9	3.7	5.6	12.2
Female	1.7	2.1	2.5	3.2	3.6	4.7	6.0	9.6	17.7
Non-Hispanic white	0.9	1.3	1.3	1.8	2.6	3.3	4.5	7.7	15.7
Male	0.7	0.9	0.8	1.2	1.9	2.3	3.2	5.0	11.4
Female	1.2	1.7	1.8	2.4	3.2	4.1	5.5	9.3	17.5
Non-Hispanic black	4.4	5.0	6.1	7.9	6.3	9.5	10.3	11.3	16.5
Male	3.8	4.6	3.7	6.1	*	9.6	*	*	*
Female	5.0	5.3	8.0	9.4	7.5	9.4	11.6	12.1	17.1

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	--------------

Other chronic lower respiratory diseases (J44,J47)

Non-Hispanic 5/	3.5	8.8	23.6	52.3	105.2	184.0	282.5	394.4	561.3
Male	4.0	9.4	25.5	56.2	115.6	215.3	350.0	510.4	800.6
Female	3.0	8.2	21.9	48.7	96.2	159.0	235.3	325.1	461.7
Non-Hispanic white	3.6	9.1	24.8	56.0	112.0	195.4	298.0	413.0	585.7
Male	4.1	9.5	26.2	58.9	120.0	223.5	361.0	525.0	823.9
Female	3.0	8.7	23.5	53.4	104.9	172.5	253.3	345.3	486.6
Non-Hispanic black	4.2	9.4	21.2	38.2	77.1	119.8	175.2	240.6	323.4
Male	4.5	11.3	27.2	51.0	107.6	180.8	288.7	414.6	592.5
Female	3.9	7.8	16.2	28.1	54.6	79.1	107.5	152.0	223.7

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Pneumonitis due to solids and liquids (J69)									
Non-Hispanic 5/	0.7	1.3	1.8	3.7	6.3	14.4	30.9	68.8	190.0
Male	0.8	1.7	2.3	4.4	8.4	19.5	41.7	99.6	271.6
Female	0.6	0.9	1.4	3.1	4.5	10.4	23.4	50.4	156.0
Non-Hispanic white	0.7	1.2	1.8	3.6	6.1	13.9	30.7	69.0	191.2
Male	0.9	1.6	2.3	4.1	8.1	18.8	41.3	100.4	276.9
Female	0.5	0.9	1.4	3.1	4.3	9.8	23.2	50.1	155.5
Non-Hispanic black	1.0	2.5	2.5	6.1	9.5	22.7	39.2	76.2	187.3
Male	*	3.5	3.2	8.4	12.8	31.3	56.9	107.9	243.7
Female	*	*	*	4.3	7.1	17.0	28.6	60.0	166.4

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	--------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Non-Hispanic 5/	2.6	4.4	7.2	12.8	23.0	38.2	62.1	89.4	142.7
Male	2.9	4.9	7.6	15.1	27.3	47.2	76.5	110.0	169.1
Female	2.4	3.9	6.8	10.7	19.2	31.0	52.0	77.1	131.8
Non-Hispanic white	2.2	3.8	6.9	12.5	23.2	39.0	63.8	92.2	145.9
Male	2.5	4.4	7.5	14.5	27.7	48.6	78.9	113.3	173.4
Female	2.0	3.3	6.3	10.6	19.1	31.2	53.1	79.4	134.5
Non-Hispanic black	5.9	9.5	11.2	17.4	24.3	36.1	50.3	67.0	110.8
Male	6.4	9.6	10.8	22.2	27.9	39.6	59.7	85.0	131.7
Female	5.5	9.4	11.5	13.6	21.7	33.7	44.7	57.9	103.1

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Peptic ulcer (K25-K28)									
Non-Hispanic 5/	0.7	0.8	1.3	2.2	3.1	4.8	8.3	16.0	33.6
Male	0.9	1.1	1.8	2.8	4.1	5.6	10.2	16.8	35.5
Female	0.5	0.6	0.8	1.6	2.3	4.3	7.0	15.5	32.8
Non-Hispanic white	0.6	0.7	1.1	2.1	3.0	4.8	8.3	16.3	34.7
Male	0.8	0.9	1.5	2.7	3.9	5.3	9.9	16.9	36.3
Female	0.5	0.6	0.8	1.5	2.3	4.5	7.1	15.9	34.0
Non-Hispanic black	1.1	1.6	2.5	3.1	3.5	5.6	8.0	10.5	20.9
Male	1.9	2.8	4.0	4.6	5.2	9.3	10.9	*	24.3
Female	*	*	*	*	*	*	6.2	9.2	19.7

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	--------------

Chronic liver disease and cirrhosis (K70,K73-K74)

Non-Hispanic 5/	16.3	19.0	19.8	23.2	27.2	29.4	29.7	27.4	21.6
Male	23.4	28.0	29.1	32.4	36.8	38.6	35.4	34.1	31.4
Female	9.3	10.3	11.1	14.8	18.8	22.0	25.7	23.4	17.5
Non-Hispanic white	16.0	18.1	19.1	23.5	27.7	30.2	30.9	28.6	22.5
Male	23.2	26.7	27.7	32.3	36.8	39.6	36.6	35.8	32.9
Female	9.0	9.7	10.8	15.3	19.6	22.6	26.8	24.3	18.2
Non-Hispanic black	19.4	27.2	28.3	23.8	27.1	23.0	20.6	14.6	10.6
Male	28.9	42.8	44.8	38.5	43.8	34.4	27.0	*	*
Female	11.3	14.0	14.8	12.3	14.8	15.4	16.7	14.6	9.8

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Alcoholic liver disease (K70)									
Non-Hispanic 5/	9.6	10.6	10.3	11.1	10.3	8.2	6.3	3.6	2.2
Male	14.0	15.7	16.3	17.3	15.9	13.6	11.0	6.3	5.1
Female	5.3	5.8	4.6	5.5	5.4	4.0	3.0	1.9	1.0
Non-Hispanic white	9.7	10.1	9.9	11.3	10.1	8.4	6.4	3.6	2.1
Male	14.3	15.1	15.6	17.4	15.4	13.6	11.2	6.4	4.8
Female	5.2	5.3	4.5	5.6	5.4	4.1	3.0	1.9	1.0
Non-Hispanic black	9.1	14.9	14.4	11.6	13.1	8.0	5.7	*	*
Male	13.3	22.5	24.0	18.8	23.4	15.5	*	*	*
Female	5.6	8.4	6.5	5.9	5.4	*	*	*	*

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	--------------

Other chronic liver disease and cirrhosis (K73-K74)

Non-Hispanic 5/	6.7	8.4	9.6	12.1	16.9	21.1	23.4	23.8	19.4
Male	9.5	12.4	12.9	15.1	20.8	25.0	24.4	27.7	26.4
Female	4.0	4.6	6.5	9.3	13.5	18.0	22.7	21.5	16.5
Non-Hispanic white	6.3	8.0	9.2	12.3	17.5	21.9	24.5	25.0	20.4
Male	8.9	11.6	12.1	14.9	21.4	25.9	25.4	29.3	28.1
Female	3.8	4.4	6.3	9.8	14.2	18.5	23.8	22.4	17.2
Non-Hispanic black	10.3	12.3	13.9	12.2	14.1	15.0	14.9	11.6	7.8
Male	15.6	20.2	20.8	19.6	20.3	18.9	18.0	*	*
Female	5.7	5.6	8.3	6.4	9.4	12.4	13.0	12.9	*

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Cholelithiasis and other disorders of gallbladder (K80-K82)									
Non-Hispanic 5/	0.3	0.4	0.6	1.3	2.0	3.6	5.8	9.8	23.4
Male	0.4	0.4	0.8	1.5	2.1	4.2	7.5	11.9	26.6
Female	0.3	0.3	0.4	1.2	1.8	3.1	4.6	8.6	22.1
Non-Hispanic white	0.3	0.3	0.5	1.2	1.7	3.4	5.7	9.6	23.6
Male	0.3	0.4	0.7	1.4	1.8	4.1	7.5	11.8	26.4
Female	*	*	0.4	1.1	1.6	2.9	4.4	8.3	22.4
Non-Hispanic black	*	*	1.5	2.0	3.5	6.3	6.2	11.6	20.0
Male	*	*	*	*	*	*	*	*	27.7
Female	*	*	*	*	*	6.4	7.1	12.9	17.1

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	--------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Non-Hispanic 5/	3.7	5.8	9.8	17.0	31.5	49.1	82.0	136.3	289.6
Male	4.5	6.8	10.8	18.1	36.5	57.0	100.3	178.2	397.6
Female	2.9	4.8	8.8	16.0	27.1	42.8	69.1	111.2	244.6
Non-Hispanic white	2.3	3.9	6.8	12.5	25.0	42.4	74.4	128.0	279.7
Male	2.7	4.8	7.8	13.9	29.9	50.9	94.0	170.0	392.6
Female	1.8	3.1	5.9	11.2	20.6	35.5	60.5	102.6	232.7
Non-Hispanic black	12.9	20.3	33.3	54.2	90.4	121.3	174.0	250.8	432.9
Male	17.0	23.6	38.6	56.2	103.4	132.6	190.6	312.3	517.4
Female	9.3	17.5	28.9	52.6	80.8	113.8	164.1	219.5	401.7

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Renal failure (N17-N19)									
Non-Hispanic 5/	3.6	5.6	9.6	16.7	30.8	48.2	80.6	134.3	285.9
Male	4.4	6.6	10.6	17.8	35.9	55.8	98.7	175.3	392.7
Female	2.8	4.7	8.6	15.8	26.5	42.1	68.0	109.7	241.5
Non-Hispanic white	2.2	3.8	6.7	12.3	24.4	41.4	73.1	126.1	276.1
Male	2.7	4.6	7.6	13.7	29.4	49.7	92.4	167.2	387.6
Female	1.8	3.0	5.8	11.1	20.0	34.7	59.4	101.3	229.7
Non-Hispanic black	12.6	19.8	33.0	53.6	88.9	120.5	172.2	248.1	428.9
Male	16.6	22.8	38.2	55.4	101.4	131.5	189.1	309.9	514.0
Female	9.2	17.2	28.8	52.2	79.6	113.1	162.1	216.6	397.4

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	--------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Non-Hispanic 5/	9.0	8.8	8.5	10.0	12.9	18.2	30.4	55.3	209.6
Male	11.6	11.8	11.1	13.2	15.7	21.1	33.8	61.6	178.2
Female	6.4	6.0	6.1	7.1	10.5	15.8	28.0	51.6	222.7
Non-Hispanic white	8.7	8.3	7.9	9.5	12.1	17.5	30.1	55.0	213.3
Male	11.2	11.1	10.0	12.1	14.4	20.5	33.9	61.6	180.4
Female	6.2	5.5	5.9	7.0	10.0	15.1	27.4	51.0	227.0
Non-Hispanic black	13.2	14.9	14.6	15.7	21.8	28.1	39.2	67.6	194.2
Male	17.4	19.2	22.2	25.9	29.7	32.0	41.2	71.1	186.0
Female	9.6	11.2	8.4	7.7	16.0	25.5	37.9	65.8	197.2

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION

BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	--------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

All origins 3/	32.1	40.9	53.2	76.4	112.0	172.3	295.5	539.3	1,476.7
Male	38.8	49.1	59.9	85.8	124.3	186.5	307.6	539.6	1,271.6
Female	25.6	33.0	46.9	67.9	101.4	161.0	287.0	539.1	1,562.6
Hispanic	22.8	30.7	42.2	59.7	83.9	136.4	209.8	396.3	991.0
Male	30.6	40.0	51.7	69.0	96.0	144.6	205.3	403.8	939.0
Female	14.9	21.9	33.5	51.7	74.0	130.2	213.0	391.7	1,016.5
Mexican	21.1	26.5	43.1	59.3	86.7	140.1	222.0	431.8	*
Male	27.5	32.2	47.8	67.3	91.2	148.7	*	*	*
Female	14.6	20.2	38.3	53.0	82.6	133.4	228.8	*	*
Puerto Rican	41.3	48.7	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Cuban	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	20.3	36.5	30.6	66.3	91.2	157.8	*	*	*
Male	28.4	51.7	*	*	*	*	*	*	*
Female	*	23.9	*	*	*	*	*	*	*

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	--------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Non-Hispanic 5/	32.9	41.5	53.8	77.5	113.6	174.0	299.1	543.9	1,491.3
Male	39.4	49.5	60.0	86.7	125.7	188.5	311.8	544.1	1,282.3
Female	26.6	33.9	47.9	69.1	103.1	162.4	290.1	543.7	1,578.3
Non-Hispanic white	29.2	37.5	49.8	73.3	110.2	170.7	296.0	545.5	1,512.9
Male	35.5	45.0	55.1	82.2	121.0	183.6	306.0	544.6	1,293.1
Female	23.0	30.2	44.7	65.1	100.7	160.1	288.9	546.0	1,604.4
Non-Hispanic black	64.1	80.9	97.2	124.2	165.8	239.3	384.5	619.4	1,431.5
Male	75.8	97.7	116.3	144.0	199.8	276.5	449.4	669.7	1,329.4
Female	54.1	66.6	81.6	108.6	140.6	214.5	345.7	593.9	1,469.3

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85+ YEARS

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

All origins 3/	36.0	31.9	29.3	31.5	36.8	49.3	80.4	130.4	276.4
Male	54.0	46.9	41.8	44.5	49.9	65.6	108.3	173.2	358.5
Female	18.5	17.5	17.6	19.6	25.5	36.3	60.9	104.8	242.0
Hispanic	32.6	30.5	29.7	31.6	32.5	49.3	62.9	92.9	166.3
Male	50.0	47.3	42.6	48.3	46.1	66.3	92.8	128.6	198.1
Female	15.2	14.6	18.1	17.2	21.5	36.3	41.6	70.8	150.8
Mexican	30.7	27.7	27.8	37.1	33.0	58.6	63.9	*	*
Male	48.3	42.6	40.5	62.5	44.1	77.4	*	*	*
Female	12.7	11.5	15.0	17.0	22.8	44.0	*	*	*
Puerto Rican	41.3	32.5	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Cuban	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	37.1	40.0	34.6	29.8	*	*	*	*	*
Male	55.9	61.0	*	*	*	*	*	*	*
Female	19.0	22.8	31.1	*	*	*	*	*	*

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	--------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Non-Hispanic 5/	36.0	31.8	29.0	31.2	36.9	49.1	81.1	131.5	279.8
Male	54.0	46.5	41.4	43.9	50.0	65.2	108.8	174.4	364.5
Female	18.7	17.8	17.4	19.7	25.6	36.2	61.8	105.8	244.6
Non-Hispanic white	34.7	30.8	28.1	30.1	36.5	48.5	81.9	134.4	289.3
Male	51.3	44.3	39.2	41.8	48.8	64.2	108.7	176.3	374.7
Female	18.3	17.5	17.4	19.4	25.6	35.9	63.0	109.1	253.8
Non-Hispanic black	50.8	45.7	41.2	43.9	45.9	58.0	79.2	105.7	183.9
Male	83.0	73.2	67.0	69.8	69.8	81.6	121.4	167.6	256.4
Female	23.0	22.3	20.0	23.5	28.3	42.2	54.0	74.1	157.0

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Transport accidents (V01-V99,Y85)									
Non-Hispanic 5/	16.2	15.4	15.0	15.0	16.3	19.2	26.0	30.1	29.0
Male	24.0	22.3	21.3	21.3	21.8	26.4	35.7	44.6	52.9
Female	8.6	8.9	9.0	9.3	11.5	13.5	19.3	21.4	19.1
Non-Hispanic white	15.7	15.4	14.8	14.7	16.2	19.2	26.5	30.8	29.9
Male	23.1	22.2	20.8	20.7	21.4	26.2	35.7	45.7	54.3
Female	8.5	8.8	9.0	9.1	11.6	13.5	20.0	21.9	19.8
Non-Hispanic black	20.7	17.2	18.1	18.9	18.9	18.7	22.3	19.9	17.8
Male	34.5	26.1	29.3	31.3	28.9	25.5	39.8	31.9	35.8
Female	8.8	9.6	8.8	9.2	11.5	14.2	11.9	13.7	11.1

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)									
Non-Hispanic 5/	14.5	13.8	13.6	13.5	14.7	17.7	24.6	28.7	28.0
Male	21.2	19.5	18.9	18.6	19.0	23.8	33.3	42.4	51.6
Female	8.0	8.3	8.6	8.8	11.0	12.9	18.4	20.4	18.2
Non-Hispanic white	14.1	13.8	13.4	13.1	14.7	17.6	25.0	29.4	29.0
Male	20.3	19.4	18.5	17.8	18.7	23.5	33.3	43.5	53.1
Female	8.0	8.2	8.6	8.8	11.1	12.8	19.2	21.0	19.0
Non-Hispanic black	18.5	15.7	16.2	17.2	16.8	18.0	20.4	19.0	16.9
Male	30.6	23.7	25.3	28.8	24.5	23.8	37.0	31.1	35.8
Female	8.1	8.9	8.7	8.1	11.2	14.2	10.5	12.9	9.8

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Nontransport accidents (W00-X59,Y86)									
Non-Hispanic 5/	19.8	16.4	14.1	16.2	20.6	29.8	55.1	101.4	250.8
Male	29.9	24.2	20.1	22.6	28.1	38.8	73.1	129.8	311.6
Female	10.1	8.9	8.4	10.5	14.1	22.7	42.5	84.4	225.5
Non-Hispanic white	19.0	15.3	13.3	15.5	20.3	29.4	55.4	103.6	259.4
Male	28.2	22.0	18.4	21.1	27.5	38.0	73.0	130.7	320.4
Female	9.8	8.7	8.4	10.3	14.0	22.3	43.0	87.2	234.0
Non-Hispanic black	30.1	28.5	23.1	24.9	27.0	39.2	56.9	85.8	166.1
Male	48.6	47.1	37.7	38.5	40.9	56.1	81.5	135.7	220.6
Female	14.2	12.7	11.2	14.3	16.7	28.0	42.2	60.4	145.9

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Falls (W00-W19)									
Non-Hispanic 5/	2.4	2.8	3.2	4.8	7.9	12.3	25.5	50.4	123.1
Male	4.0	4.5	4.9	6.9	10.7	15.8	34.6	64.9	159.0
Female	0.9	1.1	1.6	2.9	5.5	9.4	19.2	41.7	108.1
Non-Hispanic white	2.4	2.8	3.1	4.9	8.1	12.7	26.6	53.1	129.9
Male	3.8	4.4	4.7	6.9	10.9	16.3	35.7	67.5	166.0
Female	1.0	1.2	1.6	3.0	5.6	9.7	20.1	44.4	114.8
Non-Hispanic black	2.6	3.4	4.0	4.9	8.0	9.8	16.7	21.8	44.0
Male	4.9	6.1	6.4	7.5	12.0	13.4	24.6	36.8	68.1
Female	*	*	*	*	5.0	7.3	11.9	14.2	35.1

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Accidental exposure to smoke, fire and flames (X00-X09)									
Non-Hispanic 5/	1.1	1.2	1.6	1.5	2.1	2.6	3.1	3.9	5.7
Male	1.5	1.6	2.1	1.8	2.6	3.1	3.5	5.2	8.3
Female	0.6	0.7	1.1	1.2	1.6	2.2	2.8	3.1	4.6
Non-Hispanic white	0.9	1.0	1.3	1.3	1.7	2.2	2.6	3.3	4.7
Male	1.3	1.3	1.8	1.5	2.2	2.5	2.8	4.5	6.9
Female	0.6	0.7	0.9	1.1	1.4	1.9	2.5	2.6	3.8
Non-Hispanic black	2.3	2.6	3.8	3.3	5.1	7.3	8.7	11.9	17.8
Male	3.6	4.4	5.2	5.0	7.3	11.4	12.3	*	26.6
Female	*	*	2.8	*	*	4.6	6.5	10.4	14.5

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	--------------

Accidental poisoning and exposure to noxious substances (X40-X49)

Non-Hispanic 5/	11.0	6.9	3.6	2.4	1.7	1.5	2.1	2.4	3.6
Male	15.7	9.6	4.6	2.9	1.9	2.0	2.7	2.9	3.9
Female	6.5	4.3	2.6	1.9	1.5	1.0	1.7	2.2	3.4
Non-Hispanic white	10.5	6.0	3.2	2.1	1.6	1.4	2.0	2.3	3.4
Male	14.7	8.0	3.8	2.3	1.8	1.9	2.4	2.7	3.8
Female	6.3	4.0	2.7	1.9	1.5	0.9	1.8	2.0	3.2
Non-Hispanic black	18.1	15.6	7.3	5.2	2.5	*	*	*	6.6
Male	28.4	25.2	12.6	9.0	*	*	*	*	*
Female	9.2	7.5	3.0	*	*	*	*	*	*

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	--------------

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

Non-Hispanic 5/	4.0	4.5	4.6	6.4	7.9	12.4	22.9	42.9	116.6
Male	6.4	6.8	6.7	9.1	11.0	15.9	29.8	53.6	136.5
Female	1.6	2.3	2.6	3.9	5.2	9.6	18.1	36.5	108.4
Non-Hispanic white	3.8	4.4	4.5	6.0	7.8	12.0	22.7	43.2	119.5
Male	6.2	6.5	6.4	8.7	10.8	15.3	29.6	52.7	139.7
Female	1.5	2.4	2.7	3.6	5.1	9.2	17.9	37.4	111.2
Non-Hispanic black	5.6	6.0	6.1	10.5	10.5	18.9	27.5	44.7	96.5
Male	8.9	9.7	10.1	14.8	15.9	25.1	35.6	73.6	115.5
Female	2.7	2.8	2.8	7.1	6.5	14.7	22.6	30.0	89.4

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	--------------

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) 6/

Non-Hispanic 5/	16.5	15.1	14.5	12.3	13.1	14.2	16.9	19.1	17.5
Male	25.6	23.2	23.1	19.9	23.1	27.2	34.6	44.6	51.3
Female	7.8	7.4	6.5	5.5	4.3	3.8	4.4	3.8	3.5
Non-Hispanic white	18.8	17.0	16.0	13.6	14.2	15.2	18.2	20.3	18.7
Male	28.8	25.8	25.2	21.5	25.1	28.8	37.0	47.4	55.2
Female	8.8	8.3	7.2	6.2	4.6	4.1	4.8	3.9	3.5
Non-Hispanic black	6.4	6.0	5.7	4.4	4.6	6.6	5.1	6.3	*
Male	9.9	9.7	10.0	8.4	8.1	14.5	11.9	*	*
Female	3.4	2.8	*	*	*	*	*	*	*

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) /6									
Non-Hispanic 5/	6.8	5.3	4.3	3.3	3.1	2.5	2.5	2.4	2.4
Male	10.2	8.0	6.6	4.7	4.3	3.2	3.2	2.9	3.1
Female	3.5	2.7	2.2	2.1	2.1	1.9	2.0	2.2	2.1
Non-Hispanic white	5.1	4.1	3.7	2.8	2.5	2.1	2.1	2.1	2.2
Male	7.4	6.0	5.3	3.8	3.2	2.5	2.6	2.4	2.8
Female	2.9	2.3	2.0	1.8	1.9	1.7	1.7	1.9	2.0
Non-Hispanic black	18.2	13.9	10.0	7.5	7.6	6.6	7.1	6.3	*
Male	30.1	23.6	17.6	11.7	13.0	12.1	11.9	*	*
Female	7.8	5.7	3.8	4.1	*	*	*	*	*

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)									
Non-Hispanic 5/	2.6	2.0	1.5	1.4	1.1	0.9	0.9	0.7	*
Male	4.1	3.2	2.3	2.0	1.6	1.1	1.3	*	*
Female	1.1	0.9	0.8	0.8	0.7	0.7	0.6	*	*
Non-Hispanic white	1.7	1.4	1.2	1.2	1.0	0.7	0.7	0.6	*
Male	2.5	2.1	1.5	1.7	1.2	0.8	1.0	*	*
Female	0.9	0.8	0.8	0.8	0.8	0.6	0.5	*	*
Non-Hispanic black	8.5	6.6	4.4	2.9	2.8	*	*	*	*
Male	15.6	12.1	8.7	5.0	6.0	*	*	*	*
Female	2.4	2.0	*	*	*	*	*	*	*

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

45-49 YEARS 50-54 YEARS 55-59 YEARS 60-64 YEARS 65-69 YEARS 70-74 YEARS 75-79 YEARS 80-84 YEARS 85+ YEARS

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1) /6

Non-Hispanic 5/	4.2	3.3	2.8	2.0	1.9	1.6	1.6	1.7	2.1
Male	6.1	4.9	4.3	2.6	2.6	2.1	1.9	2.0	2.7
Female	2.4	1.7	1.4	1.3	1.4	1.2	1.4	1.6	1.8
Non-Hispanic white	3.4	2.7	2.5	1.6	1.5	1.3	1.3	1.5	1.9
Male	4.9	3.9	3.8	2.2	2.1	1.7	1.6	1.8	2.3
Female	2.0	1.5	1.2	1.0	1.1	1.1	1.2	1.2	1.7
Non-Hispanic black	9.6	7.3	5.7	4.5	4.9	4.1	4.8	*	*
Male	14.6	11.5	8.9	6.7	7.0	7.2	*	*	*
Female	5.4	3.7	3.0	*	*	*	*	*	*

[RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 2000 CENSUS, ESTIMATED AS OF JULY 1, 2001. RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS. FOR EXPLANATION OF ASTERISKS PRECEDING CAUSE-OF-DEATH CODES, SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001"]

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85+ YEARS
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	--------------

Complications of medical and surgical care (Y40-Y84,Y88)

Non-Hispanic 5/	0.7	0.8	1.3	1.9	2.6	3.7	5.9	8.3	12.9
Male	0.7	0.8	1.4	2.0	2.6	4.0	6.3	9.9	14.5
Female	0.6	0.8	1.2	1.8	2.5	3.5	5.6	7.4	12.2
Non-Hispanic white	0.6	0.7	1.2	1.7	2.4	3.3	5.8	8.2	12.2
Male	0.6	0.7	1.2	1.7	2.5	3.3	6.2	10.1	13.5
Female	0.5	0.7	1.2	1.6	2.3	3.2	5.5	7.1	11.6
Non-Hispanic black	1.6	1.6	2.8	4.2	4.6	8.5	8.1	11.0	23.7
Male	1.8	*	3.5	5.2	5.2	11.4	*	*	30.0
Female	*	*	*	3.5	4.2	6.7	8.2	11.2	21.4

* FIGURE DOES NOT MEET STANDARDS OF RELIABILITY OR PRECISION.

0.0 QUANTITY MORE THAN ZERO BUT LESS THAN 0.05.

... CATEGORY NOT APPLICABLE.

1/ FIGURES FOR AGE NOT STATED INCLUDED IN "ALL AGES" BUT NOT DISTRIBUTED AMONG AGE GROUPS.

2/ DEATH RATES FOR "UNDER 1 YEAR" (BASED ON POPULATION ESTIMATES) DIFFER FROM INFANT MORTALITY RATES (BASED ON LIVE BIRTHS).

3/ ALL ORIGINS INCLUDES ORIGIN NOT STATED; SPECIFIED ORIGINS EXCLUDE ORIGIN NOT STATED.

4/ INCLUDES CENTRAL AND SOUTH AMERICAN AND OTHER AND UNKNOWN HISPANIC.

5/ INCLUDES RACES OTHER THAN WHITE AND BLACK.

6/ FIGURES INCLUDE SEPTEMBER 11, 2001 RELATED DEATHS FOR WHICH DEATH CERTIFICATES WERE FILED AS OF 10/24/02; SEE [HTTP://WWW.CDC.GOV/NCHS](http://www.cdc.gov/nchs) AND REFER TO "TECHNICAL APPENDIX FROM VITAL STATISTICS OF THE UNITED STATES: MORTALITY, 2001."

SOURCE: CDC/NCHS, NATIONAL VITAL STATISTICS SYSTEM, MORTALITY.