

DESIGN SUMMARY: Variances will be computed using the Taylor  
 Linearization Method, Assuming a With Replacement (WR) Design

Sample Weight: WTMEC4YR  
 Stratification Variable(s): SDMVSTRA  
 Primary Sampling Unit: SDMVPSU

Number of observations read    : 19759      Weighted count :278652243  
 Observations in subpopulation : 9471      Weighted count :198419259  
 Denominator degrees of freedom : 29

Date: 12-13-2013

Page: 1  
 Table: 1

Variance Estimation Method: Taylor Series (WR)  
 For Subpopulation: RIDAGEYR >= 20  
 Prevalence of SPs with measured high blood pressure : NHANES 1999-2002  
 by: Variable, Gender, AGE GROUP, Race Ethnicity.

for: Variable = HBP: High BP, Gender = Total.

AGE GROUP	Race Ethnicity	Sample Size	Percent	SE
-----				
Total				
	Total	8960	29.1630	0.9817
	NH White	4439	29.6593	1.1555
	NH Black	1677	36.9562	1.5130
	Mexican American	2087	17.1412	1.2180
	Other	757	25.8660	2.2956
20-39	Total	3175	7.1830	0.8315
	NH White	1408	6.3050	0.8653
	NH Black	602	11.9044	1.3178
	Mexican American	825	5.0434	0.9021
	Other	340	8.8807	2.3944
40-59	Total	2692	29.7023	1.3820
	NH White	1287	27.4901	1.7061
	NH Black	555	46.4504	2.1302
	Mexican American	620	26.0275	1.9296
	Other	230	30.7528	4.3219
60+	Total	3093	66.7150	1.2708
	NH White	1744	65.3819	1.5779
	NH Black	520	81.4957	1.6193
	Mexican American	642	62.0765	1.4276
	Other	187	67.3857	5.0892
-----				

Variance Estimation Method: Taylor Series (WR)

For Subpopulation: RIDAGEYR >= 20

Prevalence of SPs with measured high blood pressure : NHANES 1999-2002

by: Variable, Gender, AGE GROUP, Race Ethnicity.

for: Variable = HBP: High BP, Gender = Male.

AGE GROUP	Race Ethnicity	Sample Size	Percent	SE
-----				
Total				
	Total	4228	27.2721	1.2213
	NH White	2143	27.7521	1.2987
	NH Black	800	35.3589	1.5435
	Mexican American	957	17.2746	1.6922
	Other	328	23.0655	2.8433
20-39				
	Total	1352	10.1333	1.3483
	NH White	595	9.3704	1.4354
	NH Black	265	15.4747	1.4174
	Mexican American	350	6.8777	1.6786
	Other	142	12.1588	3.8178
40-59				
	Total	1343	28.3341	1.9214
	NH White	668	26.8355	2.2446
	NH Black	283	41.5988	3.3480
	Mexican American	292	26.6817	2.5467
	Other	100	27.3367	5.6711
60+				
	Total	1533	59.4742	1.7058
	NH White	880	58.2755	1.9405
	NH Black	252	77.9679	1.8563
	Mexican American	315	57.2239	2.3411
	Other	86	55.0099	9.0828
-----				

Variance Estimation Method: Taylor Series (WR)

For Subpopulation: RIDAGEYR >= 20

Prevalence of SPs with measured high blood pressure : NHANES 1999-2002  
by: Variable, Gender, AGE GROUP, Race Ethnicity.

for: Variable = HBP: High BP, Gender = Female.

AGE GROUP Race Ethnicity	Sample Size	Percent	SE
-----			
Total			
Total	4732	30.9091	1.0690
NH White	2296	31.4636	1.3439
NH Black	877	38.2763	2.1365
Mexican American	1130	17.0021	1.5080
Other	429	28.1374	3.1949
20-39			
Total	1823	4.2766	0.6670
NH White	813	3.2251	0.6012
NH Black	337	8.9185	2.1830
Mexican American	475	3.0391	0.8401
Other	198	5.7442	2.7978
40-59			
Total	1349	31.0173	1.5526
NH White	619	28.1486	1.9675
NH Black	272	50.7866	3.3318
Mexican American	328	25.3509	2.4511
Other	130	33.2027	4.6175
60+			
Total	1560	72.2896	1.5304
NH White	864	71.0079	1.9252
NH Black	268	83.8735	2.0034
Mexican American	327	65.9937	1.8603
Other	101	75.2974	4.8470
-----			