

Significance test for difference between mean systolic blood pressure 4

The SURVEYREG Procedure

Regression Analysis for Dependent Variable BPXSAR

Estimated Regression Coefficients

| Parameter | Estimate | Standard Error | t Value | Pr > t |
|----------------|------------|-------------------|---------|---------|
| Intercept | 122.451286 | 0.53179576 | 230.26 | <.0001 |
| RIAGENDR Men | 1.357033 | 0.51399676 | 2.64 | 0.0132 |
| RIAGENDR Women | 0.000000 | 0.00000000 | . | . |

NOTE: The denominator degrees of freedom for the t tests is 29.

Matrix X'WX is singular and a generalized inverse was used to solve the normal equations. Estimates are not unique.

Significance test for difference between mean systolic blood pressure 5

The SURVEYREG Procedure

Subpop: age ge 20=1

Domain Regression Analysis for Variable BPXSAR

Estimated Regression Coefficients

| Parameter | Estimate | Standard Error | t Value | Pr > t |
|----------------|------------|----------------|---------|---------|
| Intercept | 122.451286 | 0.53179576 | 230.26 | <.0001 |
| RIAGENDR Men | 1.357033 | 0.51399676 | 2.64 | 0.0132 |
| RIAGENDR Women | 0.000000 | 0.00000000 | . | . |

NOTE: The denominator degrees of freedom for the t tests is 29.

Matrix X'WX is singular and a generalized inverse was used to solve the normal equations. Estimates are not unique.