



## Healthcare-associated Infection Surveillance Protocol for Urinary Tract Infection Events for Long-term Care Facilities

**Background:** The urinary tract is one of the most common sites of healthcare-associated infections, accounting for up to 20% of infections reported by long-term care facilities (LTCFs).<sup>1</sup> Risk factors for developing bacteriuria and UTI include age-related changes to the genitourinary tract, comorbid conditions resulting in neurogenic bladder, and instrumentation required to manage bladder voiding. The point prevalence of asymptomatic bacteriuria in LTCF residents can range from 20-50%. Although the incidence of symptomatic UTI is lower, it still comprises a significant proportion of infections manifesting in LTCF residents and results in a large amount of antibiotic use.<sup>5</sup>

Though the prevalence of indwelling urinary catheter use in LTCFs is lower than the acute care setting, catheter-associated UTI (CAUTI) can lead to complications such as cystitis, pyelonephritis, bacteremia, and septic shock. These complications can then lead to declined resident function and mobility, acute care hospitalizations, and increased mortality. Prevention of CAUTIs is discussed in the CDC/HICPAC document, *Guideline for Prevention of Catheter-associated Urinary Tract Infections*.<sup>2</sup>

Efforts to examine antibiotic use for UTI have demonstrated a discrepancy between the number UTI events identified through the application of evidence-based surveillance criteria and the numbers of clinically identified and treated UTI.<sup>3,4</sup> Consistent tracking and reporting of symptomatic UTIs using surveillance criteria will help identify opportunities to examine, understand, and address differences between surveillance events and clinically identified events.

### References:

1. Genao L, Buhr G. T. Urinary Tract Infections in Older Adults Residing in Long-Term Care Facilities. *Annals of Long-term Care*, vol. 20, no. 4, 2012, pp. 33-38.
2. Healthcare Infection Control Practices Advisory Committee (HICPAC) Approved Guidelines for the Prevention of Catheter-associated Urinary Tract Infections, 2009. Available at [www.cdc.gov/hicpac/pdf/CAUTI/CAUTIguideline2009final.pdf](http://www.cdc.gov/hicpac/pdf/CAUTI/CAUTIguideline2009final.pdf)
3. Juthani-Mehta M., et al. Diagnostic Accuracy of Criteria for Urinary Tract Infection in a Cohort of Nursing Home Residents. *Journal of the American Geriatrics Society*, vol. 55, 2007, pp. 1072-77.
4. Wang L., et al. Infection Rate and Colonization with Antibiotic-resistant Organisms in Skilled Nursing Facility Residents with Indwelling Devices. *European Journal of Clinical Microbiology & Infectious Diseases*, vol. 31, no. 8, 2012, pp. 1797-804.
5. Nace D. A., et al. Clinical Uncertainties in the Approach to Long Term Care Residents with Possible Urinary Tract Infection. *Journal of American Medical Directors Association*. vol. 15, no. 2014, 2014, pp. 133-39.



**Methods:** Facilities may choose to monitor urinary tract infections (UTIs) using healthcare-associated infection (HAI) surveillance. This surveillance method incorporates the use of laboratory data and clinical evaluation of the resident for signs and/or symptoms to monitor for catheter and non-catheter-associated urinary tract infection events.

**Settings:** UTI Event reporting is currently available for certified skilled nursing facilities/nursing homes (LTC: SKILLNURS), and intermediate/chronic care facilities for the developmentally disabled (LTC: DEVDIS). Surveillance for UTIs should be performed facility-wide (FacWideIN).

Only UTI events presenting > 2 calendar days after admission (where date of admission is equal to day 1) are considered facility onset events. If a resident is transferred from an acute care facility and develops signs/symptoms of a UTI within the first 2 calendar days of admission to the LTCF, it would be considered present at the time of transfer to the LTCF. An event present at the time of transfer should be reported back to the transferring facility and not reported to NHSN as a LTCF UTI event.

Example: NHSN Classification of reportable LTCF UTI Events				
Admission date				
June 4 <sup>th</sup>	June 5 <sup>th</sup>	June 6 <sup>th</sup>	June 7 <sup>th</sup>	June 8 <sup>th</sup>
day 1	day 2	day 3	day 4	day 5
Not a LTCF reportable UTI event		LTCF reportable UTI event		

**Requirements:** A *NHSN Monthly Reporting Plan* for the LTCF ([CDC 57.141](#)) must be completed for each calendar month in which a facility plans to enter data into the NHSN. For each participating calendar month, facilities must report numerator (catheter-associated and non-catheter-associated UTI events) and denominator data for the entire facility, referred to as facility-wide inpatient (FacWideIN). UTI surveillance should be reported for at least 6 consecutive months to provide meaningful measures.

**Definitions:**

**Date of Event:** The date when the first clinical evidence (signs/symptoms) of the UTI appeared or the date the specimen used to meet the infection criteria was collected, **whichever comes first**.

**Indwelling urinary catheter:** A drainage tube that is inserted into the urinary bladder *through the urethra*, is left in place, and is connected to a drainage bag/collection system (including leg bags); also called a Foley catheter. Indwelling urinary catheters do not include straight in-and-out catheters or suprapubic catheters.



Urinary tract infections (UTI) are defined using [Symptomatic UTI \(SUTI\)](#) criteria for residents *without* an indwelling urinary device, [Catheter-Associated Symptomatic UTI \(CA-SUTI\)](#) criteria for residents *with* an indwelling urinary device, or [Asymptomatic Bacteremic UTI \(ABUTI\)](#) criteria for residents *with or without* an indwelling urinary device.

**Symptomatic UTI (SUTI):** Events that occur when the resident manifests signs and symptoms, such as acute dysuria, new and/or marked increase in urinary frequency, suprapubic tenderness, etc., which localize the infection to the urinary tract. These events can occur in residents without urinary devices or those managed with urinary devices other than indwelling urinary catheters, such as suprapubic catheters, straight in-and-out catheters and condom catheters. Events occurring in residents with indwelling urinary catheters (defined below) are a sub-set of SUTIs referred to as Catheter-Associated SUTI (CA-SUTI) events. (See [Figure 1](#) and [Table 2](#)).

**Catheter-associated SUTIs (CA-SUTI):** Events that occur when a resident develops signs and symptoms of a UTI while having an indwelling urinary catheter in place or removed within the 2 calendar days prior to the date of event, where day of catheter removal is equal to day 1 (*urinary catheter is in place on the day of event or the day before the event*). (See [Figure 2](#) and [Table 3](#)). **Note:** to be considered a CA-UTI, the indwelling catheter must be in place for >2 calendar days on the date of event, with day of device placement being Day 1.

**EXAMPLE:** Mr. T, is a resident in your facility. On March 1<sup>st</sup>, he developed an increase in incontinence and new suprapubic pain. Later that day a Foley catheter was inserted. The following day, on March 2<sup>nd</sup>, a specimen collected from the Foley catheter was sent to the lab and subsequently tested positive for greater than 100,000 ( $\geq 10^5$ ) CFU/ml of *E. coli*. Mr. T does meet criteria for a SUTI, but it is not considered as a CA-SUTI because the Foley catheter had not been in place >2 calendar days on the date of event (March 1<sup>st</sup>).

**Asymptomatic Bacteremic UTI (ABUTI):** Events that occur when the resident has NO signs or symptoms localizing to the urinary tract but has matching urine and blood cultures positive for at least one organism (see [Table 1](#)) regardless of whether a catheter is in place or not. (See [Figure 3](#) and [Table 4](#)).

Table 1. Examples of “sameness” by organism speciation		
Culture	Companion Culture	Report as...
<i>S. epidermidis</i>	Coagulase-negative <i>staphylococcus</i>	<i>S. epidermidis</i>
<i>Klebsiella oxytoca</i>	<i>Klebsiella</i> spp.	<i>K. oxytoca</i>
<i>S. salivarius</i>	<i>Streptococcus viridans</i>	<i>S. salivarius</i>



**Key Points:**

1. An indwelling urinary catheter should be in place for > 2 calendar days on the date of event (where day of catheter insertion = Day 1) in order for the SUTI to be catheter-associated.
2. If a resident is transferred to the facility with an indwelling urinary catheter in place, and the facility replaces the catheter with a new one while the resident is in the care of the facility, then the date of insertion of the device corresponds to the date the new catheter was placed in the LTCF.
3. UTIs in residents managed with suprapubic, in and out straight catheters, or condom (males only) catheters will be captured as SUTIs, not CA-SUTIs.



**Table 2. Criteria for Symptomatic Urinary Tract Infection (SUTI)**

Criterion	<i>For residents <b>without</b> an indwelling catheter in place or removed &gt;2 calendar days prior to the date of event, where day of catheter removal is equal to day 1:</i>
1	<p><b>Either</b> of the following (Signs &amp; Symptoms):</p> <ol style="list-style-type: none"> <li>1. Acute dysuria</li> <li>2. Acute pain, swelling, or tenderness of the testes, epididymis, or prostate</li> </ol> <p><b><u>AND</u></b></p> <p><b>Either</b> of the following (Laboratory and Diagnostic Testing):</p> <ol style="list-style-type: none"> <li>1. Specimen collected from clean catch voided urine and positive culture with no more than 2 species of microorganisms, at least one of which is a bacterium of <math>\geq 10^5</math> CFU/ml</li> <li>2. Specimen collected from in/out straight catheter and positive culture with no more than 2 species of microorganisms, at least one of which is a bacterium of <math>\geq 10^5</math> CFU/ml</li> </ol>
2	<p><b>Either</b> of the following:</p> <ol style="list-style-type: none"> <li>1. Fever<sup>+</sup> [Single temperature <math>\geq 37.8^\circ\text{C}</math> (<math>&gt;100^\circ\text{F}</math>), or <math>&gt;37.2^\circ\text{C}</math> (<math>&gt;99^\circ\text{F}</math>) on repeated occasions, or an increase of <math>&gt;1.1^\circ\text{C}</math> (<math>&gt;2^\circ\text{F}</math>) over baseline]</li> <li>2. Leukocytosis (<math>&gt;14,000</math> cells/<math>\text{mm}^3</math> or Left shift [<math>&gt;6\%</math> or <math>1,500</math> bands/<math>\text{mm}^3</math>])</li> </ol> <p><b><u>AND</u></b></p> <p><b>One or more</b> of the following (New and/or marked increase):</p> <ol style="list-style-type: none"> <li>1. Costovertebral angle pain or tenderness</li> <li>2. Suprapubic tenderness</li> <li>3. Visible (Gross) hematuria</li> <li>4. Incontinence</li> <li>5. Urinary urgency</li> <li>6. Urinary frequency</li> </ol> <p><b><u>AND</u></b></p> <p><b>Either</b> of the following (Laboratory and Diagnostic Testing):</p> <ol style="list-style-type: none"> <li>1. Specimen collected from clean catch voided urine and positive culture with no more than 2 species of microorganisms, at least one of which is a bacterium of <math>\geq 10^5</math> CFU/ml</li> <li>2. Specimen collected from in/out straight catheter and positive culture with no more than 2 species of microorganisms, at least one of which is a bacterium of <math>\geq 10^5</math> CFU/ml</li> </ol>



Criterion	<i>For residents <b>without</b> an indwelling catheter in place or removed &gt;2 calendar days prior to the date of event, where day of catheter removal is equal to day 1:</i>
3	<p><b>Two or more</b> of the following (New and/or marked increase):</p> <ol style="list-style-type: none"> <li>1. Costovertebral angle pain or tenderness</li> <li>2. Incontinence</li> <li>3. Urinary urgency</li> <li>4. Urinary frequency</li> <li>5. Suprapubic tenderness</li> <li>6. Visible (gross) hematuria</li> </ol> <p><b>AND</b></p> <p><b>Either</b> of the following (Laboratory and Diagnostic Testing):</p> <ol style="list-style-type: none"> <li>1. Specimen collected from clean catch voided urine and positive culture with no more than 2 species of microorganisms, at least one of which is a bacterium of <math>\geq 10^5</math> CFU/ml</li> <li>2. Specimen collected from in/out straight catheter and positive culture with no more than 2 species of microorganisms, at least one of which is a bacterium of <math>\geq 10^5</math> CFU/ml</li> </ol> <p><b>Footnote:</b></p> <ol style="list-style-type: none"> <li>1. + Fever can be used to meet SUTI criteria even if the resident has another possible cause for the fever (for example, pneumonia).</li> </ol>



**Table 3. Criteria for Catheter-associated Symptomatic Urinary Tract Infection (CA-SUTI)**

Criterion	<i>For residents <b>with</b> an indwelling catheter in place or removed within 2 calendar days prior to event onset, where day of catheter removal is equal to day 1:</i>
	<p><b>One or more</b> of the following (Signs and Symptoms and Laboratory and Diagnostic Testing):</p> <ol style="list-style-type: none"> <li>1. Fever<sup>+</sup>[Single temperature <math>\geq 37.8^{\circ}\text{C}</math> (<math>&gt;100^{\circ}\text{F}</math>), or <math>&gt;37.2^{\circ}\text{C}</math> (<math>&gt; 99^{\circ}\text{F}</math>) on repeated occasions, or an increase of <math>&gt;1.1^{\circ}\text{C}</math> (<math>&gt;2^{\circ}\text{F}</math>) over baseline]</li> <li>2. Rigors</li> <li>3. New onset hypotension, with no alternate non-infectious cause</li> <li>4. New onset confusion/functional decline with no alternate diagnosis <b>AND</b> leukocytosis (<math>&gt;14,000</math> cells/<math>\text{mm}^3</math> or Left shift [<math>&gt;6\%</math> or <math>1,500</math> bands/<math>\text{mm}^3</math>])</li> <li>5. New or marked increase in suprapubic tenderness</li> <li>6. New or marked increase in costovertebral angle pain or tenderness</li> <li>7. Acute pain, swelling, or tenderness of the testes, epididymis, or prostate</li> <li>8. Purulent discharge from around the catheter insertion site</li> </ol> <p><b>AND</b></p> <p><b>Any</b> of the following:</p> <p><i>If urinary catheter removed within last 2 calendar days:</i></p> <ol style="list-style-type: none"> <li>1. Specimen collected from clean catch voided urine and positive culture with no more than 2 species of microorganisms, at least one of which is a bacterium of <math>\geq 10^5</math> CFU/ml</li> <li>2. Specimen collected from in/out straight catheter and positive culture with no more than 2 species of microorganisms, at least one of which is bacterium of <math>\geq 10^5</math> CFU/ml</li> </ol> <p><i>If urinary catheter in place:</i></p> <ol style="list-style-type: none"> <li>3. Specimen collected from indwelling catheter and positive with no more than 2 species of microorganisms, at least one of which is a bacterium of <math>\geq 10^5</math> CFU/ml</li> </ol> <p><b>Footnote:</b></p> <ol style="list-style-type: none"> <li>1. <sup>+</sup> Fever can be used to meet CA-SUTI criteria even if the resident has another possible cause for the fever (for example, pneumonia).</li> </ol>



**Table 4. Criteria for Asymptomatic Bacteremic Urinary Tract Infection (ABUTI)**

Criterion	<i>Resident with or without an indwelling urinary catheter</i>
	<p><b>No qualifying fever or</b> signs or symptoms (specifically, no urinary urgency, urinary frequency, acute dysuria, suprapubic tenderness, or costovertebral angle pain or tenderness). <i>If no catheter is in place, fever alone would not exclude ABUTI if other criteria are met.</i></p> <p><b><u>AND</u></b></p> <p><b>One</b> of the following:</p> <ol style="list-style-type: none"> <li>1. Specimen collected from clean catch voided urine and positive culture with no more than 2 species of microorganisms, at least one of which is a bacterium of <math>\geq 10^5</math> CFU/ml</li> <li>2. Specimen collected from in/out straight catheter and positive culture with no more than 2 species of microorganisms, at least one of which is a bacterium of <math>\geq 10^5</math> CFU/ml</li> <li>3. Specimen collected from indwelling catheter and positive culture with no more than 2 species of microorganisms, at least one of which is a bacterium of <math>\geq 10^5</math> CFU/ml</li> </ol> <p><b><u>AND</u></b></p> <p>A positive blood culture with at least 1 matching bacteria to the urine culture</p>

**Notes:**

1. “Mixed flora” is not available in the pathogen list within NSHN, and cannot be reported as a pathogen to meet the NHSN UTI criteria. Additionally, “mixed flora” often represents contamination and likely represents presence of multiple organisms in culture.
2. Yeast and other microorganisms, which are not bacteria, are not acceptable UTI pathogens.
3. To remove the subjectivity about whether a fever is attributable to a UTI event, the presence of a fever, even if due to another cause (for example, pneumonia), should still be counted as part of meeting a UTI definition.





## Numerator and Denominator Data:

**Numerator:** The *Urinary Tract Infection (UTI) for LTCF* form ([CDC 57.140](#)) is used to collect and report each UTI that is identified during the month selected for surveillance. The [Table of Instructions for Completion of a Urinary Tract Infection for LTCF form](#) include brief instructions for collection and entry of each data element on the form.

The UTI form includes resident demographic information and information on whether or not a catheter (or other urinary device) was present. Additional data include the specific clinical criteria evidence (signs and symptoms) and laboratory and diagnostic testing that were used for identifying the UTI; whether the resident developed a secondary bloodstream infection; whether the resident was transferred to an acute care facility for any reason or died from any cause within 7 days of the UTI event; and the organisms isolated from cultures and their antimicrobial susceptibilities.

**Denominator:** Includes monthly totals for resident-days, urinary catheter-days, new antibiotic starts for UTI indication, and number of urine cultured ordered. The Denominator for LTCF form ([CDC 57.142](#)) may be used to collect denominator data. The daily counts are summed and only the totals for the month are entered into the NHSN. The [Table of Instructions for Completion of the Long-term Care Facility Component-Denominators for LTCF](#) include brief instructions for collection and entry of each data element on the form.

*Catheter-days*, defined as the number of residents with an indwelling urinary (Foley) catheter, are collected daily for all residents in the facility. These daily counts are summed and only the total for the month is entered into NHSN, under Summary Data.

### Notes:

1. None of the following urinary management devices should be included when counting indwelling catheter-days: suprapubic catheters, straight in-and-out catheters, or condom catheters.
2. If a resident is transferred to an acute care facility, no additional indwelling catheter-days are reported after the day of transfer.

*Resident-days* are calculated using the daily census of residents in the facility each day of the month.

*New antibiotic starts for UTI indication* refers to a new prescription for an antibiotic ordered for a resident who is suspected of having or diagnosed with a UTI, either catheter-associated or non-catheter associated, regardless of whether that UTI meets the NHSN event definition.



**Notes:**

1. There is no minimum number of doses or days of therapy that define a new antibiotic start—count all new orders.
2. Include only antibiotics that are started while the resident is receiving care in your facility, either by clinical providers working in the facility or by outside physicians who see the resident in an outpatient clinic or emergency department.
3. Do not include antibiotic courses started by another healthcare facility prior to the resident's admission or readmission back to your facility, even if the antibiotic is continually administered while the resident is in your facility.
4. Data may be collected daily or summarized at the end of each month.

*Number of urine cultures ordered* refers to new urine cultures ordered for a resident regardless of whether the resident has a UTI meeting the NHSN event definition.

**Notes:**

1. Include only urine culture orders that are ordered while the resident is receiving care in your facility, either by clinical providers working in the facility or by outside physicians who see the resident in an outpatient clinic or Emergency department.
2. Do not include urine cultures ordered by another healthcare facility prior to the resident's admission or readmission back to your facility.
3. Data may be collected daily or summarized at the end of each month.

**Data Analyses:**

*Line listing of UTI events and UTI events by catheter status are available as part of the UTI event within the NHSN LTCF component. Below are measures and calculations that are incorporated into the analytics output.*

**Calculated UTI Rates and Metrics**

*Data will be stratified by time (month, quarter, year) and aggregated across the entire facility.*

Total UTI incidence rate/1,000 resident-days = Total Number of UTI Events (specifically, SUTI + CA-SUTI + ABUTI) / Total resident-days x 1,000.

Percent that are SUTI = Number of SUTI Events / Total number of UTI Events x 100.

Percent that are CA-SUTI = Number of CA-SUTI Events / Total number of UTI Events x 100.



Percent that are ABUTI = Number of ABUTI Events / Total number of UTI Events x 100.

SUTI incidence rate/1,000 resident-days = Number of SUTI Events / Total resident days x 1,000.

**Note:** Only SUTIs that are NOT catheter-associated will be included in the SUTI incidence rate.

CA-SUTI incidence rate/1,000 catheter-days = Number of CA-SUTI events / Total catheter days x 1,000

Urinary Catheter Utilization Ratio = Total urinary catheter days / Total resident days

Number of Urine Cultures Ordered = Number of urine cultures ordered / Total resident days x 1,000

UTI treatment ratio = New antibiotic starts for UTI / Total Number of UTI Events (SUTI + ABUTI + CA-SUTI)


**Note:**

1. When the UTI treatment ratio is <1, there are fewer reported antibiotic starts for UTI than symptomatic UTI events submitted; when the UTI treatment ratio equals 1, there are the same number of new antibiotic starts for UTI events submitted; when the UTI treatment ratio is >1, there are more reported antibiotic starts for UTI than UTI events submitted.



**Figure 1: Criteria for Defining Non-Catheter Associated Symptomatic Urinary Tract Infection (SUTI):**

**Resident *without* an indwelling catheter (Meets criteria 1 OR 2 OR 3):**



<sup>+</sup> Fever can be used to meet SUTI criteria even if the resident has another possible cause for the fever (for example, pneumonia)

<sup>a</sup> Fever: Single temperature  $\geq 37.8^{\circ}\text{C}$  ( $>100^{\circ}\text{F}$ ), or  $> 37.2^{\circ}\text{C}$  ( $>99^{\circ}\text{F}$ ) on repeated occasions, or an increase of  $>1.1^{\circ}\text{C}$  ( $>2^{\circ}\text{F}$ ) over baseline

<sup>b</sup> Leukocytosis:  $>14,000$  cells/mm<sup>3</sup>, or Left shift ( $> 6\%$  or  $1,500$  bands/mm<sup>3</sup>)



**Figure 2: Criteria for Defining Catheter Associated Symptomatic Urinary Tract Infection (CA-SUTI)**

**Resident with an indwelling urinary catheter:**

**ONE or more** of the following:

- Fever<sup>+</sup> <sup>a</sup>
- Rigors
- New onset hypotension, with no alternate noninfectious cause
- New onset confusion/functional decline with no alternate diagnosis **AND** Leukocytosis<sup>b</sup>
- New costovertebral angle pain or tenderness
- New or marked increase in suprapubic tenderness
- Acute pain, swelling or tenderness of the testes, epididymis or prostate
- Purulent discharge from around the catheter

**AND**

**Any** of the following:

*If urinary catheter removed within last 2 calendar days:*

1. Specimen collected from clean catch voided urine and positive culture with no more than 2 species of microorganisms, at least one of which is a bacterium of  $\geq 10^5$  CFU/ml
2. Specimen collected from in/out straight catheter and positive culture with no more than 2 species of microorganisms, at least one of which is a bacterium of  $\geq 10^5$  CFU/ml

*If urinary catheter in place:*

3. Specimen collected from indwelling catheter and positive culture with no more than 2 species of microorganisms, at least one of which is a bacterium of  $\geq 10^5$  CFU/ml

**CA-SUTI**

<sup>+</sup> Fever can be used to meet SUTI criteria even if the resident has another possible cause for the fever (for example, pneumonia)

<sup>a</sup> Fever: Single temperature  $\geq 37.8^\circ\text{C}$  ( $>100^\circ\text{F}$ ), or  $> 37.2^\circ\text{C}$  ( $>99^\circ\text{F}$ ) on repeated occasions, or an increase of  $>1.1^\circ\text{C}$  ( $>2^\circ\text{F}$ ) over baseline

<sup>b</sup> Leukocytosis:  $>14,000$  cells/ $\text{mm}^3$  or Left shift ( $> 6\%$  or  $1,500$  bands/ $\text{mm}^3$ )



**Figure 3: Criteria for Defining Asymptomatic Bacteremic Urinary Tract Infection (ABUTI)**

**Resident with or without an indwelling catheter:**

Resident has **no qualifying fever or localizing urinary signs or symptoms** (specifically, no urgency, frequency, acute dysuria, suprapubic tenderness, or costovertebral angle pain or tenderness). *If no catheter is in place, fever as only sign would not exclude ABUTI if other positive culture criteria are met.*

**AND**

**Any** of the following:

1. Specimen collected from clean catch voided urine and a positive culture with no more than 2 species of microorganisms, at least one of which is a bacterium of  $\geq 10^5$  CFU/ml
2. Specimen collected from in/out straight catheter and positive culture with no more than 2 species of microorganisms, at least one of which is a bacterium of  $\geq 10^5$  CFU/ml
3. Specimen collected from indwelling catheter and positive culture with no more than 2 species of microorganism, at least one of which is a bacterium of  $\geq 10^5$  CFU/ml

NOTE: Yeast and other microorganisms which are not bacteria, are not acceptable UTI pathogens

**AND**

Positive blood culture with at least 1 matching organism in urine culture

**ABUTI**