



Wyoming *State Nutrition, Physical Activity, and Obesity Profile*

Obesity has important consequences on our nation's health and economy. It is linked to a number of chronic diseases, including coronary heart disease, stroke, diabetes, and some cancers (NIH Clinical Guidelines, 1998). Among adults, the medical costs associated with obesity are estimated at 147 billion dollars (Finkelstein, 2009). Many American communities are characterized by unhealthy options when it comes to diet and physical activity. We need public health approaches that make healthy options available, accessible, and affordable for all Americans.

Wyoming - State Nutrition, Physical Activity, and Obesity Profile

CDC's Division of Nutrition and Physical, and Obesity (DNPAO) supports the nation's capacity to address public health in all policies and establish successful and sustainable interventions to support healthy eating and active living. The Division provides support (i.e., implementation and evaluation guidance, technical assistance, training, surveillance and applied research, translation and dissemination, and partnership development) to states, communities and national partners to implement policy, system, and environmental strategies. The goal is to improve dietary quality, increase physical activity and reduce obesity across multiple settings—such as child care facilities, workplaces, hospitals and medical care facilities, schools, and communities.

State Population of Wyoming

- Estimated Total Population 2010⁽¹⁾
= 563,626
- Adults age 18 and over⁽²⁾
= 76.0% of the total population in 2010
- Youth under 18 years of age⁽¹⁾
= 24.0% of the total population in 2010

⁽¹⁾ U.S. Census Bureau. State and County QuickFacts. 2011. Available online at <http://quickfacts.census.gov/qfd/index.html>

⁽²⁾ Calculated estimated = 100% minus percent of the total population under 18 years old, using State and County QuickFacts, 2010 data from the U.S. Census.

Adult Overweight and Obesity

Overweight and Obesity⁽³⁾

- 62.9% were overweight, with a Body Mass Index of 25 or greater.
- 25.1% were obese, with a Body Mass Index of 30 or greater.

Dietary Behaviors⁽⁴⁾

- 30.3% of adults reported having consumed fruits at the recommended level of 2 or more times per day.
- 26.9% of adults reported having consumed vegetables at the recommended level of 3 or more times per day.

Physical Activity⁽⁵⁾

- 51.0% of adults achieved at least 300 minutes a week of moderate-intensity aerobic physical activity or 150 minutes a week of vigorous-intensity aerobic activity (or an equivalent combination).
- 22.5% of Wyoming's adults reported that during the past month, they had not participated in any physical activity.

Source of Adult Obesity Data:

⁽³⁾ CDC. Behavioral Risk Factor Surveillance System: Prevalence and Trend Data—Overweight and Obesity, U.S. Obesity Trends, Trends by State 2010. Available online at <http://www.cdc.gov/brfss/>

Source of Adult Fruit and Vegetable Data:

⁽⁴⁾ CDC. *MMWR* September 2010 State-Specific Trends in Fruit and Vegetable Consumption Among Adults United States, 2000–2009. Available online at <http://www.cdc.gov/mmwr/pdf/wk/mm5935.pdf>

Source of Adult Physical Activity Data:

⁽⁵⁾ CDC. BRFSS Behavioral Risk Factor Surveillance System: Prevalence and Trend Data—Physical Activity, U.S. Physical Activity Trends by State 2009–2010. Available online at <http://www.cdc.gov/brfss/>

Adolescent Overweight and Obesity

Overweight and Obesity⁽⁶⁾

- 12.6% were overweight ($\geq 85^{\text{th}}$ and $< 95^{\text{th}}$ percentiles for BMI by age and sex, based on reference data)
- 9.8% were obese ($\geq 95^{\text{th}}$ percentile BMI by age and sex, based on reference data)

Unhealthy Dietary Behaviors⁽⁶⁾

- **Fruit consumption:** 73.2% ate fruits or drank 100% fruit juice less than 2 times per day during the 7 days before the survey (100% fruit juice or fruit).
- **Vegetable consumption:** 84.0% ate vegetables less than 3 times per day during the 7 days before the survey (green salad; potatoes, excluding French fries, fried potatoes, or potato chips; carrots; or other vegetables).
- **Sugar-sweetened beverage consumption:** 27.0% drank a can, bottle, or glass of soda or pop (not including diet soda or diet pop) at least one time per day during the 7 days before the survey.

Wyoming - State Nutrition, Physical Activity, and Obesity Profile

Physical Activity⁽⁶⁾

- **Achieved recommended level of activity:** Only 25.6% were physically active* for a total of at least 60 minutes per day on each of the 7 days prior to the survey.
- **Participated in daily physical education:** 22.7% of adolescents attended daily physical education classes in an average week (when they were in school).

Physical Inactivity⁽⁶⁾

- **No activity:** 13.6% did not participate in at least 60 minutes of physical activity on any day during the 7 days prior to the survey.
- **Television viewing time:** 22.0% watched television 3 or more hours per day on an average school day.

The 2010 Wyoming School Health Profiles assessed the school environment, indicating that among high schools⁽⁷⁾

- 31.9% did not sell less nutritious foods and beverages anywhere outside the school food service program
- 8.7% always offered fruits or non-fried vegetables in vending machines and school stores, canteens, or snack bars, and during celebrations whenever foods and beverages were offered
- 38.9% prohibited all forms of advertising and promotion of candy, fast food restaurants, or soft drinks in all locations. All school-related locations were defined as in school buildings; on school grounds, including on the outside of the school building, on playing fields, or other areas of the campus; on school buses or other vehicles used to transport students; and in school publications. Sources of Adolescent Obesity, Fruit and Vegetable, Sugar-sweetened Beverages, and Physical Activity Data:

* Physical activity defined as “any kind of physical activity that increases your heart rate and makes you breathe hard some of the time.”

⁽⁶⁾ CDC, Division of Adolescent and School Health. The 2009 Youth Risk Behavior Survey. Available online at <http://www.cdc.gov/HealthyYouth/yrbs/index.htm>

⁽⁷⁾ CDC, Division of Adolescent and School Health. The 2010 School Health Profiles. Available online at <http://www.cdc.gov/healthyyouth/profiles/index.htm>

Child Overweight and Obesity

Breastfeeding⁽⁸⁾

Increasing breastfeeding initiation, duration, and exclusivity is a priority strategy in CDC’s efforts to decrease the rate of childhood obesity throughout the United States.

- 83.6% of infants were Ever Breastfed.
- 48.2% of infants were Breastfed for at least 6 months.

Body Mass Index^{(9)*}

Among Wyoming’s children aged 2 years to less than 5 years*

Data Not Available

Sources of Breastfeeding Data:

⁽⁸⁾ CDC. Division of Nutrition, Physical Activity, and Obesity Breastfeeding Report Card 2011. Centers for Disease Control and Prevention National Immunization Survey, Provisional Data, 2008births. Available online at <http://www.cdc.gov/breastfeeding/data/reportcard2.htm>

Sources of Child Obesity Data:

⁽⁹⁾ CDC. Division of Nutrition, Physical Activity, and Obesity. 2010 Pediatric Nutrition Surveillance System, Table 6 (PedNSS). http://www.cdc.gov/pednss/pednss_tables/tables_health_indicators.htm

* BMI data only includes low-income children from the PedNSS sample and do not represent all children.

* BMI data is based on 2000 CDC growth chart percentiles for BMI-for-age for children 2 years of age and older.

Wyoming’s Response to Obesity

Workforce Wellness Policies

The Wyoming Department of Health focused on policies that support and promote healthy eating, breastfeeding and physical activity in the workplace. In partnership with the Diabetes Prevention & Control Program and the Heart Disease and Stroke Prevention Program, Wyoming surveyed worksites to assess the number of policies, practices, and programs in A key partner, TwoMedicine, conducted worksite wellness workshops around the state and provides a resource kit concerning how to implement a worksite wellness program. Initial reach is expected to be 2,000 employees.

Wyoming - State Nutrition, Physical Activity, and Obesity Profile

The Wyoming Department of Health conducted the 4th Annual Workforce Wellness Summit in October 2011. The purpose of the summit was to promote the building or enhancing of new and existing workforce wellness programs. The conference included sessions on tobacco cessation policy development as well as managing healthcare costs while keeping the workforce healthy and productive. A total of eighty-five companies were represented.

Breastfeeding Policy in the Workplace

Workshops were conducted throughout the state on how businesses can support breastfeeding including: offering flexible work schedules; providing lactation rooms; and accommodating infants in the workplace. The National Business Group on Health's toolkit, "Investing in Workplace Breastfeeding Programs and Policies" is provided for businesses interested in attempting to accommodate breastfeeding employees. In addition, technical assistance is provided on establishing a breastfeeding policy. Wyoming Department of Health staffers are looking at which state agencies have existing breastfeeding policies in place and how they are being implemented.

In October 2011, the Wyoming Department of Health hosted a lactation support conference. The goal of the conference was to raise awareness of breastfeeding in the workplace practices and policies and baby friendly hospitals. There were approximately 65 attendees that represented the Women, Infants, & Children Program (WIC), public health nurses, students, and hospitals from around the state of Wyoming.

Contact Information

Star Morrison, MS, RD
Program Manager
Diabetes Prevention & Control Program
6101 Yellowstone Road, Suite 510
Cheyenne, WY 82002
Phone: (307) 777-3579
Fax: (307) 777-8604
E-mail: star.morrison@wyo.gov

References

NIH. Clinical Guidelines on the Identification, Evaluation, and Treatment of Overweight and Obesity in Adults: The Evidence Report. 1998. Available online at http://www.nhlbi.nih.gov/guidelines/obesity/ob_gdlns.htm

Finkelstein, EA, Trogon, JG, Cohen, JW, and Dietz, W. Annual medical spending attributable to obesity: Payer- and service-specific estimates. *Health Affairs* 2009; 28(5): w822-w831.

