



Idaho

State Nutrition, Physical Activity, and Obesity Profile

Obesity has important consequences on our nation's health and economy. It is linked to a number of chronic diseases, including coronary heart disease, stroke, diabetes, and some cancers (NIH Clinical Guidelines, 1998). Among adults, the medical costs associated with obesity are estimated at 147 billion dollars (Finkelstein, 2009). Many American communities are characterized by unhealthy options when it comes to diet and physical activity. We need public health approaches that make healthy options available, accessible, and affordable for all Americans.

Idaho - State Nutrition, Physical Activity, and Obesity Profile

CDC's Division of Nutrition and Physical, and Obesity (DNPAO) supports the nation's capacity to address public health in all policies and establish successful and sustainable interventions to support healthy eating and active living. The Division provides support (i.e., implementation and evaluation guidance, technical assistance, training, surveillance and applied research, translation and dissemination, and partnership development) to states, communities and national partners to implement policy, system, and environmental strategies. The goal is to improve dietary quality, increase physical activity and reduce obesity across multiple settings—such as child care facilities, workplaces, hospitals and medical care facilities, schools, and communities.

State Population of Idaho

Estimated Total Population 2010⁽¹⁾
= 1,567,582

Adults age 18 and over⁽²⁾
= 72.6% of the total population in 2010

Youth under 18 years of age⁽¹⁾
= 27.4% of the total population in 2010

⁽¹⁾ U.S. Census Bureau. State and County QuickFacts. 2011. Available online at <http://quickfacts.census.gov/qfd/index.html>

⁽²⁾ Calculated estimated = 100% minus percent of the total population under 18 years old, using State and County QuickFacts, 2010 data from the U.S. Census.

Adult Overweight and Obesity

Overweight and Obesity⁽³⁾

- 62.0% were overweight, with a Body Mass Index of 25 or greater.
- 26.5% were obese, with a Body Mass Index of 30 or greater.

Dietary Behaviors⁽⁴⁾

- 32.9% of adults reported having consumed fruits at the recommended level of 2 or more times per day.
- 27.8% of adults reported having consumed vegetables at the recommended level of 3 or more times per day.

Physical Activity⁽⁵⁾

- 52.1% of adults achieved at least 300 minutes a week of moderate-intensity aerobic physical activity or 150 minutes a week of vigorous-intensity aerobic activity (or an equivalent combination).
- 21.0% of Idaho's adults reported that during the past month, they had not participated in any physical activity.

Source of Adult Obesity Data:

⁽³⁾ CDC. Behavioral Risk Factor Surveillance System: Prevalence and Trend Data—Overweight and Obesity, U.S. Obesity Trends, Trends by State 2010. Available online at <http://www.cdc.gov/brfss/>

Source of Adult Fruit and Vegetable Data:

⁽⁴⁾ CDC. *MMWR* September 2010 State-Specific Trends in Fruit and Vegetable Consumption Among Adults United States, 2000–2009. Available online at <http://www.cdc.gov/mmwr/pdf/wk/mm5935.pdf>

Source of Adult Physical Activity Data:

⁽⁵⁾ CDC. BRFSS Behavioral Risk Factor Surveillance System: Prevalence and Trend Data—Physical Activity, U.S. Physical Activity Trends by State 2009–2010. Available online at <http://www.cdc.gov/brfss/>

Adolescent Overweight and Obesity

Overweight and Obesity⁽⁶⁾

- 12.0% were overweight ($\geq 85^{\text{th}}$ and $< 95^{\text{th}}$ percentiles for BMI by age and sex, based on reference data)
- 8.8% were obese ($\geq 95^{\text{th}}$ percentile BMI by age and sex, based on reference data)

Unhealthy Dietary Behaviors⁽⁶⁾

- **Fruit consumption:** 71.1% ate fruits or drank 100% fruit juice less than 2 times per day during the 7 days before the survey (100% fruit juice or fruit) (continued on next page)

Idaho - State Nutrition, Physical Activity, and Obesity Profile

- **Vegetable consumption:** 88.9% ate vegetables less than 3 times per day during the 7 days before the survey (green salad; potatoes, excluding French fries, fried potatoes, or potato chips; carrots; or other vegetables)
- **Sugar-sweetened beverage consumption:** 18.3% drank a can, bottle, or glass of soda or pop (not including diet soda or diet pop) at least one time per day during the 7 days before the survey.

Physical Activity⁽⁶⁾

- **Achieved recommended level of activity:** Only 27.6% were physically active* for a total of at least 60 minutes per day on each of the 7 days prior to the survey.
- **Participated in daily physical education:** 23.7% of adolescents attended daily physical education classes in an average week (when they were in school).

Physical Inactivity⁽⁶⁾

- **No activity:** 10.8% did not participate in at least 60 minutes of physical activity on any day during the 7 days prior to the survey.
- **Television viewing time:** 21.9% watched television 3 or more hours per day on an average school day.

The 2010 Idaho School Health Profiles assessed the school environment, indicating that among high schools⁽⁷⁾

- 25.8% did not sell less nutritious foods and beverages anywhere outside the school food service program.
- 9.2% always offered fruits or non-fried vegetables in vending machines and school stores, canteens, or snack bars, and during celebrations whenever foods and beverages were offered.
- 41.5% prohibited all forms of advertising and promotion of candy, fast food restaurants, or soft drinks in all locations. All school-related locations were defined as in school buildings;

on school grounds, including on the outside of the school building, on playing fields, or other areas of the campus; on school buses or other vehicles used to transport students; and in school publications.

Sources of Adolescent Obesity, Fruit and Vegetable, Sugar-sweetened Beverages, and Physical Activity Data:

* Physical activity defined as "any kind of physical activity that increases your heart rate and makes you breathe hard some of the time."

⁽⁶⁾ CDC, Division of Adolescent and School Health. The 2009 Youth Risk Behavior Survey. Available online at <http://www.cdc.gov/HealthyYouth/yrbs/index.htm>

⁽⁷⁾ CDC, Division of Adolescent and School Health. The 2010 School Health Profiles. Available online at <http://www.cdc.gov/healthyouth/profiles/index.htm>

Child Overweight and Obesity

Breastfeeding⁽⁸⁾

Increasing breastfeeding initiation, duration, and exclusivity is a priority strategy in CDC's efforts to decrease the rate of childhood obesity throughout the United States.

- 84.5% of infants were Ever Breastfed.
- 61.2% of infants were Breastfed for at least 6 months.

Body Mass Index^{(9)*}

Among Idaho's children aged 2 years to less than 5 years*

- 15.3% were overweight (85th to < 95th percentile BMI-for-Age).
- 11.4% were obese (\geq 95th percentile BMI-for-Age).

Sources of Breastfeeding Data:

⁽⁸⁾ CDC. Division of Nutrition, Physical Activity, and Obesity Breastfeeding Report Card 2011. Centers for Disease Control and Prevention National Immunization Survey, Provisional Data, 2008births. Available online at <http://www.cdc.gov/breastfeeding/data/reportcard2.htm>

Sources of Child Obesity Data:

⁽⁹⁾ CDC. Division of Nutrition, Physical Activity, and Obesity. 2010 Pediatric Nutrition Surveillance System, Table 6 (PedNSS). http://www.cdc.gov/pednss/pednss_tables/tables_health_indicators.htm

* BMI data only includes low-income children from the PedNSS sample and do not represent all children.

* BMI data is based on 2000 CDC growth chart percentiles for BMI-for-age for children 2 years of age and older.

Idaho's Response to Obesity

✦ School Vending Policies

Idaho's Department of Health and Welfare (IDHW) provided mini-grants, training, and technical assistance to three school districts with interest in working on limiting unhealthy food and drink availability in schools through healthy vending policies. The three pilot sites successfully implemented changes that provided better alignment for the new federal standards for competitive foods. The districts chose to undergo a process to update their Wellness Policies and two sites have been successful at getting School Board Association approval of their new competitive standards. The third site worked with superintendent to make the changes. All three sites will be following IOM guidelines for Competitive Foods. The three action plans include focusing in on at least one area which includes ala cart, vending, and competitive foods.

The long term outcome is to establish uniform statewide vending policies that can easily be implemented.

✦ Complete Streets Policy

IDHW guided five pilot communities (*Idaho Falls, Wood River Valley, Nampa, Lewiston and Moscow*) in forming Active Living Task Forces (ALTFs) to develop the community capacity and infrastructure necessary for city transportation, planning agencies and Metropolitan Planning Organizations (MPOs) to support active community environments through complete streets policy adoption and implementation. Coupled with complete streets policies already adopted by the City of Coeur d'Alene, the Ada County Highway District (ACHD) and the Community Planning Association of Southwest Idaho (COMPASS) MPO, the potential reach is approximately 41% of Idaho's population (622,000).

IDHW, in conjunction with Idaho Smart Growth, organized and supported regional trainings throughout the state around Complete Streets and building healthy communities. These workshops provided ALTF members, state/city transportation and planning agencies; public health, and city leaders with information about preventing chronic disease and reducing health inequities through the development and implementation of complete streets and other land use policies.

Contact Information

Rebecca Lemmons, MHS
Acting Manager, Physical Activity and
Nutrition Program
Idaho Department of Health and Welfare
PO Box 83720, 6th Floor
Boise, ID 83720-0036
Office: 208-334-5610 Fax: 208-334-6573
Mobile: 208-559-4758
lemmonsr@dhw.idaho.gov

References

NIH. Clinical Guidelines Clinical Guidelines on the Identification, Evaluation, and Treatment of Overweight and Obesity in Adults: The Evidence Report. 1998. Available online at http://www.nhlbi.nih.gov/guidelines/obesity/ob_gdlns.htm

Finkelstein, EA, Trogon, JG, Cohen, JW, and Dietz, W. Annual medical spending attributable to obesity: Payer- and service-specific estimates. *Health Affairs* 2009; 28(5): w822-w831.

