



## Prevalence of Selected Maternal and Child Health Indicators\* for Georgia, Pregnancy Risk Assessment Monitoring System (PRAMS), 2012-2015

Health Indicator*	2012		2013		2014		2015		Overall 2015 <sup>¶</sup>
	n <sup>†</sup>	% <sup>§</sup> (95% CI) <sup>‡</sup>	n <sup>†</sup>	% <sup>§</sup> (95% CI) <sup>‡</sup>	n <sup>†</sup>	% <sup>§</sup> (95% CI) <sup>‡</sup>	n <sup>†</sup>	% <sup>§</sup> (95% CI) <sup>‡</sup>	% <sup>§</sup>
<b>Nutrition and Exercise</b>									
- Multivitamin use $\geq$ 4 days a week during the month before pregnancy	1109	29.1 (25.5-32.9)	733	28.0 (23.2-33.5)					38.7
- Dieting during the 12 months before pregnancy	1116	25.7 (22.3-29.4)	739	23.5 (19.1-28.7)					30.1
- Exercising $\geq$ 3 days a week during the 12 months before pregnancy	1115	47.5 (43.5-51.5)	737	40.0 (34.6-45.7)					47.3
<b>Pre-pregnancy Weight</b>									
- Underweight (BMI<18.5)	1023	3.3 (2.1-5.1)	694	1.8 (0.8-4.3)					3.9
- Overweight (BMI 25-29.9)	1023	26.8 (23.2-30.7)	694	39.5 (33.9-45.5)					24.3
- Obese (BMI $\geq$ 30)	1023	19.6 (16.5-23.2)	694	20.8 (16.4-26.1)					24.1
<b>Substance Use</b>									
- Any cigarette smoking during the 3 months before pregnancy	1112	12.1 (9.8-14.9)	729	10.3 (7.4-14.3)					19.4
- Any cigarette smoking during the last 3 months of pregnancy	1113	4.2 (2.9-6.0)	729	3.2 (1.9-5.6)					8.8
- Any cigarette smoking postpartum	1101	8.2 (6.2-10.6)	729	6.5 (4.3-9.8)					12.6
- Any alcohol use during the 3 months before pregnancy	1111	37.7 (33.8-41.7)	736	32.7 (27.6-38.2)					55.3
- Any alcohol use during the last 3 months of pregnancy	1115	3.9 (2.6-5.9)	738	4.4 (2.7-7.3)					8.0
<b>Intimate Partner Violence (IPV)</b>									
- Experienced IPV during the 12 months before pregnancy	1116	2.6 (1.6-4.2)	739	1.6 (0.6-3.9)					2.3
- Experienced IPV during pregnancy	1103	1.5 (0.8-2.7)	739	1.9 (0.8-4.5)					2.1
<b>Depression</b>									
- Told by a provider had depression before pregnancy <sup>#</sup>	1109	2.6 (1.7-3.9)	732	4.7 (2.8-7.7)					10.5
- Self-reported postpartum depressive symptoms	1108	8.0 (6.2-10.4)	737	9.2 (6.4-13.1)					12.8
<b>Pregnancy Intention and Family Planning</b>									
- Mistimed/unwanted pregnancy	945	36.8 (32.8-41.0)	594	34.6 (28.8-40.8)					34.2
- Unsure about pregnancy	1109	14.1 (11.5-17.1)	733	20.8 (16.4-25.9)					15.4
- Postpartum contraceptive use	1109	83.4 (80.2-86.2)	733	83.0 (78.3-86.9)					78.4

Health Care Services					
- Talked to provider about improving health before pregnancy <sup>#</sup>	1072	13.0 (10.5-16.1)	731	11.9 (8.7-16.1)	22.8
- Began prenatal care in 1st trimester	1093	86.1 (83.1-88.6)	721	84.4 (79.6-88.2)	83.0
- Took childbirth class <sup>#</sup>	1109	17.6 (14.7-21.0)	731	14.6 (11.0-19.0)	21.6
- Had home visitor during pregnancy <sup>#</sup>	1113	1.5 (0.8-2.7)	733	3.1 (1.6-5.9)	6.9
- Had flu shot before or during pregnancy	1078	33.5 (29.7-37.4)	722	23.7 (19.3-28.8)	57.3
- Had maternal postpartum checkup	1116	92.8 (90.5-94.5)	733	91.6 (87.8-94.3)	89.7
Oral Health					
- Teeth cleaned during 12 months before pregnancy	1116	39.7 (35.8-43.7)	738	48.1 (42.4-53.8)	56.3
- Teeth cleaned during pregnancy	1102	38.0 (34.1-42.0)	728	29.2 (24.3-34.7)	48.3
- Had dental insurance during pregnancy <sup>#</sup>	1103	51.8 (47.8-55.9)	725	39.4 (33.9-45.1)	72.6
Pre-pregnancy Health Insurance <sup>‡</sup> Status					
- Private insurance	1079	51.9 (47.8-55.9)	713	47.3 (41.6-53.1)	58.5
- Medicaid	1079	12.0 (9.7-14.6)	713	11.5 (8.3-15.7)	21.8
- No insurance	1079	35.7 (31.9-39.8)	713	39.7 (34.2-45.5)	18.7
Health Insurance <sup>‡</sup> Status During Pregnancy					
- Private insurance	1085	47.8 (43.8-51.9)	722	44.2 (38.6-50.0)	57.0
- Medicaid	1085	42.7 (38.7-46.7)	722	50.2 (44.4-55.9)	38.4
- No insurance	1110	9.6 (7.4-12.3)	734	5.2 (3.1-8.5)	2.9
Health Insurance Status Postpartum					
- Private insurance	1079	46.2 (42.2-50.3)	722	45.5 (39.8-51.3)	55.2
- Medicaid	1079	17.0 (14.3-20.1)	722	11.2 (8.2-15.3)	29.1
- No insurance	1108	35.8 (32.0-39.7)	740	41.2 (35.7-46.9)	14.2
Infant Sleep Practices					
- Baby most often laid on back to sleep	1091	46.5 (42.4-50.6)	726	43.8 (38.2-49.5)	78.4
Breastfeeding Practices					
- Ever breastfed	1091	67.8 (63.9-71.5)	730	63.9 (58.2-69.2)	87.1
- Any breastfeeding at 8 weeks	1069	50.3 (46.2-54.4)	719	48.2 (42.5-54.0)	65.6

\* Indicators and definitions available in PRAMStat (<http://www.cdc.gov/prams/pramstat/index.html>) unless otherwise indicated.

<sup>†</sup> Unweighted sample size

<sup>§</sup> Weighted percent

<sup>¥</sup> 95% Confidence Interval

<sup>¶</sup> PRAMS site aggregate for 2015: Alabama, Alaska, Arkansas, Colorado, Connecticut, Delaware, Hawaii, Illinois, Iowa, Louisiana, Maine, Maryland, Massachusetts, Michigan, Missouri, Nebraska, New Hampshire, New Jersey, New Mexico, New York City, New York State, Ohio, Oklahoma, Oregon, Pennsylvania, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, and Wyoming met the required 55% response rate threshold for inclusion.

<sup>#</sup> Indicator is new for Phase 7 (2012-2015) survey and is not currently available in PRAMStat.

<sup>‡</sup> Insurance is coded as Medicaid (Medicaid or state-named Medicaid program); Private (Private only, both Medicaid & private, any other insurance in combination with private, TRICARE or other military insurance); and No insurance (no insurance or Indian Health Service (IHS) only; in Alaska this also includes Alaska Native Health Services and Native Regional Health Corporations that are part of the IHS response option).