



## Prevalence of Selected Maternal and Child Health Indicators for Maine, Pregnancy Risk Assessment Monitoring System (PRAMS), 2016-2017

Health Indicator	2016		2017		Overall 2017 <sup>§</sup>
	N*	% (95% CI) <sup>†</sup>	N*	% (95% CI) <sup>†</sup>	% (95% CI) <sup>†</sup>
<b>Nutrition</b>					
• Multivitamin use ≥4 times a week during the month before pregnancy	817	44.7 (40.5-48.8)	870	45.9 (42.1-49.8)	40.4 (39.7-41.2)
<b>Pre-pregnancy Weight</b>					
• Underweight (Body Mass Index [BMI]<18.5 kg/m <sup>2</sup> )	806	2.3 (1.4-3.8)	854	3.3 (2.2-5.1)	3.7 (3.4-4.0)
• Overweight (BMI 25-29.9 kg/m <sup>2</sup> )	806	22.6 (19.4-26.3)	854	24.9 (21.7-28.4)	25.6 (24.9-26.3)
• Obese (BMI≥30 kg/m <sup>2</sup> )	806	28.8 (25.1-32.7)	854	25.3 (22.0-28.8)	25.3 (24.6-26.0)
<b>Substance Use</b>					
• Any cigarette smoking					
• During the 3 months before pregnancy	803	24.7 (20.9-28.8)	864	27.3 (24.0-31.0)	17.7 (17.1-18.3)
• During the last 3 months of pregnancy	802	14.3 (11.2-18.1)	864	13.8 (11.4-16.8)	8.1 (7.7-8.6)
• Postpartum	809	18.4 (15.0-22.4)	863	18.0 (15.2-21.2)	11.7 (11.2-12.2)
• Any e-cigarette use					
• During the 3 months before pregnancy	804	5.1 (3.3-7.6)	855	4.1 (2.8-6.0)	3.7 (3.4-4.0)
• During the last 3 months of pregnancy	805	1.4 (0.7-2.6)	854	2.5 (1.5-4.1)	1.1 (1.0-1.3)
• Hookah use in the last 2 years	802	2.3 (1.4-3.8)	842	2.9 (1.9-4.6)	4.8 (4.4-5.1)
• Heavy drinking (8 or more drinks a week) during the 3 months before pregnancy	814	3.1 (2.0-4.8)	863	4.2 (2.9-6.1)	2.6 (2.4-2.9)
<b>Intimate Partner Violence (IPV)<sup>¶</sup></b>					
• Experienced IPV during the 12 months before pregnancy by a husband or partner and/or an ex-husband or partner	815	3.5 (2.2-5.5)	865	3.0 (1.9-4.7)	3.0 (2.7-3.2)
• Experienced IPV during pregnancy by a husband or partner and/or an ex-husband or partner	814	2.7 (1.6-4.7)	865	1.8 (1.0-3.1)	2.2 (2.0-2.4)
<b>Depression</b>					
• Self-reported depression in the 3 months before pregnancy	814	20.5 (17.3-24.2)	860	20.1 (17.2-23.4)	12.8 (12.3-13.3)
• Self-reported depression during pregnancy	802	15.8 (12.8-19.2)	858	15.9 (13.2-18.9)	12.0 (11.5-12.5)
• Self-reported postpartum depressive symptoms <sup>‡</sup>	812	11.4 (8.8-14.7)	860	13.4 (11.0-16.4)	12.5 (12.0-13.0)
<b>Health Care Services</b>					
• Health care visit in the 12 months before pregnancy	811	74.5 (70.4-78.2)	865	74.1 (70.5-77.4)	67.7 (67.0-68.4)
• Began prenatal care in 1st trimester	803	89.4 (85.9-92.1)	865	90.3 (87.6-92.4)	86.9 (86.4-87.4)
• Had a flu shot in the 12 months before birth	797	62.3 (58.0-66.4)	863	64.0 (60.2-67.6)	58.9 (58.2-59.7)
• Had maternal postpartum checkup	810	91.4 (88.1-93.9)	865	93.7 (91.6-95.3)	90.3 (89.8-90.7)
<b>Pregnancy Intention</b>					
• Mistimed	806	16.4 (13.4-19.9)	860	15.2 (12.6-18.2)	19.5 (18.9-20.1)
• Unwanted pregnancy	806	4.3 (2.8-6.6)	860	5.2 (3.7-7.2)	6.1 (5.7-6.5)
• Unsure whether wanted pregnancy	806	16.3 (13.2-20.1)	860	14.4 (11.8-17.4)	15.5 (15.0-16.1)
• Intended pregnancy	806	63.0 (58.7-67.2)	860	65.2 (61.4-68.8)	58.9 (58.1-59.6)

Health Indicator	2016		2017		Overall 2017 <sup>§</sup>
	N*	% (95% CI) <sup>†</sup>	N*	% (95% CI) <sup>†</sup>	% (95% CI) <sup>†</sup>
<b>Postpartum<sup>§§</sup> Family Planning</b>					
• Use of any postpartum contraception**	801	76.4 (72.4-80.0)	857	74.5 (70.9-77.7)	77.0 (76.3-77.6)
• Highly effective contraceptive methods					
• Male or female sterilization	801	12.2 (9.7-15.1)	857	11.2 (9.0-13.9)	11.2 (10.7-11.6)
• Long acting reversible contraceptive method <sup>††</sup>	801	20.8 (17.5-24.5)	857	20.7 (17.7-24.0)	17.0 (16.5-17.6)
• Moderately effective contraceptive methods <sup>††</sup>	801	24.9 (21.5-28.7)	857	22.6 (19.5-26.0)	25.6 (24.9-26.3)
• Least effective contraceptive methods <sup>††</sup>	801	18.5 (15.5-21.9)	857	19.9 (17.0-23.2)	23.2 (22.5-23.8)
<b>Oral Health</b>					
• Teeth cleaned during pregnancy by a dentist or dental hygienist	800	41.9 (37.8-46.1)	862	46.3 (42.5-50.2)	46.3 (45.5-47.0)
<b>Health Insurance Status One Month Before Pregnancy<sup>††</sup></b>					
• Private insurance	795	63.1 (58.7-67.4)	838	62.7 (58.8-66.5)	63.0 (62.2-63.7)
• Medicaid	795	21.4 (17.7-25.7)	838	24.0 (20.7-27.6)	23.0 (22.3-23.6)
• No insurance	795	15.5 (12.5-18.9)	838	13.3 (10.8-16.3)	13.2 (12.6-13.7)
<b>Health Insurance Status for Prenatal Care<sup>††</sup></b>					
• Private insurance	789	60.4 (55.9-64.7)	841	60.3 (56.4-64.1)	60.7 (59.9-61.4)
• Medicaid	789	38.1 (33.9-42.6)	841	36.9 (33.2-40.8)	35.6 (34.9-36.4)
• No insurance	789	1.5 (0.8-2.8)	841	2.8 (1.7-4.5)	2.6 (2.3-2.9)
<b>Health Insurance Status Postpartum<sup>††§§</sup></b>					
• Private insurance	801	57.4 (53.0-61.7)	850	57.8 (53.8-61.6)	58.5 (57.7-59.2)
• Medicaid	801	32.4 (28.3-36.8)	850	30.6 (27.1-34.4)	29.7 (29.0-30.4)
• No insurance	801	10.2 (7.8-13.2)	850	11.6 (9.3-14.4)	10.8 (10.3-11.3)
<b>Infant Sleep Practices</b>					
• Baby most often laid on back to sleep	787	85.4 (82.1-88.2)	842	89.2 (86.5-91.4)	79.5 (78.9-80.2)
<b>Breastfeeding Practices</b>					
• Ever breastfed	790	89.7 (86.6-92.2)	842	90.7 (88.2-92.8)	87.7 (87.2-88.2)
• Any breastfeeding at 8 weeks	788	70.1 (65.8-74.0)	842	73.1 (69.5-76.4)	67.7 (67.0-68.4)

<sup>§</sup>PRAMS site aggregate for 2017: Alabama, Alaska, Colorado, Connecticut, Delaware, Georgia, Illinois, Iowa, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Missouri, Montana, New Hampshire, New Jersey, New Mexico, New York City, New York State, North Carolina, North Dakota, Oklahoma, Pennsylvania, Puerto Rico, Rhode Island, South Dakota, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, and Wyoming met the required 55% response rate threshold for inclusion.

\*Unweighted sample size

<sup>†</sup>Weighted percent (95% Confidence Interval)

<sup>††</sup>Defined as being pushed, hit, slapped, kicked, choked, or physically hurt in any way by a husband/partner and/or an ex-husband/ex-partner. In 2016 (Phase 8), the question response options were expanded to include "my ex-husband or ex-partner" in addition to "my husband or partner"

<sup>§§</sup>Postpartum is defined as the time the PRAMS survey was completed

<sup>§</sup>Defined as "always" or "often" feeling down, depressed, or hopeless or having little interest or little pleasure in doing things she usually enjoyed since delivery.

\*\*Defined as using any kind of birth control postpartum (female or male sterilization; intrauterine device (IUD); contraceptive implant; birth control pills; shots or injections; contraceptive patch; vaginal ring; condoms; rhythm method/natural family planning; withdrawal).

<sup>†††</sup>Long-acting reversible contraception (LARC) methods include Intrauterine Device (IUD) or contraceptive implant. Moderately effective methods include birth control pills, shots or injections (e.g., Depo-Provera), contraceptive patch, and vaginal ring. Least effective methods include condom, rhythm method/natural family planning, and withdrawal. Women who selected the "other" write-in option were excluded from the analysis.

<sup>§§§</sup>Insurance is coded as Medicaid (Medicaid or state-named Medicaid program); Private (Private only, any other insurance in combination with private, TRICARE or other military insurance); No insurance (no insurance or Indian Health Service (IHS) only; in Alaska this also includes Alaska Tribal Health System that are part of the IHS response option); other state-specific government plans or programs such as SCHIP/CHIP are excluded from estimates.