

Hawaii. Estimated numbers and percentages of women aged 18–49 years who were at risk for unintended pregnancy* and had ongoing or potential need for contraceptive services,† by selected sociodemographic characteristics — Behavioral Risk Factor Surveillance System, 2019

Sociodemographic characteristics	Women aged 18–49 years			
	Total no. [§]	% at risk for unintended pregnancy (95% CI)	No. and % who had ongoing or potential need for contraceptive services	
			No. (95% CI) [§]	% (95% CI)
Age group				
18-24	54,500	69.4 (62.0-76.0)	36,600 (32,400-40,300)	67.1 (59.4-73.9)
25-34	93,400	66.3 (60.0-72.1)	56,200 (50,200-62,000)	60.2 (53.7-66.4)
35-44	90,900	74.0 (68.6-78.7)	51,600 (45,800-57,200)	56.8 (50.4-62.9)
45-49	36,400	75.6 (67.9-82.0)	19,900 (16,900-22,900)	54.8 (46.3-63.0)
Race/ethnicity				
White, non-Hispanic	54,700	77.0 (70.2-82.7)	33,700 (29,500-37,600)	61.6 (53.9-68.7)
Black, non-Hispanic	¶	¶	¶	¶
Hispanic	35,000	65.7 (55.6-74.5)	20,500 (16,900-23,900)	58.6 (48.3-68.2)
Other	175,800	69.6 (65.6-73.4)	105,300 (97,700-112,700)	59.9 (55.6-64.1)
Insurance coverage				
Yes	252,600	71.0 (67.6-74.1)	150,800 (141,500-160,100)	59.7 (56.0-63.4)
No	22,200	67.5 (54.5-78.3)	12,900 (9,900-15,600)	58.0 (44.8-70.1)
Routine checkup within past year				
Yes	216,600	70.7 (67.0-74.2)	130,000 (121,100-138,600)	60.0 (55.9-64.0)
No	57,100	70.5 (63.4-76.8)	33,600 (29,400-37,700)	58.9 (51.5-66.0)
County urbanization				
Urban	275,300	70.8 (67.5-73.8)	164,400 (154,400-174,000)	59.7 (56.1-63.2)
Rural	¶	¶	¶	¶

Abbreviation: CI = confidence interval.

*Women were considered to be at risk for unintended pregnancy unless they reported 1) not being sexually active with a male partner, 2) being currently pregnant or seeking pregnancy, 3) not minding being pregnant, or 4) having had a hysterectomy.

†Women with ongoing or potential need for contraceptive services were defined as women considered to be at risk for unintended pregnancy not using permanent contraception (female sterilization or male partner vasectomy). The number of women with ongoing or potential need for contraceptive services can be used to estimate how many women might seek services.

§Weighted numbers are rounded to the nearest 100.

¶Estimate is unreliable (unweighted sample size <50 or relative standard error >30%).

