

Iowa. Estimated numbers and percentages of women aged 18–49 years who were at risk for unintended pregnancy* and had ongoing or potential need for contraceptive services,† by selected sociodemographic characteristics — Behavioral Risk Factor Surveillance System, 2019

Sociodemographic characteristics	Women aged 18–49 years			
	Total no. [§]	% at risk for unintended pregnancy (95% CI)	No. and % who had ongoing or potential need for contraceptive services	
			No. (95% CI) [§]	% (95% CI)
Age group				
18-24	160,000	77.0 (70.9-82.2) [¶]	119,500 (109,300-128,300)	74.7 (68.3-80.2) [¶]
25-34	189,500	76.4 (71.8-80.5)	113,300 (103,500-122,600)	59.8 (54.6-64.7)
35-44	187,500	84.2 (80.6-87.2)	85,100 (76,700-93,800)	45.4 (40.9-50.0)
45-49	76,400	80.0 (74.3-84.7)	30,800 (26,000-35,800)	40.3 (34.0-46.8)
Race/ethnicity				
White, non-Hispanic	497,500	80.6 (78.1-82.9)	283,600 (268,200-298,500)	57.0 (53.9-60.0)
Black, non-Hispanic	29,500	70.7 (54.0-83.3)	16,200 (11,500-20,700)	54.9 (38.9-70.0)
Hispanic	48,300	73.5 (64.0-81.2)	27,000 (22,400-31,500)	56.0 (46.4-65.2)
Other	34,700	75.2 (59.5-86.2)	19,000 (13,700-24,000)	54.7 (39.4-69.1)
Insurance coverage				
Yes	554,300	79.7 (77.1-82.0)	314,800 (298,200-330,900)	56.8 (53.8--59.7)
No	53,100	75.1 (65.8-82.5)	28,500 (23,300-33,600)	53.6 (43.8-63.2)
Routine checkup within past year				
Yes	480,000	78.9 (76.2-81.5)	271,200 (255,800-286,600)	56.5 (53.3-59.7)
No	124,800	80.2 (74.6-84.8)	69,800 (61,900-77,500)	55.9 (49.6-62.1)
County urbanization				
Urban	462,900	79.1 (76.2-81.7)	264,800 (249,500-280,100)	57.2 (53.9-60.5)
Rural	150,500	80.2 (75.6-84.1)	83,500 (75,300-91,700)	55.5 (50.0-60.9)

Abbreviation: CI = confidence interval.

*Women were considered to be at risk for unintended pregnancy unless they reported 1) not being sexually active with a male partner, 2) being currently pregnant or seeking pregnancy, 3) not minding being pregnant, or 4) having had a hysterectomy.

†Women with ongoing or potential need for contraceptive services were defined as women considered to be at risk for unintended pregnancy not using permanent contraception (female sterilization or male partner vasectomy). The number of women with ongoing or potential need for contraceptive services can be used to estimate how many women might seek services.

[§]Weighted numbers are rounded to the nearest 100.

[¶]p<0.05 for chi-squared test comparing the distribution of the outcome by the sociodemographic characteristic.

