

Louisiana. Estimated numbers and percentages of women aged 18–49 years who were at risk for unintended pregnancy* and had ongoing or potential need for contraceptive services,† by selected sociodemographic characteristics — Behavioral Risk Factor Surveillance System, 2019

Sociodemographic characteristics	Women aged 18–49 years			
	Total no. [§]	% at risk for unintended pregnancy (95% CI)	No. and % who had ongoing or potential need for contraceptive services	
			No. (95% CI) [§]	% (95% CI)
Age group				
18-24	217,400	75.5 (66.7-82.7)	155,900 (135,000-173,100)	71.7 (62.1-79.6) [¶]
25-34	330,900	82.1 (75.8-86.9)	239,900 (216,400-260,400)	72.5 (65.4-78.7)
35-44	296,100	81.1 (74.6-86.2)	168,800 (147,200-189,500)	57.0 (49.7-64.0)
45-49	124,100	78.2 (70.2-84.5)	64,300 (53,400-75,200)	51.8 (43.0-60.6)
Race/ethnicity				
White, non-Hispanic	492,200	77.8 (72.9-82.0)	285,500 (258,900-311,100)	58.0 (52.6-63.2) [¶]
Black, non-Hispanic	351,400	81.4 (75.5-86.1)	255,100 (231,600-275,500)	72.6 (65.9-78.4)
Hispanic	**	**	**	**
Other	42,100	84.7 (69.9-92.9)	28,400 (20,100-34,600)	67.4 (47.8-82.3)
Insurance coverage				
Yes	844,600	79.8 (76.1-83.0)	542,200 (505,900-576,900)	64.2 (59.9-68.3)
No	114,900	79.2 (67.9-87.2)	76,600 (62,400-88,600)	66.7 (54.3-77.1)
Routine checkup within past year				
Yes	753,400	80.2 (76.3-83.6)	483,700 (449,000-516,100)	64.2 (59.6-68.5)
No	190,700	79.9 (71.9-86.1)	130,100 (113,500-144,400)	68.2 (59.5-75.7)
County urbanization				
Urban	901,900	80.3 (76.8-83.3)	589,800 (552,900-625,000)	65.4 (61.3-69.3)
Rural	66,500	72.6 (52.7-86.3)	34,300 (23,000-45,400)	51.6 (34.6-68.2)

Abbreviation: CI = confidence interval.

*Women were considered to be at risk for unintended pregnancy unless they reported 1) not being sexually active with a male partner, 2) being currently pregnant or seeking pregnancy, 3) not minding being pregnant, or 4) having had a hysterectomy.

†Women with ongoing or potential need for contraceptive services were defined as women considered to be at risk for unintended pregnancy not using permanent contraception (female sterilization or male partner vasectomy). The number of women with ongoing or potential need for contraceptive services can be used to estimate how many women might seek services.

§Weighted numbers are rounded to the nearest 100.

¶p<0.05 for chi-squared test comparing the distribution of the outcome by the sociodemographic characteristic.

**Estimate is unreliable (unweighted sample size <50 or relative standard error >30%).

