

**South Dakota. Estimated numbers and percentages of women aged 18–49 years who were at risk for unintended pregnancy\* and had ongoing or potential need for contraceptive services,† by selected sociodemographic characteristics — Behavioral Risk Factor Surveillance System, 2019**

Sociodemographic characteristics	Women aged 18–49 years			
	Total no. <sup>§</sup>	% at risk for unintended pregnancy (95% CI)	No. and % who had ongoing or potential need for contraceptive services	
			No. (95% CI) <sup>§</sup>	% (95% CI)
<b>Age group</b>				
18-24	41,500	90.8 (83.1-95.2) <sup>¶</sup>	37,400 (34,100-39,300)	90.0 (82.1-94.7) <sup>¶</sup>
25-34	53,900	73.9 (63.6-82.2)	31,300 (25,400-36,800)	58.1 (47.2-68.2)
35-44	51,300	81.6 (72.0-88.5)	26,700 (21,500-31,900)	52.1 (42.0-62.1)
45-49	23,600	84.2 (72.7-91.4)	9,700 (7,000-12,700)	41.0 (29.5-53.7)
<b>Race/ethnicity</b>				
White, non-Hispanic	136,200	82.3 (77.3-86.5)	80,600 (71,800-89,100)	59.2 (52.7-65.4)
Black, non-Hispanic	**	**	**	**
Hispanic	**	**	**	**
Other	21,100	88.1 (78.4-93.8)	16,400 (13,500-18,300)	77.5 (64.0-86.9)
<b>Insurance coverage</b>				
Yes	140,600	82.3 (76.9-86.7)	86,000 (77,000-94,500)	61.2 (54.8-67.2)
No	27,000	78.2 (63.7-88.0)	15,400 (11,100-19,400)	57.2 (41.0-71.9)
<b>Routine checkup within past year</b>				
Yes	119,100	81.0 (74.8-85.9)	74,800 (66,600-82,500)	62.8 (55.9-69.3)
No	48,300	82.3 (72.4-89.2)	26,600 (21,000-32,000)	55.1 (43.5-66.2)
<b>County urbanization</b>				
Urban	132,000	81.9 (76.0-86.6)	81,300 (72,100-89,900)	61.6 (54.6-68.1)
Rural	38,200	79.6 (69.5-87.0)	21,700 (17,500-25,600)	56.8 (45.9-67.0)

Abbreviation: CI = confidence interval.

\*Women were considered to be at risk for unintended pregnancy unless they reported 1) not being sexually active with a male partner, 2) being currently pregnant or seeking pregnancy, 3) not minding being pregnant, or 4) having had a hysterectomy.

†Women with ongoing or potential need for contraceptive services were defined as women considered to be at risk for unintended pregnancy not using permanent contraception (female sterilization or male partner vasectomy). The number of women with ongoing or potential need for contraceptive services can be used to estimate how many women might seek services.

§Weighted numbers are rounded to the nearest 100.

¶p<0.05 for chi-squared test comparing the distribution of the outcome by the sociodemographic characteristic.

\*\*Estimate is unreliable (unweighted sample size <50 or relative standard error >30%).

