

# Insufficient Sleep Among Idaho Adults

Sleep, like food and water, is essential for life.

## Consequences of insufficient sleep.

Insufficient sleep has been linked to the onset of and correlates with a number of chronic diseases and conditions, including diabetes, cardiovascular disease, obesity, and depression. Insufficient sleep also contributes to motor vehicle crashes and machinery-related accidents, causing substantial injury and disability each year.<sup>1</sup>

## How much sleep do we need?

Although how much sleep is needed varies between individuals, most adults need 7–9 hours of sleep each night.<sup>2</sup> More than a third of U.S. adults report sleeping less than 7 hours per night.<sup>3</sup>

## Why don't we get the sleep we need?

Causes of insufficient sleep include lifestyle and occupational factors (e.g., access to technology and work hours).<sup>1</sup> In addition, some medical conditions, medications, and sleep disorders affect the quantity and quality of sleep.<sup>1</sup>

## Getting the sleep we need.

Good sleep practices are important for achieving healthy sleep.

### Sleep hygiene tips:

- Go to bed at the same time each night and rise at the same time each morning.
- Moderate physical activity may help promote sleep, but avoid vigorous exercise in the few hours before going to bed.
- Avoid large meals before bedtime.
- Avoid caffeine and alcohol close to bedtime.
- Avoid nicotine.

### The sleep environment:


- Your bedroom should be a quiet, dark, and relaxing environment, that is neither too hot nor too cold.
- Remove all TVs, computers, and other “gadgets” from the bedroom.
- Your bed should be comfortable and used only for sleeping and not for other activities, such as reading, watching TV, or listening to music.

1 Institute of Medicine. *Sleep Disorders and Sleep Deprivation: An Unmet Public Health Problem*. Washington, DC: The National Academies Press; 2006.

2 National Sleep Foundation. How much sleep do we really need? Washington, DC: National Sleep Foundation; 2010. Available at <http://www.sleepfoundation.org/article/how-sleep-works/how-much-sleep-do-we-really-need>.

3 CDC. Effect of short sleep duration on daily activities—United States, 2005–2008. *MMWR* 2011;60:239–42.


## Prevalence of Insufficient Sleep\* Among Idaho Adults by Region



\*Percentage of adult population that reported  $\geq 14$  days of insufficient sleep in the past 30 days.

For 2008–2009, 27.1% of Idaho adults reported not getting enough sleep on  $\geq 14$  days in the past 30 days. The map above presents the prevalence of insufficient sleep among Idaho adults by state region. For comparison, the national map below shows state-by-state adult prevalence of insufficient sleep.

## Adult Prevalence of Insufficient Sleep\* by State or Territory, 2008–2009



\*Percentage of adult population that reported  $\geq 14$  days of insufficient sleep in the past 30 days.

## How's your sleep?

You may suffer from a sleep disorder if:

1. You frequently have difficulty sleeping (e.g., trouble falling asleep or staying asleep, feeling unrefreshed after sleep).
2. You snore loudly or you or others have observed that you stop breathing or gasp for breath during sleep.
3. You suffer from excessive sleepiness during the day.
4. You have unpleasant, tingling, creeping feelings or nervousness in your legs when trying to sleep.

## What to do if you have trouble sleeping.

- Practice good sleep hygiene.
- Consult your physician to discuss any of the problems above.
- Keep a sleep diary to discuss with your physician or sleep specialist.

For more information, go to

<http://www.cdc.gov/sleep>

The table to the right breaks down the prevalence of insufficient sleep among Idaho adults by sex, age, race/ethnicity, education, employment status, marital status, presence of children in the home, and body mass index (a measure of excess weight).

The prevalence of self-reported insufficient sleep was **higher** among women (29.9%) than men (24.3%) ( $p < 0.05$ ).

Respondents were **more likely** ( $p < 0.05$ ) to report insufficient sleep if they:

- Were aged 25–34 (32.8%) or 35–44 years (32.5%) compared to  $\geq 55$  years
- Were of other race or multiracial (40.6%) compared to white individuals (26.9%)
- Were unable to work (48.4%) compared to other employment status categories
- Had a child living in the home (33.1%)
- Were obese (34.5%) compared to other body mass index categories

Respondents were **less likely** ( $p < 0.05$ ) to report insufficient rest or sleep if they:

- Were retired (14.4%) compared to other employment status categories
- Did not have a child living in the home (21.8%)

There were no statistically significant differences in the prevalence of self-reported insufficient sleep for groups defined by marital status.

## Prevalence of Insufficient Rest or Sleep ( $\geq 14$ days in past 30 days) Among Idaho Adults, 2008–2009 BRFSS\*

(N= 10,305)

	%	(95% CI)
<b>Total</b>	27.1	(25.9 – 28.3)
<b>Sex</b>		
Men	24.3	(22.4 – 26.1)
Women	29.9	(28.3 – 31.5)
<b>Age</b>		
18–24	29.4	(23.8 – 35.0)
25–34	32.8	(29.5 – 36.2)
35–44	32.5	(29.6 – 35.4)
45–54	28.1	(25.7 – 30.4)
55–64	22.6	(20.6 – 24.6)
$\geq 65$	14.6	(13.2 – 16.1)
<b>Race/Ethnicity</b>		
White	26.9	(25.6 – 28.2)
Hispanic	26.1	(20.5 – 31.6)
Black	-	- -
Asian	-	- -
American Indian/Alaska Native	33.1	(21.7 – 44.6)
Native Hawaiian/Pacific Islander	-	- -
Other/Multiracial	40.6	(28.7 – 52.5)
<b>Employment Status</b>		
Employed	28.0	(26.4 – 29.7)
Unemployed	31.8	(25.7 – 37.9)
Retired	14.4	(12.8 – 16.0)
Unable to work	48.4	(42.7 – 54.1)
Homemaker or student	28.6	(25.0 – 32.2)
<b>Marital Status</b>		
Married	26.1	(24.7 – 27.5)
Divorced, widowed, separated	28.3	(25.9 – 30.6)
Member of unmarried couple	25.5	(17.5 – 33.5)
Never married	30.9	(26.1 – 35.8)
<b>Children in Home</b>		
No	21.8	(20.4 – 23.2)
Yes	33.1	(31.1 – 35.2)
<b>Body Mass Index</b>		
Underweight	17.9	(10.5 – 25.2)
Normal	24.5	(22.3 – 26.7)
Overweight	24.5	(22.6 – 26.5)
Obese	34.5	(32.0 – 37.1)

\*Data source: Behavioral Risk Factor Surveillance Survey (BRFSS) for 2008 and 2009. As part of the phone survey, respondents were asked, "During the past 30 days, for about how many days have you felt you did not get enough rest or sleep?" Dashes (-) indicate a cell size  $< 50$ . For information about BRFSS methodology, go to <http://www.cdc.gov/BRFSS>.