

Rhode Island, 2007

Primary and secondary syphilis

- **CASES:** Rhode Island reported 36 cases in 2007.
- **RATES:** Rhode Island ranked 20 among 50 states, DC, and 3 territories with 3.4 cases per 100,000 population compared to the total rate of 3.8 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Rhode Island, the rate among males was 6.6 per 100,000 population compared to the total male rate of 6.6 per 100,000.
- The rate among females was 0.4 per 100,000 compared to the total female rate of 1.1 per 100,000.

Rates of primary and secondary syphilis, by race/ethnicity


In Rhode Island, the race/ethnicity adjusted rates per 100,000 population were 3.0 among whites, 6.9 among blacks, 2.5 among Hispanics, 3.3 among Asian/Pacific Islanders, and 19.0 among American Indians/Alaska Natives.

- The rate among blacks was 2.3 times that of whites.
- The rate among Asian/Pacific Islanders was 1.1 times that of whites.
- The rate among American Indians/Alaska Natives was 6.3 times that of whites.

Congenital syphilis

- **CASES:** Rhode Island reported 0 cases in 2007.
- **RATES:** Rhode Island had a rate of 0.0 cases per 100,000 live births compared to the total rate of 10.5 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Rhode Island, 1998-2007



Total P&S Syphilis Rates by County, Rhode Island, 2007

