Alabama, 2010

Primary and secondary syphilis

- CASES: Alabama reported 260 cases in 2010.
- RATES: Alabama ranked 12 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 5.5 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Alabama, the rate among males was 8.1 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 3.1 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.

Rates of primary and secondary syphilis, by race/ethnicity

- In Alabama, the race/ethnicity adjusted rates per 100,000 population were 1.1 among whites, 17.6 among blacks, 2.6 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 16 times that of whites.
- The rate among Hispanics was 2.4 times that of whites.

Congenital syphilis

- CASES: Alabama reported 9 cases in 2010.
- RATES: Alabama ranked 9 among among 30 areas (28 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 13.9 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Alabama, 2001-2010



Total P&S Syphilis Rates by County, Alabama, 2010



Rate (per 100,000 population)

<=0.4 | 0.41-2.2 | >2.2