

# District of Columbia, 2010

## Primary and secondary syphilis

- **CASES:** District of Columbia reported 134 cases in 2010.
- **RATES:** District of Columbia ranked 1 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 22.3 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In District of Columbia, the rate among males was 46.6 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.6 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


## Rates of primary and secondary syphilis, by race/ethnicity

- In District of Columbia, the race/ethnicity adjusted rates per 100,000 population were 22.1 among whites, 22.1 among blacks, 20.7 among Hispanics, 19.5 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.

## Congenital syphilis

- **CASES:** District of Columbia reported 1 case in 2010.
- **RATES:** District of Columbia ranked 13 among among 30 areas (28 states; Washington, DC; and 1 territory) reporting at least one case of congenital syphilis with a rate of 11.3 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, District of Columbia, 2001-2010



Total P&S Syphilis Rate, District of Columbia, 2010

