

Kentucky, 2010

Primary and secondary syphilis

- **CASES:** Kentucky reported 139 cases in 2010.
- **RATES:** Kentucky ranked 24 among 52 areas (49 states; Washington, DC; and 2 territories) reporting at least one case of P&S syphilis with 3.2 cases per 100,000 population compared to the U.S. rate of 4.5 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Kentucky, the rate among males was 6.2 per 100,000 population compared to the U.S. male rate of 7.9 per 100,000.
- The rate among females was 0.4 per 100,000 compared to the U.S. female rate of 1.1 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Kentucky, the race/ethnicity adjusted rates per 100,000 population were 2.2 among whites, 12.7 among blacks, 6.9 among Hispanics, 2.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 5.8 times that of whites.
- The rate among Hispanics was 3.1 times that of whites.


Congenital syphilis

- **CASES:** Kentucky reported 0 cases in 2010.
- **RATES:** Kentucky had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.7 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Kentucky, 2001-2010



Total P&S Syphilis Rates by County, Kentucky, 2010



Rate (per 100,000 population)

